

MULTIPLICATIVE FUNCTIONS COMMUTABLE WITH SUMS OF SQUARES

POO-SUNG PARK

ABSTRACT. Let k be an integer greater than or equal 3. We show that if a multiplicative function f satisfies

$$f(x_1^2 + x_2^2 + \cdots + x_k^2) = f(x_1)^2 + f(x_2)^2 + \cdots + f(x_k)^2$$

for all positive integers x_i , then f is the identity function.

1. INTRODUCTION

In 1992, Claudia Spiro [15] proved that if a multiplicative function $f : \mathbb{N} \rightarrow \mathbb{C}$ satisfies $f(p+q) = f(p) + f(q)$ for all primes p and q , then $f(n) = n$ under the condition $f(p_0) \neq 0$ for some prime p_0 . The term *multiplicative function* means $f(1) = 1$ and $f(mn) = f(m)f(n)$ for all coprime integers m and n .

It has influenced lots of mathematicians to develop her problem into diverse variations. Fang [9] showed a similar result for $f(p+q+r) = f(p) + f(q) + f(r)$ and Dubickas and Šarka [7] extended those results to $f(p_1 + p_2 + \cdots + p_k) = f(p_1) + f(p_2) + \cdots + f(p_k)$. Chung [4] classified multiplicative functions satisfying the equation $f(m^2 + n^2) = f(m^2) + f(n^2)$ for all $m, n \in \mathbb{N}$. Recently, Bašić [1] characterized all arithmetic functions such that $f(m^2 + n^2) = f(m)^2 + f(n)^2$, which is slightly different from Chung's condition. Many related problems are found in [2, 3, 4, 5, 6, 12, 13, 14]. We would call such problems *Spiro problems*.

In this article, we add new results to the list of Spiro problems by proving the following: the condition

$$f(x_1^2 + x_2^2 + \cdots + x_k^2) = f(x_1)^2 + f(x_2)^2 + \cdots + f(x_k)^2$$

for all positive integers x_i forces a multiplicative function f to be the identity function for $k \geq 3$.

2. RESULTS

For the consistency, we mention the sum of two squares.

Theorem 1. *If a multiplicative function f satisfies*

$$f(x^2 + y^2) = f(x)^2 + f(y)^2$$

for all positive integers x and y , then $f(n^2) = f(n)^2 = n^2$.

Proof. This is a consequence of Bašić's characterization of arithmetic functions satisfying the abovementioned condition [1]. \square

Key words and phrases. multiplicative function; sum of squares.

This work was supported by Kyungnam University Foundation Grant, 2016.

To prove the main theorem, we need lemmas about sums of three or more nonvanishing squares. See [10, Chapter 6], [11], [8].

Lemma 1 (Hurwitz). *The only squares that are not sums of three nonvanishing squares are the integers 4^s and $25 \cdot 4^s$ with $s \geq 0$.*

Lemma 2 (Dubouis). *Every integer n can be represented by the sum of k nonvanishing squares except*

$$n = \begin{cases} 1, 3, 5, 9, 11, 17, 29, 41, 2 \cdot 4^m, 6 \cdot 4^m, 14 \cdot 4^m & (m \geq 0) & \text{if } k = 4, \\ 1, 2, 3, 4, 6, 7, 9, 10, 12, 15, 18, 33 & & \text{if } k = 5, \\ 1, 2, \dots, k-1, k+1, k+2, k+4, k+5, k+7, k+10, k+13 & & \text{if } k > 5. \end{cases}$$

Since sums of k nonvanishing squares represent almost all positive integers for $k \geq 5$, we separate the problem into three cases: $k = 3$, $k = 4$, and $k \geq 5$.

Theorem 2. *If a multiplicative function f satisfies*

$$f(x^2 + y^2 + z^2) = f(x)^2 + f(y)^2 + f(z)^2$$

for all positive integers x, y and z , then f is the identity function.

Proof. First, we compute $f(n)$ for $n \leq 15$ and $n = 25$.

Since f is multiplicative, $f(1) = 1$. Clearly, $f(3) = f(1^2 + 1^2 + 1^2) = 3$. To find $f(2)$ let us consider

$$f(6) = f(1^2 + 1^2 + 2^2) = f(1)^2 + f(1)^2 + f(2)^2 = 2 + f(2)^2.$$

Since $f(6) = f(2)f(3) = 3f(2)$, we find that $f(2) = 1$ or $f(2) = 2$ from the equation $2 + f(2)^2 = 3f(2)$. To determine which value $f(2)$ is we need more relations.

From $f(12) = f(2^2 + 2^2 + 2^2) = 3f(2)^2 = f(3)f(4) = 3f(4)$ we have $f(4) = f(2)^2$.

Furthermore, from $f(14) = f(1^2 + 2^2 + 3^2) = 10 + f(2)^2 = f(2)f(7)$ we observe that $f(7) = f(2) + \frac{10}{f(2)}$.

To find another relation between $f(2)$ and $f(7)$, consider

$$f(21) = f(1^2 + 2^2 + 4^2) = 1 + f(2)^2 + f(2)^4 = f(3)f(7) = 3f(7).$$

From the equation

$$1 + f(2) + f(2)^2 = 3 \left(f(2) + \frac{10}{f(2)} \right)$$

we can conclude that $f(2) = 2$ and thus $f(4) = 4$, $f(6) = 6$, $f(7) = 7$, $f(12) = 12$, and $f(14) = 14$.

To find $f(5)$ we examine $f(27)$ and $f(30)$. Since

$$\begin{aligned} f(27) &= f(1^2 + 1^2 + 5^2) = 2 + f(5)^2 \\ &= f(3^2 + 3^2 + 3^2) = 3f(3)^2 = 27, \end{aligned}$$

we have $f(5) = \pm 5$. On the other hand,

$$\begin{aligned} f(30) &= f(1^2 + 2^2 + 5^2) = 5 + f(5)^2 \\ &= f(5)f(6) = 6f(5). \end{aligned}$$

Thus, we have that $f(5) = 5$.

Since

$$\begin{aligned} f(50) &= f(3^2 + 4^2 + 5^2) = 50 \\ &= f(2)f(25) = 2f(25), \end{aligned}$$

we have $f(25) = 25$ and $f(50) = 50$.

Now, to find $f(8)$ consider $f(24)$. Since

$$\begin{aligned} f(24) &= f(2^2 + 2^2 + 4^2) = 24 \\ &= f(3)f(8) = 3f(8), \end{aligned}$$

we have that $f(8) = 8$.

Similarly, it follows from

$$\begin{aligned} f(26) &= f(1^2 + 3^2 + 4^2) = 26 \\ &= f(2)f(13) = 2f(13) \end{aligned}$$

that $f(13) = 13$.

Other $f(n)$'s can be calculated as follows:

$$\begin{aligned} f(9) &= f(1^2 + 2^2 + 2^2) = 9, & f(10) &= f(2)f(5) = 10, \\ f(11) &= f(1^2 + 1^2 + 3^2) = 11, & f(15) &= f(3)f(5) = 15. \end{aligned}$$

Now consider $n > 15$. We use induction. Assume that $f(k) = k$ for all positive integers $k < n$.

If $n = a^2 + b^2 + c^2$ with $a, b, c \geq 1$, then

$$f(n) = f(a^2 + b^2 + c^2) = f(a)^2 + f(b)^2 + f(c)^2 = a^2 + b^2 + c^2 = n$$

since $f(a) = a$, $f(b) = b$, $f(c) = c$ by induction hypothesis.

If n cannot be represented as a sum of three nonvanishing squares, this means two cases:

- (1) if $n = a^2 + b^2 + c^2$ for some $a, b, c \in \mathbb{Z}$, then at least one of a, b, c should vanish.
- (2) $n \neq a^2 + b^2 + c^2$ for any $a, b, c \in \mathbb{Z}$.

Consider case (1): Suppose $n = a^2 + b^2$ with $3 \leq a \leq b$. Note that

$$5a < 2a^2 \leq a^2 + b^2 = n, \quad 3b < 4b \leq b^2 + 4 < b^2 + a^2 = n.$$

If $5 \nmid n$, then

$$\begin{aligned} f(25n) &= f(25a^2 + 25b^2) = f((5a)^2 + (3b)^2 + (4b)^2) \\ &= f(5a)^2 + f(3b)^2 + f(4b)^2 \\ &= (5a)^2 + (3b)^2 + (4b)^2 = 25n \end{aligned}$$

by the induction hypothesis. Since n is assumed to be indivisible by 5, $f(25n) = f(25)f(n) = 25f(n)$ and $f(n) = n$.

If $n = a^2 + b^2$ with $1 \leq a \leq 2$ and $4 \leq b$, then

$$5a \leq 10 < n, \quad 3b < 4b \leq b^2 < a^2 + b^2 = n$$

and thus $f(n) = n$ by the same reasoning.

It is already checked that $f(n) = n$ for $a \leq 2$ and $b \leq 3$ in the previous step.

If $n = a^2 + b^2$ is divisible by 5, let $n = 5^s k$ with $s \geq 1$ and $5 \nmid k$. We have shown that $f(5) = 5$ and $f(25) = 25$. If $s > 2$ is even, 5^s can be represented as a sum of three nonvanishing squares by Lemma 1. So $f(5^s) = 5^s$.

If $s \geq 3$ is odd, then $s - 3$ is even and

$$5^s = 5^{s-3} \cdot 125 = 5^{s-3}(3^2 + 4^2 + 10^2).$$

Thus, $f(5^s) = 5^s$.

We can conclude that $f(n) = f(5^s k) = f(5^s)f(k) = 5^s k = n$.

Now we should investigate the case of $n = a^2$. By Lemma 1 it suffices to check whether $f(4^s) = 4^s$. Note that

$$\begin{aligned} f(3 \cdot 4^s) &= f((2^s)^2 + (2^s)^2 + (2^s)^2) = f(2^s)^2 + f(2^s)^2 + f(2^s)^2 = 3 \cdot 4^s \\ &= f(3)f(4^s) = 3f(4^s). \end{aligned}$$

Thus $f(4^s) = 4^s$ and we can conclude that $f(n) = n$ for the case (1).

Now consider the case (2): It is well known that n cannot be represented as a sum of three squares if and only if n is of the form $4^s(8t+7)$ with $s, t \geq 0$. We may assume that $n = 8t+7$. Then, since $2(8t+7) = 8(2t+1)+6$ can be represented as a sum of three squares of integers, $f(2(8t+7)) = 2(8t+7)$ and thus $f(8t+7) = 8t+7$ by canceling $f(2) = 2$.

From the case (1) and (2) we can conclude that $f(n) = n$. \square

By interweaving the method in the above proof with Fermat's Two-Square Theorem, we can also prove Theorem 1 in the similar way.

Theorem 3. *If a multiplicative function f satisfies*

$$f(x_1^2 + x_2^2 + x_3^2 + x_4^2) = f(x_1)^2 + f(x_2)^2 + f(x_3)^2 + f(x_4)^2$$

for all positive integers x_i , then f is the identity function.

Proof. By Lemma 2 every positive integer can be represented as sums of four non-vanishing squares except for

$$1, 3, 5, 9, 11, 17, 29, 41, 2 \cdot 4^m, 6 \cdot 4^m, 14 \cdot 4^m \quad (m \geq 0).$$

Note that $f(1) = 1$ and $f(4) = 4$.

Since

$$\begin{aligned} f(12) &= f(3)f(4) = 4f(3) \\ &= f(1)^2 + f(1)^2 + f(1)^2 + f(3)^2 = 3 + f(3)^2, \end{aligned}$$

we obtain that $f(3) = 1$ or $f(3) = 3$.

Let us consider $f(5)$. Since

$$\begin{aligned} f(20) &= f(4)f(5) = 4f(5) \\ &= f(1)^2 + f(1)^2 + f(3)^2 + f(3)^2 = 2 + 2f(3)^2, \end{aligned}$$

$f(5) = 1$ or $f(5) = 5$ according to the value of $f(3)$.

Similarly, from

$$\begin{aligned} f(28) &= f(4)f(7) = 4f(7) \\ &= f(1)^2 + f(3)^2 + f(3)^2 + f(3)^2 = 1 + 3f(3)^2 \end{aligned}$$

we have that $f(7) = 1$ or $f(7) = 7$ according to the value of $f(3)$.

Note that

$$\begin{aligned} f(35) &= f(5)f(7) \\ &= f(1)^2 + f(3)^2 + f(3)^2 + f(4)^2 = 17 + 2f(3)^2. \end{aligned}$$

Hence we conclude that $f(3) = 3$, $f(5) = 5$, and $f(7) = 7$. Then $f(2) = 2$ also follows from

$$\begin{aligned} f(10) &= f(2)f(5) \\ &= f(1)^2 + f(1)^2 + f(2)^2 + f(2)^2 = 2 + 2f(2)^2 \end{aligned}$$

and

$$f(7) = f(1)^2 + f(1)^2 + f(1)^2 + f(2)^2 = 3 + f(2)^2.$$

If we consider

$$\begin{aligned} f(18) &= f(2)f(9) = 2f(9) \\ &= f(1)^2 + f(2)^2 + f(2)^2 + f(3)^2 = 18, \end{aligned}$$

we have that $f(9) = 9$. The similar observation gives us

$$f(11) = 11, \quad f(17) = 17, \quad f(29) = 29, \quad f(41) = 41.$$

Now suppose that $f(2 \cdot 4^r) = 2 \cdot 4^r$ for $r < m$. Note that

$$f(6 \cdot 4^r) = f(3)f(2 \cdot 4^r) = 6 \cdot 2^r \quad \text{and} \quad f(14 \cdot 4^r) = f(7)f(2 \cdot 4^r) = 14 \cdot 2^r.$$

Thus $f(n) = n$ for $n < 2 \cdot 4^m$ by Lemma 2. Since $5 \cdot 2 \cdot 4^m$ can be represented as a sum of squares of four positive integers less than $2 \cdot 4^m$, we can conclude that $f(5 \cdot 2 \cdot 4^m) = 5 \cdot 2 \cdot 4^m$. On the other hand,

$$f(5 \cdot 2 \cdot 4^m) = f(5)f(2 \cdot 4^m) = 5f(2 \cdot 4^m)$$

and thus $f(2 \cdot 4^m) = 2 \cdot 4^m$.

Now consider $6 \cdot 4^m$. Since $f(6 \cdot 4^m) = f(3)f(2 \cdot 4^m)$, we obtain that $f(6 \cdot 4^m) = 6 \cdot 4^m$. Similarly, $f(14 \cdot 4^m) = f(7)f(2 \cdot 4^m) = 14 \cdot 4^m$. \square

Theorem 4. *Let k be an integer ≥ 5 . If a multiplicative function f satisfies*

$$f(x_1^2 + x_2^2 + \cdots + x_k^2) = f(x_1)^2 + f(x_2)^2 + \cdots + f(x_k)^2$$

for all positive integers x_i , then f is the identity function.

Proof. After we prove for $k = 5, 6, 7$ in the way similar to those above, the case of $k \geq 8$ will be proved finally.

i) Assume that $k = 5$. By Lemma 2 every positive integer can be represented by the sum of five nonvanishing squares except for

$$1, 2, 3, 4, 6, 7, 9, 10, 12, 15, 18, 33.$$

We have $f(1) = 1$ and $f(5) = 5$.

From

$$\begin{aligned} f(20) &= f(4)f(5) = 5f(4) \\ &= f(1)^2 + f(1)^2 + f(1)^2 + f(1)^2 + f(4)^2 = 4 + f(4)^2 \\ &= f(2)^2 + f(2)^2 + f(2)^2 + f(2)^2 + f(2)^2 = 5f(2)^2 \end{aligned}$$

we obtain $f(4) = 1$ or $f(4) = 4$. Also, $f(2) = \pm 1$ or $f(2) = \pm 2$ according to the value of $f(4)$.

From

$$\begin{aligned} f(29) &= f(1)^2 + f(1)^2 + f(1)^2 + f(1)^2 + f(5)^2 = 29 \\ &= f(1)^2 + f(1)^2 + f(3)^2 + f(3)^2 + f(3)^2 = 2 + 3f(3)^2 \\ &= f(1)^2 + f(2)^2 + f(2)^2 + f(2)^2 + f(4)^2 = 1 + 3f(2)^2 + f(4)^2 \end{aligned}$$

we obtain that $f(29) = 29$, $f(2) = \pm 2$, and $f(3) = \pm 3$. Also, this fixes $f(4) = 4$, $f(8) = 8$, and $f(11) = 11$.

Then, it follows from

$$\begin{aligned} f(2 \cdot 11) &= f(2)f(11) = 11f(2) \\ &= f(1)^2 + f(2)^2 + f(2)^2 + f(2)^2 + f(3)^2 = 22 \end{aligned}$$

and

$$\begin{aligned} f(3 \cdot 8) &= f(3)f(8) = 8f(2) \\ &= f(1)^2 + f(1)^2 + f(2)^2 + f(2)^2 + f(4)^2 = 24 \end{aligned}$$

that $f(2) = 2$, $f(3) = 3$, and $f(6) = 6$.

Now, we use induction to show that $f(n) = n$ for $n \geq 7$. Assume that $f(m) = m$ for $m < n$. Then $n(n-1) \geq 42$ and it can be represented as a sum of squares of five positive integers less than n . Thus

$$f(n(n-1)) = n(n-1).$$

Since n and $n-1$ are relatively prime,

$$f(n(n-1)) = f(n)f(n-1) = f(n)(n-1)$$

by induction hypothesis and thus $f(n) = n$.

ii) Assume that $k = 6$. By Lemma 2 every positive integer can be represented by the sum of six nonvanishing squares except for

$$1, 2, 3, 4, 5, 7, 8, 10, 11, 13, 16, 19.$$

The proof is similar to the previous case. Since $n(n-1) > 19$ for $n \geq 6$, it suffices to show that $f(n) = n$ for $n \leq 5$. Clearly, $f(1) = 1$ and $f(6) = 6$. From the equalities

$$\begin{aligned} f(30) &= f(5)f(6) = 6f(5) \\ &= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 5^2) = 5 + f(5)^2 \\ &= f(1^2 + 1^2 + 1^2 + 3^2 + 3^2 + 3^2) = 3 + 3f(3)^2 \\ &= f(1^2 + 1^2 + 2^2 + 2^2 + 2^2 + 4^2) = 2 + 3f(2)^2 + f(4)^2, \\ f(41) &= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 6^2) = 41 \\ &= f(1^2 + 1^2 + 1^2 + 2^2 + 3^2 + 5^2) = 3 + f(2)^2 + f(3)^2 + f(5)^2, \\ f(21) &= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 4^2) = 5 + f(4)^2 \\ &= f(1^2 + 2^2 + 2^2 + 2^2 + 2^2 + 2^2) = 1 + 5f(2)^2 \end{aligned}$$

we deduce that

$$f(2) = \pm 2, \quad f(3) = \pm 3, \quad f(4) = \pm 4, \quad f(5) = 5.$$

Since $f(17) = 17$, $f(2 \cdot 17) = 2 \cdot 17$, $f(3 \cdot 17) = 3 \cdot 17$, and $f(4 \cdot 17) = 4 \cdot 17$ by Lemma 2, we obtain that $f(n) = n$ for $n \leq 5$. If $n \geq 6$, $n(n-1) > 19$ and thus we can show $f(n) = n$.

iii) Assume that $k = 7$. By Lemma 2 every positive integer can be represented by the sum of seven nonvanishing squares except for

$$1, 2, 3, 4, 5, 6, 8, 9, 11, 12, 14, 17, 20.$$

We have $f(1) = 1$ and $f(7) = 7$. Note that

$$\begin{aligned}
f(31) &= f(1^2 + 1^2 + 1^2 + 1^2 + 3^2 + 3^2 + 3^2) = 4 + 3f(3)^2 \\
&= f(1^2 + 1^2 + 1^2 + 2^2 + 2^2 + 2^2 + 4^2) = 3 + 3f(2)^2 + f(4)^2 \\
&= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 5^2) = 6 + f(5)^2, \\
f(42) &= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 2^2 3^2) = 6 + f(2)^2 f(3)^2 \\
&= f(1^2 + 2^2 + 2^2 + 2^2 + 2^2 + 3^2 + 4^2) = 1 + 4f(2)^2 + f(3)^2 + f(4)^2 \\
&= f(1^2 + 1^2 + 1^2 + 1^2 + 2^2 + 3^2 + 5^2) = 4 + f(2)^2 + f(3)^2 + f(5)^2, \\
f(55) &= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 7^2) = 55 \\
&= f(1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 5^2 + 5^2) = 5 + 2f(5)^2.
\end{aligned}$$

Thus, we can find that

$$f(2) = \pm 2, \quad f(3) = \pm 3, \quad f(4) = \pm 4, \quad f(5) = \pm 5.$$

Note that $n(n-1) > 20$ for $n \geq 6$. We can conclude that $f(n) = n$ for every positive integer by the similar way.

iv) Assume that $k \geq 8$. By Lemma 2 every positive integer can be represented by the sum of k nonvanishing squares except for

$$1, 2, \dots, k-1, k+1, k+2, k+4, k+5, k+7, k+10, k+13.$$

Since $f(1) = 1$, $f(k) = k$, and

$$\begin{aligned}
f(k(k-1)) &= f(k)f(k-1) = kf(k-1) \\
&= f(k-1 + (k-1)^2) = \underbrace{f(1)^2 + \dots + f(1)^2}_{k-1 \text{ summands}} + f(k-1)^2,
\end{aligned}$$

the value of $f(k-1)$ is either 1 or $k-1$.

Since

$$\begin{aligned}
40 &= 1^2 + 1^2 + 1^2 + 1^2 + 6^2 \\
&= 2^2 + 3^2 + 3^2 + 3^2 + 3^2,
\end{aligned}$$

we obtain that

$$(k-5)f(1)^2 + 4f(1)^2 + (f(2)f(3))^2 = (k-5)f(1)^2 + f(2)^2 + 4f(3)^2$$

or

$$4 + (f(2)f(3))^2 = f(2)^2 + 4f(3)^2.$$

Also, from

$$\begin{aligned}
32 &= 1^2 + 1^2 + 1^2 + 1^2 + 1^2 + 3^2 + 3^2 + 3^2 \\
&= 2^2 + 2^2 + 2^2 + 2^2 + 2^2 + 2^2 + 2^2 + 2^2,
\end{aligned}$$

we obtain that $5 + 3f(3)^2 = 8f(2)^2$.

By solving the system of these two equations we can find the following solutions:

$$f(2) = \pm 1, f(3) = \pm 1 \quad \text{or} \quad f(2) = \pm 2, f(3) = \pm 3.$$

Now, let $3a+8b=2k-1$ with some nonnegative integers a and b . This is possible unless $2k-1=1, 2, 4, 5, 7, 10, 13$. That is, if $2k-1 > 13$, then it is represented in such a way. The number 13 is called the *Frobenius number* of 3 and 8. Note that

$$a+b < \frac{2k-1}{3} + \frac{2k-1}{8} = \frac{11}{24}(2k-1) < k-1$$

and

$$(k-1) + (2k-1) + (k-1)^2 = k^2 + k - 1.$$

Thus, since $2^2a + 3^2b = (a+b) + (3a+8b) = (a+b) + (2k-1)$,

$$\begin{aligned} f(k^2 + k - 1) &= f(k)^2 + (k-1)f(1)^2 = k^2 + k - 1 \\ &= f\left(\underbrace{1^2 + \cdots + 1^2}_{k-a-b-1 \text{ summands}} + \underbrace{2^2 + \cdots + 2^2}_a + \underbrace{3^2 + \cdots + 3^2}_b + (k-1)^2\right) \\ &= (k-a-b-1) + af(2)^2 + bf(3)^2 + f(k-1)^2. \end{aligned}$$

Therefore

$$f(2) = \pm 2, \quad f(3) = \pm 3, \quad f(k-1) = k-1.$$

Since

$$\begin{aligned} 4^2 + 2^2 + 2^2 + 2^2 &= 1^2 + 3^2 + 3^2 + 3^2, & 5^2 + 1^2 + 1^2 &= 3^2 + 3^2 + 3^2, \\ 6 &= 2 \cdot 3, & 7^2 + 1^2 &= 5^2 + 5^2, \\ 8^2 + 1^2 &= 4^2 + 7^2, & 9^2 + 2^2 &= 6^2 + 7^2, \\ 10 &= 2 \cdot 5, \end{aligned}$$

we obtain that $f(n) = \pm n$ for $n \leq 10$.

Note that

$$(2\ell+1)^2 + (\ell-2)^2 = (2\ell-1)^2 + (\ell+2)^2 \text{ with } \ell \geq 5$$

and

$$(2\ell)^2 + (\ell-5)^2 = (2\ell-4)^2 + (\ell+3)^2 \text{ with } \ell \geq 6.$$

Hence, by induction, $f(n) = \pm n$ for every positive integer n .

Thus, by Lemma 2, $f(n) = n$ for all $n > k+13$. For each $1 < n \leq k+13$, let $m = n(k+13) + 1$, we have $f(m) = m$ and $f(nm) = nm$ since $nm > m > k+13$. It follows from $(n, m) = 1$ that $f(n) = n$.

□

ACKNOWLEDGMENTS

The author would like to thank referees for their kind and valuable comments.

REFERENCES

- [1] B. Bašić, Characterization of arithmetic functions that preserve the sum-of-squares operation, *Acta Math. Sin.* **30** (2014), 689–695.
- [2] K.-K. Chen and Y.-G. Chen, On $f(p) + f(q) = f(p+q)$ for all odd primes p and q , *Publ. Math. Debrecen* **76** (2010), 425–430.
- [3] Y.-G. Chen, J.-H. Fang, P. Yuan, and Y. Zheng, On multiplicative functions with $f(p+q+n_0) = f(p) + f(q) + f(n_0)$, *J. Number Theory* **165** (2016), 270–289.
- [4] P. V. Chung, Multiplicative functions satisfying the equation $f(m^2 + n^2) = f(m^2) + f(n^2)$, *Math. Slovaca* **46** (1996), 165–171.

- [5] P. V. Chung and B. M. Phong, Additive uniqueness sets for multiplicative functions, *Publ. Math. Debrecen* **55** (1999), 237–243.
- [6] J.-M. De Koninck, I. Kátai, and B. M. Phong, A new characteristic of the identity function, *J. Number Theory* **63** (1997), 325–338.
- [7] A. Dubickas and P. Šarka, On multiplicative functions which are additive on sums of primes, *Aequat. Math.* **86** (2013), 81–89.
- [8] E. Dubouis, Solution of a problem of J. Tannery, *Intermédiaire Math.* **18** (1911), 55–56.
- [9] J.-H. Fang, A characterization of the identity function with equation $f(p + q + r) = f(p) + f(q) + f(r)$, *Combinatorica*, **31** (2011), 697–701.
- [10] E. Grosswald, *Representations of Integers as Sums of Squares*, Springer, 2011.
- [11] A. Hurwitz, *Intérmediaire Math.* 14 (1907), 107; *Werke*, vol.2, 751.
- [12] K.-H. Indlekofer and B. M. Phong, Additive uniqueness sets for multiplicative functions, *Ann. Univ. Sci. Budapest. Sect. Comput.* **26** (2006), 65–77.
- [13] B. M. Phong, On sets characterizing the identity function, *Ann. Univ. Sci. Budapest. Sect. Comput.* **24** (2004), 295–306.
- [14] B. M. Phong, A characterization of the identity function with an equation of Hosszú type, *Publ. Math. Debrecen* **69** (2006), 219–226.
- [15] C. A. Spiro, Additive uniqueness sets for arithmetic functions, *J. Number Theory* 42 (1992), 232–246.

DEPARTMENT OF MATHEMATICS EDUCATION, KYUNGNAM UNIVERSITY, CHANGWON, REPUBLIC OF KOREA

Email address: pspark@kyungnam.ac.kr