

The Lawrence-Krammer representation is a quantization of the symmetric square of the Burau representation ¹

A.V. Kosyak

Institute of Mathematics, Ukrainian National Academy of Sciences, 3 Tereshchenkivs'ka Str., Kyiv, 01601, Ukraine

Royal Institution, London Institute for Mathematical Sciences, 21 Albemarle St, London W1S 4BS, UK

Contents

1	Introduction	2
2	Connection between the Lawrence–Krammer and the Burau representations	2
2.1	The Burau representation	2
2.2	The Burau representation of B_∞	3
2.3	The Krammer representation of the braid group B_n , different notations	4
2.4	Representations of B_{n+1} via representations of the Lie algebra \mathfrak{gl}_n . . .	4
2.5	Connection between the Krammer and the reduced Burau representation	5
2.6	Lie algebra \mathfrak{sl}_2 , its universal enveloping algebra $U(\mathfrak{sl}_2)$ and $U(\mathfrak{sl}_2)$ -modules	6
2.7	Pascal triangle and representations of B_3 and Lie algebra \mathfrak{sl}_2	7
2.8	q -Pascal triangle and representations of B_3	7
2.9	Quantization of the Pascal triangle	9
2.10	Symmetric basis in $V \otimes V$	10
2.11	Proof of Theorem 2.2	10
3	How to construct new representations of the braid group B_n	16
3.1	Quantization of the m^{th} symmetric power of the Burau representation .	16
3.2	Representations of a Lie algebra \mathfrak{sl}_{n+1}	17
4	Some polynomial invariants for knots	18
4.1	Some polynomial invariants for the trefoil, see [4]	18
4.2	The Burau representation and the Alexander polynomial	19
4.3	The Krammer representation and the corresponding rational function .	19
4.4	Alexander polynomial for the trefoil	20
4.5	Krammer representation and the trefoil	20

¹To all fearless Ukrainians defending not only their country, but the whole civilization against putin's [rashism](#)

1. Introduction

In 1936 W. Burau introduced [11] the family of representations of the braid group B_n as the *deformation* of the *standard representation* of the symmetric group S_n (see (1)). The Burau representation for $n = 2, 3$ has been known to be faithful for some time. Indeed, in 1969 Magnus and Peluso [23] showed that the Burau and *Gasner representations* are faithful for $n = 2, 3$. See also J. Birman [9, Theorem 3.15]. Moody [24] in 1991 showed that the Burau representation is not faithful for $n \geq 9$; this result was improved to $n \geq 6$ by Long and Paton in 1992, [22]. The non-faithfulness for $n = 5$ was shown by Bigelow in 1999, [5]. The faithfulness of the Burau representation for $n = 4$ is an *open problem*.

In 1990 R. Lawrence [21] have constructed two-parameter family of representation of the braid group B_n , using the homological methods. In 2000 D. Krammer [19] using completely algebraic methods have constructed the same representations and showed that this representation is faithful for B_4 . One year later, in 2001, S. Bigelow showed that the Lawrence-Krammer representation is faithful for all B_n . Next year, D. Krammer [20] proved the same result by a different method.

The Burau representation appears as a summand of the *Jones representation* [14], and for $n = 4$, the faithfulness of the Burau representation is equivalent to that of the Jones representation, which on the other hand is related to the question of whether or not the Jones polynomial is an *unknot detector* [8].

In [13] Jackson and Kerler established an isomorphism between Lawrence-Krammer representation and the submodule of the R -matrix representations on $V^{\otimes n}$ for the *generic Verma module* V of the quantum group $U_q(\mathfrak{sl}_2)$.

From [13]: “In [33] Zinno manages to find a different identification of the Lawrence-Krammer representation with a quantum algebraic object, namely the quotient of the Birman-Wenzl-Murakami algebra similarly defined over $\mathbb{Z}[q^\pm, t^\pm]$. This representation can, by [30], be understood as the one arising from the quantum orthogonal groups $U_q(\mathfrak{so}(k+1))$ acting on the n -fold tensor product of the fundamental representation.” See [10, n 4.6, p.71] for more details about connections between the Lawrence-Krammer and other representations of braid groups.

2. Connection between the Lawrence–Krammer and the Burau representations

2.1. The Burau representation

The Burau representation $\rho : B_n \rightarrow \mathrm{GL}_n(\mathbb{Z}[t, t^{-1}])$ is defined for a non-zero complex number t by

$$\sigma_i \mapsto I_{i-1} \oplus \begin{pmatrix} 1-t & t \\ 1 & 0 \end{pmatrix} \oplus I_{n-i-1} \quad (1)$$

where $1-t$ is the (i, i) entry. For $t = 1$ this is the *standard representation* of the symmetric group S_n interchanging by matrix $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ two neighbors basic elements e_i and e_{i+1} in the space \mathbb{C}^n . The vector $e = e_1 + \dots + e_n$ is invariant, therefore the representation ρ splits into 1-dimensional and an $n-1$ -dimensional irreducible representations, known as the *reduced Burau representation* $\rho_n^{(t)} : B_n \rightarrow \mathrm{GL}_{n-1}(\mathbb{Z}[t, t^{-1}])$

$$\sigma_1 \mapsto \begin{pmatrix} -t & 0 \\ -1 & 1 \end{pmatrix} \oplus I_{n-3}, \quad \sigma_{n-1} \mapsto I_{n-3} \oplus \begin{pmatrix} 1 & -t \\ 0 & -t \end{pmatrix}, \quad (2)$$

$$\sigma_i \mapsto I_{i-2} \oplus \begin{pmatrix} 1 & -t & 0 \\ 0 & -t & 0 \\ 0 & -1 & 1 \end{pmatrix} \oplus I_{n-i-2}, \quad 2 \leq i \leq n-2. \quad (3)$$

We use the following equivalent form of the reduced Burau representation $\rho_n^{(t)} : B_n \mapsto \text{GL}_{n-1}(\mathbb{Z}[t, t^{-1}])$, compare with (2) and (3):

$$\sigma_1 \mapsto \begin{pmatrix} -t & t \\ 0 & 1 \end{pmatrix} \oplus I_{n-3}, \quad \sigma_{n-1} \mapsto I_{n-3} \oplus \begin{pmatrix} 1 & 0 \\ 1 & -t \end{pmatrix}, \quad (4)$$

$$\sigma_i \mapsto I_{i-2} \oplus \begin{pmatrix} 1 & 0 & 0 \\ 1 & -t & t \\ 0 & 0 & 1 \end{pmatrix} \oplus I_{n-i-2}, \quad 2 \leq i \leq n-2. \quad (5)$$

The equivalence follows from the following:

$$J^{-1} \begin{pmatrix} 1 & -t & 0 \\ 0 & -t & 0 \\ 0 & -1 & 1 \end{pmatrix} J = \begin{pmatrix} 1 & 1 & 0 \\ 0 & -t & 0 \\ 0 & t & 1 \end{pmatrix} \xrightarrow{(\cdot)^t} \begin{pmatrix} 1 & 0 & 0 \\ 1 & -t & t \\ 0 & 0 & 1 \end{pmatrix}, \quad \text{where } J = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

2.2. The Burau representation of B_∞

Define the group B_∞ as follows

$$B_\infty = \langle (\sigma_i)_{i \in \mathbb{Z}} \mid \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \quad i \in \mathbb{Z}, \quad \sigma_i \sigma_j = \sigma_i \sigma_j, \quad |i - j| \geq 2 \rangle. \quad (6)$$

Consider a complex Hilbert space

$$H = l_2(\mathbb{Z}) = \{x = (x_k)_{k \in \mathbb{Z}} \mid \|x\|^2 = \sum_{k \in \mathbb{Z}} |x_k|^2 < \infty\} \quad (7)$$

with its standard orthonormal basis $e = (e_k)_{k \in \mathbb{Z}}$ defined by $e_k = (\delta_{k,n})_{n \in \mathbb{Z}}$. Define the reduced Burau representation of the group B_∞ in the space H as follows

$$\sigma_i \mapsto b_{i,\infty} = I_\infty \oplus \begin{pmatrix} 1 & 0 & 0 \\ 1 & -t & t \\ 0 & 0 & 1 \end{pmatrix} \oplus I_\infty, \quad i \in \mathbb{Z}, \quad (8)$$

where $-t$ is the (i, i) entry. It is clear that $b_{i+1,\infty}$ is obtained from $b_{i,\infty}$ by shifting the basis $e : e_n \rightarrow e_{n+1}$. More precisely, if we denote by J the unitary operator on $l_2(\mathbb{Z})$ defined as follows: $Je_n = e_{n+1}$, $n \in \mathbb{Z}$ we get

$$Jb_{i,\infty}J^{-1} = b_{i+1,\infty}. \quad (9)$$

Next, denote by P_n the orthogonal projector of H on the subspace H_n generated by the first n vectors, i.e., $H_n = \langle e_k \mid 1 \leq k \leq n \rangle$. Then we get

$$P_n b_{k,\infty} P_n = \rho_{n+1}^{(t)}(\sigma_k), \quad 1 \leq k \leq n. \quad (10)$$

Denote by J_n the unitary operator on H_n defined as follows

$$J_n e_k = e_{k+1}, \quad 1 \leq k \leq n-1, \quad J_n e_n = e_1, \quad (11)$$

and by i_n the natural embedding of $\text{GL}(n-1, \mathbb{C})$ into $\text{GL}(n, \mathbb{C})$ defined as follows:

$$\text{GL}(n-1, \mathbb{C}) \ni x \rightarrow i_n(x) = x + E_{nn} \in \text{GL}(n, \mathbb{C}). \quad (12)$$

Then we get

$$J_n i_n(\rho_n^{(t)}(\sigma_k)) J_n^{-1} = \rho_{n+1}^{(t)}(\sigma_k), \quad 2 \leq k \leq n. \quad (13)$$

2.3. The Krammer representation of the braid group B_n , different notations

The *Lawrence-Krammer representation* $K_n^{(t,q)}$ of the Braid group B_n was introduced by Lawrence in [21] and studied by Krammer in [19, 20] and Bigelow in [6, 7]. To give formulas, we follow Bigelow [6] and the corrected version in [7]. $K_n^{(t,q)}$ is the following action of B_n on a free module V of rank $\binom{n}{2} = \frac{n!}{2!(n-2)!}$ with basis $\{F_{i,j} : 1 \leq i < j \leq n\}$:

$$\sigma_i(F_{j,k}) = \begin{cases} F_{j,k} & i \notin \{j-1, j, k-1, k\}, \\ qF_{i,k} + q(q-1)F_{i,j} + (1-q)F_{j,k} & i = j-1, \\ F_{j+1,k} & i = j \neq k-1, \\ qF_{j,i} + (1-q)F_{j,k} + q(1-q)tF_{i,k} & i = k-1 \neq j, \\ F_{j,k+1} & i = k, \\ -tq^2F_{j,k} & i = j = k-1. \end{cases} \quad (14)$$

From [7]: “This action is given in [19], except that Krammer’s $-t$ is my t . It is also in [6], but with a sign error. The name “Krammer representation” was chosen because Krammer seems to have initially found this independently of Lawrence and without any use of homology”.

We would like to change notations and define the *Krammer representation* $k_n^{(t,q)}$ as follows:

$$\sigma_i(F_{j,k}) = \begin{cases} F_{j,k} & i \notin \{j-1, j, k-1, k\}, \\ tF_{i,k} + t(t-1)F_{i,j} + (1-t)F_{j,k} & i = j-1, \\ F_{j+1,k} & i = j \neq k-1, \\ tF_{j,i} + (1-t)F_{j,k} + t(t-1)qF_{i,k} & i = k-1 \neq j, \\ F_{j,k+1} & i = k, \\ qt^2F_{j,k} & i = j = k-1. \end{cases} \quad (15)$$

Connection between two representations is as follows:

$$K_n^{(-q,t)}(\sigma_r) = k_n^{(t,q)}(\sigma_r), \quad 1 \leq r \leq n-1. \quad (16)$$

We think that this choice of the parameters reflects better the situation since, historically Burau used parameter t , and q , in our notations, is really a *parameter of quantization* (see (24)).

2.4. Representations of B_{n+1} via representations of the Lie algebra \mathfrak{gl}_n

We can generalize the reduced Burau representation of B_n just observing (see [17]) that $\rho_n^{(t)}$ is a product of several \exp 's of an appropriate *Serre generators* of the *natural* (or *fundamental*) representation of the Lie algebra \mathfrak{sl}_n (in fact of \mathfrak{gl}_n) see (23). To be more precise, we show that the reduced Burau representation for B_3

$$\sigma_1 \mapsto \begin{pmatrix} -t & t \\ 0 & 1 \end{pmatrix}, \quad \sigma_2 \mapsto \begin{pmatrix} 1 & 0 \\ 1 & -t \end{pmatrix}, \quad (17)$$

can be rewritten as follows:

$$\sigma_1 \mapsto \begin{pmatrix} -t & t \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} -t & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \exp(sE_{11}) \exp(-E_{12}), \quad (18)$$

$$\sigma_2 \mapsto \begin{pmatrix} 1 & 0 \\ 1 & -t \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -t \end{pmatrix} = \exp(E_{21}) \exp(sE_{22}), \quad (19)$$

where E_{kn} are matrix unities and $s = \ln(-t)$ for negative t .

For $n = 4$ the reduced Burau representation is defined as follows:

$$\begin{aligned}
\rho_4^{(t)} : B_4 &\rightarrow \mathrm{GL}_3(\mathbb{Z}[t, t^{-1}]), \\
\sigma_1 &\mapsto \begin{pmatrix} -t & t & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \sigma_2 \mapsto \begin{pmatrix} 1 & 0 & 0 \\ 0 & -t & t \\ 0 & 0 & 1 \end{pmatrix}, \quad \sigma_3 \mapsto \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & -t \end{pmatrix}, \\
\sigma_1 &\mapsto \begin{pmatrix} -t & t & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} -t & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \exp(sE_{11}) \exp(-E_{12}), \\
\sigma_2 &\mapsto \begin{pmatrix} 1 & 0 & 0 \\ 0 & -t & t \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -t & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix} = \exp(E_{21}) \exp(sE_{22}) \exp(-E_{23}), \\
\sigma_3 &\mapsto \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & -t \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -t \end{pmatrix} = \exp(E_{32}) \exp(sE_{33}),
\end{aligned} \tag{20}$$

where $s = \ln(-t)$ when $t < 0$. Finally, we get

$$\sigma_1 = \exp(sE_{11}) \exp(-X_1), \quad \sigma_2 = \exp(Y_1) \exp(sE_{22}) \exp(-X_2), \tag{21}$$

$$\sigma_3 = \exp(Y_2) \exp(sE_{33}). \tag{22}$$

In the general case, the formulas are similar, see (23). Below we give the *Cartan matrix* A corresponding to a Lie algebra \mathfrak{sl}_n and “the rule to obtain” the right formulas, where $X_k = E_{kk+1}$, $Y_k = E_{k+1k}$ and $H_k = E_{kk} - E_{k+1,k+1}$, $1 \leq k \leq n-1$, are the *Serre generators* of \mathfrak{sl}_n :

$$A = \begin{pmatrix} 2 & -1 & 0 & \dots & 0 & 0 & 0 \\ -1 & 2 & -1 & \dots & 0 & 0 & 0 \\ 0 & -1 & 2 & \dots & 0 & 0 & 0 \\ 0 & 0 & 0 & \dots & 2 & -1 & 0 \\ 0 & 0 & 0 & \dots & -1 & 2 & 0 \end{pmatrix}, \quad \begin{pmatrix} E_{11} & -X_1 & & & & \\ Y_1 & E_{22} & -X_2 & & & \\ & Y_2 & E_{33} & -X_3 & & \\ & & Y_3 & & & \\ & & & \dots & & \\ & & & & -X_{n-2} & \\ & & & & E_{n-1,n-1} & -X_{n-1} \\ & & & & Y_{n-1} & E_{nn} \end{pmatrix}.$$

Lemma 2.1. *Let $\pi : \mathfrak{gl}_n \rightarrow \mathrm{End}(V)$ be an arbitrary representation of the Lie algebra \mathfrak{gl}_n in the space V . Then the following formulas*

$$\begin{aligned}
\rho_{n+1}^\pi(\sigma_1) &\mapsto \exp(s\pi(E_{11})) \exp(-\pi(X_1)), \\
\rho_{n+1}^\pi(\sigma_k) &\mapsto \exp(\pi(Y_{k-1})) \exp(s\pi(E_{kk})) \exp(-\pi(X_k)), \\
\rho_{n+1}^\pi(\sigma_n) &\mapsto \exp(\pi(Y_{n-1})) \exp(s\pi(E_{nn}))
\end{aligned} \tag{23}$$

gives us representation $\rho_{n+1}^\pi : B_{n+1} \rightarrow \mathrm{GL}(V)$ of the braid group B_{n+1} in the space V . For π being the natural representation of the Lie algebra \mathfrak{sl}_n we get the reduced Burau representation $\rho_{n+1}^{(t)}$.

It is sufficient to use the fact that formulas (2) and (3) define representation of the group B_n .

2.5. Connection between the Krammer and the reduced Burau representation

Theorem 2.2. *The Krammer representation $k_n^{(t,q)}$ defined by (15) is equivalent with the quantization of the symmetric square of the reduced Burau representation:*

$$k_n^{(t,q)} = [S^2(\rho_n^{(t)})]_q. \tag{24}$$

The spectrum of the reduced Burau representation is as follows:

$\text{Sp} \rho_n^{(t)}(\sigma_k) = \{-t, \underbrace{1, \dots, 1}_{n-1 \text{ times}}\}$. The spectrum of its symmetric square is:

$$\text{Sp} S^2(\rho_n^{(t)}(\sigma_k)) = \{t^2, \underbrace{-t}_{n-2 \text{ times}}, \dots, \underbrace{-t, 1, \dots, 1}_{(n-1)(n-2)/2 \text{ times}}\}. \quad (25)$$

In [26, Claim 5.8] (see also [27, Lemma 5]) the following fact is proved

Lemma 2.3. *The spectrum of the Krammer representation $k_n^{(t,q)}$ of the group B_n , i.e., $\text{Sp}(\sigma_1^K)$ is as follows:*

$$\text{Sp}(\sigma_1^K) = \{qt^2, \underbrace{-t}_{n-2 \text{ times}}, \dots, \underbrace{-t, 1, \dots, 1}_{(n-1)(n-2)/2 \text{ times}}\}. \quad (26)$$

If we set $q = 1$ in (26) we get (25). Moreover, in [19, Proposition 3.2] the following connection between the Krammer V module and the Burau W module was proved:

Proposition 2.4. *If $t = 1$ and 2 is invertible in R , then there is an isomorphism of B_n -modules $V \rightarrow S^2 W$ (symmetric square of W) given by $v_{ij} \mapsto w_{ij}^2$ and, more generally, $v(T) \mapsto w(T)^2$.*

Remark 1. *In fact, Krammer showed that the Lawrence-Krammer representation is a deformation of the symmetric square of the Burau representation. We show more, i.e., that this deformation is in fact quantization.*

2.6. Lie algebra \mathfrak{sl}_2 , its universal enveloping algebra $U(\mathfrak{sl}_2)$ and $U(\mathfrak{sl}_2)$ -modules

Recall that the Lie algebra \mathfrak{sl}_2 is the Lie algebra of 2×2 real matrices with trace equals to zero. The standard basis X, Y, H in a Lie algebra \mathfrak{sl}_2 is as follows:

$$X = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad Y = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (27)$$

This natural representations $\pi_2 : \mathfrak{sl}_2 \rightarrow \text{End}(\mathbb{C}^2)$ is called *fundamental representation* of the Lie algebra \mathfrak{sl}_2 .

All finite-dimensional U -module V being the highest weight module of highest weight λ are of the following form (see Kassel, [15, Theorem V.4.4.])

$$\rho(n)(X) = S^n(\pi_2(X)) = \begin{pmatrix} 0 & n & 0 & \dots & 0 \\ 0 & 0 & n-1 & \dots & 0 \\ 0 & 0 & 0 & \dots & 1 \\ 0 & 0 & 0 & \dots & 0 \end{pmatrix}, \quad \rho(n)(Y) = S^n(\pi_2(Y)) = \begin{pmatrix} 0 & 0 & \dots & 0 & 0 \\ 1 & 0 & \dots & 0 & 0 \\ 0 & 2 & \dots & 0 & 0 \\ 0 & 0 & \dots & n & 0 \end{pmatrix}, \quad (28)$$

$$\rho(n)(H) = S^n(\pi_2(H)) = \begin{pmatrix} n & 0 & \dots & 0 & 0 \\ 0 & n-2 & \dots & 0 & 0 \\ \dots & \dots & \dots & -n+2 & 0 \\ 0 & 0 & \dots & 0 & -n \end{pmatrix}, \quad (29)$$

where $\lambda = \dim(V) - 1 \in \mathbb{N}$.

2.7. Pascal triangle and representations of B_3 and Lie algebra \mathfrak{sl}_2

The connection of the Pascal triangle with representations of B_3 was firstly noticed by Humphry in [12]. Set in (35) $\Lambda_n = I$ and $q = 1$, then

$$\sigma_1 \mapsto \sigma_1(1, n), \quad \sigma_2 \mapsto \sigma_2(1, n), \quad \text{where} \quad \sigma_2(1, n) = (\sigma_1(1, n)^{-1})^\sharp \quad (30)$$

is an irreducible representations of the group B_3 in \mathbb{C}^{n+1} . Here the *Pascal triangle* denoted by $\sigma_1(1, n)$, for $n - 1 \in \mathbb{N}$, is defined as follows:

$$\sigma_1(1, 1) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad \sigma_1(1, 2) = \begin{pmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \quad \sigma_1(1, 3) = \begin{pmatrix} 1 & 3 & 3 & 1 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (31)$$

Theorem 2.5. *Let π_2 be the fundamental representation of \mathfrak{sl}_2 then*

$$\sigma_1(1, n) = \exp(\rho(X)) = \exp(S^n(\pi_2(X))), \quad (32)$$

$$\sigma_2(1, n) = \exp(\rho(-Y)) = \exp(S^n(\pi_2(-Y))), \quad (33)$$

i.e., the Humphry representation of B_3 in dimension $n + 1$ is the n^{th} symmetric power of the Burau representation ρ_3^{-1} .

2.8. q -Pascal triangle and representations of B_3

We define the family of representations of the group B_3 using q -Pascal triangle [17]. For $q \in \mathbb{C}^\times := \mathbb{C} \setminus \{0\}$ and a matrix

$$\Lambda_n = \text{diag}(\lambda_r)_{r=0}^n \quad \text{such that} \quad \lambda_r \lambda_{n-r} = \text{const} \quad (34)$$

set

$$\sigma_1^\Lambda(q, n) := \sigma_1(q, n) D_n^\sharp(q) \Lambda_n, \quad \sigma_2^\Lambda(q, n) := \Lambda_n^\sharp D_n(q) \sigma_2(q, n), \quad (35)$$

$$D_n(q) = \text{diag}(q_r)_{r=0}^n, \quad q_r = q^{\frac{(r-1)r}{2}}. \quad (36)$$

Here the *central symmetry* $A \mapsto A^\sharp$, for the matrix $A \in \text{Mat}(n+1, \mathbb{C})$, $A = (a_{km})_{0 \leq k, m \leq n}$ is defined as follows:

$$A^\sharp = (a_{km}^\sharp)_{0 \leq k, m \leq n}, \quad a_{km}^\sharp = a_{n-k, n-m}, \quad (37)$$

and $\sigma_1(q, n)$ is the q -Pascal triangle obtained from the Pascal triangle by replacing natural numbers n by q -natural numbers $(n)_q$ (see (42)). More precisely, define $\sigma_1(q, n)$ as follows $\sigma_1(q, n) = (\sigma_1(q, n)_{km})_{0 \leq k, m \leq n}$, where

$$\sigma_1(q, n)_{km} = \binom{n-k}{n-m}_q, \quad 0 \leq k, m \leq n, \quad (38)$$

$$\sigma_2(q, n) := \sigma_1^{-1}(q^{-1}, n)^\sharp, \quad (39)$$

$$(\sigma_1^{-1}(q^{-1}, n))_{km}^\sharp = \begin{cases} 0, & \text{if } 0 < k < m \leq n \\ (-1)^{k+m} q_{k-m}^{-1} C_k^m(q^{-1}), & \text{if } 0 \leq m \leq k \leq n \end{cases}. \quad (40)$$

Here the q -binomial coefficients or *Gaussian polynomials* are defined as follows:

$$C_n^k(q) := \binom{n}{k}_q := \frac{(n)!_q}{(k)!_q (n-k)!_q}, \quad C_n^k[q] := [n]_q := \frac{[n]!_q}{[k]!_q [n-k]!_q} \quad (41)$$

corresponding to two forms of q -natural numbers, defined by

$$(n)_q := \frac{q^n - 1}{q - 1}, \quad [n]_q := \frac{q^n - q^{-n}}{q - q^{-1}}. \quad (42)$$

Define q -Pochhammer symbol

$$(a; q)_n = \prod_{k=0}^{n-1} (1 - aq^k) = (1 - a)(1 - aq)(1 - aq^2) \cdots (1 - aq^{n-1}). \quad (43)$$

We have (see [2])

$$(1 + x)_q^k := (-x; q)_k = \sum_{r=0}^k q^{r(r-1)/2} C_k^r(q) x^r = \sum_{r=0}^k q^{r(r-1)/2} \binom{k}{r}_q x^r. \quad (44)$$

Theorem 2.6 ([1, 17]). Formulas $\sigma_1 \mapsto \sigma_1^\Lambda(q, n)$, $\sigma_2 \mapsto \sigma_2^\Lambda(q, n)$ (see (35)) define family of representation of the group B_3 in dimension $n + 1$.

Remark 2. Let $\pi : B_3 \rightarrow \text{GL}(m, \mathbb{C})$ be a representation. Fix two matrices $\Lambda_1, \Lambda_2 \in \text{GL}(m, \mathbb{C})$ and define $\pi^\Lambda(\sigma_1) = \Lambda_1 \pi(\sigma_1)$, $\pi^\Lambda(\sigma_2) = \pi(\sigma_2) \Lambda_2$. When π^Λ is again a representation of B_3 ? Under the additional conditions $\Lambda_1 \Lambda_2 = cI$, $c \in \mathbb{C}$ and $\Lambda_1 = \text{diag}(\lambda_k)_{k=0}^n$ the criteria for π^Λ to be representation is the following:

$$\Lambda_1 \pi(\sigma_1 \sigma_2 \sigma_1) = \pi(\sigma_1 \sigma_2 \sigma_1) \Lambda_2. \quad (45)$$

This implies condition (34), i.e., $\lambda_r \lambda_{n-r} = c$.

Theorem 2.7 ([28, 1, 17]). All representations of the group B_3 in dimension ≤ 5 are given by the formulas (35).

In dimension 2 we get for $\Lambda_1 = \text{diag}(\lambda_0, \lambda_1)$

$$\sigma_1^{\Lambda_1}(q, 1) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \lambda_0 & 0 \\ 0 & \lambda_1 \end{pmatrix}, \quad \sigma_2^{\Lambda_1}(q, 1) = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}. \quad (46)$$

In dimension 3, we get for $\Lambda_2 = \text{diag}(\lambda_0, \lambda_1, \lambda_2)$ with $\lambda_0 \lambda_2 = \lambda_1^2$

$$\sigma_1^{\Lambda_2}(q, 2) = \begin{pmatrix} \lambda_0 q & (1+q)\lambda_1 & \lambda_2 \\ 0 & \lambda_1 & \lambda_2 \\ 0 & 0 & \lambda_2 \end{pmatrix}, \quad \sigma_2^{\Lambda_2}(q, 2) = \begin{pmatrix} \lambda_2 & 0 & 0 \\ -\lambda_1 & \lambda_1 & 0 \\ \lambda_0 & -\lambda_0(1+q) & \lambda_0 q \end{pmatrix}. \quad (47)$$

In dimension 4, we get for $\Lambda_3 = \text{diag}(\lambda_0, \lambda_1, \lambda_2, \lambda_3)$ with $\lambda_0 \lambda_3 = \lambda_1 \lambda_2$:

$$\sigma_1^{\Lambda_3}(q, 3) = \begin{pmatrix} 1 & (1+q+q^2) & (1+q+q^2) & 1 \\ 0 & 1 & (1+q) & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} D_3^\sharp(q) \Lambda_3, \quad (48)$$

$$\sigma_2^\Lambda(q, 3) = \Lambda_3^\sharp \begin{pmatrix} -1 & 0 & 0 & 0 \\ 1 & -(1+q) & 1 & 0 \\ -1 & (1+q+q^2) & -(1+q+q^2) & 1 \end{pmatrix} D_4(q). \quad (49)$$

In dimension 5, we get for $\Lambda_3 = \text{diag}(\lambda_r)_{r=0}^4$ with $\lambda_0 \lambda_4 = \lambda_1 \lambda_3 = \lambda_2^2$:

$$\sigma_1^\Lambda(q, 4) = \begin{pmatrix} 1 & (1+q)(1+q^2) & (1+q^2)(1+q+q^2) & (1+q)(1+q^2) & 1 \\ 0 & 1 & 1+q+q^2 & 1+q+q^2 & 1 \\ 0 & 0 & 1 & 1+q & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} D_4^\sharp(q) \Lambda_4, \quad (50)$$

$$\sigma_2^\Lambda(q, 4) = \Lambda_4^\sharp \begin{pmatrix} -1 & 0 & 0 & 0 & 0 \\ 1 & -(1+q) & 1 & 0 & 0 \\ -1 & (1+q+q^2) & -(1+q+q^2) & 1 & 0 \\ 1 & -(1+q)(1+q^2) & (1+q^2)(1+q+q^2) & -(1+q)(1+q^2) & 1 \end{pmatrix} D_4(q). \quad (51)$$

2.9. Quantization of the Pascal triangle

Remark 3. The following procedure of quantisation (“Quant”) is based on the quantization of the Pascal triangle explained in [1] and [17], formulas (10) and (11). To be more precise, we change the binomial coefficients $\binom{n}{k}$ of the Pascal triangle by q -binomial coefficients $\binom{n}{k}_q = \frac{(n)!_q}{(k)!_q(n-k)!_q}$.

Set $J_n = \sum_{r=0}^n E_{r,n-r}$ and introduce representations $T_{3,n+1}^{\Lambda_n,q}$ of the braid group B_3 equivalent to (35) in dimension $n+1$ as follows:

$$T_{3,n+1}^{\Lambda_n,q,\pi_2}(\sigma_r) = (\sigma_r^\Lambda(q, n))^\sharp = J_n \sigma_r^\Lambda(q, n) J_n^{-1}, \quad r = 1, 2. \quad (52)$$

Remark 4. Using [29], see also [17, n7, n8], we conclude that all representation of B_3 in dimension 2, 3, 4 and 5 are equivalent with the following ones, where $\Lambda_1 = s \operatorname{diag}(-t, 1)$ and $s, t \in \mathbb{C}^\times$:

$$T_{32}^{\Lambda_1,1,\pi_2}(\sigma_1) = s \begin{pmatrix} -t & t \\ 0 & 1 \end{pmatrix} = s \begin{pmatrix} -t & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}, \quad T_{32}^{\Lambda_1,1,\pi_2}(\sigma_2) = s \begin{pmatrix} 1 & 0 \\ 1 & -t \end{pmatrix} = s \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -t \end{pmatrix}. \quad (53)$$

All representation of B_3 in dimension 3 are given by the following formulas, where $\Lambda_2 = \operatorname{diag}(st^2, -st, s) = s \operatorname{diag}(t^2, -t, 1)$ and $s, t, q \in \mathbb{C}^\times$:

$$T_{33}^{\Lambda_2,q,\pi_2}(\sigma_1) = s \begin{pmatrix} t^2q & -t^2(1+q) & t^2 \\ 0 & -t & t \\ 0 & 0 & 1 \end{pmatrix} = s \begin{pmatrix} t^2 & 0 & 0 \\ 0 & -t & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & -(1+q) & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} q & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (54)$$

$$T_{33}^{\Lambda_2,q,\pi_2}(\sigma_2) = s \begin{pmatrix} 1 & 0 & 0 \\ 1 & -t & t^2q \\ 1 & -t(1+q) & t^2q \end{pmatrix} = s \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & (1+q) & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & q \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -t & 0 \\ 0 & 0 & t^2 \end{pmatrix}. \quad (55)$$

Remark 5. If we are sufficiently attentive, we can recognize here the Lawrence-Krammer representation of B_3 .

All representation of B_3 in dimension 4 are given by the following formulas, where $\Lambda_3 = s \operatorname{diag}(-t^3, ut^2, -u^{-1}t, t)$:

$$T_{34}^{\Lambda_3,q}(\sigma_1) = \Lambda_3 \begin{pmatrix} 1 & -(1+q+q^2) & (1+q+q^2) & -1 \\ 0 & 1 & -(1+q) & 1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{pmatrix} D_3^\sharp(q), \quad (56)$$

$$T_{34}^{\Lambda_3,q}(\sigma_2) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & (1+q) & 1 & 0 \\ 1 & (1+q+q^2) & (1+q+q^2) & 1 \end{pmatrix} D_3(q) \Lambda_3^\sharp. \quad (57)$$

For $n = 5$ we get

$$T_{35}^{\Lambda_4,q}(\sigma_1) = \Lambda_4 \begin{pmatrix} 1 & -(1+q)(1+q^2) & (1+q^2)(1+q+q^2) & -(1+q)(1+q^2) & -1 \\ 0 & 1 & -(1+q+q^2) & (1+q+q^2) & -1 \\ 0 & 0 & 1 & -(1+q) & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} D_4^\sharp(q), \quad (58)$$

$$T_{35}^{\Lambda_4,q}(\sigma_2) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 1 & (1+q) & 1 & 0 & 0 \\ 1 & (1+q+q^2) & (1+q+q^2) & 1 & 0 \\ 1 & (1+q)(1+q^2) & (1+q^2)(1+q+q^2) & (1+q)(1+q^2) & 1 \end{pmatrix} D_4(q) \Lambda_4^\sharp. \quad (59)$$

The general formulas are

$$T_{3,n+1}^{\Lambda_n,q}(\sigma_1) = \Lambda_n D_n^\sharp(q) (\sigma_1^{-1}(q^{-1}, n)), \quad T_{3,n+1}^{\Lambda_n,q}(\sigma_2) = (\sigma_1(q, n))^\sharp D_n(q) \Lambda_n^\sharp. \quad (60)$$

Remark 6. The irreducibility criteria for the representation (35) of B_3 in dimensions $n \leq 5$ were obtained in [28, 29].

Remark 7. For dimensions $n \geq 6$ formulas (35) gives us some family of representations of B_3 but not all, see [29, Remark 2.11.3]. For more representations of B_3 see, e.g., Westbury, [32]. The question of the irreducibility of representations (35) is open for $n \geq 6$.

2.10. Symmetric basis in $V \otimes V$

Let V be a linear space with the basis $(e_k)_{k=1}^n$. The *tensor product* $V \otimes V$ is a linear space generated by the basis $(e_k \otimes e_r)_{k,r=1}^n$. The *symmetric square* $S^2(V)$ of the space V is a subspace of $V \otimes V$ generated by one of the equivalent *symmetric basis* $e = (e_{kn}^s)_{kn}$ or $v = (v_{kn})_{kn}$, defined as follows:

$$e_{kk}^s = e_k \otimes e_k, \quad 1 \leq k \leq n, \quad e_{kr}^s = e_k \otimes e_r + e_r \otimes e_k, \quad 1 \leq k < r \leq n, \quad (61)$$

$$v_{kr} = (e_k + \cdots + e_{r-1}) \otimes (e_k + \cdots + e_{r-1}), \quad 1 \leq k < r \leq n. \quad (62)$$

We fix a *lexicographic order* on the set (k, r) with $1 \leq k \leq r \leq n$. Let an operator A acts on a space V . Denote by $A \otimes A$ its tensor product on a space $V \otimes V$ and by $S^2(A)$ its *symmetric square*.

2.11. Proof of Theorem 2.2

We show that $[S^2(\rho_n^{(t)})_e]_q$, the quantization of the symmetric square $S^2(\rho_n^{(t)})$ of the Burau representations $\rho_n^{(t)}$ for the group B_n , calculated in the basis v , coincide with the Lawrence-Krammer representation $k_n^{(t,q)}$ in notations (15), i.e., we show that

$$C_n[S^2(\rho_n^{(t)}(\sigma_r))_e]_q C_n^{-1} = k_n^{(t,q)}(\sigma_r), \quad \text{for } 1 \leq r \leq n-1, \quad (63)$$

where C_n is the *change-of-basis matrix* from the basis e defined by (61) to the basis v defined by (62) for the vector space $S^2(\mathbb{C}^{n-1})$.

The Burau representation $\rho_3^{(t)}$ for B_3 on the space $V = \mathbb{C}^2$ is as follows:

$$\sigma_1 \mapsto \begin{pmatrix} -t & t \\ 0 & 1 \end{pmatrix}, \quad \sigma_2 \mapsto \begin{pmatrix} 1 & 0 \\ 1 & -t \end{pmatrix}. \quad (64)$$

We fix the standard basis $e = (e_k)_{k=1}^n$ in \mathbb{C}^n , namely set

$$e_k = (0, \dots, 0, 1, 0, \dots, 0), \quad 1 \leq k \leq n. \quad (65)$$

In the symmetric square $S^2(\mathbb{C}^2)$ the basis e and v are as follows:

$$e_{11}^s = e_1 \otimes e_1, \quad e_{12}^s = e_1 \otimes e_2 + e_2 \otimes e_1, \quad e_{22}^s = e_2 \otimes e_2, \quad (66)$$

$$v_{12} = e_1 \otimes e_1, \quad v_{13}^s = (e_1 + e_2) \otimes (e_1 + e_2), \quad v_{23} = e_2 \otimes e_2. \quad (67)$$

The *quantization* of the symmetric square $[S^2(\sigma_k)]_q$ for $k = 1, 2$ of the Burau representation $\rho_3^{(t)}$ have the following form in the basis e :

$$\sigma_1 \mapsto \begin{pmatrix} -t & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \xrightarrow{\text{Sym}^2} \begin{pmatrix} t^2 & 0 & 0 \\ 0 & -t & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & -2 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{\text{Quant}}$$

$$\begin{pmatrix} t^2 & 0 & 0 \\ 0 & -t & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & -(1+q) & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} q & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} t^2q & -t^2(1+q) & t^2 \\ 0 & -t & t \\ 0 & 0 & 1 \end{pmatrix}. \quad (68)$$

We denote

$$D_{1,2}(q) = \text{diag}(q, 1, 1) \quad \text{and} \quad D_{2,2}(q) = \text{diag}(1, 1, q). \quad (69)$$

Similarly, we get

$$\begin{aligned} \sigma_2 &\mapsto \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -t \end{pmatrix} \xrightarrow{\text{Sym}^2} \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -t & 0 \\ 0 & 0 & t^2 \end{pmatrix} \xrightarrow{\text{Quant}} \\ &\begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & (1+q) & q \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & q \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -t & 0 \\ 0 & 0 & t^2 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 1 & -t & 0 \\ 1 & -t(1+q) & t^2q \end{pmatrix}. \end{aligned} \quad (70)$$

Finally, we get

$$[S^2(\sigma_1)_e]_q := \begin{pmatrix} t^2q & -t^2(1+q) & t^2 \\ 0 & -t & t \\ 0 & 0 & 1 \end{pmatrix}, \quad [S^2(\sigma_2)_e]_q := \begin{pmatrix} 1 & 0 & 0 \\ 1 & -t & 0 \\ 1 & -t(1+q) & t^2q \end{pmatrix}. \quad (71)$$

To enumerate the elements of the basis e and v by the same set of indices, set

$$w_{ij} = v_{i,j-1}. \quad (72)$$

Since the basis e and w (hence v) are connected as follows:

$$\begin{aligned} e_{11}^s &= w_{11} & w_{11} &= e_{11}^s \\ e_{12}^s &= -w_{11} + w_{12} - w_{22}, & w_{12} &= e_{11}^s + e_{12}^s + e_{22}^s, \\ e_{22}^s &= w_{22} & w_{22} &= e_{22}^s \end{aligned} \quad (73)$$

the *change-of-basis matrix* C_3 in the space $S^2(\mathbb{C}^2)$ will be the following:

$$C_3 = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{pmatrix}, \quad C_3^{-1} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix}. \quad (74)$$

We show that

$$C_3[S^2(\sigma_r)_e]_q C_3^{-1} = k^{(t,q)}(\sigma_r), \quad \text{for } r = 1, 2. \quad (75)$$

Indeed, we have

$$[S^2(\sigma_1)_v]_q := C_3 \begin{pmatrix} t^2q & -t^2(1+q) & t^2 \\ 0 & -t & t \\ 0 & 0 & 1 \end{pmatrix} C_3^{-1} = \begin{pmatrix} t^2q & 0 & t(t-1) \\ 0 & 0 & t \\ 0 & 1 & 1-t \end{pmatrix}, \quad (76)$$

$$[S^2(\sigma_2)_v]_q := C_3 \begin{pmatrix} 1 & 0 & 0 \\ 1 & -t & 0 \\ 1 & -t(1+q) & t^2q \end{pmatrix} C_3^{-1} = \begin{pmatrix} 0 & t & 0 \\ 1 & 1-t & 0 \\ 0 & qt(t-1) & t^2q \end{pmatrix}. \quad (77)$$

This coincides with the *Krammer representation* $k_3^{(t,q)}$ for B_3 in notations (15):

$$k_3^{(t,q)}(\sigma_1) = \begin{pmatrix} t^2q & 0 & t(t-1) \\ 0 & 0 & t \\ 0 & 1 & 1-t \end{pmatrix}, \quad k_3^{(t,q)}(\sigma_2) = \begin{pmatrix} 0 & t & 0 \\ 1 & 1-t & 0 \\ 0 & qt(t-1) & t^2q \end{pmatrix}. \quad (78)$$

The quantization of the symmetric square $S^2(\rho_{n+1}^{(t)})$ of the Burau representations $\rho_{n+1}^{(t)}$ for the group B_{n+1} will be defined as follows:

$$\sigma_1 \rightarrow [S^2(\sigma_1)]_q = S^2(\exp(sE_{11}))[S^2(\exp(-E_{12}))]_q D_{1,n}(q), \quad (79)$$

$$\sigma_n \rightarrow [S^2(\exp(E_{n-1,n}))]_q D_{n,n}(q) S^2(\exp(sE_{nn})), \quad (80)$$

$$\sigma_k \rightarrow [S^2(\exp(E_{k,k-1}))]_q D_{k,n}(q) S^2(\exp(sE_{kk}))[S^2(\exp(-E_{kk+1}))]_q, \quad (81)$$

We rewrite (84) and (85) in the basis v and compare with the expressions $k_4^{(t,q)}(\sigma_r)$, for $r=1, 2, 3$. As before, we keep the lexicographic order to enumerate the elements of the basis e and w in $S^2(\mathbb{C}^3)$:

$$e = (e_{11}^s, e_{12}^s, e_{22}^s, e_{13}^s, e_{23}^s, e_{33}^s), \quad w = (w_{11}, w_{12}, w_{22}, w_{13}, w_{23}, w_{33}).$$

By Lemma 2.8, the change-of-basis matrices C_4 and C_4^{-1} in the space $S^2(\mathbb{C}^4)$ are the following:

$$C_4 = \begin{pmatrix} 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 0 \\ 0 & -1 & 1 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 \end{pmatrix}, \quad C_4^{-1} = \begin{pmatrix} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 \end{pmatrix}. \quad (86)$$

We calculate $[S^2(\sigma_r)_v]_q = C_4[S^2(\sigma_r)_e]_q C_4^{-1}$ for $r = 1, 2, 3$. We have

$$[S^2(\sigma_1)_v]_q = C_4 \begin{pmatrix} t^2 q & -t^2(1+q) & t^2 & 0 & 0 & 0 \\ 0 & -t & t & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -t & t & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} C_4^{-1} = \begin{pmatrix} qt^2 & 0 & t(t-1) & 0 & t(t-1) & 0 \\ 0 & 0 & t & 0 & 0 & 0 \\ 0 & 1 & 1-t & 0 & 0 & 0 \\ 0 & 0 & 0 & t & 0 & 0 \\ 0 & 0 & 0 & 1 & 1-t & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad (87)$$

$$[S^2(\sigma_2)_v]_q = C_4 \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & -t & 0 & t & 0 & 0 \\ 1 & -t(1+q) & t^2 q & t(1+q) & -t^2(1+q) & t^2 \\ 0 & 0 & 0 & 1 & -t & t \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} C_4^{-1} = \begin{pmatrix} 0 & t & 0 & 0 & 0 & 0 \\ 1 & 1-t & 0 & 0 & 0 & 0 \\ 0 & qt(t-1) & qt^2 & 0 & 0 & t(t-1) \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & t \\ 0 & 0 & 0 & 0 & 0 & 1-t \end{pmatrix},$$

$$[S^2(\sigma_3)_v]_q = C_4 \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -t & 0 & 0 \\ 0 & 0 & 1 & 0 & -t & 0 \\ 0 & 0 & 1 & 0 & -t(1+q) & t^2 q \end{pmatrix} C_4^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & t & 0 & 0 \\ 0 & 0 & 0 & 0 & t & 0 \\ 0 & 1 & 0 & 1-t & 0 & 0 \\ 0 & 0 & 1 & 0 & 1-t & 0 \\ 0 & 0 & 0 & qt(t-1) & qt(t-1) & qt^2 \end{pmatrix}. \quad (88)$$

This coincide with the Krammer representation $k_4^{(t,q)}(\sigma_r)$ for B_4 in notations (15):

$$k_4^{(t,q)}(\sigma_1) = \begin{pmatrix} qt^2 & 0 & t(t-1) & 0 & t(t-1) & 0 \\ 0 & 0 & t & 0 & 0 & 0 \\ 0 & 1 & 1-t & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & t & 0 \\ 0 & 0 & 0 & 1 & 1-t & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad k_4^{(t,q)}(\sigma_2) = \begin{pmatrix} 0 & t & 0 & 0 & 0 & 0 \\ 1 & 1-t & 0 & 0 & 0 & 0 \\ 0 & qt(1-t) & qt^2 & 0 & 0 & t(t-1) \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1-t \end{pmatrix},$$

$$k_4^{(t,q)}(\sigma_3) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & t & 0 & 0 \\ 0 & 0 & 0 & 0 & t & 0 \\ 0 & 1 & 0 & 1-t & 0 & 0 \\ 0 & 0 & 1 & 0 & 1-t & 0 \\ 0 & 0 & 0 & qt(t-1) & qt(t-1) & qt^2 \end{pmatrix}.$$

Remark 8. Rinat Kashaev pointed out to the author during his visit to Geneva in May 2022, that the formulas (87)–(88) are not correct.

Lemma 2.8. The matrices C_{n+1}^{-1} and C_{n+1} are defined as follows. The columns of the matrix C_{n+1}^{-1} are defined by the following relations:

$$w_{ij} = \sum_{i \leq k \leq r \leq j} e_{kr}^s, \quad 1 \leq i \leq j \leq n. \quad (89)$$

The columns of the matrix C_{n+1} are defined by the following relations:

$$e_{ij}^s = \begin{cases} (-1)^{i+j} \sum_{i \leq k \leq r \leq j} (-1)^{k+r} w_{kr}, & \text{if } 0 \leq j-i \leq 1, \\ \sum_{r=i+1}^{j-1} (-w_{ir} + w_{i+1,r}) + w_{ij} - w_{i+1,j}, & \text{if } i=1, \quad 2 \leq j-i, \\ -w_{ii} + \sum_{r=i+1}^{j-1} (-w_{ir} + w_{i+1,r}) + w_{ij} - w_{i+1,j}, & \text{if } i>1, \quad 2 \leq j-i. \end{cases} \quad (90)$$

Indeed, to prove (74), (86) and (94) we note that matrix $E_n = C_n^{-1}$ is block upper-triangular

$$E_n = \begin{pmatrix} a_n & b_n \\ 0 & c_n \end{pmatrix}, \quad \text{therefore} \quad C_n = E_n^{-1} = \begin{pmatrix} a_n^{-1} & -a_n^{-1}b_n c_n^{-1} \\ 0 & c_n^{-1} \end{pmatrix}. \quad (91)$$

For $n = 3$ we have

$$E_3 = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} a_3 & b_3 \\ 0 & c_3 \end{pmatrix}, \quad E_3^{-1} = \begin{pmatrix} a_3^{-1} & -a_3^{-1}b_3 c_3^{-1} \\ 0 & c_3^{-1} \end{pmatrix}$$

where

$$a_3 = (1), \quad b_3 = (1 \ 0), \quad c_3 = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}.$$

We have $c_3^{-1} = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}$, therefore $a_3^{-1}b_3 c_3^{-1} = (1)(1 \ 0) \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} = (1 \ 0)$, hence $C_3 = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{pmatrix}$. For $n = 4$ we get

$$E_4 = \begin{pmatrix} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 \end{pmatrix} = \begin{pmatrix} a_4 & b_4 \\ 0 & c_4 \end{pmatrix}, \quad E_4^{-1} = \begin{pmatrix} a_4^{-1} & -a_4^{-1}b_4 c_4^{-1} \\ 0 & c_4^{-1} \end{pmatrix}$$

where

$$a_4 = E_3, \quad b_4 = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 1 & 0 \end{pmatrix}, \quad c_4 = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}.$$

Since $c_4^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{pmatrix}$ we get

$$a_4^{-1}b_4 c_4^{-1} = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ -1 & 1 & 0 \end{pmatrix}, \quad C_4 = \begin{pmatrix} 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 0 \\ 0 & -1 & 1 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 \end{pmatrix}.$$

For $n = 5$ we have

$$E_5 = C_5^{-1} = \begin{pmatrix} 1 & 1 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \end{pmatrix} = \begin{pmatrix} a_5 & b_5 \\ 0 & c_5 \end{pmatrix}, \quad E_5^{-1} = \begin{pmatrix} a_5^{-1} & -a_5^{-1}b_5 c_5^{-1} \\ 0 & c_5^{-1} \end{pmatrix},$$

where

$$a_5 = E_4, \quad b_5 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \end{pmatrix}, \quad c_5 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{pmatrix}.$$

Since $c_5^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & -1 & 1 \end{pmatrix}$ we get

$$a_5^{-1}b_5 c_5^{-1} = \begin{pmatrix} 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 0 \\ 0 & -1 & 1 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & -1 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \end{pmatrix},$$

and

$$C_5 = \begin{pmatrix} 1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & -1 & 1 & 1 & -1 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 \end{pmatrix}. \quad \square$$

Remark 9. The matrix $E_n = C_n^{-1}$ and $E_n^{-1} = C_n$ have the following form

$$E_n = \begin{pmatrix} e_{11} & e_{12} & e_{13} & \cdots & e_{1n-1} \\ 0 & e_{22} & e_{23} & \cdots & e_{2n-1} \\ 0 & 0 & e_{33} & \cdots & e_{3n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & e_{n-1,n-1} \end{pmatrix}, \quad E_n^{-1} = \begin{pmatrix} e_{11}^{-1} & e_{12}^{-1} & e_{13}^{-1} & \cdots & e_{1n-1}^{-1} \\ 0 & e_{22}^{-1} & e_{23}^{-1} & \cdots & e_{2n-1}^{-1} \\ 0 & 0 & e_{33}^{-1} & \cdots & e_{3n-1}^{-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & e_{n-1,n-1}^{-1} \end{pmatrix}, \quad (92)$$

where $e_{kr} \in \text{Mat}(k \times r, \mathbb{C})$ for $1 \leq k \leq r \leq n-1$ are as follows:

$$e_{11} = (1), \quad e_{22} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \quad e_{33} = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}, \quad e_{kk} = (I - e_k)^{-1}, \quad e_k = \sum_{r=1}^{k-1} E_{r+1,r},$$

$e_{kr} = (e_{kk}, 0_{k,r-k})$, where $0_{k,r} = 0$ in $\text{Mat}(k \times r, \mathbb{C})$. For E_n^{-1} we have

$$e_{11}^{-1} = (1), \quad e_{22}^{-1} = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}, \quad e_{33}^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{pmatrix}, \quad e_{kk} = (I - e_k),$$

and $e_{1r}^{-1} = 0$ for $2 \leq r \leq n-1$, $e_{kr}^{-1} := (-e_{kk}^{-1}, 0_{k,r-k})$ for $2 \leq k < r \leq n-1$. Notation e_{kr}^{-1} for $k < r$ does not mean the inverse matrix to e_{kr} !

To prove (63) for general B_{n+1} , it is sufficient to consider only three cases $n = 3, 4$ and $n = 5$ for σ_3 . Indeed, we can use the *invariance* of the reduced Burau representation of B_∞ and B_n , (see (9), (10) and (13)) and invariance of the Lawrence-Krammer representation (15).

The Burau representation $\rho_5^{(t)}$ for B_5 is as follows:

$$\sigma_1 \mapsto \begin{pmatrix} -t & t & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \sigma_2 \mapsto \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & -t & t & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \sigma_3 \mapsto \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & -t & t \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \sigma_4 \mapsto \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & -t \end{pmatrix}. \quad (93)$$

By Lemma 2.8, the change-of-basis matrix C_5 and C_5^{-1} in the space $S^2(\mathbb{C}^5)$ are the following:

$$C_5 = \begin{pmatrix} 1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & -1 & 1 & 1 & -1 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 \end{pmatrix}, \quad C_5^{-1} = \begin{pmatrix} 1 & 1 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \end{pmatrix}. \quad (94)$$

Using (15) we get

$$k_5^{(t,q)}(\sigma_3) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & t & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1-t & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1-t & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & qt(t-1) & 0 & 0 & qt(t-1) & 0 & qt^2 & 0 & t(t-1) \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & t \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1-t \end{pmatrix}.$$

Indeed, since

$$\sigma_3 \mapsto \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & -t & t \\ 0 & 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -t & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{pmatrix} = \\ \exp(E_{32}) \exp(sE_{33}) \exp(-E_{34}),$$

$$S^2(e^{E_{32}}) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad S^2(e^{-E_{34}}) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$
[illegible]
$$C_5[S^2(\rho_5^{(t)}(\sigma_3)_e)_q]C_5^{-1} =$$

$$\begin{pmatrix} 1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 0 & -1 & 0 & 0 \\ 0 & -1 & 1 & 1 & -1 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & t & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -t & 0 & 0 & t & 0 & 0 \\ 0 & 0 & 1 & 0 & -t & 0 & 0 & t & 0 \\ 0 & 0 & 1 & 0 & -t(1+q) & qt^2 & 0 & t(1+q) & -t^2(1+q) \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \times$$

$$\begin{pmatrix} 1 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & t & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1-t & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -t & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -t & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & qt(t-1) & 0 & 0 & t(t-1) & 0 & qt^2 & 0 & t(t-1) & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & t & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1-t & 0 \end{pmatrix} \stackrel{(15)}{=} k_5^{(t,q)}(\sigma_3).$$

3. How to construct new representations of the braid group B_n

We start with the well-known representation (in fact, reduced Burau representation for $t = -1$) $T_{32} := \rho_3^{(-1)} : B_3 \rightarrow \mathrm{SL}(2, \mathbb{Z})$ defined by (64)

16

This representation is just exp of the fundamental (or natural) representations π_2 of the Lie algebra \mathfrak{sl}_2 (see (18) and (19)).

Remark 10. We have several possibility to generalize representation T_{32} of the group B_3 . First, we can generalize T_{32} for an arbitrary B_n (denote by T_{n2}), second, we can generalize T_{32} of the same group B_3 but for an arbitrary dimension m (denote by T_{3m}), and finally, we can generalize representations T_{n2} of B_n for higher dimension m , (denote by T'_{nm}) and extend representation T_{3m} of B_3 for other groups B_n (denote by T''_{nm}). Finally, we have two family of representations

$$T'_{nm}, T''_{nm} : B_n \rightarrow \mathrm{GL}(m, \mathbb{C}).$$

To be more precise, the *first step* gives us all Burau's representations $\rho_{n+1}^{(t)} : B_{n+1} \rightarrow \mathrm{GL}_n(\mathbb{Z}[t, t^{-1}])$, due to Theorem 2.1, if we start with $\rho_3^{(t)}$ and take for π the natural representation π_n of \mathfrak{sl}_n .

On the *second step*, we obtain the Humphry representation of B_3 in all dimension $m + 1$ as the m^{th} symmetric power of the Burau representation ρ_3^{-1} , due to Theorem 2.5. Remarkably, that *quantization* and *diagonal deformation* (governed by the matrix $\Lambda_n = \mathrm{diag}(\lambda_r)_{r=0}^n$, see Remark 2) of the Humphry representations (see (35)) include all irreducible representations of B_3 in dimension ≤ 5 , due to results of Tuba and Wensel [28, 29] and our equivalent description of their representations [1, 17], see Theorem 2.7. Our approach using q -Pascal triangle allows us, for free, extend representation T_{32} of B_3 for arbitrary dimension (see Theorem 2.6).

On the *third step*, we obtain first, the Lawrence-Krammer representation as quantization of the symmetric square of the Burau representation, Theorem 2.2. Due to formulas (54), (55) and Theorem 2.2, the Lawrence-Krammer representations of B_3 describe all irreducible representations, up to the scalar factor, in dimensions 3.

Remark 11. It would be nice to give explicit formulas for quantization of the symmetric cube and higher symmetric power of the Burau representation.

$V = \mathbb{C}^n$	B_3	B_n	type of representation
2	$T_{3,2}^{\Lambda_{1,1}, \pi_2}$	$\rho_n^{(t)}$	reduced Burau representation
3	$T_{3,3}^{\Lambda_{2,q}, \pi_2}$	$k_n^{(t,q)}$	Lawrence–Krammer representation
4	$T_{3,4}^{\Lambda_{3,q}, \pi_2}$	$[S^3(\rho_n^{(t)})]_q$	Symmertic cube
5	$T_{3,5}^{\Lambda_{4,q}, \pi_2}$	$[S^4(\rho_n^{(t)})]_q$	4^{th} Symmertic power
$m + 1$	$T_{3,m+1}^{\Lambda_{m,q}, \pi_2}$	$[S^m(\rho_n^{(t)})]_q$	m^{th} Symmertic power

3.2. Representations of a Lie algebra \mathfrak{sl}_{n+1}

We recall, following to J.P.Serre [25], Part I, Ch. VII, §1; Part III, Ch. VII, §6, that all irreducible finite-dimensional representations of a *simple* Lie algebra \mathfrak{sl}_{n+1} are *highest weight representations* and are contained in tensor powers of the *natural representation* π_{n+1} [31].

The *Cartan subalgebra* \mathfrak{h} of \mathfrak{sl}_{n+1} consists of all matrices of the form $H = \mathrm{diag}(\lambda_1, \dots, \lambda_{n+1})$ with $\sum_k \lambda_k = 0$. The *roots* of a Lie algebra \mathfrak{sl}_{n+1} are linear functionals $\alpha_{i,j} \in \mathfrak{h}^*$, $i \neq j$, of the form

$$\alpha_{i,j}(H) = \lambda_i - \lambda_j, \quad \text{where } H = \mathrm{diag}(\lambda_1, \dots, \lambda_{n+1}).$$

The *fundamental wights* ω_k are defined by

$$\omega_k(H) = \lambda_1 + \cdots + \lambda_k, \quad \text{where } H = \text{diag}(\lambda_1, \dots, \lambda_{n+1}).$$

The fundamental weight ω_1 is the fundamental weight of the *standard representation* π_{n+1} of the Lie algebra \mathfrak{sl}_{n+1} in $E = \mathbb{C}^{n+1}$. Moreover, the fundamental weight ω_k is the highest weight of the representation $\wedge^k(\pi_{n+1})$ of \mathfrak{sl}_{n+1} in the space $\wedge^k E$ (see [25, 1.7.4.6]), where $\wedge^k(A)$ is k^{th} *exteriour power* of A .

In particular case, all irreducible representations of \mathfrak{sl}_2 are $\text{Sym}^n(\pi_2)$ in $\text{Sym}^n(\mathbb{C}^2)$, $n \in \mathbb{N}$. For \mathfrak{sl}_3 all irreducible representations are contained in tensor product of π_3 and $\wedge^2(\pi_3)$, corresponding to fundamental weights ω_1 and ω_2 . All irreducible representations of \mathfrak{sl}_{n+1} are contained in tensor products of $\wedge^k(\pi_{n+1})$, $1 \leq k \leq n$.

Let $\pi_n : \mathfrak{sl}_n \rightarrow \text{End}(\mathbb{C}^m)$ be any finite-dimensional representation of a Lie algebra \mathfrak{sl}_n then $\exp(\pi_n) : \text{SL}(n, \mathbb{C}) \rightarrow \text{GL}(m, \mathbb{C})$ is representation of $\text{SL}(n, \mathbb{C})$. We can define representations of B_{n+1} as follows

$$\sigma_k \rightarrow S^m(\exp(\pi_n) \circ \rho_n^{(t)}(\sigma_k)), \quad \sigma_k \rightarrow \wedge^m(\exp(\pi_n) \circ \rho_n^{(t)}(\sigma_k)). \quad (96)$$

Consider the exterior square $\wedge^2(\rho_{2,4}^{(t)})$ of the Burau representation $\rho_{2,4}^{(t)}$ of the group B_4 defined by (4) and (5) in the basis e_{ij}^\wedge , $1 \leq i < j \leq 3$ of the space $\wedge^2(\mathbb{C}^3)$

$$e_{12}^\wedge = e_1 \otimes e_2 - e_2 \otimes e_1, \quad e_{13}^\wedge = e_1 \otimes e_3 - e_3 \otimes e_1, \quad e_{23}^\wedge = e_2 \otimes e_3 - e_3 \otimes e_2.$$

Lemma 3.1. *Representation $\wedge^2(\rho_{2,4}^{(t)})$ of the group B_4 is defined by*

$$\sigma_1 \rightarrow \begin{pmatrix} -t & 0 & 0 \\ 0 & -t & t \\ 0 & 0 & 1 \end{pmatrix}, \quad \sigma_2 \rightarrow \begin{pmatrix} -t & t & 0 \\ 0 & 1 & 0 \\ 0 & 1 & -t \end{pmatrix}, \quad \sigma_3 \rightarrow \begin{pmatrix} 1 & 0 & 0 \\ 0 & -t & 0 \\ 0 & 0 & -t \end{pmatrix}. \quad (97)$$

We have

$$\wedge^2 \rho_{2,4}^{(t)} \not\sim \rho_4^{(t)}, \quad \text{for } t \neq -1, \quad \text{but } \wedge^2 \rho_{2,4}^{(t)} = -t S_3 \rho_{1,4}^{(-t^{-1})} S_3 \quad (98)$$

where $\rho_{1,4}^{(t)}$ is defined by (2) and (3) and $S_3 = \sum_{k=1}^2 E_{k,4-k} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$.

The first part of (98) follows from comparison of two spectra: $\text{Sp} \rho_{24}^{(t)}(\sigma_1) = \{-t, 1, 1\}$ and $\text{Sp}(\wedge^2 \rho_{24}^{(t)}(\sigma_1)) = \{-t, 1, -t\} = -t\{1, -t^{-1}, 1\}$. The second part follows from (2) and (3).

Problem 3.2. *To quantize the representation of B_n defined by (96).*

$$\sigma_k \rightarrow [S^m(\exp(\pi_n) \circ \rho_n^{(t)}(\sigma_k))]_q, \quad \sigma_k \rightarrow [\wedge^m(\exp(\pi_n) \circ \rho_n^{(t)}(\sigma_k))]_q. \quad (99)$$

4. Some polynomial invariants for knots

4.1. *Some polynomial invariants for the trefoil, see [4]*

The Alexander polynomial $\Delta(x)$, BLM/Ho polynomial $Q(x)$, Conway polynomial $\nabla(x)$, HOMFLY polynomial $P(l, m)$, Jones polynomial $V(t)$, and Kauffman polynomial $F(a, z)$ of the trefoil knot are

$$\Delta(x) = x - 1 + x^{-1}, \quad (100)$$

$$Q(x) = 2x^2 + 2x - 3, \quad (101)$$

$$\nabla(x) = x^2 + 1, \quad (102)$$

$$P(l, m) = -l^4 + m^2 l^2 - 2l^2, \quad (103)$$

$$V(t)_{3_1} = t + t^3 - t^4, \quad V(t)_{3_2} = t^{-1} + t^{-3} - t^{-4}, \quad (104)$$

$$F(a, z) = -a^4 - 2a^2 + (a^4 + a^2)z^2 + (a^5 + a^3)z. \quad (105)$$

4.2. The Burau representation and the Alexander polynomial

Let K be some link, by *Alexander's theorem* this link can be obtained as the closure of some braid $X \in B_n$, notation $K = Cl(X)$. The *Alexander polynomial* $\Delta_K(t)$ can be obtained using the reduced Burau representation by the following formula, [10, (4.3), p.64]:

$$\Delta_{Cl(X)}(t) = \frac{(1-t)\det(\rho_n^{(t)}(X) - I_{n-1})}{1-t^n} = \frac{\det(\rho_n^{(t)}(X) - I_{n-1})}{\det(\rho_n^{(t)}(\sigma_1 \dots \sigma_{n-1}) - I_{n-1})}. \quad (106)$$

4.3. The Krammer representation and the corresponding rational function

Definition 1. By analogy with the definition of the Alexander polynomial we can introduce the following rational function $k_n(t, q)$ connected with the Lawrence-Krammer representation:

$$k_{n, Cl(X)}(t, q) = \frac{\det(k_n^{(t, q)}(X) - I_{n(n-1)/2})}{\det(k_n^{(t, q)}(\sigma_1 \dots \sigma_{n-1}) - I_{n(n-1)/2})}. \quad (107)$$

Remark 12. Recall that some polynomial $P(x, y, \dots)$ defined on the set of all knots is knot invariant if and only if it respect the the first and the second Markov moves (see, e.g., [16, 2.5. Markov's theorem, p.67]), i.e., for any $X \in B_n$ the following properties hold:

$$P_{Cl(wXw^{-1})}(x, y, \dots) = P_{Cl(X)}(x, y, \dots) \quad \text{for all } w \in B_n, \quad (108)$$

$$P_{Cl(X\sigma_n^{\pm 1})}(x, y, \dots) = P_{Cl(X)}(x, y, \dots). \quad (109)$$

Theorem 4.1. The rational function $k_{n, Cl(X)}(t, q)$ defined by (107) respect the first Markov's move, i.e., depends only on conjugacy classes in B_n .

We show that

$$k_{Cl(wXw^{-1})}(t, q) = k_{Cl(X)}(t, q) \quad \text{for all } w \in B_n, \quad (110)$$

Since $\det(AB) = \det(A)\det(B)$ we get by definition

$$\begin{aligned} \det(k_n^{(t, q)}(wXw^{-1}) - I_{n(n-1)/2}) &= \det(k_n^{(t, q)}(wXw^{-1}) - k_n^{(t, q)}(ww^{-1})) \\ &= \det\left(k_n^{(t, q)}(w)(k_n^{(t, q)}(X) - I_{n(n-1)/2})k_n^{(t, q)}(w)^{-1}\right) \\ &= \det(k_n^{(t, q)}(X) - I_{n(n-1)/2}), \end{aligned}$$

that proves (110).

Problem 4.2. Study the behaviour of the rational function $k_n(t, q)$ with respect to the second Markov's move.

4.4. Alexander polynomial for the trefoil

To calculate the Alexander polynomial for the trefoil knot $3_1 = Cl(\sigma_1^3) = Cl(\sigma_1^3\sigma_2)$ we get for the representation $\rho_2^{(t)}(\sigma_1) = -t$ of the group B_2

$$\Delta_{3_1}(t) = \frac{\det(\rho_2^{(t)}(\sigma_1^3) - I_1)}{\det(\rho_2^{(t)}(\sigma_1) - I_1)} = \frac{-t^3 - 1}{-(t + 1)} = t^2 - t + 1. \quad (111)$$

The second Markov move (see (109)). For the Burau representation $\rho_3^{(t)}$ of the group B_3

$$\rho_3^{(t)}(\sigma_1) = \begin{pmatrix} -t & t \\ 0 & 1 \end{pmatrix}, \quad \rho_3^{(t)}(\sigma_2) = \begin{pmatrix} 1 & 0 \\ 1 & -t \end{pmatrix}$$

we have

$$\rho_3^{(t)}(\sigma_1\sigma_2^3) = \begin{pmatrix} -t & t \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & -t \end{pmatrix}^3 = \begin{pmatrix} t^3 - t^2 & -t^4 \\ t^2 - t + 1 & -t^3 \end{pmatrix}.$$

Hence,

$$\begin{aligned} \det(\rho_3^{(t)}(\sigma_1\sigma_2^3) - I_2) &= \det \begin{pmatrix} t^3 - t^2 - 1 & -t^4 \\ t^2 - t + 1 & -t^3 - 1 \end{pmatrix} = \\ &= -(t^3 + 1)(t^3 - t^2 - 1) + t^4(t^2 - t + 1) = t^4 + t^2 + 1, \end{aligned}$$

and we obtain the same result as in (111):

$$\Delta_{3_1}(t) = \frac{\det(\rho_3^{(t)}(\sigma_1^3\sigma_2) - I_2)}{\det(\rho_3^{(t)}(\sigma_1\sigma_2) - I_2)} = \frac{t^4 + t^2 + 1}{t^2 + t + 1} = t^2 - t + 1. \quad (112)$$

4.5. Krammer representation and the trefoil

The Krammer representation for the group B_2 is as follows (see (15)) $\sigma_1 \rightarrow t^2q$, therefore, using (107) we get for the trefoil knot $3_1 = Cl(\sigma_1^3)$ (see (111))

$$k_{2,3_1}(t, q) = \Delta_{3_1}(t^2q) = (t^2q)^2 - t^2q + 1 = t^4q^2 - t^2q + 1. \quad (113)$$

Remark 13. We have

$$k_{2,3_1}(1, q) = q^2 - q + 1 = \Delta_{3_1}(q), \quad k_{2,3_1}(t, 1) = t^4 - t^2 + 1 = \Delta_{3_1}(t^2).$$

Calculate now $k_{3,3_1}(t, q)$ for the trefoil knot $3_1 = Cl(\sigma_1^3\sigma_2)$ using formulas (71), (76), (77) and (78).

Lemma 4.3. We have

$$\det(k_3^{(t,q)}(\sigma_1\sigma_2) - I_3) = t^6q^2 - 1, \quad (114)$$

$$\det(k_3^{(t,q)}(\sigma_1^3\sigma_2) - I_3) = t^{12}q^4 - t^6q(1 - q)(1 - t)(1 - tq) - 1, \quad (115)$$

$$k_{3,3_1}(t, q) = \frac{t^{12}q^4 - t^6q(1 - q)(1 - t)(1 - tq) - 1}{t^6q^2 - 1}. \quad (116)$$

Remark 14. The rational function $k_n(t, q)$ does not respect the second Markov's move. Indeed, in the particular case of $q = 1$ we get

$$k_{3,3_1}(t, 1) = \frac{t^{12} - 1}{t^6 - 1} = t^6 + 1 = (t^2 + 1)(t^4 - t^2 + 1) = (t^2 + 1)k_{2,3_1}(t, 1). \quad (117)$$

Remark 15. We have for $q = 1$

$$k_{3,3_1}(t, 1) = (t^2 + 1)k_{2,3_1}(t, 1) = (t^2 + 1)\Delta(t^2), \quad \text{the Alexander polynomial.} \quad (118)$$

In the particular case of $t = 1$ we get

$$k_{3,3_1}(1, q) = \frac{q^4 - 1}{q^2 - 1} = q^2 + 1 = \nabla(q), \quad \text{the Conway polynomial.} \quad (119)$$

To prove (114) and (115), using (71) we have

$$\sigma_1 \mapsto \begin{pmatrix} t^2q & -t^2(1+q) & t^2 \\ 0 & -t & t \\ 0 & 0 & 1 \end{pmatrix}, \quad \sigma_2 \mapsto \begin{pmatrix} 1 & 0 & 0 \\ 1 & -t & 0 \\ 1 & -t(1+q) & t^2q \end{pmatrix}, \quad (120)$$

therefore

$$\sigma_1\sigma_2 \mapsto \begin{pmatrix} 0 & 0 & t^4q \\ 0 & -tq^2 & t^3q \\ 1 & -t(1+q) & t^2q \end{pmatrix} := B(t, q).$$

To calculate $\det(B(t, q) - I_3)$ we use the following lemma.

Lemma 4.4 (Lemma 1.4.5 in [18]). For the generalized characteristic polynomial $P_C^g(\lambda)$ of a matrix $C \in \text{Mat}(m, \mathbb{C})$ defined by (121) and $\lambda \in \mathbb{C}^m$ we have

$$P_C^g(\lambda) = \det \left(C + \sum_{k=1}^m \lambda_k E_{kk} \right) = \quad (121)$$

$$\det C + \sum_{r=1}^m \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq m} \lambda_{i_1} \lambda_{i_2} \dots \lambda_{i_r} A_{i_1 i_2 \dots i_r}^{i_1 i_2 \dots i_r}(C), \quad (122)$$

where $A_{i_1 i_2 \dots i_r}^{i_1 i_2 \dots i_r}(C)$ are cofactors of C .

By Lemma 4.4 we have for $A = B(t, q)$ and $\lambda_k = -1$, $k = 1, 2, 3$

$$\begin{aligned} \det(B(t, q) - 1) &= \det A(t) - (A_1^1 + A_2^2 + A_3^3) + (A_{12}^{12} + A_{13}^{13} + A_{23}^{23}) - A_{123}^{123} = \\ &= t^6 q^2 - (t^4 q - t^4 q) + (t^2 q - t^2 q) - 1 = t^6 q^2 - 1. \end{aligned}$$

To prove (115) we have

$$\begin{aligned} \sigma_1^3 \sigma_2 &= \sigma_1^2 \sigma_1 \sigma_2 \mapsto \begin{pmatrix} t^4 q^2 & t^3(1+q)(1-tq) & t^2[t^2 q - t(1+q) + 1] \\ 0 & t^2 & -t(t-1) \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 & t^4 q \\ 0 & -tq^2 & t^3 q \\ 1 & -t(1+q) & t^2 q \end{pmatrix} = \\ &= \begin{pmatrix} t^2[t^2 q - t(1+q) + 1] & (1+q)t^3(tq-1)(t^2 q - t + 1) & t^4 q[t^4 q^2 + t^2(1+q)(1-tq) + t^2 q - t(1+q) + 1] \\ -t(t-1) & -t^4 q + t^2(1+q)(t-1) & t^3 q(t^2 - t + 1) \\ 1 & -t(1+q) & t^2 q \end{pmatrix} = A(t, q). \end{aligned} \quad (123)$$

For $A = A(t, q)$ and $\lambda_k = -1$, $k = 1, 2, 3$ we get

$$\begin{aligned} \det(A(t, q) - 1) &= \det A(t, q) - (A_1^1 + A_2^2 + A_3^3) + (A_{12}^{12} + A_{13}^{13} + A_{23}^{23}) - A_{123}^{123} \\ &= (t^3 q)^4 - t^6 q(1 - [t^2 q^2 - q(1+q)t + (1+q)] + [t^2 q - (1+q)t + 1]) \\ &\quad + t^2([t^2 q - (1+q)t + 1] - [t^2 q - (1+q)t + (1+q)] + q) - 1 = \\ &= t^{12} q^4 - t^6 q(1 - q)[t^2 q - (1+q)t + 1] - 1 = t^{12} q^4 - t^6 q(1 - q)(1 - t)(1 - tq) - 1. \end{aligned}$$

Acknowledgements 2017. The author expresses his deep gratitude to the Max–Planck–Institute of Mathematics for the financial grant and the hospitality in 2016–2017. Sergei Chmutov have read the first version of this article and made a lot of useful remarks, that improved considerable the content. I am very grateful to him for his valuable help.

Acknowledgements 2023. The author is very grateful to Prof. K.-H. Neeb, Prof. M. Smirnov and Dr P. Moree for their personal efforts to make academic stays possible at their respective institutes. The author visited: MPIM from March to April 2022 and from January to April 2023, University of Augsburg from June to July 2022, and University of Erlangen-Nuremberg from August to December 2022, all during the Russian invasion in Ukraine. Also, Prof. R. Kashaev kindly invited him to Geneva.

Further, he would like to pay his respect to Prof. P. Teichner at MPIM, for his immediate efforts started to help mathematicians in Ukraine after the Russian invasion.

Since the spring of 2023 A. Kosyak is an Arnold Fellow at the London Institute for Mathematical Sciences, and he would like to express his gratitude to Mrs S. Myers Cornaby to Miss A. Ker Mercer and to Dr M. Hall and especially to the Director of LIMS Dr T. Fink and Prof. Y.-H. He.

References

- [1] S. Albeverio and A. Kosyak, q -Pascal’s triangle and irreducible representations of the braid group B_3 in arbitrary dimension, arXiv:math.QA(RT)/0803.2778v2.
- [2] G.E. Andrews, The Theory of Partitions. Encyclopedia of Mathematics and its Applications, Addison–Wesley Publishing Company, Cambridge, Massachusetts, 1976.
- [3] E. Artin, Theorie des Zöpfe, Abh. Math. Sem. Hamburg. Univ 4 (1926) 47–72.
- [4] D. Bar-Natan and S. Morrison, and et al., The *Knot Atlas*, <http://katlas.org>
- [5] S. Bigelow, The Burrau representation of the braid group B_n is not faithful for $n=5$, Geometry and Topology 3 (1999), 397–404.
- [6] S. Bigelow, Braid groups are linear, J. Amer. Math. Soc. 14 No 2, (2001), 471–486.
- [7] S. Bigelow, The Lawrence-Krammer representation. arXiv:math.GT/0204057.
- [8] S. Bigelow, Does the Jones polynomial detect the unknot?, Journal of Knot Theory and Its Ramifications. **11** (4) (2002) 493–505.
- [9] J.S Birman, Braids, links and mapping class groups, Ann. Math. Studies, 82, Princeton Univ. Press, 1974.
- [10] J.S Birman and T.E Brendel, Braids: A survey. In Handbook of Knot Theory (Ed. W. Menasco and T. Thistlethwaite), Elsevier, 2005.

- [11] W. Burau, Über Zopfgruppen und gleichsinnig verdrillte Verkettungen. Abh. Math. Sem. Univ. Hamburg. **11** (1936) 179–186.
- [12] S.P. Humphries, Some linear representations of braid groups, Journ. Knot Theory and Its Ramifications. 9, no. 3 (2000) 341–366.
- [13] C. Jackson and T. Kerler, The Lawrence–Krammer–Bigelow representations of the braid groups via $U_q(\mathfrak{sl}_2)$, Adv. Math. 228 (2011) 1689–1717.
- [14] V. F. R. Jones, Hecke algebra representations of braid groups and link polynomials, Annals of Math. 126 (1987) 335–388.
- [15] C. Kassel, Quantum Groups, Springer-Verlag, 1995.
- [16] C. Kassel, V. Turaev, Braid groups, Graduate Texts in Mathematics, vol. 247, Springer, New York, 2008 (340 p. + xii).
- [17] A.V. Kosyak, Representations of the braid group B_n and the highest weight modules of $U(\mathfrak{sl}_{n-1})$ and $U_q(\mathfrak{sl}_{n-1})$, arXiv:math.QA(RT)/0803.2785v2.
- [18] A.V. Kosyak, Regular, quasiregular and induced representations of infinite-dimensional groups, EMS (Europ. Math. Soc), 536 p., to appear in Europ. Math. Soc. 2017, (submitted for publication).
- [19] D. Krammer, The braid group B_4 is linear, Invent. Math. 142 (2000), no. 3, 451–486.
- [20] D. Krammer, Braid groups are linear, Ann. of Math. (2) 155 No. 1, (2002), 131–156.
- [21] R. Lawrence, Homological Representations of the Hecke Algebra, Commun. Math. Phys. 135 (1990) 141–191.
- [22] D. Long and M. Paton, The Burrau representation of the braid group B_n is not faithful for $n \geq 6$, Topology 32 (1993), 439–447.
- [23] W. Magnus and A. Peluso, On a theorem of V.I. Arnold, Comm Pure Appl. Math, 22 (1969) 683–692.
- [24] J. Moody, The Burrau representation of the braid group B_n is not faithful for large n , Bull. Math. Soc. 25 (1991), 379–384.
- [25] J.-P. Serre, Lie Algebras and Lie Groups, 1964 Lectures Given at Harvard University, Part of Lect. Notes Math. 1500.
- [26] R. Smeltzer, Linear representations of braid groups. A thesis for the degree of the master of sciences, McMaster University, July 2003. 65 p.
- [27] A. Stoimenov, The density of Lawrence–Krammer and non-conjugate braid representation of links, arXiv:math.GR/ 0809.0033 v2.
- [28] I. Tuba, Low-dimensional unitary representations of B_3 , Proc. Amer. Math. Soc. 129 (2001) 2597–2606.

- [29] I. Tuba, H. Wenzl, Representations of the braid group B_3 and of $SL(2, \mathbb{Z})$, Pacific J. Math. 197, No.2 (2001) 491–510.
- [30] H. Wenzl, Quantum groups and subfactors of type B , C , and D , Comm. Math. Phys. 133 (1990) 383–432.
- [31] H. Weyl, The Classical Groups, Their Invariants and Representations, (First Edition, 1939, Second Edition with supplements, 1949) Princeton University Press.
- [32] B. Westbury, On the character varieties of the modular group, preprint Nottingham, 1995.
- [33] M. Zinno, On Krammer’s representation of the braid group, Math. Ann. 321 (2001) 197–211.