A Wald-type test statistic for testing linear hypothesis in logistic regression models based on minimum density power divergence estimator

Basu, A.¹; Ghosh, A.²; Mandal, A.³; Martin, N.⁴ and Pardo, L.⁴ ¹Indian Statistical Institute, Kolkata 700108, India

¹Indian Statistical Institute, Kolkata 700108, India ²University of Oslo, Oslo, Norway ³University of Pittsburgh, Pittsburgh, PA 15260, USA ⁴Complutense University of Madrid, 28040 Madrid, Spain

January 2, 2022

Abstract

In this paper a robust version of the classical Wald test statistics for linear hypothesis in the logistic regression model is introduced and its properties are explored. We study the problem under the assumption of random covariates although some ideas with non random covariates are also considered. The family of tests considered is based on the minimum density power divergence estimator instead of the maximum likelihood estimator and it is referred to as the Wald-type test statistic in the paper. We obtain the asymptotic distribution and also study the robustness properties of the Wald type test statistic. The robustness of the tests is investigated theoretically through the influence function analysis as well as suitable practical examples. It is theoretically established that the level as well as the power of the Wald-type tests are stable against contamination, while the classical Wald type test breaks down in this scenario. Some classical examples are presented which numerically substantiate the theory developed. Finally a simulation study is included to provide further confirmation of the validity of the theoretical results established in the paper.

MSC: 62F35, 62F05

Keywords: Influence function, Logistic regression, Minimum density power divergence estimators, Random explatory variables, Robustness, Wald-type test statistics.

1 Introduction

Experimental settings often include dichotomous response data, wherein a Bernoulli model may be assumed for the independence response variables $Y_1, ..., Y_n$, with

$$\Pr(Y_i = 1) = \pi_i \text{ and } \Pr(Y_i = 0) = 1 - \pi_i, \ i = 1, ..., n.$$

In many cases, a series of explanatory variables $x_{i0}, ..., x_{ik}$ may be associated with each Y_i ($x_{i0} = 1$, $x_{ij} \in \mathbb{R}, i = 1, ..., n, j = 1, ..., k, k < n$). We shall assume that the binomial parameter, π_i , is linked to the linear predictor $\sum_{j=0}^{k} \beta_j x_{ij}$ via the logit function, i.e.,

$$\operatorname{logit}\left(\pi_{i}\right) = \sum_{j=0}^{k} \beta_{j} x_{ij},\tag{1}$$

where logit(p) = log(p/(1-p)). In the following, we shall denote the binomial parameter π_i , by

$$\pi_i = \pi(\boldsymbol{x}_i^T \boldsymbol{\beta}) = \frac{e^{\boldsymbol{x}_i^T \boldsymbol{\beta}}}{1 + e^{\boldsymbol{x}_i^T \boldsymbol{\beta}}}, \ i = 1, ..., n,$$
(2)

where $\boldsymbol{x}_i^T = (x_{i0}, ..., x_{ik})$ and $\boldsymbol{\beta} = (\beta_0, ..., \beta_k)^T$ is a (k+1)-dimensional vector of unknown parameters with $\beta_i \in (-\infty, \infty)$. The "design matrix", $\mathbb{X} = (\boldsymbol{x}_1, ..., \boldsymbol{x}_n)^T$, is assumed to be full rank $(rank(\mathbb{X}) = k+1)$, without any loss of generality.

Let M be any matrix of r rows and k + 1 columns with rank(M) = r, and m a vector of order r with specified constants such that rank $(M^T, m) = r$. If we are interested in testing

$$H_0: \boldsymbol{M}^T \boldsymbol{\beta} = \boldsymbol{m},\tag{3}$$

the Wald test statistic is usually used in which β is estimated using the maximum likelihood estimator (MLE). Notice that if we consider $M = I_{k+1}$ and $m = \beta_0$, we get the Wald-type test statistic presented by Bianco and Martinez (2009) based on a weighted Bianco and Yohai (1996) estimator. It is well known that the MLE of β can be severely affected by outlying observations. Croux et al. (2002) discuss the breakdown behavior of the MLE in the logistic regression model and show that the MLE breaks down when several outliers are added to a data set. In the recdent years several authors have attempted to derive robust estimates of the parameters in the logistic regression model; see for instance Pregibon (1982), Morgenthaler (1992), Carrol and Pedersen (1993), Cristmann (1994), Bianco and Yohai (1996), Croux and Haesbroeck (2003), Bondell (2005, 2008) and Hobza et al. (2012). Our interest in this paper is to present a family of Wald-type test statistics based on the robust minimum density power divergence estimator for testing the general linear hypothesis given in (3).

In Section 2 we present the minimum density power divergence estimator for β . The Wald-type test statistics, based on the minimum density power divergence estimator, are presented in Section 3, as well as their asymptotic properties. The theoretical robustness properties are presented in Section 4 and finally, Section 5 and 6 are devoted to present a simulation study and real data examples, respectively.

2 Minimum density power divergence estimator

If we denote by $y_1, ..., y_n$ the observed values of the random variables $Y_1, ..., Y_n$, the likelihood function for the logistic regression model is given by

$$\mathcal{L}(\boldsymbol{\beta}) = \prod_{i=1}^{n} \pi^{y_i}(\boldsymbol{x}_i^T \boldsymbol{\beta}) \left(1 - \pi(\boldsymbol{x}_i^T \boldsymbol{\beta})\right)^{1-y_i}.$$
(4)

So the MLE of β , $\hat{\beta}$, is obtained minimizing log-likelihood function almost surely over β belonging to

$$\Theta = \left\{ (\beta_0, ..., \beta_k)^T : \beta_j \in (-\infty, \infty), \ j = 0, ..., k \right\} = \mathbb{R}^{k+1}$$

We consider the probability vectors,

$$\widehat{p} = \left(\frac{y_1}{n}, \frac{1-y_1}{n}, \frac{y_2}{n}, \frac{1-y_2}{n}, \dots, \frac{y_n}{n}, \frac{1-y_n}{n}\right)^T$$

and

$$\boldsymbol{p}\left(\boldsymbol{\beta}\right) = \left(\pi(\boldsymbol{x}_{1}^{T}\boldsymbol{\beta})\frac{1}{n}, \left(1 - \pi(\boldsymbol{x}_{1}^{T}\boldsymbol{\beta})\right)\frac{1}{n}, \dots, \pi(\boldsymbol{x}_{n}^{T}\boldsymbol{\beta})\frac{1}{n}, \left(1 - \pi(\boldsymbol{x}_{n}^{T}\boldsymbol{\beta})\right)\frac{1}{n}\right)^{T}$$

The Kullback-Leibler divergence measure between the probability vectors \hat{p} and $p(\beta)$ is given by

$$d_{KL}\left(\widehat{\boldsymbol{p}}, \boldsymbol{p}\left(\boldsymbol{\beta}\right)\right) = \sum_{i=1}^{n} \sum_{j=1}^{2} \frac{y_{ij}}{n} \log \frac{y_{ij}}{\pi_j(\boldsymbol{x}_i^T \boldsymbol{\beta})},\tag{5}$$

where

$$\pi_1(\boldsymbol{x}_i^T\boldsymbol{\beta}) = \pi(\boldsymbol{x}_i^T\boldsymbol{\beta}), \ \pi_2(\boldsymbol{x}_i^T\boldsymbol{\beta}) = 1 - \pi(\boldsymbol{x}_i^T\boldsymbol{\beta}), \ y_{i1} = y_i \text{ and } y_{i2} = 1 - y_i.$$

It is not difficult to establish that

$$d_{KL}\left(\widehat{\boldsymbol{p}}, \boldsymbol{p}\left(\boldsymbol{\beta}\right)\right) = c - \frac{1}{n} \log \mathcal{L}\left(\boldsymbol{\beta}\right).$$
(6)

Therefore, the MLE of β can be defined by

$$\widehat{\boldsymbol{\beta}} = \arg\min_{\boldsymbol{\beta}\in\Theta} d_{KL}\left(\widehat{\boldsymbol{p}}, \boldsymbol{p}\left(\boldsymbol{\beta}\right)\right).$$
(7)

Based on (7) we can use any divergence measure $d(\hat{p}, p(\beta))$ in order to define a minimum divergence estimator for β . In this paper we shall use the density power divergence measure defined by Basu et al. (1998) because the minimum density power divergence estimators have excellent robustness properties, see for instance Basu et al. (2011, 2013, 2015, 2016), Ghosh et al. (2015, 2016). The density power divergence between the probability vectors \hat{p} and $p(\beta)$ is given by

$$d_{\lambda}\left(\widehat{\boldsymbol{p}},\boldsymbol{p}\left(\boldsymbol{\beta}\right)\right) = \frac{1}{n^{1+\lambda}} \left\{ \sum_{i=1}^{n} \left(\sum_{j=1}^{2} \pi_{j}^{1+\lambda}(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}) - \left(1 + \frac{1}{\lambda}\right) \sum_{j=1}^{2} y_{ij}\pi_{j}^{\lambda}(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}) \right) + \frac{n}{\lambda} \right\}$$
(8)

for $\lambda > 0$. For $\lambda = 0$, we have

$$d_{0}\left(\widehat{\boldsymbol{p}},\boldsymbol{p}\left(\boldsymbol{\beta}
ight)
ight)=\lim_{\lambda
ightarrow0}d_{\lambda}\left(\widehat{\boldsymbol{p}},\boldsymbol{p}\left(\boldsymbol{\beta}
ight)
ight)=d_{KL}\left(\widehat{\boldsymbol{p}},\boldsymbol{p}\left(\boldsymbol{\beta}
ight)
ight).$$

Based on (7) and (8), we shall define the minimum density power divergence estimator in the following way.

Definition 1 The minimum density power divergence estimator for the parameter β , $\hat{\beta}_{\lambda}$, in the logistic regression model is given by

$$\widehat{\boldsymbol{\beta}}_{\lambda} = \arg\min_{\boldsymbol{\beta}\in\Theta} d_{\lambda}\left(\widehat{\boldsymbol{p}}, \boldsymbol{p}\left(\boldsymbol{\beta}\right)\right),$$

where $d_{\lambda}\left(\widehat{\boldsymbol{p}}, \boldsymbol{p}\left(\boldsymbol{\beta}\right)\right)$ was defined in (8).

In order to obtain the estimating equations we must get the derivative of (8) with respect to β . First we are going to write expression (8) in the following way,

$$d_{\lambda}\left(\widehat{\boldsymbol{p}},\boldsymbol{p}\left(\boldsymbol{\beta}\right)\right) = \frac{1}{n^{1+\lambda}} \left\{ \sum_{i=1}^{n} \left(\pi^{1+\lambda} \left(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}\right) + \left(1 - \pi\left(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}\right)\right)^{1+\lambda} - \left(1 + \frac{1}{\lambda}\right) \left(y_{i}\pi^{\lambda}\left(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}\right) + \left(1 - y_{i}\right) \left(1 - \pi\left(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}\right)\right)^{\lambda}\right) \right) + \frac{n}{\lambda} \right\}.$$

Now, taking into account that

$$\frac{\partial \pi(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta})}{\partial \boldsymbol{\beta}} = \pi(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}) \left(1 - \pi(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta})\right) \boldsymbol{x}_{i} \text{ and } \frac{\partial \left(1 - \pi(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta})\right)}{\partial \boldsymbol{\beta}} = -\pi(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}) \left(1 - \pi(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta})\right) \boldsymbol{x}_{i}$$

and after some algebra, we get

$$\frac{\partial d_{\lambda}\left(\widehat{\boldsymbol{p}},\boldsymbol{p}\left(\boldsymbol{\beta}\right)\right)}{\partial \boldsymbol{\beta}} = \frac{1+\lambda}{n^{\lambda+1}} \sum_{i=1}^{n} \left(e^{\lambda \boldsymbol{x}_{i}^{T}\boldsymbol{\beta}} + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}}\right) \frac{e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}} - y_{i}\left(1 + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}}\right)}{\left(1 + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}}\right)^{\lambda+2}} \boldsymbol{x}_{i}$$

Therefore, the estimating equations for $\lambda > 0$ are given by

$$\sum_{i=1}^{n} \frac{e^{\lambda \boldsymbol{x}_{i}^{T}\boldsymbol{\beta}} + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}}}{(1 + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}})^{\lambda+1}} \left(\pi(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}) - y_{i}\right) \boldsymbol{x}_{i} = \boldsymbol{0},\tag{9}$$

where $\pi(\boldsymbol{x}_i^T\boldsymbol{\beta})$ is (2). Based on the previous results we have established the following theorem.

Theorem 2 The minimum density power divergence estimator of β , $\hat{\beta}_{\lambda}$, can be obtained as the solution of the system of equations given in (9).

If we consider $\lambda = 0$ in (9), we get the estimating equations for the MLE as

$$\sum_{i=1}^{n} \left(\pi(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}) - y_{i} \right) \boldsymbol{x}_{i} = \boldsymbol{0}.$$

Based on expression (9), we can write the MDPDE for the logistic regression model by

$$\sum_{i=1}^{n} oldsymbol{\Psi}_{\lambda}\left(oldsymbol{x}_{i},y_{i},oldsymbol{eta}
ight) = oldsymbol{0},$$

with

$$\Psi_{\lambda}\left(\boldsymbol{x}_{i}, y_{i}, \boldsymbol{\beta}\right) = \left(e^{\lambda \boldsymbol{x}_{i}^{T}\boldsymbol{\beta}} + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}}\right) \frac{e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}} - y_{i}(1 + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}})}{(1 + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}})^{\lambda+2}} \boldsymbol{x}_{i}.$$
(10)

In order to get the asymptotic distribution of the MDPDE of β , $\hat{\beta}_{\lambda}$, we are going to assume that not only the explanatory variables are random but are also identically distributed and moreover

$$\left(\boldsymbol{X}_{1},Y_{1}\right),...,\left(\boldsymbol{X}_{n},Y_{n}\right)$$

are independent and identically distributed. We shall assume that $X_1, ..., X_n$ is a random sample from a random variable X with marginal distribution function H(x). By following the method given in Maronna et al. (2006), the asymptotic variance covariance matrix of $\sqrt{n} \hat{\beta}_{\lambda}$ is

$$\boldsymbol{J}_{\lambda}^{-1}\left(\boldsymbol{\beta}_{0}\right)\boldsymbol{K}_{\lambda}\left(\boldsymbol{\beta}_{0}\right)\boldsymbol{J}_{\lambda}^{-1}\left(\boldsymbol{\beta}_{0}\right),$$

where

$$\boldsymbol{K}_{\lambda}(\boldsymbol{\beta}) = E\left[\boldsymbol{\Psi}_{\lambda}\left(\boldsymbol{X}, \boldsymbol{Y}, \boldsymbol{\beta}\right) \boldsymbol{\Psi}_{\lambda}^{T}\left(\boldsymbol{X}, \boldsymbol{Y}, \boldsymbol{\beta}\right)\right] = \int_{\mathcal{X}} E\left[\boldsymbol{\Psi}_{\lambda}\left(\boldsymbol{x}, \boldsymbol{Y}, \boldsymbol{\beta}\right) \boldsymbol{\Psi}_{\lambda}^{T}\left(\boldsymbol{x}, \boldsymbol{Y}, \boldsymbol{\beta}\right)\right] dH(\boldsymbol{x}),$$

 \mathcal{X} is the support of \boldsymbol{X} , and

$$\boldsymbol{J}_{\lambda}(\boldsymbol{\beta}) = E\left[\frac{\partial \boldsymbol{\Psi}_{\lambda}(\boldsymbol{X}, \boldsymbol{Y}, \boldsymbol{\beta})}{\partial \boldsymbol{\beta}^{T}}\right] = \int_{\mathcal{X}} E\left[\frac{\partial \boldsymbol{\Psi}_{\lambda}(\boldsymbol{x}, \boldsymbol{Y}, \boldsymbol{\beta})}{\partial \boldsymbol{\beta}^{T}}\right] dH(\boldsymbol{x})$$

In relation to the matrix $\boldsymbol{K}_{\lambda}(\boldsymbol{\beta}_{0})$, we have

$$E\left[\Psi_{\lambda}\left(\boldsymbol{x},Y,\boldsymbol{\beta}\right)\Psi_{\lambda}^{T}\left(\boldsymbol{x},Y,\boldsymbol{\beta}\right)\right] = \frac{\left(e^{\lambda\boldsymbol{x}^{T}\boldsymbol{\beta}} + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}\right)^{2}}{\left(1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}\right)^{2(\lambda+2)}}E\left[\left(e^{\boldsymbol{x}^{T}\boldsymbol{\beta}} - Y(1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})\right)^{2}\right]\boldsymbol{x}\boldsymbol{x}^{T},$$

but $E\left[Y^2\right] = \pi(\boldsymbol{x}^T\boldsymbol{\beta})$ and

$$E\left[\left(e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}-Y(1+e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})\right)^{2}\right]=e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}$$

Therefore

$$\boldsymbol{K}_{\lambda}(\boldsymbol{\beta}) = E\left[\boldsymbol{\Psi}_{\lambda}\left(\boldsymbol{X}, \boldsymbol{Y}, \boldsymbol{\beta}\right) \boldsymbol{\Psi}_{\lambda}^{T}\left(\boldsymbol{X}, \boldsymbol{Y}, \boldsymbol{\beta}\right)\right] = \int_{\mathcal{X}} \frac{(e^{\lambda \boldsymbol{x}^{T}\boldsymbol{\beta}} + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})^{2}}{(1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})^{2(\lambda+2)}} e^{\boldsymbol{x}^{T}\boldsymbol{\beta}} \boldsymbol{x} \boldsymbol{x}^{T} dH(\boldsymbol{x}).$$
(11)

An estimator of $\boldsymbol{K}_{\lambda}\left(\boldsymbol{\beta}\right)$ will be

$$\widehat{oldsymbol{K}}_{\lambda}\left(oldsymbol{eta}
ight) = \int_{\mathcal{X}} rac{(e^{\lambda oldsymbol{x}^Toldsymbol{eta}} + e^{oldsymbol{x}^Toldsymbol{eta}})^2}{(1 + e^{oldsymbol{x}^Toldsymbol{eta}})^{2(\lambda+2)}} e^{oldsymbol{x}^Toldsymbol{eta}} oldsymbol{x}^Toldsymbol{eta}} H_n(oldsymbol{x}),$$

where $H_n(\boldsymbol{x})$ the empirical distribution function associated with the sample $\boldsymbol{x}_1,...,\boldsymbol{x}_n$. Then

$$\widehat{\boldsymbol{K}}_{\lambda}\left(\boldsymbol{\beta}\right) = \frac{1}{n} \sum_{i=1}^{n} \frac{(e^{\lambda \boldsymbol{x}^{T}\boldsymbol{\beta}} + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})^{2}}{(1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})^{2(\lambda+2)}} e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}} \boldsymbol{x}_{i} \boldsymbol{x}_{i}^{T}.$$
(12)

It is interesting to observe that for $\lambda = 0$ we get

$$\begin{aligned} \widehat{\boldsymbol{K}}_{0}\left(\boldsymbol{\beta}\right) &= \frac{1}{n} \sum_{i=1}^{n} \frac{(1 + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}})^{2}}{(1 + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}})^{4}} e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}} \boldsymbol{x}_{i} \boldsymbol{x}_{i}^{T} &= \frac{1}{n} \mathbb{X}^{T} \operatorname{diag}\left(\pi_{i}(\boldsymbol{x}^{T}\boldsymbol{\beta})\left(1 - \pi_{i}(\boldsymbol{x}^{T}\boldsymbol{\beta})\right)\right)_{i=1,\dots,n} \mathbb{X} \\ &= \boldsymbol{I}_{F}\left(\boldsymbol{\beta}\right), \end{aligned}$$

with $I_{F}(\boldsymbol{\beta}_{0})$ being the Fisher information matrix associated to the logistic regression model.

To compute the matrix $\boldsymbol{J}_{\lambda}(\boldsymbol{\beta}_{0})$, first we need to calculate

$$\frac{\partial \Psi_{\lambda}(\boldsymbol{x}, y, \boldsymbol{\beta})}{\partial \boldsymbol{\beta}^{T}} = L_{1}(\boldsymbol{x}, y, \boldsymbol{\beta}) + L_{2}(\boldsymbol{x}, y, \boldsymbol{\beta}),$$

where

$$L_1(\boldsymbol{x}, y, \boldsymbol{\beta}) = (\lambda e^{\lambda \boldsymbol{x}^T \boldsymbol{\beta}} + e^{\boldsymbol{x}^T \boldsymbol{\beta}}) \frac{e^{\boldsymbol{x}^T \boldsymbol{\beta}} - y(1 + e^{\boldsymbol{x}^T \boldsymbol{\beta}})}{(1 + e^{\boldsymbol{x}^T \boldsymbol{\beta}})^{\lambda + 2}} \boldsymbol{x} \boldsymbol{x}^T$$

and

$$L_{2}(\boldsymbol{x},\boldsymbol{y},\boldsymbol{\beta}) = (e^{\lambda \boldsymbol{x}^{T}\boldsymbol{\beta}} + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}) \left(\frac{\left(e^{\boldsymbol{x}^{T}\boldsymbol{\beta}} - \boldsymbol{y}e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}\right)(1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})^{\lambda+2}}{(1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})^{2(\lambda+2)}} - \frac{(\lambda+2)\left((1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})^{\lambda+1}\right)e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}\left(e^{\boldsymbol{x}^{T}\boldsymbol{\beta}} - \boldsymbol{y}(1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})\right)}{(1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})^{2(\lambda+2)}}\right)\boldsymbol{x}\boldsymbol{x}^{T},$$

and hence

$$E\left[\frac{\partial \Psi_{\lambda}\left(\boldsymbol{x},Y,\boldsymbol{\beta}\right)}{\partial \boldsymbol{\beta}^{T}}\right] = E\left[L_{1}(\boldsymbol{x},Y,\boldsymbol{\beta})\right] + E\left[L_{2}(\boldsymbol{x},Y,\boldsymbol{\beta})\right].$$

But

$$E\left[e^{\boldsymbol{x}^T\boldsymbol{\beta}} - Y(1+e^{\boldsymbol{x}^T\boldsymbol{\beta}})\right] = e^{\boldsymbol{x}^T\boldsymbol{\beta}} - \frac{e^{\boldsymbol{x}^T\boldsymbol{\beta}}}{1+e^{\boldsymbol{x}^T\boldsymbol{\beta}}}(1+e^{\boldsymbol{x}^T\boldsymbol{\beta}}) = 0.$$

Therefore

$$E[L_1(\boldsymbol{x}, Y, \boldsymbol{\beta})] = \mathbf{0}_{(k+1)(k+1)}.$$

On the other hand

$$E\left[L_{2}(\boldsymbol{x}, \boldsymbol{Y}, \boldsymbol{\beta})\right] = \frac{e^{\lambda \boldsymbol{x}^{T}\boldsymbol{\beta}} + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}}{(1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})^{2(\lambda+2)}} \left((1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})^{\lambda+2}E\left[e^{\boldsymbol{x}^{T}\boldsymbol{\beta}} - \boldsymbol{Y}e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}\right] \\ + (\lambda+2)\left(1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}\right)^{\lambda+1}e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}E\left[e^{\boldsymbol{x}^{T}\boldsymbol{\beta}} - \boldsymbol{Y}(1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})\right]\right)\boldsymbol{x}\boldsymbol{x}^{T} \\ = \frac{e^{\lambda \boldsymbol{x}^{T}\boldsymbol{\beta}} + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}}{(1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})^{\lambda+3}}e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}\boldsymbol{x}\boldsymbol{x}^{T}.$$

Finally,

$$\boldsymbol{J}_{\lambda}(\boldsymbol{\beta}) = \int_{\mathcal{X}} E\left[\frac{\partial \boldsymbol{\Psi}_{\lambda}(\boldsymbol{x}, Y, \boldsymbol{\beta})}{\partial \boldsymbol{\beta}^{T}}\right] dH(\boldsymbol{x}) \qquad (13)$$

$$= \int_{\mathcal{X}} \frac{e^{\lambda \boldsymbol{x}^{T}\boldsymbol{\beta}} + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}}}{(1 + e^{\boldsymbol{x}^{T}\boldsymbol{\beta}})^{\lambda+3}} e^{\boldsymbol{x}^{T}\boldsymbol{\beta}} \boldsymbol{x} \boldsymbol{x}^{T} dH(\boldsymbol{x}),$$

and an estimator of $\boldsymbol{J}_{\lambda}\left(\boldsymbol{\beta}_{0}\right)$ is given by

$$\widehat{\boldsymbol{J}}_{\lambda}\left(\boldsymbol{\beta}\right) = \frac{1}{n} \sum_{i=1}^{n} \frac{e^{\lambda \boldsymbol{x}_{i}^{T}\boldsymbol{\beta}} + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}}}{(1 + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}})^{\lambda+3}} e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}} \boldsymbol{x}_{i} \boldsymbol{x}_{i}^{T}.$$
(14)

In particular, for $\lambda = 0$, we have

$$\widehat{\boldsymbol{J}}_{0}(\boldsymbol{\beta}) = \frac{1}{n} \mathbb{X}^{T} \operatorname{diag} \left(\pi_{i}(\boldsymbol{x}^{T}\boldsymbol{\beta}) \left(1 - \pi_{i}(\boldsymbol{x}^{T}\boldsymbol{\beta}) \right) \right)_{i=1,\dots,n} \mathbb{X}$$
$$= \boldsymbol{I}_{F}(\boldsymbol{\beta}).$$

From the sequence of above results, the next theorem follows.

Theorem 3 The asymptotic distribution of the MDPDE for β , $\hat{\beta}_{\lambda}$, is given by

$$\sqrt{n}(\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{\beta}_{0}) \xrightarrow[n \to \infty]{\mathcal{L}} \mathcal{N}(\boldsymbol{0}, \boldsymbol{\Sigma}_{\lambda}(\boldsymbol{\beta}_{0}))$$

where

$$\boldsymbol{\Sigma}_{\lambda}\left(\boldsymbol{\beta}_{0}\right) = \boldsymbol{J}_{\lambda}^{-1}\left(\boldsymbol{\beta}_{0}\right) \boldsymbol{K}_{\lambda}\left(\boldsymbol{\beta}_{0}\right) \boldsymbol{J}_{\lambda}^{-1}\left(\boldsymbol{\beta}_{0}\right)$$

and the matrices $J_{\lambda}(\beta_0)$ and $K_{\lambda}(\beta_0)$ where defined in (13) and (11), respectively.

Remark 4 We have considered that the covariates are random, a crutial assumption to get the asymptotic distribution of the MDPDE by using, "in part", the standard asymptotic theory for M-estimators. It is interesting to highlight that whenever the covariates were non-stochastic (fixed design case), the asymptotic distribution of the MDPDE could be obtained from Ghosh and Basu (2013) without using the standard asymptotic theory of M-estimators. In order to present the results in the most general setting, we shall assume that the random variables Y_i with i = 1, ..., I, are binomial with parameters n_i and $\pi_i = \pi(\mathbf{x}_i^T \boldsymbol{\beta})$ instead of Bernoulli random variables. We shall denote by $N = \sum_{i=1}^{I} n_i$ and let n_{i1} denotes the observed value of Y_i . We will assume that I is fixed and for each i = 1, ..., I, construct the independent and identically distributed latent observations $z_{i1}, ..., z_{in_i}$ each following a Bernoulli distribution with probability π and $n_{i1} = \sum_{j=1}^{n_i} z_{ij}$. Then, N random observations $z_{11}, ..., z_{1n_1}, z_{21}, ..., z_{2n_2}, ..., z_{I_1}, ..., z_{In_I}$ are independent but have possibly different distribution with $z_{ij} \sim Ber(\pi_i)$. This falls under the general set-up of independent but non-homogeneous observations as considered in Ghosh

and Basu (2013) and hence it is immediately seen that the corresponding estimating equations for the MDPDE, $\hat{\beta}^*_{\lambda}$ in this context, for $\lambda > 0$ are given by

$$\sum_{i=1}^{I} \frac{e^{\lambda \boldsymbol{x}_{i}^{T}\boldsymbol{\beta}} + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}}}{(1 + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}})^{\lambda+1}} \left(n_{i}\pi(\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}) - n_{i1} \right) \boldsymbol{x}_{i} = \boldsymbol{0}$$

and for $\lambda = 0$, by

$$\sum_{i=1}^{I} \left(n_i \pi(\boldsymbol{x}_i^T \boldsymbol{\beta}) - n_{i1} \right) \boldsymbol{x}_i = \boldsymbol{0}.$$
(15)

Now, assuming

$$\lim_{N \to \infty} \frac{n_i}{N} = \alpha_i \in (0, 1), \ i = 1, ..., I,$$

and following Ghosh and Basu (2013), we get the asymptotic distribution of the MDPDE of β , $\hat{\beta}^*_{\lambda}$, as given by

$$\sqrt{N}(\widehat{\boldsymbol{\beta}}_{\lambda}^{*} - \boldsymbol{\beta}_{0}) \xrightarrow[N \to \infty]{\mathcal{L}} \mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma}^{*}(\boldsymbol{\beta}_{0}))$$
(16)

where

$$\boldsymbol{\Sigma}^{*}\left(\boldsymbol{\beta}_{0}\right) = \boldsymbol{J}^{*-1}\left(\boldsymbol{\beta}_{0}\right)\boldsymbol{K}^{*}\left(\boldsymbol{\beta}_{0}\right)\boldsymbol{J}^{*-1}\left(\boldsymbol{\beta}_{0}\right)$$

Here, the matrices $J^*(\beta_0)$ and $K^*(\beta_0)$ can be obtained directly from the general results of Ghosh and Basu (2013) or from the simplified results in the context of Bernoulli logistic regression with fixed design in Ghosh and Basu (2015) and are given by

$$\boldsymbol{J}^{*}\left(\boldsymbol{\beta}_{0}\right) = \sum_{i=1}^{I} \alpha_{i} e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}} \frac{e^{\lambda \boldsymbol{x}_{i}^{T}\boldsymbol{\beta}} + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}}}{(1 + e^{\boldsymbol{x}_{i}^{T}\boldsymbol{\beta}})^{\lambda+3}} \boldsymbol{x}_{i} \boldsymbol{x}_{i}^{T},$$

and

$$\boldsymbol{K}^{*}(\boldsymbol{\beta}_{0}) = \sum_{i=1}^{I} \alpha_{i} e^{\boldsymbol{x}_{i}^{T} \boldsymbol{\beta}} \frac{(e^{\lambda \boldsymbol{x}_{i}^{T} \boldsymbol{\beta}} + e^{\boldsymbol{x}_{i}^{T} \boldsymbol{\beta}})^{2}}{(1 + e^{\boldsymbol{x}_{i}^{T} \boldsymbol{\beta}})^{2(\lambda+2)}} \boldsymbol{x}_{i} \boldsymbol{x}_{i}^{T}.$$

For $\lambda = 0$, it is clear, based on (15), that we get the classical likelihood estimator. We can observe that in this situation

$$oldsymbol{J}^{*}\left(oldsymbol{eta}_{0}
ight)=oldsymbol{K}^{*}\left(oldsymbol{eta}_{0}
ight)=oldsymbol{I}_{F}\left(oldsymbol{eta}_{0}
ight)$$

and we get the classical result,

$$\sqrt{N}(\widehat{\boldsymbol{\beta}}_{\lambda=0}^{*}-\boldsymbol{\beta}_{0}) \xrightarrow[N \to \infty]{\mathcal{L}} \mathcal{N}\left(\boldsymbol{0}, \boldsymbol{I}_{F}^{-1}\left(\boldsymbol{\beta}_{0}\right)\right).$$

3 Wald type test statistic for testing linear hypothesis

Based on the asymptotic distribution of $\hat{\beta}_{\lambda}$ we are going to define a family of Wald-type test statistics for testing the null hypothesis

$$H_0: \boldsymbol{M}^T \boldsymbol{\beta} = \boldsymbol{m},\tag{17}$$

where M^T is any matrix of r rows and k+1 columns and m a vector of order r of specified constant. We assume that the matrix M^T has full row rank, i.e., rank (M) = r.

Definition 5 Let $\hat{\boldsymbol{\beta}}_{\lambda}$ be the minimum power divergence estimator. The family of Wald type test statistics for testing the null hypothesis given in (17) is given by

$$W_{n} = n(\boldsymbol{M}^{T}\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{m})^{T} \left(\boldsymbol{M}^{T}\boldsymbol{J}_{\lambda}^{-1}(\widehat{\boldsymbol{\beta}}_{\lambda})\boldsymbol{K}_{\lambda}(\widehat{\boldsymbol{\beta}}_{\lambda})\boldsymbol{J}_{\lambda}^{-1}(\widehat{\boldsymbol{\beta}}_{\lambda})\boldsymbol{M} \right)^{-1} (\boldsymbol{M}^{T}\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{m})$$
$$= n(\boldsymbol{M}^{T}\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{m})^{T} (\boldsymbol{M}^{T}\boldsymbol{\Sigma}_{\lambda}(\widehat{\boldsymbol{\beta}}_{\lambda})\boldsymbol{M})^{-1} (\boldsymbol{M}^{T}\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{m}).$$
(18)

In the particular case of $\lambda = 0$, i.e. $\hat{\beta}$ is the MLE, we get the classical Wald test statistic because in this case

$$\boldsymbol{J}_{\lambda=0}^{-1}\left(\boldsymbol{\beta}_{0}\right)\boldsymbol{K}_{\lambda=0}\left(\boldsymbol{\beta}_{0}\right)\boldsymbol{J}_{\lambda=0}^{-1}\left(\boldsymbol{\beta}_{0}\right)=\boldsymbol{I}_{F}^{-1}\left(\boldsymbol{\beta}_{0}\right).$$

Theorem 6 The asymptotic distribution of the Wald type test statistic, W_n , defined in (18), under the null hypothesis given in (17), is a chi-square distribution with r degrees of freedom.

Proof. We have $M^T \hat{\boldsymbol{\beta}}_{\lambda} - m = M^T (\hat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{\beta}_0)$ and $\sqrt{n} (\hat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{\beta}_0) \xrightarrow[n \to \infty]{\mathcal{L}} \mathcal{N}(\boldsymbol{0}, \boldsymbol{\Sigma}_{\lambda}(\boldsymbol{\beta}_0))$. Therefore

$$\sqrt{n}(\boldsymbol{M}^{T}\widehat{\boldsymbol{eta}}_{\lambda}-\boldsymbol{m}) \stackrel{\mathcal{L}}{\underset{n o \infty}{\longrightarrow}} \mathcal{N}\left(\boldsymbol{0}, \boldsymbol{M}^{T}\boldsymbol{\Sigma}_{\lambda}\left(\boldsymbol{eta}_{0}
ight) \boldsymbol{M}
ight)$$

and since $M^T \Sigma_{\lambda}(\beta_0) M \left(M^T J_{\lambda}^{-1}(\beta_0) K_{\lambda}(\beta_0) J_{\lambda}^{-1}(\beta_0) M \right)^{-1} = I_{r \times r}$, the asymptotic distribution of W_n is a chi-square distribution with r degrees of freedom.

Remark 7 If we consider

$$\boldsymbol{M}^{T} = \begin{pmatrix} \boldsymbol{0}_{k \times 1} & \boldsymbol{I}_{k \times k} \end{pmatrix}_{k \times (k+1)}$$
(19)

we have

$$M^T \beta = 0,$$

if and only if $\beta_i = 0$, i = 1, ..., k. Therefore, we can consider the Wald-type test statistics with \mathbf{M}^T defined in (19) for testing

$$H_0: \beta_1 = \beta_2 = \dots = \beta_k = 0.$$

In this case, the asymptotic distribution of the Wald type test statistic is a chi square distribution with k degrees of freedom. If we consider \mathbf{M}^T to be a vector with all elements equal zero except for the (i + 1)-th term, equals 1, we can test

$$H_0:\beta_i=0.$$

Based on the previous theorem the null hypothesis given in (17) will be rejected if we have that

$$W_n > \chi^2_{r,\alpha},\tag{20}$$

where $\chi^2_{r,\alpha}$ is the quantile of order $1 - \alpha$ for a chi-square with r degrees of freedom Let us consider $\beta^* \in \Theta$ such that $M^T \beta^* \neq m$, i.e., β^* does not belong to the null hypothesis. We denote

$$q_{\boldsymbol{\beta}_{1}}(\boldsymbol{\beta}_{2}) = \left(\boldsymbol{M}^{T}\boldsymbol{\beta}_{1} - \boldsymbol{m}\right)^{T} \left(\boldsymbol{M}^{T}\boldsymbol{\Sigma}_{\lambda}\left(\boldsymbol{\beta}_{2}\right)\boldsymbol{M}\right)^{-1} \left(\boldsymbol{M}^{T}\boldsymbol{\beta}_{1} - \boldsymbol{m}\right)$$

and we are going to get an approximation to the power function for the test statistics given in (20).

Theorem 8 Let $\beta^* \in \Theta$, with $M^T \beta^* \neq m$, be the true value of the parameter so that $\hat{\beta}_{\lambda} \xrightarrow[n \to \infty]{P} \beta^*$. The power function of the test statistic given in (20), in β^* , is given by

$$\pi\left(\boldsymbol{\beta}^{*}\right) = 1 - \Phi_{n}\left(\frac{1}{\sigma\left(\boldsymbol{\beta}^{*}\right)}\left(\frac{\chi_{r,\alpha}^{2}}{\sqrt{n}} - \sqrt{n}q_{\boldsymbol{\beta}^{*}}\left(\boldsymbol{\beta}^{*}\right)\right)\right),\tag{21}$$

where $\Phi_n(x)$ tends uniformly to the standard normal distribution $\Phi(x)$ and $\sigma(\beta^*)$ is given by

$$\sigma^{2}(\boldsymbol{\beta}^{*}) = \frac{\partial q_{\boldsymbol{\beta}}(\boldsymbol{\beta}^{*})}{\partial \boldsymbol{\beta}^{T}}\Big|_{\boldsymbol{\beta}=\boldsymbol{\beta}^{*}} \boldsymbol{\Sigma}_{\lambda}(\boldsymbol{\beta}_{0}) \frac{\partial q_{\boldsymbol{\beta}}(\boldsymbol{\beta}^{*})}{\partial \boldsymbol{\beta}}\Big|_{\boldsymbol{\beta}=\boldsymbol{\beta}^{*}}$$

Proof. We have

$$\pi\left(\boldsymbol{\beta}^{*}\right) = \Pr\left(W_{n} > \chi_{r,\alpha}^{2}\right) = \Pr\left(n\left(q_{\widehat{\boldsymbol{\beta}}_{\lambda}}(\widehat{\boldsymbol{\beta}}_{\lambda}) - q_{\boldsymbol{\beta}^{*}}(\boldsymbol{\beta}^{*})\right) > \chi_{r,\alpha}^{2} - nq_{\boldsymbol{\beta}^{*}}(\boldsymbol{\beta}^{*})\right)$$
$$= \Pr\left(\sqrt{n}\left(q_{\widehat{\boldsymbol{\beta}}_{\lambda}}(\widehat{\boldsymbol{\beta}}_{\lambda}) - q_{\boldsymbol{\beta}^{*}}(\boldsymbol{\beta}^{*})\right) > \frac{\chi_{r,\alpha}^{2}}{\sqrt{n}} - \sqrt{n}q_{\boldsymbol{\beta}^{*}}(\boldsymbol{\beta}^{*})\right).$$

Now we are going to get the asymptotic distribution of the random variable $\sqrt{n}(q_{\hat{\beta}_{\lambda}}(\hat{\beta}_{\lambda}) - q_{\beta^*}(\beta^*))$. It is clear that $q_{\hat{\beta}_{\lambda}}(\hat{\beta}_{\lambda})$ and $q_{\hat{\beta}_{\lambda}}(\beta^*)$ have the same asymptotic distribution because $\hat{\beta}_{\lambda} \xrightarrow{P}_{n \to \infty} \beta^*$. A first order Taylor expansion of $q_{\hat{\beta}_{\lambda}}(\beta^*)$ at $\hat{\beta}_{\lambda}$ around β^* gives

$$q_{\widehat{\boldsymbol{\beta}}_{\lambda}}(\boldsymbol{\beta}^{*}) - q_{\boldsymbol{\beta}^{*}}(\boldsymbol{\beta}^{*}) = \frac{\partial q_{\boldsymbol{\beta}}(\boldsymbol{\beta}^{*})}{\partial \boldsymbol{\beta}^{T}} \Big|_{\boldsymbol{\beta} = \boldsymbol{\beta}^{*}} \left(\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{\beta}^{*}\right) + o_{p}\left(\left\|\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{\beta}^{*}\right\|\right).$$

Therefore it holds

$$\sqrt{n}\left(q_{\widehat{\boldsymbol{\beta}}_{\lambda}}(\widehat{\boldsymbol{\beta}}_{\lambda}) - q_{\boldsymbol{\beta}^{*}}(\boldsymbol{\beta}^{*})\right) \xrightarrow[n \to \infty]{\mathcal{L}} \mathcal{N}\left(0, \sigma^{2}\left(\boldsymbol{\beta}^{*}\right)\right),$$

where

$$\sigma^{2}(\boldsymbol{\beta}^{*}) = \frac{\partial q_{\boldsymbol{\beta}}(\boldsymbol{\beta}^{*})}{\partial \boldsymbol{\beta}^{T}}\Big|_{\boldsymbol{\beta}=\boldsymbol{\beta}^{*}} \boldsymbol{J}_{\lambda}^{-1}(\boldsymbol{\beta}_{0}) \boldsymbol{K}_{\lambda}(\boldsymbol{\beta}_{0}) \boldsymbol{J}_{\lambda}^{-1}(\boldsymbol{\beta}_{0}) \frac{\partial q_{\boldsymbol{\beta}}(\boldsymbol{\beta}^{*})}{\partial \boldsymbol{\beta}}\Big|_{\boldsymbol{\beta}=\boldsymbol{\beta}^{*}}.$$

Now the result follows. \blacksquare

Remark 9 Based on the previous theorem we can obtain the sample size necessary to get a fix power $\pi(\beta^*) = \pi_0$. From (21), we must solve the equation

$$1 - \pi_0 = \Phi\left(\frac{1}{\sigma\left(\boldsymbol{\beta}^*\right)}\left(\frac{\chi^2_{r,\alpha}}{\sqrt{n}} - \sqrt{n}q_{\boldsymbol{\beta}^*}(\boldsymbol{\beta}^*)\right)\right)$$

and we get that $n = [n^*] + 1$ with

$$n^* = \frac{A + B + \sqrt{A(A + 2B)}}{2q_{\boldsymbol{\beta}^*}^2(\boldsymbol{\beta}^*)}$$

being

$$A = \sigma^{2} \left(\beta^{*} \right) \left(\Phi^{-1} \left(1 - \pi_{0} \right) \right)^{2} \text{ and } B = 2q_{\beta^{*}} \left(\beta^{*} \right) \chi^{2}_{r,\alpha}.$$

In the following theorem we present an approximation to the power function at the contiguous alternative hypothesis

$$\boldsymbol{\beta}_n = \boldsymbol{\beta}_0 + n^{-1/2} \boldsymbol{d}, \tag{22}$$

with \boldsymbol{d} satisfying $\boldsymbol{\beta}_0 + n^{-1/2} \boldsymbol{d} \in \Theta$.

Theorem 10 An approximation of the power function for the test statistic given in (20), in $\beta_n = \beta_0 + n^{-1/2} d$ is given by

$$\pi\left(\boldsymbol{\beta}_{n}\right) = 1 - F_{\chi_{r}^{2}\left(\delta\right)}\left(\chi_{r,\alpha}^{2}\right),$$

where $F_{\chi_r^2(\delta)}$ is the distribution function of a non-central chi-square with p degrees of freedom and non-centrality parameter δ given by $\delta = \mathbf{d}^T \Sigma_{\lambda}(\boldsymbol{\beta}_0) \mathbf{d}$.

4 Robustness Analysis

4.1 Influence function of the MDPDE

We will consider the influence function analysis of Hampel et al. (1986) to study the robustness of our proposed MDPDE and the corresponding Wald-type test of general linear hypothesis in the logistic regression model. Since the MDPDE can be written in term of a *M*-estimator as shown in Section 2 with ψ -function given by (10), we can apply directly the results of the M-estimation theory of Hampel et al. (1986) in order to get the influence function of the proposed MDPDE.

However, we first need to re-define the minimum density power divergence estimator β_{λ} from Definition 1 in terms of a statistical functional. Let us assume the stochastic nature of the covariates X and that the observations $(X_1, Y_1), \ldots, (X_n, Y_n)$ are i.i.d. with some joint distribution G. Then we define the required statistical functional corresponding to $\hat{\beta}_{\lambda}$ as follows.

Definition 11 The minimum DPD functional $T_{\lambda}(G)$, corresponding to the minimum DPD estimator $\hat{\beta}_{\lambda}$, at the joint distribution G is defined as the solution of the system of equations

$$E_G[\Psi_\lambda(\boldsymbol{X},\boldsymbol{Y},\boldsymbol{\beta})] = \mathbf{0}$$

with respect to β , whenever the solution exists.

Now, if G_0 denotes the joint model distribution with the true parameter value β_0 under which

$$P_{G_0}(Y_i = 1 | \boldsymbol{X}_i = \boldsymbol{x}_i) = \pi(\boldsymbol{x}_i^T \boldsymbol{\beta}_0),$$

then it is easy to see that $E_{G_0}[\Psi_{\lambda}(X, Y, \beta_0)] = 0$ and hence $T_{\lambda}(G_0) = \beta_0$. Therefore, the minimum DPD functional T_{λ} is Fisher consistent.

Next, we can easily obtain the influence function for our MDPDE at the model distribution G_0 as presented in the following theorem. This can be derived either through a straightforward calculation or by applying the corresponding results from M-estimation theory of Hampel et al. (1986) and hence the proof of the theorem is omitted.

Theorem 12 The influence function of the minimum DPD functional T_{λ} , as defined in Definition 11 with tuning parameter λ , at the model distribution G_0 is given by

$$\mathcal{IF}((\boldsymbol{x}_t, y_t), T_{\lambda}, G_0) = \boldsymbol{J}_{\lambda}^{-1}(\boldsymbol{\beta}_0) \left(\boldsymbol{\Psi}_{\lambda}(\boldsymbol{x}_t, y_t, \boldsymbol{\beta}_0) - E_{G_0}[\boldsymbol{\Psi}_{\lambda}(\boldsymbol{X}, Y, \boldsymbol{\beta}_0)] \right) \\ = \boldsymbol{J}_{\lambda}^{-1}(\boldsymbol{\beta}_0) \boldsymbol{\Psi}_{\lambda}(\boldsymbol{x}_t, y_t, \boldsymbol{\beta}_0),$$

where $J_{\lambda}(\beta)$ is as defined in Section 2 of the paper and (x_t, y_t) is the point of contamination.

Before studying the above influence function, let us first recall different types of outliers in logistic regression model following the discussion in Croux and Haesbroeck (2003). A contamination point (x_t, y_t) will be a leverage point if x_t is outlying in the covariates space and will be a vertical outlier (in response) if it is not a leverage point but the residual $y_t - \pi(\boldsymbol{x}_t^T \boldsymbol{\beta})$ is large. Croux and Haesbroeck (2003) also noted that, for the maximum likelihood estimator of $\boldsymbol{\beta}$, a vertical outlier or a "good" leverage point (for which the residual is small) has bounded influence whereas a bad leverage point (e.g., misclassified observation etc.) has infinite influence for $||\boldsymbol{x}_t|| \to \infty$.

Next, in order to study the similar nature of the influence function of the MDPDE having different λ , note that the influence function given in Theorem 12 can be factored into two components as

$$\mathcal{IF}((\boldsymbol{x}_t, y_t), T_{\lambda}, G_0) = \widetilde{\Psi}_{\lambda}(\boldsymbol{x}_t^T \boldsymbol{\beta}_0, y_t) \boldsymbol{J}_{\lambda}^{-1}(\boldsymbol{\beta}_0) \boldsymbol{x}_t,$$

where the first part $\widetilde{\Psi}_{\lambda}$ depends on the score, $s = \boldsymbol{x}_t^T \boldsymbol{\beta}_0$, and the response, y_t , and is defined as

$$\widetilde{\Psi}_{\lambda}(s,y) = \frac{\left(e^{\lambda s} + e^{s}\right)\left(e^{s} - y(1+e^{s})\right)}{\left(1+e^{s}\right)^{\lambda+2}}.$$

Figure 1 shows the nature of this function over the score input at y = 0, 1 for different values of λ . Clearly, the function $\tilde{\Psi}_{\lambda}$ corresponding to $\lambda = 0$ (MLE) is unbounded as $s \to \infty$, illustrating the well-known non-robust nature of the MLE. However, for $\lambda > 0$ the function $\tilde{\Psi}_{\lambda}$ is bounded in s and becomes more re-descending as λ increase, which implies the increasing robustness of our proposed MDPDEs with increasing $\lambda > 0$.



Figure 1: Plots of $\widetilde{\Psi}_{\lambda}(s; y)$ over s for different λ and y = 0, 1.

Further, to examine the effect of different types of leverage points more clearly, following Croux and Haesbroeck (2003), in Figure 2, we present the influence function of the MDPDE of the first slope parameter β_1 over the covariates values in a logistic regression model with two independent standard normal covariates and $\beta_0 = (0, 1, 1)^T$ fixing $y_t = 0$ (without loss of generality). We can see that when both covariates tends to $-\infty$ the influence function becomes zero for all MDPDEs including the MLE (at $\lambda = 0$). These are the "good" leverage points, as noted in Croux and Haesbroeck (2003), and all MDPDEs are robust with respect to such good leverages as in the case of MLE. However, when the covariates approaches to ∞ they yield bad leverage points (generally corresponding to misclassified points) and have large influence for the MLE ($\lambda = 0$). But the influence function of the MDPDEs with $\lambda > 0$ are quite small even for these bad leverages and become even smaller as λ increases. This again proves the greater robustness of our proposed MDPDEs with larger positive λ .

Remark 13 Under the set-up of Remark 4 with non-stochastic covariate also, we can derive the influence function of the corresponding MDPDE, $\hat{\beta}^*_{\lambda}$, following Ghosh and Basu (2013). Whenever the covariates \mathbf{x}_i s are fixed, the contamination need to be considered over the conditional distribution of response given covariates which are not identical for each group with given fixed covariates. Hence, as in Ghosh and Basu (2013), we can consider the contamination in any one group or in all the group. Following the results in Ghosh and Basu (2013) or by direct calculation, we get the influence function of $\hat{\beta}^*_{\lambda}$ under contamination only in one group (i₀-th, say) with covariate \mathbf{x}_{i_0} as given by

$$\mathcal{IF}_{i_0}(y_{t_{i_0}}, T_{\lambda}, G_0) = \boldsymbol{J}_{\lambda}^{*-1}(\boldsymbol{\beta}_0) \boldsymbol{\Psi}_{\lambda}(\boldsymbol{x}_{i_0}, y_{t_{i_0}}, \boldsymbol{\beta}_0),$$



Figure 2: Influence function of the MDPDE of the first slope parameter β_1 for different λ ($y_t = 0$).

where $y_{t_{i_0}}$ is the contamination point in the contaminated distribution of Y given $\mathbf{X} = \mathbf{x}_{i_0}$. Similarly, if there is contamination in all the groups with covariates $\mathbf{x}_1, \ldots, \mathbf{x}_I$ respectively at the contamination points y_{t_1}, \ldots, y_{t_I} , then the resulting influence function has the form

$$\mathcal{IF}((y_{t_1},\ldots,y_{t_l}),T_{\lambda},G_0)=\boldsymbol{J}_{\lambda}^{*-1}(\boldsymbol{\beta}_0)\sum_{i=1}^{I}\boldsymbol{\Psi}_{\lambda}(\boldsymbol{x}_i,y_{t_i},\boldsymbol{\beta}_0),$$

Note that, since the response in a logistic regression takes only values 0 and 1, the y_{t_i} contamination points all take values only in {0,1} (misclassification errors) and hence all the above influence functions are bounded with respect to contamination in response for all $\lambda \geq 0$. Hence, the effect of these (misclassification) error in response cannot be clearly inferred only from these influence functions; see Pregibon (1982), Copas (1988) and Victoria-Feser (2000) for more such analysis of misclassification error in logistic regression with fixed design. However, the above influence functions are bounded in the values of given fixed covariates only for $\lambda > 0$, implying the robustness of the MDPDEs with $\lambda > 0$ and non-robust nature of MLE (at $\lambda = 0$) with respect to the extreme values of the fixed design in any one group.

4.2 Influence function of the Wald-Type Test Statistics

We will now study the robustness of the proposed Wald-type test of Section 3 through the influence function of the corresponding test statistics W_n defined in Definition 5. Ignoring the multiplier n, let us define the associated statistical functional for the test statistics W_n evaluated at any joint distribution G as given by

$$W_{\lambda}(G) = \left(\boldsymbol{M}^{T}\boldsymbol{T}_{\lambda}(G) - \boldsymbol{m}\right)^{T} \left(\boldsymbol{M}^{T}\boldsymbol{\Sigma}_{\lambda}(\widehat{\boldsymbol{\beta}}_{\lambda})\boldsymbol{M}\right)^{-1} \left(\boldsymbol{M}^{T}\boldsymbol{T}_{\lambda}(G) - \boldsymbol{m}\right).$$
(23)

Now, considering the ε -contaminated joint distribution $G_{\varepsilon} = (1 - \varepsilon)G + \varepsilon \wedge w$ with respect to the point mass contamination distribution \wedge_w at the contamination point $w = (x_t, y_t)$, the influence function of $W_{\lambda}(\cdot)$ is defined as

$$\mathcal{IF}(\boldsymbol{w}, W_{\lambda}, G) = \frac{\partial W_{\lambda}(G_{\varepsilon})}{\partial \varepsilon} \Big|_{\varepsilon=0}$$

= $\left(\boldsymbol{M}^{T} T_{\lambda}(G) - \boldsymbol{m}\right)^{T} \left(\boldsymbol{M}^{T} \boldsymbol{\Sigma}_{\lambda}\left(\boldsymbol{\beta}_{0}\right) \boldsymbol{M}\right)^{-1} \boldsymbol{M}^{T} \mathcal{IF}(\boldsymbol{w}, T_{\lambda}, G).$

Now, assuming the null hypothesis to be true, let G_0 denote the joint model distribution with true parameter value β_0 satisfying $M^T \beta_0 = m$. Then, under G_0 , we have $T_{\lambda}(G_0) = \beta_0$ and hence $\mathcal{IF}(\boldsymbol{w}, W_{\lambda}, G_0) = \mathbf{0}$. Therefore, the first order influence function analysis is not adequate to quantify the robustness of the proposed Wald-type test statistics W_{λ} . It is bounded in the contamination points $\boldsymbol{w} = (\boldsymbol{x}_t, y_t)$ for all $\lambda \geq 0$ but does not necessarily imply the robustness of the tests since it includes the well-known non-robust MLE based Wald-test at $\lambda = 0$. This fact is consistent with the robustness analysis of different other Wald-type tests under different set-ups (See, for example, Rousseeuw and Ronchetti, 1979; Toma and Broniatowski, 2011; Ghosh et al., 2016 etc.) and we need to consider the second order influence analysis to asses the robustness of W_{λ} .

The second order influence function of the Wald-type test statistics W_n at the joint distribution G is defined as

$$\begin{split} \mathcal{IF}_{2}(\boldsymbol{w}, W_{\lambda}, G) &= \left. \frac{\partial^{2} W_{\lambda}(G_{\varepsilon})}{\partial \varepsilon^{2}} \right|_{\varepsilon=0} \\ &= \left(\boldsymbol{M}^{T} \boldsymbol{T}_{\lambda}(G) - \boldsymbol{m} \right)^{T} \left(\boldsymbol{M}^{T} \boldsymbol{\Sigma}_{\lambda} \left(\boldsymbol{\beta} \right) \boldsymbol{M} \right)^{-1} \boldsymbol{M}^{T} \mathcal{IF}_{2}(\boldsymbol{w}, \boldsymbol{T}_{\lambda}, G) \\ &+ \mathcal{IF}^{T}(\boldsymbol{w}, \boldsymbol{T}_{\lambda}, G) \boldsymbol{M} \left(\boldsymbol{M}^{T} \boldsymbol{\Sigma}_{\lambda} \left(\boldsymbol{\beta} \right) \boldsymbol{M} \right)^{-1} \boldsymbol{M}^{T} \mathcal{IF}(\boldsymbol{w}, \boldsymbol{T}_{\lambda}, G). \end{split}$$

Again, under the null hypothesis H_0 with β_0 being the corresponding true parameter value, this second order influence function simplifies further as presented in the following theorem and yields the possibility to study the robustness of our proposed tests through its boundedness.

Theorem 14 The second order influence function of the proposed Wald-type test statistics W_n , given in Definition 5, at the null model distribution G_0 having true parameter value β_0 is given by

$$\begin{split} \mathcal{IF}_{2}(\boldsymbol{w}, W_{\lambda}, G_{0}) \\ &= \mathcal{IF}^{T}(\boldsymbol{w}, \boldsymbol{T}_{\lambda}, G_{0}) \boldsymbol{M} \left(\boldsymbol{M}^{T} \boldsymbol{\Sigma}_{\lambda} \left(\boldsymbol{\beta}_{0} \right) \boldsymbol{M} \right)^{-1} \boldsymbol{M}^{T} \mathcal{IF}(\boldsymbol{w}, \boldsymbol{T}_{\lambda}, G_{0}). \\ &= \widetilde{\Psi}_{\lambda}^{2}(\boldsymbol{x}_{t}^{T} \boldsymbol{\beta}_{0}, y_{t}) \boldsymbol{x}_{t}^{T} \boldsymbol{J}_{\lambda}^{-1}(\boldsymbol{\beta}_{0}) \boldsymbol{M} \left(\boldsymbol{M}^{T} \boldsymbol{\Sigma}_{\lambda} \left(\boldsymbol{\beta}_{0} \right) \boldsymbol{M} \right)^{-1} \boldsymbol{M}^{T} \boldsymbol{J}_{\lambda}^{-1}(\boldsymbol{\beta}_{0}) \boldsymbol{x}_{t}. \end{split}$$

Note that, the influence function of the Wald-type test statistic is directly a quadratic function of the corresponding MDPDE used. Hence, as described in the previous subsection, the influence function for the proposed tests with $\lambda > 0$ will be small and bounded for all kinds of outliers in a logistic regression model, whereas the classical MLE based Wald-type test will have an unbounded influence function for large "bad" leverage points. Figure 3 shows the plots of this second order influence functions for the Wald-type test statistics for different λ for testing the significance of the first slope parameter in a logistic regression model with with two independent standard normal covariates and $\beta_0 = (0, 1, 1)^T$ fixing $y_t = 0$. The behavior of the influence functions are again similar to those observed for the corresponding MDPDE in Figure 3, which shows the greater robustness of our proposal at larger positive λ over the non-robust MLE based Wald test at $\lambda = 0$.

4.3 Level and Power Influence Functions

We now study the robustness of the proposed tests through the stability of their Type-I and Type-II error which are two basic components for measuring the performance of any testing procedure. In particular, we will study eth local stability of level and power of the proposed tests through corresponding influence function analysis. Note that the finite sample level and power of our proposed Wald-type tests are difficult to compute and has no general form; on the other hand, the tests are consistent having asymptotic power as one against any fixed alternative. So, we will study the influence function of the asymptotic level under the null $\beta = \beta_0$ and asymptotic power under the sequence of contiguous alternatives $\beta_n = \beta_0 + n^{-1/2} d$ as defined in, for example, Hampel et al. (1986) and Ghosh et al. (2016) among others. In particular, assuming the contamination proportion tends to zero at the same rate as the contiguous alternatives approaches to the null, here we consider the following contaminated joint distribution for the power stability calculation as

$$G_{n,\varepsilon,\boldsymbol{w}}^{P} = (1 - \frac{\varepsilon}{\sqrt{n}})G_{\boldsymbol{\beta}_{n}} + \frac{\varepsilon}{\sqrt{n}}\wedge_{\boldsymbol{w}},$$
(24)

where \boldsymbol{w} denote the contamination point $\boldsymbol{w} = (\boldsymbol{x}_t^T, y_t)^T$, and $G_{\boldsymbol{\beta}_n}$ denote the joint model distribution with true parameter value $\boldsymbol{\beta} = \boldsymbol{\beta}_n$. The contamination distribution to be considered for the level stability check can be obtained by substituting $\boldsymbol{d} = \boldsymbol{0}$ in (24), which yields

$$G_{n,\varepsilon,\boldsymbol{w}}^P = (1 - \frac{\varepsilon}{\sqrt{n}})G_{\boldsymbol{\beta}_0} + \frac{\varepsilon}{\sqrt{n}} \wedge_{\boldsymbol{w}}.$$

Then, the level and power influence functions are defined in terms of the following quantities

$$\alpha(\varepsilon, \boldsymbol{w}) = \lim_{n \to \infty} P_{G_{n,\varepsilon,\boldsymbol{w}}^{L}}(W_n > \chi_{r,\alpha}^2),$$

and

$$\pi(\boldsymbol{\beta}_n, \varepsilon, \boldsymbol{x}) = \lim_{n \to \infty} P_{G_{n,\varepsilon,\boldsymbol{w}}^P}(W_n > \chi_{r,\alpha}^2).$$



Figure 3: Second order Influence function of the Wald-type test statistics for testing significance of the first slope parameter β_1 for different λ ($y_t = 0$).

Definition 15 The level influence function (LIF) and the power influence function (PIF) for the Wald-type test statistics W_n are defined respectively as

$$\mathcal{LIF}(\boldsymbol{w}; W_n, G_{\boldsymbol{\beta}_0}) = \left. \frac{\partial}{\partial \varepsilon} \alpha(\varepsilon, \boldsymbol{w}) \right|_{\varepsilon=0}, \quad \mathcal{PIF}(\boldsymbol{x}; W_n, G_{\boldsymbol{\beta}_0}) = \left. \frac{\partial}{\partial \varepsilon} \pi(\boldsymbol{\beta}_n, \varepsilon, \boldsymbol{w}) \right|_{\varepsilon=0}$$

See Ghosh et al. (2016) for an extensive discussion on the interpretations of the level and power influence functions and their relations with the influence function of the test statistics in the context of a general Wald-type test.

Next, we will derive the forms of the LIF and PIF for our proposed tests in logistic regression model assuming the conditions required for the derivation of asymptotic distributions of the MDPDE hold.

Theorem 16 Assume that the conditions of Theorem 6 holds and consider the contiguous alternatives $\beta_n = \beta_0 + n^{-1/2} d$ along with the contaminated model in (24). Then we have the following results:

(i) The asymptotic distribution of the test statistics W_n under $G_{n,\varepsilon,\boldsymbol{w}}^P$ is non-central chi-square with r degrees of freedom and the non-centrality parameter

$$\delta = \widetilde{\boldsymbol{d}}_{arepsilon, \boldsymbol{w}, \lambda}^T (oldsymbol{eta}_0) \boldsymbol{M} \left(\boldsymbol{M}^T \boldsymbol{\Sigma}_{\lambda} \left(oldsymbol{eta}_0
ight) \boldsymbol{M}
ight)^{-1} \boldsymbol{M}^T \widetilde{\boldsymbol{d}}_{arepsilon, \boldsymbol{w}, \lambda} (oldsymbol{eta}_0),$$

where $\widetilde{d}_{\varepsilon, w, \lambda}(\beta_0) = d + \varepsilon \mathcal{IF}(w, T_{\lambda}, G_{\beta_0}).$

(ii) The asymptotic power under $G_{n,\varepsilon,w}^P$ can be approximated as

$$\pi(\boldsymbol{\beta}_{n},\varepsilon,\boldsymbol{w}) \cong P\left(\chi_{r}^{2}(\delta) > \chi_{r,\alpha}^{2}\right)$$
$$\cong \sum_{v=0}^{\infty} C_{v}\left(\boldsymbol{M}^{T}\widetilde{\boldsymbol{d}}_{\varepsilon,\boldsymbol{w},\lambda}(\boldsymbol{\beta}_{0}), \left(\boldsymbol{M}^{T}\boldsymbol{\Sigma}_{\lambda}\left(\boldsymbol{\beta}_{0}\right)\boldsymbol{M}\right)^{-1}\right) P\left(\chi_{r+2v}^{2} > \chi_{r,\alpha}^{2}\right), \quad (25)$$

where

$$C_{v}\left(\boldsymbol{t},\boldsymbol{A}\right) = \frac{\left(\boldsymbol{t}^{T}\boldsymbol{A}\boldsymbol{t}\right)^{v}}{v!2^{v}}e^{-\frac{1}{2}\boldsymbol{t}^{T}\boldsymbol{A}\boldsymbol{t}}$$

 $\chi_p^2(\delta)$ denotes a non-central chi-square random variable with p degrees of freedom and δ as non-centrality parameter and $\chi_q^2 = \chi_q^2(0)$ denotes a central chi-square random variable having degrees of freedom q.

Proof. Let us denote $\boldsymbol{\beta}_n^* = \boldsymbol{T}_{\lambda}(G_{n,\varepsilon,\boldsymbol{w}}^P)$. Then, we get

$$W_{n} = n(\boldsymbol{M}^{T}\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{m})^{T} \left(\boldsymbol{M}^{T}\boldsymbol{\Sigma}_{\lambda}\left(\boldsymbol{\beta}_{0}\right)\boldsymbol{M}\right)^{-1} \left(\boldsymbol{M}^{T}\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{m}\right)$$

$$= n\left(\boldsymbol{M}^{T}\boldsymbol{\beta}_{n}^{*} - \boldsymbol{m}\right)^{T} \left(\boldsymbol{M}^{T}\boldsymbol{\Sigma}_{\lambda}\left(\boldsymbol{\beta}_{0}\right)\boldsymbol{M}\right)^{-1} \left(\boldsymbol{M}^{T}\boldsymbol{\beta}_{n}^{*} - \boldsymbol{m}\right)$$

$$+ n(\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{\beta}_{n}^{*})^{T}\boldsymbol{M} \left(\boldsymbol{M}^{T}\boldsymbol{\Sigma}_{\lambda}\left(\boldsymbol{\beta}_{0}\right)\boldsymbol{M}\right)^{-1} \boldsymbol{M}^{T}(\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{\beta}_{n}^{*})$$

$$+ n(\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{\beta}_{n}^{*})^{T}\boldsymbol{M} \left(\boldsymbol{M}^{T}\boldsymbol{\Sigma}_{\lambda}\left(\boldsymbol{\beta}_{0}\right)\boldsymbol{M}\right)^{-1} \left(\boldsymbol{M}^{T}\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{m}\right)$$

$$= S_{1,n} + S_{2,n} + S_{3,n}.$$
 (26)

Next, one can show that

$$\sqrt{n}(\boldsymbol{\beta}_{n}^{*}-\boldsymbol{\beta}_{0}) = \boldsymbol{d} + \varepsilon \mathcal{IF}\left(\boldsymbol{w}, \boldsymbol{T}_{\lambda}, \boldsymbol{G}_{\boldsymbol{\beta}_{0}}\right) + o_{p}(\boldsymbol{1}_{p}) \\
= \widetilde{\boldsymbol{d}}_{\varepsilon,\boldsymbol{w},\lambda}(\boldsymbol{\theta}_{0}) + o_{p}(\boldsymbol{1}_{p}).$$
(27)

Thus, we get

$$\sqrt{n}(\boldsymbol{M}^{T}\boldsymbol{\beta}_{n}^{*}-\boldsymbol{m}) = \boldsymbol{M}^{T}\widetilde{\boldsymbol{d}}_{\varepsilon,\boldsymbol{w},\lambda}(\boldsymbol{\theta}_{0}) + o_{p}(\boldsymbol{1}_{p}).$$
(28)

Further, under $G_{n,\varepsilon,w}^P$, the asymptotic distribution of MDPDE yields

$$\sqrt{n}(\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{\beta}_{n}^{*}) \xrightarrow[n \to \infty]{\mathcal{L}} \mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma}_{\lambda}(\boldsymbol{\beta}_{0})).$$
(29)

Thus, we get

$$S_{3,n} \xrightarrow[n \to \infty]{\mathcal{L}} \chi_r^2.$$

Combining (26), (28) and (29), we get

$$W_n = \boldsymbol{Z}_n^T \left(\boldsymbol{M}^T \boldsymbol{\Sigma}_{\lambda} \left(\boldsymbol{\beta}_0 \right) \boldsymbol{M} \right)^{-1} \boldsymbol{Z}_n + o_p(1),$$

where

$$\boldsymbol{Z}_n = \sqrt{n} \boldsymbol{M}^T (\widehat{\boldsymbol{\beta}}_{\lambda} - \boldsymbol{\beta}_n^*) + \boldsymbol{M}^T \widetilde{\boldsymbol{d}}_{\varepsilon, \boldsymbol{w}, \lambda}(\boldsymbol{\theta}_0).$$

By (29),

$$oldsymbol{Z}_n \stackrel{\mathcal{L}}{\underset{n o \infty}{\longrightarrow}} \mathcal{N}\left(oldsymbol{M}^T \widetilde{oldsymbol{d}}_{arepsilon,oldsymbol{w},\lambda}(oldsymbol{ heta}_0), oldsymbol{M}^T oldsymbol{\Sigma}_\lambda\left(oldsymbol{eta}_0
ight) oldsymbol{M}
ight),$$

and hence we get that

$$W_n \xrightarrow[n \to \infty]{\mathcal{L}} \chi_r^2(\delta),$$

where δ is as defined in Part (i) of the theorem.

Part (ii) of the theorem follows from Part (i) using the infinite series expansion of a non-central distribution function in terms of that of the central chi-square variables:

$$\pi(\boldsymbol{\beta}_{n},\varepsilon,\boldsymbol{w}) = \lim_{n \to \infty} P_{G_{n,\varepsilon,\boldsymbol{w}}^{P}}(W_{n} > \chi_{r,\alpha}^{2}) \cong P(\chi_{r,\delta}^{2} > \chi_{r,\alpha}^{2})$$
$$= \sum_{v=0}^{\infty} C_{v} \left(\boldsymbol{M}^{T} \widetilde{\boldsymbol{d}}_{\varepsilon,\boldsymbol{w},\lambda}(\boldsymbol{\beta}_{0}), \left(\boldsymbol{M}^{T} \boldsymbol{\Sigma}_{\lambda}\left(\boldsymbol{\beta}_{0}\right) \boldsymbol{M}\right)^{-1}\right) P\left(\chi_{r+2v}^{2} > \chi_{r,\alpha}^{2}\right).$$

Corollary 17 Putting $\varepsilon = 0$ in Theorem 16, we get the asymptotic power of the proposed Wald-type tests under the contiguous alternative hypotheses $\beta_n = \beta_0 + n^{-1/2} d$ as

$$\pi(\boldsymbol{\beta}_n) = \pi(\boldsymbol{\beta}_n, 0, \boldsymbol{w}) \cong \sum_{v=0}^{\infty} C_v \left(\boldsymbol{M}^T \boldsymbol{d}, \left(\boldsymbol{M}^T \boldsymbol{\Sigma}_{\lambda}(\boldsymbol{\beta}_0) \, \boldsymbol{M} \right)^{-1} \right) P \left(\chi_{r+2v}^2 > \chi_{r,\alpha}^2 \right).$$

This is identical with the results obtained earlier in Theorem 10 independently.

Corollary 18 Putting d = 0 in Theorem 16, we get the asymptotic distribution of W_n under $G_{n,\varepsilon,w}^L$ as the non-central chi-square distribution having r degrees of freedom and non-centrality parameter

$$arepsilon^2 \mathcal{IF}(oldsymbol{w};oldsymbol{T}_{\lambda},G_{oldsymbol{eta}_0})^Toldsymbol{M}\left(oldsymbol{M}^Toldsymbol{\Sigma}_{\lambda}\left(oldsymbol{eta}_0
ight)oldsymbol{M}
ight)^{-1}oldsymbol{M}^T\mathcal{IF}(oldsymbol{w};oldsymbol{T}_{\lambda},G_{oldsymbol{eta}_0}).$$

Then, the asymptotic level under contiguous contamination is given by

$$\begin{aligned} \alpha(\varepsilon, \boldsymbol{w}) &= \pi(\boldsymbol{\beta}_0, \varepsilon, \boldsymbol{w}) \\ &\cong \sum_{v=0}^{\infty} C_v \left(\varepsilon \boldsymbol{M}^T \mathcal{IF}(\boldsymbol{w}; \boldsymbol{T}_{\lambda}, \boldsymbol{G}_{\boldsymbol{\beta}_0}), \left(\boldsymbol{M}^T \boldsymbol{\Sigma}_{\lambda} \left(\boldsymbol{\beta}_0 \right) \boldsymbol{M} \right)^{-1} \right) P \left(\chi_{r+2v}^2 > \chi_{r, \alpha}^2 \right). \end{aligned}$$

In particular, as $\varepsilon \to 0, \beta_n^* \to \beta_0$ and the non-centrality parameter of the above asymptotic distribution tends to zero leading to the null distribution of W_n .

Now we can easily obtain the power and level influence functions of the Wald-type test statistics from Theorem 16 and Corollary 18 and these have been presented in the following theorem.

Theorem 19 Under the assumptions of Theorem 16, the power and level influence functions of the proposed Wald-type test statistic W_n is given by

$$\mathcal{PIF}(\boldsymbol{w}, W_n, G_{\boldsymbol{\beta}_0}) \cong K_r^* \left(\boldsymbol{s}^T \left(\boldsymbol{\beta}_0 \right) \mathbf{d} \right) \boldsymbol{s}^T (\boldsymbol{\beta}_0) \mathcal{IF}(\boldsymbol{w}, \boldsymbol{T}_\lambda, G_{\boldsymbol{\beta}_0}), \tag{30}$$

with $\boldsymbol{s}^{T}(\boldsymbol{\beta}_{0}) = \boldsymbol{d}^{T}\boldsymbol{M}\left(\boldsymbol{M}^{T}\boldsymbol{\Sigma}_{\lambda}\left(\boldsymbol{\beta}_{0}\right)\boldsymbol{M}\right)^{-1}\boldsymbol{M}^{T}$ and

$$K_r^*(s) = e^{-\frac{s}{2}} \sum_{v=0}^{\infty} \frac{s^{v-1}}{v! 2^v} \left(2v - s\right) P\left(\chi_{r+2v}^2 > \chi_{r,\alpha}^2\right),$$

and

$$\mathcal{LIF}(\boldsymbol{w}, W_n, G_{\boldsymbol{\beta}_0}) = 0.$$

Further, the derivative of $\alpha(\varepsilon, \boldsymbol{w})$ of any order with respect to ε will be zero at $\varepsilon = 0$, implying that the level influence function of any order will be zero.

Proof. We start with the expression of $\pi(\boldsymbol{\beta}_n, \varepsilon, \boldsymbol{w})$ from Theorem 16. Clearly, by definition of PIF and using the chain rule of derivatives, we get

$$\begin{aligned} \mathcal{PIF}(\boldsymbol{w}, W_n, G_{\boldsymbol{\beta}_0}) &= \frac{\partial}{\partial \varepsilon} \left. \pi(\boldsymbol{\beta}_n, \varepsilon, \boldsymbol{w}) \right|_{\varepsilon=0} \\ &\cong \sum_{v=0}^{\infty} \frac{\partial}{\partial \varepsilon} \left. C_v \left(\boldsymbol{M}^T \widetilde{\boldsymbol{d}}_{\varepsilon, \boldsymbol{w}, \lambda}(\boldsymbol{\beta}_0), \left(\boldsymbol{M}^T \boldsymbol{\Sigma}_{\lambda} \left(\boldsymbol{\beta}_0 \right) \boldsymbol{M} \right)^{-1} \right) \right|_{\varepsilon=0} P \left(\chi_{r+2v}^2 > \chi_{r, \alpha}^2 \right) \\ &\cong \sum_{v=0}^{\infty} \frac{\partial}{\partial t^T} \left. C_v \left(\boldsymbol{M}^T \boldsymbol{t}, \left(\boldsymbol{M}^T \boldsymbol{\Sigma}_{\lambda} \left(\boldsymbol{\beta}_0 \right) \boldsymbol{M} \right)^{-1} \right) \right|_{\boldsymbol{t}=\widetilde{\boldsymbol{d}}_{0, \boldsymbol{w}, \lambda}(\boldsymbol{\beta}_0)} \left. \frac{\partial}{\partial \varepsilon} \left. \widetilde{\boldsymbol{d}}_{\varepsilon, \boldsymbol{w}, \lambda}(\boldsymbol{\beta}_0) \right|_{\varepsilon=0} P \left(\chi_{r+2v}^2 > \chi_{r, \alpha}^2 \right). \end{aligned}$$

Now $d_{0,w,\lambda}(\boldsymbol{\beta}_0) = d$ and standard differentiations give

$$\frac{\partial}{\partial \varepsilon} \widetilde{\boldsymbol{d}}_{\varepsilon, \boldsymbol{w}, \lambda}(\boldsymbol{\beta}_0) = \mathcal{IF}(\boldsymbol{w}, \boldsymbol{T}_{\lambda}, \boldsymbol{G}_{\boldsymbol{\beta}_0}),$$

and

$$rac{\partial}{\partial t}C_{v}\left(t,oldsymbol{A}
ight)=rac{\left(t^{T}oldsymbol{A}t
ight)^{v-1}}{v!2^{v}}\left(2v-t^{T}oldsymbol{A}t
ight)oldsymbol{A}te^{-rac{1}{2}t^{T}oldsymbol{A}t}.$$

Combining above results and simplifying, we get the required expression of PIF as presented in the theorem. \blacksquare

It is clear from the above theorem that, the asymptotic level of the proposed Wald-type test statistic will be unaffected by a contiguous contamination for any values of the tuning parameter λ , whereas the power influence function will be bounded whenever the influence function of the MDPDE is bounded (which happens for all $\lambda > 0$). Thus, the robustness of the power of the proposed tests again turns out to be directly dependent on the robustness of the MDPDE β_{λ} used in constructing the test. In particular, the asymptotic contiguous power of the classical MLE based Wald-type test (at $\lambda = 0$) will be non-robust whereas that for the Wald-type tests with $\lambda > 0$ will be robust under contiguous contaminations and this robustness increases as λ increases further.



Figure 4: (a) Simulated levels of different tests for pure data; (b) simulated levels of different tests for contaminated data; (c) simulated powers of different tests for pure data; (d) simulated powers of different tests for contaminated data.

5 Simulation study

In this section we have empirically demonstrated some of the strong robustness properties of the density power divergence tests for the logistic regression model. We considered two explanatory variables x_1 and x_2 in this study, so k = 2. These two variables are distributed according a standard normal distribution $\mathcal{N}(\mathbf{0}, \mathbf{I}_{2\times 2})$. The response variables Y_i are generated following the logit model as given in (1). The true value of the parameter is taken as $\boldsymbol{\beta}_0 = (0, 1, 1)^T$. We considered the null hypothesis $H_0: (\beta_1, \beta_2)^T = (1, 1)^T$. It can be written in the form of the general hypothesis given in (3), where $\boldsymbol{m} = (1, 1)^T$ and

$$\boldsymbol{M} = \left(\begin{array}{cc} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{array} \right).$$

Our interest was in studying the observed level (measured as the proportion of test statistics exceeding the corresponding chi-square critical value in a large number – here 1000 – of replications) of the test under the correct null hypothesis. The result is given in Figure 4(a) where the sample size n varies from 20 to 100. We have used several Wald-type test statistics, corresponding to different minimum density power divergnece estimators. We have used, $\lambda = 0$, 0.1, 0.5 and 1, in this particular study. As it is previously mentioned, $\lambda = 0$ is the classical Wald test for the logistic regression model. The horizontal lines in the figure represents the nominal level of 0.05. It may be noticed that all the tests are slightly conservative for small sample sizes and lead to somewhat deflated observed levels. In particular, the Wald-type tests with higher values of λ are relatively more conservative. However, this discrepancy decreases rapidly as sample size increases.

To evaluate the stability of the level of the tests under contamination, we repeated the tests for the same null hypothesis by adding 3% outliers in the data. For the outlying observations we first introduced the leverage points where x_1 and x_2 are generated from $\mathcal{N}(\boldsymbol{\mu}_c, \sigma \boldsymbol{I}_{2\times 2})$ with $\boldsymbol{\mu}_c = (5, 5)^T$ and $\sigma = 0.01$. Then the values of the response variable corresponding to those leverage points were altered to produce vertical outliers ($y_t = 1$ was converted to $y_t = 0$). Figure 4(b) shows that the levels of the classical Wald test as well as DPD(0.1) test break down, whereas Wald-type test statistics for $\lambda = 0.5$ and $\lambda = 1$ present highly stable levels.

To investigate the power of the tests we changed the null hypothesis to $H_0^* : (\beta_1, \beta_2)^T = (0, 0)^T$, and kept the data generating distributions as before, as well as the true value of the parameter as $\beta_0 = (0, 1, 1)^T$. In terms of the null hypothesis in (3) the value of \boldsymbol{m} is changed to $(0, 0)^T$ whereas \boldsymbol{M} remained unchanged from the previous experiment. The empirical power functions are calculated in the same manner as the levels of the tests, and plotted in Figure 4(c). The Wald test is the most powerful under pure data. The power of the Wald-type test statistic for $\lambda = 0.1$ almost coincide with the classical Wald test in this case. The performances of the Wald-type test statisdtics for $\lambda = 0.5$ and $\lambda = 1$ are relatively poor, however, as the sample size increases to 60 and beyond, the powers are practically identical.

Finally, we calculated the power functions under contamination for the above hypothesis under the same setup as of the level contamination. The observed powers of that the tests are given in Figure 4(d). The Wald-type test statistics for $\lambda = 0.5$ and $\lambda = 1$ show stable powers under contamination, but the classical Wald test and the Wald-type test for $\lambda = 0.1$ exhibit a drastic loss in power. In very small sample sizes the classical Wald test and the Wald-type test for $\lambda = 0.1$ have slightly higher power than the other tests, but this must be a consequence of the observed levels of these tests being higher than the latter for such sample sizes. On the whole, the proposed Wald-type test statistics corresponding to moderately large λ appear to be quite competitive to the classical Wald test for pure normal data, but they are far better in terms of robustness properties under contaminated data.

6 Real Data Examples

In this section we will explore the performance of the proposed Wald-type tests in logistic regression models by applying it on different interesting real data sets. The estimators are computed by minimizing the corresponding density power divergence through the software R, and the minimization is performed using "optim" function.

6.1 Students Data

As an interesting data example leading to the logistic regression model, we consider the students data set from Muñoz-Garcia et al. (2006). The data set consists of 576 students of the University of Seville. The response variable is the students aim to graduate after three years. The explanatory variables are gender $(x_{i1} = 0 \text{ if male}; x_{i1} = 1 \text{ if female})$, entrance examination (EE) in University $(x_{i2} = 1 \text{ if the first time}; x_{i2} = 0 \text{ otherwise})$ and sum of marks (x_{i3}) obtained for the courses of first term. There were 61 distinct cases (i.e. n = 61) in this study. We assume that the response variable follows a binomial logistic regression model as mentioned in Remark 4. We are interested to test the null hypothesis that the gender of student does not play any role on their aim. So the null hypothesis is given by $H_0: \beta_1 = 0$. Figure 5 shows *p*-values of Wald-type tests for different values of λ . Muñoz-Garcia et al. (2006) mentioned that 32nd observation is the most influential point as it has a large residual and a high leverage value. If we use the classical Wald test or Wald-type tests with small λ under the full data, the null hypothesis is rejected at 10% level of significance. But this result is clearly a false positive as the outlie

with large λ give rol



Figure 5: P-values of Wald-type tests for $^{\lambda}$ testing $H_0: \beta_1 = 0$ in Students data.

6.2 Lymphatic Cancer Data

Brown (1980), Martín and Pardo (2009) and Zelterman (2005, Section 3.3) studied the data that focused on the evidence of lymphatic cancer in prostate cancer patients for predicting lymph nodal involvement of cancer. There were five covariates (three dichotomous and two continuous): the X-ray finding ($x_{i1} = 1$ if present; $x_{i1} = 0$ if absent), size of the tumor by palpation ($x_{i2} = 1$ if serious; $x_{i2} = 0$ if not serious), pathology grade by biopsy ($x_{i3} = 1$ if serious; $x_{i3} = 0$ if not serious), the age of the patient at the time of diagnosis (x_{i4}) and serum acid phosphatase level (x_{i5}). The diagnostics was associated with 53 individuals. An ordinary logistic model is assumed here. We are interested to test the significance of the size of the tumor on the response variable, so the null hypothesis is taken as $H_0: \beta_2 = 0$. The *p*-values of Wald-type tests for different values of λ are given in Figure 6. Martín and Pardo (2009) noticed that the 24th observation is an influential point. The *p*-value of the classical Wald test under the full data is 0.0430, but if the outlier is deleted it becomes 0.0668. So if we consider a test at outlying observation en we delete just one duce high *p*-values.

regibon



Figure 6: P-values of Wald-type tests for testing $H_0: \beta_2 = 0$ in Lymphatic Cancer data.

6.3 Vasoconstriction Data

Finney (1947), Pregibon (1981) and Martín and Pardo (2009) studied the data where the interest is on the occurrence of vasoconstriction in the skin of the finger. The covariates of the study were the

logarithm of volume (x_{i1}) and the (1981) has shown that two observes they have large residuals. Howeves the y-space and are not leverage period to be the null hypothesis is given by the full data is 0.0194, and in the Wald-type tests with large λ processing the second secon



Figure 7: P-values of Wald-type tests for testing $H_0: \beta_1 = \beta_2 = 0$ in Vasoconstriction data.

6.4 Leukemia Data

The data set consists of 33 cases on the survival of individuals diagnosed with leukemia. The explanatory variables are white blood cell count (x_{i1}) and another variable which indicates the presence or absence of a certain morphologic characteristic in the white cells $(x_{i2} = 1 \text{ if present}; x_{i2} = 0 \text{ if absent})$. This data set was also studied by Cook and Weisberg (1982), Johnson (1985) and Martín and Pardo (2009). They defined a success to be patient survival in excess of 52 weeks. We are interested to test the significance of two covariates, i.e. the null hypothesis is $H_0: \beta_1 = \beta_2 = 0$. The plot of the *p*-values of Wald-type tests for different values of λ is given in Figure 8. Martín and Pardo (2009) noticed that the 15th observation is an influential point. The *p*-value of the classical Wald test under the full data is 0.0226, but if the outlier is deleted it becomes 0.0683. Thus, at 5% level of significance, the decision of the test α Vald-type tests with

high values of λ alwa



Figure 8: P-values of Wald-type tests for testing $H_0: \beta_1 = \beta_2 = 0$ in Leukemia data.

7 Concluding Remarks

Logistic regression for binary outcomes is one of the most popular and successful tools in the statisticians toolbox. It is frequently used by applied scientists of many disciplines to solve problems of real interest in their doman of application. However, in the present age of big data, the need for protection against data contamination and other modelling errors is paramount, and, wherever possible, strong robustness qualities should be a default requirement for statistical methods used in practice. In this paper we have presented one such class of inference procedures. We have provided a thorough theoretical evaluation of the proposted class of tests for testing the linear hypothesis in the logistic regression model highlighting their robustness advantages. We have also produced substantial numerical evidence, including simulation results and a large number of real problems, to demonstrate how these theoretical advantages translate in practice to real gains. On the whole, we feel that the proposed tests will turn out to be an useful method with significant practical application.

Acknowledgement. This research is partially supported by Grants MTM2015-67057 and ECO2015-66593-P from Ministerio de Economia y Competitividad (Spain).

References

- Basu, A., Harris, I. R., Hjort, N. L. and Jones, M. C. (1998). Robust and efficient estimation by minimizing a density power divergence. *Biometrika*, 85, 549–559.
- [2] Basu, A., Shioya, H. and Park, C. (2011). The minimum distance approach. Monographs on Statistics and Applied Probability. CRC Press, Boca Raton.
- [3] Basu, A., Mandal, A., Martín, N. and Pardo, L. (2013). Testing statistical hypotheses based on the density power divergence. Annals of the Institute of Statistical Mathematics, 65, 319–348.
- [4] Basu, A., Mandal, A., Martín, N. and Pardo, L. (2015). Robust tests for the equality of two normal means based on the density power divergence. *Metrika*, 78, 611–634.
- [5] Basu, A., Mandal, A., Martín, N., Pardo, L. (2016). Generalized Wald-type tests based on minimum density power divergence estimators. *Statistics*, 50, 1–26.

- [6] Bianco, A. M. and Martinez, E. (2009). Robust testing in the logistic regression model. Computational Statistics and Data Analysis, 53, 4095–4105.
- [7] Bianco, A. M., and Yohai, V. J. (1996). Robust Estimation in the Logistic Regression Model, in Robust Statistics, Data Analysis, and Computer Intensive Methods, 17–34; Lecture Notes in Statistics 109, Springer Verlag, Ed. H. Rieder. New York
- [8] Bondell, H. D. (2005). Minimum distance estimation for the logistic regression model. *Biometrika*, 92, 724-731.
- [9] Bondell, H. D. (2008). A characteristic function approach to the biased sampling model, with application to robust logistic regression. *Journal of Statistical Planning and Inference*, 138, 742-755.
- Brown, B.W. (1980). Prediction analysis for binary data, in *Biostatistics Casebook*, R.G. Miller,
 B. Efron, B.W. Brown and L.E. Moses, eds., John Wiley and Sons, New York, pp. 3–18.
- [11] Carroll, R. J. and Pederson, S. (1993). On Robustness in the logistic regression model. Journal of the Royal Statistical Society: Series B, 55, 669-706.
- [12] Croux, C. and Haesbroeck, G. (2003). Implementing the Bianco and Yohai estimator for logistic regression. *Computational Statistics and Data Analysis*, 44, 273–295.
- [13] Christmann, A. (1994). Least Median of Weighted Squares in Logistic Regression with Large Strata. *Biometrika*, 81, 413-417.
- [14] Christmann, A. and Rousseeuw, P.J. (2001). Measuring overlap in binary regression, Comp. Statistics & Data Analysis, 37, 65-75.
- [15] Cook, R.D. and Weisberg, S. (1982). Residuals and Influence in Regression, Chapman & Hall, London.
- [16] Feigl, P. and Zelen, M. (1965). Estimation of exponential probabilities with concomitant information. *Biometrics*, 21, 826–838.
- [17] Finney, D.J. (1947). The estimation from individual records of the relationship between dose and quantal response. *Biometrika*, 34, 320–334.
- [18] Johnson, W. (1985). Influence measures for logistic regression: Another point of view. *Biometrics*, 72, 59–65.
- [19] Ghosh, A., Basu, A. (2013). Robust Estimation for Independent but Non-Homogeneous Observations using Density Power Divergence with application to Linear Regression. *Electronic Journal* of Statistics, 7, 2420–2456.
- [20] Ghosh, A., Basu, A. and Pardo, L. (2015). On the robustness of a divergence based test of simple statistical hypotheses, *Journal of Statistical Planning and Inference*, 161, 91–108.
- [21] Ghosh, A., Mandal, A., Martín, N. and Pardo, L. (2016). Influence Analysis of Robust Wald-type Tests. Journal of Multivariate Analysis, 147, 102–126.
- [22] Greene, W. H. (2003). Econometric Analysis. Upper Saddle River: Prentice Hall Inc
- [23] Hampel, F. R., Ronchetti, E. M., Rousseeuw, P. J., and Stahel, W. A., (1986). Robust statistics: The approach based on influence functions. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York.

- [24] Hobza, T., Pardo, L. and I. Vajda (2008). Robust median estimator in logistic regression. Journal of Statistical Planning and Inference, 138, 3822-3840.
- [25] Maronna, R. A., Martin, R. D. and Yohai, V. J. (2006). Robust Statistics. Theory and Methods. Wiley Series in Probability and Statistics.
- [26] Martín, N. and Pardo, L. (2009). On the asymptotic distribution of Cook's distance in logistic regression models. *Journal of Applied Statistics*, 36, 1119–1146.
- [27] Muñoz-Garcia, J., Muñoz-Pichardo, J.M. and Pardo, L. (2006). Cressie and Read powerdivergences as influence measures for logistic regression models. *Comput. Statist. Data Anal.*, 50, 3199–3221.
- [28] Morgenthaler, S. (1992), Least-absolute-deviations fits for generalized linear models. *Biometrika*, 79, 747-754.
- [29] Pregibon, D. (1981). Logistic regression diagnostics. Annals of Statistics, 9, 705–724.
- [30] Pregibon, D. (1982), Resistant lits for some commonly used logistic models with medical applications, Biometrics, **38**, 485-498.
- [31] Rousseeuw, P. J. and Christmann, A. (2003), Robustness against separation and outliers in logistic regression. *Computational Statistics and Data Analysis*, **43**, 315–332.
- [32] Rousseeuw, P. J. and Ronchetti, E. (1979) The influence curve for tests. *Research Report* 21, Fachgruppe fur Statistik, ETH Zurich.
- [33] Toma, A. and Broniatowski, M. (2011). Dual divergence estimators and tests: Robustness results. Journal of Multivariate Analysis 102, 20–36.
- [34] Yohai, V.J. (1987). High breakdown-point and high efficiency robust estimates for regression. Annals Statistics, 15, 692–656.
- [35] Zelterman, D. (2005). Models for Discrete Data. Oxford University Press, New York.