

# Uniformly most powerful unbiased test for conditional independence in Gaussian graphical model

Koldanov Petr <sup>1</sup>, Koldanov Alexander, Kalyagin Valeriy; Pardalos Panos

*National Research University Higher School of Economics, Laboratory of Algorithms and Technologies for Network Analysis, Rodionova 136, Nizhny Novgorod, Russia; Preeminent Professor in Industrial and Systems Engineering, Director, CAO Industrial and Systems Engineering, University of Florida, Gainesville*

---

## Abstract

Model selection for Gaussian concentration graph is based on multiple testing of pairwise conditional independence. In practical applications partial correlation tests are widely used. However it is not known whether partial correlation test is uniformly most powerful for pairwise conditional independence testing. This question is answered in the paper. Uniformly most powerful unbiased test of Neymann structure is obtained. It turns out, that this test can be reduced to usual partial correlation test. It implies that partial correlation test is uniformly most powerful unbiased one.

*Keywords:* Conditional independence, Exponential families, Multivariate normal distribution, Sample partial correlation test, Tests of Neyman structure, Uniformly most powerful unbiased tests.

---

## 1. Introduction.

Let  $X = (X_1, X_2, \dots, X_N)$  be a random vector with multivariate Gaussian distribution. Concentration graph is defined as follows: nodes of the graph are associated with random variables  $X_1, X_2, \dots, X_N$ , edge  $(i, j)$  is included in the graph iff random variables  $X_i, X_j$  are conditionally dependent  
 5 [Lauritzen 1996], [Anderson 2003]. Model selection for Gaussian concentration

---

<sup>1</sup>Corresponding author, pkoldanov@hse.ru

graph consists of identification of concentration graph from observations. This problem has a practical importance in biology and genetics [Edwards 2000], [Drton and Perlman 2004]. Common approach for model selection is based on multiple testing of individual hypotheses of pairwise conditional independence [Drton and Perlman 2007], [Drton and Perlman 2008].

Conditional independence of  $X_i, X_j$  given  $X_k, k \in N(i, j) = \{1, 2, \dots, N\} \setminus \{i, j\}$  is equivalent to the equation  $\rho_{i,j \bullet N(i,j)} = 0$ , where  $\rho_{i,j \bullet N(i,j)}$  is the partial correlation of  $X_i$  and  $X_j$  given  $X_k, k \in N(i, j)$ . For testing hypothesis  $\rho_{i,j \bullet N(i,j)} = 0$  for multivariate normal distributions the test of sample partial correlation is largely used [Anderson 2003]. At the same time, as far as we know, there are no results concerning uniformly most powerful unbiased (UMPU) tests for conditional independence. Such test is of own interest and could improve multiple testing procedures for model selection. In the present paper we construct a uniformly most powerful unbiased test of Neyman structure for testing pairwise conditional independence. It turns out that this test can be reduced to the sample partial correlation test. Therefore the sample partial correlation test is uniformly most powerful unbiased one. This fact has some important consequences for multiple testing with additive loss function.

The paper is organized as follows. Section 2 contains basic definition and problem statement. In Section 3 a general description of the tests of Neyman structure is given. In Section 4 the UMPU test for testing pairwise conditional independence is constructed. In Section 5 it is proved that the UMPU test can be reduced to the sample partial correlation test.

## 2. Basic notations and problem statement.

Let random vector  $X = (X_1, X_2, \dots, X_N)$  have a multivariate normal distribution  $N(\mu, \Sigma)$ , where  $\mu = (\mu_1, \mu_2, \dots, \mu_N)$  is the vector of means and  $\Sigma = (\sigma_{i,j})$  is the covariance matrix,  $\sigma_{i,j} = \text{cov}(X_i, X_j), i, j = 1, 2, \dots, N$ . Let  $x(t), t = 1, 2, \dots, n$  be a sample of the size  $n$  from the distribution of  $X$  and

$$s_{i,j} = \frac{1}{n} \sum_{t=1}^n (x_i(t) - \bar{x}_i)(x_j(t) - \bar{x}_j),$$

be the sample covariance between  $X_i, X_j$ , where  $\bar{x}_i = (1/n) \sum_{t=1}^n x_i(t)$ . Denote by  $S = (s_{i,j})$  the matrix of sample covariances.

The inverse matrix for  $\Sigma$ ,  $\Sigma^{-1} = (\sigma^{i,j})$  is known as the concentration or precision matrix for the distribution of  $X$ . For simplicity we use the notation  $\rho^{i,j} = \rho_{i,j \bullet N(i,j)}$ . The problem of pairwise conditional independence testing has the form:

$$h_{i,j} : \rho^{i,j} = 0 \text{ vs } k_{i,j} : \rho^{i,j} \neq 0 \quad (1)$$

According to [Lauritzen 1996] the partial correlation can be calculated as

$$\rho^{i,j} = -\frac{\sigma^{i,j}}{\sqrt{\sigma^{i,i}\sigma^{j,j}}}$$

Therefore the problem of pairwise conditional independence testing (1) can be formulated as

$$h_{i,j} : \sigma^{i,j} = 0, \text{ vs } k_{i,j} : \sigma^{i,j} \neq 0. \quad (2)$$

### 3. Test of Neyman structure.

To construct UMPU test for the problem (2) we use a test of Neyman structure for natural parameters of exponential family [Lehmann and Romano 2005]. Let  $f(x; \theta)$  be the density of the exponential family:

$$f(x; \theta) = c(\theta) \exp \left( \sum_{j=1}^M \theta_j T_j(x) \right) m(x) \quad (3)$$

where  $c(\theta)$  is a function defined in the parameters space,  $m(x), T_j(x)$  are functions defined in the sample space, and  $T_j(X)$  are the sufficient statistics for  $\theta_j, j = 1, \dots, M$ .

Suppose that hypothesis has the form:

$$h_j : \theta_j = \theta_j^0 \text{ vs } k_j : \theta_j \neq \theta_j^0, \quad (4)$$

where  $\theta_j^0$  is fixed.

The UMPU test for hypotheses (4) is (see [Lehmann and Romano 2005], Ch. 4, theorem 4.4.1):

$$\varphi_j = \begin{cases} 0, & \text{if } c'_j(t_1, \dots, t_{j-1}, t_{j+1}, \dots, t_M) < t_j < c''_j(t_1, \dots, t_{j-1}, t_{j+1}, \dots, t_M) \\ 1, & \text{otherwise} \end{cases} \quad (5)$$

50 where  $t_i = T_i(x), i = 1, \dots, M$ . The constants  $c'_j, c''_j$  are defined from the equations

$$\int_{c'_j}^{c''_j} f(t_j; \theta_j^0 | T_i = t_i, i = 1, \dots, M; i \neq j) dt_j = 1 - \alpha \quad (6)$$

and

$$\begin{aligned} & \int_{-\infty}^{c'_j} t_j f(t_j; \theta_j^0 | T_i = t_i, i = 1, \dots, M; i \neq j) dt_j + \\ & + \int_{c''_j}^{+\infty} t_j f(t_j; \theta_j^0 | T_i = t_i, i = 1, \dots, M; i \neq j) dt_j = \\ & = \alpha \int_{-\infty}^{+\infty} t_j f(t_j; \theta_j^0 | T_i = t_i, i = 1, \dots, M; i \neq j) dt_j \end{aligned} \quad (7)$$

where  $f(t_j; \theta_j^0 | T_i = t_i, i = 1, \dots, M; i \neq j)$  is the density of conditional distribution of statistic  $T_j$  given  $T_i = t_i, i = 1, 2, \dots, N, i \neq j$ , and  $\alpha$  is the significance  
55 level of the test.

#### 4. Uniformly most powerful unbiased test for conditional independence.

Now we construct the UMPU test for testing hypothesis of conditional independence (2). Consider statistics

$$S_{k,l} = \frac{1}{n} \sum_{t=1}^n (X_i(t) - \bar{X}_i)(X_j(t) - \bar{X}_j),$$

Joint distribution of statistics  $S_{k,l}, k, l = 1, 2, \dots, N, n > N$  is given by Wishart density function [Anderson 2003]:

$$f(\{s_{k,l}\}) = \frac{[\det(\sigma^{k,l})]^{n/2} \times [\det(s_{k,l})]^{(n-N-2)/2} \times \exp[-(1/2) \sum_k \sum_l s_{k,l} \sigma^{k,l}]}{2^{(Nn/2)} \times \pi^{N(N-1)/4} \times \Gamma(n/2) \Gamma((n-1)/2) \cdots \Gamma((n-N+1)/2)}$$

if the matrix  $S = (s_{k,l})$  is positive definite, and  $f(\{s_{k,l}\}) = 0$  otherwise. It implies that statistics  $S_{k,l}$  are sufficient statistics for natural parameters  $\sigma^{k,l}$ .

Wishart density function can be written as:

$$f(\{s_{k,l}\}) = C(\{\sigma^{k,l}\}) \exp[-\sigma^{i,j} s_{i,j} - \frac{1}{2} \sum_{(k,l) \neq (i,j); (k,l) \neq (j,i)} s_{k,l} \sigma^{k,l}] m(\{s_{k,l}\})$$

where

$$\begin{aligned} C(\{\sigma^{k,l}\}) &= c_1^{-1} [\det(\sigma^{k,l})]^{n/2} \\ c_1 &= 2^{(Nn/2)} \times \pi^{N(N-1)/4} \times \Gamma(n/2) \Gamma((n-1)/2) \cdots \Gamma((n-N+1)/2) \\ m(\{s_{k,l}\}) &= [\det(s_{k,l})]^{(n-N-2)/2} \end{aligned}$$

According to (5) the UMPU test for hypothesis (2) has the form:

$$\varphi_{i,j}(\{s_{k,l}\}) = \begin{cases} 0, & \text{if } c'_{i,j}(\{s_{k,l}\}) < s_{i,j} < c''_{i,j}(\{s_{k,l}\}), (k,l) \neq (i,j) \\ 1, & \text{if } s_{i,j} \leq c'_{i,j}(\{s_{k,l}\}) \text{ or } s_{i,j} \geq c''_{i,j}(\{s_{k,l}\}), (k,l) \neq (i,j) \end{cases} \quad (8)$$

where the critical values  $c'_{i,j}, c''_{i,j}$  are defined from the equations (according to

60 (6),(7))

$$\frac{\int_{I \cap [c'_{i,j}; c''_{i,j}]} [\det(s_{k,l})]^{(n-N-2)/2} ds_{i,j}}{\int_I [\det(s_{k,l})]^{(n-N-2)/2} ds_{i,j}} = 1 - \alpha \quad (9)$$

$$\begin{aligned} & \int_{I \cap (-\infty; c'_{i,j}]} s_{i,j} [\det(s_{k,l})]^{(n-N-2)/2} ds_{i,j} + \\ & + \int_{I \cap [c''_{i,j}; +\infty)} s_{i,j} [\det(s_{k,l})]^{(n-N-2)/2} ds_{i,j} = \\ & = \alpha \int_I s_{i,j} [\det(s_{k,l})]^{(n-N-2)/2} ds_{i,j} \end{aligned} \quad (10)$$

where  $I$  is the interval of values of  $s_{i,j}$  such that the matrix  $S = (s_{k,l})$  is positive definite and  $\alpha$  is the significance level of the test.

Let  $S = (s_{k,l})$  be positive definite (this is true with probability 1 if  $n > N$ ).

65 Consider  $\det(s_{k,l})$  as a function of the variable  $s_{i,j}$  only, when fixing the values of all others  $\{s_{k,l}\}$ . This determinant is a quadratic polynomial of  $s_{i,j}$ :

$$\det(s_{k,l}) = -as_{i,j}^2 + bs_{i,j} + c \quad (11)$$

Let  $K = (n-N-2)/2$ . Denote by  $x_1, x_2$  ( $x_1 < x_2$ ) the roots of the equation  $-ax^2 + bx + c = 0$ . One has

$$\int_f^d (ax^2 - bx - c)^K dx = (-1)^K a^K (x_2 - x_1)^{2K+1} \int_{\frac{f-x_1}{x_2-x_1}}^{\frac{d-x_1}{x_2-x_1}} u^K (1-u)^K du$$

Therefore the equation (9) takes the form:

$$\int_{\frac{c'-x_1}{x_2-x_1}}^{\frac{c''-x_1}{x_2-x_1}} u^K (1-u)^K du = (1-\alpha) \int_0^1 u^K (1-u)^K du \quad (12)$$

or

$$\frac{\Gamma(2K+2)}{\Gamma(K+1)\Gamma(K+1)} \int_{\frac{c'-x_1}{x_2-x_1}}^{\frac{c''-x_1}{x_2-x_1}} u^K (1-u)^K du = (1-\alpha) \quad (13)$$

It means that conditional distribution of  $S_{i,j}$  when all other  $S_{k,l}$  are fixed,  
<sup>70</sup>  $S_{k,l} = s_{k,l}$  is the beta distribution  $Be(K+1, K+1)$ .

Beta distribution  $Be(K+1, K+1)$  is symmetric with respect to the point  $\frac{1}{2}$ . Therefore the significance level condition (9) and unbiasedness condition (10) are satisfied if and only if:

$$\frac{c''-x_1}{x_2-x_1} = 1 - \frac{c'-x_1}{x_2-x_1}$$

Let  $q$  be the  $\frac{\alpha}{2}$ -quantile of beta distribution  $Be(K+1, K+1)$ , i.e.  $F_{Be}(q) = \frac{\alpha}{2}$ . Then thresholds  $c'$ ,  $c''$  are defined by:

$$\begin{aligned} c' &= x_1 + (x_2 - x_1)q \\ c'' &= x_2 - (x_2 - x_1)q \end{aligned} \quad (14)$$

Finally, the UMPU test for testing conditional independence of  $X_i, X_j$  has the form

$$\varphi_{i,j} = \begin{cases} 0, & 2q-1 < \frac{as_{i,j} - \frac{b}{2}}{\sqrt{\frac{b^2}{4} + ac}} < 1-2q \\ 1, & \text{otherwise} \end{cases} \quad (15)$$

<sup>75</sup> where  $a, b, c$  are defined in (11).

## 5. Sample partial correlation test.

It is known [Lauritzen 1996] that hypothesis  $\sigma^{i,j} = 0$  is equivalent to the hypothesis  $\rho^{i,j} = 0$ , where  $\rho^{i,j}$  is the partial correlation between  $X_i$  and  $X_j$  given  $X_k, k \in N(i, j) = \{1, 2, \dots, N\} \setminus \{i, j\}$ :

$$\rho^{i,j} = -\frac{\sigma^{i,j}}{\sqrt{\sigma^{i,i}\sigma^{j,j}}} = \frac{-\Sigma^{i,j}}{\sqrt{\Sigma^{i,i}\Sigma^{j,j}}}$$

where for a given matrix  $A = (a_{k,l})$  we denote by  $A^{i,j}$  the cofactor of the element  $a_{i,j}$ . Denote by  $r^{i,j}$  sample partial correlation

$$r^{i,j} = \frac{-S^{i,j}}{\sqrt{S^{i,i}S^{j,j}}}$$

where  $S^{i,j}$  is the cofactor of the element  $s_{i,j}$  in the matrix  $S$  of sample covariances.

Well known sample partial correlation test for testing hypothesis  $\rho^{i,j} = 0$  has the form [Anderson 2003]:

$$\varphi_{i,j} = \begin{cases} 0, & |r^{i,j}| \leq c_{i,j} \\ 1, & |r^{i,j}| > c_{i,j} \end{cases} \quad (16)$$

where  $c_{i,j}$  is  $(1 - \alpha/2)$ -quantile of the distribution with the following density function

$$f(x) = \frac{1}{\sqrt{\pi}} \frac{\Gamma(n - N + 1)/2}{\Gamma((n - N)/2)} (1 - x^2)^{(n - N - 2)/2}, \quad -1 \leq x \leq 1$$

Note, that in practical applications the following Fisher transformation is applied:

$$z_{i,j} = \frac{\sqrt{n}}{2} \ln \left( \frac{1 + r^{i,j}}{1 - r^{i,j}} \right)$$

Under condition  $\rho^{i,j} = 0$  statistic  $Z_{i,j}$  has asymptotically standard normal distribution. That is why the following test is largely used in applications [Drton and Perlman 2004], [Drton and Perlman 2007], [Drton and Perlman 2008]:

$$\varphi_{i,j} = \begin{cases} 0, & |z_{i,j}| \leq c_{i,j} \\ 1, & |z_{i,j}| > c_{i,j} \end{cases} \quad (17)$$

where the constant  $c_{i,j}$  is  $(1 - \alpha/2)$ -quantile of standard normal distribution.

In this section we prove that the UMPU test (15) can be reduced to the sample partial correlation test (16), and therefore the well known sample partial correlation test for conditional independence is the UMPU one.

**Theorem:** *Sample partial correlation test (16) is equivalent to UMPU test (15) for testing hypothesis  $\rho^{i,j} = 0$  vs  $\rho^{i,j} \neq 0$ .*

90 *Proof:* it is sufficient to prove that

$$\frac{S^{i,j}}{\sqrt{S^{i,i}S^{j,j}}} = \frac{aS_{i,j} - \frac{b}{2}}{\sqrt{\frac{b^2}{4} + ac}} \quad (18)$$

To prove this equation we introduce some notations. Let  $A = (a_{k,l})$  be an  $(N \times N)$  symmetric matrix. Fix  $i < j$ ,  $i, j = 1, 2, \dots, N$ . Denote by  $A(x)$  the matrix obtained from  $A$  by replacing the elements  $a_{i,j}$  and  $a_{j,i}$  by  $x$ . Denote by  $A^{i,j}(x)$  the cofactor of the element  $(i, j)$  in the matrix  $A(x)$ . Then the following  
95 statement is true

**Lemma:** *One has  $[\det A(x)]' = -2A^{i,j}(x)$ .*

*Proof of the Lemma:* one has from the general Laplace decomposition of  $\det A(x)$  by two rows  $i$  and  $j$ :

$$\begin{aligned} \det(A(x)) &= \det \begin{pmatrix} a_{i,i} & x \\ x & a_{j,j} \end{pmatrix} A^{\{i,j\},\{i,j\}} + \sum_{k < j, k \neq i} \det \begin{pmatrix} a_{i,k} & x \\ a_{j,k} & a_{j,j} \end{pmatrix} A^{\{i,j\},\{k,j\}} + \\ &+ \sum_{k > j} \det \begin{pmatrix} x & a_{i,k} \\ a_{j,j} & a_{j,k} \end{pmatrix} A^{\{i,j\},\{j,k\}} + \sum_{k < i} \det \begin{pmatrix} a_{i,k} & a_{i,i} \\ a_{j,k} & x \end{pmatrix} A^{\{i,j\},\{k,i\}} + \\ &\sum_{k > i, k \neq j} \det \begin{pmatrix} a_{i,i} & a_{i,k} \\ x & a_{j,k} \end{pmatrix} A^{\{i,j\},\{i,k\}} + \sum_{k < l, k, l \neq i, j} \det \begin{pmatrix} a_{i,k} & a_{i,l} \\ a_{j,k} & a_{j,l} \end{pmatrix} A^{\{i,j\},\{k,l\}} \end{aligned}$$

where  $A^{\{i,j\},\{k,l\}}$  is the cofactor of the matrix  $\begin{pmatrix} a_{i,k} & a_{i,l} \\ a_{j,k} & a_{j,l} \end{pmatrix}$  in the matrix  $A$ .

Taking the derivative of  $\det A(x)$  one get

$$\begin{aligned} [\det(A(x))]' &= -2xA^{\{i,j\},\{i,j\}} - \sum_{k < j, k \neq i} a_{j,k} A^{\{i,j\},\{k,j\}} + \sum_{k > j} a_{j,k} A^{\{i,j\},\{j,k\}} + \\ &+ \sum_{k < i} a_{k,i} A^{\{i,j\},\{k,i\}} - \sum_{k > i, k \neq j} a_{k,i} A^{\{i,j\},\{k,i\}} = -2A^{i,j}(x) \end{aligned}$$

The last equation follows from the symmetry conditions  $a_{k,l} = a_{l,k}$  and from Laplace decompositions of  $A^{i,j}(x)$  by the row  $j$  and the column  $i$ . Lemma is proved. Note, that similar result is proved in ([Anderson 2003], Appendix A).

Now we come back to the proof of the theorem. One has  $\det(S(x)) = -ax^2 + bx + c$ , where  $a, b, c$  are the same as in (11). Therefore by Lemma



one has  $[\det S(x)]' = -2ax + b = -2S^{i,j}(x)$ , i.e.  $S^{i,j}(x) = ax - b/2$ . Let  $x = s_{i,j}$  then  $as_{i,j} - \frac{b}{2} = S^{i,j}$ . To prove the theorem it is sufficient to prove that  $\sqrt{S^{i,i}S^{j,j}} = \sqrt{\frac{b^2}{4} + ac}$ . Let  $x_2 = \frac{b + \sqrt{b^2 + 4ac}}{2a}$  be the maximum root of equation  $ax^2 - bx - c = 0$ . Then  $ax_2 - \frac{b}{2} = \sqrt{\frac{b^2}{4} + ac}$ . Consider

$$r^{i,j}(x) = \frac{-S^{i,j}(x)}{\sqrt{S^{i,i}S^{j,j}}}$$

According to Silvester determinant identity one can write :

$$S^{\{i,j\},\{i,j\}} \det S(x) = S^{i,i}S^{j,j} - [S^{i,j}(x)]^2$$

Therefore for  $x = x_1$  and  $x = x_2$  one has

$$S^{i,i}S^{j,j} - [S^{i,j}(x)]^2 = 0$$

That is for  $x = x_1$  and  $x = x_2$  one has  $r^{i,j}(x) = \pm 1$ . The equation  $S^{i,j}(x) = ax - \frac{b}{2}$  implies that when  $x$  is increasing from  $x_1$  to  $x_2$  then  $r^{i,j}(x)$  is decreasing from 1 to -1. That is  $r^{i,j}(x_2) = -1$ , i.e.  $ax_2 - \frac{b}{2} = \sqrt{S^{i,i}S^{j,j}}$ . Therefore

$$\sqrt{S^{i,i}S^{j,j}} = \sqrt{\frac{b^2}{4} + ac}$$

100 The Theorem is proved.

Finally, the UMPU test for testing conditional independence of  $X_i$  and  $X_j$  can be written in the following form

$$\varphi_{i,j} = \begin{cases} 0, & 2q - 1 < r^{i,j} < 1 - 2q \\ 1 & \text{otherwise} \end{cases} \quad (19)$$

where  $r^{i,j}$  is the sample partial correlation, and  $q$  is the  $\frac{\alpha}{2}$ -quantile of beta distribution  $Be(\frac{n-N}{2}, \frac{n-N}{2})$ ,

## 105 6. Concluding remarks

In general optimality of tests for individual hypotheses testing does not imply optimality of multiple testing procedures. However if the losses from false decisions are supposed to be additive then it is possible to prove optimality from decision-theoretic point of view of some multiple testing procedures

110 [Lehmann 1957], [Koldanov *et al* 2013], [Hochberg 1987]. Application of this  
approach for Gaussian graphical model selection is a subject of further investi-  
gations.

**Acknowledgment:** The work was conducted at National Research University  
Higher School of Economics, Laboratory of Algorithms and Technologies for  
115 Network Analysis. Partly supported by RFFI 14-01-00807.

## References

## References

- [Anderson 2003] Anderson T. (2003). An introduction to multivariate statistical  
analysis.3-d edition. Wiley-Interscience, New York.
- 120 [Drton and Perlman 2004] Drton M. Perlman M.(2004). Model selection for  
Gaussian concentration graph. *Biometrika* **91(3)**, 591–602.
- [Drton and Perlman 2007] Drton M. Perlman M.(2007). Multiple Testing and  
Error Control in Gaussian Graphical Model Selection. *Statistical Science*  
**22(3)**, 430–449.
- 125 [Drton and Perlman 2008] Drton M. Perlman M.(2008). A SINful approach to  
Gaussian graphical model selection. *Journal of Statistical Planning and  
Inference* **138**, 1179–1200.
- [Edwards 2000] Edwards D.(2000). Introduction to graphical modeling.  
Springer-Verlag New York, Inc.
- 130 [Lauritzen 1996] Lauritzen S.L. (1996). Graphical models. Oxford university  
press.
- [Lehmann and Romano 2005] Lehmann E.L. Romano J.P. (2005). Testing sta-  
tistical hypotheses. 3-d edition. Springer, New York.
- [Lehmann 1957] Lehmann E.L. (1957). A theory of some multiple decision prob-  
135 lems, I. *The Annals of Mathematical Statistics*, 1-25.

[Koldanov *et al* 2013] Koldanov A.P., Koldanov P.A., Kalyagin V.A., Pardalos P.M. (2013) Statistical procedures for the market graph construction. *Computational Statistics & Data Analysis* **68**, 17–29, Elsevier.

[Hochberg 1987] Hochberg, Y., Tamhane, A. Multiple comparison procedures. John Wiley & Sons, Inc. 1987.

140