

Closed-form REML estimators and sample size determination for mixed effects models for repeated measures under monotone missingness

Yongqiang Tang

We derive the closed-form restricted maximum likelihood (REML) estimator and Kenward-Roger's variance estimator for fixed effects in the mixed effects model for repeated measures (MMRM) when the missing data pattern is monotone. As an important application of the analytic result, we present the formula for calculating the power of treatment comparison using the Wald t test with the Kenward-Roger adjusted variance estimate in MMRM. It allows adjustment for baseline covariates without the need to specify the covariate distribution in randomized trials. A simple two-step procedure is proposed to determine the sample size needed to achieve the targeted power. The proposed method performs well for both normal and moderately nonnormal data even in small samples ($n = 20$) in simulations. An anti-depressant trial is analyzed for illustrative purposes. Copyright © 2015 John Wiley & Sons, Ltd.

Keywords: Kenward-Roger's variance estimator; Power and sample size; Restricted maximum likelihood

1. Introduction

In clinical trials, the mixed effects model for repeated measures (MMRM) is commonly used to analyze longitudinal continuous outcomes collected at a number of fixed time points [1, 2, 3]. In MMRM, there is no random effect, and the within-subject dependence is modeled by an unstructured covariance matrix. It assumes that the post-baseline outcome $\mathbf{y}_i = (y_{i1}, \dots, y_{ip})'$ follows a p -dimensional multivariate normal distribution

$$\mathbf{y}_i \sim N_p[(\boldsymbol{\alpha}'_1 \mathbf{x}_i, \dots, \boldsymbol{\alpha}'_p \mathbf{x}_i)', \Sigma], \quad (1)$$

where \mathbf{x}_i is the baseline covariate (including intercept and treatment status) for subject i , and $\boldsymbol{\alpha}_j$ is the $q \times 1$ vector of covariate and treatment effects at visit j . In clinical trials, the primary analysis model is generally pre-specified in the protocol. While it is difficult, if not impossible, to correctly pre-specify the covariance structure, the use of an unstructured covariance matrix usually provides reasonable control of the type I error rate [1]. Any stronger assumptions on the mean response or covariance structure can be difficult to be accepted by regulatory agencies without rigid justifications.

Shire, 300 Shire Way, Lexington, MA 02421, USA

* Correspondence to: E-mail: yongqiang_tang@yahoo.com

Some theories for model (1) can be presented in a simple and elegant way [4] for monotone missing data. Closed-form maximum likelihood (ML) estimators [5] and Bayes estimators [6] can be obtained by factoring the observed data distribution as $f(y_{i1}, \dots, y_{ij}) = f(y_{i1}) \prod_{k=2}^j f(y_{ik} | y_{i1}, \dots, y_{i(k-1)})$. These analytic results are of great practical interest since the real clinical data are often monotone or approximately monotone. For near-monotone data, monotone pattern can be obtained by removing subjects with intermittent missing data [7], by ignoring data collected after the first missing visit, or by filling in the intermittent missing data using a single imputation method. A numerical example is provided in Section 2.4. Alternatively, the intermittent missing data can be handled using the monotone expectation-maximization (MEM) algorithm [8] or monotone data augmentation (MDA) algorithm [6, 9] whenever appropriate.

In Section 2, we derive the closed-form restricted maximum likelihood (REML) estimator and assess the biases of ML and REML estimators for model (1) under monotone missingness. We obtain the closed-form Kenward-Roger (KR) adjusted variance estimator [10] and compare it with the delta variance for inference on fixed effects. An anti-depressant trial [11] is analyzed for illustrative purposes.

Sample size calculation is critical in planning a clinical trial. If the sample size is too small, the trial may have little chance to detect a clinically meaningful treatment effect, whereas an unnecessarily large sample may waste time, money and resources. Sample size determination methods were previously developed for MMRM [3, 12] based on the asymptotic variance of the treatment effect and/or normal approximation. Since the use of the asymptotic variance underestimates the true variance of the estimated treatment effect [10], and the normal distribution can not adequately approximate the t distribution, the methods in [3, 12] tend to underestimate the required size especially in small samples.

In Section 3, we derive the power formula for the KR adjusted Wald t test comparing two treatments in MMRM. The power is computed based on the KR adjusted variance and t test. Therefore, the proposed method tends to be more accurate than the methods of [3, 12], but the corresponding formulae have simpler expressions. One advantage of our method is that it allows adjustment for covariates, but there is no need to specify the covariate distribution. Information about the covariate distribution is usually not available at the design stage since subjects will be enrolled into the trials only if they meet certain inclusion/exclusion criteria. There is no closed-form solution for the sample size needed to achieve a targeted power. We propose a two-step procedure to approximate the required size and assess its performance for both normal and non-normal continuous data in small to moderate samples via simulation.

We focus on the monotone missing data. Section 5 will discuss how to apply the sample size calculation method to non-monotone missing data.

2. REML estimator under monotone missingness

2.1. Closed-form REML estimator

We assume the missing data pattern is monotone in the sense that if y_{ij} is observed, then y_{it} 's are observed for all $t \leq j$. Let r_i denote the dropout pattern according to the last observation. A subject is in pattern $r_i = s$ if y_{ij} is observed for $j \leq s$, $r_i = 0$ if y_i is missing, and $r_i = p$ if the subject completes all p visits. Without loss of generality, we sort the data so that subjects in pattern s are before subjects in pattern t if $s > t$. For notational simplicity, we assume there are two treatment groups ($g_i = 1$ for active treatment, $g_i = 0$ for placebo). Let \mathcal{B}_{gs} denote the subset of subjects retained at visit s (y_{is} is observed) in group g , n_{gs} the number of subjects in \mathcal{B}_{gs} , and $n_s = \sum_{g=0}^1 n_{gs}$. Let x_{bi} denote the baseline covariates excluding the intercept term and g_i . The baseline outcome y_{i0} is usually included as a covariate in x_{bi} . Let \bar{x}_{gs} be the average of x_{bi} among subjects in \mathcal{B}_{gs} .

The clinical data are often analyzed by model (1), where $x_i = (1, x'_{bi}, g_i)'$ and $\alpha_j = (\mu_j, \alpha'_{jb}, \tau_j)'$. The ML estimator of the variance parameters is biased downward since it does not take into account of the loss in degrees of freedom (d.f.) from the estimation of α_j 's [13]. The REML procedure is generally employed to reduce the bias in estimating the variance parameters, and inference for fixed effects is often made based on the KR variance estimate especially in

small samples. Let W_{ij} be a $pq \times 1$ vector whose $[(j-1)q+1]$ -th through $[jq]$ -th entries are given by \mathbf{x}_i while other entries are 0's. Let $W_i = (W_{i1}, \dots, W_{ir_i})'$ and $\mathbf{y}_{io} = (y_{i1}, \dots, y_{ir_i})'$. Let $W = (W'_1, \dots, W'_n)'$ and $Y = (\mathbf{y}'_{1o}, \dots, \mathbf{y}'_{no})'$. The restricted likelihood is the marginal likelihood of KY , where $N = \sum_{i=1}^n r_i$ and K is any $(N-pq) \times N$ matrix of full rank satisfying $KW = \mathbf{0}$. The restricted log-likelihood ℓ_r does not depend on α and the choice of K , and can be expressed as [14, 15]

$$2\ell_r = - \sum_{i=1}^n \log |\Sigma_{r_i}| - \log \left| \sum_{i=1}^n W'_i \Sigma_{r_i}^{-1} W_i \right| - \left[\sum_{i=1}^n \mathbf{y}'_{io} \Sigma_{r_i}^{-1} \mathbf{y}_{io} - \left(\sum_{i=1}^n \mathbf{y}'_{io} \Sigma_{r_i}^{-1} W_i \right) \left(\sum_{i=1}^n W'_i \Sigma_{r_i}^{-1} W_i \right)^{-1} \left(\sum_{i=1}^n W'_i \Sigma_{r_i}^{-1} \mathbf{y}_{io} \right) \right], \quad (2)$$

where Σ_k is the leading $k \times k$ submatrix of Σ .

Let $\Sigma = L\Lambda L'$ be the LDL decomposition of Σ , where $U = \begin{bmatrix} 1 & 0 & \dots & 0 \\ -\beta_{21} & 1 & \dots & 0 \\ & \dots & \dots & 0 \\ -\beta_{p1} & \dots & -\beta_{p,p-1} & 1 \end{bmatrix}$, $L = U^{-1}$ and $\Lambda =$

$\text{diag}(\sigma_1^2, \dots, \sigma_p^2)$. Let l_{jt} be the (j, t) entry of L . The decomposition admits nice interpretation. The entries $\beta_j = (\beta_{j1}, \dots, \beta_{jj-1})'$ are the regression parameters of y_{ij} on $\vec{\mathbf{y}}_{j-1} = (y_{i1}, \dots, y_{i,j-1})'$ since model (1) can be written as

$$(\underline{\mathbf{y}}_{i1}, \dots, \underline{\mathbf{y}}_{ip})' \sim N_p[(\underline{\alpha}'_1 \mathbf{x}_i, \dots, \underline{\alpha}'_p \mathbf{x}_i)', \Lambda], \quad (3)$$

which can be reorganized as [9]

$$y_{ij} = \mathbf{x}'_i \underline{\alpha}_j + \vec{\mathbf{y}}'_{i,j-1} \beta_j + \varepsilon_{ij} = \mathbf{z}'_{i,j-1} \boldsymbol{\theta}_j + \varepsilon_{ij} \text{ for } j \leq p, \quad (4)$$

where $\underline{\mathbf{y}}_{ij} = y_{ij} - \sum_{t=1}^{j-1} \beta_{jt} y_{it}$, $\underline{\alpha}_j = \alpha_j - \sum_{t=1}^{j-1} \beta_{jt} \alpha_t$, $\boldsymbol{\theta}_j = (\underline{\alpha}'_j, \beta'_j)'$, $\mathbf{z}_{ij} = (\mathbf{x}'_i, \vec{\mathbf{y}}'_{i,j-1})'$, $\varepsilon_{ij} \sim N(0, \sigma_j^2)$, and ε_{ij} 's are independent.

Let $X_{o_j} = [\mathbf{x}_1, \dots, \mathbf{x}_{n_j}]'$, $Z_{o_j} = [\mathbf{z}_{1j}, \dots, \mathbf{z}_{n_j j}]'$, and $Y_{o_j} = (y_{1j}, \dots, y_{n_j j})'$. The least square (LS) estimator for the parameters in model (4) is

$$\hat{\boldsymbol{\theta}}_j = (Z'_{o_j} Z_{o_j})^{-1} Z'_{o_j} Y_{o_j} \text{ and } \hat{\sigma}_j^2 = \hat{S}_j / (n_j - q - j + 1),$$

where $\hat{S}_j = \sum_{i=1}^{n_j} (y_{ij} - \mathbf{z}'_{i,j-1} \hat{\boldsymbol{\theta}}_j)^2$. The ML estimator for the parameters in model (4) is given by

$$\hat{\boldsymbol{\theta}}_{j,ml} = \hat{\boldsymbol{\theta}}_j \text{ and } \hat{\sigma}_{j,ml}^2 = \hat{S}_j / n_j.$$

Lemma 1 derives the closed-form REML estimator for the parameters in model (1). Its proof is given in the appendix.

Lemma 1 (a) The asymptotic variance of the fixed effect estimate $(\hat{\alpha}'_1, \dots, \hat{\alpha}'_p)'$ is $(\sum_{i=1}^n W'_i \Sigma_{r_i}^{-1} W_i)^{-1} = (L \otimes I_q) V_w (L' \otimes I_q)$, where I_q is the $q \times q$ identity matrix, and $V_w = \text{diag}[\sigma_1^2 (X'_{o_1} X_{o_1})^{-1}, \dots, \sigma_p^2 (X'_{o_p} X_{o_p})^{-1}]$.

(b) The restricted log-likelihood (2) can be decomposed as the sum of p independent log-likelihoods $\ell_r = \sum_{j=1}^p \ell_{r_j} + \text{constant}$, where

$$\ell_{r_j} = \frac{q - n_j}{2} \log(\sigma_j^2) - \frac{(Y_{o_j} - \vec{Y}_{o_j} \beta_j)' Q_j (Y_{o_j} - \vec{Y}_{o_j} \beta_j)}{2\sigma_j^2}, \quad (5)$$

$Q_j = I - X_{o_j} (X'_{o_j} X_{o_j})^{-1} X_{o_j}$, and \vec{Y}_{o_j} is a $n_j \times (j-1)$ matrix whose (i, t) entry is y_{it} .

(c) The REML estimator is $\hat{\beta}_j = (\vec{Y}'_{o_j} Q_j \vec{Y}_{o_j})^{-1} \vec{Y}'_{o_j} Q_j Y_{o_j}$, $\hat{\underline{\alpha}}_j = (X'_{o_j} X_{o_j})^{-1} X'_{o_j} (Y_{o_j} - \vec{Y}_{o_j} \hat{\beta}_j)$ and $\hat{\sigma}_{j,re}^2 = \hat{S}_j / (n_j - q)$. Furthermore, $\hat{\boldsymbol{\theta}}_{j,re} = (\hat{\underline{\alpha}}'_j, \hat{\beta}'_j)' = \hat{\boldsymbol{\theta}}_j$.

(d) The asymptotic variance of $\hat{\beta}_j$ based on the observed or expected information matrix is given respectively by

$$\text{var}_o(\hat{\beta}_j) = \sigma_j^2 (\vec{Y}'_{o_j} Q_j \vec{Y}_{o_j})^{-1} \text{ and } \text{var}_E(\hat{\beta}_j) = \sigma_j^2 [\mathbb{E}(\vec{Y}'_{o_j} Q_j \vec{Y}_{o_j})]^{-1} = \frac{\sigma_j^2}{n_j - q} \Sigma_{j-1}^{-1}.$$

Remarks:

- 1) Lemma 1(a) is also valid for models with structured covariance matrix;
- 2) We set $\vec{Y}_{o_j} \beta_j = 0$ at $j = 1$ since β_j (\vec{Y}_{o_j}) is an empty vector (matrix);
- 3) $(\vec{Y}'_{o_j} Q_j \vec{Y}_{o_j})^{-1}$ is identical to the lower-right $(j - 1) \times (j - 1)$ submatrix of $(Z'_{o_j} Z_{o_j})^{-1}$.

Lemma 1(b) can also be derived from the Bayesian viewpoint based on the fact [14] that the restricted likelihood is proportional to the marginal posterior density over the variance parameters under a flat prior for (α, Σ) . We have to ignore the Jacobian correction factor in the posterior density under reparameterization since the REML estimators are invariant under reparameterization. Under a flat prior, (θ_j, σ_j^2) 's are independent in the posterior distribution, and the posterior density of (θ_j, σ_j^2) is proportional to the likelihood function for model (4), which implies that

$$\begin{aligned} \sigma_j^2 | Y &\sim \hat{S}_j / \chi_{n_j - q - j - 1}^2, \\ \beta_j | \sigma_j^2, Y &\sim N_{j-1}(\hat{\beta}_j, \sigma_j^2 [\vec{Y}'_{o_j} Q_j \vec{Y}_{o_j}]^{-1}), \\ \alpha_j | \sigma_j^2, \beta_j, Y &\sim N_q[(X'_{o_j} X_{o_j})^{-1} X'_{o_j} (Y_{o_j} - \vec{Y}_{o_j} \beta_j), \sigma_j^2 (X'_{o_j} X_{o_j})^{-1}]. \end{aligned}$$

The marginal posterior distribution of (β_j, σ_j^2) is given by

$$f(\sigma_j^2, \beta_j | Y) = f(\sigma_j^2 | Y) f(\beta_j | \sigma_j^2, Y) \propto \exp(\ell_{r_j}).$$

2.2. Bias in parameter estimates

The LS, ML and REML methods produce identical estimate of θ_j 's. Since $\hat{\theta}_j$ is an unbiased estimate of θ_j , it is easy to show by induction that $(\hat{l}_{j1}, \dots, \hat{l}_{jj})$ and $\hat{\alpha}_j = \sum_{t=1}^j \hat{l}_{jt} \hat{\alpha}_t$ are unbiased respectively for (l_{j1}, \dots, l_{jj}) and $\alpha_j = \sum_{t=1}^j l_{jt} \alpha_t$.

For σ_j^2 , the LS estimator is unbiased. The bias $\mathbb{E}(\hat{\sigma}_{j,re}^2) - \sigma_j^2 = -(j - 1)\sigma_j^2 / (n_j - q)$ of the REML estimator is smaller than that $\mathbb{E}(\hat{\sigma}_{j,ml}^2) - \sigma_j^2 = -(q + j - 1)\sigma_j^2 / n_j$ of the ML estimator.

The bias is not invariant to reparameterization. In REML, the bias in $\hat{\Sigma} = \hat{L} \hat{\Lambda} \hat{L}'$ is of the order of $O(n^{-2})$. It can be derived by the Cox-Snell [16] method (using the equation on line 21, page 2586 of [17] and equation (9.62) of [18]), or based on the second-order Taylor series expansion of $\hat{\Sigma} - \Sigma = \hat{L} \hat{\Lambda} \hat{L}' - L \Lambda L'$ around (β_j, σ_j^2) 's. In ML, the bias in $\hat{\Sigma}_{ML}$ is $O(n^{-1})$ since $\mathbb{E}(\hat{\Sigma}_{re} - \hat{\Sigma}_{ML}) = \mathbb{E}[\hat{L} \text{diag}(\varphi_1, \dots, \varphi_p) \hat{L}']$, where $\varphi_j = \mathbb{E}(\hat{\sigma}_{j,re}^2 - \hat{\sigma}_{j,ml}^2) = O(n^{-1})$. The bias in the LS estimator of Σ is also $O(n^{-1})$.

2.3. Variance of fixed effects

The fixed effect estimate is $\hat{\alpha} = (\hat{\alpha}_1, \dots, \hat{\alpha}_p)' = \hat{L}(\hat{\alpha}_1, \dots, \hat{\alpha}_p)'$, where $\hat{\alpha}_j$ is defined in Lemma 1c. The asymptotic variance of $\hat{\alpha}$ given in Lemma 1a is in fact the variance of the estimate $(\tilde{\alpha}'_1, \dots, \tilde{\alpha}'_p)'$ assuming that the variance parameters (β_j, σ_j^2) 's and hence L are known, where $(\tilde{\alpha}_1, \dots, \tilde{\alpha}_p)' = L(\tilde{\alpha}_1, \dots, \tilde{\alpha}_p)'$, $\tilde{\alpha}_j = (X'_{o_j} X_{o_j})^{-1} X'_{o_j} (Y_{o_j} - \vec{Y}_{o_j} \beta_j)$, and $\text{var}(\tilde{\alpha}_j) = \sigma_j^2 (X'_{o_j} X_{o_j})^{-1}$. Let $\Phi_p = \text{var}(\tilde{\alpha}_p) = \sum_{j=1}^p l_{pj}^2 \sigma_j^2 (X'_{o_j} X_{o_j})^{-1}$.

In the appendix, we show $\text{cov}(\hat{\alpha}_p - \tilde{\alpha}_p, \tilde{\alpha}_p) = \mathbf{0}$, and derive the variance of $\hat{\alpha}_p - \tilde{\alpha}_p$

$$\Psi_p = \text{var}(\hat{\alpha}_p - \tilde{\alpha}_p) = \sum_{j=2}^p l_{pj}^2 (V_{dj} + V_{ej}) \approx \sum_{j=2}^p l_{pj}^2 V_{dj}, \quad (6)$$

where L_j is the leading $j \times j$ submatrix of L , $(\omega_{j1}, \dots, \omega_{j,j-1})$ are the diagonal elements of $L'_{j-1} \text{var}(\hat{\beta}_j) L_{j-1}$ for $j \geq 2$, $D_{jt} = (X'_{o_j} X_{o_j})^{-1} - (X'_{o_t} X_{o_t})^{-1}$, $V_{dj} = \sum_{t=1}^{j-1} \omega_{jt} \sigma_t^2 D_{jt}$, $V_{e1} = \mathbf{0}$, $V_{ej} = \sum_{t=1}^{j-1} \omega_{jt} (V_{dt} + V_{et})$ at $j \geq 2$, V_{ej} is of lower order than V_{dj} and can be ignored. Thus

$$\text{var}(\hat{\alpha}_p) = \Phi_p + \Psi_p = \sum_{j=1}^p l_{pj}^2 \sigma_j^2 (X'_{o_j} X_{o_j})^{-1} + \sum_{j=2}^p l_{pj}^2 V_{dj}. \quad (7)$$

Although Ψ_p is of lower order than Φ_p , it is not negligible in small samples. The variance at other visits can be derived similarly, and would not be presented here.

Let $\Delta_s = \bar{\mathbf{x}}_{1s} - \bar{\mathbf{x}}_{0s}$, $S_{x_s} = \sum_{g=0}^1 \sum_{i \in \mathcal{B}_{gs}} (\mathbf{x}_{bi} - \bar{\mathbf{x}}_{gs})^{\otimes 2}$ and $V_{x_s} = n_{1s}^{-1} + n_{0s}^{-1} + \Delta'_s S_{x_s}^{-1} \Delta_s$, where $a^{\otimes 2} = aa'$. By equation (7), the variance of the treatment effect estimate $\hat{\tau}_p$ is given by

$$\text{var}(\hat{\tau}_p) = \sum_{j=1}^p l_{pj}^2 \sigma_j^2 V_{x_j} + \sum_{j=2}^p l_{pj}^2 \sum_{t=1}^{j-1} \omega_{jt} \sigma_t^2 (V_{x_j} - V_{x_t}). \quad (8)$$

2.3.1. Kenward-Roger (KR) variance estimate The KR variance estimate has been widely used in practice particularly when the sample size is small. The KR approach not only takes into account of the variability in the variance parameter estimate, but also adjusts for the bias in $\hat{\Phi}_p$. In the appendix, we show the bias in $\hat{\Phi}_p$ is

$$\Psi_p^* = E(\hat{\Phi}_p) - \Phi_p = \sum_{j=2}^p \sum_{t=1}^{j-1} l_{pj}^2 \omega_{jt} \sigma_t^2 (X'_{o_t} X_{o_t})^{-1} - \sum_{j=2}^p \frac{j-1}{n_j - q} \sigma_j^2 l_{pj}^2 (X'_{o_j} X_{o_j})^{-1}. \quad (9)$$

If $\text{var}_E(\hat{\beta}_j)$ in Lemma 1d is used, then $\omega_{jt} \sigma_t^2 = \sigma_j^2 / (n_j - q)$, $\Psi_p = \sum_{j=2}^p l_{pj}^2 \sigma_j^2 [\sum_{t=1}^{j-1} D_{jt}] / (n_j - q)$, and $\Psi_p^* = -\Psi_p$. The KR variance evaluated at the REML estimator is given by

$$\widehat{\text{var}}_{\text{kr}}(\hat{\alpha}_p) = \hat{\Phi}_p + \hat{\Psi}_p - \hat{\Psi}_p^* = \hat{\Phi}_p + 2\hat{\Psi}_p, \quad (10)$$

and it provides a roughly unbiased estimate of the variance $\Phi_p + \Psi_p$ of $\hat{\alpha}_p$ while the lower order terms are dropped. The KR variance estimate is invariant under reparameterization of an unstructured covariance matrix [17]. Although the derivation of the KR variance estimate relies on $\text{var}_E(\hat{\beta}_j)$, equation (10) yields the same result as Kenward-Roger's [10] formula if the calculation is based on $\text{var}_O(\hat{\beta}_j)$.

2.3.2. Delta variance estimate An alternative variance estimate for $\hat{\alpha}_p$ can be obtained from the delta method

$$V_d = \sum_{j=1}^p \frac{\partial \alpha_p}{\partial \theta_j} \text{var}(\hat{\theta}_j) \left(\frac{\partial \alpha_p}{\partial \theta_j} \right)' = \sum_{j=1}^p l_{pj}^2 \sigma_j^2 \ddot{\alpha}_j (Z'_{o_j} Z_{o_j})^{-1} \ddot{\alpha}'_j,$$

where $\partial \alpha_p / \partial \theta_j = l_{pj} \ddot{\alpha}_j$ and $\ddot{\alpha}_j = (I_q, \alpha_1, \dots, \alpha_{j-1})$.

By equation (19) in the appendix, the delta variance evaluated at the LS estimator can be written as

$$\hat{V}_d = \sum_{j=1}^p \hat{l}_{pj}^2 \hat{\sigma}_j^2 (X'_{o_j} X_{o_j})^{-1} + \sum_{j=2}^p \hat{l}_{pj}^2 (\hat{\alpha}_j - \hat{\alpha}_j^*) V_{\beta_j} (\hat{\alpha}_j - \hat{\alpha}_j^*)',$$

where $\hat{\alpha}_j = (\hat{\alpha}_1, \dots, \hat{\alpha}_{j-1})$ and $\hat{\alpha}_j^* = (X'_{o_j} X_{o_j})^{-1} X'_{o_j} \bar{Y}_{o_j}$. The delta variance estimate tends to be more conservative than the KR variance estimate since the latter is a roughly unbiased variance estimate for $\hat{\alpha}_p$, while the former

Table 1: Treatment effect estimate and its standard error at week 6 in an antidepressant trial

Method	Treatment effect	Standard Error		
		Asymptotic	KR	Delta
MMRM	-2.799	1.114	1.116	1.122
cLDA	-2.799	1.098	1.108	-

overestimates the variance of $\hat{\alpha}_p$ by

$$E(\hat{V}_d) - \Phi_p - \Psi_p \approx \sum_{j=1}^p \text{var}(\hat{l}_{pj}) \sigma_j^2 (X'_{o_j} X_{o_j})^{-1} + \sum_{j=2}^p \text{var}(\hat{l}_{pj}) \left[\sum_{t=1}^{j-1} D_{jt} \omega_{jt} \sigma_t^2 \right].$$

The delta variance is closely related to the variance from the multiple imputation (MI) inference [19]. One may compare the KR variance estimate given in equation (10) with the MI variance derived by [19]. In general, the MI inference tends to be slightly more conservative than the REML inference [2].

2.4. Analysis of an antidepressant trial

To illustrate the analytic result derived in this section, we analyze an antidepressant trial reported by [11]. The Hamilton 17-item rating scale for depression (HAMD₁₇) is collected at baseline and weeks 1, 2, 4, 6. The dataset consists of 84 subjects on the active treatment, and 88 subjects on placebo. The number of subjects in patterns 0 to 4 is respectively (0, 7, 5, 11, 65) in the placebo arm, and (0, 6, 5, 9, 64) in the active arm. The missing data are mainly due to dropout. Only one subject has an intermittent missing value at visit 2. We impute the intermittent missing value as 11.70432 based on the linear regression of y_{i2} on (y_{i0}, y_{i1}, g_i) , where y_{i0} is the baseline HAMD₁₇. A better approach would be to replace the missing value with multiple plausible values to reflect uncertainty about the missing data via the MI procedure [20, 9]. However, due to the small amount of non-monotone missing data, the result from this single imputation approach is very close to that from the MI inference [20].

Table 1 reports the result from the MMRM analysis. The model includes visit, baseline \times visit and treatment \times visit interactions as the fixed effects, and an unstructured covariance matrix is used to model the within-patient errors. This corresponds to $\mathbf{x}_i = (1, y_{i0}, g_i)'$ in terms of model (1). SAS PROC MIXED yields the same result as our analytic formulae if a more stringent convergence criterion (CONVG= 1E-10 option) than the default is used. The KR variance estimate is smaller than the delta variance, but slightly larger than the asymptotic variance. The p -value for testing the treatment effect at week 6 is 0.0132 based on the KR variance estimate.

Table 1 also reports the result from the constrained longitudinal data analysis (cLDA), in which the baseline outcome is treated as a response variable instead of a covariate, and constrained to have equal mean across treatment groups [21, 22]. For monotone missing pattern, the REML estimate and the associated variance estimate can be derived using essentially the same technique described in Sections 2.1 and 2.3, and the details are omitted here. The REML estimates are in fact identical, and the corresponding KR variance estimates are very close in MMRM and cLDA. This is consistent with the simulation result reported in [22]. The cLDA is useful if there is a non-negligible amount of missing data at baseline. However, for monotone missing data, the MMRM is generally preferred since there is no need to assume that y_{i0} 's are normally distributed and that the baseline means are equal across treatment groups in MMRM.

3. Power and sample size formulae for MMRM

3.1. Power and sample size formulae

Throughout Section 3, we assume the hypothesis of interest is to test the treatment effect at the last visit. Let γ_g be the proportion of subjects randomized to group g , π_{gt} the retention rate at visit t in group g , and $\bar{\pi}_t = \sum_{g=0}^1 \gamma_g \pi_{gt}$ the pooled retention rate at visit t . Then $n_{gt} = n\gamma_g \pi_{gt}$. In the variance expression (8), x_{bi} 's are assumed to be fixed in the analysis, but unknown at the design stage. We will replace V_{x_j} 's in equation (8) by their expected values. Suppose the covariates are balanced between two arms. We don't require the covariates to be continuous, or follow a specific distribution in randomized trials. Let q^* be the dimension of x_{bi} , $q = q^* + 2$ and $\varpi_{\pi_t} = \sum_{g=0}^1 (\gamma_g \pi_{gt})^{-1}$. It is easy to show that $n_{0t}n_{1t}(n_t - q^* - 1)/(n_t q^*) \Delta_t' S_{x_t}^{-1} \Delta_t$ approximately follows a F distribution with q^* and $n_t - q^* - 1$ d.f. (this holds exactly if x_{bi} is normally distributed). We have

$$E(\Delta_t' S_{x_t}^{-1} \Delta_t) \approx \frac{\varpi_{\pi_t}}{n} \varpi_{\delta_t}(n) \text{ and } E(V_{x_t}) \approx \frac{\varpi_{x_t}(n)}{n}. \tag{11}$$

where $\varpi_{\delta_t}(n) = q^*/(n\bar{\pi}_t - q - 1)$ and $\varpi_{x_t}(n) = \varpi_{\pi_t}[1 + \varpi_{\delta_t}(n)]$ are functions of n .

If we use $\text{var}_E(\beta_j)$, then $\omega_{jt}\sigma_t^2 = \sigma_j^2/(n_j - q)$. The (expected) variance of $\hat{\tau}_p$ is $\varpi_{\tau}(n)/n$, where

$$\varpi_{\tau}(n) = \sum_{j=1}^p l_{pj}^2 \sigma_j^2 \varpi_{x_j}(n) + \sum_{j=2}^p l_{pj}^2 \sigma_j^2 \frac{\sum_{t=1}^{j-1} (\varpi_{x_j}(n) - \varpi_{x_t}(n))}{n\bar{\pi}_j - q}.$$

The power of the Wald test at a two-sided significance level of α is given by

$$P = \Pr\left(\left|\frac{\hat{\tau}_p}{\sqrt{\widehat{\text{var}}_{\text{kr}}(\hat{\tau}_p)}}\right| \geq t_{f, 1-\frac{\alpha}{2}}\right) = \Pr(t(f, \sqrt{n}\lambda) \geq t_{f, 1-\frac{\alpha}{2}}) + \Pr(t(f, \sqrt{n}\lambda) \leq -t_{f, 1-\frac{\alpha}{2}}) \approx \Pr(t(f, \sqrt{n}|\lambda|) \geq t_{f, 1-\frac{\alpha}{2}}), \tag{12}$$

where $\widehat{\text{var}}_{\text{kr}}(\hat{\tau}_p)$ is the (q, q) entry of $\widehat{\text{var}}_{\text{kr}}(\hat{\alpha}_p)$ defined in equation (10), $\lambda = \tau_p/\sqrt{\varpi_{\tau}(n)}$, $t(f, \nu)$ is distributed as a non-central t distribution with f d.f. and non-central parameter ν , and $t_{f,p}$ is the p -th percentile of t_f . To simplify the calculation, we approximate the power by

$$P = \Pr(\hat{\tau}_p^* \geq t_{f, 1-\frac{\alpha}{2}} - \sqrt{n}\lambda) + \Pr(\hat{\tau}_p^* \leq -t_{f, 1-\frac{\alpha}{2}} - \sqrt{n}\lambda) \approx \Pr(\hat{\tau}_p^* \leq \sqrt{n}|\lambda| - t_{f, 1-\frac{\alpha}{2}}), \tag{13}$$

where $\hat{\tau}_p^* = (\hat{\tau}_p - \tau_p)/\sqrt{\widehat{\text{var}}_{\text{kr}}(\hat{\tau}_p)}$ approximately follows a t distribution with f d.f., and

$$\sqrt{n}\lambda = \frac{\tau_p}{\sqrt{\widehat{\text{var}}_{\text{kr}}(\hat{\tau}_p)}} \approx \frac{\tau_p}{\sqrt{n^{-1}\varpi_{\tau}(n)}}.$$

For a scalar fixed effect, Kenward and Roger [10] used the Satterthwaite-type d.f., which is a random variable (could be larger than $n_1 - q$), and difficult to evaluate at the design stage. We approximate f by

$$f(n) = (n_1 - q)f_o = (n\bar{\pi}_1 - q)f_o,$$

where

$$f_o = \frac{(\sum_{j=1}^p l_{pj}^2 \sigma_j^2) \varpi_{x_1}}{\sum_{j=1}^p l_{pj}^2 \sigma_j^2 \varpi_{x_j}} \approx \frac{(\sum_{j=1}^p l_{pj}^2 \sigma_j^2) \varpi_{\pi_1}}{\sum_{j=1}^p l_{pj}^2 \sigma_j^2 \varpi_{\pi_j}}$$

can be roughly interpreted as the fraction of observed information among subjects retained at visit 1.

There is no closed-form solution for the sample size needed to achieve a desired power P . We propose a two-step procedure to approximate the required size. Inverting equation (13) yields

$$n_{\text{ur}} = (t_{f,1-\frac{\alpha}{2}} + t_{f,P})^2 \frac{\varpi_{\tau}(n)}{\delta_p^2} = \frac{(t_{f,1-\frac{\alpha}{2}} + t_{f,P})^2}{\lambda^2}. \quad (14)$$

In large samples, we can approximate $t_{f,p}$ by the p th percentile z_p of $N(0, 1)$, and approximate $\varpi_{\tau}(n)$ by $\varpi_{\tau}^*[1 + q^*(\sum_{j=1}^p b_j/\bar{\pi}_j)/n]$, where $\varpi_{\tau}^* = \sum_{j=1}^p l_{pj}^2 \sigma_j^2 \varpi_{\pi_j}$ and $b_j = l_{pj}^2 \sigma_j^2 \varpi_{\pi_j} / \varpi_{\tau}^*$. In step 1, we find the sample size based on the normal approximation

$$n_l = (z_{1-\frac{\alpha}{2}} + z_P)^2 \frac{\varpi_{\tau}^*}{\delta_p^2} + q^* \sum_{j=1}^p \frac{b_j}{\bar{\pi}_j} \approx (z_{1-\frac{\alpha}{2}} + z_P)^2 \frac{\varpi_{\tau}^*}{\delta_p^2} + \frac{q^*}{\bar{\pi}_p}, \quad (15)$$

and it generally provides a sharp lower bound on the required size. If the retention rate is low (e.g. $\bar{\pi}_p < 0.5$), we use the first equality in equation (15). In the implementation, we round n_l up to the nearest integer. In step 2, we use the approximation

$$n_u = (t_{f(n_l),1-\frac{\alpha}{2}} + t_{f(n_l),P})^2 \frac{\varpi_{\tau}(n_l)}{\delta_p^2}, \quad (16)$$

and it usually provides a very good estimate of the required size. We may approximate n_u by

$$n_u^* \approx (t_{f(n_l),1-\frac{\alpha}{2}} + t_{f(n_l),P})^2 \frac{\varpi_{\tau}^*}{\delta_p^2} + q^* \sum_{j=1}^p \frac{b_j}{\bar{\pi}_j} \approx (t_{f(n_l),1-\frac{\alpha}{2}} + t_{f(n_l),P})^2 \frac{\varpi_{\tau}^*}{\delta_p^2} + \frac{q^*}{\bar{\pi}_p}. \quad (17)$$

We recommend using equation (16) or equation (17) to estimate the sample size, and using equation (12) to evaluate the nominal power at a given sample size since equation (12) is more accurate than equation (13) when f is extremely small (e.g. $f < 12$). If the nominal power at n_u or n_u^* is not close to the targeted power, one may increase or decrease the sample size to achieve the desired power.

We also try a slight variation of the proposed size estimation procedure by increasing n_l defined in equation (15) by 2, but all other steps remain the same. More details are provided in simulation 1 reported in Section 3.2.1 below.

3.2. Numerical examples

3.2.1. Simulation 1. In an antidepressant trial analyzed in Section 2.4, the active treatment is significantly better than placebo in reducing depression. It might be of interest to design a new study to assess the effect of a similar compound on HAMD₁₇. Suppose $p = 4$, $y_{i0} \stackrel{i.i.d}{\sim} N(17.9, 5.5^2)$, $\mu_{i1} = 3.3 + 0.72y_0 + 0.1g_i$, $\mu_{i2} = 2.7 + 0.69y_0 - 1.5g_i$, $\mu_{i3} = 2.9 + 0.61y_0 - 2.3g_i$, $\mu_{i4} = 1 + 0.67y_0 + \tau_4 g_i$,

$$\begin{pmatrix} y_{i1} \\ y_{i2} \\ y_{i3} \\ y_{i4} \end{pmatrix} \sim N_4 \left(\begin{pmatrix} \mu_{i1} \\ \mu_{i2} \\ \mu_{i3} \\ \mu_{i4} \end{pmatrix}, \begin{pmatrix} 19.68 & 16.45 & 15.39 & 16.36 \\ 16.45 & 34 & 25.34 & 26.13 \\ 15.39 & 25.34 & 38.44 & 33.91 \\ 16.36 & 26.13 & 33.91 & 45.28 \end{pmatrix} \right),$$

and the retention rate is $(\pi_{01}, \dots, \pi_{04}) = (1, 0.92, 0.86, 0.74)$ and $(\pi_{11}, \dots, \pi_{14}) = (1, 0.93, 0.87, 0.76)$. These parameters are specified based roughly on the MMRM analysis of the antidepressant trial. Note that the sample size depends on the retention rates, the treatment effect τ_4 at the last visit, and Σ . Other parameters are specified in order to simulate the data. For the purpose of illustration, we set $\tau_4 = -4, -8$ or -12 , and three alternative covariance structures are also considered: 1) a compound symmetry (CS) structure: $\Sigma_{kk} = 45$ and $\Sigma_{kj} = 15$ if $k \neq j$; 2) an autoregressive (AR(1)) structure $\Sigma_{jk} = 45 \times 0.8^{|j-k|}$; 3) a Toeplitz structure $\Sigma_{jk} = 40 - 6|j - k|$. For CS and AR(1) structures, the analytic

Table 2: Calculated sample sizes and simulated power based on 10,000 replications for testing $H_0 : \tau_4 = 0$ in simulation 1: ^a The sample size is evaluated using equation (16). The size estimates from equations (15) and (17) are presented for the purpose of comparison; ^b The difference in sample size between two arms is 0 for even n and 1 for odd n ; ^c The nominal power is evaluated as $\Pr(t(f(n), \sqrt{n}|\tau_4|/\sqrt{\varpi_\tau(n_l)}) \geq t_{f(n),0.975})$ based on equation (12). The difference in nominal power estimated by equation (12) and equation (13) is $< 0.2\%$ in all cases.

true covariance structure	τ_4	estimated sample size ^a			total size n^b	power (%)	
		n_l	n_u^*	n_u		nominal ^c	simulated
1: Unstructured	-12	17	19.4	20.4	21	91.86	91.48
	-8	36	38.1	38.6	39	90.49	89.99
	-4	139	140.8	141.0	142	90.22	90.47
2: CS	-12	19	21.1	22.7	23	91.36	90.84
	-8	40	41.8	42.7	43	90.36	89.74
	-4	153	154.8	155.5	156	90.11	90.22
3: AR(1)	-12	17	19.6	20.7	21	91.53	91.10
	-8	36	38.4	39.0	39	90.24	89.75
	-4	140	141.8	142.1	143	90.21	90.87
4: Toeplitz	-12	15	17.6	18.7	19	91.81	91.86
	-8	32	33.9	34.3	35	90.81	90.31
	-4	122	123.7	123.9	124	90.04	89.71

expressions for the LDL decomposition provided in the appendix can be used in the calculation.

We calculate the sample size necessary to achieve 90% power at a two-sided significance level of $\alpha = 0.05$ using equation (16). In each case, 10000 datasets are simulated and analyzed using model (1) and KR variance estimate. There is about 95% chance that the simulated power lies within 0.6% (standard error $\approx \sqrt{0.9 * 0.1/10000} = 0.3\%$) of the true power. The result is summarized in Table 2. The difference between the simulated and nominal power is $< 1\%$ in all cases.

Empirical evidence indicates that the normal approximation based on equation (15) usually underestimates the sample size by at least 2. We try a slight variation of the sample size estimation procedure by increasing n_l by 2 (all other steps remain the same). This modification makes a difference mainly in small samples (i.e. $\tau_4 = -12$) in that the sample size estimate is reduced by 1. We still recommend the procedure described in Section 3.1 since it is generally desirable to use a conservative size estimate.

3.2.2. *Simulation 2.* This simulation illustrates that the power and sample size depend on the number of covariates. The covariates consist of the baseline outcome y_{i0} , treatment status g_i and a categorical prognostic factor A with three levels. Suppose the status of the prognostic factor is A_i for subject i , $y_{i0} \stackrel{i.i.d}{\sim} N(\eta_{A_i} + 17.9, 5.5^2)$, and

$$\begin{bmatrix} y_{i1} \\ y_{i2} \\ y_{i3} \\ y_{i4} \end{bmatrix} \sim N_4 \left(\begin{bmatrix} \eta_{A_i} + \mu_{i1} \\ \eta_{A_i} + \mu_{i2} \\ \eta_{A_i} + \mu_{i3} \\ \eta_{A_i} + \mu_{i4} \end{bmatrix}, \Sigma \right),$$

where $\eta_1 = 0$, $\eta_2 = -0.5$, $\eta_3 = 0.5$, and μ_{ij} 's are defined in simulation 1. We assume that the effect of the prognostic factor is constant over time, and that each subject is in level 1, 2 and 3 of the prognostic factor with probability 0.3, 0.4, and 0.3 respectively. Other assumptions are the same as that in simulation 1. In the MMRM analysis, the fixed effects include visit, $y_{i0} \times$ visit, $A \times$ visit and treatment \times visit interactions.

The result is displayed in Table 3. The required size in simulation 2 is larger than that in simulation 1 under the same assumption on the treatment effect and residual covariance matrix Σ due to the increase in the number of baseline

Table 3: Calculated sample sizes and simulated power based on 10,000 replications for testing $H_0 : \tau_4 = 0$ in simulation 2: ^a The sample size is evaluated using equation (16). The size estimates from equations (15) and (17) are presented for the purpose of comparison; ^b The difference in sample size between two arms is 0 for even n and 1 for odd n ; ^c The nominal power is evaluated as $\Pr(t(f(n), \sqrt{n}|\tau_4|/\sqrt{\varpi_\tau(n_l)}) \geq t_{f(n),0.975})$ based on equation (12).

true covariance structure	τ_4	estimated sample size ^a			total size n ^b	power (%)	
		n_l	n_u^*	n_u		nominal ^c	simulated
1: Unstructured	-12	20	21.9	23.5	24	91.51	91.67
	-8	39	40.7	41.2	42	90.70	90.91
	-4	142	143.4	143.4	144	90.13	89.65
2: CS	-12	21	23.8	27.1	28	92.30	92.97
	-8	42	44.5	45.9	46	90.27	90.01
	-4	155	157.5	158.2	159	90.17	89.99
3: AR(1)	-12	20	22.0	23.9	24	91.09	91.12
	-8	39	41.0	41.7	42	90.42	90.67
	-4	143	144.4	144.5	145	90.12	89.74
4: Toeplitz	-12	18	20.0	21.9	22	91.31	91.52
	-8	35	36.4	36.9	37	90.24	89.96
	-4	125	126.4	126.3	127	90.17	90.52

covariates. The difference between the simulated and nominal power is $< 1\%$ in all cases. Overall, the sample size is underestimated by the normal approximation approach given in equation (15) especially at $\tau_4 = -12$, and the proposed two-step sample size calculation method performs well when the d.f. in equation (16) is larger than $f(n_l) \geq 12$.

3.2.3. Simulation 3. So far, we assume the data are normally distributed. The third simulation is conducted to assess the performance of the proposed method for non-normal data in small to moderate samples ($\tau_4 = -8$ or -12). The data are simulated from the multivariate skew-normal distribution [23] and multivariate t distribution. For the purpose of comparison, we set the covariance matrix of (y_{i1}, \dots, y_{i4}) and the treatment effect at each visit to be identical to that in simulation 1. The nominal power and sample size estimates will be same as that in simulation 1.

In the multivariate t distribution, we generate the data as $y_{ij} = \mu_{ij} + \epsilon_{ij} / \sqrt{u_i/d}$, where $u_i \stackrel{i.i.d.}{\sim} \chi_d^2$ follows a chi-square distribution with d d.f., and $(\epsilon_{i1}, \dots, \epsilon_{i4})'$ is normally distributed with zero mean vector and covariance matrix $\frac{d-2}{d}\Sigma$. The covariance matrix of (y_{i1}, \dots, y_{i4}) is Σ . We set $d = 6$ or 10 . The skewness of y_{ij} is 0. The (excess) kurtosis of y_{ij} is $6/(d-4) = 3$ at $d = 6$ and 1 at $d = 10$.

The skew-normal distribution [23] can be used to model data that are mildly to moderately skewed. Let $R = D^{-1}\Sigma D^{-1}$ be the correlation matrix corresponding to Σ , and $R_2 = (1 - \kappa^2)^{-1}[aR - b\mathbf{1}_p\mathbf{1}'_p]$, where κ is a prespecified scalar value in the interval $(-1, 1)$, $a = 1 - 2\kappa^2/\pi$, $b = (1 - 2/\pi)\kappa^2$, and $D = \text{diag}(\Sigma_{11}, \dots, \Sigma_{pp})$. The data are generated as $y_{ij} = \mu_{ij} + \sqrt{\Sigma_{jj}/a}\epsilon_{ij}$, where $\epsilon_{ij} = \kappa e_i + \sqrt{1 - \kappa^2}\tilde{\epsilon}_{ij}$, $e_i \stackrel{i.i.d.}{\sim} N(0, 1)$, and $(\tilde{\epsilon}_{i1}, \dots, \tilde{\epsilon}_{ip})' \sim N_p(\mathbf{0}, R_2)$. The covariance matrix of $(y_{i1}, \dots, y_{i4})'$ is Σ . The marginal distribution of $\tilde{\epsilon}_{ij}$ is $N(0, 1)$, and the skewness of ϵ_{ij} (y_{ij} has the same skewness as ϵ_{ij}) ranges from -0.995 to 0.995 as κ changes from -1 to 1 . We set $\kappa = 0.8$ and 0.9 . No result is produced for the CS covariance structure at $\kappa = 0.9$ since the corresponding R_2 is not positive definite.

The result is presented in Table 4. In general, the type I error is close to the nominal 5% level except in few cases under the t distribution with 6 d.f., where the t test with KR adjusted variance provides slightly conservative control of the type I error. The performance on the power estimate is almost as good as that under the normality assumption in simulation 1. The simulated power is within 1.2% of the nominal power in all cases. Additional simulation (results not shown) is conducted under the skew-normal distribution at $\kappa = -0.8, -0.9$ and ± 0.95 , and different covariance matrices may be used to ensure that R_2 is positive definite. All simulations indicate that the proposed power and sample size estimation procedure is fairly robust to mild or moderate deviations from non-normality even when $n = 20$.

Table 4: Calculated sample sizes and simulated power based on 10,000 replications for testing $H_0 : \tau_4 = 0$ under multivariate t and skew-normal distributions in simulation 3: ^a The sample size and nominal power estimates are identical to that in simulation 1; ^b Type I error (Type1) is evaluated at $\tau_4 = 0$; ^c Simulated power (SIM) is evaluated at $\tau_4 = -8$ and -12 .

true covariance structure	τ_4	total size ^a n	nominal power ^a (%)	skew-normal distribution (%)				multivariate t distribution (%)			
				$\kappa = 0.8$		$\kappa = 0.9$		10 d.f.		6 d.f.	
				Type1 ^b	SIM ^c	Type1	SIM	Type1	SIM	Type1	SIM
1: Unstructured	-12	21	91.86	4.97	91.26	4.84	91.02	5.21	90.80	4.39	90.87
	-8	39	90.49	5.05	90.07	5.10	90.32	4.97	89.52	4.83	90.25
2: CS	-12	23	91.36	4.86	90.56	—	—	5.09	90.30	4.72	90.18
	-8	43	90.36	4.84	89.66	—	—	5.28	89.63	5.06	89.82
3: AR(1)	-12	21	91.53	4.89	90.91	4.99	90.72	5.13	90.54	4.35	90.45
	-8	39	90.24	5.06	89.92	4.93	89.91	4.92	89.28	4.86	90.09
4: Toeplitz	-12	19	91.81	5.24	90.80	4.93	90.88	4.91	91.07	5.11	91.13
	-8	35	90.81	5.01	90.26	4.82	90.84	5.30	89.69	4.99	90.36

4. Discussion

We derive the closed-form REML estimator and compare several variance estimators for fixed effects in MMRM when the missing data pattern is monotone. The bias in the ML and REML parameter estimators is assessed. For monotone missing data, the MMRM yields the same treatment effect as the cLDA [22], but it makes less assumption on the distribution of baseline outcomes. One referee raises concern over the terminology *MMRM* since there is no random effect in model (1). We partially agree with the referee, but this terminology is commonly used in the pharmaceutical industry [1, 2, 3].

A two-step sample size determination procedure is proposed for the KR adjusted t test for comparing two treatments in MMRM. Simulation demonstrates its good performance even in small samples. When the d.f. given in equation (16) is $f(n_i) < 12$, the power formula (12) still works reasonably well, but the required size can be overestimated. The main reason is because the normal approximation approach generally underestimates the required size and hence the d.f., leading to non-negligible inflation in $t_{f, 1-\alpha/2} + t_{f, p}$ and $\varpi_\tau(n)$ in small samples. When $f(n_i) < 12$, we would recommend using a numerical (e.g. bisection) method to find the size based on equation (12), which is the smallest integer at which the power reaches the pre-specified level. Simulation indicates that the proposed method is quite robust to mild or moderate deviations from non-normality. However, this robustness property may not hold under severe non-normality even for the one sample t test [24]. It is always prudent to run a simulation study to verify the power and sample size estimate especially when the data are highly non-normal or when the sample size is small. We assume balanced baselines across groups. If the baselines are imbalanced, we may replace equation (11) by

$$E(\Delta'_t S_{x_t}^{-1} \Delta_t) \approx \frac{\varpi_{\delta_t}(n) \varpi_{\pi_t}}{n} + \frac{D}{n\bar{\pi}_t - q - 1} \text{ and } E(V_{x_t}) = \frac{\varpi_{x_t}(n)}{n} + \frac{D}{n\bar{\pi}_t - q - 1},$$

where μ_d is the difference in mean of x_{bi} between two arms, Σ_x is the covariance matrix of x_{bi} and $D = \mu'_d \Sigma_x^{-1} \mu_d$.

In a companion paper, we investigate the power of treatment comparison in an unstratified randomized trial, and the power for testing treatment effect and treatment by stratum interaction in a stratified trial using analysis of covariance (ANCOVA). For the MMRM analysis of a stratified trial with h strata, we may replace equation (11) by

$$\varpi_{\delta_t}(n) = \frac{q^*}{n\bar{\pi}_t - q - 1} \text{ and } E(V_{x_t}) \approx \frac{\varpi_{x_t}(n)}{n} = \frac{\varpi_{\pi_t}[1 + \varpi_{\delta_t}(n)]}{n}, \tag{18}$$

where q^* is the dimension of covariates excluding intercept, treatment status, and pre-stratification factors, $(x_{i_1}, \dots, x_{i_{h-1}})$ are the indicator variables for strata, $\mathbf{x}_i = (1, x_{i_1}, \dots, x_{i_{h-1}}, x'_{bi}, g_i)'$ and $q = q^* + h + 1$. Equation (18) is derived by

assuming constant treatment effect, treatment allocation ratio and retention rate across strata.

Simulation 2 indicates that including the correction term ϖ_{δ_t} in ϖ_{x_t} in equations (12) and (13) and the last term in equations (15) and (16) can greatly improve the accuracy of the power and sample size estimate when q^* is large or n is relatively small. Covariate selection is critical in small trials. In the power calculation, $\Sigma = L\Lambda L'$ is the variance of the outcome unexplained by the covariates. Inclusion of important covariates can reduce the residual variance Σ , and increase the power of the analysis. However, if the covariates are unrelated or only weakly related to the outcome, both the precision of the treatment effect estimate (by equation (11)) and the d.f. in the t test may decrease, resulting in reduced power.

It can be challenging to specify Σ at the design stage of a trial. One strategy is to assume a structured covariance matrix that represents the best guess about the true covariance matrix to reduce the number of nuisance parameters in the sample size calculation [12], but the data are still analyzed by model (1). Furthermore, blinded size reassessment procedure may be used to re-estimate the variance parameters and adjust the size based on interim blinded data, in which the treatment effect is generally small and thus ignored in estimating the variance parameters [25].

We assume monotone missing pattern in the power and sample size calculation. In practice, the trials generally contain only a small amount of intermittent missing data. For example, in the anti-depressant trial analyzed in Section 2.4, only one out of 172 subjects has intermittent missing values. In the presence of non-monotone missing data, sample size can be calculated in a slightly conservative way by excluding subjects with intermittent missing data or by excluding data collected after the first missing visit. One needs to adjust the retention rates before applying the power and sample size formulae derived in section 3.1. If a large amount of intermittent missing data is expected, one may use simulation to find the appropriate sample size. The above approach provides an upper size bound, and a lower size bound can be obtained by pretending there are no intermittent missing data.

One limitation of the proposed method is that the data are assumed to be missing at random in the MMRM. Sensitivity analyses under the nonignorable missingness have become increasingly popular in new drug applications [26, 20, 9]. Sample size formulae for such sensitivity analysis may be obtained by using the analytical expressions for the MI treatment effect estimate and the associated variance derived in [19].

We have focused on the longitudinal continuous outcome. For other types of longitudinal outcomes (e.g. binary or ordinal), one may use the methods described in [27, 28] to compute the sample size for the tests based on the generalized estimating equations (GEE). However, these methods may not be suitable for small samples [29].

The analytic REML solution is useful in statistical computing. For example, we find the default REML estimation algorithm used by SAS Proc mixed fails to converge in about 5% monotone datasets at $n = 15$ in a simulation, but the REML estimate can be easily obtained using Lemma 1. For near-monotone datasets, the convergence of the REML algorithm may be accelerated by using the REML estimate excluding subjects with intermittent missing data as the initial parameter values. For non-monotone data, Liu [8] proposed a MEM algorithm to find the ML estimate. It is also possible to develop a MEM algorithm for the REML estimation, and the details will be presented elsewhere.

Acknowledgements

We would like to thank the associate editor and two referees for their helpful suggestions that improve the quality of the manuscript.

A. Appendix: technical details

Proof of Lemma 1: (a) Let U_k , $L_k = U_k^{-1}$, Λ_k and $\Sigma_k = L_k\Lambda_kL'_k$ denote respectively the leading $k \times k$ submatrix of U , $L = U^{-1}$, Λ and $\Sigma = L\Lambda L'$. Then $W_i = \tilde{I}_{r_i} \otimes \mathbf{x}_i$, $W'_i\Sigma_{r_i}^{-1}W_i = (\sum_{s=1}^{r_i} \tilde{\beta}'_s\tilde{\beta}_s/\sigma_s^2) \otimes (\mathbf{x}'_i\mathbf{x}_i)$, $\sum_i W'_i\Sigma_{r_i}^{-1}W_i = \sum_{s=1}^p [(\tilde{\beta}'_s\tilde{\beta}_s) \otimes (X'_{o_s}X_{o_s}/\sigma_s^2)]$, $(L' \otimes I_p)(\sum_i W'_i\Sigma_{r_i}^{-1}W_i)(L \otimes I_p) = V_w^{-1}$, where \tilde{I}_k is formed by the first k row of I_p , and $\tilde{\beta}_s = (-\beta_{s1}, \dots, -\beta_{s,s-1}, 1, 0, \dots, 0)'$. Thus (a) holds.

(b) We can prove (b) by showing $\sum_i \mathbf{y}_i \Sigma_{r_i}^{-1} \mathbf{y}_i = \sum_{j=1}^p (\sigma_j^{-2} \sum_{k=1}^{n_j} \underline{y}_{ij}^2)$, $\sum_i \log |\Sigma_{r_i}| - \log |\sum_i W_i' \Sigma_{r_i}^{-1} W_i| = \sum_{j=1}^p (n_j - q) \log(\sigma_j^2)$, $\sum_{i=1}^n W_i' \Sigma_{r_i}^{-1} \mathbf{y}_{io} = (U' \otimes I_p) \begin{bmatrix} \sigma_1^{-2} \sum_{i=1}^{n_1} \mathbf{x}'_i \underline{y}_{i1} \\ \dots \\ \sigma_p^{-2} \sum_{i=1}^{n_p} \mathbf{x}'_i \underline{y}_{ip} \end{bmatrix}$, and the last term of equation (2) equals $-\sum_{j=1}^p [\sigma_j^{-2} (\sum_{i=1}^{n_j} \mathbf{x}'_i \underline{y}_{ij})' (X'_{o_j} X_{o_j})^{-1} (\sum_{i=1}^{n_j} \mathbf{x}'_i \underline{y}_{ij})]$.

(c) Maximizing equation (5) yields $(\hat{\beta}_j, \hat{\sigma}_{j,re}^2)$'s, and $\underline{\alpha}_j$'s are evaluated at the REML estimator $(\hat{\beta}_j, \hat{\sigma}_{j,re}^2)$'s. We can prove $\hat{\theta}_{j,re} = \hat{\theta}_j$ by using

$$\begin{bmatrix} x'x & x'z \\ z'x & z'z \end{bmatrix}^{-1} = \begin{bmatrix} (x'x)^{-1} + \eta V_z \eta' & -\eta V_z \\ -V_z \eta' & V_z \end{bmatrix}, \tag{19}$$

where $V_z = [z'(I - x(x'x)^{-1}x')z]^{-1}$, and $\eta = (x'x)^{-1}x'z$.

(d) Let $\vec{Y}_{o_j} = (\underline{Y}_{o_{j1}}, \dots, \underline{Y}_{o_{j,j-1}}) = \vec{Y}_{o_j} U_{j-1}'$. By equation (3), $\underline{Y}_{o_{jk}} \sim N(X_{o_j} \underline{\alpha}_k, \sigma_k^2 I_{n_j})$, and $\underline{Y}_{o_{jk}}$'s ($k = 1, \dots, j - 1$) are independent. Lemma 1(e) holds since $E(\vec{Y}_{o_j}' Q_j \vec{Y}_{o_j}) = (n_j - q) \Lambda_{j-1}$ and $E(\vec{Y}_{o_j}' Q_j \vec{Y}_{o_j}) = (n_j - q) \Sigma_{j-1}$.

Proof of equation (6): A little algebra shows that $\hat{\alpha}_k = \sum_{t=1}^{k-1} \hat{\beta}_{kt} \hat{\alpha}_t + \hat{\alpha}_k$, $\tilde{\alpha}_k = \sum_{t=1}^{k-1} \beta_{kt} \tilde{\alpha}_t + \tilde{\alpha}_k$, $\hat{\alpha}_k - \tilde{\alpha}_k = d_k + e_k + \sum_{t=1}^{k-1} \beta_{kt} (\hat{\alpha}_t - \tilde{\alpha}_t)$, and

$$(\hat{\alpha}_1 - \tilde{\alpha}_1, \dots, \hat{\alpha}_k - \tilde{\alpha}_k)' = L_k (d_1 + e_1, \dots, d_k + e_k)', \tag{20}$$

where $\hat{\alpha}_{kt}^* = (X'_{o_k} X_{o_k})^{-1} X'_{o_k} \underline{Y}_{o_{kt}}$, $\zeta_k = L'_{k-1} (\hat{\beta}_k - \beta_k)$, $\tilde{\alpha}_k - \hat{\alpha}_k = (X'_{o_k} X_{o_k})^{-1} X'_{o_k} \vec{Y}_{o_k} (\hat{\beta}_k - \beta_k) = (\hat{\alpha}_{k1}^*, \dots, \hat{\alpha}_{kk-1}^*) \zeta_k$, $d_1 = e_1 = \mathbf{0}$, $d_k = \sum_{t=1}^{k-1} (\hat{\beta}_{kt} - \beta_{kt}) \tilde{\alpha}_t + (\hat{\alpha}_k - \tilde{\alpha}_k) = (\tilde{\alpha}_1 - \hat{\alpha}_{k1}^*, \dots, \tilde{\alpha}_{k-1} - \hat{\alpha}_{kk-1}^*) \zeta_k$ and $e_k = \sum_{t=1}^{k-1} (\hat{\beta}_{kt} - \beta_{kt}) (\hat{\alpha}_t - \tilde{\alpha}_t) = (d_1 + e_1, \dots, d_{k-1} + e_{k-1}) \zeta_k$ at $k \geq 2$. Note that d_k 's, e_k 's and $\tilde{\alpha}_k$'s are independent, $\text{var}(d_k) = \sum_{t=1}^{k-1} \sigma_t^2 \omega_{kt} D_{kt}$, and $E(d_k) = E(e_k) = \mathbf{0}$. Thus $\text{cov}(\tilde{\alpha}, \hat{\alpha} - \tilde{\alpha}) = \mathbf{0}$. Let $\epsilon_j = \hat{\alpha}_j - \tilde{\alpha}_j$ and $\epsilon = (\epsilon'_1, \dots, \epsilon'_p)'$. By equation (20), we have $\text{var}(\epsilon) = (L \otimes I_q) V_{de} (L \otimes I_q)'$ and $\text{var}(\hat{\alpha}_p - \tilde{\alpha}_p) = \sum_{j=2}^p l_{pj}^2 (V_{dj} + V_{ej})$, where $V_{de} = \text{diag}(\mathbf{0}, V_{d2} + V_{e2}, \dots, V_{dp} + V_{ep})$.

Proof of equation (9): Let $g_j = \sigma_j^2 l_{pj}^2$ and $c_j = E(\hat{g}_j) - g_j$. Then $E(\hat{\Phi}_p) - \Phi_p = \sum_j c_j (X'_{o_j} X_{o_j})^{-1}$. We evaluate c_j using the second-order Taylor series approximation

$$c_j \approx \frac{\partial g_j}{\partial \sigma_j^2} E(\hat{\sigma}_j^2 - \sigma_j^2) + \frac{1}{2} \frac{\partial^2 g_j}{\partial l_j \partial l_j} E(\hat{l}_j - l_j)^2 = \sigma_j^2 \text{var}(l_{pj}) - \frac{j-1}{n_j - q} \sigma_j^2 l_{pj}^2,$$

by noting $E(\hat{l}_j - l_j) = 0$, $E(\hat{l}_j - l_j)(\hat{\sigma}_j^2 - \sigma_j^2) = 0$ and $\partial^2 g_j / \partial \sigma_j^2 \partial \sigma_j^2 = 0$. Since $\partial(l_{pj}) / \partial \beta_k = l_{pk} (l_{1j}, \dots, l_{k-1,j})'$ [it is 0 if $j \geq k$] and $(l_{1j}, \dots, l_{k-1,j}) \text{var}(\beta_k) (l_{1j}, \dots, l_{k-1,j})' = \omega_{kj}$, we have $\text{var}(l_{pj}) = \sum_{k=j+1}^p l_{pk}^2 \omega_{kj}$ and equation (9) holds.

Analytic expressions for LDL decomposition of CS and AR(1) covariance

For an AR(1) covariance matrix ($\Sigma_{jk} = h\rho^{|j-k|}$), the LDL decomposition satisfies $l_{jk} = \rho^{|j-k|}$ if $k \leq j$, $l_{jk} = 0$ if $j > k$, $\sigma_1^2 = h$, and $\sigma_k^2 = h(1 - \rho^2)$ for $k \geq 2$.

For a CS covariance matrix ($\Sigma_{jk} = h\rho$ if $j \neq k$, h if $j = k$), we have $l_{jk} = 1$ if $k = j$, $\frac{\rho}{1+(k-1)\rho}$ if $k < j$, and 0 otherwise, and $\sigma_1^2 = h$, $\sigma_2^2 = h(1 - \rho^2)$, and $\sigma_k^2 = \sigma_{k-1}^2 [1 - (\frac{\rho}{1+(k-2)\rho})^2]$ for $k \geq 2$.

References

- Siddiqui O, Hung JHM, O'Neill R. MMRM vs. LOCF: A comprehensive comparison based on simulation study and 25 NDA data sets. *Journal of Biopharmaceutical Statistics* 2009; **19**:227–246.
- Siddiqui O. MMRM versus MI in dealing with missing data comparison based on 25 NDA data sets. *Journal of Biopharmaceutical Statistics* 2011; **21**:423 – 436.

3. Lu K, Luo X, Chen P. Sample size estimation for repeated measures analysis in randomized clinical trials with missing data. *The International Journal of Biostatistics* 2008; **4**:article 9.
4. Little RJ. In praise of simplicity not mathematistry! ten simple powerful ideas for the statistical scientist. *Journal of the American Statistical Association* 2013; **108**:359 – 369.
5. Anderson TW. Maximum likelihood estimates for a multivariate normal distribution when some observations are missing. *Journal of the American Statistical Association* 1957; **52**:200 – 203.
6. Tang Y. An efficient monotone data augmentation algorithm for Bayesian analysis of incomplete longitudinal data. *Statistics and Probability Letters* 2015; **104**:146 – 152.
7. Little R, Rubin DB. *Statistical Analysis with Missing Data*. John Wiley Sons: Hoboken, NJ, 2002.
8. Liu C. Efficient ML estimation of the multivariate normal distribution from incomplete data. *Journal of Multivariate Analysis* 1999; **69**:206 – 217.
9. Tang Y. An efficient monotone data augmentation algorithm for multiple imputation in a class of pattern mixture models. *Journal of Biopharmaceutical Statistics* doi:10.1080/10543406.2016.1167075 2016; .
10. Kenward MG, Roger JH. Small sample inference for fixed effects from restricted maximum likelihood. *Biometrics* 1997; **53**:983 – 997.
11. Mallinckrodt C, Roger J, Chuang-stein C, Molenberghs G, Lane P, Okelly M, Ratitch B, Xu L, Gilbert S, Mehrotrak D, *et al.*. Missing data: Turning guidance into action. *Statistics in Biopharmaceutical Research* 2013; **5**:369–382.
12. Lu K, Mehrotra DV, Liu G. Sample size determination for constrained longitudinal data analysis. *Statistics in Medicine* 2009; **28**:679 – 699.
13. Harville DA. Maximum-likelihood approaches to variance component estimation and to related problems. *Journal of the American Statistical Association* 1977; **72**:320 – 337.
14. Harville DA. Bayesian inference for variance components using only error contrasts. *Biometrika* 1974; **61**:383 – 385.
15. Verbyla AP. A conditional derivation of residual maximum likelihood. *Australian Journal of Statistics* 1990; **61**:227 – 230.
16. Cox DR, Snell EJ. A general definition of residuals. *Journal of the Royal Statistical Society: Series B* 1968; **30**:248 – 275.
17. Kenward MG, Roger JH. An improved approximation to the precision of fixed effects from restricted maximum likelihood. *Computational Statistics and Data Analysis* 2009; **53**:2583 – 2595.
18. Pase L, Salvani A. *Principles of Statistical Inference*. New Jersey: World Scientific, 1997.
19. Tang Y. On the multiple imputation variance estimator for control-based and delta-adjusted pattern mixture models. *Revision submitted* 2017; .
20. Tang Y. An efficient multiple imputation algorithm for control-based and delta-adjusted pattern mixture models using SAS. *Statistics in Biopharmaceutical Research* doi:10.1080/19466315.2016.1225595 2016; .
21. Liu GF, Lu K, Mogg R, Mallick M, Mehrotra DV. Should baseline be a covariate or dependent variable in analyses of change from baseline in clinical trials? *Statistics in medicine* 2009; **28**:2509 – 2530.
22. Lu K. On efficiency of constrained longitudinal data analysis versus longitudinal analysis of covariance. *Biometrics* 2010; **66**:891 – 896.
23. AZZALINI A, VALLE AD. The multivariate skew-normal distribution. *Biometrika* 1996; **83**:715–726.
24. Boos DD, Hughes-Oliver JM. How large does n have to be for z and t intervals? *The American Statistician* 2000; **54**:121–128.
25. Tang Y. Sample size estimation for negative binomial regression comparing rates of recurrent events with unequal follow-up time. *Journal of Biopharmaceutical Statistics* 2015; **25**:1100 – 1113.
26. Kenward MG. Controlled multiple imputation methods for sensitivity analyses in longitudinal clinical trials with dropout and protocol deviation. *Clinical Investigation* 2015; **5**:311–20.
27. Liu G, Liang K. Sample size calculations for studies with correlated observations. *Biometrics* 1997; **53**:937947.
28. Rochon J. Application of GEE procedures for sample size calculations in repeated measures experiments. *Statistics in Medicine* 1998; **17**:1643–1658.
29. Lu B, Preisser JS, Qaqish BF, Suchindran C, Bangdiwala SI, Wolfson M. A comparison of two bias-corrected covariance estimators for generalized estimating equations. *Biometrics* 2007; **63**:935–941.