

Reply to Comment on “Anomalous Edge State in a Non-Hermitian Lattice”

Tony E. Lee

Department of Physics, Indiana University Purdue University Indianapolis (IUPUI), Indianapolis, Indiana 46202, USA
(Dated: November 13, 2018)

The Comment by Xiong *et al.* (arXiv:1610.06275) criticizing my Letter [Phys. Rev. Lett. 116, 133903 (2016)] was rejected by Physical Review Letters. In this Reply, I show that all their claims are wrong.

Xiong *et al.* posted a Comment [1] that was rejected by Physical Review Letters. They dispute the conclusions of my Letter [2]. They make three claims: (i) the winding number, calculated using Berry phase, is not 1/2, (ii) the real spectrum of an open chain is a finite-size effect, and (iii) the breakdown of the bulk-boundary correspondence is not due to the defectiveness of the Hamiltonian. In this Reply, I show that they are wrong on every point.

I. WINDING NUMBER

In my Letter [2], I said that the winding number has a fractional value of 1/2 because k must sweep through 4π in order for $(\langle\sigma_x\rangle, \langle\sigma_z\rangle)$ to make one complete loop around the origin. Xiong *et al.* claim that according to the Berry phase, the winding number is not 1/2. Here, I show that the Berry phase does in fact verify that the winding number is 1/2.

Consider the momentum-space Hamiltonian H_k given in Eq. (1) of the Letter [2]. When H_k encircles an exceptional point, one needs to be careful when calculating the winding number, since the eigenvectors are 4π -periodic in k . The proper expression for the winding number in this case is

$$w = \frac{1}{2\pi}\gamma_B, \quad (1)$$

where γ_B is the Berry phase when k increases by 4π :

$$\gamma_B = i \int_0^{4\pi} dk \frac{\langle\langle u|\partial_k|u\rangle\rangle}{\langle\langle u|u\rangle\rangle}, \quad (2)$$

where $|u\rangle\rangle$ and $|u\rangle$ are left and right eigenvectors of H_k . It is important to integrate over 4π in k so that the eigenvectors return to their original values (see below). To account for this, there is a factor of 1/2 in Eq. (1) because the Brillouin zone is swept through twice.

In Ref. [3], it was shown analytically that $\gamma_B = \pi$ when H_k circles an exceptional point. Thus, the winding number is $w = 1/2$. In other words, one must sweep through the Brillouin zone *twice* in order to pick up a Berry phase of π .

Xiong *et al.* used an unphysical expression for the winding number:

$$w = \frac{1}{\pi}\tilde{\gamma}_B, \quad (3)$$

where $\tilde{\gamma}_B$ is the Berry phase when k increases by 2π :

$$\tilde{\gamma}_B = i \int_0^{2\pi} dk \frac{\langle\langle u|\partial_k|u\rangle\rangle}{\langle\langle u|u\rangle\rangle}. \quad (4)$$

It is well known that a Berry phase is physically meaningful only if the eigenvector makes a closed loop during the integration; otherwise, the Berry phase will be gauge dependent. When circling an exceptional point, the eigenvectors are 4π -periodic in k . Since Xiong *et al.* in Eq. (4) integrated over only 2π in k , it is no surprise that they find a gauge-dependent Berry phase and winding number. To get physically meaningful results, one needs to integrate over 4π in k so that the eigenvectors make a closed loop. That is what I did in Eq. (2).

Xiong *et al.*'s argument about the “demo model” is nonsense. Their Eq. (A6) is wrong and should be replaced with Eqs. (1) and (2) (using the usual inner product). If one decides to double the Brillouin zone to 4π , one should integrate over 4π in k . Furthermore, $A = 2$ since the Brillouin zone is swept through twice. Then one finds a winding number of 1, as expected. The important point is: *for Hermitian systems, Eqs. (1) and (3) give the same result, but when an exceptional point is involved, one needs to use Eq. (1) to get physically meaningful results.*

II. REAL SPECTRUM

In my Letter [2], I said that a chain with open boundary conditions has real eigenvalues for $|v| \geq \gamma/2$. Xiong *et al.* claim that this is a finite-size effect [1]. As evidence, they present their Fig. 1(e), which supposedly shows complex eigenvalues for large N . However, their result is an artifact due to flawed numerics.

As discussed in my Letter [2], the Hamiltonian H has two symmetries: chiral symmetry and \mathcal{PT} symmetry. Due to chiral symmetry, if E is an eigenvalue of H , $-E$ is also an eigenvalue. Due to \mathcal{PT} symmetry, if E is an eigenvalue of H , E^* is also an eigenvalue. As a result, when the eigenvalues are plotted in the complex plane, they should be symmetric around both the real and imaginary axes. This is a simple sanity check.

The eigenvalues calculated by Xiong *et al.* in their Fig. 1(e) do not obey these symmetries [1]: their eigenvalues are not symmetric around the real or imaginary axes. Thus, their calculation is flawed, and their observation of complex eigenvalues is nonsense. I do not know




FIG. 1. Eigenvalues for chains with open boundary conditions and different sizes N . The parameters are the same as Fig. 1(e) of Ref. [1]: $\gamma = 1$, $r = 0.5$, $v = 0.52$.

how they screwed up, but when I (correctly) calculate the eigenvalues, the eigenvalues are purely real, even for large N . The correct eigenvalues are shown in Fig. 1.

III. DEFECTIVENESS

In my Letter [2], I said that the bulk-boundary correspondence is broken because H is defective, such that there is only one $E = 0$ edge state, as opposed to the even number of edge states found in typical topological insulators. Xiong *et al.* claim that the breakdown of the bulk-boundary correspondence is not due to the defectiveness of H [1]. But my Letter showed analytically that the $E = 0$ eigenvalue is defective, leading to only one $E = 0$ edge state. This by itself shows the breakdown of the bulk-boundary correspondence.

Xiong *et al.* claim that the breakdown is instead due to the “bulk” (in-gap) states being localized on the edge [1]. But the in-gap states are localized on the edge precisely because H is defective. As mentioned in the Letter, when $v = \gamma/2$, the eigenvalues at $E = \pm r$ (corresponding to in-gap states) are highly defective [2]. One can show analytically that the corresponding eigenvectors are

$$\left(\pm 1 + \frac{i\gamma}{r}, \pm i + \frac{\gamma}{r}, i, 1, 0, 0, \dots\right)^T, \quad (5)$$

so all the in-gap states are localized on the left edge. What is amusing is that even if Xiong *et al.* are right about the in-gap states, the breakdown is still due to the defectiveness of H .

[1] Y. Xiong, T. Wang, X. Wang, and P. Tong, arXiv:1610.06275.

[2] T. E. Lee, Phys. Rev. Lett. **116**, 133903 (2016).

[3] F. Keck, H. J. Korsch, and S. Mossmann, J. Phys. A **36**, 2125 (2003).