A new expression for the Moore–Penrose inverse of a class of matrices

Xuefeng Xu

LSEC, Institute of Computational Mathematics and Scientific/Engineering Computing, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China. E-mail address: xuxuefeng@lsec.cc.ac.cn

Abstract

An expression for the Moore–Penrose inverse of a matrix of the form M = XNY, where X and Y are nonsingular, has been recently established by Castro-González et al. [1, Theorem 2.2]. The expression plays an essential role in developing explicit expressions for the Moore–Penrose inverse of a two-by-two block matrix. In this paper, we present a new expression for the Moore– Penrose inverse of this class of matrices, which improves the result in [1].

Keywords: Moore–Penrose inverse; Matrix product; Orthogonal projector

1. Introduction

We first introduce some notations and concepts which are frequently used in the subsequent content. Let \mathbb{N}^+ and \mathbb{C} denote the set of all positive integers and the field of complex numbers, respectively. Let $\mathbb{C}^{m \times n}$ be the set of all $m \times n$ complex matrices. The identity matrix of order nis denoted by I_n or I when its size is clear in the context. For a matrix $A \in \mathbb{C}^{m \times n}$, A^* denotes the conjugate transpose of A. We denote by $\mathcal{R}(A)$ and $\mathcal{N}(A)$ the range and null space of A, respectively, namely, $\mathcal{R}(A) := \{y \in \mathbb{C}^m : y = Ax, x \in \mathbb{C}^n\}$ and $\mathcal{N}(A) := \{x \in \mathbb{C}^n : Ax = 0\}$. The *Moore–Penrose inverse* of A is denoted by A^{\dagger} , which is defined as the unique matrix $Z \in \mathbb{C}^{n \times m}$ satisfying the following equations:

(a)
$$AZA = A$$
, (b) $ZAZ = Z$, (c) $(AZ)^* = AZ$, (d) $(ZA)^* = ZA$.

The symbols $E_A := I - AA^{\dagger}$ and $F_A := I - A^{\dagger}A$ stand for the orthogonal projectors onto $\mathcal{N}(A^*)$ and $\mathcal{N}(A)$, respectively. A matrix $Z \in \mathbb{C}^{n \times m}$ is called an *inner inverse* of A if it satisfies the equality (a).

For a matrix $M \in \mathbb{C}^{m \times n}$ which can be decomposed as M = XNY, where $X \in \mathbb{C}^{m \times m}$ and $Y \in \mathbb{C}^{n \times n}$ are nonsingular, the equality $M^{\dagger} = Y^{-1}N^{\dagger}X^{-1}$ may fail. Several conditions validating $M^{\dagger} = Y^{-1}N^{\dagger}X^{-1}$ are presented in [2]. Recently, Castro-González et al. [1] obtained an explicit expression for M^{\dagger} , provided that $XE_N = E_N$ and $F_NY = F_N$. More concretely, it is proved by Castro-González et al. [1, Theorem 2.2] that

$$M^{\dagger} = (I + L_0^*)(I + L_0 L_0^*)^{-1} Y^{-1} N^{\dagger} X^{-1} (I + R_0^* R_0)^{-1} (I + R_0^*), \qquad (1.1)$$

where $R_0 := E_N(I - X^{-1})$ and $L_0 := (I - Y^{-1})F_N$. The expression (1.1) is a crucial result in [1], which can be exploited to establish explicit expressions for the Moore–Penrose inverse of a two-by-two block matrix. Assume that the singular value decomposition (SVD) of $N \in \mathbb{C}^{m \times n}$ is $N = U \begin{pmatrix} \Sigma & 0 \\ 0 & 0 \end{pmatrix} V^*$, where $\Sigma \in \mathbb{C}^{r \times r}$ is a diagonal matrix with positive diagonal entries, r is the rank of N, and both $U \in \mathbb{C}^{m \times m}$ and $V \in \mathbb{C}^{n \times n}$ are unitary. Let $X \in \mathbb{C}^{m \times m}$ and $Y \in \mathbb{C}^{n \times n}$. We now give two assumptions \mathbf{A}_1 and \mathbf{A}_2 as follows:

$$\mathbf{A}_1: \ X = U \begin{pmatrix} X_1 & 0 \\ X_2 & X_4 \end{pmatrix} U^*,$$

where $X_1 \in \mathbb{C}^{r \times r}$, $X_2 \in \mathbb{C}^{(m-r) \times r}$, and $X_4 \in \mathbb{C}^{(m-r) \times (m-r)}$;

$$\mathbf{A}_2: \ Y = V \begin{pmatrix} Y_1 & Y_3 \\ 0 & Y_4 \end{pmatrix} V^*,$$

where $Y_1 \in \mathbb{C}^{r \times r}$, $Y_3 \in \mathbb{C}^{r \times (n-r)}$, and $Y_4 \in \mathbb{C}^{(n-r) \times (n-r)}$.

In this paper, we further investigate explicit expressions for the Moore–Penrose inverse of this class of matrices. A new expression under weakened conditions for M^{\dagger} is derived, which has enhanced the expression (1.1). More specifically, if the assumptions \mathbf{A}_1 and \mathbf{A}_2 are satisfied, then we have

$$M^{\dagger} = (I + L^{*})(I + LL^{*})^{-1}N^{\dagger}N(Y^{-1}N^{\dagger}X^{-1})NN^{\dagger}(I + R^{*}R)^{-1}(I + R^{*}), \qquad (1.2)$$

where $R := X E_N X^{-1} (E_N - I)$ and $L := (F_N - I) Y^{-1} F_N Y$.

The rest of this paper is organized as follows. In Section 2, we first introduce a useful lemma which gives an explicit expression for the Moore–Penrose inverse of a two-by-two block matrix, and then give some specific conditions to validate \mathbf{A}_1 and \mathbf{A}_2 . In Section 3, we present a new and improved expression (i.e., (1.2)) for M^{\dagger} based on the assumptions \mathbf{A}_1 and \mathbf{A}_2 .

2. Preliminaries

In this section, we first introduce a useful lemma, which provides an explicit expression for the Moore–Penrose inverse of a two-by-two block matrix; see [3]. It is worth mentioning that some improved results of this lemma can be found in [1].

Lemma 2.1. Let M be a two-by-two block matrix as the form $M = \begin{pmatrix} A & C \\ B & D \end{pmatrix}$. Assume that $\mathcal{R}(B^*) \subseteq \mathcal{R}(A^*), \ \mathcal{R}(C) \subseteq \mathcal{R}(A), \ and \ D - BA^{\dagger}C = 0$. Then M^{\dagger} can be given by

$$M^{\dagger} = \begin{pmatrix} I \\ (A^{\dagger}C)^* \end{pmatrix} \Psi A^{\dagger} \Phi \begin{pmatrix} I & (BA^{\dagger})^* \end{pmatrix},$$

where $\Phi = \left(I + (BA^{\dagger})^* BA^{\dagger}\right)^{-1}$ and $\Psi = \left(I + A^{\dagger} C (A^{\dagger} C)^*\right)^{-1}$.

Next, we give several specific conditions to guarantee the assumptions A_1 and A_2 .

Lemma 2.2. Let $N \in \mathbb{C}^{m \times n}$ have the singular value decomposition $N = U \begin{pmatrix} \Sigma & 0 \\ 0 & 0 \end{pmatrix} V^*$, where $\Sigma \in \mathbb{C}^{r \times r}$ is a diagonal matrix with positive diagonal entries, r is the rank of N, and both

 $U \in \mathbb{C}^{m \times m}$ and $V \in \mathbb{C}^{n \times n}$ are unitary. Let $X \in \mathbb{C}^{m \times m}$ be an arbitrary matrix. Suppose that one of the following conditions holds:

 $\mathbf{C}_1: NN^*X \text{ is normal};$

- \mathbf{C}_2 : For any $0 \neq c_1 \in \mathbb{C}$, there exists $k_1 \in \mathbb{N}^+$ such that $(NN^*X)^{k_1} = c_1 NN^{\dagger}$;
- \mathbf{C}_3 : For any $0 \neq c_2 \in \mathbb{C}$ and $\ell \in \mathbb{N}^+$, there exists $k_2 \in \mathbb{N}^+$ such that $(NN^*X)^{k_2} = c_2(NN^*)^{\ell}$;
- $\mathbf{C}_4: XE_N \text{ is normal};$
- \mathbf{C}_5 : For any $0 \neq c_3 \in \mathbb{C}$, there exists $k_3 \in \mathbb{N}^+$ such that $(XE_N)^{k_3} = c_3E_N$;
- $\mathbf{C}_6: NN^{\dagger}XE_N = 0;$
- \mathbf{C}_7 : There exists $k_4 \in \mathbb{N}^+$ such that $(NN^*)^{k_4}XE_N = 0$.

Then X must be of the form

$$X = U \begin{pmatrix} X_1 & 0 \\ X_2 & X_4 \end{pmatrix} U^*,$$

where $X_1 \in \mathbb{C}^{r \times r}$, $X_2 \in \mathbb{C}^{(m-r) \times r}$, and $X_4 \in \mathbb{C}^{(m-r) \times (m-r)}$.

Proof. Based on the SVD of N, the expressions of N^{\dagger} and E_N can be given by

$$N^{\dagger} = V \begin{pmatrix} \Sigma^{-1} & 0 \\ 0 & 0 \end{pmatrix} U^*$$
 and $E_N = U \begin{pmatrix} 0 & 0 \\ 0 & I \end{pmatrix} U^*.$

Partition U^*XU as $U^*XU = \begin{pmatrix} X_1 & X_3 \\ X_2 & X_4 \end{pmatrix}$, where $X_1 \in \mathbb{C}^{r \times r}$, $X_2 \in \mathbb{C}^{(m-r) \times r}$, $X_3 \in \mathbb{C}^{r \times (m-r)}$, and $X_4 \in \mathbb{C}^{(m-r) \times (m-r)}$. Then $X = U \begin{pmatrix} X_1 & X_3 \\ X_2 & X_4 \end{pmatrix} U^*$.

(i) The condition \mathbf{C}_1 states that

$$NN^*X = U \begin{pmatrix} \Sigma^2 X_1 & \Sigma^2 X_3 \\ 0 & 0 \end{pmatrix} U^*$$

is normal, which yields that $\Sigma^2 X_1$ is normal and $\Sigma^2 X_3 = 0$. It follows from the non-singularity of Σ that $X_3 = 0$.

(ii) We have known that $NN^*X = U\begin{pmatrix} \Sigma^2 X_1 & \Sigma^2 X_3 \\ 0 & 0 \end{pmatrix} U^*$. Then, for any $k_1 \in \mathbb{N}^+$, we have

$$(NN^*X)^{k_1} = U \begin{pmatrix} (\Sigma^2 X_1)^{k_1} & (\Sigma^2 X_1)^{k_1 - 1} \Sigma^2 X_3 \\ 0 & 0 \end{pmatrix} U^*$$

In addition, it is easy to see that

$$c_1 N N^{\dagger} = U \begin{pmatrix} c_1 I & 0 \\ 0 & 0 \end{pmatrix} U^*.$$

Hence, \mathbf{C}_2 implies that $(\Sigma^2 X_1)^{k_1} = c_1 I$ and $(\Sigma^2 X_1)^{k_1 - 1} \Sigma^2 X_3 = 0$. Due to the facts that $c_1 \neq 0$ and Σ is nonsingular, it follows that X_1 is nonsingular and $X_3 = 0$.

(iii) Direct calculation yields

$$(NN^*X)^{k_2} = U \begin{pmatrix} (\Sigma^2 X_1)^{k_2} & (\Sigma^2 X_1)^{k_2 - 1} \Sigma^2 X_3 \\ 0 & 0 \end{pmatrix} U^*,$$
$$c_2(NN^*)^{\ell} = U \begin{pmatrix} c_2 \Sigma^{2\ell} & 0 \\ 0 & 0 \end{pmatrix} U^*.$$

Because $c_2 \neq 0$ and Σ is nonsingular, we deduce from \mathbf{C}_3 that $\Sigma^2 X_1$ is nonsingular and $(\Sigma^2 X_1)^{k_2-1} \Sigma^2 X_3 = 0$. Hence, $X_3 = 0$.

(iv) Straightforward calculation shows

$$XE_N = U \begin{pmatrix} 0 & X_3 \\ 0 & X_4 \end{pmatrix} U^*.$$

If XE_N is normal, then we get that X_4 is normal and $X_3 = 0$.

(v) Direct computation yields

$$(XE_N)^{k_3} = U \begin{pmatrix} 0 & X_3 X_4^{k_3 - 1} \\ 0 & X_4^{k_3} \end{pmatrix} U^*.$$

It follows from C_5 that $X_4^{k_3} = c_3 I$ and $X_3 X_4^{k_3-1} = 0$. By $c_3 \neq 0$, we derive that X_4 is nonsingular. Hence, we obtain from $X_3 X_4^{k_3-1} = 0$ that $X_3 = 0$.

(vi) It is easy to compute that

$$NN^{\dagger}XE_N = U \begin{pmatrix} 0 & X_3 \\ 0 & 0 \end{pmatrix} U^*.$$

Therefore, $NN^{\dagger}XE_N = 0$ if and only if $X_3 = 0$.

(vii) Direct calculation yields

$$(NN^*)^{k_4} X E_N = U \begin{pmatrix} 0 & \Sigma^{2k_4} X_3 \\ 0 & 0 \end{pmatrix} U^*.$$

Due to the fact that Σ is nonsingular, it follows that $(NN^*)^{k_4}XE_N = 0$ is equivalent to $X_3 = 0$.

Consequently, if one of the conditions C_1 - C_7 holds, then X must be of the from

$$X = U \begin{pmatrix} X_1 & 0 \\ X_2 & X_4 \end{pmatrix} U^*,$$

which completes the proof.

Analogously, we can prove the following lemma. Its detailed proof is omitted due to limited space.

Lemma 2.3. Let $Y \in \mathbb{C}^{n \times n}$ and let $N \in \mathbb{C}^{m \times n}$ be the same as in Lemma 2.2. Assume that one of the following conditions holds:

 \mathbf{C}'_1 : YN*N is normal;

 $\begin{aligned} \mathbf{C}_{2}': For \ any \ 0 \neq c_{1}' \in \mathbb{C}, \ there \ exists \ k_{1}' \in \mathbb{N}^{+} \ such \ that \ (YN^{*}N)^{k_{1}'} &= c_{1}'N^{\dagger}N; \\ \mathbf{C}_{3}': For \ any \ 0 \neq c_{2}' \in \mathbb{C} \ and \ \ell' \in \mathbb{N}^{+}, \ there \ exists \ k_{2}' \in \mathbb{N}^{+} \ such \ that \ (YN^{*}N)^{k_{2}'} &= c_{2}'(N^{*}N)^{\ell'}; \\ \mathbf{C}_{4}': F_{N}Y \ is \ normal; \\ \mathbf{C}_{5}': For \ any \ 0 \neq c_{3}' \in \mathbb{C}, \ there \ exists \ k_{3}' \in \mathbb{N}^{+} \ such \ that \ (F_{N}Y)^{k_{3}'} &= c_{3}'F_{N}; \\ \mathbf{C}_{6}': F_{N}YN^{\dagger}N &= 0; \\ \mathbf{C}_{7}': There \ exists \ k_{4}' \in \mathbb{N}^{+} \ such \ that \ F_{N}Y(N^{*}N)^{k_{4}'} &= 0. \end{aligned}$ Then Y must be of the form

$$Y = V \begin{pmatrix} Y_1 & Y_3 \\ 0 & Y_4 \end{pmatrix} V^*,$$

where $Y_1 \in \mathbb{C}^{r \times r}$, $Y_3 \in \mathbb{C}^{r \times (n-r)}$, and $Y_4 \in \mathbb{C}^{(n-r) \times (n-r)}$.

Remark 2.4. Notice that Lemma 2.2 (resp., Lemma 2.3) does not need the non-singularity of X (resp., Y). In addition, the reader can give other conditions to ensure that A_1 and A_2 hold.

3. Main results

In order to prove our main result, we first consider explicit expressions for $(XN)^{\dagger}$ and $(NY)^{\dagger}$. The following theorem provides two applicable formulas for M_1^{\dagger} and M_2^{\dagger} , where $M_1 = XN$ and $M_2 = NY$.

Theorem 3.1. Let $N \in \mathbb{C}^{m \times n}$, $X \in \mathbb{C}^{m \times m}$, $Y \in \mathbb{C}^{n \times n}$, $M_1 = XN$, and $M_2 = NY$. Suppose that X and Y are nonsingular.

(1) If the assumption A_1 holds, then

$$M_1^{\dagger} = N^{\dagger} X^{-1} N N^{\dagger} (I + R^* R)^{-1} (I + R^*), \qquad (3.1)$$

where $R = XE_N X^{-1}(E_N - I)$. (2) If the assumption \mathbf{A}_2 holds, then

$$M_2^{\dagger} = (I + L^*)(I + LL^*)^{-1}N^{\dagger}NY^{-1}N^{\dagger},$$

where $L = (F_N - I)Y^{-1}F_NY$.

Proof. (1) The assumption \mathbf{A}_1 reads $X = U \begin{pmatrix} X_1 & 0 \\ X_2 & X_4 \end{pmatrix} U^*$, where $X_1 \in \mathbb{C}^{r \times r}$ and r is the rank of N. It follows from the non-singularity of X that both $X_1 \in \mathbb{C}^{r \times r}$ and $X_4 \in \mathbb{C}^{(m-r) \times (m-r)}$ are nonsingular. We define $R := X E_N X^{-1}(E_N - I)$. By simple computation, we can get

$$R = U \begin{pmatrix} 0 & 0 \\ X_2 X_1^{-1} & 0 \end{pmatrix} U^* = U \begin{pmatrix} 0 & 0 \\ G & 0 \end{pmatrix} U^*,$$

where $G := X_2 X_1^{-1}$. Because U and V are unitary matrices and

$$M_1 = XN = U \begin{pmatrix} X_1 \Sigma & 0 \\ X_2 \Sigma & 0 \end{pmatrix} V^*,$$

we obtain

$$M_1^{\dagger} = V \begin{pmatrix} X_1 \Sigma & 0 \\ X_2 \Sigma & 0 \end{pmatrix}^{\dagger} U^*.$$

Note that $X_1\Sigma$ is nonsingular. Using Lemma 2.1, we obtain

$$\begin{pmatrix} X_1 \Sigma & 0 \\ X_2 \Sigma & 0 \end{pmatrix}^{\dagger} = \begin{pmatrix} I \\ 0 \end{pmatrix} \Sigma^{-1} X_1^{-1} (I + G^* G)^{-1} \begin{pmatrix} I & G^* \end{pmatrix}.$$

Hence,

$$M_1^{\dagger} = V \begin{pmatrix} I \\ 0 \end{pmatrix} \Sigma^{-1} X_1^{-1} (I + G^* G)^{-1} \begin{pmatrix} I & G^* \end{pmatrix} U^*.$$

Straightforward computation yields

$$\begin{split} N^{\dagger}X^{-1} &= V \begin{pmatrix} I \\ 0 \end{pmatrix} \Sigma^{-1}X_1^{-1} \begin{pmatrix} I & 0 \end{pmatrix} U^*, \\ NN^{\dagger}(I+R^*R)^{-1}(I+R^*) &= U \begin{pmatrix} I \\ 0 \end{pmatrix} (I+G^*G)^{-1} \begin{pmatrix} I & G^* \end{pmatrix} U^*. \end{split}$$

It can be easily seen that $M_1^{\dagger} = N^{\dagger} X^{-1} N N^{\dagger} (I + R^* R)^{-1} (I + R^*)$ holds.

(2) Applying the formula (3.1) to the matrix Y^*N^* , we obtain

$$(M_2^*)^{\dagger} = (N^*)^{\dagger} (Y^*)^{-1} N^* (N^*)^{\dagger} (I + \widehat{R}^* \widehat{R})^{-1} (I + \widehat{R}^*),$$

where

$$\widehat{R} = Y^* E_{N^*} (Y^*)^{-1} (E_{N^*} - I) = Y^* (F_N)^* (Y^{-1})^* (F_N - I)^*.$$

We define $L := (F_N - I)Y^{-1}F_NY$. Then,

$$(M_2^{\dagger})^* = (M_2^*)^{\dagger} = (N^{\dagger})^* (Y^{-1})^* N^* (N^{\dagger})^* (I + LL^*)^{-1} (I + L).$$

Therefore, we drive that $M_2^{\dagger} = (I + L^*)(I + LL^*)^{-1}N^{\dagger}NY^{-1}N^{\dagger}$.

Using Theorem 3.1, we can easily obtain the following expressions for the orthogonal projectors onto $\mathcal{R}(M_1)$ and $\mathcal{R}(M_2^*)$.

Corollary 3.2. Under the same conditions as in Theorem 3.1. (1) If the assumption A_1 is valid, then

$$M_1 M_1^{\dagger} = (I+R)NN^{\dagger}(I+R^*R)^{-1}(I+R^*).$$
(3.2)

(2) If the assumption \mathbf{A}_2 is valid, then

$$M_2^{\dagger}M_2 = (I + L^*)(I + LL^*)^{-1}N^{\dagger}N(I + L).$$
(3.3)

Proof. According to the equality (3.1), it follows that

$$M_1 M_1^{\dagger} = X N N^{\dagger} X^{-1} N N^{\dagger} (I + R^* R)^{-1} (I + R^*).$$
(3.4)

Notice that

$$(I+R)NN^{\dagger} = \left(I - NN^{\dagger} + XNN^{\dagger}X^{-1}NN^{\dagger}\right)NN^{\dagger} = XNN^{\dagger}X^{-1}NN^{\dagger}.$$
 (3.5)

Inserting (3.5) into (3.4) gives $M_1 M_1^{\dagger} = (I+R)NN^{\dagger}(I+R^*R)^{-1}(I+R^*)$. Similarly, we can prove the equality (3.3).

Based on the expressions (3.2) and (3.3) for orthogonal projectors $M_1 M_1^{\dagger}$ and $M_2^{\dagger} M_2$, we can establish the following main result.

Theorem 3.3. Let $N \in \mathbb{C}^{m \times n}$, $X \in \mathbb{C}^{m \times m}$, $Y \in \mathbb{C}^{n \times n}$, and M = XNY. Assume that X and Y are nonsingular. If the assumptions \mathbf{A}_1 and \mathbf{A}_2 are satisfied, then

$$M^{\dagger} = (I + L^{*})(I + LL^{*})^{-1}N^{\dagger}N(Y^{-1}N^{\dagger}X^{-1})NN^{\dagger}(I + R^{*}R)^{-1}(I + R^{*}),$$

where $R = XE_N X^{-1}(E_N - I)$ and $L = (F_N - I)Y^{-1}F_N Y$.

Proof. Note that $Y^{-1}N^{\dagger}X^{-1}$ is an inner inverse of M. Then we have

$$M^{\dagger} = M^{\dagger} M (Y^{-1} N^{\dagger} X^{-1}) M M^{\dagger}.$$

Let $M_1 = XN$ and $M_2 = NY$. We claim that $MM^{\dagger} = M_1M_1^{\dagger}$ and $M^{\dagger}M = M_2^{\dagger}M_2$. In fact, it is clear that MM^{\dagger} is the orthogonal projector onto $\mathcal{R}(M)$. Because Y is nonsingular and $M = M_1Y$, it follows that $\mathcal{R}(M) = \mathcal{R}(M_1)$. Hence, MM^{\dagger} is also an orthogonal projector onto $\mathcal{R}(M_1)$. Using the uniqueness of orthogonal projectors, we get that $MM^{\dagger} = M_1M_1^{\dagger}$. Similarly, we can verify that $M^{\dagger}M = M_2^{\dagger}M_2$. Therefore, we have

$$M^{\dagger} = M_2^{\dagger} M_2 (Y^{-1} N^{\dagger} X^{-1}) M_1 M_1^{\dagger}.$$

Under the assumptions of this theorem, by Corollary 3.2, we have

$$M^{\dagger} = (I + L^{*})(I + LL^{*})^{-1}N^{\dagger}N(I + L)Y^{-1}N^{\dagger}X^{-1}(I + R)NN^{\dagger}(I + R^{*}R)^{-1}(I + R^{*}).$$

Using $R = XE_N X^{-1}(E_N - I)$ and $L = (F_N - I)Y^{-1}F_N Y$, we obtain

$$\begin{split} (I+L)Y^{-1}N^{\dagger}X^{-1}(I+R) &= Y^{-1}N^{\dagger}X^{-1} + Y^{-1}N^{\dagger}X^{-1}R + LY^{-1}N^{\dagger}X^{-1} + LY^{-1}N^{\dagger}X^{-1}R \\ &= Y^{-1}N^{\dagger}X^{-1}, \end{split}$$

where we have applied the facts that $N^{\dagger}E_N = 0$ and $F_N N^{\dagger} = 0$. Consequently, we infer that

$$M^{\dagger} = (I + L^{*})(I + LL^{*})^{-1}N^{\dagger}N(Y^{-1}N^{\dagger}X^{-1})NN^{\dagger}(I + R^{*}R)^{-1}(I + R^{*}).$$

This completes the proof.

Corollary 3.4. Under the same conditions as in Theorem 3.3. If both XE_N and F_NY are Hermitian, then

$$M^{\dagger} = (I + L^{*})(I + LL^{*})^{-1}Y^{-1}N^{\dagger}X^{-1}(I + R^{*}R)^{-1}(I + R^{*}).$$
(3.6)

Proof. Because XE_N and F_NY are Hermitian, by Lemmas 2.2 and 2.3, the assumptions A_1 and A_2 are clearly satisfied. An application of Theorem 3.3 gives

$$M^{\dagger} = (I + L^{*})(I + LL^{*})^{-1}N^{\dagger}N(Y^{-1}N^{\dagger}X^{-1})NN^{\dagger}(I + R^{*}R)^{-1}(I + R^{*}).$$
(3.7)

Due to both XE_N and F_NY are Hermitian, it follows that $XE_N = E_NX^*$ and $F_NY = Y^*F_N$. Then, $E_N(X^*)^{-1} = X^{-1}E_N$ and $(Y^*)^{-1}F_N = F_NY^{-1}$. Notice that

$$N^{\dagger}N(Y^{-1}N^{\dagger}X^{-1})NN^{\dagger} = Y^{-1}N^{\dagger}X^{-1} - F_NY^{-1}N^{\dagger}X^{-1} - Y^{-1}N^{\dagger}X^{-1}E_N + F_NY^{-1}N^{\dagger}X^{-1}E_N.$$

Using $(Y^*)^{-1}F_N = F_NY^{-1}$ and $F_NN^{\dagger} = 0$, we can derive that $F_NY^{-1}N^{\dagger}X^{-1} = 0$. By $E_N(X^*)^{-1} = X^{-1}E_N$ and $N^{\dagger}E_N = 0$, we have $Y^{-1}N^{\dagger}X^{-1}E_N = 0$. Consequently,

$$N^{\dagger}N(Y^{-1}N^{\dagger}X^{-1})NN^{\dagger} = Y^{-1}N^{\dagger}X^{-1}.$$
(3.8)

By substituting (3.8) into (3.7), we obtain the formula (3.6).

Remark 3.5. If $XE_N = E_N$ and $F_NY = F_N$, the conditions in Corollary 3.4 are obviously satisfied because E_N and F_N are orthogonal projectors. In this case,

$$R = XE_N X^{-1}(E_N - I) = E_N (X^{-1}E_N - X^{-1}) = E_N (E_N - X^{-1}) = E_N (I - X^{-1}) = R_0,$$

$$L = (F_N - I)Y^{-1}F_N Y = (F_N Y^{-1} - Y^{-1})F_N = (F_N - Y^{-1})F_N = (I - Y^{-1})F_N = L_0,$$

where R_0 and L_0 are defined as in expression (1.1). Therefore, Corollary 3.4 has extended the expression (1.1).

Acknowledgements

The author would like to thank Professor Chen-Song Zhang for his helpful suggestions. This work was supported partially by the National Natural Science Foundation of China Grant 91430215 and 91530323.

References

[1] N. Castro-González, M. F. Martínez-Serrano, J. Robles, Expressions for the Moore–Penrose inverse of block matrices involving the Schur complement, Linear Algebra Appl. 471 (2015) 353–368.

[2] Y. Tian, The Moore–Penrose inverses of $m \times n$ block matrices and their applications, Linear Algebra Appl. 283 (1998) 35–60.

[3] B. Noble, A method for computing the generalized inverse of a matrix, SIAM J. Numer. Anal. 3 (1966) 582–584.