MULTIPLE SOLUTIONS FOR A CLASS OF NONHOMOGENEOUS FRACTIONAL SCHRÖDINGER EQUATIONS IN \mathbb{R}^N

VINCENZO AMBROSIO AND HICHEM HAJAIEJ

ABSTRACT. This paper is concerned with the following fractional Schrödinger equation

$$\begin{cases} (-\Delta)^s u + u = k(x)f(u) + h(x) \text{ in } \mathbb{R}^N\\ u \in H^s(\mathbb{R}^N), u > 0 \text{ in } \mathbb{R}^N, \end{cases}$$

where $s \in (0,1)$, N > 2s, $(-\Delta)^s$ is the fractional Laplacian, k is a bounded positive function, $h \in L^2(\mathbb{R}^N)$, $h \not\equiv 0$ is nonnegative and f is either asymptotically linear or superlinear at infinity. By using the *s*-harmonic extension technique and suitable variational methods, we prove the existence of at least two positive solutions for the problem under consideration, provided that $|h|_2$ is sufficiently small.

1. INTRODUCTION

In this paper we deal with the existence of positive solutions for the following nonlinear fractional equation

$$\begin{cases} (-\Delta)^s u + u = k(x)f(u) + h(x) \text{ in } \mathbb{R}^N\\ u \in H^s(\mathbb{R}^N), u > 0 \text{ in } \mathbb{R}^N, \end{cases}$$
(1.1)

with $s \in (0,1)$, N > 2s, k is a bounded positive function, $h \in L^2(\mathbb{R}^N)$, $h \ge 0$, $h \not\equiv 0$, and the nonlinearity $f : \mathbb{R} \to \mathbb{R}$ is a smooth function which can be either asymptotically linear or superlinear at infinity. Here, $(-\Delta)^s$ is the fractional Laplacian which can be defined for $u : \mathbb{R}^N \to \mathbb{R}$ belonging to the Schwartz space $\mathcal{S}(\mathbb{R}^N)$ of rapidly decaying C^{∞} -functions in \mathbb{R}^N , by setting

$$(-\Delta)^{s}u(x) = -\frac{C(N,s)}{2} \int_{\mathbb{R}^{N}} \frac{u(x+y) + u(x-y) - 2u(x)}{|y|^{N+2s}} dy \quad (x \in \mathbb{R}^{N})$$

where

$$C(N,s) = \left(\int_{\mathbb{R}^N} \frac{1 - \cos(x_1)}{|x|^{N+2s}} \, dx\right)^{-1};$$

see for instance [17].

A basic motivation to study (1.1), comes from the nonlinear fractional Schrödinger equation

$$i\frac{\partial\psi}{\partial t} = (-\Delta)^s \psi + V(x)\psi - g(x,|\psi|) \quad (t,x) \in \mathbb{R} \times \mathbb{R}^N,$$
(1.2)

when we are interested in standing wave solutions, namely solutions of the form $\psi(t, x) = u(x)e^{-ict}$. Indeed, it is easily observed that a function $\psi(t, x)$ of this form satisfies (1.2) if and only if u is a solution of

$$(-\Delta)^{s}u + V(x)u = g(x, u) \text{ in } \mathbb{R}^{N}$$

$$(1.3)$$

with V(x) = 1 + c and g(x, u) = k(x)f(u) + h(x).

The equation (1.2) has been proposed by Laskin in [32, 33] and it is fundamental in quantum mechanics, because it appears in problems involving nonlinear optics, plasma physics and condensed

²⁰¹⁰ Mathematics Subject Classification. 35A15, 35J60, 35R11, 45G05.

Key words and phrases. Fractional Laplacian; mountain pass theorem; extension method; positive solutions.

matter physics. When s = 1, the equation (1.3) becomes the classical Schrödinger equation

$$-\Delta u + V(x)u = g(x, u)$$
 in \mathbb{R}^N

which has been widely investigated by many authors, particularly on the existence of ground state solutions, positive solutions, sign-changing solutions and multiplicity of standing wave solutions. Since we cannot review the huge bibliography here, we just mention the works [6, 7, 8, 23, 40].

Recently, a considerable attention has been focused on the study of problems involving fractional and non-local operators. This interest is motivated by both the theoretical research and the large number of applications such as phase transitions, flames propagation, chemical reaction in liquids, population dynamics, American options in finance, crystal dislocation, obstacle problems, soft thin films, semipermeable membranes, conservation laws, quasi-geostrophic flows, and so on.

Since the current literature is really too wide to attempt any reasonable account here, we derive the interested reader to [17, 37], where an elementary introduction to this topic is given.

Concerning the fractional Schrödinger equation (1.3), in the last decade, many several existence and multiplicity results have been obtained by using different variational methods.

Felmer et al. [21] dealt with the existence and the symmetry of positive solutions for (1.3) with V(x) = 1 and involving a superlinear nonlinearity g(x, u) satisfying the Ambrosetti-Rabinowitz condition. Secchi [41] obtained via minimization on Nehari manifold, the existence of ground state solutions to (1.3) when the nonlinearity is superlinear and subcritical, and the potential V(x)satisfies suitable assumptions as $|x| \to \infty$. Chang and Wang [12] (see also [3]) investigated the existence of nontrivial solutions to (1.3) with V(x) = 1, q is autonomous and verifies Berestycki-Lions type assumptions. Frank et al. [24] showed uniqueness and nondegeneracy of ground state solutions for (1.3) with V(x) = 1 and $q(x, u) = |u|^q u$ for all admissible exponents q. Cho et al. [14] obtained the existence of standing waves using the method of concentration-compactness by studying the associated constrained minimization problem, and showed the orbital stability of standing waves which are the minimizers of the associate variational problem. Pucci et al. [38] established via Mountain Pass Theorem and Ekeland variational principle, the existence of multiple solutions for a Kirchhoff fractional Schrödinger equation involving a nonlinearity satisfying the Ambrosetti-Rabinowitz condition, a positive potential V(x) verifying suitable assumptions, and in presence of a perturbation term. Figueiredo and Siciliano [22] established a multiplicity result for a fractional Schrödinger equation via Ljusternick- Schnirelmann and Morse theory. We also mention the papers [1, 2, 4, 13, 14, 20, 29, 30, 31, 36, 42, 44] where further results related to (1.3) are given.

After an accurate bibliographic review, we have realised that there are only few papers concerning the existence and the multiplicity of solutions for nonhomogeneous problems in non-local setting [16, 43, 47], and this is surprising, because, in the classic framework, such type of problems have been extensively investigated by many authors [5, 11, 25, 45, 49, 50].

Strongly motivated by this fact, the purpose of this work is to investigate the existence and the multiplicity of positive solutions for the nonhomogeneous equation (1.1), under the effect of a small perturbation $h \in L^2(\mathbb{R}^N)$, and requiring suitable assumptions on the nonlinearity f. More precisely, along the paper, we assume that f satisfies the following conditions: (f1) $f \in C^1(\mathbb{R}, \mathbb{R}^+)$, f(0) = 0 and f(t) = 0 for $t \leq 0$; (f2) $\lim_{t \to 0^+} \frac{f(t)}{t} = 0$; (f3) there exists $p \in (1, \frac{N+2s}{N-2s})$ such that $\lim_{t \to +\infty} \frac{f(t)}{t^p} = 0$; (f4) there exists $l \in (0, \infty]$ such that $\lim_{t \to +\infty} \frac{f(t)}{t} = l$. Let us note that (f_1) - (f_3) , yield that for any $\varepsilon > 0$ there exists $C_{\varepsilon} > 0$ such that

$$|F(t)| \le \frac{\varepsilon}{2}t^2 + \frac{1}{p+1}C_{\varepsilon}|t|^{p+1} \text{ for all } t \in \mathbb{R},$$
(1.4)

while (f4) implies that f is asymptotically linear if $l < \infty$, or superlinear when $l = \infty$. Then, under the above assumptions, we are able to prove the existence of weak solutions to (1.1). We recall that the precise meaning of solution to (1.1) is the following:

Definition 1.1. We say that $u \in H^{s}(\mathbb{R}^{N})$ is a weak solution to (1.1) if verifies the following condition

$$\iint_{\mathbb{R}^{2N}} \frac{(u(x) - u(y))}{|x - y|^{N+2s}} (\varphi(x) - \varphi(y)) \, dx \, dy + \int_{\mathbb{R}^N} u\varphi \, dx = \int_{\mathbb{R}^N} [k(x)f(u) + h(x)]\varphi \, dx$$

for any $\varphi \in H^{s}(\mathbb{R}^{N})$.

Here

$$H^{s}(\mathbb{R}^{N}) = \left\{ u \in L^{2}(\mathbb{R}^{N}) : \frac{|u(x) - u(y)|}{|x - y|^{\frac{N+2s}{2}}} \in L^{2}(\mathbb{R}^{2N}) \right\}.$$

is the fractional Sobolev space [17, 37]; see Section 2 below.

Due to the presence of the fractional Laplacian, which is a nonlocal operator, we prefer to analyze (1.1) by using the s-harmonic extension method [10]. This procedure is commonly used to study fractional problems, since it allows us to write a given nonlocal equation in a local way and to adapt known techniques of the Calculus of Variations to these kind of problems.

Hence, instead of (1.1), we can consider the following degenerate elliptic equation with a nonlinear Neumann boundary condition

$$\begin{cases} \operatorname{div}(y^{1-2s}\nabla U) = 0 & \operatorname{in} \mathbb{R}^{N+1}_+ = \{(x,y) \in \mathbb{R}^N \times \mathbb{R} : y > 0\} \\ \frac{\partial U}{\partial \nu^{1-2s}} = \kappa_s [-U + k(x)f(U) + h(x)] & \operatorname{on} \partial \mathbb{R}^{N+1}_+ = \mathbb{R}^N \times \{0\} \end{cases},$$
(1.5)

where κ_s is a suitable constant; see [10]. Taking into account this fact, we are able to resemble some variational techniques developed in the papers [27, 35, 46, 48], dealing with asymptotically or superlinear classical problems, by introducing the following functional

$$I(U) = \frac{1}{2} \Big[\kappa_s^{-1} \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U|^2 \, dx \, dy + \int_{\mathbb{R}^N} U(x,0)^2 \, dx \Big] - \int_{\mathbb{R}^N} k(x) F(U(x,0)) \, dx - \int_{\mathbb{R}^N} h(x) U(x,0) \, dx + \int_{\mathbb{R}^N} h(x) \, dx + \int_{\mathbb{R}^N}$$

defined on the weight Sobolev space $X^{s}(\mathbb{R}^{N+1}_{+})$ consisting of functions $U: \mathbb{R}^{N+1}_{+} \to \mathbb{R}$ such that

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U|^2 \, dx \, dy + \int_{\mathbb{R}^N} U(x,0)^2 \, dx < \infty$$

Clearly, this functional simplification creates some additional difficulties to overcome such as, for instance, some weighted embedding are needed (see Theorem 2.2) to obtain some convergence results (compare with Lemma 3.1). Moreover, the arguments used in [27, 35] to prove the non-existence of solutions for certain eigenvalues problems, have to be handled carefully in order to take care the trace of the involved functions (see Lemma 3.5).

Now, we state our first main result concerning the existence of positive solutions to (1.1) in the asymptotically linear case, that is $l < \infty$.

Theorem 1.1. Let $s \in (0,1)$ and N > 2s. Assume that $h \in L^2(\mathbb{R}^N)$, $h(x) \ge 0$, $h(x) \ne 0$ and $k \in L^{\infty}(\mathbb{R}^N, \mathbb{R}_+)$ verifies the following condition:

(K) there exists $R_0 > 0$ such that

$$\sup\left\{\frac{f(t)}{t}: t > 0\right\} < \inf\left\{\frac{1}{k(x)}: |x| \ge R_0\right\}.$$
(1.6)

Let us suppose that f verifies (f1)-(f4) and $\mu^* \in (l, \infty)$ where

$$\mu^* = \inf\left\{\int_{\mathbb{R}^N} (|(-\Delta)^{\frac{s}{2}}u|^2 + u^2) \, dx : u \in H^s(\mathbb{R}^N), \int_{\mathbb{R}^N} k(x)u^2 \, dx = 1\right\}.$$
(1.7)

Let us assume that

$$|h|_{2} < m := \max_{t \ge 0} \left[\left(\frac{1}{2} - \frac{\varepsilon}{2} |k|_{L^{\infty}(\mathbb{R}^{N})} \right) t - \frac{C_{\varepsilon}}{p+1} S_{*}^{p+1} t^{p} |k|_{L^{\infty}(\mathbb{R}^{N})} \right],$$
(1.8)

where $\varepsilon \in (0, |k|_{L^{\infty}(\mathbb{R}^N)}^{-1})$ is fixed and S_* is the best Sobolev constant of the embedding $H^s(\mathbb{R}^N) \subset L^{2^*_s}(\mathbb{R}^N)$. Then, the problem (1.1) possesses at least two positive solutions $u_1, u_2 \in H^s(\mathbb{R}^N)$ with the property that $E(u_1) < 0 < E(u_2)$. Here $E : H^s(\mathbb{R}^N) \to \mathbb{R}$ is the energy functional associated to (1.1), that is

$$E(u) = \frac{1}{2} \int_{\mathbb{R}^N} (|(-\Delta)^{\frac{s}{2}}u|^2 + u^2) \, dx - \int_{\mathbb{R}^N} k(x)F(u) \, dx - \int_{\mathbb{R}^N} h(x)u \, dx.$$

Remark 1.1. The assumption on the size of h is a necessary condition to find a solution to (1.1). In fact, proceeding as in [11], one can obtain a non-existence result to (1.1) when $|h|_2$ is sufficiently large.

The proof of the above theorem goes as follows: under the assumption $l < \infty$, we first use the Ekeland variational principle to prove that for $|h|_2$ small enough, there exists a positive solution to (1.5) such that $I(U_0) < 0$. Then, we use a variant of Mountain Pass Theorem [19], to find a Cerami sequence which converges strongly in $X^s(\mathbb{R}^{N+1}_+)$ to a solution U_1 of (1.5) with $I(U_1) > 0$. Clearly, these two solutions U_0 and U_1 are different.

Our second result deals with the existence of positive solutions to (1.1) in the superlinear case $l = \infty$.

Theorem 1.2. Let $s \in (0,1)$ and N > 2s. Assume that f verifies (f1)-(f4) with $l = \infty$. We also suppose that $k(x) \equiv 1$, $h \in C^1(\mathbb{R}^N) \cap L^2(\mathbb{R}^N)$ is a radial function such that $h(x) \ge 0$, $h(x) \ne 0$ and $(H) \ x \cdot \nabla h(x) \in L^1(\mathbb{R}^N) \cap L^\infty(\mathbb{R}^N)$ and

$$x \cdot \nabla h(x) \ge 0$$
 for all $x \in \mathbb{R}^N$.

Let us assume that

$$|h|_{2} < m_{1} := \max_{t \ge 0} \left[\left(\frac{1}{2} - \frac{\varepsilon}{2} \right) t - \frac{C_{\varepsilon}}{p+1} S_{*}^{p+1} t^{p} \right],$$

$$(1.9)$$

where $\varepsilon \in (0,1)$ is fixed. Then, (1.1) admits two positive solutions $u_3, u_4 \in H^s_r(\mathbb{R}^N)$ such that $E(u_3) < 0 < E(u_4)$.

Due to the presence of radial functions k(x) = 1 and h = h(|x|), we work in the subspace $X_r^s(\mathbb{R}^{N+1}_+)$ of the weight space $X^s(\mathbb{R}^{N+1}_+)$, involving the functions which are radial with respect to $x \in \mathbb{R}^N$. We point out that the methods used to study the asymptotically linear case do not work any more. Indeed, to prove that a Palais-Smale sequence converges to a second solution different from the first one, we have to use the concentration-compactness principle which seems very hard to apply without requiring further assumptions on k(x) and f(t).

This time, we use the compactness of $X_r^s(\mathbb{R}^{N+1})$ into $L^q(\mathbb{R}^N)$ for any $q \in (2, 2_s^*)$, and the Ekeland principle, to get a first solution to (1.5) with negative energy, provided that $|h|_2$ is sufficiently small. The existence of a second solution with positive energy is obtained by combining a generic result due to Jeanjean [26], which allows us to prove the existence of bounded Palais-Smale sequences for parametrized functionals, the Pohozaev identity for the fractional Laplacian and the assumption (H), which guarantee the existence of a bounded Palais-Smale sequence for I, which converges to a radial positive solution to (1.5).

MULTIPLE SOLUTIONS

The plan of the paper is the following: in section 2 we give some preliminaries which will be useful along the paper. In section 3 we consider the asymptotically linear case and we prove the existence of two positive solutions via mountain pass theorem. In section 4 we study the superlinear case, and we provide the proof of Theorem 1.2. Finally, as applications of our results, we present some concrete examples.

2. Preliminaries

In this section, we briefly recall some properties of the fractional Sobolev spaces, and we introduce some notations which we will used along the paper.

For any $s \in (0,1)$, we define $\mathcal{D}^{s,2}(\mathbb{R}^N)$ as the completion of $C_0^{\infty}(\mathbb{R}^N)$ with respect to

$$[u]^{2} = \iint_{\mathbb{R}^{2N}} \frac{|u(x) - u(y)|^{2}}{|x - y|^{N + 2s}} \, dx \, dy = |(-\Delta)^{\frac{s}{2}} u|^{2}_{L^{2}(\mathbb{R}^{N})},$$

that is

$$\mathcal{D}^{s,2}(\mathbb{R}^N) = \left\{ u \in L^{2^*_s}(\mathbb{R}^N) : [u] < \infty \right\}.$$

Now, let us introduce the fractional Sobolev space

$$H^{s}(\mathbb{R}^{N}) = \left\{ u \in L^{2}(\mathbb{R}^{N}) : \frac{|u(x) - u(y)|}{|x - y|^{\frac{N+2s}{2}}} \in L^{2}(\mathbb{R}^{2N}) \right\}$$

endowed with the natural norm

$$||u||_{H^s(\mathbb{R}^N)} = \sqrt{[u]^2 + |u|^2_{L^2(\mathbb{R}^N)}}$$

For the convenience of the reader we recall the following fundamental embeddings:

Theorem 2.1. [17] Let $s \in (0, 1)$ and N > 2s. Then there exists a sharp constant $S_* = S(N, s) > 0$ such that for any $u \in H^s(\mathbb{R}^N)$

$$u|_{L^{2^*_s}(\mathbb{R}^N)}^2 \le S_*[u]^2.$$
(2.1)

Moreover $H^s(\mathbb{R}^N)$ is continuously embedded in $L^q(\mathbb{R}^N)$ for any $q \in [2, 2_s^*]$ and compactly in $L^q_{loc}(\mathbb{R}^N)$ for any $q \in [2, 2_s^*)$.

We also define the space of radial functions in $H^{s}(\mathbb{R}^{N})$

$$H_r^s(\mathbb{R}^N) = \left\{ u \in H^s(\mathbb{R}^N) : u(x) = u(|x|) \right\}.$$

Related to this space, the following compactness result due to Lions [34] holds:

Theorem 2.2. [34] Let $s \in (0,1)$ and $N \geq 2$. Then $H^s_r(\mathbb{R}^N)$ is compactly in $L^q(\mathbb{R}^N)$ for any $q \in (2, 2^*_s)$.

We also state the following useful result obtained in [12]:

Lemma 2.1. [12] Let $(X, \|\cdot\|)$ be a Banach space such that X is embedded respectively continuously and compactly into $L^q(\mathbb{R}^N)$ for $q \in [q_1, q_2]$ and $q \in (q_1, q_2)$, where $q_1, q_2 \in (0, \infty)$. Assume that $(u_n) \subset X, u : \mathbb{R}^N \to \mathbb{R}$ is a measurable function and $P \in C(\mathbb{R}, \mathbb{R})$ is such that

(i)
$$\lim_{|t| \to 0} \frac{P(t)}{|t|^{q_1}} = 0,$$

 $P(t)$

(*ii*)
$$\lim_{|t| \to \infty} \frac{1}{|t|^{q_2}} = 0,$$

(*iii*) $\sup_{t \to \infty} \|u_t\|_{q_2} \le 0.$

(*iii*)
$$\sup_{n\in\mathbb{N}} \|u_n\|_X < \infty$$
,

(iv)
$$\lim_{n \to \infty} P(u_n(x)) = u(x)$$
 for a.e. $x \in \mathbb{R}^N$.

Then, up to a subsequence, we have

$$\lim_{n \to \infty} \|P(u_n) - u\|_{L^1(\mathbb{R}^N)} = 0.$$

Now, let us denote by $\mathcal{D}(\mathbb{R}^{N+1}_+, y^{1-2s})$ the completion of $C_0^{\infty}(\overline{\mathbb{R}^{N+1}_+})$ under the norm

$$||U||_{\mathcal{D}(\mathbb{R}^{N+1}_+, y^{1-2s})}^2 = \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U|^2 \, dx dy.$$

It is known [10] that for any $U \in \mathcal{D}(\mathbb{R}^{N+1}_+, y^{1-2s})$, its trace U(x, 0) belongs $\mathcal{D}^{s,2}(\mathbb{R}^N)$ and that it is possible to define a trace continous map

$$[U(\cdot,0)] \le C \|U\|_{\mathcal{D}(\mathbb{R}^{N+1}_{+},y^{1-2s})}$$
(2.2)

By combining (2.1) and (2.2), we can derive the following Sobolev inequality

$$|U(\cdot,0)|_{L^{2^*}_s(\mathbb{R}^N)} \le C ||U||_{\mathcal{D}(\mathbb{R}^{N+1}_+,y^{1-2s})}$$
(2.3)

for any $U \in \mathcal{D}(\mathbb{R}^{N+1}_+, y^{1-2s})$.

Then, as proved in [10], for any $u \in \mathcal{D}^{s,2}(\mathbb{R}^N)$ there exists $U \in \mathcal{D}(\mathbb{R}^{N+1}_+, y^{1-2s})$, called the *s*-harmonic extension of *u*, such that

$$\begin{cases} \operatorname{div}(y^{1-2s}\nabla U) = 0 & \operatorname{in} \mathbb{R}^{N+1}_+\\ U(x,0) = u(x) & \operatorname{on} \mathbb{R}^N \end{cases}$$
(2.4)

Moreover,

$$\frac{\partial U}{\partial \nu^{1-2s}} = -\lim_{y \to 0^+} y^{1-2s} \frac{\partial U}{\partial y} = (-\Delta)^s u(x)$$

and

$$\int_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U|^2 \, dx \, dy = \kappa_s [u]^2_{H^s(\mathbb{R}^N)}.$$

Therefore, we can reformulate (1.1) in a local way, and we can investigate the following extended problem in \mathbb{R}^{N+1}_+

$$\begin{cases} \operatorname{div}(y^{1-2s}\nabla U) = 0 & \operatorname{in} \mathbb{R}^{N+1}_+\\ \frac{\partial U}{\partial \nu^{1-2s}} = \kappa_s[-U(x,0) + k(x)f(U(x,0)) + h(x)] & \operatorname{on} \mathbb{R}^N \end{cases}.$$

$$(2.5)$$

Qualitatively, the result of [10] states that one can localize the fractional Laplacian by adding an additional variable. This argument is fundamental to apply known variational methods. At this point, we introduce the following functional space

$$X^{s}(\mathbb{R}^{N+1}_{+}) = \{ U \in \mathcal{D}(\mathbb{R}^{N+1}_{+}, y^{1-2s}) : \int_{\mathbb{R}^{N}} |U(x,0)|^{2} \, dx < \infty \}$$

endowed with the norm

$$\|U\|_{X^{s}(\mathbb{R}^{N+1}_{+})}^{2} = \int_{\mathbb{R}^{N+1}_{+}} y^{1-2s} |\nabla U|^{2} \, dx \, dy + \int_{\mathbb{R}^{N}} |U(x,0)|^{2} \, dx$$

We recall that $X^{s}(\mathbb{R}^{N+1}_{+})$ is locally compactly embedded in the weight space $L^{2}(\mathbb{R}^{N+1}_{+}, y^{1-2s})$ endowed with the norm

$$||U||_{L^2(\mathbb{R}^{N+1}_+, y^{1-2s})} = \int_{\mathbb{R}^{N+1}_+} y^{1-2s} |U|^2 \, dx dy.$$

More precisely, we have

Lemma 2.2. [18] Let R > 0 and let \mathcal{T} be a subset of $\mathcal{D}(\mathbb{R}^{N+1}_+, y^{1-2s})$ such that

$$\sup_{U \in \mathcal{T}} \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U|^2 \, dx \, dy < \infty$$

Then \mathcal{T} is pre-compact in $L^2(B_R^+, y^{1-2s})$, where $B_R^+ = \{(x, y) \in \mathbb{R}^{N+1}_+ : |(x, y)| < R\}$. There exists a constant $C_0 > 0$ such that for all $U \in \mathcal{D}(\mathbb{R}^{N+1}_+, y^{1-2s})$ it holds

$$\left(\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |U|^{2\gamma} \, dx dy\right)^{\frac{1}{2\gamma}} \le C_0 \left(\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U|^2 \, dx dy\right)^{\frac{1}{2\gamma}}$$

where $\gamma = 1 + \frac{2}{N-2s}$.

Remark 2.1. With abuse of notation, we will denote by u the trace of a function $U \in \mathcal{D}(\mathbb{R}^{N+1}_+, y^{1-2s})$. Moreover, we denote by $|u|_p$ the L^p -norm of a function u belonging to $L^p(\mathbb{R}^N)$.

In what follows, for simplicity, we will omit the constant κ_s appearing in the extended problem (2.5).

3. Asymptotically linear case: proof of Theorem 1.1

In this section we discuss the existence of positive solutions to (1.1) under the assumption that f is asymptotically linear. Taking into account the results presented in Section 2, we can consider the following degenerate elliptic problem

$$\begin{cases} \operatorname{div}(y^{1-2s}\nabla U) = 0 & \operatorname{in} \mathbb{R}^{N+1}_+\\ \frac{\partial U}{\partial \nu^{1-2s}} = -u + k(x)f(u) + h(x) & \operatorname{on} \mathbb{R}^N \end{cases}$$
(3.1)

where k(x) is a bounded positive function, $h \in L^2(\mathbb{R}^N)$, $h \ge 0$ $(h \ne 0)$ and f satisfies (f1)-(f4) with $l < +\infty$.

Since the proof of Theorem 1.1, consists of several steps, we first collect some useful lemmas. We begin proving the following result:

Lemma 3.1. Suppose that (f_1) - (f_4) with $l < +\infty$ hold. Let $h \in L^2(\mathbb{R}^N)$, k satisfies (1.6), and $\{U_n\} \subset X^s(\mathbb{R}^{N+1}_+)$ be a bounded (PS) sequence of I. Then $\{U_n\}$ has a strongly convergent subsequence in $X^s(\mathbb{R}^{N+1}_+)$.

Proof. Firstly, we show that for any $\varepsilon > 0$, there exist $R(\varepsilon) > R_0$ (where R_0 is given by (K)) and $n(\varepsilon) > 0$ such that

$$\iint_{\mathbb{R}^{N+1}_+ \setminus B^+_R} y^{1-2s} |\nabla U_n|^2 \, dx \, dy + \int_{\mathbb{R}^N \setminus B_R} u_n^2 \, dx \le \varepsilon, \quad \forall R \ge R(\varepsilon) \text{ and } n \ge n(\varepsilon). \tag{3.2}$$

Let $\Psi_R \in C^{\infty}(\mathbb{R}^{N+1}_+)$ be a smooth function such that $0 \leq \Psi_R \leq 1$,

$$\Psi_R(x,y) = \begin{cases} 0 & (x,y) \in B_{\frac{R}{2}}^+ \\ 1 & (x,y) \notin B_{R}^+. \end{cases}$$
(3.3)

and

$$|\nabla \Psi_R(x,y)| \le \frac{C}{R} \text{ for all } (x,y) \in \mathbb{R}^{N+1}_+$$
(3.4)

for some positive constant C independent of R.

Then, we can observe that for any $U \in X^{s}(\mathbb{R}^{N+1}_{+})$ and all $R \geq 1$, there exists a constant $C_{1} > 0$ such that

$$\|\Psi_R U\|_{X^s(\mathbb{R}^{N+1}_+)} \le C_1 \|U\|_{X^s(\mathbb{R}^{N+1}_+)}.$$

Indeed, by using Young inequality and Lemma 2.2, we can see that

$$\begin{split} &\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla (U\Psi_R)|^2 \, dx dy + \int_{\mathbb{R}^N} |u\psi_R|^2 \, dx \\ &\leq 2 \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U|^2 \Psi_R^2 \, dx dy + 2 \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla \Psi_R|^2 U^2 \, dx dy + \int_{\mathbb{R}^N} |u|^2 \, dx \\ &\leq 2 \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U|^2 \, dx dy + \frac{2C}{R^2} \iint_{B^+_R \setminus B^+_R} y^{1-2s} U^2 \, dx dy + \int_{\mathbb{R}^N} |u|^2 \, dx \\ &\leq 2 \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U|^2 \, dx dy + \int_{\mathbb{R}^N} |u|^2 \, dx + \frac{2C}{R^2} \Big(\iint_{B^+_R \setminus B^+_R} y^{1-2s} |\nabla U|^{2\gamma} \, dx dy \Big)^{\frac{1}{\gamma}} \Big(\iint_{B^+_R \setminus B^+_R} y^{1-2s} \, dx dy \Big)^{\frac{\gamma-1}{\gamma}} \\ &\leq 2 (1+C) \, \|U\|^2_{X^s(\mathbb{R}^{N+1}_+)} \leq C_1 \|U\|^2_{X^s(\mathbb{R}^{N+1}_+)}, \end{split}$$

where we used the facts

$$\iint_{B_R^+ \setminus B_{\frac{R}{2}}^+} y^{1-2s} \, dx \, dy \le C R^{N+2-2s} \text{ and } \frac{\gamma-1}{\gamma} = \frac{2}{N+2-2s}.$$

Since $I'(U_n) \to 0$ as $n \to \infty$ and $\{U_n\}$ is bounded in $X^s(\mathbb{R}^{N+1}_+)$, we know that, for any $\varepsilon > 0$, there exists $n(\varepsilon) > 0$ such that

$$\langle I'(U_n), \Psi_R U_n \rangle \le C_1 \|I'(U_n)\| \|U_n\|_{X^s(\mathbb{R}^{N+1}_+)} \le \frac{\varepsilon}{4}, \text{ for } n \ge n(\varepsilon).$$

Equivalently, for all $n \ge n(\varepsilon)$, we get

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U_n|^2 \Psi_R \, dx \, dy + \int_{\mathbb{R}^N} u_n^2 \psi_R \, dx$$

$$\leq \int_{\mathbb{R}^N} (k(x)f(u_n) + h(x)) u_n \psi_R \, dx - \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} \nabla U_n \nabla \psi_R U_n \, dx \, dy + \frac{\varepsilon}{4}. \tag{3.5}$$

Now, by using (f1) and (1.6), we obtain that there exists $0 < \theta < 1$ such that

$$k(x)f(u_n)u_n \le \theta u_n^2 \quad \text{ for } |x| \ge R_0.$$
(3.6)

Since $h \in L^2(\mathbb{R}^N)$ and $||U_n||_{X^s(\mathbb{R}^{N+1}_+)} \leq C$ for some constant C > 0, it follows from (3.3) there exists $R(\varepsilon) > R_0$ such that

$$\int_{\mathbb{R}^N} h(x) u_n \psi_R \, dx \le |h(x)\psi_R|_2 |u_n|_2 \le \frac{\varepsilon}{4}, \quad \text{for } R \ge R(\varepsilon).$$
(3.7)

Due to the boundedness of $\{U_n\}$ in $X^s(\mathbb{R}^{N+1}_+)$, we may assume, up to a subsequence, that there exists $U \in X^s(\mathbb{R}^{N+1}_+)$ such that $U_n \to U$ in $X^s(\mathbb{R}^{N+1}_+)$, $u_n \to u$ in $L^q_{loc}(\mathbb{R}^N)$ for any $q \in [2, 2^*_s)$ and $u_n \to u$ a.e. in \mathbb{R}^N .

Therefore, (3.4), $||U_n||_{X^s(\mathbb{R}^{N+1}_+)} \leq C$, Hölder inequality and Lemma 2.2 yield

$$\begin{split} \lim_{R \to \infty} \limsup_{n \to \infty} \left| \iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} \nabla U_n \nabla \Psi_R U_n \, dx \, dy \right| \\ &\leq \lim_{R \to \infty} \limsup_{n \to \infty} \frac{C}{R} \left(\iint_{B_R^+ \setminus B_{\frac{R}{2}}^+} y^{1-2s} |\nabla U_n|^2 \, dx \, dy \right)^{\frac{1}{2}} \left(\iint_{B_R^+ \setminus B_{\frac{R}{2}}^+} y^{1-2s} |U_n|^2 \, dx \, dy \right)^{\frac{1}{2}} \\ &\leq \lim_{R \to \infty} \frac{C}{R} \left(\iint_{B_R^+ \setminus B_{\frac{R}{2}}^+} y^{1-2s} |U|^2 \, dx \, dy \right)^{\frac{1}{2}} \\ &\leq \lim_{R \to \infty} \frac{C}{R} \left(\iint_{B_R^+ \setminus B_{\frac{R}{2}}^+} y^{1-2s} |U|^{2\gamma} \, dx \, dy \right)^{\frac{1}{2\gamma}} \left(\iint_{B_R^+ \setminus B_{\frac{R}{2}}^+} y^{1-2s} \, dx \, dy \right)^{\frac{\gamma-1}{2\gamma}} \\ &\leq C \lim_{R \to \infty} \left(\iint_{B_R^+ \setminus B_{\frac{R}{2}}^+} y^{1-2s} |U|^{2\gamma} \, dx \, dy \right)^{\frac{1}{2\gamma}} = 0. \end{split}$$

$$(3.8)$$

Then, putting together (3.5), (3.6), (3.7) and (3.8), we have for any $R \ge R(\varepsilon)$ and $n \ge n(\varepsilon)$ sufficiently large

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U_n|^2 \Psi_R \, dx \, dy + \int_{\mathbb{R}^N} (1-\theta) u_n^2 \psi_R \, dx \le \varepsilon \,. \tag{3.9}$$

From $\theta \in (0, 1)$ and (3.3), we can deduce that (3.9) implies (3.2).

Now, we exploit the relation (3.2) in order to prove the existence of a convergent subsequence for $\{U_n\}$. By using the fact that $I'(U_n) = 0$ and $\{U_n\}$ is bounded in $X^s(\mathbb{R}^{N+1}_+)$, we can see that

$$\langle I'(U_n), U_n \rangle = \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U_n|^2 \, dx \, dy + \int_{\mathbb{R}^N} u_n^2 \, dx - \int_{\mathbb{R}^N} k(x) f(u_n) u_n \, dx - \int_{\mathbb{R}^N} h(x) u_n \, dx = o(1)$$
(3.10)

and

$$\langle I'(U_n), U \rangle = \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} \nabla U_n \nabla U \, dx \, dy + \int_{\mathbb{R}^N} u_n u \, dx - \int_{\mathbb{R}^N} k(x) f(u_n) u \, dx - \int_{\mathbb{R}^N} h(x) u \, dx = o(1).$$

$$(3.11)$$

Hence, in order to prove our Lemma, it is enough to prove that $||U_n||_{X^s(\mathbb{R}^{N+1}_+)} \to ||U||_{X^s(\mathbb{R}^{N+1}_+)}$ as $n \to \infty$. In view of (3.10) and (3.11), this is equivalent to show that

$$\int_{\mathbb{R}^N} k(x) f(u_n)(u_n - u) \, dx + \int_{\mathbb{R}^N} h(x)(u_n - u) \, dx = o(1).$$
(3.12)

Clearly, by using the facts $k \in L^{\infty}(\mathbb{R}^N)$, $h \in L^2(\mathbb{R}^N)$ and $u_n \to u$ in $L^2(B_R)$ for any R > 0, we can see that

$$\int_{B_R} k(x) f(u_n)(u_n - u) \, dx + \int_{B_R} h(x)(u_n - u) \, dx = o(1). \tag{3.13}$$

On the other hand, by using (3.2), we know that for any $\varepsilon > 0$ there exists $R(\varepsilon) > 0$ such that

$$\int_{|x|\geq R(\varepsilon)} k(x)f(u_n)(u_n-u)\,dx + \int_{\mathbb{R}^N} h(x)(u_n-u)\,dx$$

$$\leq \left(\int_{|x|\geq R(\varepsilon)} k(x)|f(u_n)|^2\,dx\right)^{\frac{1}{2}} \left(\int_{|x|\geq R(\varepsilon)} k(x)|u_n-u|^2\,dx\right)^{\frac{1}{2}}$$

$$+ \left(\int_{|x|\geq R(\varepsilon)} |h(x)|^2\,dx\right)^{\frac{1}{2}} \left(\int_{|x|\geq R(\varepsilon)} |u_n-u|^2\,dx\right)^{\frac{1}{2}}$$

$$\leq C \left(\int_{|x|\geq R(\varepsilon)} |u_n|^2\,dx\right)^{\frac{1}{2}} \left(\int_{|x|\geq R(\varepsilon)} |u_n-u|^2\,dx\right)^{\frac{1}{2}} + |h|_2 \left(\int_{|x|\geq R(\varepsilon)} |u_n-u|^2\,dx\right)^{\frac{1}{2}}$$

$$\leq C \varepsilon$$
(3.14)

for n large enough. By combining (3.13) and (3.14) we obtain (3.12). This concludes the proof of lemma.

$$\Box$$

In the next Lemma we show that I is positive on the boundary of some ball in $X^{s}(\mathbb{R}^{N+1}_{+})$, provided that $|h|_{2}$ is sufficiently small. This property will be fundamental to apply Ekeland's variational principle.

Lemma 3.2. Let us assume that (f1)-(f3) hold, $h \in L^2(\mathbb{R}^N)$ such that (1.8) is satisfied, and $k \in L^{\infty}(\mathbb{R}^N)$. Then there exist $\rho, \alpha, m > 0$ such that $I(U)|_{||U||=\rho} \ge \alpha > 0$ for $|h|_2 < m$.

Proof. Fix $\varepsilon \in (0, |k|_{L^{\infty}(\mathbb{R}^N)}^{-1})$. Then, in view of Theorem 1.2 and (1.4), we get

$$I(U) \geq \frac{1}{2} \|U\|^2 - \frac{\varepsilon}{2} |k|_{L^{\infty}(\mathbb{R}^N)} \|U\|^2 - \frac{C(\varepsilon)}{p+1} |k|_{L^{\infty}(\mathbb{R}^N)} S_*^{p+1} \|U\|^{p+1} - |h|_2 \|U\|$$

$$= \|U\| \left[\left(\frac{1}{2} - C_1 \varepsilon \right) \|U\| - C_2(\varepsilon) \|U\|^p - |h|_2 \right],$$
(3.15)

where

$$C_1 := \frac{1}{2} |k|_{L^{\infty}(\mathbb{R}^N)}$$
 and $C_2(\varepsilon) := \frac{C(\varepsilon)}{p+1} |k|_{L^{\infty}(\mathbb{R}^N)} S_*^{p+1}$.

By using (1.8) and (3.15), we can infer that there exist $\rho, \alpha > 0$ such that $I(U)|_{||U||=\rho} \ge \alpha$ provided that $|h|_2 < m$.

For ρ given by Lemma 3.2, we denote $B_{\rho} = \{U \in X^{s}(\mathbb{R}^{N+1}_{+}) : ||U|| < \rho\}$ the ball in $X^{s}(\mathbb{R}^{N+1}_{+})$ with center in 0 and radius ρ . As a consequence of Ekeland's variational principle and Lemma 3.1, we can see that I has a local minimum if $|h|_{2}$ is small enough.

Theorem 3.1. Assume that (f_1) - (f_4) with $l < +\infty$ hold, $h \in L^2(\mathbb{R}^N)$, $h \ge 0$ ($h \ne 0$) and k satisfies (1.6). If $|h|_2 < m$, m is given by Lemma 3.2, then there exists $U_0 \in X^s(\mathbb{R}^{N+1}_+)$ such that

$$I(U_0) = \inf\{I(U) : U \in \overline{B}_\rho\} < 0$$

and U_0 is a positive solution of problem (3.1).

Proof. Since $h(x) \in L^2(\mathbb{R}^N)$, $h \ge 0$ and $h \ne 0$, we can choose a function $V \in X^s(\mathbb{R}^{N+1}_+)$ such that

$$\int_{\mathbb{R}^N} h(x)v(x)\,dx > 0. \tag{3.16}$$

For all t > 0, we can note that

$$\begin{split} I(tV) &= \frac{t^2}{2} \left[\int_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla V|^2 \, dx dy + \int_{\mathbb{R}^N} v^2 \, dx \right] - \int_{\mathbb{R}^N} k(x) F(tv) \, dx - t \int_{\mathbb{R}^N} h(x) v(x) \, dx \\ &\leq \frac{t^2}{2} \|V\|^2 - t \int_{\mathbb{R}^N} h(x) v(x) \, dx < 0 \text{ for } t > 0 \text{ small enough.} \end{split}$$

Then

$$c_0 := \inf\{I(U) : U \in \overline{B}_\rho\} < 0.$$

By applying the Ekeland's variational principle, we know that there exists $\{U_n\} \subset \overline{B}_{\rho}$ such that (i) $c_0 \leq I(U_n) < c_0 + \frac{1}{n}$, (ii) $I(W) \geq I(U_n) - \frac{1}{n} ||W - U_n||$ for all $W \in \overline{B}_{\rho}$. Now, our claim is to prove that $\{U_n\}$ is a bounded (PS) sequence of I.

Firstly, we show that $||U_n|| < \rho$ for a *n* large enough. If $||U_n|| = \rho$ for infinitely many *n*, then we may assume that $||U_n|| = \rho$ for all $n \ge 1$. Hence, by Lemma 3.2, we can see that $I(U_n) \ge \alpha > 0$. Taking the limit as $n \to \infty$ and by using (i), we can deduce that $0 > c_0 \ge \alpha > 0$, which is a contradiction.

Now, we show that $I'(U_n) \to 0$. Indeed, for any $U \in X^s(\mathbb{R}^{N+1}_+)$ with ||U|| = 1, let $W_n = U_n + tU$. For a fixed n, we have $||W_n|| \leq ||U_n|| + t < \rho$ when t is small enough. By using (ii), we deduce that

$$I(W_n) \ge I(U_n) - \frac{t}{n} \|U\|,$$

that is

$$\frac{I(W_n) - I(U_n)}{t} \ge -\frac{\|U\|}{n} = -\frac{1}{n}.$$

Taking the limit as $t \to 0$, we deduce that $\langle I'(U_n), U \rangle \geq -\frac{1}{n}$, which means $|\langle I'(U_n), U \rangle| \leq \frac{1}{n}$ for any $U \in X^{s}(\mathbb{R}^{N+1})$ with ||U|| = 1. This shows that $\{U_n\}$ is a bounded (PS) sequence of I. Then, by using Lemma 3.1, we can see that there exists $U_0 \in X^s(\mathbb{R}^{N+1}_+)$ such that $I'(U_0) = 0$ and $I(U_0) = c_0 < 0.$

In what follows, we show that problem (3.1) has a mountain pass type solution. In order to do this, we use the following variant of version of Mountain Pass Theorem which allows us to find a so-called Cerami sequence $\{U_n\}$. Since this type of Palais-Smale sequence enjoys of some useful properties, we are able to prove its boundedness in the asymptotically linear case.

Theorem 3.2. [19] Let X be a real Banach space with its dual X^* , and suppose that $I \in C^1(X, \mathbb{R})$ satisfies

$$\max\{I(0), I(e)\} \le \mu < \alpha \le \inf_{\|x\|=\rho} I(x),$$

for some $\mu < \alpha$, $\rho > 0$ and $e \in X$ with $||e|| > \rho$. Let $c \ge \alpha$ be characterized by

$$c = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} I(\gamma(t)),$$

where

$$\Gamma = \{ \gamma \in C([0,1], X) : \gamma(0) = 0, \gamma(1) = e \}.$$

Then, there exists a Cerami sequence $\{x_n\} \subset X$ at the level c that is

$$I(x_n) \to c \text{ and } (1 + ||x_n||) ||I'(x_n)||_* \to 0$$

as $n \to \infty$.

The below lemma, shows that I possesses a mountain pass geometry.

Lemma 3.3. Suppose that (f1)- (f4) hold and $\mu^* \in (l, +\infty)$ with μ^* given by (1.7). Then there exists $V \in X^s(\mathbb{R}^{N+1}_+)$ with $||V|| > \rho$, ρ is given by Lemma 3.2, such that I(V) < 0.

Proof. Being $l > \mu^*$, we can find a nonnegative function $W \in X^s(\mathbb{R}^{N+1}_+)$, such that

$$\int_{\mathbb{R}^N} k(x) w^2 \, dx = 1 \text{ such that } \int_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla W|^2 \, dx \, dy + \int_{\mathbb{R}^N} w^2 \, dx < l$$

By using (f4) and Fatou's lemma, we can see that

$$\lim_{t \to \infty} \frac{I(tW)}{t^2} = \frac{1}{2} \|W\|^2 - \lim_{t \to \infty} \int_{\mathbb{R}^N} k(x) \frac{F(tw)}{t^2} \, dx - \lim_{t \to \infty} \frac{1}{t} \int_{\mathbb{R}^N} h(x) w(x) \, dx \le \frac{1}{2} (\|W\|^2 - l) < 0$$

Then, we take $V = t_0 W$ with t_0 large enough.

Putting together Lemmas 3.2 and Lemma 3.3, we can see that the assumptions of Theorem 3.2 are satisfied. Then, we can find a sequence $\{U_n\} \subset X^s(\mathbb{R}^{N+1}_+)$ with the following property

$$I(U_n) \to c > 0 \text{ and } ||I'(U_n)||(1 + ||U_n||) \to 0,$$
 (3.17)

Let

$$W_n = \frac{U_n}{\|U_n\|}.$$

Obviously, $\{W_n\}$ is bounded in $X^s(\mathbb{R}^{N+1}_+)$, so there exists a $W \in X^s(\mathbb{R}^{N+1}_+)$ such that, up to a subsequence, we have

$$W_n \to W \text{ in } X^s(\mathbb{R}^{N+1}_+),$$

$$w_n \to w \text{ a.e. in } \mathbb{R}^N,$$

$$w_n \to w \text{ strongly in } L^2_{loc}(\mathbb{R}^N).$$
(3.18)

With the notation above introduced, we prove the following lemma.

Lemma 3.4. Assume that (f_1) - (f_4) and (K) hold. Let $h \in L^2(\mathbb{R}^N)$ and $\mu^* \in (l, +\infty)$ for μ^* given by (1.7). If $||U_n|| \to \infty$, then W given by (3.18) is a nontrivial nonnegative solution of

$$\begin{cases} \operatorname{div}(y^{1-2s}\nabla W) = 0 & \text{in } \mathbb{R}^{N+1}_+ \\ \frac{\partial W}{\partial \nu^{1-2s}} = -w + lk(x)w & \text{on } \mathbb{R}^N \end{cases}$$
(3.19)

Proof. Firstly, we show that $W \neq 0$. We argue by contradiction, and we assume that $W \equiv 0$. By using the Sobolev embedding, we can see that $w_n \to 0$ strongly in $L^2(B_{R_0})$ where R_0 is given by (K). On the other hand, by (f1), (f4) and $l < +\infty$, we can find C > 0 such that

$$\frac{f(t)}{t} \le C, \text{ for all } t \in \mathbb{R}.$$
(3.20)

Therefore, we can deduce that

$$\int_{|x| < R_0} k(x) \frac{f(u_n)}{u_n} w_n^2 \, dx \le C |k|_\infty \int_{|x| < R_0} w_n^2 \, dx \to 0.$$
(3.21)

By the condition (K), we can find $\eta \in (0,1)$ such that

$$\sup\left\{\frac{f(t)}{t}: t > 0\right\} < \eta \inf\left\{\frac{1}{k(x)}: |x| \ge R_0\right\},\tag{3.22}$$

so, for all $n \in \mathbb{N}$, we get

$$\int_{|x|\ge R_0} k(x) \frac{f(u_n)}{u_n} |w_n|^2 \, dx \le \eta \int_{|x|\ge R_0} |w_n|^2 \, dx \le \eta < 1.$$
(3.23)

Putting together (3.21) and (3.23), we have

$$\limsup_{n \to \infty} \int_{\mathbb{R}^N} k(x) \frac{f(u_n)}{u_n} w_n^2 \, dx < 1.$$
(3.24)

Now, by using the fact that $||U_n|| \to \infty$ and (3.17), we can see that

$$\frac{\langle I'(U_n), U_n \rangle}{\|U_n\|^2} = o(1),$$

that is

$$o(1) = \|W_n\|^2 - \int_{\mathbb{R}^N} k(x) \frac{f(u_n)}{u_n} w_n^2 \, dx = 1 - \int_{\mathbb{R}^N} k(x) \frac{f(u_n)}{u_n} w_n^2 \, dx,$$

which yields a contradiction in view of (3.24). Then, we have proved that $W \neq 0$.

In what follows, we will show that W is nonnegative, that is $W \ge 0$. Let $W_n^-(x) = \max\{-W_n(x), 0\}$, and we observe that $\{W_n^-\}$ is bounded in $X^s(\mathbb{R}^{N+1}_+)$.

Since $||U_n|| \to \infty$, we obtain that

$$\frac{\langle I'(U_n), W_n^- \rangle}{\|U_n\|} = o(1),$$

which gives

$$- \|W_n^-\|^2 = \int_{\mathbb{R}^N} k(x) \frac{f(u_n)}{\|u_n\|} w_n^- dx + o(1).$$
(3.25)

Taking into account (f1), we know that $f(t) \equiv 0$ for all $t \leq 0$, so (3.25) implies that

$$\lim_{n \to \infty} \|W_n^-\| = 0,$$

which gives $W^- = 0$ a.e. $x \in \mathbb{R}^N$, that is $W \ge 0$.

Finally, we prove that W is a solution to (3.19). By using (3.17) and $||U_n|| \to \infty$, we get

$$\frac{\langle I'(U_n), \Phi \rangle}{\|U_n\|} = o(1), \text{ for any } \Phi \in C_0^\infty(\mathbb{R}^N),$$

or explicitly

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} \nabla W_n \nabla \Phi \, dx \, dy + \int_{\mathbb{R}^N} w_n \phi \, dx = \int_{\mathbb{R}^N} k(x) \frac{f(u_n)}{u_n} w_n \phi \, dx + o(1) \tag{3.26}$$

where we have used the notation $\phi = \Phi(\cdot, 0)$. Since $W_n \to W$ in $X^s(\mathbb{R}^{N+1}_+)$ and $w_n \to w$ in $L^2_{loc}(\mathbb{R}^N)$, we can deduce that

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} \nabla W_n \nabla \Phi \, dx \, dy + \int_{\mathbb{R}^N} w_n \phi \, dx = \int_{\mathbb{R}^N} k(x) \frac{f(u_n)}{u_n} w_n \phi \, dx + o(1). \tag{3.27}$$

As a consequence, to prove that W solves (3.19), it is suffices to show that

$$\int_{\mathbb{R}^N} k(x) \frac{f(u_n)}{u_n} w_n(x) \phi(x) \, dx \to \int_{\mathbb{R}^N} lk(x) w(x) \phi(x) \, dx. \tag{3.28}$$

Firstly, we note that by (3.20) and $||W_n|| = 1$ we get

$$\int_{\mathbb{R}^{N}} \left| \frac{f(u_{n})}{u_{n}} w_{n}(x) \right|^{2} dx \leq C \int_{\mathbb{R}^{N}} w_{n}^{2} dx \leq C ||W_{n}||^{2} = C$$

that is $\{\frac{f(u_n)}{u_n}w_n\}$ is bounded in $L^2(\mathbb{R}^N)$. Now, let us define the following sets

$$\Omega_{+} = \{ x \in \mathbb{R}^{N} : w(x) > 0 \} \text{ and } \Omega_{0} = \{ x \in \mathbb{R}^{N} : w(x) = 0 \}.$$

In view of (3.18), it is clear that $u_n(x) \to +\infty$ a.e. in $x \in \Omega_+$. Then, by (f4), it follows that

$$\frac{f(u_n)}{u_n}w_n(x) \to lw(x) \text{ a.e. in } x \in \Omega_+.$$
(3.29)

Since $w_n \to 0$ a.e. in $x \in \Omega_0$, from (3.20) we obtain that

$$\frac{f(u_n)}{u_n}w_n(x) \to 0 \equiv lw(x) \text{ a.e. in } x \in \Omega_0.$$
(3.30)

Putting together (3.29) and (3.30), we can deduce that

$$\frac{f(u_n)}{u_n}w_n(x) \rightharpoonup lw(x) \text{ in } L^2(\mathbb{R}^N).$$
(3.31)

Now, by using the facts $\phi \in C_0^{\infty}(\mathbb{R}^N)$ and $k \in L^{\infty}(\mathbb{R}^N)$, we can see that $z(x) = k(x)\phi(x) \in L^2(\mathbb{R}^N)$, and this together with (3.31) implies that

$$\int_{\mathbb{R}^N} \frac{f(u_n)}{u_n} w_n(x) z(x) \to \int_{\mathbb{R}^N} lw(x) z(x) \, dx \text{ as } n \to \infty,$$

that is (3.28) holds.

Lemma 3.5. If $k \in L^{\infty}(\mathbb{R}^N, \mathbb{R}^+)$ and let μ^* be defined by (1.4) with $l \in (\mu^*, +\infty)$. Then, (3.19) has no any nontrivial nonnegative solution.

Proof. Since $l > \mu^*$, there is a constant $\delta > 0$ such that $\mu^* < \mu^* + \delta < l$. By the definition of μ^* , there exists $V_{\delta} \in X^s(\mathbb{R}^{N+1}_+)$ such that $\int_{\mathbb{R}^N} K(x) v_{\delta}^2(x) dx = 1$ and

$$\mu^* \le \|V_{\delta}\|^2 < \mu^* + \delta.$$

Since $C_0^{\infty}(\mathbb{R}^{N+1}_+)$ is dense in $X^s(\mathbb{R}^{N+1}_+)$, we may assume $V_{\delta} \in C_0^{\infty}(\mathbb{R}^{N+1}_+)$. Let R > 0 be such that $\operatorname{supp} V_{\delta} \subset B_R^+$ and define

$$\mu_R = \inf\left\{\iint_{B_R^+} y^{1-2s} |\nabla U|^2 \, dx \, dy + \int_{\Gamma_R^0} u^2 \, dx : \int_{\Gamma_R^0} K(x) u^2(x) \, dx = 1, U \in H^1_{\Gamma_R^+}(B_R^+)\right\},$$

where we used the following notations

$$\begin{split} B_R^+ &= \{(x,y) \in \mathbb{R}^{N+1} : y > 0, |(x,y)| < R\},\\ \Gamma_R^+ &= \{(x,y) \in \mathbb{R}^{N+1} : y \ge 0, |(x,y)| = R\},\\ \Gamma_R^0 &= \{(x,0) \in \partial \mathbb{R}^{N+1}_+ : |x| < R\}, \end{split}$$

and

$$H^{1}_{\Gamma^{+}_{R}}(B^{+}_{R}) = \{ V \in H^{1}(B^{+}_{R}, y^{1-2s}) : V \equiv 0 \text{ on } \Gamma^{+}_{R} \}.$$

Since $V_{\delta} \equiv 0$ on Γ_R^+ , we can infer that $V_{\delta} \in H^1_{\Gamma_R^+}(B_R^+)$ and

$$\mu_R \le \|V_\delta\|^2 < \mu^* + \delta < l.$$
(3.32)

By the compactness of the embedding $H^1_{\Gamma^+_R}(B^+_R) \subset L^2(\Gamma^0_R)$, it is not difficult to see that there exists $W_R \in H^1_{\Gamma^+_R}(B^+_R) \setminus \{0\}$ with $W_R \ge 0$ and $\int_{\Gamma^0_R} K(x) w_R^2(x) dx = 1$ such that

$$\begin{cases} \operatorname{div}(y^{1-2s}\nabla W_R) = 0 & \operatorname{in} B_R^+\\ \frac{\partial W_R}{\partial \nu^{1-2s}} = -w_R + \mu_R k(x) w_R & \operatorname{on} \Gamma_R^0\\ W^R = 0 & \operatorname{on} \Gamma_R^+. \end{cases}$$
(3.33)

It follows from the strong maximum principle [9] that $W_R > 0$ on B_R^+ . We extend $W_R = 0$ in $\mathbb{R}^{N+1} \setminus B_R^+$, so that $W_R \in X^s(\mathbb{R}^{N+1}_+)$. Therefore, if $U \neq 0$, $U \in X^s(\mathbb{R}^{N+1}_+)$ is a nonnegative solution of (3.19), then

$$\mu_R \int_{\Gamma_R^0} k(x) w_R u \, dx = \iint_{B_R^+} y^{1-2s} \nabla W_R \nabla U \, dx dy + \int_{\Gamma_R^0} w_R u \, dx$$

$$= l \int_{\Gamma_R^0} k(x) u w_R \, dx.$$
(3.34)

Using $u \ge 0$ and $u \ne 0$, we may choose R > 0 large enough such that $\int_{\Gamma_R^0} K(x) u w_R dx > 0$. Then, (3.34) implies that $\mu_R = l$, which is a contradiction in view of (3.32).

Proof of Theorem 1.1. By using Lemmas 3.4 and 3.5, it is obvious that the situation $||U_n|| \to \infty$ cannot occur. Therefore, the sequence $\{U_n\}$ is bounded in $X^s(\mathbb{R}^{N+1}_+)$. Taking into account Lemma 3.1 and the Harnack inequality [9], we can deduce that problem (3.1) admits a positive solution $U_1 \in X^s(\mathbb{R}^{N+1}_+)$ with $I(U_1) > 0$. Then, the thesis of theorem follows by Theorem 3.1.

4. Superlinear case: proof of Theorem 1.2

This section is devoted to the proof of Theorem 1.2. For simplicity, we consider problem (2.5) with $k(x) \equiv 1$, that is

$$\begin{cases} \operatorname{div}(y^{1-2s}\nabla U) = 0 & \operatorname{in} \mathbb{R}^{N+1}_+\\ \frac{\partial U}{\partial \nu^{1-2s}} = -u + f(u) + h(x) & \operatorname{on} \mathbb{R}^N \end{cases}$$
(4.1)

where $h(x) = h(|x|) \in C^1(\mathbb{R}^N) \cap L^2(\mathbb{R}^N)$, $h(x) \ge 0$, $h(x) \ne 0$ and f satisfies (f_1) - (f_4) with $l = +\infty$. Since we assume that $k(x) \equiv 1$ and h(x) is radial, it is natural to work on the space of the function belonging to $X^s(\mathbb{R}^{N+1}_+)$ which are radial with respect to x, that is

$$X_r^s(\mathbb{R}^{N+1}_+) = \left\{ U \in X^s(\mathbb{R}^{N+1}_+) : U(x,y) = U(|x|,y) \right\}.$$

We begin proving the following preliminary result

Theorem 4.1. Suppose that $h(x) = h(|x|) \in L^2(\mathbb{R}^N)$, $h(x) \ge 0$, $h(x) \ne 0$ and conditions (f1)-(f3) holds, then there exist $m_1 > 0$ and $\tilde{U}_0 \in X_r^s(\mathbb{R}^{N+1}_+)$ such that $I'(\tilde{U}_0) = 0$ and $I(\tilde{U}_0) < 0$ if $|h|_2 < m_1$.

Proof. Arguing as in the proof of Theorem 3.1, by applying the Ekeland's variational principle, we can obtain the existence of a bounded (PS) sequence $\{\tilde{U}_n\} \subset X_r^s(\mathbb{R}^{N+1}_+)$ such that

$$I(\tilde{U}_n) \to \tilde{c}_0 := \inf\{I(U) : U \in X_r^s(\mathbb{R}^{N+1}_+) \text{ and } \|U\| = \rho\} < 0,$$

where ρ is given by Lemma 3.2. We claim that such infimum is achieved. By using Theorem 2.2, we may assume that there exists $\tilde{U}_0 \in X_r^s(\mathbb{R}^{N+1}_+)$ such that $\tilde{U}_n \rightharpoonup \tilde{U}_0$ in $X_r^s(\mathbb{R}^{N+1}_+)$, $\tilde{u}_n \rightarrow \tilde{u}_0$ in $L^{p+1}(\mathbb{R}^N)$.

Taking into account (f1)-(f3), Theorem 2.1, and by exploiting the fact that $\{U_n\}$ is bounded in $X_r^s(\mathbb{R}^{N+1}_+)$, we can see that

$$\left| \int_{\mathbb{R}^N} f(\tilde{u}_n) (\tilde{u}_n - \tilde{u}_0) \, dx \right| \le \varepsilon \, |\tilde{u}_n|_2 |\tilde{u}_n - \tilde{u}_0|_2 + C_\varepsilon |\tilde{u}_n|_{p+1}^p |\tilde{u}_n - \tilde{u}_0|_{p+1} \le C \, \varepsilon + C_\varepsilon C |\tilde{u}_n - \tilde{u}_0|_{p+1}$$

Hence

$$\lim_{n \to \infty} \left| \int_{\mathbb{R}^N} f(\tilde{u}_n) (\tilde{u}_n - \tilde{u}_0) \, dx \right| \le C \, \varepsilon$$

and by the arbitrariness of ε , we deduce that

$$\int_{\mathbb{R}^N} f(\tilde{u}_n)(\tilde{u}_n - \tilde{u}_0) \, dx \to 0$$

Putting together (f_1) - (f_3) and by using Lemma 2.1, we can obtain that

$$\int_{\mathbb{R}^N} f(\tilde{u}_n) \tilde{u}_n \, dx \to \int_{\mathbb{R}^N} f(\tilde{u}_0) \tilde{u}_0 \, dx$$

Then we can infer that

$$\int_{\mathbb{R}^N} (f(\tilde{u}_n) - f(\tilde{u}_0)) \tilde{u}_0 \, dx = \int_{\mathbb{R}^N} (f(\tilde{u}_n) \tilde{u}_n - f(\tilde{u}_0) \tilde{u}_0) \, dx - \int_{\mathbb{R}^N} f(\tilde{u}_n) (\tilde{u}_n - \tilde{u}_0) \, dx \to 0$$

On the other hand, $\tilde{u}_n \rightarrow \tilde{u}_0$ in $L^2(\mathbb{R}^N)$, so by using the fact that $h \in L^2(\mathbb{R}^N)$, we also have

$$\int_{\mathbb{R}^N} h(x)\tilde{u}_n \, dx \to \int_{\mathbb{R}^N} h(x)\tilde{u}_0 \, dx.$$

Then, by combining $\langle I'(\tilde{U}_n), \tilde{U}_n \rangle \to 0$, $\langle I'(\tilde{U}_n), \tilde{U}_0 \rangle \to 0$, and the above relations, it follows that $\tilde{U}_n \to \tilde{U}_0$ strongly in $X_r^s(\mathbb{R}^{N+1}_+)$.

Therefore, we get

$$I(\tilde{U}_0) = \tilde{c}_0 < 0 \text{ and } I'(\tilde{U}_0) = 0.$$

Now, in order to prove that (4.1) has a mountain pass type solution, we use the following abstract result due to Jeanjean [26]:

Theorem 4.2. [26] Let $(X, \|\cdot\|)$ be a Banach space and $J \subset \mathbb{R}_+$ be an interval. Let $(\mathcal{I}_{\lambda})_{\lambda \in J}$ be a family of C^1 functionals on X of the form

$$\mathcal{I}_{\lambda}(u) = A(u) - \lambda B(u), \quad \text{for } \lambda \in J,$$

where $B(u) \ge 0$ for all $u \in X$, and either $A(u) \to \infty$ or $B(u) \to \infty$ as $||u|| \to \infty$. We assume that there exist $v_1, v_2 \in X$ such that

$$c_{\lambda} = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} \mathcal{I}_{\lambda}(\gamma(t)) > \max\{\mathcal{I}_{\lambda}(v_1), \mathcal{I}_{\lambda}(v_2)\}, \quad \forall \lambda \in J$$

where

$$\Gamma = \{\gamma \in C([0,1], X) : \gamma(0) = v_1, \gamma(1) = v_2\}.$$

Then, for almost every $\lambda \in J$, there is a sequence $(v_n) \subset X$ such that

(i) (v_n) is bounded;

(*ii*) $\mathcal{I}_{\lambda}(v_n) \to c_{\lambda};$ (*iii*) $\mathcal{I}_{\lambda}(v_n) \to 0 \text{ on } X^{-1}$

(iii)
$$L_{\lambda}(v_n) \to 0$$
 on X^{-1}

Moreover, the map $\lambda \mapsto c_{\lambda}$ is continuous from the left hand-side.

For any $\lambda \in [\frac{1}{2}, 1]$, we introduce the following family of functionals $I_{\lambda} : X_r^s(\mathbb{R}^{N+1}_+) \to \mathbb{R}$ defined by

$$I_{\lambda}(U) = \frac{1}{2} \left[\iint_{\mathbb{R}^{N+1}_{+}} y^{1-2s} |\nabla U|^2 \, dx \, dy + \int_{\mathbb{R}^N} u^2 \, dx \right] - \lambda \int_{\mathbb{R}^N} (F(u) + h(x)u) \, dx.$$

for any $U \in X_r^s(\mathbb{R}^{N+1}_+)$.

Next, our claim is to show that I_{λ} verifies the assumptions of Theorem 4.2.

Lemma 4.1. Assume that (f_1) - (f_4) with $l = +\infty$ hold. Then,

(i) There exists $\overline{V} \in X_r^s(\mathbb{R}^{N+1}_+) \setminus \{0\}$ such that $I_\lambda(\overline{V}) < 0$ for all $\lambda \in [\frac{1}{2}, 1]$.

(ii) For $m_1 > 0$ given in Theorem 4.1, if $|h|_2 < m_1$, then

$$c_{\lambda} = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} I_{\lambda}(\gamma(t)) > \max\{I_{\lambda}(0), I_{\lambda}(\bar{V})\} \quad \forall \lambda \in \left\lfloor \frac{1}{2}, 1 \right\rfloor,$$

where $\Gamma = \{\gamma \in C([0,1], X_r^s(\mathbb{R}^{N+1}_+))) : \gamma(0) = 0, \gamma(1) = \bar{V}\}.$

Proof. (i) For any $\delta > 0$, we can find $V \in X_r^s(\mathbb{R}^{N+1}_+) \setminus \{0\}$ and $V \ge 0$ such that

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla V|^2 \, dx < \delta \int_{\mathbb{R}^N} v^2 dx.$$

This is lawful due to

$$\inf\left\{\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U|^2 \, dx : U \in X^s_r(\mathbb{R}^{N+1}_+) \text{ and } |u|_2 = 1\right\} = 0$$

(via the Pohozaev identity, one can see that $(-\Delta)^s$ has no eigenvalues in $H^s(\mathbb{R}^N)$). By using (f4) with $l = +\infty$, and by applying Fatou's lemma, we can deduce that

$$\lim_{t \to +\infty} \int_{\mathbb{R}^N} \frac{F(tv)}{t^2} \, dx \ge (1+\delta) \int_{\mathbb{R}^N} v^2 \, dx.$$

Hence, for any $\lambda \in [\frac{1}{2}, 1]$, we get

$$\lim_{t \to +\infty} \frac{I_{\lambda}(tV)}{t^2} \le \lim_{t \to +\infty} \frac{I_{\frac{1}{2}}(tV)}{t^2} \le \frac{1}{2} \left(\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla V|^2 \, dx \, dy - \delta \int_{\mathbb{R}^N} |v|^2 \, dx \right) < 0.$$

Take $t_1 > 0$ large enough such that $I_{\frac{1}{2}}(t_1 V) < 0$, and we set $\bar{V} = t_1 V$. Then, we can see that $I_{\lambda}(\bar{V}) \leq I_{\frac{1}{2}}(\bar{V}) < 0$, that is the condition (i) is satisfied.

(*ii*) It is clear that, for any $\lambda \in [\frac{1}{2}, 1]$ and $U \in X_r^s(\mathbb{R}^{N+1}_+)$, we have

$$I_{\lambda}(U) \ge \frac{1}{2} \left[\int_{\mathbb{R}^N} y^{1-2s} |\nabla U|^2 \, dx \, dy + \int_{\mathbb{R}^N} u^2 \, dx \right] - \int_{\mathbb{R}^N} F(u) \, dx - |h|_2 |u|_2 =: J(U).$$

Then, we can proceed as in the proof of Lemma 3.2, to deduce that

$$\inf_{\gamma \in \Gamma} \max_{t \in [0,1]} J(\gamma(t)) > 0,$$

provided that $|h|_2 < m_1$, with m_1 given by Theorem 4.1. Then, for any $\lambda \in \left[\frac{1}{2}, 1\right]$, we can see that

$$c_{\lambda} = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} I_{\lambda}(\gamma(t)) \ge \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} J(\gamma(t)) > \max\{I_{\lambda}(0), I_{\lambda}(\bar{V})\}.$$

This ends the proof of the lemma.

By using Lemma 4.1 and Theorem 4.2, we can infer that there exists $\{\lambda_j\} \subset [\frac{1}{2}, 1]$ such that (i) $\lambda_j \to 1$ as $j \to +\infty$;

(*ii*) I_{λ_j} has a bounded (PS) sequence $\{U_n^j\}$ at the level c_{λ_j} . In view of Theorem 2.2, we deduce that for each $j \in \mathbb{N}$, there exists $U_j \in X_r^s(\mathbb{R}^{N+1}_+)$ such that $U_n^j \to U_n$ strongly in $X_r^s(\mathbb{R}^{N+1}_+)$ and U_j is a positive solution of

$$\begin{cases} \operatorname{div}(y^{1-2s}\nabla U_j) = 0 & \text{in } \mathbb{R}^{N+1}_+\\ \frac{\partial U_j}{\partial \nu^{1-2s}} = -u_j + \lambda_j [f(u_j) + h(x)] & \text{on } \mathbb{R}^N \end{cases}$$

Proceeding as in [1, 12], it is easy to see that each U_j satisfies the following Pohozaev identity:

$$\frac{N-2s}{2} \iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U_j|^2 \, dx \, dy + \frac{N}{2} \int_{\mathbb{R}^N} u_j^2 \, dx = N\lambda_j \int_{\mathbb{R}^N} (F(u_j) + hu_j) \, dx + \lambda_j \int_{\mathbb{R}^N} \nabla h(x) \cdot xu_j \, dx.$$

$$\tag{4.2}$$

In the next lemma, we use the condition (H) to prove the boundedness of the sequence $\{U_i\}$.

Lemma 4.2. Assume that (f_1) - (f_4) with $l = +\infty$ hold, and h satisfies (1.5) and $|h|_2 < m_1$ for m_1 given in Theorem 4.1. Then $\{U_j\} \subset X_r^s(\mathbb{R}^{N+1}_+)$ is bounded.

Proof. By using Theorem 4.2, we know that the map $\lambda \to c_{\lambda}$ is continuous from the left. Then, by Lemma 4.1 (*ii*), we can deduce that $I_{\lambda_j}(U_j) = c_{\lambda_j} \to c_1 > 0$ as $\lambda_j \to 1$.

Hence, we can find a constant K > 0 such that $I_{\lambda_j}(U_j) \leq K$ for all $j \in \mathbb{N}$. By combining this, (4.2), $u_j > 0$ and (H), we can see that

$$\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U_j|^2 \, dx dy \le \frac{KN}{s} - \frac{\lambda_j}{s} \int_{\mathbb{R}^N} \nabla h(x) \cdot x u_j \, dx \le \frac{KN}{s},$$

which together with the Sobolev inequality (2.3), implies that

$$|u_j|_{2_s^*} \le S_* \left(\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U_j|^2 \, dx \, dy \right)^{\frac{1}{2}} \le C.$$
(4.3)

Now, from $I_{\lambda_j}(U_j) \leq K$ for all $j \in \mathbb{N}$, it follows that

$$\frac{1}{2} \left[\iint_{\mathbb{R}^{N+1}_+} y^{1-2s} |\nabla U_j|^2 \, dx \, dy + \int_{\mathbb{R}^N} u_j^2 \, dx \right] - \lambda_j \int_{\mathbb{R}^N} (F(u_j) + h(x)u_j) \, dx \le K. \tag{4.4}$$

On the other hand, by using (f_2) , (f_3) , we can see that there exists a constant C > 0 such that

$$\int_{\mathbb{R}^N} F(u_j) \, dx \leq \frac{1}{4} \int_{\mathbb{R}^N} u_j^2 \, dx + C \int_{\mathbb{R}^N} u_j^{2^*} \, dx,$$

where $2_s^* = \frac{2N}{N-2s}$.

Substituting this inequality into (4.4), and by using (4.3), (2.3), we deduce that

$$\frac{1}{2} \int_{\mathbb{R}^N} u_j^2 dx \le \lambda_j \int_{\mathbb{R}^N} (F(u_j) + h(x)u_j) dx + K$$
$$\le \frac{1}{4} |u_j|_2^2 + C|u_j|_{2_s^*}^{2_s^*} + |h|_2 |u_j|_2 + K$$
$$\le \frac{1}{4} |u_j|_2^2 + \bar{C} + |h|_2 |u_j|_2 + K.$$

Then

$$\frac{1}{4} \int_{\mathbb{R}^N} u_j^2 \, dx \le \tilde{C} + |h|_2 |u_j|_2,$$

that is

$$|u_j|_2 \le C \text{ for all } j \in \mathbb{N},\tag{4.5}$$

for some positive constant C independent of j. Putting together (4.3) and (4.5), we can conclude the proof of this lemma.

Lemma 4.3. Under the assumptions of Lemma 4.2, the above sequence $\{U_j\}$ is also a (PS) sequence of I.

Proof. From the definitions of I and I_{λ_i} we can deduce that

$$I(U_j) = I_{\lambda_j}(U_j) + (\lambda_j - 1) \int_{\mathbb{R}^N} (F(u_j) + h(x)u_j) \, dx.$$
(4.6)

By using Theorem 4.2, we obtain

$$I_{\lambda_j}(U_j) = c_{\lambda_j} \to c_1 > 0$$
 as $\lambda_j \to 1$

Hence, by applying Lemma 4.2 and (4.6), we get $I(U_j) \to c_1 > 0$. Being $I'_{\lambda_j}(U_j) = 0$, we can infer that, for any $\Psi \in C_0^{\infty}(\mathbb{R}^{N+1}_+)$,

$$\langle I'(U_j), \Psi \rangle = \langle I'_{\lambda_j}(U_j), \Psi \rangle + (\lambda_j - 1) \int_{\mathbb{R}^N} (f(u_j) + h(x)) \psi \, dx \to 0$$

that is $I'(U_j) \to 0$ as $j \to \infty$ in the dual space of $X_r^s(\mathbb{R}^{N+1}_+)$.

Finally, we give the proof of the main result of this section:

Proof of Theorem 1.2. Taking into account Theorem 4.1, we know that (4.1) admits a positive solution $\tilde{U}_0 \in X_r^s(\mathbb{R}^{N+1}_+)$ such that $I(\tilde{U}_0) < 0$. On the other hand, by Lemma 4.3 and Theorem 2.2, we know that problem (4.1) has a second positive solution $\tilde{U}_1 \in X_r^s(\mathbb{R}^{N+1}_+)$ with $I(\tilde{U}_1) = c_1 > 0$. As a consequence $\tilde{U}_0 \neq \tilde{U}_1$, and this ends the proof.

5. Some examples

In this last section we provide some examples of functions f, k and h for which our main results are applicable.

Example 5.1. Let $R_0 > 0$ and let us define

$$k(x) = \begin{cases} \frac{1}{1+|x|} & \text{if } |x| < R_0\\ \frac{1}{1+R_0} & \text{if } |x| \ge R_0 \end{cases} \quad and \ f(t) = \begin{cases} \frac{R_0 t^2}{1+t} & \text{if } t > 0\\ 0 & \text{if } t \le 0 \end{cases}$$

It is clear that $|k|_{\infty} = 1$, and f satisfies (f1)-(f3) and (f4) with $l = R_0$. Moreover, we note that (K) holds because of

$$\sup\left\{\frac{f(t)}{t} : t > 0\right\} = R_0 < R_0 + 1 = \inf\left\{\frac{1}{k(x)} : |x| \ge R_0\right\}.$$

Now, to verify that $l > \mu^*$, we have to choose a special $R_0 > 0$. For R > 0, we take $\phi \in C_0^{\infty}(\mathbb{R}^N)$ such that $\phi(x) = 1$ if $|x| \le R$, $\phi(x) = 0$ if $|x| \ge 2R$ and $|\nabla \phi(x)| \le \frac{C}{R}$ for all $x \in \mathbb{R}^N$.

Since $\phi \in H^1(\mathbb{R}^N) \subset H^s(\mathbb{R}^N)$, we can see that

$$\|\phi\|_{H^s(\mathbb{R}^N)} \le C \|\phi\|_{H^1(\mathbb{R}^N)}.$$

On the other hand, for any $R_0 > 2R$, we have

$$\frac{\int_{\mathbb{R}^N} \phi^2 dx}{\int_{\mathbb{R}^N} k(x) \phi^2 dx} \le \frac{\int_{\mathbb{R}^N} \phi^2 dx}{\frac{1}{1+2R} \int_{\mathbb{R}^N} \phi^2 dx} = 1 + 2R$$

and

$$\frac{\int_{\mathbb{R}^N} |\nabla \phi|^2 dx}{\int_{\mathbb{R}^N} k(x) \phi^2 dx} \le \frac{\frac{C^2}{R^2} |B_{2R}|}{\int_{B_R} k(x) dx} \le \frac{\frac{C^2}{R^2} |B_{2R}|}{\frac{1}{1+R} |B_R|} = C_1 \frac{(1+R)}{R^2}.$$

Therefore

$$\frac{\|\phi\|_{H^s(\mathbb{R}^N)}^2}{\int_{\mathbb{R}^N} k(x)\phi^2 dx} \le \frac{C\|\phi\|_{H^1(\mathbb{R}^N)}^2}{\int_{\mathbb{R}^N} k(x)\phi^2 dx} \le C_2 \frac{(1+R)}{R^2} + C_3(1+2R)$$

where $C_2, C_3 > 0$ are constants independent of R. Choosing R > 0 such that $C_2 \frac{(1+R)}{R^2} \leq C_3$, we can infer that $\mu^* \leq 2C_3(R+1)$. Then, taking $R_0 = 2C_3(R+1) + 2R$, we have

$$\lim_{t \to +\infty} \frac{f(t)}{t} = l = R_0 > \mu^*$$

Now, fix $\varepsilon \in (0,1)$, and let $h \in L^2(\mathbb{R}^N)$ such that

$$|h|_2 < m := \max_{t \ge 0} \left[\left(\frac{1}{2} - \frac{\varepsilon}{2} \right) t - \frac{C_{\varepsilon}}{p+1} S_*^{p+1} t^p \right].$$

Then, all assumptions of Theorem 1.1 are satisfied, and we can find at least two positive solutions to (1.1).

Example 5.2. Fix $\varepsilon \in (0,1)$, and let us consider the following functions

$$h(x) = \begin{cases} 0 & \text{if } |x| < \sqrt{3} \lor |x| > 2\\ C(|x|^2 - 2)^2(|x|^2 - 3)^2(|x|^2 - 4)^2 & \text{if } \sqrt{3} \le |x| \le 2 \end{cases}$$

and

$$f(t) = \begin{cases} t \log(1+t) & \text{if } t > 0\\ 0 & \text{if } t \le 0 \end{cases}$$

where C > 0 is a constant such that

$$|h|_2 < m := \max_{t \ge 0} \left[\left(\frac{1}{2} - \frac{\varepsilon}{2} \right) t - \frac{C_{\varepsilon}}{p+1} S_*^{p+1} t^p \right].$$

It is clear that f satisfies (f1)-(f3) and (f4) with $l = \infty$, and $h \in C^1(\mathbb{R}^N) \cap L^2(\mathbb{R}^N)$. Moreover, for any $\sqrt{3} < |x| < 2$, we have

$$x \cdot \nabla h = 4C \left[|x|^2 (|x|^2 - 2)(|x|^2 - 3)(|x|^2 - 4)(3|x|^4 - 22|x|^2 + 26) \right] \ge 0,$$

so $x \cdot \nabla h \geq 0$ on \mathbb{R}^N . In particular, $x \cdot \nabla h \in L^q(\mathbb{R}^N)$ for any $q \in [1, \infty]$. Then, we can apply Theorem 1.2 to deduce that the problem (1.1) admits at least two positive solutions.

References

- V. Ambrosio, Ground states for a fractional scalar field problem with critical growth, Differential Integral Equations 30 (2017), no. 1-2, 115–132.
- [2] V. Ambrosio, Ground states for superlinear fractional Schrödinger equations in ℝ^N, Ann. Acad. Sci. Fenn. Math. 41 (2016), no. 2, 745–756.
- [3] V. Ambrosio, Multiple solutions for a nonlinear scalar field equation involving the fractional Laplacian, preprint arXiv:1603.09538.
- [4] V. Ambrosio, Multiple solutions for a fractional p-Laplacian equation with sign-changing potential, Electron. J. Differential Equations (2016), Paper No. 151, 12 pp.
- [5] A. Bahri and H. Berestycki, A perturbation method in critical point theory and applications, Trans. Amer. Math. Soc. 267 (1981), no. 1, 1–32.
- [6] T. Bartsch, Z. Liu and T. Weth, Sign changing solutions of superlinear Schrödinger equations, Comm. Partial Differential Equations 29 (2004), no. 1-2, 25–42.
- [7] T. Bartsch, Z. Q. Wang, Existence and multiplicity results for some superlinear elliptic problems on \mathbb{R}^N , Comm. Partial Differential Equations **20** (1995)1725–1741.
- [8] H. Berestycki and P.L. Lions, Nonlinear scalar field equations. I. Existence of a ground state, Arch. Rational Mech. Anal. 82 (1983), no. 4, 313–345.
- [9] X. Cabré and Y. Sire, Nonlinear equations for fractional Laplacians, I: Regularity, maximum principles, and Hamiltonian estimates, Ann. Inst. H. Poincaré Anal. Non Linéaire 31 (2014), 23–53.
- [10] L.A. Caffarelli and L.Silvestre, An extension problem related to the fractional Laplacian, Comm. Partial Differential Equations 32 (2007),1245–1260.
- [11] Dao-Min M. Cao and Huan-Song Zhou, Multiple positive solutions of nonhomogeneous semilinear elliptic equations in ℝ^N, Proc. Roy. Soc. Edinburgh Sect. A **126** (1996), no. 2, 443–463.

- [12] X. J. Chang and Z.Q. Wang, Ground state of scalar field equations involving fractional Laplacian with general nonlinearity, Nonlinearity 26, 479–494 (2013).
- [13] X. Chang, Ground state solutions of asymptotically linear fractional Schrödinger equations, J. Math. Phys. 54 (2013), no. 6, 061504, 10 pp.
- [14] Y. Cho, M.M Fall, H. Hajaiej, P. Markowich and S. Trabelsi, On the orbital stability of standing waves of a class of fractional Schrödinger equations with general Hartree-type integrand, (To appear in Analysis and Applications, DOI: 10.1142/S0219530516500056).
- [15] Y. Cho, H. Hajaiej, G. Hwang and T. Ozawa, On the Cauchy problem of fractional Schrödinger equation with Hartree type nonlinearity, Funkcial. Ekvac. 56 (2013), no. 2, 193–224.
- [16] E. Colorado, A. de Pablo, and U. Sánchez, Perturbations of a critical fractional equation, Pacific J. Math. 271 (2014), no. 1, 65–85.
- [17] E. Di Nezza, G. Palatucci and E. Valdinoci, *Hitchhiker's guide to the fractional Sobolev spaces*, Bull. Sci. math. 136 (2012), 521–573.
- [18] S. Dipierro, M. Medina, and E. Valdinoci, Fractional elliptic problems with critical growth in the whole of \mathbb{R}^N , (2015). arXiv:1506.01748vl.
- [19] I. Ekeland, Convexity Methods in Hamiltonian Mechanics, Springer-Verlag, Berlin, 1990. x+247 pp.
- [20] M. M. Fall, F. Mahmoudi and E. Valdinoci, Ground states and concentration phenomena for the fractional Schrödinger equation, Nonlinearity 28 (2015), no. 6, 1937–1961.
- [21] P. Felmer, A. Quaas and J. Tan, Positive solutions of the nonlinear Schrödinger equation with the fractional Laplacian, Proc. Roy. Soc. Edinburgh Sect. A 142 (2012), 1237–1262.
- [22] G. M. Figueiredo and G. Siciliano, A multiplicity result via Ljusternick-Schnirelmann category and Morse theory for a fractional Schrödinger equation in ℝ^N, NoDEA Nonlinear Differential Equations Appl. 23 (2016), no. 2, Art. 12, 22 pp.
- [23] A. Floer and A. Weinstein, Nonspreading wave packets for the cubic Schrdinger equation with a bounded potential, J. Funct. Anal. 69 (1986), 397–408.
- [24] R. Frank, E. Lenzmann, L. Silvestre, Uniqueness of radial solutions for the fractional Laplacian, Comm. Pure Appl. Math. 69 (2016), no. 9, 1671–1726.
- [25] L. Jeanjean, Two positive solutions for a class of nonhomogeneous elliptic equations, Differential Integral Equations 10 (1997), no. 4, 609–624.
- [26] J. Jeanjean, On the existence of bounded Palais-Smale sequences and application to a Landesman-Lazer-type problem set on \mathbb{R}^N , Proc. Roy. Soc. Edinburgh Sect. A **129** (1999), no. 4, 787–809.
- [27] L. Jeanjean, K. Tanaka, A positive solution for an asymptotically linear elliptic problem on \mathbb{R}^N autonomous at infinity, ESAIM Control Optim. Calc. Var. 7 (2002), 597–614.
- [28] H. Hajaiej, L. Molinet, T. Ozawa and B. Wang, Necessary and sufficient conditions for the fractional Gargliardo-Nirenberg inequalities and applications to Navier-Stokes and generalized boson equations, Harmonic analysis and nonlinear partial differential equations, 159–175, RIMS Kkyroku Bessatsu, B26, Res. Inst. Math. Sci. (RIMS), Kyoto, 2011.
- [29] H. Hajaiej, Existence of minimizers of functional involving the fractional gradient in the absence of compactness, symmetry and monotonicity, J. Math. Anal. Appl. 399 (2013), no. 1, 17–26.
- [30] H. Hajaiej, On the optimality of the assumptions used to prove the existence and symmetry of minimizers of some fractional constrained variational problems, Ann. Henri Poincaré 14 (2013), no. 5, 1425–1433.
- [31] H. Hajaiej, Symmetry of minimizers of some fractional problems, Applicable Analysis, 94 (2015), no. 4, 694–700.
- [32] N. Laskin, Fractional quantum mechanics and Lèvy path integrals, Phys. Lett. A 268 (2000), 298–305.
- [33] N. Laskin, Fractional Schrödinger equation, Phys. Rev. E 66 (2002), 056108.
- [34] P. L. Lions, Symetrié et compacité dans les espaces de Sobolev, J.Funct.Anal. 49, (1982), 315-334.
- [35] C. Liu, Z. Wang and H.S. Zhou, Asymptotically linear Schrödinger equation with potential vanishing at infinity, J. Differential Equations 245 (2008), no. 1, 201–222.
- [36] G. Molica Bisci and V. Rădulescu, Ground state solutions of scalar field fractional Schrödinger equations, Calc. Var. Partial Differential Equations 54 (2015), no. 3, 2985–3008.
- [37] G. Molica Bisci, V. Rădulescu and R. Servadei, Variational methods for nonlocal fractional problems, with a foreword by Jean Mawhin. Encyclopedia of Mathematics and its Applications, 162. Cambridge University Press, Cambridge, 2016. xvi+383 pp.
- [38] P. Pucci, M. Xiang, B. Zhang, Multiple solutions for nonhomogeneous Schrödinger-Kirchhoff type equations involving the fractional p-Laplacian in \mathbb{R}^N , Calc. Var. Partial Differential Equations 54 (2015), no. 3, 2785– 2806.
- [39] P. H. Rabinowitz, Multiple critical points of perturbed symmetric functionals, Trans. Amer. Math. Soc. 272 (1982), no. 2, 753–769.
- [40] P. Rabinowitz, On a class of nonlinear Schrödinger equations Z. Angew. Math. Phys. 43 (1992), no. 2, 270291.

- [41] S. Secchi, Ground state solutions for nonlinear fractional Schrödinger equations in \mathbb{R}^N , J. Math. Phys. 54 (2013), 031501.
- [42] S. Secchi, On some nonlinear fractional equations involving the Bessel potential, J. Dynam. Differential Equations (2016), doi:10.1007/s10884-016-9521-y.
- [43] R. Servadei, Infinitely many solutions for fractional Laplace equations with subcritical nonlinearity, Contemp. Math. 595, (2013), 317–340.
- [44] X. Shang and J. Zhang, Ground states for fractional Schrödinger equations with critical growth, Nonlinearity 27 (2014), no. 2, 187–207.
- [45] M. Struwe, Variational methods. Applications to nonlinear partial differential equations and Hamiltonian systems, Springer-Verlag, Berlin, (1990).
- [46] C.A. Stuart and H.S. Zhou, Applying the mountain pass theorem to an asymptotically linear elliptic equation on \mathbb{R}^N , Comm. Partial Differential Equations 24 (1999), no. 9-10, 1731–1758.
- [47] C. L. Torres, Non-homogeneous fractional Schrödinger equation, J. Fract. Calc. Appl. 6 (2015), no. 2, 108–114.
- [48] Z. Wang and H.S. Zhou, Positive solutions for a nonhomogeneous elliptic equation on \mathbb{R}^N without (AR) condition, J. Math. Anal. Appl. **353** (2009), no. 1, 470–479.
- [49] X.P. Zhu, A perturbation result on positive entire solutions of a semilinear elliptic equation, J. Differential Equations 92 (1991), no. 2, 163–178.
- [50] X.P. Zhu and H.S. Zhu, Existence of multiple positive solutions of inhomogeneous semilinear elliptic problems in unbounded domains, Proc. Roy. Soc. Edinburgh Sect. A 115 (1990), no. 3-4, 301–318.

DIPARTIMENTO DI SCIENZE PURE E APPLICATE (DISPEA), UNIVERSITÀ DEGLI STUDI DI URBINO 'CARLO BO' PIAZZA DELLA REPUBBLICA, 13 61029 URBINO (PESARO E URBINO, ITALY) *E-mail address*: vincenzo.ambrosio@uniurb.it

Department of Mathematics California State University, Los Angeles 5151 State University Drive

Los Angeles, CA 90032

E-mail address: hichem.hajaiej@gmail.com