

VARIATIONS ON A PROOF OF A BORDERLINE BOURGAIN–BREZIS SOBOLEV EMBEDDING THEOREM

SAGUN CHANILLO, JEAN VAN SCHAFTINGEN, AND PO-LAM YUNG

To Haïm Brezis in admiration and friendship

ABSTRACT. We offer a variant of a proof of a borderline Bourgain–Brezis Sobolev embedding theorem on \mathbb{R}^n . We use this idea to extend the result to real hyperbolic spaces \mathbb{H}^n .

1. INTRODUCTION

The Sobolev embedding theorem states that if $\dot{W}^{1,p}(\mathbb{R}^n)$ is the homogeneous Sobolev space, obtained by completing the set of compactly supported smooth functions $C_c^\infty(\mathbb{R}^n)$ under the norm $\|\nabla u\|_{L^p(\mathbb{R}^n)}$, then $\dot{W}^{1,p}(\mathbb{R}^n)$ embeds into $L^{p^*}(\mathbb{R}^n)$, whenever $n \geq 2$, $1 \leq p < n$ and $\frac{1}{p^*} = \frac{1}{p} - \frac{1}{n}$. This fails when $p = n$, i.e. $\dot{W}^{1,n}(\mathbb{R}^n)$ does not embed into $L^\infty(\mathbb{R}^n)$. One of the well-known remedies of this failure is to say that $\dot{W}^{1,n}(\mathbb{R}^n)$ embeds into $\text{BMO}(\mathbb{R}^n)$, the space of functions of bounded mean oscillation. In [2, 4], Bourgain and Brezis established another remedy of the failure of this Sobolev embedding for $\dot{W}^{1,n}(\mathbb{R}^n)$. They proved, among other things, that if X is a differential ℓ -form on \mathbb{R}^n with $\dot{W}^{1,n}(\mathbb{R}^n)$ coefficients, where $1 \leq \ell \leq n - 1$, then there exists a differential ℓ -form Y , whose components are all in $\dot{W}^{1,n} \cap L^\infty(\mathbb{R}^n)$, such that

$$dY = dX,$$

with

$$\|Y\|_{\dot{W}^{1,n} \cap L^\infty} \leq C \|dX\|_{L^n}.$$

(Such a theorem would have been trivial by Hodge decomposition, if $\dot{W}^{1,n}(\mathbb{R}^n)$ were to embed into $L^\infty(\mathbb{R}^n)$.) The existing proofs of the above theorem are all long and complicated. On the contrary, a weaker version of this theorem, where one replaces the space $\dot{W}^{1,n} \cap L^\infty$ by L^∞ , can be obtained from the following theorem of Van Schaftingen [9], when $\ell \leq n - 2$:

Theorem 1 (Van Schaftingen [9]). *Suppose f is a smooth vector field on \mathbb{R}^n , with*

$$\operatorname{div} f = 0.$$

Then for any compactly supported smooth vector field ϕ on \mathbb{R}^n , we have

$$(1) \quad \left| \int_{\mathbb{R}^n} \langle f, \phi \rangle \right| \leq C \|f\|_{L^1} \|\nabla \phi\|_{L^n},$$

where $\langle \cdot, \cdot \rangle$ is the pointwise Euclidean inner product of two vector fields in \mathbb{R}^n .

See e.g. Bourgain and Brezis [4], Lanzani and Stein [6]. We refer the interested reader to the survey in [10], for a more detailed account of this circle of ideas.

The original direct proof of Theorem 1 in [9] proceeds by decomposing

$$\int_{\mathbb{R}^n} \langle f, \phi \rangle = \sum_{i=1}^m \int_{\mathbb{R}} \left(\int_{\mathbb{R}^{i-1} \times \{s\} \times \mathbb{R}^{n-i}} f_i \phi_i \right) ds,$$

and by estimating first directly the innermost $(n-1)$ -dimensional integral. This gives the impression that the strategy is quite rigid. The first goal of this note is to prove Theorem 1 by averaging a suitable estimate over all *unit spheres* in \mathbb{R}^n .

In a second part of this paper, we adapt this idea of averaging over families of sets to prove an analogue of Theorem 1, in the setting where \mathbb{R}^n is replaced by the real hyperbolic space \mathbb{H}^n :

Theorem 2. *Suppose f is a smooth vector field on \mathbb{H}^n , with*

$$\operatorname{div}_g f = 0$$

where div_g is the divergence with respect to the metric g on \mathbb{H}^n . Then for any compactly supported smooth vector field ϕ on \mathbb{H}^n , we have

$$(2) \quad \left| \int_{\mathbb{H}^n} \langle f, \phi \rangle_g dV_g \right| \leq C \|f\|_{L^1(\mathbb{H}^n)} \|\nabla_g \phi\|_{L^n(\mathbb{H}^n)}.$$

where $\langle \cdot, \cdot \rangle_g$ and dV_g are the pointwise inner product and the volume measure with respect to g respectively, $\nabla_g \phi$ is the $(1,1)$ tensor given by the Levi-Civita connection of ϕ with respect to g , and

$$\|f\|_{L^1(\mathbb{H}^n)} = \int_{\mathbb{H}^n} |f|_g dV_g, \quad \|\nabla_g \phi\|_{L^n(\mathbb{H}^n)} = \left(\int_{\mathbb{H}^n} |\nabla_g \phi|_g^n dV_g \right)^{1/n}.$$

We note that the above theorem is formulated entirely geometrically on \mathbb{H}^n , without the need of specifying a choice of coordinate chart. As explained in Appendix A, Theorem 2 can be proved indirectly by

patching together known estimates on \mathbb{R}^n via a partition of unity, and by applying Hardy's inequality to get rid of lower order terms.

We shall prove Theorem 2 by averaging a suitable estimate over a family of hypersurfaces in \mathbb{H}^n , where the family of hypersurfaces is obtained from the orbit of a “vertical hyperplane” under all isometries in \mathbb{H}^n . The latter shares a similar flavour to the proof we will give below of Theorem 1. The main innovation there is in deducing Theorem 2 from Proposition 6, and in establishing Lemma 10 (see Section 3 for details).

Acknowledgments. S. Chanillo was partially supported by NSF grant DMS 1201474. J. Van Schaftingen was partially supported by the Fonds de la Recherche Scientifique-FNRS. P.-L. Yung was partially supported by a Titchmarsh Fellowship at the University of Oxford, a junior research fellowship at St. Hilda's College, a direct grant for research from the Chinese University of Hong Kong (3132713), and an Early Career Grant CUHK24300915 from the Hong Kong Research Grant Council.

2. ANOTHER PROOF OF THEOREM 1

Theorem 1 will follow from the following Proposition:

Proposition 3. *Let f, ϕ be as in Theorem 1. Write \mathbb{B}^n for the unit ball $\{x \in \mathbb{R}^n: |x| < 1\}$ in \mathbb{R}^n , and \mathbb{S}^{n-1} for the unit sphere (i.e. the boundary of \mathbb{B}^n). Also write $d\sigma$ for the standard surface measure on \mathbb{S}^{n-1} , and ν for the outward unit normal to the sphere \mathbb{S}^{n-1} . Then*

$$(3) \quad \left| \int_{\mathbb{S}^{n-1}} \langle f, \nu \rangle \langle \phi, \nu \rangle d\sigma \right| \leq C \|f\|_{L^1(\mathbb{R}^n \setminus \mathbb{B}^n)}^{\frac{1}{n}} \|f\|_{L^1(\mathbb{S}^{n-1})}^{1-\frac{1}{n}} \|\phi\|_{W^{1,n}(\mathbb{S}^{n-1})}.$$

Here $\|\phi\|_{W^{1,n}(\mathbb{S}^{n-1})} = \|\phi\|_{L^n(\mathbb{S}^{n-1})} + \|\nabla_{\mathbb{S}^{n-1}} \phi\|_{L^n(\mathbb{S}^{n-1})}$, where $\nabla_{\mathbb{S}^{n-1}} \phi$ is the (1,1) tensor on \mathbb{S}^{n-1} given by the covariant derivative of the vector field ϕ .

The proof of Proposition 3 in turn depends on the following two lemmas. The first one is a simple lemma about integration by parts:

Lemma 4. *Let f, ν be as in Proposition 3. Then for any compactly supported smooth function ψ on \mathbb{R}^n , we have*

$$\int_{\mathbb{S}^{n-1}} \langle f, \nu \rangle \psi d\sigma = - \int_{\mathbb{R}^n \setminus \mathbb{B}^n} \langle f, \nabla \psi \rangle dx.$$

The second one is a decomposition lemma for functions on the sphere \mathbb{S}^{n-1} :

Lemma 5. *Let φ be a smooth function on \mathbb{S}^{n-1} . For any $\lambda > 0$, there exists a decomposition*

$$\varphi = \varphi_1 + \varphi_2 \quad \text{on } \mathbb{S}^{n-1},$$

and an extension $\tilde{\varphi}_2$ of φ_2 to $\mathbb{R}^n \setminus \mathbb{B}^n$, such that $\tilde{\varphi}_2$ is smooth and bounded on $\mathbb{R}^n \setminus \mathbb{B}^n$, with

$$\begin{aligned} \|\varphi_1\|_{L^\infty(\mathbb{S}^{n-1})} &\leq C\lambda^{\frac{1}{n}} \|\nabla_{\mathbb{S}^{n-1}} \varphi\|_{L^n(\mathbb{S}^{n-1})}, \\ \|\nabla \tilde{\varphi}_2\|_{L^\infty(\mathbb{R}^n \setminus \mathbb{B}^n)} &\leq C\lambda^{\frac{1}{n}-1} \|\nabla_{\mathbb{S}^{n-1}} \varphi\|_{L^n(\mathbb{S}^{n-1})}. \end{aligned}$$

Here $|\nabla_{\mathbb{S}^{n-1}} \varphi|$ is the norm of the gradient of the function φ on \mathbb{S}^{n-1} . We postpone the proofs of Lemmas 4 and 5 to the end of this section.

Now we are ready for the proof of Proposition 3.

Proof of Proposition 3. Let f, ϕ be as in the statement of Theorem 1. Apply Lemma 5 to $\varphi = \langle \phi, \nu \rangle$, where $\lambda > 0$ is to be chosen. Then since

$$\|\nabla_{\mathbb{S}^{n-1}} \langle \phi, \nu \rangle\|_{L^n(\mathbb{S}^{n-1})} \leq C\|\phi\|_{W^{1,n}(\mathbb{S}^{n-1})},$$

there exists a decomposition

$$\langle \phi, \nu \rangle = \varphi_1 + \varphi_2 \quad \text{on } \mathbb{S}^{n-1},$$

and an extension $\tilde{\varphi}_2$ of φ_2 to $\mathbb{R}^n \setminus \mathbb{B}^n$, such that $\tilde{\varphi}_2 \in C^\infty \cap L^\infty(\mathbb{R}^n \setminus \mathbb{B}^n)$,

$$\|\varphi_1\|_{L^\infty(\mathbb{S}^{n-1})} \leq C\lambda^{\frac{1}{n}} \|\phi\|_{W^{1,n}(\mathbb{S}^{n-1})},$$

and

$$\|\nabla \tilde{\varphi}_2\|_{L^\infty(\mathbb{R}^n \setminus \mathbb{B}^n)} \leq C\lambda^{\frac{1}{n}-1} \|\phi\|_{W^{1,n}(\mathbb{S}^{n-1})}.$$

Now

$$\begin{aligned} \int_{\mathbb{S}^{n-1}} \langle f, \nu \rangle \langle \phi, \nu \rangle \, d\sigma &= \int_{\mathbb{S}^{n-1}} \langle f, \nu \rangle \varphi_1 \, d\sigma + \int_{\mathbb{S}^{n-1}} \langle f, \nu \rangle \varphi_2 \, d\sigma \\ &= I + II. \end{aligned}$$

In the first term, we estimate trivially

$$|I| \leq \|f\|_{L^1(\mathbb{S}^{n-1})} \|\varphi_1\|_{L^\infty(\mathbb{S}^{n-1})} \leq C\lambda^{\frac{1}{n}} \|f\|_{L^1(\mathbb{S}^{n-1})} \|\phi\|_{W^{1,n}(\mathbb{S}^{n-1})}.$$

To estimate the second term, we let θ be a smooth cut-off function with compact support on \mathbb{R}^n such that $\theta(x) = 1$ whenever $|x| \leq 1$. For $\varepsilon \in (0, 1)$, let $\theta_\varepsilon(x) = \theta(\varepsilon x)$. Then $\theta_\varepsilon = 1$ on \mathbb{S}^{n-1} , so we can rewrite II as

$$II = \int_{\mathbb{S}^{n-1}} \langle f, \nu \rangle \varphi_2 \theta_\varepsilon \, d\sigma$$

for any $\varepsilon \in (0, 1)$. We then integrate by parts using Lemma 4, with $\psi := \tilde{\varphi}_2 \theta_\varepsilon$, and obtain

$$II = - \int_{\mathbb{R}^n \setminus \mathbb{B}^n} \langle f, \nabla \tilde{\varphi}_2 \rangle \theta_\varepsilon \, dx - \int_{\mathbb{R}^n \setminus \mathbb{B}^n} \langle f, \nabla \theta_\varepsilon \rangle \tilde{\varphi}_2 \, dx.$$

(The cut-off function θ_ε is inserted so that ψ has compact support.) We now let $\varepsilon \rightarrow 0^+$. The second term then tends to 0, since it is just

$$-\varepsilon \int_{\mathbb{R}^n \setminus \mathbb{B}^n} \langle f(x), (\nabla \theta)(\varepsilon x) \rangle \tilde{\varphi}_2(x) \, dx,$$

where $f \in L^1$, $\nabla \theta(\varepsilon \cdot) \in L^\infty$ and $\tilde{\varphi}_2 \in L^\infty$ on $\mathbb{R}^n \setminus \mathbb{B}^n$. On the other hand, the first term tends to

$$-\int_{\mathbb{R}^n \setminus \mathbb{B}^n} \langle f, \nabla \tilde{\varphi}_2 \rangle \, dx$$

by dominated convergence theorem, since $f \in L^1$ and $\nabla \tilde{\varphi}_2 \in L^\infty$ on $\mathbb{R}^n \setminus \mathbb{B}^n$. As a result,

$$II = -\int_{\mathbb{R}^n \setminus \mathbb{B}^n} \langle f, \nabla \tilde{\varphi}_2 \rangle \, dx,$$

from which we see that

$$|II| \leq \|f\|_{L^1(\mathbb{R}^n \setminus \mathbb{B}^n)} \|\nabla \tilde{\varphi}_2\|_{L^\infty(\mathbb{R}^n \setminus \mathbb{B}^n)} \leq C \lambda^{\frac{1}{n}-1} \|f\|_{L^1(\mathbb{R}^n \setminus \mathbb{B}^n)} \|\phi\|_{W^{1,n}(\mathbb{S}^{n-1})}.$$

Together, by choosing $\lambda = \frac{\|f\|_{L^1(\mathbb{R}^n \setminus \mathbb{B}^n)}}{\|f\|_{L^1(\mathbb{S}^{n-1})}}$, we get

$$\left| \int_{\mathbb{S}^{n-1}} \langle f, \nu \rangle \langle \phi, \nu \rangle \, d\sigma \right| \leq C \|f\|_{L^1(\mathbb{R}^n \setminus \mathbb{B}^n)}^{\frac{1}{n}} \|f\|_{L^1(\mathbb{S}^{n-1})}^{1-\frac{1}{n}} \|\phi\|_{W^{1,n}(\mathbb{S}^{n-1})}$$

as desired. \square

We will now deduce Theorem 1 from Proposition 3. The idea is to average (3) over all unit spheres in \mathbb{R}^n .

Proof of Theorem 1. First, for each fixed $x \in \mathbb{R}^n$, we have

$$(4) \quad \langle f(x), \phi(x) \rangle = c \int_{\mathbb{S}^{n-1}} \langle f(x), \omega \rangle \langle \phi(x), \omega \rangle \, d\sigma(\omega)$$

where we are identifying $\omega \in \mathbb{S}^{n-1}$ with the corresponding unit tangent vector to \mathbb{R}^n based at the point x . Hence to estimate $\int_{\mathbb{R}^n} \langle f(x), \phi(x) \rangle \, dx$, it suffices to estimate

$$\int_{\mathbb{R}^n} \int_{\mathbb{S}^{n-1}} \langle f(x), \omega \rangle \langle \phi(x), \omega \rangle \, d\sigma(\omega) \, dx,$$

which is the same as

$$\int_{\mathbb{R}^n} \int_{\mathbb{S}^{n-1}} \langle f(z + \omega), \omega \rangle \langle \phi(z + \omega), \omega \rangle \, d\sigma(\omega) \, dz$$

by a change of variable $(x, \omega) \mapsto (z + \omega, \omega)$. Now when $z = 0$, the inner integral can be estimated by Proposition 3; for a general $z \neq 0$, one can still estimate the inner integral by Proposition 3, since the

proposition is invariant under translations. Thus the above double integral is bounded, in absolute value, by

$$C\|f\|_{L^1(\mathbb{R}^n)}^{\frac{1}{n}} \int_{\mathbb{R}^n} \|f(z + \cdot)\|_{L^1(\mathbb{S}^{n-1})}^{1-\frac{1}{n}} \|\phi(z + \cdot)\|_{W^{1,n}(\mathbb{S}^{n-1})} dz.$$

Applying Hölder's inequality to the last integral in z , one bounds this by

$$C\|f\|_{L^1(\mathbb{R}^n)}^{\frac{1}{n}} \left(\int_{\mathbb{R}^n} \|f(z + \cdot)\|_{L^1(\mathbb{S}^{n-1})} dz \right)^{1-\frac{1}{n}} \left(\int_{\mathbb{R}^n} \|\phi(z + \cdot)\|_{W^{1,n}(\mathbb{S}^{n-1})}^n dz \right)^{\frac{1}{n}}.$$

Since

$$\int_{\mathbb{R}^n} \|f(z + \cdot)\|_{L^1(\mathbb{S}^{n-1})} dz = c\|f\|_{L^1(\mathbb{R}^n)}$$

and

$$\int_{\mathbb{R}^n} \|\phi(z + \cdot)\|_{W^{1,n}(\mathbb{S}^{n-1})}^n dz \leq c \left(\|\phi\|_{L^n(\mathbb{R}^n)}^n + \|\nabla \phi\|_{L^n(\mathbb{R}^n)}^n \right),$$

we proved that under the assumption of Theorem 1, we have

$$(5) \quad \left| \int_{\mathbb{R}^n} \langle f, \phi \rangle \right| \leq C\|f\|_{L^1(\mathbb{R}^n)} \left(\|\nabla \phi\|_{L^n(\mathbb{R}^n)} + \|\phi\|_{L^n(\mathbb{R}^n)} \right).$$

This is almost the desired conclusion, except that we have an additional zeroth order term on ϕ on the right hand side of the estimate. But that can be scaled away by homogeneity. In fact, if f and ϕ satisfies the assumption of Theorem 1, then so does the dilations

$$f_\varepsilon(x) := \varepsilon^{-n} f(\varepsilon^{-1}x), \quad \phi_\varepsilon(x) := \phi(\varepsilon^{-1}x), \quad \varepsilon > 0.$$

Applying (5) to f_ε and ϕ_ε instead, we get

$$\left| \int_{\mathbb{R}^n} \langle f_\varepsilon, \phi_\varepsilon \rangle \right| \leq C\|f_\varepsilon\|_{L^1(\mathbb{R}^n)} \left(\|\nabla \phi_\varepsilon\|_{L^n(\mathbb{R}^n)} + \|\phi_\varepsilon\|_{L^n(\mathbb{R}^n)} \right),$$

i.e.

$$\left| \int_{\mathbb{R}^n} \langle f, \phi \rangle \right| \leq C\|f\|_{L^1(\mathbb{R}^n)} \left(\|\nabla \phi\|_{L^n(\mathbb{R}^n)} + \varepsilon \|\phi\|_{L^n(\mathbb{R}^n)} \right),$$

so letting $\varepsilon \rightarrow 0^+$, we get the desired conclusion of Theorem 1. \square

Proof of Lemma 4. Note that $\langle f, \nu \rangle \psi = \langle \psi f, \nu \rangle$, and ν is the inward unit normal to $\partial(\mathbb{R}^n \setminus \mathbb{B}^n)$. So by the divergence theorem on \mathbb{R}^n , we have

$$\int_{\mathbb{S}^{n-1}} \langle f, \nu \rangle \psi d\sigma = - \int_{\mathbb{R}^n \setminus \mathbb{B}^n} \operatorname{div}(\psi f) dx.$$

But since $\operatorname{div} f = 0$, we have

$$\operatorname{div}(\psi f) = \langle f, \nabla \psi \rangle + \psi \operatorname{div} f = \langle f, \nabla \psi \rangle,$$

and the desired equality follows. \square

Proof of Lemma 5. Suppose φ and λ are as in Lemma 5. We will construct first a decomposition $\varphi = \varphi_1 + \varphi_2$ on \mathbb{S}^{n-1} , so that both φ_1 and φ_2 are smooth on \mathbb{S}^{n-1} , and

$$\|\varphi_1\|_{L^\infty(\mathbb{S}^{n-1})} \leq C\lambda^{\frac{1}{n}} \|\nabla_{\mathbb{S}^{n-1}} \varphi\|_{L^n(\mathbb{S}^{n-1})},$$

$$\|\nabla_{\mathbb{S}^{n-1}} \varphi_2\|_{L^\infty(\mathbb{S}^{n-1})} \leq C\lambda^{\frac{1}{n}-1} \|\nabla_{\mathbb{S}^{n-1}} \varphi\|_{L^n(\mathbb{S}^{n-1})}.$$

(Here $\nabla_{\mathbb{S}^{n-1}} \varphi_2$ is the gradient on \mathbb{S}^{n-1} .) Once this is established, the lemma will follow, by extending φ_2 so that it is homogeneous of degree 0; in other words, we will then define

$$\tilde{\varphi}_2(x) := \varphi_2\left(\frac{x}{|x|}\right), \quad |x| \geq 1.$$

It is then straight forward to verify that

$$\|\nabla \tilde{\varphi}_2\|_{L^\infty(\mathbb{R}^n \setminus \mathbb{B}^n)} \leq C\lambda^{\frac{1}{n}-1} \|\nabla_{\mathbb{S}^{n-1}} \varphi\|_{L^n(\mathbb{S}^{n-1})},$$

since the radial derivative of $\tilde{\varphi}_2$ is zero.

To construct the desired decomposition on \mathbb{S}^{n-1} , we proceed as follows.

If $\lambda \geq 1$, we set $\varphi_2 = f_{\mathbb{S}^n} \varphi$, so that $\nabla_{\mathbb{S}^{n-1}} \varphi_2 = 0$ on \mathbb{S}^{n-1} ; then

$$\varphi_1 = \varphi - \int_{\mathbb{S}^n} \varphi,$$

and the estimate for φ_1 follows from the classical Morrey–Sobolev estimate.

If $0 < \lambda < 1$, we pick a non-negative radial cut-off function $\eta \in C_c^\infty(\mathbb{R}^n)$, with $\eta = 1$ in a neighborhood of 0, and define $\eta_\lambda(x) = \eta(\lambda^{-1}x)$ for $x \in \mathbb{R}^n$. We then consider the function

$$x \in \mathbb{R}^n \mapsto \int_{\mathbb{S}^{n-1}} \eta_\lambda(x - y) d\sigma(y).$$

When restricted to $x \in \mathbb{S}^{n-1}$, this function is a constant independent of the choice of $x \in \mathbb{S}^{n-1}$, by rotation invariance of the integral. We then write c_λ for this constant, i.e.

$$(6) \quad c_\lambda := \int_{\mathbb{S}^{n-1}} \eta_\lambda(x - y) d\sigma(y), \quad |x| = 1.$$

Note that by our choice of η , when $0 < \lambda < 1$,

$$(7) \quad c_\lambda \simeq \lambda^{n-1}.$$

Now we define, for $x \in \mathbb{S}^{n-1}$, that

$$\varphi_2(x) = c_\lambda^{-1} \int_{\mathbb{S}^{n-1}} \eta_\lambda(x - y) \varphi(y) d\sigma(y), \quad \varphi_1(x) = \varphi(x) - \varphi_2(x).$$

Then for $x \in \mathbb{S}^{n-1}$, we have

$$\varphi_1(x) = c_\lambda^{-1} \int_{\mathbb{S}^{n-1}} \eta_\lambda(x-y)[\varphi(x) - \varphi(y)] d\sigma(y)$$

by definition of c_λ . But for $x, y \in \mathbb{S}^{n-1}$, we have, by Morrey's embedding, that

$$|\varphi(x) - \varphi(y)| \leq C|x-y|^{\frac{1}{n}} \|\nabla_{\mathbb{S}^{n-1}} \varphi\|_{L^n(\mathbb{S}^{n-1})}.$$

It follows that

$$|\varphi_1(x)| \leq C \|\nabla_{\mathbb{S}^{n-1}} \varphi\|_{L^n(\mathbb{S}^{n-1})} c_\lambda^{-1} \int_{\mathbb{S}^{n-1}} \eta_\lambda(x-y) |x-y|^{\frac{1}{n}} d\sigma(y)$$

Letting $\tilde{\eta}(x) = |x|^{\frac{1}{n}} \eta(x)$, and $\tilde{\eta}_\lambda(x) = \tilde{\eta}(\lambda^{-1}x)$, we see that the right hand side above is just

$$C \lambda^{\frac{1}{n}} \|\nabla_{\mathbb{S}^{n-1}} \varphi\|_{L^n(\mathbb{S}^{n-1})} c_\lambda^{-1} \int_{\mathbb{S}^{n-1}} \tilde{\eta}_\lambda(x-y) d\sigma(y).$$

But this last integral can be estimated by

$$\int_{\mathbb{S}^{n-1}} \tilde{\eta}_\lambda(x-y) d\sigma(y) \lesssim \lambda^{n-1},$$

by the support and L^∞ bound of $\tilde{\eta}_\lambda$. Hence using also (7), we see that

$$\|\varphi_1\|_{L^\infty(\mathbb{S}^{n-1})} \leq C \lambda^{\frac{1}{n}} \|\nabla_{\mathbb{S}^{n-1}} \varphi\|_{L^n(\mathbb{S}^{n-1})}$$

as desired.

Next, suppose $x \in \mathbb{S}^{n-1}$, and v is a unit tangent vector to \mathbb{S}^{n-1} at x . Then

$$(8) \quad \partial_v \phi_2(x) = \lambda^{-1} c_\lambda^{-1} \int_{\mathbb{S}^{n-1}} \langle v, \nabla \eta \rangle (\lambda^{-1}(x-y)) \varphi(y) d\sigma(y).$$

But if we differentiate both sides of the definition (6) of c_λ with respect to ∂_v , we see that

$$(9) \quad 0 = \int_{\mathbb{S}^{n-1}} \langle v, \nabla \eta \rangle (\lambda^{-1}(x-y)) d\sigma(y).$$

Multiplying (9) by $\lambda^{-1} c_\lambda^{-1} \phi(x)$, and subtracting that from (8), we get

$$\partial_v \phi_2(x) = \lambda^{-1} c_\lambda^{-1} \int_{\mathbb{S}^{n-1}} \langle v, \nabla \eta \rangle (\lambda^{-1}(x-y)) [\varphi(y) - \varphi(x)] d\sigma(y).$$

Using Morrey's embedding again, we see that

$$|\partial_v \phi_2(x)| \leq C \lambda^{-1} \|\nabla_{\mathbb{S}^{n-1}} \varphi\|_{L^n(\mathbb{S}^{n-1})} c_\lambda^{-1} \int_{\mathbb{S}^{n-1}} |\langle v, \nabla \eta \rangle| (\lambda^{-1}(x-y)) |x-y|^{\frac{1}{n}} d\sigma(y).$$

Letting $\bar{\eta}(x) = |x|^{\frac{1}{n}} |\langle v, \nabla \eta \rangle|(x)$, and $\bar{\eta}_\lambda(x) = \bar{\eta}(\lambda^{-1}x)$, we see that the right hand side above is just

$$C\lambda^{\frac{1}{n}}\lambda^{-1}\|\nabla_{\mathbb{S}^{n-1}}\varphi\|_{L^n(\mathbb{S}^{n-1})}c_\lambda^{-1}\int_{\mathbb{S}^{n-1}}\bar{\eta}_\lambda(x-y)\,d\sigma(y).$$

But this last integral can be estimated by

$$\int_{\mathbb{S}^{n-1}}\bar{\eta}_\lambda(x-y)\,d\sigma(y)\,d\sigma(y)\lesssim\lambda^{n-1},$$

by the support and L^∞ bound of $\bar{\eta}_\lambda$. Hence using also (7), we see that

$$\|\nabla_{\mathbb{S}^{n-1}}\varphi_2\|_{L^\infty(\mathbb{S}^{n-1})}\leq C\lambda^{\frac{1}{n}-1}\|\nabla_{\mathbb{S}^{n-1}}\varphi\|_{L^n(\mathbb{S}^{n-1})}$$

as desired. \square

3. A BORDERLINE SOBOLEV EMBEDDING ON THE REAL HYPERBOLIC SPACE \mathbb{H}^n

We now turn to a corresponding result on the real hyperbolic space \mathbb{H}^n . We will first give a direct proof in this current section, in the spirit of the earlier proof of Theorem 1 using spherical averages. In the appendix we give a less direct proof, using a variant of Theorem 1 on \mathbb{R}^n .

First we need some notations. We will use the upper half space model for the hyperbolic space. In other words, we take \mathbb{H}^n to be

$$\mathbb{H}^n = \mathbb{R}_+^n = \{x = (x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R} : x_n > 0\}$$

and the metric on \mathbb{H}^n to be

$$g := \frac{|dx|^2}{x_n^2}.$$

We will use the following orthonormal frame of vector fields

$$e_i := x_n \frac{\partial}{\partial x_i}, \quad i = 1, \dots, n,$$

at every point of \mathbb{H}^n . Note that if $j \neq n$, then

$$(10) \quad \nabla_{e_n} e_j = 0.$$

(Here $\nabla = \nabla_g$ is the Levi-Civita connection with respect to the hyperbolic metric g .) In fact, since $\{e_1, \dots, e_n\}$ is an orthonormal basis, for any $k = 1, \dots, n$, we have

$$\begin{aligned} \langle \nabla_{e_n} e_j, e_k \rangle_g &= \frac{1}{2} (\langle [e_n, e_j], e_k \rangle_g - \langle [e_n, e_k], e_j \rangle_g - \langle [e_j, e_k], e_n \rangle_g) \\ &= \frac{1}{2} (\langle e_j, e_k \rangle_g - \langle (1 - \delta_{kn})e_k, e_j \rangle_g - \langle -\delta_{kn}e_j, e_n \rangle_g) = 0. \end{aligned}$$

Also, we have

$$(11) \quad \nabla_{e_n} e_n = 0.$$

This is because if $j \neq n$, then

$$\langle \nabla_{e_n} e_n, e_j \rangle_g = -\langle e_n, \nabla_{e_n} e_j \rangle_g = 0$$

by (10), and

$$\langle \nabla_{e_n} e_n, e_n \rangle_g = \frac{1}{2} e_n (\langle e_n, e_n \rangle_g) = 0.$$

To prove Theorem 2, note that we only need to consider the case $n \geq 2$, since when $n = 1$,

$$\|\phi\|_{L^\infty(\mathbb{H}^1)} \leq \int_0^\infty |\partial_y \phi(y)| dy = \|\nabla_g \phi\|_{L^1(\mathbb{H}^1)},$$

and (2) follows trivially. Hence from now on we assume $n \geq 2$.

We will deduce Theorem 2 from the following Proposition:

Proposition 6. *Assume $n \geq 2$. Let f, ϕ be as in Theorem 2. Write S for the copy of $(n-1)$ -dimensional hyperbolic space inside \mathbb{H}^n , given by*

$$S = \{x \in \mathbb{H}^n : x_1 = 0\},$$

and X for the half-space

$$X = \{x \in \mathbb{H}^n : x_1 > 0\}$$

so that S is the boundary of X . Also write dV'_g for the volume measure on S with respect to the hyperbolic metric on S , and $\nu = e_1$ for the unit normal to S . Then

$$(12) \quad \left| \int_S \langle f, \nu \rangle_g \langle \phi, \nu \rangle_g dV'_g \right| \leq C \|f\|_{L^1(X)}^{\frac{1}{n}} \|f\|_{L^1(S)}^{1-\frac{1}{n}} \|\phi\|_{W^{1,n}(S)}.$$

Here $\|\phi\|_{W^{1,n}(S)} = \|\phi\|_{L^n(S)} + \|\nabla_g \phi\|_{L^n(S)}$, and all integrals on S on the right hand side are with respect to dV'_g .

The S will be called a vertical hyperplane in \mathbb{H}^n . It is a totally geodesic submanifold of \mathbb{H}^n . We will consider all hyperbolic hyperplanes in \mathbb{H}^n , that is the image of S under all isometries of \mathbb{H}^n . The set of all such hypersurfaces in \mathbb{H}^n will be denoted by \mathcal{S} ; it will consist of all Euclidean parallel translates of S in the x' -direction, and all Euclidean northern hemispheres whose centers lie on the plane $\{x_n = 0\}$.

The proof of Proposition 6 in turn depends on the following two lemmas. The first one is a simple lemma about integration by parts, which is the counterpart of Lemma 4:

Lemma 7. *Assume $n \geq 2$. Let f, S, X, ν be as in Proposition 6. Then for any compactly supported smooth function ψ on \mathbb{H}^n , we have*

$$\int_S \langle f, \nu \rangle_g \psi \, dV'_g = - \int_X \langle f, \nabla_g \psi \rangle_g \, dV_g.$$

The second one is a decomposition lemma for functions on S , which is the counterpart of Lemma 5:

Lemma 8. *Assume $n \geq 2$. Let φ be a smooth function with compact support on S . For any $\lambda > 0$, there exists a decomposition*

$$\varphi = \varphi_1 + \varphi_2 \quad \text{on } S,$$

and an extension $\tilde{\varphi}_2$ of φ_2 to \mathbb{H}^n , such that $\tilde{\varphi}_2$ is smooth with compact support on \mathbb{H}^n , and

$$\|\varphi_1\|_{L^\infty(S)} \leq C\lambda^{\frac{1}{n}} \|\nabla_g \varphi\|_{L^n(S)},$$

with

$$\|\nabla_g \tilde{\varphi}_2\|_{L^\infty(\mathbb{H}^n)} \leq C\lambda^{\frac{1}{n}-1} \|\nabla_g \varphi\|_{L^n(S)}.$$

We postpone the proofs of Lemmas 7 and 8 to the end of this section.

Now we are ready for the proof of Proposition 6.

Proof of Proposition 6. Let f, ϕ be as in the statement of Theorem 2. Apply Lemma 8 to $\varphi = \langle \phi, \nu \rangle_g$, where $\lambda > 0$ is to be chosen. Then since

$$\|\nabla_g \langle \phi, \nu \rangle_g\|_{L^n(S)} \leq C\|\phi\|_{W^{1,n}(S)},$$

(this follows since $|e_k \langle \phi, \nu \rangle| = |\langle \nabla_{e_k} \phi, \nu \rangle + \langle \phi, \nabla_{e_k} \nu \rangle| \leq |\nabla_g \phi|_g + |\phi|_g$ for all k), there exists a decomposition

$$\langle \phi, \nu \rangle_g = \varphi_1 + \varphi_2 \quad \text{on } S,$$

and an extension $\tilde{\varphi}_2$ of φ_2 to \mathbb{H}^n , such that $\tilde{\varphi}_2 \in C_c^\infty(\mathbb{H}^n)$,

$$\|\varphi_1\|_{L^\infty(S)} \leq C\lambda^{\frac{1}{n}} \|\phi\|_{W^{1,n}(S)},$$

and

$$\|\nabla_g \tilde{\varphi}_2\|_{L^\infty(\mathbb{H}^n)} \leq C\lambda^{\frac{1}{n}-1} \|\phi\|_{W^{1,n}(S)}.$$

Now

$$\begin{aligned} \int_S \langle f, \nu \rangle_g \langle \phi, \nu \rangle_g \, dV'_g &= \int_S \langle f, \nu \rangle_g \varphi_1 \, dV'_g + \int_S \langle f, \nu \rangle_g \varphi_2 \, dV'_g \\ &= I + II. \end{aligned}$$

In the first term, we estimate trivially

$$|I| \leq \|f\|_{L^1(S)} \|\varphi_1\|_{L^\infty(S)} \leq C\lambda^{\frac{1}{n}} \|f\|_{L^1(S)} \|\phi\|_{W^{1,n}(S)}.$$

In the second term, we first integrate by parts using Lemma 7, with $\psi = \tilde{\varphi}_2$, and obtain

$$II = - \int_X \langle f, \nabla_g \tilde{\varphi}_2 \rangle_g dV_g,$$

so

$$\begin{aligned} |II| &\leq \|f\|_{L^1(X)} \|\nabla_g \tilde{\varphi}_2\|_{L^\infty(\mathbb{H}^n)} \\ &\leq C \lambda^{\frac{1}{n}-1} \|f\|_{L^1(X)} \|\phi\|_{W^{1,n}(S)}. \end{aligned}$$

Together, by choosing $\lambda = \frac{\|f\|_{L^1(X)}}{\|f\|_{L^1(S)}}$, we get

$$\left| \int_S \langle f, \nu \rangle_g \langle \phi, \nu \rangle_g dV'_g \right| \leq C \|f\|_{L^1(X)}^{\frac{1}{n}} \|f\|_{L^1(S)}^{1-\frac{1}{n}} \|\phi\|_{W^{1,n}(S)}$$

as desired. \square

We will now deduce Theorem 2 from Proposition 6. The idea is to average (12) over all images of S under isometries in \mathbb{H}^n (i.e. all hypersurfaces in the collection \mathcal{S}).

Proof of Theorem 2. First, for each fixed $x = (x', x_n) \in \mathbb{H}^n$, we have the following analogue of the identity (4), used in the proof of Theorem 1:

$$\langle f(x), \phi(x) \rangle_g = c \int_{\mathbb{S}^{n-1}} \langle f(x), x_n \omega \rangle_g \langle \phi(x), x_n \omega \rangle_g d\sigma(\omega)$$

Here we are identifying $\omega \in \mathbb{S}^{n-1}$ with the corresponding tangent vector to \mathbb{H}^n based at the point x . (Note then $x_n \omega$ has length 1 with respect to the metric g at x , so $x_n \omega$ belongs to the unit sphere bundle at x .) Furthermore, since the above integrand is even in ω , we may replace the integral over \mathbb{S}^{n-1} by the integral only over the northern hemisphere $\mathbb{S}_+^{n-1} := \{\omega \in \mathbb{S}^{n-1} : \omega_n > 0\}$. Hence to estimate $\int_{\mathbb{H}^n} \langle f(x), \phi(x) \rangle_g dV_g$, it suffices to estimate

$$(13) \quad \int_{\mathbb{R}_+^n} \int_{\mathbb{S}_+^{n-1}} \langle f(x), x_n \omega \rangle_g \langle \phi(x), x_n \omega \rangle_g d\sigma(\omega) \frac{dx}{x_n^n}.$$

We will compute this integral by making a suitable change of variables.

To do so, given $x \in \mathbb{R}_+^n$ and $\omega \in \mathbb{S}_+^{n-1}$, let $S(x, \omega)$ be the hyperbolic hypersurface in \mathcal{S} passing through x with normal vector ω at x . In other words, $S(x, \omega)$ would be an Euclidean hemisphere, with center on the plane $\{x_n = 0\}$; we denote the center of this Euclidean hemisphere by $(z, 0)$, where $z = z(x, \omega)$.

For each fixed $x \in \mathbb{R}_+^n$, the map $\omega \mapsto z(x, \omega)$ provides an invertible change of variables from \mathbb{S}_+^{n-1} to \mathbb{R}^{n-1} . Thus we are led to parametrize

the integral in (13) by z instead of ω . In order to do that we observe that the vectors $x - (z, 0)$ and ω are collinear. This implies that if $z = z(x, \omega)$, then

$$\omega = \frac{x - (z, 0)}{|x - (z, 0)|}.$$

(Here $|x - (z, 0)|$ is the Euclidean norm of $x - (z, 0)$.) Write $\Phi_x(z)$ for the right hand side of the above equation. We view Φ_x as a map $\Phi_x : \mathbb{R}^{n-1} \rightarrow \mathbb{S}_+^{n-1} \subset \mathbb{R}^n$, and compute the Jacobian of the map. We have

$$(D\Phi_x)^t(z) = \frac{1}{|x - (z, 0)|} \left((-I, 0) + \frac{(x' - z) \otimes (x - (z, 0))^t}{|x - (z, 0)|^2} \right),$$

(here we think of x, z as column vectors, and $D\Phi_x$ as an $(n-1) \times n$ matrix). Thus

$$(D\Phi_x)^t D\Phi_x(z) = \frac{1}{|x - (z, 0)|^2} \left(I - \frac{(x' - z) \otimes (x' - z)^t}{|x - (z, 0)|^2} \right).$$

By computing the determinant in a basis that contains $x' - z$, we get

$$\begin{aligned} \text{Jac } \Phi_x(z) &= \sqrt{\det[(D\Phi_x)^t D\Phi_x(z)]} \\ &= \frac{1}{|x - (z, 0)|^{n-1}} \sqrt{1 - \frac{|x' - z|^2}{|x - (z, 0)|^2}} \\ &= \frac{x_n}{|x - (z, 0)|^n}. \end{aligned}$$

By a change of variable $\omega = \Phi_x(z)$, and using Fubini's theorem, we see that (13) is equal to

$$(14) \quad \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}_+^n} \frac{\langle f(x), x_n \Phi_x(z) \rangle_g \langle \phi(x), x_n \Phi_x(z) \rangle_g}{|x - (z, 0)|^n} \frac{dx}{x_n^{n-1}} dz.$$

Now we fix $z \in \mathbb{R}^{n-1}$, and compute the inner integral over x by integrating over successive hemispheres of radius r centered at $(z, 0)$. More precisely, let $S(z, r)$ be the Euclidean northern hemisphere with center $(z, 0)$ and of radius $r > 0$. Then $S(z, r) \in \mathcal{S}$, and for any $z \in \mathbb{R}^{n-1}$, we have

$$\begin{aligned} & \int_{\mathbb{R}_+^n} \frac{\langle f(x), x_n \Phi_x(z) \rangle_g \langle \phi(x), x_n \Phi_x(z) \rangle_g}{|x - (z, 0)|^n} \frac{dx}{x_n^{n-1}} \\ &= \int_0^\infty \int_{x \in S(z, r)} \langle f(x), x_n \Phi_x(z) \rangle_g \langle \phi(x), x_n \Phi_x(z) \rangle_g \frac{d\sigma(x)}{x_n^{n-1}} \frac{dr}{r^n}, \end{aligned}$$

where $d\sigma(x)$ is the Euclidean surface measure on $S(z, r)$. However, if $x \in S(z, r)$, then $x_n \Phi_x(z)$ is precisely the upward unit normal to $S(z, r)$

at x . Also, if dV'_g is the induced surface measure on $S(z, r)$ from the hyperbolic metric on \mathbb{H}^n , then

$$dV'_g = \frac{d\sigma(x)}{x_n^{n-1}r};$$

indeed if we write $\omega = \frac{x-(z,0)}{|x-(z,0)|}$, then at $x \in S(z, r)$ we have

$$dV'_g = i_{x_n\omega} dV_g = i_{x_n\omega} \frac{dx}{x_n^n} = \frac{i_{r\omega} dx}{x_n^{n-1}r} = \frac{d\sigma(x)}{x_n^{n-1}r}.$$

(here i denotes the interior product of a vector with a differential form). Hence the integral (14) is just equal to

$$(15) \quad \int_{\mathbb{R}^{n-1}} \int_0^\infty \int_{S(z,r)} \langle f, \nu \rangle_g \langle \phi, \nu \rangle_g dV'_g \frac{dr}{r^{n-1}} dz.$$

By Proposition 6 and its invariance under isometries of the hyperbolic space \mathbb{H}^n , we have

$$\begin{aligned} & \left| \int_{S(z,r)} \langle f, \nu \rangle_g \langle \phi, \nu \rangle_g dV'_g \right| \\ & \leq C \|f\|_{L^1(\mathbb{H}^n)}^{\frac{1}{n}} \left(\int_{S(z,r)} |f|_g dV'_g \right)^{1-\frac{1}{n}} \left(\int_{S(z,r)} (|\nabla_g \phi|_g^n + |\phi|_g^n) dV'_g \right)^{\frac{1}{n}}. \end{aligned}$$

Hence by Hölder's inequality, (15) is bounded by

$$\begin{aligned} & C \|f\|_{L^1(\mathbb{H}^n)}^{\frac{1}{n}} \left(\int_{\mathbb{R}^{n-1}} \int_0^\infty \int_{S(z,r)} |f|_g dV'_g \frac{dr}{r^{n-1}} dz \right)^{1-\frac{1}{n}} \\ & \cdot \left(\int_{\mathbb{R}^{n-1}} \int_0^\infty \int_{S(z,r)} (|\nabla_g \phi|_g^n + |\phi|_g^n) dV'_g \frac{dr}{r^{n-1}} dz \right)^{\frac{1}{n}}. \end{aligned}$$

But undoing our earlier changes of variable, we see that

$$\begin{aligned} & \int_{\mathbb{R}^{n-1}} \int_0^\infty \int_{S(z,r)} |f|_g dV'_g \frac{dr}{r^{n-1}} dz \\ & = \int_{\mathbb{R}^{n-1}} \int_0^\infty \int_{x \in S(z,r)} |f(x)|_g \frac{d\sigma(x)}{x_n^{n-1}} \frac{dr}{r^n} dz \\ & = \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}_+^n} \frac{|f(x)|_g}{|x-(z,0)|^n} \frac{dx}{x_n^{n-1}} dz \\ & = \int_{\mathbb{R}_+^n} |f(x)|_g \left(\int_{\mathbb{R}^{n-1}} \frac{x_n}{|x-(z,0)|^n} dz \right) \frac{dx}{x_n^n} \\ & = C \|f\|_{L^1(\mathbb{H}^n)}. \end{aligned}$$

Similarly,

$$\int_{\mathbb{R}^{n-1}} \int_0^\infty \int_{S(z,r)} (|\nabla_g \phi|_g^n + |\phi|_g^n) dV'_g \frac{dr}{r^{n-1}} dz = C(\|\nabla_g \phi\|_{L^n(\mathbb{H}^n)}^n + \|\phi\|_{L^n(\mathbb{H}^n)}^n).$$

Altogether, (15) (and hence (13)) is bounded by

$$C\|f\|_{L^1(\mathbb{H}^n)}(\|\nabla_g \phi\|_{L^n(\mathbb{H}^n)} + \|\phi\|_{L^n(\mathbb{H}^n)}).$$

This is almost what we want, except that on the right hand side we have an additional $\|\phi\|_{L^n(\mathbb{H}^n)}$. To fix this, one applies Lemma 9 below, with $p = n$, and the desired conclusion of Theorem 2 follows. \square

Lemma 9. *Assume $n \geq 2$. For any compactly supported smooth vector field ϕ on \mathbb{H}^n , and any $1 \leq p < \infty$, we have*

$$\|\phi\|_{L^p(\mathbb{H}^n)} \leq C\|\nabla_g \phi\|_{L^p(\mathbb{H}^n)}.$$

Proof. In fact, for any function $\Phi \in C_c^\infty(\mathbb{H}^n)$, and any exponent $1 \leq p < \infty$, we have, from Hardy's inequality, that

$$(16) \quad \|\Phi\|_{L^p(\mathbb{H}^n)} \leq C\|e_n \Phi\|_{L^p(\mathbb{H}^n)}.$$

This is because

$$\int_0^\infty |\Phi(x)|^p \frac{dx_n}{x_n^n} \leq \left(\frac{p}{n-1}\right)^p \int_0^\infty |e_n \Phi|^p(x) \frac{dx_n}{x_n^n}$$

by Hardy's inequality. (16) then follows by integrating over all $x' \in \mathbb{R}^{n-1}$ with respect to dx' . Now we apply (16) to $\Phi = \langle \phi, e_j \rangle$, $1 \leq j \leq n-1$. In view of (10), we have

$$e_n \Phi = \langle \nabla_{e_n} \phi, e_j \rangle + \langle \phi, \nabla_{e_n} e_j \rangle = \langle \nabla_{e_n} \phi, e_j \rangle,$$

so we get

$$\|\langle \phi, e_j \rangle\|_{L^p(\mathbb{H}^n)} \leq C\|\nabla_g \phi\|_{L^p(\mathbb{H}^n)}.$$

Similarly, we can apply (16) to $\Phi = \langle \phi, e_n \rangle$, and use (11) in place of (10). Altogether we see that

$$\|\phi\|_{L^p(\mathbb{H}^n)} \leq C\|\nabla_g \phi\|_{L^p(\mathbb{H}^n)},$$

as desired. \square

We now turn to the proofs of Lemma 7 and 8.

Proof of Lemma 7. Note that $\langle f, \nu \rangle_g \psi = \langle \psi f, \nu \rangle_g$, and ν is the inward unit normal to ∂X . Also dV'_g agrees with the induced surface measure on S from \mathbb{H}^n . So by the divergence theorem on \mathbb{H}^n , we have

$$\int_S \langle f, \nu \rangle_g \psi dV'_g = - \int_X \operatorname{div}_g(\psi f) dV_g.$$

But since $\operatorname{div}_g f = 0$, we have

$$\operatorname{div}_g(\psi f) = \langle f, \nabla_g \psi \rangle_g + \psi \operatorname{div}_g f = \langle f, \nabla_g \psi \rangle_g,$$

and the desired equality follows. \square

The proof of Lemma 8 will be easy, once we establish the following lemma:

Lemma 10. *Let φ be a smooth function with compact support on \mathbb{H}^m , $m \geq 1$. For any $p > m$ and $\lambda > 0$, there exists a decomposition*

$$\varphi = \varphi_1 + \varphi_2 \quad \text{on } \mathbb{H}^m,$$

such that φ_2 is smooth with compact support on \mathbb{H}^m , and

$$\|\varphi_1\|_{L^\infty(\mathbb{H}^m)} \leq C\lambda^{1-\frac{m}{p}} \|\nabla_g \varphi\|_{L^p(\mathbb{H}^m)},$$

with

$$\|\nabla_g \varphi_2\|_{L^\infty(\mathbb{H}^m)} \leq C\lambda^{-\frac{m}{p}} \|\nabla_g \varphi\|_{L^p(\mathbb{H}^m)}.$$

We postpone the proof of this lemma to the end of this section.

Proof of Lemma 8. Suppose φ and λ are as in Lemma 8. We identify S with \mathbb{H}^m , where $m = n - 1$. (This is possible because the restriction of the metric of \mathbb{H}^n to S induces a metric on S that is isometric to that of \mathbb{H}^m .) Using Lemma 10, with $p = n$, we obtain a decomposition $\varphi = \varphi_1 + \varphi_2$ on S , such that

$$\|\varphi_1\|_{L^\infty(S)} \leq C\lambda^{\frac{1}{n}} \|\nabla_g \varphi\|_{L^n(S)},$$

with

$$(17) \quad \|\nabla_g \varphi_2\|_{L^\infty(S)} \leq C\lambda^{\frac{1}{n}-1} \|\nabla_g \varphi\|_{L^n(S)}.$$

We then extend φ_2 to \mathbb{H}^n by setting for $(x_1, x'', x_n) \in \mathbb{R} \times \mathbb{R}^{n-2} \times \mathbb{R}_+$

$$\tilde{\varphi}_2(x_1, x'', x_n) = \varphi_2\left(0, x'', \sqrt{x_1^2 + x_n^2}\right).$$

One immediately checks that $\tilde{\varphi}_2$ is smooth with compact support on \mathbb{H}^n , with

$$\|\nabla_g \tilde{\varphi}_2\|_{L^\infty(\mathbb{H}^n)} \leq \|\nabla_g \varphi_2\|_{L^\infty(S)}.$$

In view of (17), we obtain the desired bound for $\nabla_g \tilde{\varphi}_2$. \square

It remains to prove Lemma 10.

Proof of Lemma 10. When $m = 1$, the 1-dimensional hyperbolic space \mathbb{H}^1 is isometric to \mathbb{R} , and Lemma 10 follows from its counterpart on \mathbb{R} (see e.g. [9]). Alternatively, it will follow from our treatment in the case $m \geq 2$, $0 < \lambda < 1$ below.

So assume from now on $m \geq 2$. Suppose φ is smooth with compact support on \mathbb{H}^m , $p > m$ and $\lambda > 0$. We will construct our desired

decomposition $\varphi = \varphi_1 + \varphi_2$. Recall that since $p > m$, the Morrey inequality on \mathbb{H}^m implies that

$$(18) \quad \|\varphi\|_{L^\infty(\mathbb{H}^m)} \leq C \|\nabla_g \varphi\|_{L^p(\mathbb{H}^m)}.$$

To see this, let $\zeta \in C_c^\infty(\mathbb{R})$ is a cut-off function such that $\zeta(s) = 1$ if $|s| \leq 1/2$, and $\zeta(s) = 0$ if $|s| \geq 1$. Let $x_0 := (0, 1) \in \mathbb{H}^m$, and let $\zeta_{x_0}(x) := \zeta(d(x, x_0))$, where d is the hyperbolic distance on \mathbb{H}^m . Consider the localization $\zeta_{x_0}\varphi$ of φ , to the unit ball centered at x_0 . It satisfies

$$(19) \quad \|\zeta_{x_0}\varphi\|_{W^{1,p}(\mathbb{R}^m)} \leq C \|\nabla_g \varphi\|_{L^p(\mathbb{H}^m)},$$

where the left-hand side is a shorthand for

$$\|\zeta_{x_0}\varphi\|_{L^p(\mathbb{R}^m)} + \|\nabla_e(\zeta_{x_0}\varphi)\|_{L^p(\mathbb{R}^m)};$$

here ∇_e denotes the Euclidean gradient of a function. (19) holds because by the support of ζ_{x_0} , we have

$$\|\zeta_{x_0}\varphi\|_{L^p(\mathbb{R}^m)} \simeq \|\zeta_{x_0}\varphi\|_{L^p(\mathbb{H}^m)} \leq \|\nabla_g \varphi\|_{L^p(\mathbb{H}^m)},$$

and

$$\begin{aligned} \|\nabla_e(\zeta_{x_0}\varphi)\|_{L^p(\mathbb{R}^m)} &\leq \|(\nabla_e \zeta_{x_0})\varphi\|_{L^p(\mathbb{R}^m)} + \|\zeta_{x_0}(\nabla_e \varphi)\|_{L^p(\mathbb{R}^m)} \\ &\leq C(\|\varphi\|_{L^p(\mathbb{H}^m)} + \|\nabla_g \varphi\|_{L^p(\mathbb{H}^m)}) \\ &\leq C \|\nabla_g \varphi\|_{L^p(\mathbb{H}^m)} \end{aligned}$$

where we have used Lemma 9 in the last inequalities (note that Lemma 9 applies now since $m \geq 2$). In particular, by Morrey's inequality on \mathbb{R}^m , we get, from (19), that

$$|\zeta_{x_0}(x)\varphi(x) - \zeta_{x_0}(y)\varphi(y)| \leq C \|\nabla_g \varphi\|_{L^p(\mathbb{H}^m)} |x - y|^{1-\frac{m}{p}}$$

for all $x, y \in \mathbb{R}^m$. Taking $x = x_0$ and $y \in \mathbb{H}^m$ such that $d(y, x_0) = 2$, we get that

$$|\varphi(x_0)| \leq C \|\nabla_g \varphi\|_{L^p(\mathbb{H}^m)}.$$

Since the isometry group of \mathbb{H}^m acts transitively on \mathbb{H}^m , and since the right hand side of the above inequality is invariant under isometries, we obtain

$$|\varphi(x)| \leq C \|\nabla_g \varphi\|_{L^p(\mathbb{H}^m)}.$$

for all $x \in \mathbb{H}^m$, and hence (18).

In particular, in view of (18), when $\lambda \geq 1$, it suffices to take $\varphi_1 = \varphi$ and $\varphi_2 = 0$. We then get the desired estimates for φ_1 and φ_2 trivially.

On the other hand, suppose now $0 < \lambda < 1$. We fix a compactly supported smooth function $\eta \in C_c^\infty(\mathbb{R}^m)$, with

$$\int_{\mathbb{R}^m} \eta(v) dv = 1.$$

For $x = (x', x_m) \in \mathbb{H}^m$, we define

$$\varphi_2(x) = \int_{\mathbb{R}^m} \varphi(x' + x_m e^{v_m} v', x_m e^{v_m}) \lambda^{-m} \eta(\lambda^{-1} v) dv$$

(where we wrote $v = (v', v_m) \in \mathbb{R}^{m-1} \times \mathbb{R}$), and define

$$\varphi_1(x) = \varphi(x) - \varphi_2(x).$$

Note that φ_2 is smooth with compact support on \mathbb{H}^m , and hence so is φ_1 . Now for $i = 1, 2, \dots, m-1$, we have

$$(e_i \varphi_2)(x, y) = \int_{\mathbb{R}^m} e^{-v_m} (e_i \varphi)(x' + x_m u' e^{v_m}, x_m e^{v_m}) \lambda^{-m} \eta(\lambda^{-1} v) dv.$$

Since $v \mapsto \eta(\lambda^{-1} v)$ has support uniformly bounded with respect to $0 < \lambda < 1$, we have $e^{-v_m} \leq C$ on the support of the integral, where C is independent of $0 < \lambda < 1$. Hence by Hölder's inequality, we have

$$\begin{aligned} & \|e_i \varphi_2\|_{L^\infty(\mathbb{H}^m)} \\ & \leq C \left(\int_{\mathbb{R}^m} |e_i \varphi|^p(x' + x_m e^{v_m} v', x_m e^{v_m}) dv \right)^{\frac{1}{p}} \|\lambda^{-m} \eta(\lambda^{-1} v)\|_{L^{p'}(dv)} \\ & = C \lambda^{-\frac{m}{p}} \left(\int_{\mathbb{R}_+^m} |e_i \varphi|^p(z) \frac{dz}{z_m^m} \right)^{\frac{1}{p}}, \end{aligned}$$

the last line following from the changes of variables $z_m = e^{v_m}$, and then $z' = x' + z_m v'$. We thus see that

$$\|e_i \varphi_2\|_{L^\infty(\mathbb{H}^m)} \leq C \lambda^{-\frac{m}{p}} \|e_i \varphi\|_{L^p(\mathbb{H}^m)},$$

as desired.

Furthermore, when $i = m$,

$$\begin{aligned} (e_m \varphi_2)(x) &= \int_{\mathbb{R}^m} \left[(e_m \varphi)(x' + x_m v' e^{v_m}, x_m e^{v_m}) \right. \\ & \quad \left. + \sum_{i=1}^{m-1} v_i (e_i \varphi)(x' + x_m v' e^{v_m}, x_m e^{v_m}) \right] \lambda^{-m} \eta(\lambda^{-1} v) dv. \end{aligned}$$

Using that $|v_i| \leq C$ on the support of the integrals (uniformly in $0 < \lambda < 1$), and Hölder's inequality as above, we see that

$$\|e_m \varphi_2\|_{L^\infty(\mathbb{H}^m)} \leq C \lambda^{-\frac{m}{p}} \|\nabla_g \varphi\|_{L^p(\mathbb{H}^m)},$$

as desired.

Finally, to estimate φ_1 , note that

$$\varphi(x) = \lim_{\lambda \rightarrow 0^+} \int_{\mathbb{R}^m} \varphi(x' + x_m e^{v_m} v', x_m e^{v_m}) \lambda^{-m} \eta(\lambda^{-1} v) dv.$$

Hence

$$\begin{aligned}\varphi_1(x) &= \varphi(x) - \varphi_2(x) \\ &= - \int_0^\lambda \int_{\mathbb{R}^m} \varphi(x' + x_m e^{v_m} v', x_m e^{v_m}) \frac{d}{ds} [s^{-m} \eta(s^{-1} v)] \, dv \, ds.\end{aligned}$$

But

$$-\frac{d}{ds} [s^{-m} \eta(s^{-1} v)] = \sum_{i=1}^m \frac{\partial}{\partial v_i} [s^{-m} (v_i \eta)(s^{-1} v)]$$

so we can plug this back in the equation for φ_1 , and integrate by parts in v . Now

$$\frac{\partial}{\partial v_i} [\varphi(x' + x_m e^{v_m} v', x_m e^{v_m})] = (e_i \varphi)(x' + x_m e^{v_m} v', x_m e^{v_m}), \quad i = 1, \dots, m-1.$$

and

$$\begin{aligned}\frac{\partial}{\partial v_m} [\varphi(x' + x_m e^{v_m} v', x_m e^{v_m})] &= (e_m \varphi)(x' + x_m e^{v_m} v', x_m e^{v_m}) \\ &\quad + \sum_{i=1}^{m-1} v_i (e_i \varphi)(x' + x_m e^{v_m} v', x_m e^{v_m})\end{aligned}$$

Hence

$$\begin{aligned}\varphi_1(x) &= - \int_0^\lambda \sum_{i=1}^m \int_{\mathbb{R}^m} (e_i \varphi)(x' + x_m e^{v_m} v', x_m e^{v_m}) s^{-m} (v_i \eta)(s^{-1} v) \, dv \, ds \\ &\quad - \int_0^\lambda \sum_{i=1}^{m-1} \int_{\mathbb{R}^m} v_i (e_i \varphi)(x' + x_m e^{v_m} v', x_m e^{v_m}) s^{-m} (v_i \eta)(s^{-1} v) \, dv \, ds.\end{aligned}$$

When $0 < \lambda < 1$, the integral in v in each term can now be estimated by Hölder's inequality, yielding

$$\begin{aligned}\|\varphi_1\|_{L^\infty(\mathbb{H}^n)} &\leq \int_0^\lambda C s^{-\frac{m}{p}} \|\nabla_g \varphi\|_{L^p(\mathbb{H}^m)} \, ds \\ &= C \lambda^{1-\frac{m}{p}} \|\nabla_g \varphi\|_{L^p(\mathbb{H}^m)}.\end{aligned}$$

This completes the proof of Lemma 10. \square

APPENDIX A. INDIRECT PROOF OF THEOREM 2

We will give an alternative proof of Theorem 2 from the following variant of Theorem 1, whose proof can be found, for instance, in Van Schaftingen [9] (it can also be deduced by a small modification of the proof we gave above of Theorem 1):

Proposition 11 (Van Schaftingen [9]). *Suppose f is a smooth vector field on \mathbb{R}^n (not necessarily $\operatorname{div} f = 0$). Then for any compactly supported smooth vector field ϕ on \mathbb{R}^n , we have*

$$(20) \quad \left| \int_{\mathbb{R}^n} \langle f, \phi \rangle \right| \leq C (\|f\|_{L^1} \|\nabla \phi\|_{L^n} + \|\operatorname{div} f\|_{L^1} \|\phi\|_{L^n}),$$

where $\langle \cdot, \cdot \rangle$ is the pointwise Euclidean inner product of two vector fields in \mathbb{R}^n .

To prove Theorem 2, we consider a function $\zeta \in C_c^\infty(\mathbb{R})$ and we define for $\alpha \in \mathbb{H}^n$ the function $\zeta_\alpha : \mathbb{H}^n \rightarrow \mathbb{R}$ by

$$\zeta_\alpha(x) = \zeta(d(x, \alpha)).$$

We assume that

$$\int_{\mathbb{H}^n} \zeta_\alpha^2(x) dV_g(\alpha) = 1.$$

Now given vector fields f and ϕ as in Theorem 2, we write

$$\int_{\mathbb{H}^n} \langle f, \phi \rangle_g dV_g = \int_{\mathbb{H}^n} \int_{\mathbb{H}^n} \langle \zeta_\alpha f(x), \zeta_\alpha \phi(x) \rangle_g dV_g(z) dV_g(\alpha).$$

If $\alpha = (0, 1) \in \mathbb{H}^n$, then

$$\int_{\mathbb{H}^n} \langle \zeta_\alpha f(x), \zeta_\alpha \phi(x) \rangle_g dV_g = \int_{\mathbb{R}_+^n} \langle \zeta_\alpha f, \frac{\zeta_\alpha \phi}{x_n^{n+2}} \rangle_e dx,$$

where $\langle \cdot, \cdot \rangle_e$ is the Euclidean inner product of two vectors. Hence by Proposition 11, this last integral is bounded by

$$C \left(\|\zeta_\alpha f\|_{L^1(\mathbb{R}^n)} \left\| \nabla_e \left(\frac{\zeta_\alpha \phi}{x_n^{n+2}} \right) \right\|_{L^n(\mathbb{R}^n)} + \|\langle \nabla_e(\zeta_\alpha), f \rangle_e\|_{L^1(\mathbb{R}^n)} \left\| \frac{\zeta_\alpha \phi}{x_n^{n+2}} \right\|_{L^n(\mathbb{R}^n)} \right)$$

where we write ∇_e to emphasize that the gradients are with respect to the Euclidean metric. Now on the support of ζ_α , we have $|x_n| \simeq 1$, so altogether we get

$$(21) \quad \left| \int_{\mathbb{H}^n} \langle \zeta_\alpha f, \zeta_\alpha \phi \rangle_g dV_g \right| \leq C \left(\|\zeta_\alpha f\|_{L^1(\mathbb{H}^n)} + \|\langle \nabla_g(\zeta_\alpha), f \rangle_g\|_{L^1(\mathbb{H}^n)} \right) \left(\|\nabla_g \phi\|_{L^n(\mathbb{H}^n)} + \|\phi\|_{L^n(\mathbb{H}^n)} \right).$$

This remains true even if $\alpha \neq (0, 1)$, since there is an isometry mapping α to $(0, 1)$, and since (21) is invariant under isometries of \mathbb{H}^n . By integrating with respect to $\alpha \in \mathbb{H}^n$, we see that

$$\left| \int_{\mathbb{H}^n} \langle f, \phi \rangle_g dV_g \right| \leq C \|f\|_{L^1(\mathbb{H}^n)} \left(\|\nabla_g \phi\|_{L^n(\mathbb{H}^n)} + \|\phi\|_{L^n(\mathbb{H}^n)} \right).$$

We now use Lemma 9 to bound $\|\phi\|_{L^n(\mathbb{H}^n)}$ by $\|\nabla_g \phi\|_{L^n(\mathbb{H}^n)}$. This concludes our alternative proof of Theorem 2.

REFERENCES

- [1] J. Bourgain and H. Brezis, *Sur l'équation $\operatorname{div} u = f$* , C. R. Math. Acad. Sci. Paris **334** (2002), no. 11, 973–976.
- [2] ———, *On the equation $\operatorname{div} Y = f$ and application to control of phases*, J. Amer. Math. Soc. **16** (2003), no. 2, 393–426.
- [3] ———, *New estimates for the Laplacian, the div-curl, and related Hodge systems*, C. R. Math. Acad. Sci. Paris **338** (2004), no. 7, 539–543.
- [4] ———, *New estimates for elliptic equations and Hodge type systems*, J. Eur. Math. Soc. (JEMS) **9** (2007), no. 2, 277–315.
- [5] E. Hebey, *Nonlinear analysis on manifolds: Sobolev spaces and inequalities*, Courant Lecture Notes in Mathematics, vol. 5, New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 1999.
- [6] L. Lanzani and E. M. Stein, *A note on div curl inequalities*, Math. Res. Lett. **12** (2005), no. 1, 57–61.
- [7] P. Li, *Lecture notes on geometric analysis*, Lecture Notes Series, vol. 6, Seoul National University, Research Institute of Mathematics, Global Analysis Research Center, Seoul, 1993.
- [8] J. Van Schaftingen, *A simple proof of an inequality of Bourgain, Brezis and Mironescu*, C. R. Math. Acad. Sci. Paris **338** (2004), no. 1, 23–26.
- [9] ———, *Estimates for L^1 -vector fields*, C. R. Math. Acad. Sci. Paris **339** (2004), no. 3, 181–186.
- [10] ———, *Limiting Bourgain-Brezis estimates for systems of linear differential equations: theme and variations*, J. Fixed Point Theory Appl. **15** (2014), no. 2, 273–297.

DEPARTMENT OF MATHEMATICS, RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854, USA

E-mail address: `chanillo@math.rutgers.edu`

INSTITUT DE RECHERCHE EN MATHÉMATIQUE ET EN PHYSIQUE, UNIVERSITÉ CATHOLIQUE DE LOUVAIN, CHEMIN DU CYCLOTRON 2 BTE L7.01.01, 1348 LOUVAIN-LA-NEUVE, BELGIUM

E-mail address: `Jean.VanSchaftingen@uclouvain.be`

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG

E-mail address: `plyung@math.cuhk.edu.hk`