Lower Bounds for Differential Privacy from Gaussian Width

Assimakis Kattis $^{\ast 1}$ and Aleksandar Nikolov $^{\dagger 2}$

^{1,2}Department of Computer Science, University of Toronto.

Abstract

We study the optimal sample complexity of a given workload of linear queries under the constraints of differential privacy. The sample complexity of a query answering mechanism under error parameter α is the smallest n such that the mechanism answers the workload with error at most α on any database of size n. Following a line of research started by Hardt and Talwar [STOC 2010], we analyze sample complexity using the tools of asymptotic convex geometry. We study the sensitivity polytope, a natural convex body associated with a query workload that quantifies how query answers can change between neighboring databases. This is the information that, roughly speaking, is protected by a differentially private algorithm, and, for this reason, we expect that a "bigger" sensitivity polytope implies larger sample complexity. Our results identify the mean Gaussian width as an appropriate measure of the size of the polytope, and show sample complexity lower bounds in terms of this quantity. Our lower bounds completely characterize the workloads for which the Gaussian noise mechanism is optimal up to constants as those having asymptotically maximal Gaussian width.

Our techniques also yield an alternative proof of Pisier's Volume Number Theorem which also suggests an approach to improving the parameters of the theorem.

1 Introduction

The main goal of private data analysis is to estimate aggregate statistics while preserving individual privacy guarantees. Intuitively, we expect that, for statistics that do not depend too strongly on any particular individual, a sufficiently large database allows computing an estimate that is both accurate and private. A natural question then is to characterize the *sample complexity* under privacy constraints: the smallest database size for which we can privately estimate the answers to a given collection of queries within some allowable error tolerance. Moreover, it is desirable to identify algorithms that are simple, efficient, and have close to the best possible sample complexity. In this work, we study these questions for collections of *linear queries* under the constraints of *approximate differential privacy*.

We model a *database* \mathcal{D} of *size* n as a multiset of n elements (counted with repetition) from an arbitrary finite universe \mathcal{U} . Each element of the database corresponds to the data of a single individual. To define a privacy-preserving computation on \mathcal{D} , we use the strong notion of *differential privacy*. Informally, an algorithm is differentially private if it has almost identical behavior on any two databases \mathcal{D} and \mathcal{D}' that differ in the data of a single individual. To capture this concept formally, let us define two databases to be *neighboring* if they have symmetric difference of size at most 1 (counted with multiplicity). Then differential privacy is defined as follows:

^{*}Email:kattis@cs.toronto.edu

[†]Email:anikolov@cs.toronto.edu

Definition 1 ([DMNS06]). A randomized algorithm \mathcal{A} that takes as input a database and outputs a random element from the set Y satisfies (ε, δ) -differential privacy if for all neighboring databases $\mathcal{D}, \mathcal{D}'$ and all measurable $S \subseteq Y$ we have that:

$$\mathbb{P}[\mathcal{A}(\mathcal{D}) \in S] \le e^{\varepsilon} \mathbb{P}[\mathcal{A}(\mathcal{D}') \in S] + \delta,$$

where probabilities are taken with respect to the randomness of \mathcal{A} .

One of the most basic primitives in private data analysis, and data analysis in general, are counting queries and, slightly more generally, linear queries. While interesting and natural in themselves, they are also quite powerful: any statistical query (SQ) learning algorithm can be implemented using noisy counting queries as a black box [Kea98]. In our setting, we specify a linear query by a function $q: \mathcal{U} \to [0, 1]$ (given by its truth table). Slightly abusing notation, we define the value of the query as $q(\mathcal{D}) = \frac{1}{n} \sum_{e \in \mathcal{D}} q(e)$, where the elements of \mathcal{D} are counted with multiplicity. For example, when $q: \mathcal{U} \to \{0, 1\}$, we can think of q as a property defined on \mathcal{U} and $q(\mathcal{D})$ as the fraction of elements of \mathcal{D} that satisfy the property: this is a counting query. We call a set \mathcal{Q} of linear queries a workload and an algorithm that answers a query workload a mechanism. We denote by $\mathcal{Q}(\mathcal{D}) = (q(\mathcal{D}))_{q \in \mathcal{Q}}$ the vector of answers to the queries in \mathcal{Q} . Throughout the paper, we will use the letter m for the size of a workload \mathcal{Q} .

Starting from the work of Dinur and Nissim [DN03], it is known that we cannot hope to answer too many linear queries too accurately while preserving even a very weak notion of privacy. For this reason, we must allow our private mechanisms to make some error. We focus on *average error* (in an L_2 sense). We define the average error of an algorithm \mathcal{A} on a query workload \mathcal{Q} and databases of size at most n as:

$$\operatorname{err}(\mathcal{Q}, \mathcal{A}, n) = \max_{\mathcal{D}} \left(\mathbb{E} \sum_{q \in \mathcal{Q}} \frac{(\mathcal{A}(\mathcal{D})_q - q(D))^2}{|\mathcal{Q}|} \right)^{1/2} = \max_{\mathcal{D}} \left(\mathbb{E} \frac{1}{m} \|\mathcal{A}(\mathcal{D}) - \mathcal{Q}(\mathcal{D})\|_2^2 \right)^{1/2},$$

where the maximum is over all databases \mathcal{D} of size at most n, $\mathcal{A}(\mathcal{D})_q$ is the answer to query q given by the algorithm \mathcal{A} on input \mathcal{D} , and expectations are taken with respect to the random choices of \mathcal{A} . This is a natural notion of error that also works particularly well with the geometric tools that we use.

In this work we study *sample complexity*: the smallest database size which allows us to answer a given query workload with error at most α . The sample complexity of an algorithm \mathcal{A} with error α is defined as:

$$\operatorname{sc}(\mathcal{Q}, \mathcal{A}, \alpha) = \min\{n : \operatorname{err}(\mathcal{Q}, \mathcal{A}, n) \le \alpha\}.$$

The sample complexity of answering the linear queries Q with error α under (ε, δ) -differential privacy is defined by:

$$\operatorname{sc}_{\varepsilon,\delta}(\mathcal{Q},\alpha) = \inf \{\operatorname{sc}(\mathcal{Q},\mathcal{A},\alpha) : \mathcal{A} \text{ is } (\epsilon,\delta) \text{-differentially private} \}.$$

The two main questions we are interested in are:

- 1. Can we characterize $\mathrm{sc}_{\varepsilon,\delta}(\mathcal{Q},\alpha)$ in terms of a natural property of the workload \mathcal{Q} ?
- 2. Can we identify conditions under which simple and efficient (ε, δ)-differentially private mechanisms have nearly optimal sample complexity?

We make progress on both questions. We identify a geometrically defined property of the workload that gives lower bounds on the sample complexity. The lower bounds also *characterize* when one of the simplest differentially private mechanisms, the Gaussian noise mechanism, has nearly optimal sample complexity in the regime of constant α .

Before we can state our results, we need to define a natural geometric object associated with a workload of linear queries. This object has been important in applying geometric techniques to differential privacy [HT10, BDKT12, NTZ13, Nik15].

Definition 2. The sensitivity polytope K of a workload Q of m linear queries is equal to $K = \text{conv}\{\pm Q(D) : D \text{ is a database of size } 1\}.$

From the above definition, we see that K is a symmetric (i.e. K = -K) convex polytope in \mathbb{R}^m . The importance of K lies in the fact that it captures how query answers can change between neighboring databases: for any two neighboring databases \mathcal{D} and \mathcal{D}' of size n and n' respectively, $n\mathcal{Q}(\mathcal{D}) - n'\mathcal{Q}(\mathcal{D}') \in K$. This is exactly the information that a differentially private algorithm is supposed to hide. Intuitively, we expect that the larger K is, the larger $\operatorname{sc}_{\varepsilon,\delta}(K, \alpha)$ should be.

We give evidence for the above intuition, and propose the width of K in a random direction as a measure of its "size". Let h_K be the support function of K: $h_K(y) = \max_{x \in K} \langle x, y \rangle$. For a unit vector y, $h_K(y) + h_K(-y)$ is the width of K in the direction of y; for arbitrary y, $h_K(ty)$ scales linearly with t (and is, in fact, a norm). We define the ℓ^* -norm of K, also known as its *Gaussian* mean width, as $\ell^*(K) = \mathbb{E}[h_K(g)]$, where g is a standard Gaussian random vector in \mathbb{R}^m . The following theorem captures our main result.

Theorem 1. Let \mathcal{Q} be a workload of m linear queries, and let K be its sensitivity polytope. The following holds for all $\varepsilon = O(1)$, $2^{-\Omega(n)} \leq \delta \leq 1/n^{1+\Omega(1)}$, and any $\alpha \leq \frac{\ell^*(K)}{Cm(\log 2m)^2}$, where C is an absolute constant, and $\sigma(\varepsilon, \delta) = (0.5\sqrt{\varepsilon} + \sqrt{2\log(1/\delta)})/\varepsilon$:

$$sc_{\varepsilon,\delta}(\mathcal{Q},\alpha) = O\left(\min\left\{\frac{\sigma(\varepsilon,\delta)\ell^*(K)}{\sqrt{m\alpha^2}}, \frac{\sigma(\varepsilon,\delta)\sqrt{m}}{\alpha}\right\}\right);$$

$$sc_{\varepsilon,\delta}(\mathcal{Q},\alpha) = \Omega\left(\frac{\sigma(\varepsilon,\delta)\ell^*(K)^2}{m^{3/2}(\log 2m)^4\alpha}\right).$$

The upper bound on sample complexity is achieved by a mechanism running in time polynomial in $m, n, and |\mathcal{U}|$. Moreover, if $\ell^*(K) = \Omega(m)$, then $\operatorname{sc}_{\varepsilon,\delta}(\mathcal{Q}, \alpha) = \Theta\left(\frac{\sigma(\varepsilon,\delta)\sqrt{m}}{\alpha}\right)$ for any $\alpha \leq 1/C$, where C is an absolute constant.

The sample complexity upper bounds in the theorem above are known from prior work: one is given by the projection mechanism from [NTZ13], with the sample complexity upper bound in terms of $\ell^*(K)$ shown in [DNT14]; the other upper bound is given by the Gaussian noise mechanism [DN03, DN04, DMNS06]. The main new contribution in this work are the lower bounds on sample complexity. The gap between upper and lower bounds is small when $\ell^*(K)$ is close to its maximal value of m. Indeed, when $\ell^*(K) = \Theta(m)$, our results imply that the Gaussian noise mechanism has optimal sample complexity up to constants. This is, to the best of our knowledge, the first example of a general geometric condition under which a simple and efficient mechanism has optimal sample complexity up to *constant* factors. Moreover, in the constant error regime this condition is also *necessary* for the Gaussian mechanism to be optimal up to constants: when $\ell^*(K) = o(m)$ and $\alpha = \Omega(1)$, the projection mechanism has asymptotically smaller sample complexity than the Gaussian mechanism.

We can prove somewhat stronger results for another natural problem in private data analysis, which we call the *mean point problem*. In this problem, we are given a closed convex set $K \subset \mathbb{R}^m$,

and we are asked to approximate the mean $\overline{\mathcal{D}}$ of the database \mathcal{D} , where $\mathcal{D} = \{x_1, \ldots, x_n\}$ is a multiset of points in K and $\overline{\mathcal{D}} = \frac{1}{n} \sum_{i=1}^{n} x_i$. This problem, which will be the focus for most of this paper, has a more geometric flavor, and is closely related to the query release problem for linear queries. In fact, Theorem 1 will essentially follow from a reduction from the results below for the mean point problem.

With respect to the mean point problem, we define the error of an algorithm \mathcal{A} as:

$$\operatorname{err}(K, \mathcal{A}, n) = \sup_{\mathcal{D}} (\mathbb{E} \| \mathcal{A}(\mathcal{D}) - \overline{\mathcal{D}} \|_2^2)^{1/2}$$

where the supremum is over databases \mathcal{D} consisting of at most n points from K, and the expectation is over the randomness of the algorithm. The sample complexity of an algorithm \mathcal{A} with error α is defined as:

$$\operatorname{sc}(K, \mathcal{A}, \alpha) = \min\{n : \operatorname{err}(K, \mathcal{A}, n) \le \alpha\}.$$

The sample complexity of solving the mean point problem with error α over K is defined by:

$$\operatorname{sc}_{\varepsilon,\delta}(K,\alpha) = \min\{\operatorname{sc}(K,\mathcal{A},\alpha) : \mathcal{A} \text{ is } (\varepsilon,\delta) \text{-differentially private}\}$$

Our main result for the mean point problem is given in the following theorem:

Theorem 2. Let K be a symmetric convex body contained in the unit Euclidean ball B_2^m in \mathbb{R}^m . The following holds for all $\varepsilon = O(1)$, $2^{-\Omega(n)} \leq \delta \leq 1/n^{1+\Omega(1)}$, and any $\alpha \leq \frac{\ell^*(K)}{C\sqrt{m}(\log 2m)^2}$, where C is an absolute constant, and $\sigma(\varepsilon, \delta) = (0.5\sqrt{\varepsilon} + \sqrt{2\log(1/\delta)})/\varepsilon$:

$$sc_{\varepsilon,\delta}(K,\alpha) = O\left(\min\left\{\frac{\sigma(\varepsilon,\delta)\ell^*(K)}{\alpha^2}, \frac{\sigma(\varepsilon,\delta)\sqrt{m}}{\alpha}\right\}\right);$$

$$sc_{\varepsilon,\delta}(K,\alpha) = \Omega\left(\frac{\sigma(\varepsilon,\delta)\ell^*(K)}{(\log 2m)^2\alpha}\right).$$

The upper bound on sample complexity is achieved by a mechanism running in time polynomial in $m, n, and |\mathcal{U}|$. Moreover, when $\ell^*(K) = \Omega(\sqrt{m})$, then $\operatorname{sc}_{\varepsilon,\delta}(\mathcal{Q}, \alpha) = \Theta\left(\frac{\sigma(\varepsilon,\delta)\sqrt{m}}{\alpha}\right)$ for any $\alpha \leq 1/C$, where C is an absolute constant.

The upper bounds again follow from prior work, and in fact are also given by the projection mechanism and the Gaussian noise mechanism, which can be defined for the mean point problem as well. Notice that the gap between the upper and the lower bound is on the order $\frac{(\log 2m)^2}{\alpha}$. If the lower bound was valid for all values of the error parameter α less than a fixed constant, rather than for $\alpha \leq \frac{\ell^*(K)}{C\sqrt{m}(\log 2m)^2}$, Theorem 2 would nearly characterize the optimal sample complexity for the mean point problem for all constant α . Unfortunately, the restriction on α is, in general, necessary (up to the logarithmic terms) for lower bounds on sample complexity in terms of $\ell^*(K)$. For example, we can take $K = \gamma B_2^m$, i.e. a Euclidean ball in \mathbb{R}^m with radius γ . Then, $\ell^*(K) = \Theta(\gamma\sqrt{m})$, but the sample complexity is 0 when $\alpha > \gamma$, since the trivial algorithm which ignores the database and outputs 0 achieves error γ . Thus, a more sensitive measure of the size of K is necessary to prove optimal lower bounds. We do, nevertheless, trust that the techniques introduced in this paper bring us closer to this goal.

We conclude this section with a high-level overview of our techniques. Our starting point is a recent tight lower bound on the sample complexity of a special class of linear queries: the 1-way marginal queries. These queries achieve the worst case sample complexity for a family of m linear queries: $\Omega(\sqrt{m}/\alpha)$ [BUV14, SU15]. The sensitivity polytope of the 1-way marginals is the cube

 $[-1, 1]^m$, and it can be shown that the lower bound on the sample complexity of 1-way marginals implies an analogous lower bound on the sample complexity of the mean point problem with $K = Q^m = [-1/\sqrt{m}, 1/\sqrt{m}]^m$. For the mean point problem, it is easy to see that when $K' \subseteq K$, the sample complexity for K' is no larger than the sample complexity for K. Moreover, we can show that the sample complexity of any projection of K is no bigger than the sample complexity of K itself. So, our strategy then is to find a large scaled copy of $Q^{m'}$, $m' \leq m$, inside a projection of K onto a large dimensional subspace whenever $\ell^*(K)$ is large. We solve this geometric problem using deep results from asymptotic convex geometry, namely the Dvoretzky criterion, the low M^* estimate, and the MM^* estimate.

Our techniques also yield an alternative proof of the volume number theorem of Milman and Pisier [MP87]. Besides avoiding the quotient of subspace theorem, our proof yields an improvement in the volume number theorem, conditional on the well-known conjecture that any symmetric convex body K has a position (affine image) TK for which $\ell^*(TK)\ell(TK) = O(m\sqrt{\log 2m})$, where $\ell(K)$ is the expected K-norm of a standard Gaussian. More details about this connection are given in Section 7.

1.1 Prior Work

Most closely related to our work are the results of Nikolov, Talwar, and Zhang [NTZ13], who gave a private mechanism (also based on the projection mechanism, but more involved) which has nearly optimal sample complexity (with respect to average error), up to factors polynomial in $\log m$ and $\log |\mathcal{U}|$. This result was subsequently improved by Nikolov [Nik15], who showed that the $\log m$ factors can be replaced by $\log n$. While these results are nearly optimal for subconstant values of the error parameter α , i.e. the optimality guarantees do not depend on $1/\alpha$, factors polynomial in $\log |\mathcal{U}|$ can be prohibitively large. Indeed, in many natural settings, such as that of marginal queries, $|\mathcal{U}|$ is exponential in the number of queries m, so the competitiveness ratio can be polynomial in m.

The line of work that applies techniques from convex geometry to differential privacy started with the beautiful paper of Hardt and Talwar [HT10], whose results were subsequently strengthened in [BDKT12]. These papers focused on the "large database" regime (or, in our language, the setting of subconstant error), and pure differential privacy ($\delta = 0$).

2 Preliminaries

We begin with the introduction of some notation. Throughout the paper we use C, C_1 , etc., for absolute constants, whose value may change from line to line. We use $\|\cdot\|_2$ for the Euclidean norm, and $\|\cdot\|_1$ for the ℓ_1 norm. We define B_1^m and B_2^m to be the ℓ_1 and ℓ_2 unit balls in \mathbb{R}^m respectively, while $Q^m = [-\frac{1}{\sqrt{m}}, \frac{1}{\sqrt{m}}]^m \subseteq \mathbb{R}^m$ will refer to the *m*-dimensional hypercube, normalized to be contained in the unit Euclidean ball. We use I_m for the identity operator on \mathbb{R}^m , as well as for the $m \times m$ identity matrix. For a given subspace E, we define $\Pi_E \colon \mathbb{R}^m \to \mathbb{R}^m$ as the orthogonal projection operator onto E. Moreover, when $T \colon E \to F$ is a linear operator between the subspaces $E, F \subseteq \mathbb{R}^m$, we define $\|T\| = \max\{\|Tx\|_2 \colon \|x\|_2 = 1\}$ as its operator norm, which is also equal to its largest singular value $\sigma_1(T)$. For the diameter of a set K we use the nonstandard, but convenient, definition diam $K = \max\{\|x\|_2 : x \in K\}$. For sets symmetric around 0, this is equivalent to the standard definition, but scaled up by a factor of 2. We use $N(\mu, \Sigma)$ to refer to the Gaussian distribution with mean μ and covariance Σ , and we use the notation $x \sim N(\mu, \Sigma)$ to denote that xis distributed as a Gaussian random variable with mean μ and covariance Σ . For a $m \times m$ matrix (or equivalently an operator from ℓ_2^m to ℓ_2^m) A we use $A \succeq 0$ to denote that A is positive semidefinite. For positive semidefinite matrices/operators A, B, we use the notation $A \preceq B$ to denote $B - A \succeq 0$.

3 Probability Theory

We make use of some basic comparison theorems from the theory of stochastic processes. First we state the well-known symmetrization lemma. We also give the short proof for completeness. Recall that ξ_1, \ldots, ξ_n are a sequence of *Rademacher random variables* if each ξ_i is uniformly and independently distributed in $\{-1, 1\}$.

Lemma 1 (Symmetrization). Let p > 1, and let $\|\cdot\|$ be a norm on \mathbb{R}^m . Then, for any sequence x_1, \ldots, x_n of independent random variables in \mathbb{R}^m such that $\mathbb{E}\|x_i\|^p$ is finite for every *i*, we have

$$\mathbb{E} \left\| \sum_{i} x_{i} - \mathbb{E} \sum_{i} x_{i} \right\|^{p} \leq 2^{p} \mathbb{E} \left\| \sum_{i} \xi_{i} x_{i} \right\|^{p},$$

where ξ_1, \ldots, ξ_n are Rademacher random variables, independent of x_1, \ldots, x_n . Each expectation above is with respect to all random variables involved.

Proof. Let x'_1, \ldots, x'_n be independent copies of x_1, \ldots, x_n . Then, $\mathbb{E} \sum_i x_i = \mathbb{E} \sum_i x'_i$, and, by convexity of the function $\|\cdot\|^p$ and Jensen's inequality,

$$\mathbb{E} \left\| \sum_{i} x_{i} - \mathbb{E} \sum_{i} x_{i} \right\|^{p} = \mathbb{E} \left\| \sum_{i} x_{i} - \mathbb{E} \sum_{i} x_{i}' \right\|^{p} \leq \mathbb{E} \left\| \sum_{i} x_{i} - x_{i}' \right\|^{p}.$$

Because x_i and x'_i are independent and identically distributed, the random variables $x_i - x'_i$ and $x'_i - x_i$ are also identically distributed. Therefore, $\mathbb{E} \| \sum_i x_i - x'_i \|^p = \mathbb{E} \| \sum_i \xi_i (x_i - x'_i) \|^p$, where ξ_1, \ldots, ξ_n are independent Rademacher random variables, as in the statement of the lemma. Finally, by Minkowski's inequality (i.e. triangle inequality for L_p), we have

$$\mathbb{E}\left\|\sum_{i}\xi_{i}(x_{i}-x_{i}')\right\|^{p} \leq \left(\left(\mathbb{E}\left\|\sum_{i}\xi_{i}x_{i}\right\|^{p}\right)^{1/p} + \left(\mathbb{E}\left\|\sum_{i}\xi_{i}x_{i}'\right\|^{p}\right)^{1/p}\right)^{p} = 2^{p}\mathbb{E}\left\|\sum_{i}\xi_{i}x_{i}\right\|^{p},$$

as desired.

Next we state a simple comparison theorem for Gaussian random variables.

Lemma 2. Let $x \sim N(0, \Sigma)$ and $x' \sim N(0, \Sigma')$ be Gaussian random variables in \mathbb{R}^m , and assume $\Sigma \leq \Sigma'$. Then, for any norm $\|\cdot\|$ on \mathbb{R}^m , we have:

$$\mathbb{E}\|x\| \le \mathbb{E}\|x'\|.$$

Proof. Couple x and x' so that they are independent, and define a new random variable $y \sim N(0, \Sigma' - \Sigma)$, independent of x and x'. Then the random variables x + y and x - y are distributed identically to x', and, by linearity of expectation and the triangle inequality we have

$$\mathbb{E}\|x'\| = \mathbb{E}\left[\frac{\|x+y\| + \|x-y\|}{2}\right] \ge \mathbb{E}\|x\|.$$

This completes the proof.

Note that the same conclusion follows under weaker assumptions from Slepian's lemma.

3.1 Convex Geometry

In this section, we outline the main geometric tools we use in later sections. For a more detailed treatment, we refer to the lecture notes by Vershynin [Ver09] and the books by Pisier [Pis89] and Artstein-Avidan, Giannopoulos, and Milman [AAGM15].

Throughout, we define a *convex body* K as a compact subset of \mathbb{R}^m with non-empty interior. A convex body K is *(centrally) symmetric* if and only if K = -K. We define the *polar body* K° of K as: $K^\circ = \{y : \langle x, y \rangle \leq 1 \ \forall x \in K\}$. The following basic facts are easy to verify and very useful.

Fact 1. For convex bodies $K, L \subseteq \mathbb{R}^m, K \subseteq L \Leftrightarrow L^{\circ} \subseteq K^{\circ}$.

Fact 2 (Section/Projection Duality). For a convex body $K \subseteq \mathbb{R}^m$ and a subspace $E \subseteq \mathbb{R}^m$:

- 1. $(K \cap E)^{\circ} = \prod_{E} (K^{\circ});$
- 2. $(\Pi_E(K))^\circ = K^\circ \cap E.$

In both cases, the polar is taken in the subspace E.

Fact 3. For any invertible linear map T and any convex body K, $T(K)^{\circ} = T^{-*}(K^{\circ})$, where T^{-*} is the inverse of the adjoint operator T^* .

A simple special case of Fact 3 is that, for any convex body K, $(rK)^{\circ} = \frac{1}{r}K^{\circ}$. Using this property alongside Fact 1, we have the following useful corollary.

Corollary 1. For a convex body $K \subseteq \mathbb{R}^m$ and $E \subseteq \mathbb{R}^m$ a subspace with $k = \dim E$, the following two statements are equivalent:

- 1. $\Pi_E(rB_2^m) \subseteq \Pi_E(K);$
- 2. $K^{\circ} \cap E \subseteq \frac{1}{r}(B_2^m \cap E),$

where, as before, taking the polar set is considered in the subspace E. Notice that the second statement is also equivalent to diam $(K^{\circ} \cap E) \leq \frac{1}{r}$.

Our work relies on appropriately quantifying the "size" of (projections and sections of) a convex body. It turns out that, for our purposes, the right measure of size is related to the notion of width, captured by the support function. Recall from the introduction that the support function of a convex body $K \subset \mathbb{R}^m$ is given by $h_K(y) = \max_{x \in K} \langle x, y \rangle$ for every $y \in \mathbb{R}^m$. The support function is intimately related to the Minkowski norm $\|\cdot\|_K$, defined for a symmetric convex body $K \subseteq \mathbb{R}^m$ by $\|x\|_K = \min\{r \in \mathbb{R} : x \in rK\}$, for every $x \in \mathbb{R}^m$. It is easy to verify that $\|\cdot\|_K$ is indeed a norm. The support function h_K is identical to the Minkowski norm of the polar body K° (which is also the dual norm to $\|\cdot\|_K$): $h_K(y) = \|y\|_{K^\circ}$ for every $y \in \mathbb{R}^m$.

Now we come to the measure of the "size" of a convex body which will be central to our results: the Gaussian mean width of the body, defined next.

Definition 3. The Gaussian mean width and Gaussian mean norm of a symmetric convex body $K \subseteq \mathbb{R}^m$ are defined respectively as:

$$\ell^*(K) = \mathbb{E} \|g\|_{K^\circ} = \mathbb{E}[h_K(g)], \qquad \qquad \ell(K) = \mathbb{E} \|g\|_K,$$

where $g \sim N(0, I_m)$ is a standard Gaussian random variable.

The next lemma gives an estimate of how the mean width changes when applying a linear transformation to K.

Lemma 3. For any symmetric convex body $K \subset \mathbb{R}^m$, and any linear operator $T: \ell_2^m \to \ell_2^m$:

$$\ell^*(T(K)) \le ||T|| \ell^*(K).$$

Proof. Notice that, for a standard Gaussian $g \sim N(0, I_m)$,

$$\ell^*(T(K)) = \mathbb{E} \sup_{x \in K} \langle T(x), g \rangle = \mathbb{E} \sup_{x \in K} \langle x, T^*(g) \rangle = \mathbb{E} \| T^*(g) \|_{K^\circ}.$$

Treating T^*T as an $m \times m$ matrix in the natural way, we see that $T^*(g) \sim N(0, T^*T)$. By applying Lemma 2 to $\Sigma = T^*T$ and $\Sigma' = ||T||^2 I_m$, we have that

$$\ell^*(T(K)) = \mathbb{E} \|T^*(g)\|_{K^\circ} \le \|T\| \cdot \mathbb{E} \|g\|_{K^\circ} = \|T\|\ell^*(K).$$

This finishes the proof of the lemma.

Similar to approaches in previous works ([HT10], [NTZ13]), we exploit properties inherent to a specific position of K to prove lower bounds on its sample complexity.

Definition 4 (ℓ -position). A convex body $K \subseteq \mathbb{R}^m$ is in ℓ -position if for all linear operators $T: \ell_2^m \to \ell_2^m$:

$$\ell^*(K) \cdot \ell(K) \le \ell^*(T(K)) \cdot \ell(T(K)).$$

Clearly, K is in ℓ -position if and only if K° is in ℓ -position, since $\ell^*(K) = \ell(K^{\circ})$ for any convex body K. Note further that the product $\ell^*(K) \cdot \ell(K)$ is scale-invariant, in the sense that $\ell^*(rK) \cdot \ell(rK) = \ell^*(K) \cdot \ell(K)$ for any real r. This is because, for any $x, y \in \mathbb{R}^m$, $||x||_{rK} = \frac{1}{r} ||x||_K$, and $h_{rK}(y) = rh_K(y)$, so $\ell^*(rK) = r\ell^*(K)$ and $\ell(rK) = \frac{1}{r}\ell(K)$.

We will relate the Gaussian mean width of K to another measure of its size, and the size of its projections and sections, known as Gelfand width. A definition follows.

Definition 5 (Gelfand width). For two symmetric convex bodies $K, L \subset \mathbb{R}^m$, the Gelfand width of order k of K with respect to L is defined as:

$$c_k(K,L) = \inf_E \inf\{r : K \cap E \subseteq r(L \cap E)\} = \inf_E \sup\{\|x\|_L : x \in K \cap E\},\$$

where the first infimum is over subspaces $E \subseteq \mathbb{R}^m$ of co-dimension at most k-1 (i.e. of dimension at least m-k+1). When k > m, we define $c_k(K,L) = 0$. We denote $c_k(K) = c_k(K, B_2^m)$, and we call $c_k(K)$ simply the Gelfand width of K of order k.

Note that $c_k(K) = \inf_E \operatorname{diam}(K \cap E)$, where the infimum is over subspaces $E \subseteq \mathbb{R}^m$ of codimension at most k-1. Observe also that for any K and L, $c_k(K, L)$ is non-increasing in k. It is well-known that the infimum in the definition is actually achieved [Pin85].

3.2 Composition of Differential Privacy

One of the most important properties of differential privacy is that it behaves nicely under (adaptively) composing mechanisms.

Lemma 4 (Composition). For randomized algorithms \mathcal{A}_1 and \mathcal{A}_2 satisfying $(\varepsilon_1, \delta_1)$ - and $(\varepsilon_2, \delta_2)$ differential privacy respectively, the algorithm $\mathcal{A}(\mathcal{D}) = (\mathcal{A}_1(\mathcal{D}), \mathcal{A}_2(\mathcal{A}_1(\mathcal{D}), \mathcal{D}))$ satisfies $(\varepsilon_1 + \varepsilon_2, \delta_1 + \delta_2)$ -differential privacy.

3.3 Known Bounds

In this section, we recall some known differentially private mechanisms, with bounds on their sample complexity, as well as a lower bound on the optimal sample complexity. We start with the lower bound:

Theorem 3 ([BUV14, SU15]). For all $\varepsilon = O(1)$, $2^{-\Omega(n)} \leq \delta \leq 1/n^{1+\Omega(1)}$ and $\alpha \leq 1/10$:

$$\operatorname{sc}_{\varepsilon,\delta}\left(Q^m,\alpha\right) = \Omega\left(\frac{\sqrt{m\log 1/\delta}}{\alpha\varepsilon}\right).$$
 (1)

Next we recall one of the most basic mechanisms in differential privacy, the Gaussian mechanism. A proof of the privacy guarantee, with the constants given below, can be found in [NTZ13].

Theorem 4 (Gaussian Mechanism [DN03, DN04, DMNS06]). Let $\mathcal{D} = \{x_1, \ldots, x_n\}$ be such that $\forall i : \|x_i\|_2 \leq \sigma$. If $w \sim N(0, \sigma(\varepsilon, \delta)^2 \sigma^2 I_m)$, $\sigma(\varepsilon, \delta) = (0.5\sqrt{\varepsilon} + \sqrt{2\log(1/\delta)})/\varepsilon$ and $I_m \in \mathbb{R}^{m \times m}$ is the identity matrix, then the algorithm \mathcal{A}_{GM} defined by $\mathcal{A}_{GM}(\mathcal{D}) = \overline{\mathcal{D}} + \frac{1}{n}w$ is (ε, δ) -differentially private.

Corollary 2. For any symmetric convex $K \subseteq B_2^m$:

$$\operatorname{sc}_{\varepsilon,\delta}(K,\alpha) = O\left(\frac{\sqrt{m\log 1/\delta}}{\alpha\varepsilon}\right)$$

In the rest of the paper we will use the notation $\sigma(\varepsilon, \delta) = \frac{0.5\sqrt{\varepsilon} + \sqrt{2\log(1/\delta)}}{\varepsilon}$ from the theorem statement above.

Finally, we also present the projection mechanism from [NTZ13], which post-processes the output of the Gaussian mechanism by projecting onto K.

Theorem 5 (Projection Mechanism [NTZ13, DNT14]). Let $K \subseteq B_2^m$ be a symmetric convex body, and define \mathcal{A}_{PM} to be the algorithm that, on input $\mathcal{D} = \{x_1, \ldots, x_n\} \subset K$, outputs:

$$\hat{y} = \arg\min\{\|\hat{y} - \tilde{y}\|_2^2 : \hat{y} \in K\},\$$

where $\tilde{y} = \overline{D} + \frac{1}{n}w$, $w \sim N(0, \sigma(\varepsilon, \delta)^2 I_m)$. Then \mathcal{A}_{PM} satisfies (ε, δ) -differential privacy and has sample complexity:

$$\operatorname{sc}(K, \mathcal{A}_{PM}, \alpha) = O\left(\frac{\sigma(\varepsilon, \delta)\ell^*(K)}{\alpha^2}\right).$$

Corollary 3. For any symmetric convex $K \subseteq B_2^m$:

$$\operatorname{sc}_{\varepsilon,\delta}(K,\alpha) = O\left(\frac{\sigma(\varepsilon,\delta)\ell^*(K)}{\alpha^2}\right).$$

4 Basic Properties of Sample Complexity

In this section, we prove some fundamental properties of sample complexity that will be extensively used in later sections.

Lemma 5. $L \subseteq K \Rightarrow \forall \alpha \in (0,1) : \operatorname{sc}_{\varepsilon,\delta}(L,\alpha) \leq \operatorname{sc}_{\varepsilon,\delta}(K,\alpha).$

Proof. Observe first that for any algorithm \mathcal{A} and any n, $\operatorname{err}(L, \mathcal{A}, n) \leq \operatorname{err}(K, \mathcal{A}, n)$, because $\operatorname{err}(K, \mathcal{A}, n)$ is a supremum over a larger set than $\operatorname{err}(L, \mathcal{A}, n)$. This implies that $\operatorname{sc}(K, \mathcal{A}, \alpha) \leq \operatorname{sc}(L, \mathcal{A}, \alpha)$ holds for any algorithm \mathcal{A} , and, in particular, for the (ε, δ) -differentially private algorithm \mathcal{A}^* that achieves $\operatorname{sc}_{\varepsilon,\delta}(K, \alpha)$. Then, we have:

$$\operatorname{sc}_{\varepsilon,\delta}(L,\alpha) \leq \operatorname{sc}(L,\mathcal{A}^*,\alpha) \leq \operatorname{sc}(K,\mathcal{A}^*,\alpha) = \operatorname{sc}_{\varepsilon,\delta}(K,\alpha),$$

as desired.

Corollary 4. For all $\varepsilon = O(1)$, $2^{-\Omega(n)} \le \delta \le 1/n^{1+\Omega(1)}$ and $\alpha \le 1/10$:

$$\operatorname{sc}_{\varepsilon,\delta}(B_2^m, \alpha) = \Omega\left(\frac{\sqrt{m\log 1/\delta}}{\alpha\varepsilon}\right).$$

Proof. Since $Q^m \subseteq B_2^m$, this follows directly from Lemma 5 and Lemma 3.

Lemma 6. For any $\alpha \in (0,1)$, any linear operator $T \colon \mathbb{R}^m \to \mathbb{R}^m$ and any symmetric convex body $K \subset \mathbb{R}^m$:

$$\operatorname{sc}_{\varepsilon,\delta}(K,\alpha) \ge \operatorname{sc}_{\varepsilon,\delta}(T(K),\alpha \cdot ||T||).$$

Proof. Let \mathcal{A} be an (ε, δ) -differentially private algorithm that achieves $\mathrm{sc}_{\varepsilon,\delta}(K, \alpha)$. Fix a function $f: T(K) \to K$ so that for every $x' \in T(K)$, $f(x') \in T^{-1}(x') \cap K$. We define a new algorithm \mathcal{A}' that takes as input $\mathcal{D}' = \{x'_1, ..., x'_n\} \subset T(K)$ and outputs $\mathcal{A}'(\mathcal{D}') = T(\mathcal{A}(\mathcal{D}))$, where $\mathcal{D} = f(\mathcal{D}') = \{f(x'_1), ..., f(x'_n)\}$. We claim that \mathcal{A}' is (ε, δ) -differentially private and that $\mathrm{err}(T(K), \mathcal{A}', n) \leq \|T\| \cdot \mathrm{err}(K, \mathcal{A}, n)$ holds for every n. This claim is sufficient to prove the lemma, because it implies:

 $\mathrm{sc}_{\varepsilon,\delta}(T(K),\alpha \cdot \|T\|) \leq \mathrm{sc}(T(K),\mathcal{A}',\alpha \cdot \|T\|) \leq \mathrm{sc}(K,\mathcal{A},\alpha) = \mathrm{sc}_{\varepsilon,\delta}(K,\alpha).$

To show the claim, first observe that, by linearity, $\overline{\mathcal{D}'} = T(\overline{\mathcal{D}})$. We get:

$$\mathbb{E} \|\mathcal{A}'(\mathcal{D}') - \overline{\mathcal{D}'}\|_{2}^{2} = \mathbb{E} \|T(\mathcal{A}(\mathcal{D})) - T(\overline{\mathcal{D}})\|_{2}^{2}$$

$$= \mathbb{E} \|T(\mathcal{A}(\mathcal{D}) - \overline{\mathcal{D}})\|_{2}^{2}$$

$$\leq \|T\|^{2} \cdot \mathbb{E} \|\mathcal{A}(\mathcal{D}) - \overline{\mathcal{D}}\|_{2}^{2} \leq \|T\|^{2} \operatorname{err}(K, \mathcal{A}, n)^{2},$$

where the first inequality follows by the definition of the operator norm. Since this holds for arbitrary n and $\mathcal{D}' \subset T(K)$ of size n, it implies the claim on the error bound of \mathcal{A}' . It remains to show that \mathcal{A}' is (ε, δ) -differentially private. Note that for every two neighboring databases \mathcal{D}'_1 and \mathcal{D}'_2 of points in T(K), the corresponding databases $\mathcal{D}_1 = f(\mathcal{D}'_1)$ and $\mathcal{D}_2 = f(\mathcal{D}'_2)$ of points in K are also neighboring. Then $\mathcal{A}(\mathcal{D}) = \mathcal{A}(f(\mathcal{D}'))$ is (ε, δ) -differentially private as a function of \mathcal{D}' , and the privacy of \mathcal{A}' follows from Lemma 4.

Corollary 5. For any t > 0:

$$\operatorname{sc}_{\varepsilon,\delta}(tK, t\alpha) = \operatorname{sc}_{\varepsilon,\delta}(K, \alpha).$$

Proof. Taking $T = tI_m$ in Lemma 6, where I_m is the identity on \mathbb{R}^m , the lemma implies $\mathrm{sc}_{\varepsilon,\delta}(tK, t\alpha) \leq \mathrm{sc}_{\varepsilon,\delta}(K, \alpha)$. Since this inequality holds for any t and K, we may apply it to K' = tK and t' = 1/t, and we get $\mathrm{sc}_{\varepsilon,\delta}(K, \alpha) = \mathrm{sc}_{\varepsilon,\delta}((1/t)tK, (1/t)t\alpha) \leq \mathrm{sc}_{\varepsilon,\delta}(tK, t\alpha)$.

Since for any subspace E of \mathbb{R}^m , the corresponding orthogonal projection Π_E has operator norm 1, we also immediately get the following corollary of Lemma 6:

Corollary 6. For any subspace E:

$$\operatorname{sc}_{\varepsilon,\delta}(K,\alpha) \ge \operatorname{sc}_{\varepsilon,\delta}(\Pi_E(K),\alpha)$$

In the next theorem, we combine the lower bound in Corollary 4 and the properties we proved above in order to give a lower bound on the sample complexity of an arbitrary symmetric convex body K in terms of its geometric properties. In the following sections we will relate this geometric lower bound to the mean Gaussian width of K.

Theorem 6 (Geometric Lower Bound). For all $\varepsilon = O(1)$, $2^{-\Omega(n)} \leq \delta \leq 1/n^{1+\Omega(1)}$, any convex symmetric body $K \subseteq \mathbb{R}^m$, any $1 \leq k \leq m$ and any $\alpha \leq 1/(10c_k(K^\circ))$:

$$\operatorname{sc}_{\varepsilon,\delta}(K,\alpha) = \Omega\left(\frac{\sqrt{\log 1/\delta}}{\alpha\varepsilon} \cdot \frac{\sqrt{m-k+1}}{c_k(K^\circ)}\right)$$

Proof. Let us fix k and let E be the subspace that achieves $c_k(K^\circ)$, i.e. $\operatorname{diam}(\Pi_E(K^\circ)) = c_k(K^\circ)$ and $d_E = \dim E \ge m - k + 1$. By Corollary 1, we have $c_k(K^\circ)^{-1}\Pi_E(B_2^m) \subseteq \Pi_E(K)$. Applying Corollary 6, Lemma 5, and Corollary 5 in sequence, we get:

$$sc_{\varepsilon,\delta}(K,\alpha) \ge sc_{\varepsilon,\delta}(\Pi_E(K),\alpha) \ge sc_{\varepsilon,\delta}(c_k(K^\circ)^{-1}\Pi_E(B_2^m),\alpha)$$
$$= sc_{\varepsilon,\delta}(\Pi(B_2^m),\alpha c_k(K^\circ)).$$

Notice that $\Pi_E(B_2^m)$ is the Euclidean unit ball in the subspace E, and, therefore:

$$\operatorname{sc}_{\varepsilon,\delta}(\Pi(B_2^m), \alpha c_k(K^\circ)) = \operatorname{sc}_{\varepsilon,\delta}(B_2^{d_E}, \alpha c_k(K^\circ)).$$

Finally, by Corollary 4, we get the following lower bound, as long as $\alpha c_k(K^{\circ}) \leq 1/10$:

$$\mathrm{sc}_{\varepsilon,\delta}(B_2^{d_E},\alpha c_k(K^\circ)) = \Omega\left(\frac{\sqrt{d_E \log 1/\delta}}{\alpha c_k(K^\circ)\varepsilon}\right) = \Omega\left(\frac{\sqrt{\log 1/\delta}}{\alpha\varepsilon} \cdot \frac{\sqrt{m-k+1}}{c_k(K^\circ)}\right).$$

Combining the inequalities completes the proof.

5 Optimality of the Gaussian Mechanism

In this section, we present the result that the Gaussian mechanism is optimal, up to constant factors, when $K \subseteq B_2^m$ is sufficiently large. More specifically, if the Gaussian mean width of K is asymptotically maximal, then we can get a tight lower bound on the sample complexity of the Gaussian mechanism. This is summarized in the theorem below.

Theorem 7. For all $\varepsilon < O(1)$, $2^{-\Omega(n)} \le \delta \le 1/n^{1+\Omega(1)}$, sufficiently small constant α , and any symmetric convex body $K \subseteq B_2^m$, if

$$\ell^*(K) = \Omega(\sqrt{m}),$$

then:

$$\operatorname{sc}_{\varepsilon,\delta}(K,\alpha) = \Theta\left(\frac{\sqrt{m\log 1/\delta}}{\alpha\varepsilon}\right),$$

and $\operatorname{sc}_{\varepsilon,\delta}(K,\alpha)$ is achieved, up to constants, by the Gaussian mechanism.

By Corollary 2 we have an upper bound for the Gaussian mechanism defined previously. To prove its optimality, we use a classical result from convex geometry, known as Dvoretzky's criterion, to show a matching lower bound for the sample complexity. This result relates the existence of a nearly-spherical section of a given convex body to the Gaussian mean norm. It was a key ingredient in Milman's probabilistic proof of Dvoretzky's theorem: see Matoušek's book [Mat02] for an exposition.

Theorem 8 ([Mil71]; Dvoretzky's Criterion). For every symmetric convex body $K \subseteq \mathbb{R}^m$ such that $B_2^m \subseteq K$, and every $\beta < 1$, there exists a constant $c(\beta)$ and a subspace E with dimension $\dim E \ge c(\beta)\ell(K)^2$ for which:

$$(1-\beta)\frac{\ell(K)}{\sqrt{m}}B_2^m \cap E \subseteq K \cap E \subseteq (1+\beta)\frac{\ell^*(K)}{\sqrt{m}}B_2^m \cap E.$$

Proof of Theorem 7. Given the matching upper bound on sample complexity in Corollary 2, it suffices to show the equivalent lower bound, namely that:

$$\operatorname{sc}_{\varepsilon,\delta}(K,\alpha) = \Omega\left(\frac{\sqrt{m\log 1/\delta}}{\alpha\varepsilon}\right).$$

To this end, we will show that there exists a $k \leq (1-c)m+1$, for an absolute constant c, so that $c_k(K^\circ) = O(1)$. Then the lower bound will follow directly from Theorem 6.

We will prove the claim above by applying Dvoretzky's criterion to K° . By Fact 1, $K \subseteq B_2^m \Rightarrow B_2^m \subseteq K^{\circ}$. We can then apply Dvoretzky's criterion with $\beta = 1/2$, ensuring that there exists a subspace E of dimension dim $E \ge c(1/2)\ell(K^{\circ})^2$ for which:

$$K^{\circ} \cap E \subseteq \frac{\ell(K^{\circ})}{2\sqrt{m}} B_2^m \cap E.$$

Let us define $k = m - \dim E + 1$; then $k \le m - c(1/2)\ell(K^{\circ})^2 + 1 = m - c(1/2)\ell^*(K)^2 + 1$. Since, by assumption $\ell(K^*) = \Omega(m)$, there exists a constant c so that $k \le (1 - c)m + 1$. Finally, by the definition of Gelfand width, $c_k(K^{\circ}) \le \frac{\ell(K^{\circ})}{2\sqrt{m}} = O(1)$, as desired. This completes the proof.

6 Gaussian Width Lower Bounds in ℓ -position

In Section 5 we showed that the Gaussian Mechanism is optimal when the Gaussian mean width of K is asymptotically as large possible. Our goal in this and the following section is to show general lower bounds on sample complexity in terms of $\ell^*(K)$. This is motivated by the sample complexity upper bound in terms of $\ell^*(K)$ provided by the projection mechanism.

It is natural to follow the strategy from Section 5: use Dvoretzky's criterion to find a nearlyspherical projection of K of appropriate radius and dimension. An inspection of the proof of 7 shows that the sample complexity lower bound we get this way is $\Omega\left(\frac{\ell^*(K)^2}{\sqrt{m}}\right)$ (ignoring the dependence on ε , δ , and α here, and in the rest of this informal discussion). Recall that we are aiming for a lower bound of of $\Omega(\ell^*(K))$, so we are off by a factor of $\frac{\ell^*(K)}{\sqrt{m}}$. Roughly speaking, the problem is that Dvoretzky's criterion does too much: it guarantees a spherical section of K° , while we only need a bound on the diameter of the section. In order to circumvent this difficulty, we use a different result from asymptotic convex geometry, the low M^* -estimate, which bounds the diameter of a random section of K° , without also producing a large ball contained inside the section. A technical difficulty is that the resulting upper bound on the diameter is in terms of the Gaussian mean K-norm, rather than the (reciprocal of the) mean width. When K is in ℓ -position, this is not an issue, because results of Pisier, Figiel, and Tomczak-Jaegermann show that in that case $\ell(K)\ell^*(K) = O(\log m)$. In this section we assume that K is in ℓ -position, and we remove this requirement in the subsequent section.

The main result of this section is summarized below.

Theorem 9. For all $\varepsilon = O(1)$, $2^{-\Omega(n)} \leq \delta \leq 1/n^{1+\Omega(1)}$, all symmetric convex bodies $K \subseteq \mathbb{R}^m$ in ℓ -position, and for $\alpha \leq \frac{\ell^*(K)}{C\sqrt{m}\log 2m}$, where C is an absolute constant:

$$\operatorname{sc}_{\varepsilon,\delta}(K,\alpha) = \Omega\left(\frac{\sqrt{\log 1/\delta}}{\alpha\varepsilon} \cdot \frac{\ell^*(K)}{\log 2m}\right).$$

The following two theorems are the main technical ingredients we need in the proof of Theorem 9.

Theorem 10 ([FTJ79], [Pis80]; MM^* Bound). There exists a constant C such that for every symmetric convex body $K \subset \mathbb{R}^m$ in ℓ -position:

$$\ell(K) \cdot \ell^*(K) \le C \cdot m \log 2m.$$

It is an open problem whether this bound can be improved to $m\sqrt{\log 2m}$. This would be tight for the cube Q^m . This improvement would lead to a corresponding improvement in our bounds.

Theorem 11 ([PTJ86]; Low M^* estimate). There exists a constant C such that for every symmetric convex body $K \subset \mathbb{R}^m$ there exists a subspace $E \subseteq \mathbb{R}^m$ with dim E = m - k for which:

diam
$$(K \cap E) \le C \cdot \frac{\ell^*(K)}{\sqrt{k}}$$

Combining Theorems 10 and 11, we get the following key lemma.

Lemma 7. There exists a constant C such that for every symmetric convex body $K \subset \mathbb{R}^m$ in ℓ -position, and every $\beta \in (0, 1-1/m)$, there exists a subspace E of dimension at least βm satisfying:

diam
$$(K \cap E) \le C \frac{\sqrt{m \log 2m}}{\sqrt{1 - \beta} \cdot \ell(K)}$$

Proof. Let $k = \lfloor (1 - \beta)m \rfloor \ge 1$. Using the low M^* estimate on K, there exists a subspace E with dim $E = m - k = \lceil \beta m \rceil$ for which:

diam
$$(K \cap E) \le C_1 \cdot \frac{\ell^*(K)}{\sqrt{k}}.$$

By the MM^* upper bound, since K is in ℓ -position, we have that:

$$\ell^*(K) \le C_2 \cdot \frac{m \log 2m}{\ell(K)},$$

and, combining the two inequalities, we get that:

diam
$$(K \cap E) \le C_1 C_2 \frac{m \log 2m}{\sqrt{k} \cdot \ell(K)} \le C \frac{\sqrt{m} \log m}{\sqrt{1 - \beta} \cdot \ell(K)},$$

for an appropriate constant C. This completes the proof.

The proof of the desired lower bound now follows easily from this lemma.

Proof of Theorem 9. By Theorem 6, it suffices to show that

$$\max_{k=1}^{m} \frac{\sqrt{m-k+1}}{c_k(K^\circ)} = \Omega\left(\frac{\ell^*(K)}{\log 2m}\right).$$
(2)

Indeed, if k^* is the value of k for which the maximum on the left hand side is achieved, then $\frac{\sqrt{m-k^*+1}}{c_{k^*}(K^\circ)}$ is a lower bound on the sample complexity for all $\alpha \leq 1/(10c_{k^*}(K^\circ))$, and by (2):

$$\frac{1}{10c_{k^*}(K^\circ)} = \Omega\left(\frac{\ell^*(K)}{\sqrt{m-k+1}\cdot\log 2m}\right) = \Omega\left(\frac{\ell^*(K)}{\sqrt{m}\cdot\log 2m}\right).$$

In the rest of the proof, we establish (2).

Since K (and thus also K°) are in ℓ -position by assumption, from Lemma 7 applied to K° we have that there exists a subspace E such that dim $E \ge m/2$ and:

diam
$$(K^{\circ} \cap E) = O\left(\frac{\sqrt{m}\log 2m}{\ell(K^{\circ})}\right) = O\left(\frac{\sqrt{m}\log 2m}{\ell^{*}(K)}\right).$$

Setting $k_E = m - \dim E + 1 \le m/2 + 1$, and because $c_{k_E}(K^\circ) \le \operatorname{diam}(K^\circ \cap E)$ by definition, we get that:

$$\max_{k=1}^{m} \frac{\sqrt{m-k+1}}{c_k(K^\circ)} \ge \frac{\sqrt{\dim E}}{\operatorname{diam}(K^\circ \cap E)} = \Omega\left(\frac{\ell^*(K)}{\log 2m}\right),$$

as desired.

7 Gaussian Width Lower Bounds for Arbitrary Bodies

In this section, we remove the assumption that K is in ℓ -position from the previous section. Instead, we use a recursive charging argument in order to reduce to the ℓ -position case. The resulting guarantee is worse than the one we proved for bodies in ℓ -position by a logarithmic factor.

The main lower bound result of this section is the following theorem.

Theorem 12. For all $\varepsilon = O(1)$, $2^{-\Omega(n)} \leq \delta \leq 1/n^{1+\Omega(1)}$, any symmetric convex body $K \subset \mathbb{R}^m$, and any $\alpha \leq \frac{\ell^*(K)}{C\sqrt{m}(\log 2m)^2}$, where C is an absolute constant:

$$\operatorname{sc}_{\varepsilon,\delta}(K,\alpha) = \Omega\left(\frac{\sigma(\varepsilon,\delta)\ell^*(K)}{(\log 2m)^2\alpha}\right)$$

The lower bound follows from the geometric lemma below, which is interesting in its own right.

Lemma 8. There exists a constant C such that, for any symmetric convex body $K \subset \mathbb{R}^m$,

$$\ell^*(K) \le C(\log 2m) \left(\sum_{i=1}^m \frac{1}{\sqrt{i} \cdot c_{m-i+1}(K^\circ)} \right).$$
 (3)

Lemma 8 is closely related to the volume number theorem of Milman and Pisier [MP87], which states that the inequality (3) holds with $\frac{1}{c_{m-i+1}(K^{\circ})}$ replaced by the volume number $v_i(K)$, defined as:

$$v_i(K) = \sup_{E:\dim E=i} \frac{\operatorname{vol}(\Pi_E(K))^{1/i}}{(\operatorname{vol}(\Pi_E(B_2^m))^{1/i})},$$

where the supremum is over subspaces E of \mathbb{R}^m . Inequality (3) is stronger than the volume number theorem, because $\frac{1}{c_{m-i+1}(K^\circ)} \leq v_i(K)$. Indeed, setting $r = \frac{1}{c_{m-i+1}(K^\circ)}$, by Corollary 1 and the definition of Gelfand width we have that there exists a subspace E of dimension i such that $r\Pi_E(B_2^m) \subseteq \Pi_E K$. Therefore, $\operatorname{vol}(\Pi_E(K)) \geq r^i \operatorname{vol}(\Pi_E(B_2^m))$, which implies the desired inequality.

Even though the volume number theorem is weaker than (3), the proof given by Pisier in his book [Pis89], with minor modifications, appears to yield the stronger inequality we need. Rather than repeat this argument, we give a self-contained and slightly different proof below. Our proof only uses the low M^* estimate, the MM^* estimate, and elementary linear algebra, while Pisier's proof uses Milman's quotient of subspace theorem. Moreover, if the MM^* estimate can be improved to $O(m\sqrt{\log 2m})$, our argument would imply a corresponding improvement of the logarithmic factor in (3) from $\log 2m$ to $\sqrt{\log 2m}$. This does not appear to be the case in Pisier's proof, where the logarithmic factor comes from the (tight) upper estimate of the K-convexity constant of mdimensional Banach spaces.

To prove Lemma 8 we will first establish an auxiliary lemma.

Lemma 9. There exists a constant C such that, for any symmetric convex body $K \subset \mathbb{R}^m$ and $m \ge 4$, there exists a subspace $E \subseteq \mathbb{R}^m$ such that dim $E \ge \lfloor m/4 \rfloor$ and:

$$\ell^*(\Pi_E(K)) \le C(\log 2m) \frac{\sqrt{m}}{c_{2\lfloor m/4 \rfloor + 1}(K^\circ)}.$$

Proof. Let T be a linear operator such that T(K) is in ℓ -position. By the rotational invariance of Gaussians, $\ell^*(UT(K)) = \ell^*(T(K))$ and $\ell(UT(K)) = \ell(T(K))$ for any orthogonal transformation U; so, we may assume that T is self-adjoint and positive definite.

Let $\lambda_1 \geq \ldots \geq \lambda_m > 0$ be the eigenvalues of T. Let us set $k = \lfloor m/4 \rfloor$ and define E to be the subspace spanned by the eigenvectors of T corresponding to $\lambda_1, \ldots, \lambda_k$. Observe that Π_E and T commute because they are simultaneously diagonalized by the eigenvectors of T.

Using Lemma 3, we calculate:

$$\ell^*(\Pi_E(K)) = \ell^*(\Pi_E T^{-1}T(K)) \le \|\Pi_E T^{-1}\|\ell^*(T(K)).$$

By the definition of E, the singular values of $\Pi_E T^{-1}$ are $\lambda_k^{-1} \ge \ldots \ge \lambda_1^{-1}$, and therefore the operator norm of $\Pi_E T^{-1}$ is $\|\Pi_E T^{-1}\| = \lambda_k^{-1}$. Thus we have:

$$\ell^*(\Pi_E(K)) \le \frac{\ell^*(T(K))}{\lambda_k}.$$
(4)

Since T(K), and thus $T(K)^{\circ}$, is in ℓ -position, we can apply Lemma 7 to $T(K)^{\circ}$ and $\beta = \frac{m-k}{m} \ge \frac{1}{4}$ and get that there exists a subspace F of dimension at least m - k so that:

$$\ell^*(T(K)) \le C_1(\log 2m) \frac{\sqrt{m}}{\operatorname{diam}(T(K)^\circ \cap F)},$$

where C_1 is an absolute constant.

Let $G = E^{\perp}$ be the orthogonal complement of the subspace E, i.e. the space spanned by the eigenvectors of T corresponding to the eigenvalues $\lambda_{k+1}, \ldots, \lambda_m$. The restriction $T|_G : G \to \mathbb{R}^m$ of T to G has eigenvalues $\lambda_{k+1} \ge \ldots \ge \lambda_m$. Therefore $||T|_G|| \le \lambda_{k+1} \le \lambda_k$. Because G is spanned by eigenvectors of T, it is invariant under action by T, i.e. T(G) = G, and, equivalently, $T^{-1}(G) = G$. Therefore $T^{-1}(K^{\circ}) \cap G = T^{-1}(K^{\circ} \cap G)$ and $K^{\circ} \cap G = (T|_G)T^{-1}(K^{\circ} \cap G)$. It follows that:

$$\operatorname{diam}(K^{\circ} \cap F \cap G) = \operatorname{diam}((T|_G)T^{-1}(K^{\circ} \cap G) \cap F)$$
$$\leq ||T|_G|| \operatorname{diam}(T^{-1}(K^{\circ} \cap G) \cap F)$$
$$\leq \lambda_k \operatorname{diam}(T^{-1}(K^{\circ}) \cap F)$$
$$= \lambda_k \operatorname{diam}(T(K)^{\circ} \cap F).$$

Setting $H = F \cap G$ and combining the inequalities, we get that:

$$\ell^*(T(K)) \le C_1(\log 2m) \frac{\lambda_k \sqrt{m}}{\operatorname{diam}(K^\circ \cap H)} \le C_1(\log 2m) \frac{\lambda_k \sqrt{m}}{c_{2k+1}(K^\circ)},$$

where the final inequality follows from the definition of Gelfand width, since the co-dimension of H is at most 2k. Sibstituting this inequality into the right hand side of (4) finishes the proof of the lemma.

We are now ready to prove Lemma 8.

Proof of Lemma 8. Let us first establish the lemma for $m \leq 16$. Notice that:

$$c_m(K^{\circ}) = \min_{\theta: \|\theta\|_2 = 1} \operatorname{diam}(K^{\circ} \cap \{t\theta : t \in \mathbb{R}\}).$$

But diam $(K^{\circ} \cap \{t\theta : t \in \mathbb{R}\}) = \frac{1}{\|\theta\|_{K^{\circ}}}$ for any unit vector θ , by the definition of Minkowski norm. Therefore, $1/c_m(K^{\circ}) = \max\{\|\theta\|_{K^{\circ}} : \|\theta\|_2 = 1\}$. It follows that, for a standard Gaussian g in \mathbb{R}^m :

$$\ell^*(K) = \mathbb{E} \|g\|_{K^\circ} \le \frac{\mathbb{E} \|g\|_2}{c_m(K^\circ)} \le \frac{(\mathbb{E} \|g\|_2^2)^{1/2}}{c_m(K^\circ)} = \frac{\sqrt{m}}{c_m(K^\circ)} \le \frac{4}{c_m(K^\circ)}.$$
(5)

This establishes the lemma for $m \leq 16$.

For m > 16, we will use an induction argument. We will strengthen the induction hypothesis to:

$$\ell^*(K) \le C(\log 2k) \left(\sum_{i=1}^{\lceil k/2 \rceil} \frac{1}{\sqrt{i} \cdot c_{k-i+1}(K^\circ)} \right).$$
(6)

Assume the inequality holds for a sufficiently large absolute constant C and all symmetric convex bodies $K \subseteq \mathbb{R}^k$, in any dimension k < m. We will show that the inequality then holds in dimension k = m as well. The inequality (5), provides the base case for the induction $(k \le 16)$.

We proceed with the inductive step. By Lemma 9 there exists a subspace E of dimension at least |m/4| such that:

$$\ell^*(\Pi_E(K)) \le C_1(\log 2m) \frac{\sqrt{m}}{c_{2\lfloor m/4 \rfloor + 1}(K^\circ)}.$$

Observe that, since $2\lfloor m/4 \rfloor \leq \lfloor m/2 \rfloor$, and $c_j(K^\circ)$ is monotone non-increasing in j, we have that $c_{2\lfloor m/4 \rfloor+1}(K^\circ) \geq c_{m-i+1}(K^\circ)$ for any $i \leq \lceil m/2 \rceil$. Moreover,

$$\sum_{i=\lceil 3m/8\rceil+1}^{\lfloor m/2 \rfloor} \frac{1}{\sqrt{i}} \ge \frac{1}{C_2} \sqrt{m},$$

for all m > 16 and an absolute constant C_2 . These observations together imply that

$$\ell^*(\Pi_E(K)) \le C_1(\log 2m) \frac{\sqrt{m}}{c_{2\lfloor m/4 \rfloor + 1}(K^\circ)} \le C_1 C_2(\log 2m) \sum_{i = \lceil 3m/8 \rceil + 1}^{\lfloor m/2 \rfloor} \frac{1}{\sqrt{i} \cdot c_{m-i+1}(K^\circ)}.$$
 (7)

For a standard Gaussian $g \sim N(0, I_m)$, using the triangle inequality and the fact that $I_m = \Pi_E + \Pi_{E^{\perp}}$, we have:

$$\ell^*(K) = \mathbb{E} \max_{x \in K} |\langle x, g \rangle|$$

= $\mathbb{E} \max_{x \in K} |\langle \Pi_E x, g \rangle + \Pi_{E^{\perp}} x, g \rangle|$
 $\leq \mathbb{E} \max_{x \in K} |\langle \Pi_E x, g \rangle| + \mathbb{E} \max_{x \in K} |\langle \Pi_{E^{\perp}} x, g \rangle|$
= $\ell^*(\Pi_E K) + \ell^*(\Pi_{E^{\perp}} K).$

Then, the inductive hypothesis follows by (7) and by the inductive hypothesis (6) applied to $\ell^*(\Pi_{E^{\perp}})$. This finishes the proof of the lemma.

Proof of Theorem 12. As in the proof of Theorem 9, it is sufficient to prove that:

$$\max_{k=1}^{m} \frac{\sqrt{m-k+1}}{c_k(K^{\circ})} = \Omega\left(\frac{\ell^*(K)}{(\log 2m)^2}\right).$$
(8)

But this inequality follows easily from Lemma 8 and the trivial case of Hölder's inequality:

$$\begin{split} \ell^*(K) &\leq C(\log 2m) \left(\sum_{i=1}^m \frac{1}{\sqrt{i} \cdot c_{m-i+1}(K^\circ)} \right) \leq C(\log 2m) \left(\sum_{i=1}^m \frac{1}{i} \right) \cdot \left(\max_{i=1}^m \frac{\sqrt{i}}{c_{m-i+1}(K^\circ)} \right) \\ &= O((\log 2m)^2) \cdot \left(\max_{i=1}^m \frac{\sqrt{i}}{c_{m-i+1}(K^\circ)} \right). \end{split}$$

Then, the proof of the theorem follows from (8) analogously to the proof of Theorem 9.

 \Box

We now have everything in place to prove our main result for the mean point problem.

Proof of Theorem 2. The upper bounds on sample complexity follow from Theorem 4, Corollary 2, Theorem 5, and Corollary 3. The lower bounds follow from Theorem 12. The statement after "moreover" follows from Corollary 2 and Theorem 7.

8 From Mean Point to Query Release

All the bounds we proved so far were for the mean point problem. In this section we show reductions between this problem, and the query release problem, which allow us to translate our lower bounds to the query release setting and prove Theorem 1. We will show that the problem of approximating $\mathcal{Q}(\mathcal{D})$ for a query workload \mathcal{Q} under differential privacy is nearly equivalent to approximating the mean point problem with universe $K' = \frac{1}{\sqrt{m}}K$, where K is the sensitivity polytope of \mathcal{Q} . The main technical lemma follows.

Lemma 10. Let \mathcal{Q} be a workload of m linear queries over the universe \mathcal{U} with sensitivity polytope K. Define $K' = \frac{1}{\sqrt{m}}K$. Then, we have the inequalities:

$$\operatorname{sc}_{\varepsilon,\delta}(\mathcal{Q},\alpha) \le \operatorname{sc}_{\varepsilon,\delta}(K',\alpha);$$
(9)

$$\operatorname{sc}_{\varepsilon,\delta}(K',\alpha) \le \max\left\{\operatorname{sc}_{2\varepsilon,2\delta}(\mathcal{Q},\alpha/4), \frac{16\operatorname{diam}(K')}{\alpha^2}\right\}.$$
 (10)

Moreover, we can use an (ε, δ) -differentially private algorithm \mathcal{A}' as a black box to get an (ε, δ) differentially private algorithm \mathcal{A} such that $\operatorname{sc}(\mathcal{Q}, \mathcal{A}, \alpha) = \operatorname{sc}(K', \mathcal{A}', \alpha)$. \mathcal{A} makes a single call to \mathcal{A}' , and performs additional computation of worst-case complexity O(mn), where n is the size of the database.

Proof. Note that inequality (9) is implied by the statement after "moreover". We prove this claim first. The algorithm \mathcal{A} uses \mathcal{D} to form a database \mathcal{D}' of elements from K' which contains, for each $e \in \mathcal{D}$, a copy of $\frac{1}{\sqrt{m}}\mathcal{Q}(\{e\})$. This transformation clearly takes time O(mn). Then \mathcal{A} simply outputs $\sqrt{m}\mathcal{A}'(\mathcal{D}')$. Because any two neighboring databases \mathcal{D}_1 , \mathcal{D}_2 drawn from \mathcal{U} are transformed into neighboring databases \mathcal{D}'_1 , \mathcal{D}'_2 of points from K', and \mathcal{A}' was assumed to be (ε, δ) -differentially private, we have that \mathcal{A} is (ε, δ) -differentially private as well. Also, since $\overline{\mathcal{D}}' = \frac{1}{\sqrt{m}}\mathcal{Q}(\mathcal{D})$, $\operatorname{err}(\mathcal{Q}, \mathcal{A}, n) = \operatorname{err}(K', \mathcal{A}, n)$ by definition, which implies that $\operatorname{sc}(\mathcal{Q}, \mathcal{A}, \alpha) = \operatorname{sc}(K', \mathcal{A}', \alpha)$.

The second inequality (10) is more challenging. We will show that for any (ε, δ) -differentially private algorithm \mathcal{A} there exists an $(2\varepsilon, 2\delta)$ -differentially private algorithm \mathcal{A}' such that:

$$\operatorname{err}(K', \mathcal{A}', n) \le 2 \operatorname{err}(\mathcal{Q}, \mathcal{A}, n) + \frac{2 \operatorname{diam}(K')}{\sqrt{n}}$$

A simple calculation then shows that this implies the desired inequality. In constructing \mathcal{A}' we will run \mathcal{A} on a database formed by sampling from the vertices of K' (which correspond to universe elements) so that the true query answers are preserved in expectation. The analysis uses symmetrization.

Let \mathcal{A}' be given the input $\mathcal{D}' = \{x_1, \ldots, x_n\}$, which is a multiset of points from K'. \mathcal{A}' will randomly construct two databases \mathcal{D}_+ and \mathcal{D}_- (i.e. multisets of elements from \mathcal{U}), and output $\frac{1}{\sqrt{m}}(\mathcal{A}(\mathcal{D}_+) - \mathcal{A}(\mathcal{D}_-))$.

Next we describe how \mathcal{D}_+ and \mathcal{D}_- are sampled. Observe that, for each i, x_i is a convex combination of vertices of K'. Therefore, by Caratheodory's theorem, there exist universe elements $e_{i,1}, \ldots, e_{i,k_i} \in \mathcal{U}$, where $k_i \leq m+1$, such that $x_i = \frac{1}{\sqrt{m}} \sum_{j=1}^{k_i} \alpha_{i,j} \mathcal{Q}(e_{i,j})$ for some $\alpha_{i,1}, \ldots, \alpha_{i,k_i} \in [-1, 1]$ satisfying $\sum_{j=1}^{k_i} |\alpha_i| = 1$. We would like to fix a unique way to pick the $e_{i,j}$ and $\alpha_{i,j}$ for each x_i , and indeed for any point in K'. To this end, fix an arbitrary order on \mathcal{U} , and for each x_i choose a minimal sequence $e_{i,1}, \ldots, e_{i,k_i}$ that satisfies the conditions above and which comes earliest in the lexicographic order induced by the order on \mathcal{U} . Once we have chosen the $e_{i,j}$ and $\alpha_{i,j}$, we construct \mathcal{D}_+ and \mathcal{D}_- using the following sampling procedure: for each $i \in [n]$, we independently sample j_i from $[k_i]$, s.t. $\mathbb{P}[j_i = j] = |\alpha_{i,j}|$; we add e_{i,j_i} to \mathcal{D}_+ if $\alpha_{i,j_i} \geq 0$, and to \mathcal{D}_- otherwise. Then, as mentioned above, \mathcal{A}' outputs $\frac{1}{\sqrt{m}}(\mathcal{A}(\mathcal{D}_+) - \mathcal{A}(\mathcal{D}_-))$.

First, we show that \mathcal{A}' is $(2\varepsilon, 2\delta)$ -differentially private. Let $\mathcal{D}'_1 = \{x_1, \ldots, x_n\}$ and $\mathcal{D}'_2 = \{x_1, \ldots, x_n, x_{n+1}\}$ be two neighboring databases and let $\{e_{i,j}\}$ and $\{\alpha_{i,j}\}$ be defined as above, where $1 \leq i \leq n+1$ and for each $i, 1 \leq j \leq k_i$. Let $\mathcal{D}^{(1)}_+, \mathcal{D}^{(1)}_-$ be the random databases sampled by \mathcal{A}' on input \mathcal{D}'_1 , and let $\mathcal{D}^{(2)}_+, \mathcal{D}^{(2)}_-$ be the random databases sampled on input \mathcal{D}'_2 . Let us couple $\mathcal{D}^{(1)}_+$ and $\mathcal{D}^{(2)}_+$ so that they have symmetric difference at most 1 and $\mathcal{D}^{(2)}_+$ is a superset of $\mathcal{D}^{(1)}_+$. We

can achieve this by sampling $j_i \in [k_i]$ as described above for $1 \leq i \leq n$ and adding e_{i,j_i} to both $\mathcal{D}^{(1)}_+$ and $\mathcal{D}^{(2)}_+$ if $\alpha_{i,j_i} \geq 0$ (or to neither otherwise) and sampling $j_{n+1} \in [k_{n+1}]$ and adding $e_{n+1,j_{n+1}}$ to $\mathcal{D}^{(2)}_+$ if $\alpha_{n+1,j_{n+1}} \geq 0$. Because, with this coupling, $\mathcal{D}^{(1)}_+$ and $\mathcal{D}^{(2)}_+$ are always neighboring, by the (ε, δ) -differential privacy guarantee for \mathcal{A} we have that, for any measurable S in the range of \mathcal{A} :

$$\mathbb{P}[\mathcal{A}(\mathcal{D}^{(1)}_+) \in S] \le e^{\varepsilon} \mathbb{P}[\mathcal{A}(\mathcal{D}^{(2)}_+) \in S] + \delta,$$
$$\mathbb{P}[\mathcal{A}(\mathcal{D}^{(1)}_+) \in S] \ge e^{-\varepsilon} \mathbb{P}[\mathcal{A}(\mathcal{D}^{(2)}_+) \in S] - \delta,$$

where the probabilities are taken over the random choices of $\mathcal{D}^{(1)}_+$ and $\mathcal{D}^{(2)}_+$, and over the randomness of \mathcal{A} . An analogous argument holds for $\mathcal{D}^{(1)}_-$ and $\mathcal{D}^{(2)}_-$. Therefore, each of the two calls of \mathcal{A} made by \mathcal{A}' , composed with the sampling procedure that produces the input to \mathcal{A} , is (ε, δ) -differentially private with respect to the input to \mathcal{A}' . The privacy claim follows by composition (Lemma 4).

Finally we analyze the error of \mathcal{A}' . By Minkowski's inequality, we have:

$$\begin{aligned} (\mathbb{E} \|\mathcal{A}'(\mathcal{D}') - \overline{\mathcal{D}}'\|_{2}^{2})^{1/2} &\leq \left(\mathbb{E} \left\|\mathcal{A}(\mathcal{D}_{+}) - \frac{1}{\sqrt{m}}\mathcal{Q}(\mathcal{D}_{+})\right\|_{2}^{2}\right)^{1/2} + \left(\mathbb{E} \left\|\mathcal{A}(\mathcal{D}_{-}) - \frac{1}{\sqrt{m}}\mathcal{Q}(\mathcal{D}_{-})\right\|_{2}^{2}\right)^{1/2} \\ &+ \left(\mathbb{E} \left\|\frac{1}{\sqrt{m}}(\mathcal{Q}(\mathcal{D}_{+}) - \mathcal{Q}(\mathcal{D}_{-})) - \overline{\mathcal{D}}'\right\|_{2}^{2}\right)^{1/2} \\ &\leq 2\operatorname{err}(\mathcal{Q}, \mathcal{A}, n) + \left(\mathbb{E} \left\|\frac{1}{\sqrt{m}}(\mathcal{Q}(\mathcal{D}_{+}) - \mathcal{Q}(\mathcal{D}_{-})) - \overline{\mathcal{D}}'\right\|_{2}^{2}\right)^{1/2}. \end{aligned}$$

We proceed to bound the second term on the right hand side. Observe that, by the way we sample \mathcal{D}_+ and \mathcal{D}_- , linearity of expectation, and the linearity of $\mathcal{Q}(\cdot)$, $\overline{\mathcal{D}}' = \mathbb{E}\frac{1}{\sqrt{m}}(\mathcal{Q}(\mathcal{D}_+) - \mathcal{Q}(\mathcal{D}_-))$. Let, for each $i \in [n]$, $j_i \in [k_i]$ be sampled as above, and let $y_i = \frac{1}{\sqrt{m}}\operatorname{sign}(\alpha_{i,j})\mathcal{Q}(e_{i,j_i})$. Then $\frac{1}{\sqrt{m}}(\mathcal{Q}(\mathcal{D}_+) - \mathcal{Q}(\mathcal{D}_-)) = \frac{1}{n}\sum_{i=1}^n y_i$, so $\mathbb{E}\frac{1}{n}\sum_{i=1}^n y_i = \overline{\mathcal{D}}'$. The vectors y_1, \ldots, y_n are independent and all belong to $K' \subseteq R \cdot B_2^m$, where $R = \operatorname{diam}(K')$. Let ξ_1, \ldots, ξ_n be Rademacher random variables, independent from everything else; by Lemma 1 and the parallelogram identity:

$$\mathbb{E} \left\| \frac{1}{\sqrt{m}} (\mathcal{Q}(\mathcal{D}_{+}) - \mathcal{Q}(\mathcal{D}_{-})) - \overline{\mathcal{D}}' \right\|_{2}^{2} = \mathbb{E} \left\| \frac{1}{n} \sum_{i=1}^{n} y_{i} - \mathbb{E} \frac{1}{n} \sum_{i=1}^{n} y_{i} \right\|_{2}^{2}$$
$$\leq \frac{4}{n^{2}} \mathbb{E} \left\| \sum_{i=1}^{n} \xi_{i} y_{i} \right\|_{2}^{2} = \frac{4}{n^{2}} \sum_{i=1}^{n} \|y_{i}\|_{2}^{2} \leq \frac{4R^{2}}{n}.$$

With this, we have established the desired bound on $\operatorname{err}(K', \mathcal{A}', n)$, and, therefore, the lemma.

We will also use a simple lemma that relates the sample complexity at an error level α to the sample complexity at a lower error level $\alpha' < \alpha$. The proof is a padding argument and can be found in [SU15].

Lemma 11. For any workload Q, any $0 < \alpha' < \alpha < 1$, and any privacy parameters ε, δ , we have

$$\operatorname{sc}_{\varepsilon,\delta}(\mathcal{Q},\alpha') = \Omega\left(\frac{\alpha}{C\alpha'}\right) \cdot \operatorname{sc}_{\varepsilon,\delta}(\mathcal{Q},\alpha),$$

for an absolute constant C.

We are now ready to finish the proof of our main result for query release.

Proof of Theorem 1. The upper bounds on sample complexity follow from the upper bounds in Theorem 2 together with Lemma 10.

Denote $K' = \frac{1}{\sqrt{m}}K$, and let $\alpha_0 = \frac{\ell^*(K')}{C\sqrt{m}(\log 2m)^2} = \frac{\ell^*(K)}{Cm(\log 2m)^2}$ be the smallest error parameter for which Theorem 12 holds. Then, by Theorem 12 and Lemma 10:

$$\max\left\{\mathrm{sc}_{\varepsilon,\delta}(\mathcal{Q},\alpha_0),\frac{\mathrm{diam}(K)}{\sqrt{m\alpha_0^2}}\right\} = \Omega\left(\frac{\sigma(\varepsilon,\delta)\ell^*(K)}{\sqrt{m}(\log 2m)^2\alpha_0}\right).$$

It is easy to show that $\operatorname{sc}_{\varepsilon,\delta}(\mathcal{Q},\alpha_0) = \Omega(\operatorname{diam}(K)/(\alpha_0\sqrt{m}))$ for all sufficiently small ε and δ . Therefore, we have:

$$\operatorname{sc}_{\varepsilon,\delta}(\mathcal{Q},\alpha_0) = \Omega\left(\frac{\sigma(\varepsilon,\delta)\ell^*(K)}{\sqrt{m}(\log 2m)^2}\right).$$

By Lemma 11, we get that for any $\alpha \leq \alpha_0$ the sample complexity is at least:

$$\mathrm{sc}_{\varepsilon,\delta}(\mathcal{Q},\alpha) = \Omega\left(\frac{\sigma(\varepsilon,\delta)\ell^*(K)\alpha_0}{\sqrt{m}(\log 2m)^2\alpha}\right) = \Omega\left(\frac{\sigma(\varepsilon,\delta)\ell^*(K)^2}{m^{3/2}(\log 2m)^4\alpha}\right).$$

An analogous proof, with $\alpha_0 = 1/C$ set to the smallest error parameter for which Theorem 7 holds, establishes the statement after "moreover".

References

- [AAGM15] Shiri Artstein-Avidan, Apostolos Giannopoulos, and Vitali D. Milman. Asymptotic geometric analysis. Part I, volume 202 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2015.
- [BDKT12] Aditya Bhaskara, Daniel Dadush, Ravishankar Krishnaswamy, and Kunal Talwar. Unconditional differentially private mechanisms for linear queries. In *Proceedings of the* 44th symposium on Theory of Computing, STOC '12, pages 1269–1284, New York, NY, USA, 2012. ACM.
- [BUV14] Mark Bun, Jonathan Ullman, and Salil Vadhan. Fingerprinting codes and the price of approximate differential privacy. In *Proceedings of the 46th Annual ACM Symposium* on Theory of Computing, pages 1–10. ACM, 2014.
- [DMNS06] Cynthia Dwork, Frank McSherry, Kobbi Nissim, and Adam Smith. Calibrating noise to sensitivity in private data analysis. In *Theory of Cryptography Conference*, pages 265–284. Springer, 2006.
- [DN03] Irit Dinur and Kobbi Nissim. Revealing information while preserving privacy. In Proceedings of the 22nd ACM SIGMOD-SIGACT-SIGART Symposium on Principles of Database Systems, pages 202–210. ACM, 2003.
- [DN04] Cynthia Dwork and Kobbi Nissim. Privacy-preserving datamining on vertically partitioned databases. In Annual International Cryptology Conference, pages 528–544. Springer, 2004.
- [DNT14] Cynthia Dwork, Aleksandar Nikolov, and Kunal Talwar. Using convex relaxations for efficiently and privately releasing marginals. In 30th Annual Symposium on Computational Geometry, SOCG'14, Kyoto, Japan, June 08 - 11, 2014, page 261. ACM, 2014.

- [FTJ79] T. Figiel and Nicole Tomczak-Jaegermann. Projections onto Hilbertian subspaces of Banach spaces. Israel J. Math., 33(2):155–171, 1979.
- [HT10] Moritz Hardt and Kunal Talwar. On the geometry of differential privacy. In Proceedings of the 42nd ACM Symposium on Theory of Computing, STOC '10, pages 705–714, New York, NY, USA, 2010. ACM.
- [Kea98] Michael Kearns. Efficient noise-tolerant learning from statistical queries. J. ACM, 45(6):983–1006, 1998.
- [Mat02] Jiří Matoušek. Lectures on discrete geometry, volume 212 of Graduate Texts in Mathematics. Springer-Verlag, New York, 2002.
- [Mil71] V. D. Milman. A new proof of A. Dvoretzky's theorem on cross-sections of convex bodies. *Funkcional. Anal. i Priložen.*, 5(4):28–37, 1971.
- [MP87] V. D. Milman and G. Pisier. Gaussian processes and mixed volumes. Ann. Probab., 15(1):292–304, 1987.
- [Nik15] Aleksandar Nikolov. An improved private mechanism for small databases. In Automata, Languages, and Programming - 42nd International Colloquium, ICALP 2015, Kyoto, Japan, July 6-10, 2015, Proceedings, Part I, volume 9134 of Lecture Notes in Computer Science, pages 1010–1021. Springer, 2015.
- [NTZ13] Aleksandar Nikolov, Kunal Talwar, and Li Zhang. The geometry of differential privacy: the sparse and approximate cases. In *Symposium on Theory of Computing Conference*, *STOC'13*, *Palo Alto, CA, USA, June 1-4, 2013*, pages 351–360. ACM, 2013.
- [Pin85] Allan Pinkus. n-widths in approximation theory, volume 7 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin, 1985.
- [Pis80] G. Pisier. Sur les espaces de Banach K-convexes. In Seminar on Functional Analysis, 1979–1980 (French), pages Exp. No. 11, 15. École Polytech., Palaiseau, 1980.
- [Pis89] Gilles Pisier. The volume of convex bodies and Banach space geometry, volume 94 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1989.
- [PTJ86] Alain Pajor and Nicole Tomczak-Jaegermann. Subspaces of small codimension of finitedimensional Banach spaces. Proc. Amer. Math. Soc., 97(4):637–642, 1986.
- [SU15] Thomas Steinke and Jonathan Ullman. Between pure and approximate differential privacy. *CoRR*, abs/1501.06095, 2015.
- [Ver09] R. Vershynin. Lectures in geometric functional analysis. http://www-personal.umich.edu/ romanv/papers/GFA-book/GFA-book.pdf, 2009.