

# On the generalized porous medium equation in Fourier-Besov spaces

Weiliang Xiao<sup>1</sup> Xuhuan Zhou<sup>2\*</sup>

*School of Applied Mathematics, Nanjing University of Finance and Economics, Nanjing, 210023, China;*

*E-mail: xwltc123@163.com*

*Department of Information Technology, Nanjing Forest Police College, Nanjing, 210023, China;*

*E-mail: zhouxuhuan@163.com*

**Abstract** We study a kind of generalized porous medium equation with fractional Laplacian and abstract pressure term. For a large class of equations corresponding to the form:  $u_t + \nu \Lambda^\beta u = \nabla \cdot (u \nabla P u)$ , we get their local well-posedness in Fourier-Besov spaces for large initial data. If the initial data is small, then the solution becomes global. Furthermore, we prove a blowup criterion for the solutions.

**Keywords** porous medium equation, well-posedness, blowup criterion, Fourier-Besov spaces

**MR (2010)** 42B37, 35Q35, 35K55, 76S05

## 1 Introduction

In this paper, we study the nonlinear nonlocal equation in  $\mathbb{R}^n$  of the form

$$\begin{cases} u_t + \nu \Lambda^\alpha u = \nabla \cdot (u \nabla P u); \\ u(0, x) = u_0. \end{cases} \quad (1.1)$$

Usually,  $u = u(t, x)$  is a real-valued function, represents a density or concentration. The dissipative coefficient  $\nu > 0$  corresponds to the viscous case, while  $\nu = 0$  corresponds to the inviscid case. In this paper we study the viscous case and take  $\nu = 1$  for simplicity. The fractional operator  $\Lambda^\alpha$  is defined by Fourier transform as  $(\Lambda^\alpha u)^\wedge = |\xi|^\alpha \hat{u}$ .  $P$  is an abstract operator.

Equation (1.1) here comes from the same proceeding with that of the fractional porous medium equation (FPME) introduced by Caffarelli and Vázquez [5]. In fact, equation (1.1) comes into being by adding the fractional dissipative term  $\nu \Lambda^\alpha u$  to the continuity equation  $u_t + \nabla \cdot (uV) = 0$ , where the velocity  $V = -\nabla p$  and the velocity potential or pressure  $p$  is related to  $u$  by an abstract operator  $p = Pu$ .

The abstract form pressure term  $Pu$  gives a good suitability in many cases. The simplest case comes from a model in groundwater in filtration [1, 19]:  $u_t = \Delta u^2$ , that is:  $\nu = 0, Pu =$

---

The research was supported by the NNSF of China under grant #11601223 and # 11626213.

$u$ . A more general case appears in the fractional porous medium equation [5] when  $\nu = 0$  and  $Pu = \Lambda^{-2s}u, 0 < s < 1$ . In the critical case when  $s = 1$ , it is the mean field equation first studied by Lin and Zhang [15]. Some studies on the well-posedness and regularity on those equations we refer to [4, 6, 7, 17, 18, 20, 23] and the references therein.

In the FPME, the pressure can also be represented by Riesz potential as  $Pu = \Lambda^{-2s}u = \mathcal{K} * u$ , with kernel  $\mathcal{K} = c_{n,s}|y|^{2s-n}$ . Replacing the kernel  $\mathcal{K}$  by other functions in this form:  $Pu = \mathcal{K} * u$ , equation (1.1) also appears in granular flow and biological swarming, named aggregation equation. The typical kernels are the Newton potential  $|x|^\gamma$  and the exponent potential  $-e^{-|x|}$ .

One of concerned problems on this equation is the singularity of the potential  $Pu$  which holds the well-posedness or leads to the blowup solution. Generally, smooth kernels at origin  $x = 0$  lead to the global in time solution [3], meanwhile nonsmooth kernels may lead to blowup phenomenon [14]. Li and Rodrigo [13, 14] studied the well-posedness and blowup criterion of equation (1.1) with the pressure  $Pu = \mathcal{K} * u$ , where  $\mathcal{K}(x) = e^{-|x|}$  in Sobolev spaces. Wu and Zhang [21] generalize their work to require  $\nabla\mathcal{K} \in W^{1,1}$  which includes the case  $\mathcal{K}(x) = e^{-|x|}$ . They take advantage of the controllability in Besov spaces of the convolution  $\mathcal{K} * u$  under this condition, as well as the controllability of its gradient  $\nabla\mathcal{K} * u$ .

In this article we study the well-posedness and blowup criterion of equation (1.1) in Fourier-Besov spaces under an abstract pressure condition

$$\|\widehat{\Delta_k(\nabla Pu)}\|_{L^p} \leq C2^{k\sigma} \|\widehat{\Delta_k u}\|_{L^p}. \quad (1.2)$$

In Fourier-Besov spaces, it is the localization express of the norm estimate

$$\|\nabla Pu\|_{F\dot{B}_{p,q}^s} \leq C\|u\|_{F\dot{B}_{p,q}^{s+\sigma}}. \quad (1.3)$$

Corresponding to the FPME, i.e.  $Pu = \Lambda^{-2s}$ , we get  $\sigma = 1 - 2s$  obviously. And if  $Pu = \mathcal{K} * u$ ,  $\mathcal{K} \in W^{1,1}$  in the aggregation equation, we get  $\sigma = 1$  when  $\mathcal{K} \in L^1$  and  $\sigma = 0$  when  $\nabla\mathcal{K} \in L^1$ .

The Fourier-Besov spaces we use here come from Konieczny and Yoneda [11] when deal with the Navier-Stokes equation (NSE) with Coriolis force. Besides, Fourier-Besov spaces have been widely used to study the well-posedness, singularity, self-similar solution, etc. of Fluid Dynamics in various of forms. For instance, the early pseudomeasure spaces  $PM^\alpha$  in which Cannone and Karch studied the smooth and singular properties of Navier-Stokes equations [8]. The Lei-Lin spaces  $\mathcal{X}^\sigma$  deal with global solutions to the NSE [12] and to the quasi-geostrophic equations (QGE) [2]. The Fourier-Herz spaces  $\mathcal{B}_q^\sigma$  in the Keller-Segel system [9], in the NSE with Coriolis force [10] and in the magneto-hydrodynamic equations (MHD) [16].

In the case of Besov spaces, we gain some well-posedness and blow-up results of equation (1.1) under an corresponding condition to (1.3) in [24]. Due to the difficulty in deal with the nonlinear term  $\nabla \cdot (u\nabla Pu)$ , in that case we need a little strict initial condition:  $u_0 \in \dot{B}_{p,1}^{n/p+\sigma-\beta} \cap \dot{B}_{p,1}^{n/p+\sigma-\beta+1}$ . However, in this paper, we find that Fourier-Besov spaces are

powerful in deal with the nonlinear term, by a very different method used in [24], we prove the following theorems:

**Theorem 1.1.** *Let  $p, q \in [1, \infty]$ ,  $\max\{1, \sigma + 1\} < \alpha < n(1 - \frac{1}{p}) + \sigma + 2$ . Then for any  $u_0 \in \dot{F}B_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}$ , equation (1.1) admits a unique mild solution  $u$  and*

$$u \in C([0, T]; \dot{F}B_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}) \cap \tilde{L}^1([0, T]; \dot{F}B_{p,q}^{n(1-\frac{1}{p})+\sigma+1}).$$

Moreover, there exists a constant  $C_0 = C_0(\alpha, p, q)$  such that for  $u_0$  satisfying  $\|u_0\|_{\dot{F}B_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}} \leq C_0$ , the solution  $u$  is global, and

$$\|u\|_{L_T^\infty(\dot{F}B_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1})} + \|u\|_{\tilde{L}_T^1(\dot{F}B_{p,q}^{n(1-\frac{1}{p})+\sigma+1})} \leq 2C\|u_0\|_{\dot{F}B_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}}.$$

**Theorem 1.2.** *Let  $T^*$  denote the maximal time of existence of  $u$  in  $L_T^\infty(\dot{F}B_{p,q}^\beta) \cap \tilde{L}_T^1(\dot{F}B_{p,q}^{\beta+\alpha})$ . Here  $\beta = n(1 - \frac{1}{p}) - \alpha + \sigma + 1$ . If  $T^* < \infty$ , then*

$$\|u\|_{\tilde{L}^1([0, T^*]; \dot{F}B_{p,q}^{\beta+\alpha})} = \infty.$$

## 2 Preliminaries

Let us introduce some basic knowledge on Littlewood-Paley theory and Fourier-Besov spaces. Let  $\varphi \in C_c^\infty(\mathbb{R}^n)$  be a radial positive function such that

$$\text{supp } \varphi \subset \{\xi \in \mathbb{R}^n : \frac{3}{4} \leq |\xi| \leq \frac{8}{3}\}, \quad \sum_{j \in \mathbb{Z}} \varphi(2^{-j}\xi) = 1 \text{ for any } \xi \neq 0.$$

Define the frequency localization operators as follows:

$$\Delta_j u = \varphi_j(D)u; \quad S_j u = \psi_j(D)u,$$

here  $\varphi_j(\xi) = \varphi(2^{-j}\xi)$  and  $\psi_j = \sum_{k \leq j-1} \varphi_k$ .

By Bony's decomposition we can split the product  $uv$  into three parts:

$$uv = T_u v + T_v u + R(u, v),$$

with

$$T_u v = \sum_j S_{j-1} u \Delta_j v, \quad R(u, v) = \sum_j \Delta_j u \tilde{\Delta}_j v, \quad \tilde{\Delta}_j v = \Delta_{j-1} v + \Delta_j v + \Delta_{j+1} v.$$

Let us now define the Fourier-Besov space as follows.

**Definition 2.1.** For  $\beta \in \mathbb{R}, p, q \in [1, \infty]$ , we define the Fourier-Besov space  $\dot{F}B_{p,q}^\beta$  as

$$\dot{F}B_{p,q}^\beta = \{f \in \mathcal{S}'/\mathbb{P} : \|f\|_{\dot{F}B_{p,q}^\beta} = \left( \sum_{j \in \mathbb{Z}} 2^{j\beta q} \|\varphi_j \hat{f}\|_{L^p}^q \right)^{1/q} < \infty\}.$$

Here the norm changes normally when  $p = \infty$  or  $q = \infty$ , and  $\mathbb{P}$  is the set of all polynomials.

**Definition 2.2.** In this paper, we need two kinds of mixed time-space norm defined as follows: For  $s \in \mathbb{R}, 1 \leq p, q \leq \infty, I = [0, T), T \in (0, \infty]$ , and let  $X$  be a Banach space with norm  $\|\cdot\|_X$ ,

$$\begin{aligned} \|f(t, x)\|_{L^r(I; X)} &:= \left( \int_I \|f(\tau, \cdot)\|_X^r d\tau \right)^{\frac{1}{r}}, \\ \|f(t, x)\|_{\tilde{L}^r(I; \dot{F}B_{p,q}^\beta)} &:= \left( \sum_{j \in \mathbb{Z}} 2^{j\beta q} \|\varphi_j \hat{f}\|_{L^r(I; L^p)}^q \right)^{\frac{1}{q}}. \end{aligned}$$

By Minkowski' inequality, there holds

$$L^r(I; \dot{F}B_{p,q}^s) \hookrightarrow \tilde{L}^r(I; \dot{F}B_{p,q}^s), \text{ if } r \leq q \quad \text{and} \quad \tilde{L}^r(I; \dot{F}B_{p,q}^s) \hookrightarrow L^r(I; \dot{F}B_{p,q}^s), \text{ if } r \geq q. \quad (2.1)$$

**Lemma 2.1.** Let  $X$  be a Banach space with norm  $\|\cdot\|_X$  and  $B : X \times X \mapsto X$  be a bounded bilinear operator satisfying

$$\|B(u, v)\|_X \leq \eta \|u\|_X \|v\|_X,$$

for all  $u, v \in X$  and a constant  $\eta > 0$ . Then for any fixed  $y \in X$  satisfying  $\|y\|_X < \epsilon < \frac{1}{4\eta}$ , the equation  $x := y + B(x, x)$  has a solution  $\bar{x}$  in  $X$  such that  $\|\bar{x}\|_X \leq 2\|y\|_X$ . Also, the solution is unique in  $\bar{B}(0, 2\epsilon)$ . Moreover, the solution depends continuously on  $y$  in the sense: if  $\|y'\|_X < \epsilon, x' = y' + B(x', x'), \|x'\|_X < 2\epsilon$ , then

$$\|\bar{x} - x'\|_X \leq \frac{1}{1 - 4\epsilon\eta} \|y - y'\|_X.$$

**Lemma 2.2.** [22] If  $\frac{1}{r} = \frac{1-\theta}{r_1} + \frac{\theta}{r_2}$ , then

$$\|u\|_{\tilde{L}_T^{r_1}(\dot{F}B_{p,q}^{\beta+\theta\alpha})} \leq \|u\|_{\tilde{L}_T^{r_1}(\dot{F}B_{p,q}^\beta)}^{1-\theta} \|u\|_{\tilde{L}_T^{r_2}(\dot{F}B_{p,q}^{\beta+\alpha})}^\theta.$$

Now we prove a priori estimate which will be used in our proof. Consider the following dissipative equation:

$$\partial_t u + \Lambda^\alpha u = f(t, x), \quad u(0, x) = u_0(x), \quad t > 0, x \in \mathbb{R}^n. \quad (2.2)$$

**Lemma 2.3.** Let  $0 < T \leq \infty, \beta \in \mathbb{R}$  and  $1 \leq r \leq \infty$ . Assume  $u_0 \in \dot{F}B_{p,q}^\beta, f \in \tilde{L}^1(I; \dot{F}B_{p,q}^\beta)$ . Then the solution  $u(t, x)$  to (2.2) satisfies

$$\|u\|_{\tilde{L}^r(I; \dot{F}B_{p,q}^{\beta+\frac{\alpha}{r}})} \leq C(\|u_0\|_{\dot{F}B_{p,q}^\beta} + \|f\|_{\tilde{L}^1(I; \dot{F}B_{p,q}^\beta)}) \quad (2.3)$$

*Proof.* Consider the integral form of (2.2)

$$u(t, x) = e^{-t\Lambda^\alpha} u_0 + \int_0^t e^{-(t-\tau)\Lambda^\alpha} f(\tau, x) d\tau := Lu_0 + Gf.$$

For the linear part,

$$\|\varphi_j \mathcal{F}(Lu_0)\|_{L^p} \leq e^{-t2^{j\alpha}(3/4)^\alpha} \|\varphi_j \widehat{u_0}\|_{L^p}.$$

Hence

$$\|Lu_0\|_{\tilde{L}_T^r(FB_{p,q}^{\beta+\frac{\alpha}{r}})} \leq \|2^{j(\beta+\frac{\alpha}{r})}\| e^{-t2^{j\alpha}(3/4)^\alpha} \|L_T^r \|\varphi_j \widehat{u_0}\|_{L^p}\|_{l^q} \leq C \|u_0\|_{FB_{p,q}^\beta}.$$

On the other hand, for the integral part,

$$\|\varphi_j \mathcal{F}(Gf)\|_{L^p} \leq \int_0^t e^{-(t-\tau)(\frac{3}{4}2^j)^\alpha} \|\varphi_j \hat{f}\|_{L^p} d\tau$$

Taking  $L^r$ -norm with respect to time, by Minkowskii's inequality

$$\|\varphi_j \mathcal{F}(Gf)\|_{L_T^r(L^p)} \leq C 2^{-\frac{j\alpha}{r}} \|\varphi_j \hat{f}\|_{L_T^1(L^p)}.$$

Hence

$$\|Gf\|_{\tilde{L}_T^r(FB_{p,q}^{\beta+\frac{\alpha}{r}})} = \|2^{j(\beta+\frac{\alpha}{r})}\| \|\varphi_j \mathcal{F}(Gf)\|_{L_T^r(L^p)}\|_{l^q} \leq C \|f\|_{\tilde{L}_T^1(FB_{p,q}^\beta)}.$$

Combine the above estimates, we obtain our desire inequality.  $\square$

### 3 Local and global well-posedness

In this section we prove our main Theorem. First we know that the integral form of  $u$  is as follows

$$\begin{aligned} u &= e^{-t\Lambda^\alpha} u_0 + \int_0^t e^{-(t-\tau)\Lambda^\alpha} \nabla \cdot (u(\tau) \nabla P u(\tau)) d\tau \\ &:= S(t)u_0 + H(u, u). \end{aligned} \tag{3.1}$$

Here  $S(t)u_0 = \mathcal{F}^{-1}(e^{-t|\xi|^\alpha} \hat{u}_0)$ , and  $H(u, v) = \int_0^t e^{-(t-\tau)\Lambda^\alpha} \nabla \cdot (u(\tau) \nabla P v(\tau)) d\tau$ . Now we get the following estimate

**Proposition 3.1.** *Let  $\gamma, p, q \geq 1$ ,  $\epsilon > \max\{0, -\sigma\}$ ,  $\beta > 0$ ,  $\frac{1}{\gamma} = \frac{1}{\gamma_1} + \frac{1}{\gamma_2}$ , there holds*

$$\begin{aligned} \|u \partial_i P v\|_{\tilde{L}_t^\gamma(FB_{p,q}^\beta)} &\leq C \|u\|_{\tilde{L}_t^{\gamma_1}(FB_{p,q}^{n(1-\frac{1}{p})-\epsilon})} \|v\|_{\tilde{L}_t^{\gamma_2}(FB_{p,q}^{\beta+\sigma+\epsilon})} \\ &\quad + C \|v\|_{\tilde{L}_t^{\gamma_1}(FB_{p,q}^{n(1-\frac{1}{p})-\epsilon})} \|u\|_{\tilde{L}_t^{\gamma_2}(FB_{p,q}^{\beta+\sigma+\epsilon})}. \end{aligned}$$

*Proof.* By the Bony's decomposition, it is easy to get that

$$\Delta_j(u\partial_i P v) = \sum_{|k-j|\leq 4} \Delta_j(S_{k-1}u\Delta_k(\partial_i P v)) \quad (3.2)$$

$$+ \sum_{|k-j|\leq 4} \Delta_j(S_{k-1}(\partial_i P v)\Delta_k u) + \sum_{k\geq j-2} \Delta_j(\Delta_k u\tilde{\Delta}_k(\partial_i P v)) \quad (3.3)$$

We can get the following estimates:

$$\begin{aligned} & 2^{j\beta} \left\| \sum_{|k-j|\leq 4} \varphi_j \mathcal{F}(S_{k-1}u\Delta_k(\partial_i P v)) \right\|_{L_t^\gamma(L^p)} \\ & \leq 2^{j\beta} \sum_{|k-j|\leq 4} \left\| \mathcal{F}(S_{k-1}u) * \varphi_k \mathcal{F}(\partial_i P v) \right\|_{L_t^\gamma(L^p)} \\ & \leq C 2^{j\beta} \sum_{|k-j|\leq 4} \left\| \sum_{l\leq k-2} 2^{ln(1-\frac{1}{p})} \|\varphi_l \hat{u}\|_{L^p} 2^{k\sigma} \|\varphi_k \hat{v}\|_{L^p} \right\|_{L_t^\gamma} \\ & \leq C 2^{j\beta} \sum_{|k-j|\leq 4} \left\| \sum_{l\leq k-2} 2^{(l-k)\epsilon} 2^{nl(1-\frac{1}{p})-\epsilon l} \|\varphi_l \hat{u}\|_{L^p} 2^{k(\sigma+\epsilon)} \|\varphi_k \hat{v}\|_{L^p} \right\|_{L_t^\gamma} \\ & \leq C \|u\|_{\tilde{L}_t^{\gamma_1}(FB_{p,q}^{n(1-\frac{1}{p})-\epsilon})} 2^{j(\beta+\sigma+\epsilon)} \|\varphi_j \hat{v}\|_{L_t^{\gamma_2}(L^p)}. \end{aligned}$$

Similarly, we can estimate

$$\begin{aligned} & 2^{j\beta} \left\| \sum_{|k-j|\leq 4} \varphi_j \mathcal{F}(S_{k-1}(\partial_i P v)\Delta_k u) \right\|_{L_t^\gamma(L^p)} \\ & \leq C \|v\|_{\tilde{L}_t^{\gamma_1}(FB_{p,q}^{n(1-\frac{1}{p})-\epsilon})} 2^{j(\beta+\sigma+\epsilon)} \|\varphi_k \hat{u}\|_{L_t^{\gamma_2}(L^p)}. \end{aligned}$$

And when  $\beta > 0$ , we can also get that for some  $\|a_j\|_{l^q} = 1$ ,

$$\begin{aligned} & 2^{j\beta} \left\| \mathcal{F} \left( \sum_{k\geq j-2} \Delta_j(\Delta_k(\partial_i P v)\tilde{\Delta}_k u) \right) \right\|_{L_t^\gamma(L^p)} \\ & \leq 2^{j\beta} \left\| \sum_{k\geq j-2} \varphi_j \mathcal{F}(\Delta_k(\partial_i P v)) * (\tilde{\varphi}_k \hat{u}) \right\|_{L_t^\gamma(L^p)} \\ & \leq C 2^{j\beta} \left\| \sum_{k\geq j-2} 2^{kn(1-\frac{1}{p})+k\sigma} \|\varphi_k \hat{v}\|_{L^p} \|\tilde{\varphi}_k \hat{u}\|_{L^p} \right\|_{L_t^\gamma} \\ & \leq C \sum_{k\geq j-2} 2^{(j-k)\beta} \left\| 2^{kn(1-\frac{1}{p})-\epsilon k} \|\varphi_k \hat{v}\|_{L^p} 2^{k(\beta+\sigma+\epsilon)} \|\tilde{\varphi}_k \hat{u}\|_{L^p} \right\|_{L_t^\gamma} \\ & \leq C \sum_{k\geq j-2} 2^{(j-k)\beta} \|v\|_{\tilde{L}_t^{\gamma_1}(FB_{p,q}^{n(1-\frac{1}{p})-\epsilon})} 2^{k(\beta+\sigma+\epsilon)} \|\tilde{\varphi}_k u\|_{\tilde{L}_t^{\gamma_2}(L^p)} \\ & \leq C a_j \|v\|_{\tilde{L}_t^{\gamma_1}(FB_{p,q}^{n(1-\frac{1}{p})-\epsilon})} \|u\|_{\tilde{L}_t^{\gamma_2}(FB_{p,q}^{\beta+\sigma+\epsilon})}. \end{aligned}$$

Combine the above estimate, we obtain the following inequality

$$\begin{aligned} \|u\partial_i P v\|_{\tilde{L}_t^\gamma(FB_{p,q}^\beta)} &\leq C\|u\|_{\tilde{L}_t^{\gamma_1}(FB_{p,q}^{n(1-\frac{1}{p})-\epsilon})} \|v\|_{\tilde{L}_t^{\gamma_2}(FB_{p,q}^{\beta+\sigma+\epsilon})} \\ &\quad + C\|v\|_{\tilde{L}_t^{\gamma_1}(FB_{p,q}^{n(1-\frac{1}{p})-\epsilon})} \|u\|_{\tilde{L}_t^{\gamma_2}(FB_{p,q}^{\beta+\sigma+\epsilon})}. \end{aligned}$$

□

**Proposition 3.2.** *Let  $p, q \geq 1$ ,  $\epsilon > \max\{0, -\sigma\}$ ,  $\beta > -1$ . If  $u, v \in \tilde{L}_T^2(FB_{p,q}^{n(1-\frac{1}{p})-\epsilon}) \cap \tilde{L}_T^2(FB_{p,q}^{\beta+\sigma+\epsilon+1})$ , we have*

$$\begin{aligned} &\|H(u, v)\|_{\tilde{L}_T^\infty(FB_{p,q}^\beta)} + \|H(u, v)\|_{\tilde{L}_T^1(FB_{p,q}^{\beta+\alpha})} \\ &\leq C\|u\|_{\tilde{L}_T^2(FB_{p,q}^{n(1-\frac{1}{p})-\epsilon})} \|v\|_{\tilde{L}_T^2(FB_{p,q}^{\beta+\sigma+\epsilon+1})} + C\|v\|_{\tilde{L}_T^2(FB_{p,q}^{n(1-\frac{1}{p})-\epsilon})} \|u\|_{\tilde{L}_T^2(FB_{p,q}^{\beta+\sigma+\epsilon+1})}. \end{aligned}$$

*Proof.* We note that  $H(u, v)$  is a solution to equation (2.2) with  $u_0 = 0, f = \nabla \cdot (u\nabla P v)$ . So by Lemma 2.3 there holds

$$\|H(u, v)\|_{\tilde{L}_T^\infty(FB_{p,q}^\beta)} + \|H(u, v)\|_{\tilde{L}_T^1(FB_{p,q}^{\beta+\alpha})} \leq C\|\nabla \cdot (u\nabla P v)\|_{\tilde{L}_T^1(FB_{p,q}^\beta)}.$$

Then Proposition 3.1 gives the estimate. □

**Theorem 3.1.** *Let  $p, q \in [1, \infty]$ ,  $2 \max\{1, \sigma + 1\} < \alpha < n(1 - \frac{1}{p}) + \sigma + 2$ . Then for any  $u_0 \in FB_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}$ , equation (1.1) admits a unique mild solution  $u$  and*

$$u \in C([0, T]; FB_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}) \cap \tilde{L}^1([0, T]; FB_{p,q}^{n(1-\frac{1}{p})+\sigma+1}).$$

Moreover, there exists a constant  $C_0 = C_0(\alpha, p, q)$  such that for  $u_0$  satisfying  $\|u_0\|_{FB_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}} \leq C_0$ , the solution  $u$  is global, and

$$\|u\|_{L_T^\infty(FB_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1})} + \|u\|_{\tilde{L}_T^1(FB_{p,q}^{n(1-\frac{1}{p})+\sigma+1})} \leq 2C\|u_0\|_{FB_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}}.$$

*Proof.* First suppose  $t \in [0, T]$ ,  $T$  fixed. Let  $\epsilon = \frac{\alpha}{2} - \sigma - 1$ ,  $\beta = n(1 - \frac{1}{p}) - \alpha + \sigma + 1$ , by the above proposition

$$\begin{aligned} &\|H(u, v)\|_{L_T^\infty(FB_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1})} + \|H(u, v)\|_{\tilde{L}_T^1(FB_{p,q}^{n(1-\frac{1}{p})+\sigma+1})} \\ &\leq C\|u\|_{\tilde{L}_T^2(FB_{p,q}^{n(1-\frac{1}{p})-\frac{\alpha}{2}+\sigma+1})} \|v\|_{\tilde{L}_T^2(FB_{p,q}^{n(1-\frac{1}{p})-\frac{\alpha}{2}+\sigma+1})}. \end{aligned}$$

Define  $X = \tilde{L}_T^2(FB_{p,q}^{n(1-\frac{1}{p})-\frac{\alpha}{2}+\sigma+1})$ , by Lemma 2.2

$$\|H(u, v)\|_X \leq C\|u\|_X \|v\|_X.$$

By Lemma 2.1 we know that if  $\|e^{-t\Lambda^\alpha} u_0\|_X < \frac{1}{4C}$ , then (3.1) has a unique solution in  $B(0, \frac{1}{2C})$ . Here  $B(0, \frac{1}{2C}) := \{x \in X : \|x\|_X \leq \frac{1}{2C}\}$ .

To conclude  $\|e^{-t\Lambda^\alpha} u_0\|_X < \frac{1}{4C}$ , we first note that  $e^{-t\Lambda^\alpha} u_0$  is the solution to (2.2) with  $f = 0, u_0 = u_0$ , by Lemma 2.3, we can obtain

$$\|e^{-t\Lambda^\alpha} u_0\|_X \leq C \|u_0\|_{\dot{F}B_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}}. \quad (3.4)$$

Hence if  $\|u_0\|_{\dot{F}B_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}} \leq \frac{1}{4C^2}$ , (3.1) has a unique global solution in  $X$ . Moreover,  $\|u\|_X \leq 2C \|u_0\|_{\dot{F}B_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}}$ .

On the other hand, denote  $u_0 = \mathcal{F}^{-1} \chi_{\{|\xi| \leq \lambda\}} \widehat{u}_0 + \mathcal{F}^{-1} \chi_{\{|\xi| > \lambda\}} \widehat{u}_0 := u_0^1 + u_0^2$ , where  $\lambda = \lambda(u_0) > 0$  is a real number determined later. Since  $u_0^2$  converges to 0 in  $\dot{F}B_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}$  as  $\lambda \rightarrow +\infty$ . By (3.4) there exists  $\lambda$  large enough such that

$$\|e^{-t\Lambda^\alpha} u_0^2\|_X \leq \frac{1}{8C}.$$

For  $u_0^1$ ,

$$\begin{aligned} \|e^{-t\Lambda^\alpha} u_0^1\|_X &= \|2^{j(n(1-\frac{1}{p})-\frac{\alpha}{2}+\sigma+1)} \|\varphi_j e^{-t|\xi|^\alpha} \chi_{\{|\xi| \leq \lambda\}} \widehat{u}_0\|_{L_T^2(L^p)}\|_{l^q} \\ &\leq \|2^{j(n(1-\frac{1}{p})-\frac{\alpha}{2}+\sigma+1)} \|\sup_{|\xi| \leq \lambda} e^{-t|\xi|^\alpha} |\xi|^{\frac{\alpha}{2}}\|_{L_T^2} \|\varphi_j |\xi|^{-\frac{\alpha}{2}} \widehat{u}_0\|_{L^p}\|_{l^q} \\ &\leq C \lambda^{\frac{\alpha}{2}} T^{\frac{1}{2}} \|u_0\|_{\dot{F}B_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}}. \end{aligned}$$

Thus for arbitrary  $u_0$  in  $\dot{F}B_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}$ , (3.1) has a unique local solution in  $X$  on  $[0, T)$  where

$$T \leq \left( \frac{1}{8C^2 \lambda^{\frac{\alpha}{2}} \|u_0\|_{\dot{F}B_{p,q}^{n(1-\frac{1}{p})-\alpha+\sigma+1}}} \right)^2.$$

The continuity with respect to time is standard.  $\square$

Next we give a blowup criterion as following:

**Theorem 3.2.** *Let  $T^*$  denote the maximal time of existence of  $u$  in  $L_T^\infty(\dot{F}B_{p,q}^\beta) \cap \tilde{L}_T^1(\dot{F}B_{p,q}^{\beta+\alpha})$ . Here  $\beta = n(1 - \frac{1}{p}) - \alpha + \sigma + 1$ . If  $T^* < \infty$ , then*

$$\|u\|_{\tilde{L}^1([0, T^*]; \dot{F}B_{p,q}^{\beta+\alpha})} = \infty. \quad (3.5)$$

*Proof.* Supposing  $\|u\|_{\tilde{L}^1([0, T^*]; \dot{F}B_{p,q}^{\beta+\alpha})} dt < \infty$ , then we can find  $0 < T_0 < T^*$  satisfying

$$\|u\|_{\tilde{L}^1([T_0, T^*]; \dot{F}B_{p,q}^{\beta+\alpha})} < \frac{1}{2}.$$



For  $t \in [T_0, T^*]$ ,  $s \in [T_0, t]$ , by Lemma 2.3

$$\begin{aligned} \|u(s)\|_{\dot{F}B_{p,q}^\beta} + \|u\|_{\tilde{L}^1([T_0,s];\dot{F}B_{p,q}^{\beta+\alpha})} &\leq \|u(T_0)\|_{\dot{F}B_{p,q}^\beta} + \|u\|_{L^\infty([T_0,s];\dot{F}B_{p,q}^\beta)} \|u\|_{\tilde{L}^1([T_0,s];\dot{F}B_{p,q}^{\beta+\alpha})} \\ &\leq \|u(T_0)\|_{\dot{F}B_{p,q}^\beta} + \frac{1}{2} \|u\|_{L^\infty([T_0,s];\dot{F}B_{p,q}^\beta)}. \end{aligned}$$

So,

$$\sup_{T_0 \leq s \leq t} \|u(s)\|_{\dot{F}B_{p,q}^\beta} \leq \|u(T_0)\|_{\dot{F}B_{p,q}^\beta} + \frac{1}{2} \|u\|_{L^\infty([T_0,t];\dot{F}B_{p,q}^\beta)}.$$

Put

$$M = \max(2\|u(T_0)\|_{\dot{F}B_{p,q}^\beta}, \max_{t \in [0, T_0]} \|u\|_{\dot{F}B_{p,q}^\beta}),$$

we can get

$$\|u(t)\|_{\dot{F}B_{p,q}^\beta} \leq M, \quad \forall t \in [0, T^*].$$

On the other side,

$$\begin{aligned} u(t_2) - u(t_1) &= e^{-t_2|D|^\alpha} u_0 - e^{-t_1|D|^\alpha} u_0 + \int_0^{t_2} e^{-(t_2-\tau)|D|^\alpha} \nabla \cdot (uPu)(\tau) d\tau \\ &\quad - \int_0^{t_1} e^{-(t_1-\tau)|D|^\alpha} \nabla \cdot (uPu)(\tau) d\tau \\ &= [e^{-t_2|D|^\alpha} u_0 - e^{-t_1|D|^\alpha} u_0] + \left[ \int_0^{t_1} e^{-(t_1-\tau)|D|^\alpha} (e^{-(t_2-t_1)|D|^\alpha} - 1) \nabla \cdot (uPu)(\tau) d\tau \right] \\ &\quad + \left[ \int_{t_1}^{t_2} e^{-(t_2-\tau)|D|^\alpha} \nabla \cdot (uPu)(\tau) d\tau \right] \\ &:= L_1 + L_2 + L_3. \end{aligned}$$

$$\begin{aligned} \|L_1\|_{\dot{F}B_{p,q}^\beta} &= \|2^{j\beta} \|\varphi_j(e^{-t_2|\xi|^\alpha} - e^{-t_1|\xi|^\alpha}) \hat{u}_0\|_{L^p}\|_{l^q} \\ &\leq \|2^{j\beta} \|\varphi_j(e^{-(t_2-t_1)|\xi|^\alpha} - 1) \hat{u}_0\|_{L^p}\|_{l^q} \\ &\leq \|u_0\|_{\dot{F}B_{p,q}^\beta}. \end{aligned}$$

$$\begin{aligned} \|L_2\|_{\dot{F}B_{p,q}^\beta} &\leq \|2^{j\beta} \int_0^{t_1} \|\varphi_j e^{-(t_2-\tau)|\xi|^\alpha} (1 - e^{-(t_2-t_1)|\xi|^\alpha}) \mathcal{F}(\nabla \cdot (uPu)(\tau))\|_{L^p} d\tau\|_{l^q} \\ &\leq \|2^{j(\beta+1)} \int_0^{t_1} \|\varphi_j (e^{-(t_2-t_1)|\xi|^\alpha} - 1) \mathcal{F}(uPu)(\tau)\|_{L^p} d\tau\|_{l^q}. \end{aligned}$$

$$\|L_3\|_{\dot{F}B_{p,q}^\beta} \leq \|2^{j\beta} \int_{t_1}^{t_2} \|\varphi_j e^{-(t_2-\tau)|\xi|^\alpha} \mathcal{F}(\nabla \cdot (uPu)(\tau))\|_{L^p} d\tau\|_{l^q}$$

$$\leq \left\| 2^{j(\beta+1)} \int_{t_1}^{t_2} \|\varphi_j \mathcal{F}(uPu)(\tau)\|_{L^p} d\tau \right\|_{l^q}.$$

By the dominated convergence theorem, we can get

$$\limsup_{t_1, t_2 \nearrow T^*, t_1 < t_2} \|u(t_2) - u(t_1)\|_{\dot{F}B_{p,q}^\beta} = 0.$$

Then there is an element  $u^*$  of  $\dot{F}B_{p,q}^\beta$  such that

$$\lim_{t \nearrow T^*} \|u(t)\|_{\dot{F}B_{p,q}^\beta} = u^*.$$

Now set  $u(T^*) = u^*$  and consider the equation starting by  $u^*$ , by the well-posedness we obtain a solution existing on a larger time interval than  $[0, T^*)$ , which is a contradiction.  $\square$

## 4 Improvement of the index

The index in Theorem 3.1 is not a nature one since  $\alpha > 2$  is a very strong condition. However, the method used here can in fact gain some better index range by a slight modification. The key point is that we can seek the solution firstly in the space  $X := \tilde{L}_T^r(\dot{F}B_{p,q}^{\beta+\frac{\alpha}{r}}) \cap \tilde{L}_T^{r'}(\dot{F}B_{p,q}^{\beta+\frac{\alpha}{r'}})$  instead of  $\tilde{L}_T^2(\dot{F}B_{p,q}^{\beta+\frac{\alpha}{2}})$ , with  $\beta = n(1 - \frac{1}{p}) - \alpha + \sigma + 1$ . Since the proof are very similar, we list the key steps.

**Step 1:** Taking respectively  $r$  and  $r'$  in Lemma 2.3, and using the proof of Proposition 3.2, we get for  $\epsilon > \max\{0, -\sigma\}$ ,  $\beta > -1$ ,

$$\begin{aligned} & \|H(u, v)\|_{\tilde{L}_T^r(\dot{F}B_{p,q}^{\beta+\frac{\alpha}{r}})} + \|H(u, v)\|_{\tilde{L}_T^{r'}(\dot{F}B_{p,q}^{\beta+\frac{\alpha}{r'}})} \\ & \leq C \|u\|_{\tilde{L}_T^r(\dot{F}B_{p,q}^{n(1-\frac{1}{p})-\epsilon})} \|v\|_{\tilde{L}_T^{r'}(\dot{F}B_{p,q}^{\beta+\sigma+\epsilon+1})} + C \|v\|_{\tilde{L}_T^r(\dot{F}B_{p,q}^{n(1-\frac{1}{p})-\epsilon})} \|u\|_{\tilde{L}_T^{r'}(\dot{F}B_{p,q}^{\beta+\sigma+\epsilon+1})}. \end{aligned}$$

Set  $\epsilon = \frac{\alpha}{r'} - \sigma - 1$ ,  $\beta = n(1 - \frac{1}{p}) - \alpha + \sigma + 1$ , we then gain the important bilinear estimate  $H(u, v) \leq C \|u\|_X \|v\|_X$  under the condition:

$$r' \max\{1, \sigma + 1\} < \alpha < n(1 - \frac{1}{p}) + \sigma + 2.$$

Thus by Lemma 2.1 we know that if  $\|e^{-t\Lambda^\alpha} u_0\|_X < \frac{1}{4C}$ , then equation (3.1) admits a unique solution in  $X$ .

**Step 2:** Now we need to derive  $\|e^{-t\Lambda^\alpha} u_0\|_X < \frac{1}{4C}$ . Since  $e^{-t\Lambda^\alpha} u_0$  is the solution to (2.2) with  $f = 0$ ,  $u_0 = u_0$ , by Lemma 2.3 we gain the global solution in  $X$  for small initial data. On the other hand, we can also obtain the local solution on  $[0, T)$  in  $X$  by the same method in Theorem 3.1 for arbitrary initial data, where

$$T \leq \min \left\{ \left( \frac{1}{16C^2 \lambda^{\frac{\alpha}{r}} \|u_0\|_{\dot{F}B_{p,q}^\beta}} \right)^r, \left( \frac{1}{16C^2 \lambda^{\frac{\alpha}{r'}} \|u_0\|_{\dot{F}B_{p,q}^\beta}} \right)^{r'} \right\}.$$

**Step 3:** We have proved equation has an unique solution in  $X$  under the condition:  $r' \max\{1, \sigma + 1\} < \alpha < n(1 - \frac{1}{p}) + \sigma + 2$ ,  $1 < r < \infty$ . Using the integral form (3.1) and Lemma 2.3, we can deduce

$$\|u\|_{\tilde{L}_T^\infty(FB_{p,q}^\beta)} + \|u\|_{\tilde{L}_T^1(FB_{p,q}^{\beta+\alpha})} \leq C(\|u_0\|_{FB_{p,q}^\beta} + \|u\|_X).$$

Hence  $u$  is also belongs to  $C([0, T]; FB_{p,q}^\beta) \cap \tilde{L}_T^1(FB_{p,q}^{\beta+\alpha})$ . Since  $1 < r < \infty$  in  $X$  can be chose to be a sufficiently large number, we in fact improve the index in Theorem 3.1 to

$$\max\{1, \sigma + 1\} < \alpha < n(1 - \frac{1}{p}) + \sigma + 2.$$

Besides, this improvement make no difference to the proof of Theorem 3.2.

## References

- [1] J. Bear, *Dynamics of Fluids in Porous Media*, Dover, New York, 1972.
- [2] J. Benameur, M. Benhamed, *Global existence of the two-dimensional QGE with sub-critical dissipation*, J. Math. Anal. Appl., 423(2015), 1330-1347.
- [3] A. L. Bertozzi, J. A. Carrillo, T. Laurent, *Blow-up in multidimensional aggregation equations with mildly singular interaction kernels*, Nonlinearity, 22 (2009), 683-710.
- [4] P. Biler, C. Imbert, G. Karch, *The nonlocal porous medium equation: Barenblatt profiles and other weak solutions*, Arch. Ration. Mech. Anal. 215 (2015), 497-529.
- [5] L. Caffarelli, J. L. Vázquez, *Nonlinear porous medium flow with fractional potential pressure*. Arch. Ration. Mech. Anal. 202 (2011), 537-565.
- [6] L. Caffarelli, J. L. Vázquez, *Asymptotic behaviour of a porous medium equation with fractional diffusion*, Discrete Contin. Dyn. Syst. 29 (2011), 1393-1404.
- [7] L. Caffarelli, F. Soria, J. L. Vázquez, *Regularity of solutions of the fractional porous medium flow*, J. Eur. Math. Soc. (JEMS) 15 (2013), 1701-1746.
- [8] M. Cannone, K. Grzegorz, *Smooth or singular solutions to the NavierStokes system?* Journal of Differential Equations 197.2 (2004): 247-274.
- [9] T. Iwabuchi, *Global well-posedness for Keller-Segel system in Besov type spaces*, J. Math. Anal. Appl. 379 (2011), 930-948.
- [10] T. Iwabuchi, R. Takada, *Global well-posedness and ill-posedness for the Navier-Stokes equations with the Coriolis force in function spaces of Besov type*, J. Funct. Anal. 267 (2014), 1321-1337.

- [11] P. Konieczny, T. Yoneda, *On dispersive effect of the Coriolis force for the stationary Navier-Stokes equations*, J. Differential Equations, 250 (2011), 3859-3873.
- [12] Z. Lei, F. Lin, *Global mild solutions of Navier-Stokes equations*, Commun. Pure Appl. Math. 64 (2011), 1297-1304.
- [13] D. Li, J. Rodrigo, *Finite-time singularities of an aggregation equation in  $\mathcal{R}^n$  with fractional dissipation*, Commun. Math. Phys., 287 (2009), 687-703.
- [14] D. Li, J. Rodrigo, *Well-posedness and regularity of solutions of an aggregation equation*, Rev. Mat. Iberoam., 26 (2010), 261-294.
- [15] F. Lin, P. Zhang, *On the hydrodynamic limit of Ginzburg-Landau wave vortices*, Comm. Pure Appl. Math. 55 (2002), 831-856.
- [16] Q. Liu, J. Zhao, *Global well-posedness for the generalized magneto-hydrodynamic equations in the critical Fourier-Herz spaces*, J. Math. Anal. Appl. 420 (2014), 1301-1315.
- [17] S. Serfaty, J. L. Vázquez, *A mean field equation as limit of nonlinear diffusions with fractional Laplacian operators*, Calc. Var. Partial Differential Equations 49 (2014), 1091-1120.
- [18] D. Stan, F. del Teso, J. L. Vázquez, *Finite and infinite speed of propagation for porous medium equations with fractional pressure*, C. R. Math. Acad. Sci. Paris 352 (2014), 123-128.
- [19] J. L. Vázquez, *The Porous Medium Equation. Mathematical theory*. Oxford Mathematical Monographs. The Clarendon Press/Oxford University Press, Oxford, 2007.
- [20] J. L. Vázquez, *Recent progress in the theory of nonlinear diffusion with fractional Laplacian operators*, Discrete Contin. Dyn. Syst. Ser. S 7 (2014), 857-885.
- [21] G. Wu, Q. Zhang, *Global well-posedness of the aggregation equation with supercritical dissipation in Besov spaces*, Z. Angew. Math. Mech. 93 (2013), 882-894.
- [22] W. Xiao, J. Chen, D. Fan, X. Zhou, *Global Well-Posedness and Long Time Decay of Fractional Navier-Stokes Equations in Fourier-Besov Spaces*, Abstract and Applied Analysis., 2014 (2014).
- [23] X. Zhou, W. Xiao, J. Chen, *Fractional porous medium and mean field equations in Besov spaces*, Electron. J. Differential Equations, 2014 (2014), 1-14.
- [24] X. Zhou, W. Xiao, T. Zheng, *Well-posedness and blowup criterion of generalized porous medium equation in Besov spaces*, Electron. J. Differential Equations, 2015(2015), 1-14.