# MULTI-BUMP SOLUTIONS FOR FRACTIONAL NIRENBERG PROBLEM

CHUNGEN LIU<sup>1</sup> AND QIANG REN

ABSTRACT. We consider the multi-bump solutions of the following fractional Nirenberg problem

 $(-\Delta)^s u = K(x)u^{\frac{n+2s}{n-2s}}, \quad u > 0 \quad \text{in } \mathbb{R}^n, \tag{0.1}$ 

where  $s \in (0, 1)$  and n > 2+2s. If K is a periodic function in some k variables with  $1 \le k < \frac{n-2s}{2}$ , we proved that (0.1) has multi-bump solutions with bumps clustered on some lattice points in  $\mathbb{R}^k$ via Lyapunov-Schmidt reduction. It is also established that the equation (0.1) has an infinitemany-bump solutions with bumps clustered on some lattice points in  $\mathbb{R}^n$  which is isomorphic to  $\mathbb{Z}_+^k$ .

### 1. INTRODUCTION AND MAIN RESULTS

The classic Nirenberg problem asks that on the standard sphere  $(\mathbb{S}^n, g_{\mathbb{S}^n})$  with  $n \geq 2$ , whether there exists a function w such that the scalar curvature (Gauss curvature in the dimension 2) of the conformal metric  $g = e^w g_{\mathbb{S}^n}$  equals to a prescribed function  $\tilde{K}$ . This problem is equivalent to solving the following equations

$$-\Delta_{q_{\mathbb{S}^n}} w + 1 = \tilde{K} e^{2w} \quad \text{on } \quad \mathbb{S}^2 \tag{1.1}$$

and

$$-\Delta_{g_{\mathbb{S}^n}}v + \frac{n-2}{4(n-1)}R_{g_{\mathbb{S}^n}}v = \frac{n-2}{4(n-1)}\tilde{K}v^{\frac{n+2}{n-2}} \text{ on } \mathbb{S}^n \text{ for } n \ge 3,$$
(1.2)

where  $R_{g_{\mathbb{S}^n}} = n(n-1)$  is the scalar curvature of  $(\mathbb{S}^n, g_{\mathbb{S}^n})$  and  $v = e^{\frac{n-2}{4}w}$ . The linear operators defined on left-hand side of the equation (1.1) and (1.2) are called the conformal Laplacian on  $\mathbb{S}^n$ .

For any Riemannian manifold (M, g), the conformal Laplacian is defined by  $P_1^g = -\Delta_g + \frac{n-2}{4(n-1)}R_g$ , where  $R_g$  is the scalar curvature of (M, g). Let u > 0 and  $h = u^{\frac{4}{n-2}}g$ , the conformal Laplacian has the following conformally invariant property

$$P_1^g(u\phi) = u^{\frac{n+2}{n-2}} P_1^h(\phi) \text{ for } \phi \in C^{\infty}(M).$$

The Paneitz operator  $P_2^g$  is another interesting conformal invariant operator. It was defined in [27] by

$$P_2^g = \Delta_g^2 + div_g (a_n R_g Id - b_n \mathcal{R}ic_g) \nabla_g + \frac{n-4}{2} Q_g,$$

Date: Completed: December 27, 2021.

<sup>2010</sup> Mathematics Subject Classification. 35R11, 35B25, 35B33.

Key words and phrases. Fractional partial differential equations, Lyapunov-Schmidt reduction, critical exponents.

<sup>&</sup>lt;sup>1</sup>The first author is partially supported by the NSF of China (11471170).

where  $a_n = \frac{(n-2)^2+4}{2(n-1)(n-2)}$ ,  $b_n = -\frac{4}{n-2}$ ,  $\mathcal{R}ic: TM \to TM$  is a (1,1)-tensor operator defined by  $\mathcal{R}ic_i^j = g^{jk}Ric_{ki}$  and  $Q_g = -\frac{2}{(n-2)^2}|Ric_g|^2 + \frac{n^3-4n^2+16n-16}{8(n-1)^2(n-2)^2}R_g^2 - \frac{1}{2(n-1)}\Delta_g R_g$  which is called the Q-curvature of (M, g).

Later on, more conformally covariant elliptic operators were found. The operator  $P_1^g$  and  $P_2^g$  were generalized by Graham, Jenne, Mason and Sparling in [16] to a sequence of integer order conformally covariant elliptic operators  $P_k^g$  for  $k \in \mathbb{N}_+$  if n is odd; and  $k \in \{1, \ldots, \frac{n}{2}\}$  if n is even. Furthermore, any real number order conformally covariant pseudo-differential operator was intrinsically defined by Peterson in [28]. Graham and Zworski in [17] proved that the operators  $P_k^g$  can be considered as the residue of a meromorphic family of scattering operators S(s) at  $s = \frac{n}{2} + k$ . Then a family of non-integer order conformally covariant pseudo-differential operators  $P_s^g$  ( $0 < s < \frac{n}{2}$ ) were naturally defined. Using the localization method in [5], Chang and González [6] showed that for any  $s \in (0, \frac{n}{2})$ , the operator  $P_s^g$  can also be defined as a Dirichlet-to-Neumann operator of a conformally compact Einstein manifold.

The conformally covariant law for  $P_s^g$  means that for any Riemannian manifold (M, g) and a conformal transformation  $h = v^{\frac{4}{n-2s}}g$ , v > 0, there holds

$$P_s^g(v\phi) = v^{\frac{n+2s}{n-2s}} P_s^h(\phi) \quad \text{for } \phi \in C^\infty(M).$$

Especially,  $P_s^g(1)$  is called the  $Q_s$  curvature or s-curvature of (M, g) (see [6] and [18] for example).

The fractional Nirenberg problem was naturally raised on  $Q_s$  curvature, it asks that on the standard sphere  $\mathbb{S}^n$ , whether there exists a function v > 0 such that the  $Q_s$  curvature of the conformal metric  $g = v^{\frac{4}{n-s}}g_{\mathbb{S}^n}$  equals to a prescribed function  $\tilde{K}$ . It can be reduced to the existence of the solution of the following equation

$$P_s^{g_{\mathbb{S}^n}}(v) = \tilde{K}v^{\frac{n+2s}{n-2s}}, \quad v > 0 \quad \text{on } \quad \mathbb{S}^n, \tag{1.3}$$

where  $s \in (0, 1)$ , n > 2s and  $\tilde{K}$  is a given positive function.

It was shown in [4] that the operator  $P_s^{g_{\mathbb{S}^n}}$  is an intertwining operator and can be expressed as

$$P_s^{g_{\mathbb{S}^n}} = \frac{\Gamma(B + \frac{1}{2} + s)}{\Gamma(B + \frac{1}{2} - s)}, \quad B = \sqrt{-\Delta_{\mathbb{S}^n} + \left(\frac{n-1}{2}\right)^2},$$

where  $\Delta_{\mathbb{S}^n}$  is the Beltrami-Laplacian operator. What is more,  $P_s^{g_{\mathbb{S}^n}}$  is more concrete under the stereographic projection. Let

$$F: \mathbb{R}^n \to \mathbb{S}^n \setminus \{\mathcal{N}\}, \quad x \mapsto \left(\frac{2x}{|x|^2 + 1}, \frac{|x|^2 - 1}{|x|^2 + 1}\right)$$

be the inverse of stereographic projection, where  $\mathcal{N}$  is the north pole of  $\mathbb{S}^n$ . Then it holds that

$$P_s^{g_{\mathbb{S}^n}}(\phi) \circ F = |J_F|^{-\frac{n+2s}{2n}} (-\Delta)^s (|J_F|^{\frac{n-2s}{2n}} (\phi \circ F)),$$

where  $(-\Delta)^s$  is the fractional Laplacian defined by

$$(-\Delta)^{s}\phi(x) = C(n,s)P.V. \int_{\mathbb{R}^{n}} \frac{\phi(x) - \phi(y)}{|x - y|^{n + 2s}} dy, \quad \text{where} \ \phi \in C^{\infty}(\mathbb{R}^{n}).$$

If we write  $u = |J_F|^{\frac{n-2s}{2n}} (v \circ F)$  and  $K = \tilde{K} \circ F$ , the equation (1.3) is transformed into

$$(-\Delta)^{s} u = K(x) u^{\frac{n+2s}{n-2s}}, \quad u > 0 \text{ in } \mathbb{R}^{n},$$
 (1.4)

where  $s \in (0, 1)$  and n > 2s. The existence of the solutions to the problem (1.3) has been proved under various conditions (see for exmaple [1,2,7,9–11,18,19]). The compactness of the solutions to (1.3) was studied in [18]. Chen and Zheng [10] found a 2-peak solution when  $K(x) = 1 + \varepsilon \tilde{K}(x)$ has at least two critical points and satisfies some local conditions. What is more, Liu in [25] constructed infinitely many 2-peak solutions when K has a sequence of strictly local maximum points moving to infinity. When K is a radial symmetric function, in [24] and [26] it was showed that (1.4) has infinitely many non-radial solutions.

In this paper, we continue to study the bump solutions or peak solutions of (1.4). Assume that K satisfies the following conditions

 $(H_1) \ 0 < \inf_{\mathbb{R}^n} K \leq \sup_{\mathbb{R}^n} K < \infty;$ 

 $(H_2)$  K(x) is a  $C^{1,1}$  function, and 1-periodic in the first k variables  $x_1, \dots, x_k$ ;

 $(H_3)$  0 is a critical point of K, and in a neighborhood of 0, there is a number  $\beta \in (n-2s,n)$  such that

$$K(x) = K(0) + \sum_{i=1}^{n} a_i |x_i|^{\beta} + R(x),$$

where  $a_i \neq 0$  for i = 1, ..., n,  $\sum_{i=1}^n a_i < 0$ ,  $R(y) \in C^{[\beta]-1,1}$  and  $\sum_{j=0}^{[\beta]} |\nabla^j R(y)| |y|^{-\beta+j} = o(1)$  as  $y \to 0$ . Here  $\nabla^j R(y)$  denote all of the possible derivatives of R(y) of the order j.

We note that the conditions  $(H_1)$ - $(H_3)$  and the condition  $(\mathcal{K})$  in [25] has some intersection. When K satisfies both the condition  $(\mathcal{K})$  in [25] and our conditions  $(H_1)$ - $(H_3)$ , the equation (1.4) has infinitely many 2-peak solutions according to [25].

In this paper, we will show that equation (1.4) has solutions with large number bumps and its bumps located near some lattice points in  $\mathbb{R}^k$  with  $1 \le k < \frac{n-2s}{2}$ .

Let  $Q_m := ([0, m+1)^k \times \mathbf{0}) \cap \mathbb{Z}^n$ , where  $m \in \mathbb{N}_+ \cup \{\infty\}$  and  $\mathbf{0}$  is a zero vector in  $\mathbb{R}^{n-k}$ .

**Theorem 1.1.** Suppose n > 2s + 2,  $m \in \mathbb{N}_+ \cup \{\infty\}$  and  $1 \le k < \frac{n-2s}{2}$ . If K satisfies the conditions  $(H_1)$ - $(H_3)$ , there exists an integer  $l_0 \in \mathbb{N}$ , such that for any integer  $l > l_0$ , the equation (1.4) has a solution with its bumps clustered on  $lQ_m$ .

Notice that  $Q_{\infty}$  is an infinite lattice which isomorphic to  $\mathbb{Z}^k_+$ . So we get an infinite-many-bump solution of the equation (1.5) via Theorem 1.1.

In order to prove Theorem 1.1, we assume K(0) = 1 with no loss of generality. For any positive integer l, define  $\lambda = l^{\frac{n-2s}{\beta-n+2s}}$ . Then we have  $\lambda^{\beta} = (\lambda l)^{n-2s}$ . Using the transformation  $u(x) \mapsto \lambda^{-\frac{n-2s}{2}} u(\frac{x}{\lambda})$ , we can change the equation (1.4) into

$$(-\Delta)^s u = K(\frac{x}{\lambda}) u^{\frac{n+2s}{n-2s}}, \quad u > 0, \quad \text{in } \mathbb{R}^n.$$

$$(1.5)$$

The functional corresponding to equation (1.5) is

$$I(u) = \frac{1}{2} \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u|^2 - \frac{n-2s}{2n} \int_{\mathbb{R}^n} K\left(\frac{x}{\lambda}\right) (u_+)^{\frac{2n}{n-2s}}, \quad u \in \dot{H}^s(\mathbb{R}^n),$$

where  $u_{+} = \max\{u, 0\}$ . The Hilbert space  $\dot{H}^{s}(\mathbb{R}^{n})$  is the completion of  $C_{0}^{\infty}(\mathbb{R}^{n})$  under the Gagliardao semi-norm (cf. [15] for detail)

$$[u]_{\dot{H}^{s}(\mathbb{R}^{n})} := \left( \int_{\mathbb{R}^{n}} \int_{\mathbb{R}^{n}} \frac{|u(x) - u(y)|^{2}}{|x - y|^{n + 2s}} \right)^{\frac{1}{2}} = \left( 2C(n,s) \int_{\mathbb{R}^{n}} |(-\Delta)^{\frac{s}{2}}|^{2} \right)^{\frac{1}{2}},$$

where C(n, s) is a constant depending on n and s. It is well known that  $\dot{H}^{s}(\mathbb{R}^{n})$  can be imbedded into  $L^{2^{*}(s)}(\mathbb{R}^{n})$  and the following Hardy-Littlewood-Sobolev inequality holds

$$S\left(\int_{\mathbb{R}^n} |u|^{2^*(s)}\right)^{\frac{2}{2^*(s)}} \le \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}}u|^2, \quad \text{for } u \in C_0^\infty(\mathbb{R}^n), \tag{1.6}$$

where  $s \in (0,1)$ , n > 2s and  $2^*(s) = \frac{2n}{n-2s}$ . Lieb [23] proved that the extremals corresponding to the best constant S of (1.6) are of the form

$$U_{\xi,\Lambda,C_0} = C_0 \left(\frac{\Lambda}{1 + \Lambda^2 |x - \xi|^2}\right)^{\frac{n-2s}{2}}$$

where  $C_0, \Lambda \in \mathbb{R}_+$  and  $\xi \in \mathbb{R}^n$ .

Choosing a suitable constant  $C_0 = C_0(n, s)$ , we see that the function  $U_{\xi,\Lambda} := U_{\xi,\Lambda,C_0}$  solves the equation

$$(-\Delta)^s u = u^{\frac{n+2s}{n-2s}}, \quad u > 0 \text{ in } \mathbb{R}^n.$$

$$(1.7)$$

Under some decay assumptions, [8, 20, 22] proved that all the solutions of (1.7) are only of the form  $U_{\xi,\Lambda}$ . Furthermore, it was proved in [13] that the solution  $U_{\xi,\Lambda}$  of the equation (1.7) is nondegenerate, *i.e.* any bounded solution of the equation  $(-\Delta)^s \phi = \frac{n+2s}{n-2s} U_{\xi,\Lambda}^{\frac{4s}{n-2s}} \phi$  is a linear combination of  $\frac{\partial U_{\xi,\Lambda}}{\partial \Lambda}$  and  $\frac{\partial U_{\xi,\Lambda}}{\partial \xi_i}$ , i = 1, 2..., n. We will use the functions  $U_{\xi,\Lambda}$  to construct the approximate solutions of the equation (1.5).

We will use the functions  $U_{\xi,\Lambda}$  to construct the approximate solutions of the equation (1.5). We define  $X_{l,m} = \{\lambda lx | x \in Q_m\}$  and arrange it in any way as a sequence  $X_{l,m} = \{X^i\}_{i=1}^{(m+1)^k}$ . Let  $P^i \in B_{\frac{1}{2}}(X^i) = \{X \in \mathbb{R}^n | |X - X^i| < \frac{1}{2}\}, \Lambda_i \in [C_1, C_2]$ , for  $i = 1, 2..., (m+1)^k$ , where  $C_1$ and  $C_2$  are some positive numbers to be defined later( see (3.6)). Let

$$W_m(x) := \sum_{i=1}^{(m+1)^k} U_{P^i,\Lambda_i}(x)$$

to be an approximate solution of the problem (1.5).

**Theorem 1.2.** Under the same conditions of Theorem 1.1, there exists an interger  $l_0 > 0$ , such that for any integer  $l > l_0$ , equation (1.5) has a  $C_{loc}^2$  solution  $u_m$  of the form

$$u_m = W_m + \phi_m$$

where  $m \in \mathbb{N}_+ \cup \{\infty\}, \ |\phi|_{L^{\infty}(\mathbb{R}^n)} \to 0 \ and \max_{i=1,\dots,(m+1)^k} \{|P^i - X^i|\} \to 0 \ as \ l \to \infty.$ 

As a consequence of Theorem 1.2, we have

**Corollary 1.3.** Under the same conditions of Theorem 1.1, the equation (1.4) has infinitely many multi-bump solutions.

Theorem 1.1 follows from Theorem 1.2. So we only need to prove Theorem 1.2. In this article we use l as the pertubation parameter and follow the methods developed in [21, 29]. In the section 2, we carry out the Liapunov-Schmidt reduction. Theorem 1.2 is proved in the section 3. Some useful estimations are presented in Appendix A. The expansions of the functional  $\frac{\partial}{\partial \Lambda_i} I(W_m)$  and  $\frac{\partial}{\partial P^i} I(W_m)$  are shown in Appendix B.

In this article, C denotes a varying constant independent of m.

## 2. FINITE DIMENSIONAL REDUCTION

In this section, we will carry out the Lyapunov-Schmidt reduction in the case of  $m < \infty$ . We define two weighted norms

$$||u||_* = \sup_{y \in \mathbb{R}^n} \left( \gamma(y) \sum_{i=1}^{(m+1)^k} \frac{1}{(1+|y-X^i|)^{\frac{n-2s}{2}+\tau}} \right)^{-1} |u(y)|,$$

and

$$||u||_{**} = \sup_{y \in \mathbb{R}^n} \left( \gamma(y) \sum_{i=1}^{(m+1)^k} \frac{1}{(1+|y-X^i|)^{\frac{m+2s}{2}+\tau}} \right)^{-1} |u(y)|,$$

where

$$\gamma(y) = \min\left\{\min_{i=1,\dots,(m+1)^k} \left(\frac{1+|y-X^i|}{\lambda}\right)^{\tau-s}, 1\right\},\,$$

and  $\tau \in (k, \frac{n-2s}{2})$  is a constant. Consider the following equation

$$(-\Delta)^{s}\phi - \frac{n+2s}{n-2s}K\left(\frac{x}{\lambda}\right)W_{m}^{\frac{4s}{n-2s}}\phi = g \text{ in } \mathbb{R}^{n}.$$
(2.1)

**Lemma 2.1.** Let  $\phi$  be a solution of the equation (2.1), then we have the following estimate

$$\left(\gamma(y)\sum_{h=1}^{(m+1)^{k}} \frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+\tau}}\right)^{-1} |\phi(y)|$$

$$\leq C \|g\|_{**} + C \|\phi\|_{*} \left(\frac{1}{(\lambda l)^{\frac{4s}{n-2s}k}} \frac{\sum_{h=1}^{(m+1)^{k}} \frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+\tau+\theta}}}{\sum_{h=1}^{(m+1)^{k}} \frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+\tau}}}\right), \quad (2.2)$$

where  $\theta > 0$  is a constant and C is independent of m.

*Proof.* We rewrite the equation (2.1) into an integral equation

$$\phi(y) = C_1(n,s) \int_{\mathbb{R}^n} \frac{1}{|y-z|^{n-2s}} \left( \frac{n+2s}{n-2s} K(\frac{z}{\lambda}) W_m^{\frac{4s}{n-2s}}(z) \phi(z) + g(z) \right) dz,$$
(2.3)

where the constant  $C_1(n,s)$  is defined in the Green function of  $(-\Delta)^s$  on  $\mathbb{R}^n$  (cf. [5]).

From Lemma A.5, we get

$$\begin{aligned} \left| \int_{\mathbb{R}^{n}} \frac{1}{|y-z|^{n-2s}} K(\frac{z}{\lambda}) W_{m}^{\frac{4s}{n-2s}}(z) \phi(z) dz \right| \\ &\leq C \|\phi\|_{*} \int_{\mathbb{R}^{n}} \frac{1}{|y-z|^{n-2s}} W_{m}^{\frac{4s}{n-2s}}(z) \gamma(z) \sum_{h=1}^{(m+1)^{k}} \frac{1}{(1+|z-X^{h}|)^{\frac{n-2s}{2}+\tau}} dz \\ &\leq C \|\phi\|_{*} \left( \gamma(y) \sum_{h=1}^{(m+1)^{k}} \frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+\tau+\theta}} \right. \\ &+ \frac{1}{(\lambda l)^{\frac{4s}{n-2s}k}} \gamma(y) \sum_{h=1}^{(m+1)^{k}} \frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+\tau}} \right). \end{aligned}$$
(2.4)

For the second term on the right-hand side of (2.3), we have

$$\int_{\mathbb{R}^n} \frac{1}{|y-z|^{n-2s}} |g(z)| dz \le \|g\|_{**} \int_{\mathbb{R}^n} \frac{1}{|y-z|^{n-2s}} \gamma(z) \sum_{h=1}^{(m+1)^k} \frac{1}{(1+|z-X^h|)^{\frac{n+2s}{2}+\tau}} dz.$$
(2.5)

Using Lemma A.2, we obtain

$$\int_{\mathbb{R}^n} \frac{1}{|y-z|^{n-2s}} \sum_{h=1}^{(m+1)^k} \frac{1}{(1+|z-X^h|)^{\frac{n+2s}{2}+\tau}} dz \le C \sum_{h=1}^{(m+1)^k} \frac{1}{(1+|z-X^h|)^{\frac{n-2s}{2}+\tau}}.$$
 (2.6)

Define  $B_i := B_{\lambda l}(X^i), B_{i,m} := B_{r_0}(X^i)$  with  $r_0 = \max\{\frac{m}{4}, 1\}$  and

$$\Omega_i := \{ z \in \mathbb{R}^n : |z - X^i| = \min_{j=1,\dots,(m+1)^k} |z - X^j| \}.$$

Without loss of generality, we assume  $y \in \Omega_1$ . Make use of Lemma A.3 under different cases, we have

$$\frac{1}{\lambda^{\tau-s}} \sum_{h=1}^{(m+1)^k} \frac{1}{(1+|y-X^h|)^{\frac{n}{2}}} \le C\left(\frac{1+|y-X^1|}{\lambda}\right)^{\tau-s} \sum_{h=1}^{(m+1)^k} \frac{1}{(1+|y-X^h|)^{\frac{n-2s}{2}+\tau}}.$$
 (2.7)

Using Lemma A.2 and (2.7), we have

$$\int_{\mathbb{R}^{n}} \frac{1}{|y-z|^{n-2s}} \min_{i=1,\dots,(m+1)^{k}} \left\{ \left( \frac{1+|z-X^{i}|}{\lambda} \right)^{\tau-s} \right\}^{(m+1)^{k}} \sum_{h=1}^{(m+1)^{k}} \frac{1}{(1+|z-X^{h}|)^{\frac{n+2s}{2}+\tau}} dz \\
\leq \frac{C}{\lambda^{\tau-s}} \int_{\mathbb{R}^{n}} \frac{1}{|y-z|^{n-2s}} \sum_{h=1}^{(m+1)^{k}} \frac{1}{(1+|z-X^{h}|)^{\frac{n}{2}+2s}} dz \\
\leq \frac{C}{\lambda^{\tau-s}} \sum_{h=1}^{(m+1)^{k}} \frac{1}{(1+|y-X^{h}|)^{\frac{n}{2}}} \\
\leq C \left( \frac{1+|y-X^{1}|}{\lambda} \right)^{\tau-s} \sum_{h=1}^{(m+1)^{k}} \frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+\tau}}.$$
(2.8)

From the definition of  $\gamma(y)$ , (2.5), (2.6) and (2.8), we know

$$\int_{\mathbb{R}^n} \frac{1}{|y-z|^{n-2s}} |g(z)| dz \le C ||g||_{**} \gamma(y) \sum_{h=1}^{(m+1)^k} \frac{1}{(1+|y-X^h|)^{\frac{n-2s}{2}+\tau}}.$$
(2.9)

Now (2.2) follows from (2.3), (2.4) and (2.9).

Consider the following problem

$$\begin{cases} (-\Delta)^{s}\phi - \frac{n+2s}{n-2s}K(\frac{x}{\lambda})W_{m}^{\frac{4s}{n-2s}}\phi = g + \sum_{i=1}^{(m+1)^{k}}\sum_{j=1}^{n+1}c_{ij}^{(m)}U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}}Z_{i,j}, \\ \int_{\mathbb{R}^{n}}U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}}Z_{i,j}\phi dx = 0, \quad \phi \in \dot{H}^{s}(\mathbb{R}^{n}), \quad i = 1, \dots, (m+1)^{k}, \quad j = 1, \dots, n+1, \end{cases}$$

$$(2.10)$$

where  $Z_{i,j} = \frac{\partial U_{P^i,\Lambda_i}}{\partial P_j^i}$  for  $j = 1, \dots, n$  and  $Z_{i,n+1} = \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i}$ .

**Lemma 2.2.** Assume  $\phi$  solves the problem (2.10), there is exists  $l_0 > 0$ , such that for all  $l > l_0$ , we have  $\|\phi\|_* \leq C \|g\|_{**}$ , where C is independent of m.

*Proof.* If this lemma is not right, then there would be sequences  $\{g_l\}_{l=1}^{\infty}$  and  $\{\phi_l\}_{l=1}^{\infty}$  satisfying (2.10) with  $\|\phi_l\|_* = 1$  and  $\|g_l\|_{**} \to 0$  as  $l \to +\infty$ . For notation simplicity, we suppress l in the argument below.

First, we give an estimate of the parameters  $c_{ij}^{(m)}$ . Multiplying (2.10) with  $Z_{r,t}$  and integrating on both sides, we get

$$-\frac{n+2s}{n-2s}\int_{\mathbb{R}^n}K(\frac{x}{\lambda})W_m^{\frac{4s}{n-2s}}\phi Z_{r,t}dx = \int_{\mathbb{R}^n}gZ_{r,t}dx + \sum_{i=1}^{(m+1)^k}\sum_{j=1}^{n+1}c_{ij}^{(m)}\int_{\mathbb{R}^n}U_{P^i,\Lambda_i}^{\frac{4s}{n-2s}}Z_{i,j}Z_{r,t}dx.$$
 (2.11)

For the first term on the right hand side of (2.11), using Lemma A.2, we have

$$\begin{aligned} \left| \int_{\mathbb{R}^{n}} g Z_{r,t} dx \right| &\leq C \|g\|_{**} \int_{\mathbb{R}^{n}} \frac{1}{(1+|x-X^{r}|)^{n-2s}} \gamma(x) \sum_{j=1}^{(m+1)^{k}} \frac{1}{(1+|x-X^{j}|)^{\frac{n+2s}{2}+\tau}} dx \\ &\leq \frac{C}{\lambda^{\tau-s}} \|g\|_{**} \int_{\mathbb{R}^{n}} \frac{1}{(1+|x-X^{r}|)^{n-2s}} \sum_{j=1}^{(m+1)^{k}} \frac{1}{(1+|x-X^{j}|)^{\frac{n}{2}+2s}} dx \\ &\leq C \frac{\|g\|_{**}}{\lambda^{\tau-s}} \left( \int_{\mathbb{R}^{n}} \frac{1}{(1+|x-X^{r}|)^{n+\frac{n}{2}}} dx + \sum_{j\neq r} \frac{1}{|X^{j}-X^{r}|^{\frac{n}{2}}} \right) \\ &\leq C \frac{\|g\|_{**}}{\lambda^{\tau-s}}, \end{aligned}$$

where we have used the fact that

$$\sum_{j \neq r} \frac{1}{|X^j - X^r|^{\frac{n}{2}}} \text{ converges for } \frac{n}{2} > k.$$
(2.12)

Since the left hand side of the equation (2.11) is estimated in Lemma A.6, we have

$$\sum_{i=1}^{(m+1)^k} \sum_{j=1}^{n+1} c_{ij}^{(m)} \int_{\mathbb{R}^n} U_{P^i,\Lambda_i}^{\frac{4s}{n-2s}} Z_{i,j} Z_{r,t} dx = \frac{1}{\lambda^{\tau-s}} O\left( \|g\|_{**} + \frac{\|\phi\|_*}{(\lambda l)^{\frac{n}{2}}} \right).$$

As we know  $\int_{\mathbb{R}^n} U_{P^i,\Lambda_i}^{\frac{4s}{n-2s}} Z_{i,j} Z_{i,t} dx = C \delta_{jt}$  and  $\int_{\mathbb{R}^n} |U_{P^i,\Lambda_i}^{\frac{4s}{n-2s}} Z_{i,j} Z_{r,t}| dx \leq \frac{C}{|X^i - X^r|^{n-2s}}$  for  $i \neq r$ , we obtain

$$\max_{i,j}\{|c_{ij}^{(m)}|\} = \frac{1}{\lambda^{\tau-s}}O\left(\|g\|_{**} + \frac{\|\phi\|_{*}}{(\lambda l)^{\frac{n}{2}}}\right).$$

An argument similar to the one used in (2.7) yields

$$\begin{aligned} \left| \sum_{i=1}^{(m+1)^{k}} \sum_{j=1}^{n+1} c_{ij}^{(m)} U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}} Z_{i,j} \right| &\leq \frac{C}{\lambda^{\tau-s}} \left( \|g\|_{**} + \frac{\|\phi\|_{*}}{(\lambda l)^{\frac{n}{2}}} \right) \sum_{i=1}^{(m+1)^{k}} \frac{1}{(1+|y-X^{i}|)^{n+2s}} \\ &\leq C \left( \|g\|_{**} + \frac{\|\phi\|_{*}}{(\lambda l)^{\frac{n}{2}}} \right) \gamma(y) \sum_{i=1}^{(m+1)^{k}} \frac{1}{(1+|y-X^{i}|)^{\frac{n+2s}{2}+\tau}} \end{aligned}$$

From the definition of the norm  $\|\cdot\|_{**}$ , we have

$$\|\sum_{i=1}^{(m+1)^{k}}\sum_{j=1}^{n+1}c_{ij}^{(m)}U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}}Z_{i,j}\|_{**} \le C\left(\|g\|_{**} + \frac{\|\phi\|_{*}}{(\lambda l)^{\frac{n}{2}}}\right)$$

Applying Lemma 2.1 to the first equation of the system (2.10), one get

$$\begin{split} & \left(\gamma(y)\sum_{h}\frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+\tau}}\right)^{-1}|\phi(y)| \\ & \leq C\left(\|g\|_{**}+\|\sum_{i=1}^{(m+1)^{k}}\sum_{j=1}^{n+1}c_{ij}^{(m)}U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}}Z_{i,j}\|_{**}+\left(\frac{1}{(\lambda l)^{\frac{4s}{n-2s}k}}+\frac{\sum_{h}\frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+\tau+\theta}}}{\sum_{h}\frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+\tau}}}\right)\|\phi\|_{*}\right) \\ & \leq C\left(\|g\|_{**}+\frac{1}{(\lambda l)^{\frac{4s}{n-2s}k}}+\frac{\sum_{h}\frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+\tau+\theta}}}{\sum_{h}\frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+\tau}}}\right). \end{split}$$

As a result, there exist a number  $i_0 \in \mathbb{N}$  and a large constant R > 0, such that

$$1 = \|\phi\|_{*} = \sup_{B_{R}(X^{i_{0}})} (\gamma(y) \sum_{h=1}^{(m+1)^{k}} \frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+\tau}})^{-1} |\phi(y)|.$$
(2.13)

Hence there is a constant  $c_0 > 0$  such that  $|\lambda^{\tau-s}\phi|_{L^{\infty}(B_R(X^{i_0}))} \ge c_0$ .

Applying Lemma A.8 to the equation (2.10), we know  $\lambda^{\tau-s}\phi$  is equi-continuous. Also  $\lambda^{\tau-s}|\phi(\cdot)|$  is uniformly bounded. In fact, we assume that  $y \in \Omega_1$  with no loss of generality. From the fact (2.12), we have

$$\lambda^{\tau-s}|\phi(y)| \le \|\phi\|_* \sum_h \frac{1}{(1+|y-X^h|)^{\frac{n}{2}}} \le C + \sum_{h\neq 1} \frac{1}{|X^h - X^1|^{\frac{n}{2}}} \le C.$$

Then the Arzelà-Ascoli Theorem yields that there is a function  $\tilde{\phi}$ , such that  $\lambda^{\tau-s}\phi(\cdot + P^{i_0})$  convergent to  $\tilde{\phi}$  uniformly on compact sets. Then

$$|\phi|_{L^{\infty}(B_{R+1}(0))} \ge c_0. \tag{2.14}$$

Using a similar argument as in [12, Lemma 7.3], we know  $\tilde{\phi}$  satisfies

$$\begin{cases} (-\Delta)^s \tilde{\phi} - \frac{n+2s}{n-2s} U_{0,\Lambda_{i_0}}^{\frac{4s}{n-2s}} \tilde{\phi} = 0, \\ \int_{\mathbb{R}^n} U_{0,\Lambda_{i_0}}^{\frac{4s}{n-2s}} \frac{\partial U_{0,\Lambda_{i_0}}}{\partial \Lambda_{i_0}} \tilde{\phi} = 0, \quad \int_{\mathbb{R}^n} U_{0,\Lambda_{i_0}}^{\frac{4s}{n-2s}} \frac{\partial U_{0,\Lambda_{i_0}}}{\partial P_j^{i_0}} \tilde{\phi} = 0, \quad j = 1, \dots, n. \end{cases}$$

Then  $\tilde{\phi} = 0$  by nondegeneracy, which is contradict to (2.14). Hence the solution  $\phi$  of the equation (2.10) satisfies  $\|\phi\|_* \leq C \|g\|_{**}$ .

Combining Lemma 2.2, Lemma A.8 and the argument of [14, Proposition 4.1] (cf. [24, Proposition 2.2]), we have

**Proposition 2.3.** For any g satisfying  $||g||_{**} < +\infty$ , (2.10) has a unique solution  $\phi = L_m(g) \in \dot{H}^s(\mathbb{R}^n) \cap C^{0,\alpha}(\mathbb{R}^n)$  with  $\alpha = \min\{2s,1\}$ , such that  $||L_m(g)||_* \leq C||g||_{**}$ . The constant  $c_{ij}^{(m)}$  satisfies  $|c_{ij}^{(m)}| \leq \frac{C}{\lambda^{\tau-s}} ||g||_{**}$ .

Since we are interested in the solution of the form  $W_m + \phi_m$  of the equation (1.5), we now consider the following problem

$$\begin{cases} (-\Delta)^{s}\phi - \frac{n+2s}{n-2s}K(\frac{x}{\lambda})W_{m}^{\frac{4s}{n-2s}}\phi = N(\phi) + l_{m} + \sum_{i=1}^{(m+1)^{k}}\sum_{j=1}^{n+1}c_{ij}^{(m)}U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}}Z_{i,j}, \\ \int_{\mathbb{R}^{n}}U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}}Z_{i,j}\phi dx = 0, \quad \phi \in \dot{H}^{s}(\mathbb{R}^{n}), \quad i = 1,\ldots,(m+1)^{k}, \quad j = 1,\ldots,n+1, \end{cases}$$
(2.15)

where

$$N(\phi) = K(\frac{x}{\lambda}) \left( (W_m + \phi)_+^{\frac{n+2s}{n-2s}} - W_m^{\frac{n+2s}{n-2s}} - \frac{n+2s}{n-2s} W_m^{\frac{4s}{n-2s}} \phi \right)$$

and

$$l_m = K(\frac{x}{\lambda}) W_m^{\frac{n+2s}{n-2s}} - \sum_{i=1}^{(m+1)^k} U_{P^i,\Lambda_i}^{\frac{n+2s}{n-2s}}.$$

**Lemma 2.4.** For the terms  $N(\phi)$  and  $l_m$  defined above, we have the following estimates

$$\|N(\phi)\|_{**} \le C \|\phi\|_{*}^{\min\{2,2^{*}(s)-1\}},$$
$$\|l_{m}\|_{**} \le \frac{C}{\lambda^{\frac{n+2s}{2}-\tau}}.$$

*Proof.* The proof of the first estimation is rather standard(cf. [29, Lemma 2.4] for ideas). We only prove the second estimate.

Without loss of generality, we assume  $x \in \Omega_1$ . Then

$$l_{m} = K(\frac{x}{\lambda})W_{m}^{\frac{n+2s}{n-2s}} - U_{P^{1},\Lambda_{1}}^{\frac{n+2s}{n-2s}} - \sum_{h\neq 1} U_{P^{h},\Lambda_{h}}^{\frac{n+2s}{n-2s}}$$
$$= \left(K(\frac{x}{\lambda}) - 1\right)U_{P^{1},\Lambda_{1}}^{\frac{n+2s}{n-2s}} + O\left(\left(\sum_{h\neq 1} U_{P^{h},\Lambda_{h}}\right)^{\frac{n+2s}{n-2s}} + U_{P^{1},\Lambda_{1}}^{\frac{4s}{n-2s}}\sum_{h\neq 1} U_{P^{h},\Lambda_{h}}\right). \quad (2.16)$$

The two error terms in (2.16) can be estimated by using Lemma A.3 under different cases. Case 1:  $x \in \Omega_1 \cap B_1^c \cap B_{1,m}$ , we have  $\gamma(x) = 1$ . Using Lemma A.3, we have

$$\left(\sum_{h\neq 1} U_{P^{h},\Lambda_{h}}\right)^{\frac{n+2s}{n-2s}} \leq \frac{C}{(\lambda l)^{\frac{n+2s}{n-2s}k}} \frac{1}{(1+|x-X^{1}|)^{n+2s-\frac{n+2s}{n-2s}k}} \\ \leq \frac{C}{(\lambda l)^{\frac{n+2s}{2}-\tau+k}} \frac{1}{(1+|x-X^{1}|)^{\frac{n+2s}{2}+\tau-k}} \\ \leq \frac{C}{(\lambda l)^{\frac{n+2s}{2}-\tau}} \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\frac{n+2s}{2}+\tau}}.$$

**Case 2:**  $x \in \Omega_1 \cap B_1$ , it holds that  $|x - X^i| \ge \frac{1}{2}|X^i - X^1| \ge \frac{1}{2}\lambda l$  for  $i \ne 1$ . From Lemma A.3,

$$\left(\sum_{h\neq 1} U_{P^{h},\Lambda_{h}}\right)^{\frac{n+2s}{n-2s}} \leq \begin{cases} \frac{C}{(\lambda l)^{\frac{n+2s}{2}-\tau}} \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\frac{n+2s}{2}+\tau}}, & \text{if } x \in \Omega_{1} \cap B_{1} \cap B_{\lambda}^{c}(X_{1}), \\ C\left(\frac{1+|x-X^{1}|}{\lambda}\right)^{\tau-s} \frac{\lambda^{\tau-s}}{(\lambda l)^{\frac{n}{2}}} \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\frac{n+2s}{2}+\tau}}, & \text{if } x \in \Omega_{1} \cap B_{1} \cap B_{\lambda}(X^{1}). \end{cases}$$

**Case 3:**  $x \in \Omega_1 \cap B_{1,m}^c$ , we can get

$$\left(\sum_{h\neq 1} U_{P^h,\Lambda_h}\right)^{\frac{n+2s}{n-2s}} \le \frac{Cm^{\frac{n+2s}{n-2s}k}}{(1+|x-X^1|)^{n+2s}} \le \frac{Cm^{\frac{4s}{n-2s}k}}{(1+|x-X^1|)^{2s}} \frac{(1+C[\frac{m}{2}])^k}{(1+|x-X^1|)^n}.$$

Following the proof of Lemma A.3, we have

$$\begin{split} \sum_{h} \frac{1}{(1+|x-X^{h}|)^{n}} &\geq \frac{C}{(1+|x-X^{1}|)^{n}} \left(1+2^{-k} \int_{[0,[\frac{m}{2}]+1]^{k} \setminus [0,1]^{k}} \frac{1}{(1+\frac{\lambda l}{1+|x-X^{1}|}|z|)^{n}} dz \right) \\ &\geq \frac{(1+C[\frac{m}{2}])^{k}}{(1+|x-X^{1}|)^{n}}. \end{split}$$

Since in the domain  $\Omega_1 \cap B_{1,m}^c$ , we can get  $|x - X^h| \ge \frac{1}{2}\lambda l$  for  $h = 1, \dots, (m+1)^k$ , then

$$\left(\sum_{h\neq 1} U_{P^h,\Lambda_h}\right)^{\frac{n+2s}{n-2s}} \le \frac{C}{(\lambda l)^{2s}} \sum_h \frac{1}{(1+|x-X^h|)^n} \le \frac{C}{(\lambda l)^{\frac{n+2s}{2}-\tau}} \sum_h \frac{1}{(1+|x-X^h|)^{\frac{n+2s}{2}+\tau}}.$$

Combining these three cases above, we have  $\|(\sum_{h\neq 1} U_{P^h,\Lambda_h})^{\frac{n+2s}{n-2s}}\|_{**} \leq \frac{C}{(\lambda l)^{\frac{n+2s}{2}-\tau}}$ . By the same procedure, we can also get the estimation  $\|U_{P^1,\Lambda_1}^{\frac{4s}{n-2s}}\sum_{h\neq 1}U_{P^h,\Lambda_h}\|_{**} \leq \frac{C}{(\lambda l)^{\frac{n+2s}{2}-\tau}}$ .

At last, we estimate the first term in (2.16).

In the case of  $|x - X^1| \ge \lambda$ , we have  $\gamma(x) = 1$ . Then

$$\left| K(\frac{x}{\lambda}) - 1 \right| U_{P^1,\Lambda_1}^{\frac{n+2s}{n-2s}} \le \frac{C}{(1+|x-X^1|)^{n+2s}} \le \frac{C}{\lambda^{\frac{n+2s}{2}-\tau}} \gamma(x) \sum_{h=1}^{(m+1)^k} \frac{1}{(1+|x-X^h|)^{\frac{n+2s}{2}+\tau}}.$$
 (2.17)

In the case of  $|x - X^1| < \lambda$ , it holds  $\frac{1 + |x - X^1|}{\lambda} \leq C$ . The condition (H<sub>3</sub>) yields

$$\left| K(\frac{x}{\lambda}) - 1 \right| U_{P^{1},\Lambda_{1}}^{\frac{n+2s}{n-2s}} \leq C \frac{|x - X^{1}|^{\beta}}{\lambda^{\beta}} \frac{1}{(1 + |x - X^{1}|)^{n+2s}} \\ \leq \frac{C}{\lambda^{\frac{n+2s}{2}-\tau}} \left( \frac{1 + |x - X^{1}|}{\lambda} \right)^{\tau-s} \sum_{h=1}^{(m+1)^{k}} \frac{1}{(1 + |x - X^{h}|)^{\frac{n+2s}{2}+\tau}} \\ \leq \frac{C}{\lambda^{\frac{n+2s}{2}-\tau}} \gamma(y) \sum_{h=1}^{(m+1)^{k}} \frac{1}{(1 + |x - X^{h}|)^{\frac{n+2s}{2}+\tau}}.$$
(2.18)

Summarizing (2.17) and (2.18), we have

$$\|\left(K(\frac{x}{\lambda})-1\right)U_{P^1,\Lambda_1}^{\frac{n+2s}{n-2s}}\|_{**} \leq \frac{C}{\lambda^{\frac{n+2s}{2}-\tau}}$$

Hence this lemma follows.

**Proposition 2.5.** For  $\lambda$  large enough, the problem (2.15) has a unique solution  $\phi_m \in \dot{H}^s(\mathbb{R}^n) \cap C^{0,\alpha}(\mathbb{R}^n)$  with  $\alpha = \min\{2s,1\}$ , such that  $\|\phi_m\|_* \leq \frac{C}{\lambda^{\frac{n+2s}{2}-\tau}}$ . The constants  $c_{ij}^{(m)}$  satisfy  $|c_{ij}^{(m)}| \leq \frac{C}{\lambda^{\frac{n+2s}{2}-\tau}}$ .  $C\lambda^{-\frac{n}{2}}$ .

Proof. We define

$$E = \left\{ \varphi \in \dot{H}(\mathbb{R}^n) \cap C(\mathbb{R}^n) : \|\varphi\|_* \le \frac{1}{\lambda^{\frac{n+2s}{2}-\tau-\epsilon_1}}, \quad \int_{\mathbb{R}^n} U_{P^i,\Lambda_i}^{\frac{4s}{n-2s}} Z_{i,j}\varphi = 0, \quad \substack{i=1,\ldots,(m+1)^k, \\ j=1,\ldots,n+1} \right\},$$

where  $\epsilon_1 = \min\{\frac{1}{4}, \frac{2s}{n+2s}\}(\frac{n+2s}{2} - \tau)$ . Notice that  $(E, \|\cdot\|_*)$  is a metric space. In order to use the contraction map theorem, we define  $A\varphi := L_m(N(\varphi) + l_m)$ , where  $L_m$  is an operator defined in Proposition 2.3.

Firstly, we show that A maps E into itself for  $\lambda$  large. Combining Proposition 2.3 and Lemma 2.4, we have  $\forall \varphi \in E$ ,

$$||A\varphi||_* \le C(||N(\varphi)||_{**} + ||l_m||_{**}) \le C(||\varphi||_*^{\min\{2,2^*(s)-1\}} + ||l_m||_{**}) \le \frac{1}{\lambda^{\frac{n+2s}{2}-\tau-\epsilon_1}}.$$

Secondly, we prove A is an contraction map for  $\lambda$  large.

Choose  $\varphi_1, \varphi_2 \in E$  with  $\varphi_1 \neq \varphi_2$ . If  $N \geq 6s$ , we have

$$|N(\varphi_{1}) - N(\varphi_{2})| = |N'(t\varphi_{1} + (1 - t)\varphi_{2})(\varphi_{1} - \varphi_{2})|$$

$$\leq C(|\varphi_{1}|^{\frac{4s}{n-2s}}| + |\varphi_{2}|^{\frac{4s}{n-2s}}|)(|\varphi_{1} - \varphi_{2}|)$$

$$\leq C(||\varphi_{1}||^{\frac{4s}{n-2s}} + ||\varphi_{2}||^{\frac{4s}{n-2s}})||\varphi_{1} - \varphi_{2}||_{*} \left(\gamma(x)\sum_{h=1}^{(m+1)^{k}} \frac{1}{(1 + |x - X^{h}|)^{\frac{n-2s}{2} + \tau}}\right)^{\frac{n+2s}{n-2s}}$$

$$\leq C(||\varphi_{1}||^{\frac{4s}{n-2s}} + ||\varphi_{2}||^{\frac{4s}{n-2s}})||\varphi_{1} - \varphi_{2}||_{*} \left(\gamma(x)\sum_{h=1}^{(m+1)^{k}} \frac{1}{(1 + |x - X^{h}|)^{\frac{n+2s}{2} + \tau}}\right).$$

We remind that in the last inequality, we have used the Hölder inequality. Hence  $||N(\varphi_1) - N(\varphi_2)||_{**} \leq C(||\varphi_1||_*^{\frac{4s}{n-2s}} + ||\varphi_2||_*^{\frac{4s}{n-2s}})||\varphi_1 - \varphi_2||_*.$ 

In the case of  $N \leq 6s$ , we also have  $||N(\varphi_1) - N(\varphi_2)||_{**} \leq C(||\varphi_1||_*^{\min\{\frac{4s}{n-2s},1\}} + ||\varphi_2||_*^{\min\{\frac{4s}{n-2s},1\}})||\varphi_1 - \varphi_2||_*$  by a similar argument.

Then there hold

$$\begin{aligned} \|A\varphi_{1} - A\varphi_{2}\|_{*} &\leq C \|N(\varphi_{1}) - N(\varphi_{2})\|_{**} \\ &\leq C(\|\varphi_{1}\|_{*}^{\min\{\frac{4s}{n-2s},1\}} + \|\varphi_{2}\|_{*}^{\min\{\frac{4s}{n-2s},1\}}) \|\varphi_{1} - \varphi_{2}\|_{*} \\ &\leq \frac{C}{\lambda^{(\frac{n+2s}{2} - \tau - \epsilon_{1})\min\{\frac{4s}{n-2s},1\}}} \|\varphi_{1} - \varphi_{2}\|_{*}. \end{aligned}$$

For  $\lambda$  large enough, we get  $||A\varphi_1 - A\varphi_2||_* \leq \frac{1}{2} ||\varphi_1 - \varphi_2||_*$ .

Using the contracting map theorem, we know that there is a unique  $\phi_m \in E$ , such that  $A(\phi_m) = \phi_m$ , *i.e.*  $\phi_m$  is a unique solution of (2.15) in E. From Proposition 2.3 and Lemma 2.4, we know  $\phi_m \in \dot{H}^s(\mathbb{R}^n) \cap C^{0,\alpha}(\mathbb{R}^n)$  satisfying  $\|\phi_m\|_* = \|A\phi_m\|_* \leq \frac{C}{\lambda^{\frac{n+2s}{2}-\tau}}$  and  $|c_{ij}^{(m)}| \leq C\lambda^{-\frac{n}{2}}$ , since  $(\frac{n+2s}{2}-\tau-\epsilon_1)\min\{2,2^*(s)-1\} > \frac{n+2s}{2}-\tau$ .

### 3. Proof of the main theorem

Let  $\Lambda := (\Lambda_1, \ldots, \Lambda_{(m+1)^k}) \in \mathbb{R}^{(m+1)^k}_+$  and  $P := (P^1, \ldots, P^{(m+1)^k}) \in \mathbb{R}^{n \times (m+1)^k}$ , in which  $P^i = (P_1^i, \ldots, P_n^i) \in \mathbb{R}^n$  for  $i = 1, 2, \ldots, (m+1)^k$ . We define  $J(P, \Lambda) = I(W_m + \phi_m)$ , where  $\phi_m$  is a unique small solution obtained by Proposition 2.5. A standard argument shows that from a critical point of J, we can get a critical point of I of the form  $W_m + \phi_m$  (for example, cf. [14, Lemma 6.1] for ideas).

**Proposition 3.1.** For  $\lambda$  large, we have the following expansions

$$\frac{\partial J}{\partial P_j^i}(P,\Lambda) = -\frac{c_3 a_j}{\Lambda_i^{\beta-2} \lambda^{\beta}} (P_j^i - X_j^i) + O\left(\frac{|P^i - X^i|^2}{\lambda^{\beta}}\right) + o(\lambda^{-\beta}), \tag{3.1}$$

and

$$\frac{\partial J}{\partial \Lambda_i}(P,\Lambda) = -\frac{c_1}{\Lambda_i^{\beta+1}\lambda^{\beta}} + \sum_{h\neq i} \frac{c_2}{\Lambda_i(\Lambda_i\Lambda_h)^{\frac{n-2s}{2}} |X^i - X^h|^{n-2s}} + O\left(\frac{|P^i - X^i|^{\min\{2,\beta-1\}}}{\lambda^{\beta}}\right) + o(\lambda^{-\beta}),$$
(3.2)

where  $i = 1, ..., (m+1)^k$  and j = 1, ..., n and the constant  $c_1, c_2, c_3$  are positive. Proof. A simple calculation yields

$$\begin{aligned} \frac{\partial J}{\partial P_{j}^{i}}(P,\Lambda) &= \langle I'(W_{m} + \phi_{m}), \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial P_{j}^{i}} + \frac{\partial \phi_{m}}{\partial P_{j}^{i}} \rangle \\ &= \frac{\partial I}{\partial P_{j}^{i}}(W_{m}) + \int_{\mathbb{R}^{n}} K(\frac{x}{\lambda})(W_{m}^{\frac{n+2s}{n-2s}} - (W_{m} + \phi_{m})_{+}^{\frac{n+2s}{n-2s}}) \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial P_{j}^{i}} \\ &+ \sum_{t=1}^{(m+1)^{k}} \sum_{h=1}^{n+1} c_{th}^{(m)} \int_{\mathbb{R}^{n}} U_{P^{t},\Lambda_{t}}^{\frac{4s}{n-2s}} Z_{t,h} \frac{\partial \phi_{m}}{\partial P_{j}^{i}}. \end{aligned}$$

The functional  $\frac{\partial I}{\partial P_j^i}(W_m)$  is expanded in the Proposition B.6. So we only need to estimate the last two terms in the equility above.

From Lemma A.4, we see that for  $\lambda$  large enough,  $\{x : W_m \leq -\phi_m\} \subset \{x : \frac{1}{2}W_m \leq |\phi_m|\} \subset \cup_h(\Omega_h \cap B_h^c)$ . Then we have

$$\begin{split} &\int_{\mathbb{R}^n} K(\frac{x}{\lambda})((W_m + \phi_m)_+^{\frac{n+2s}{n-2s}} - W_m^{\frac{n+2s}{n-2s}}) \frac{\partial U_{P^i,\Lambda_i}}{\partial P_j^i} dx \\ &= \int_{\mathbb{R}^n} K(\frac{x}{\lambda})((W_m + \phi_m)^{\frac{n+2s}{n-2s}} - W_m^{\frac{n+2s}{n-2s}}) \frac{\partial U_{P^i,\Lambda_i}}{\partial P_j^i} dx - \int_{-\phi_m \ge W_m} K(\frac{x}{\lambda})(W_m + \phi_m)^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial P_j^i} dx \\ &= \frac{n+2s}{n-2s} \int_{\mathbb{R}^n} K(\frac{x}{\lambda}) W_m^{\frac{4s}{n-2s}} \phi_m \frac{\partial U_{P^i,\Lambda_i}}{\partial P_j^i} dx + O\left(\int_{|\phi_m| \ge \frac{1}{2}W_m} |\phi_m|^{\frac{n+2s}{n-2s}} U_{P^i,\Lambda_i} dx \right. \\ &+ \int_{|\phi_m| < \frac{1}{2}W_m} W_m^{\frac{6s-n}{n-2s}} \phi_m^2 \frac{\partial U_{P^i,\Lambda_i}}{\partial P_j^i} dx \right) \\ &= \frac{n+2s}{n-2s} \int_{\mathbb{R}^n} K(\frac{x}{\lambda}) W_m^{\frac{4s}{n-2s}} \phi_m \frac{\partial U_{P^i,\Lambda_i}}{\partial P_j^i} dx + O\left(\int_{\cup_h (\Omega_h \cap B_h^c)} |\phi_m|^{\frac{n+2s}{n-2s}} U_{P^i,\Lambda_i} dx \right. \\ &+ \int_{\mathbb{R}^n} W_m^{\frac{6s-n}{n-2s}} \phi_m^2 U_{P^i,\Lambda_i} dx \right). \end{split}$$

Using Proposition 2.5, Lemma A.6 and Lemma A.7, we have

$$\int_{\mathbb{R}^n} K(\frac{x}{\lambda}) \left( (W_m + \phi_m)_+^{\frac{n+2s}{n-2s}} - W_m^{\frac{n+2s}{n-2s}} \right) \frac{\partial U_{P^i,\Lambda_i}}{\partial P_j^i} dx = O\left(\lambda^{-n}\right) = o\left(\lambda^{-\beta}\right).$$

By using the orthogonal condition of (2.15) and Lemma A.1, we have

$$\begin{vmatrix} (m+1)^{k} \sum_{t=1}^{n+1} c_{th} \int_{\mathbb{R}^{n}} U_{P^{t},\Lambda_{t}}^{\frac{4s}{n-2s}} Z_{t,h} \frac{\partial \phi_{m}}{\partial P_{j}^{i}} dx \end{vmatrix} = \left| \sum_{h=1}^{n+1} c_{ih} \int_{\mathbb{R}^{n}} \frac{\partial}{\partial P_{j}^{i}} (U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}} Z_{i,h}) \phi_{m} dx \right| \\ \leq \frac{C}{\lambda^{\frac{n}{2}}} \frac{\|\phi_{m}\|_{*}}{\lambda^{\tau-s}} \int_{\mathbb{R}^{n}} \frac{1}{(1+|x-X^{i}|)^{n+2s}} \sum_{r=1}^{(m+1)^{k}} \frac{1}{(1+|x-X^{r}|)^{\frac{n}{2}}} dx \leq C\lambda^{-n}.$$

Hence we can get (3.1). The estimation (3.2) can be derived by the same procedure along with Proposition B.5.

**Remark 3.2.** From Proposition 3.1, we know that there exist bounded functions  $\Xi_j^i = \Xi_j^i(P, \Lambda, \lambda)$ and  $\Theta_j^i = \Theta_j^i(P, \Lambda, \lambda), i = 1, \dots, (m+1)^k; j = 1, \dots, n+1$  satisfying

 $|\Xi_j^i| \leq C$ , where C is constant independent of m

and

 $|\Theta_j^i| \leq C_{\lambda}$ , where  $C_{\lambda}$  is a constant only depend on  $\lambda$ , and  $C_{\lambda} \to 0$  as  $\lambda \to \infty$ such that

$$\frac{\partial J}{\partial P_j^i}(P,\Lambda) = -\frac{c_3 a_j}{\Lambda_i^{\beta-2} \lambda^{\beta}} (P_j^i - X_j^i) + \frac{|P^i - X^i|^2}{\lambda^{\beta}} \Xi_j^i + \lambda^{-\beta} \Theta_j^i,$$

and

$$\frac{\partial J}{\partial \Lambda_i}(P,\Lambda) = -\frac{c_1}{\Lambda_i^{\beta+1}\lambda^{\beta}} + \sum_{h \neq i} \frac{c_2}{\Lambda_i(\Lambda_i\Lambda_h)^{\frac{n-2s}{2}} |X^i - X^h|^{n-2s}} - \frac{|P^i - X^i|^{\min\{2,\beta-1\}}}{\lambda^{\beta}} \Xi_{n+1}^i - \lambda^{-\beta} \Theta_{n+1}^i - \frac{|P^i - X^i|^{\min\{2,\beta-1\}}}{\lambda^{\beta}} = \frac{|P^i - X^i|^$$

Proof of Theorem 1.2. Firstly we look for the solution of (1.5) of the form  $W_m + \phi_m$ ,  $m < \infty$ . It is equivalent to solving the system  $\begin{cases}
\frac{\partial J}{\partial P_j^i}(P,\Lambda) = 0, \\ i = 1, 2, \dots, (m+1)^k; \quad j = 1, 2, \dots, n, \\
\frac{\partial J}{\partial \Lambda_i}(P,\Lambda) = 0,
\end{cases}$ 

that is

$$\begin{cases} \frac{c_{3}a_{j}}{\Lambda_{i}^{\beta-2}\lambda^{\beta}}(P_{j}^{i}-X_{j}^{i}) = \frac{|P^{i}-X^{i}|^{2}}{\lambda^{\beta}}\Xi_{j}^{i} + \lambda^{-\beta}\Theta_{j}^{i};\\ -\frac{c_{1}}{\Lambda_{i}^{\beta+1}\lambda^{\beta}} + \sum_{h\neq i}\frac{c_{2}}{\Lambda_{i}(\Lambda_{i}\Lambda_{h})^{\frac{n-2s}{2}}|X^{i}-X^{h}|^{n-2s}} = \frac{|P^{i}-X^{i}|^{\min\{2,\beta-1\}}}{\lambda^{\beta}}\Xi_{n+1}^{i} + \lambda^{-\beta}\Theta_{n+1}^{i}. \end{cases}$$
(3.3)

To simplify the equations (3.3), we denote  $d_j = \Lambda_j^{-\frac{n-2s}{2}}$  and  $A_{ih} = \begin{cases} 0, & \text{if } i = h, \\ \frac{(\lambda l)^{n-2s}}{|X^i - X^h|^{n-2s}}, & \text{if } i \neq h. \end{cases}$ The equations (3.3) can be written as

$$\begin{cases} P_{j}^{i} - X_{j}^{i} = \frac{\Lambda_{i}^{\beta-2}\Xi_{j}^{i}}{c_{3}a_{j}}|P^{i} - X^{i}|^{2} + \frac{\Lambda_{i}^{\beta-2}}{c_{3}a_{j}}\Theta_{j}^{i}, \\ c_{2}\sum_{h\neq i}A_{ih}d_{h} - c_{1}d_{i}^{\frac{2\beta}{n-2s}-1} = \Lambda_{i}^{\frac{n-2s}{2}+1}\Xi_{n+1}^{i}|P^{i} - X^{i}|^{\min\{2,\beta-1\}} + \Lambda_{i}^{\frac{n-2s}{2}+1}\Theta_{n+1}^{i}, \end{cases}$$
(3.4)

where  $i = 1, 2, ..., (m+1)^k$  and j = 1, 2, ..., n.

Define a function  $F(z) := \frac{c_2}{2} \sum_{h \neq i} A_{ih} z_i z_h - \frac{(n-2s)c_1}{2\beta} \sum_h z_h^{\frac{2\beta}{n-2s}}$ , where  $z = (z_1, z_2, \dots, z_{(m+1)^k}) \in \mathbb{R}^{(m+1)^k}$ . Obviously, F(z) has a maximum point  $b = (b_1, b_2, \dots, b_{(m+1)^k}) \in \mathbb{R}^{(m+1)^k}_+$ . It holds that

$$c_2 \sum_{h \neq i} A_{ih} b_h - c_1 b_i^{\frac{2\beta}{n-2s}-1} = 0, \quad i = 1, \dots, (m+1)^k.$$
(3.5)

**Claim:** Each component  $b_i$  of b satisfies  $0 < C'_1 \le b_i \le C'_2$  for some constant  $C'_1$  and  $C'_2$ . Suppose that  $b_1 \le b_i \le b_2$ . Using the definition of  $A_{ih}$ , we know  $\sum_{h \ne i} A_{ih}$  is bounded. From (3.5), we can get

$$c_1 b_2^{\frac{2\beta}{n-2s}-1} = c_2 \sum_{h \neq 2} A_{2h} b_h \le C_3 b_2,$$

which tell us  $b_2$  is bounded from above.

Using (3.5) again, we have

$$c_1 b_1^{\frac{2\beta}{n-2s}-1} = c_2 \sum_{h \neq 1} A_{1h} b_h \ge c_2 \sum_{h \neq 1} A_{1h} b_1 \ge c_2 A_{12} b_1,$$

which implies  $b_1$  is bounded from below, away from zero. Hence the Claim follows.

We can choose a small  $\delta_0 > 0$  such that  $b_2^{-\frac{2}{n-2s}} - \delta_0 > 0$ . The constant  $C_1$  and  $C_2$  in the introduction can be defined by

$$C_1 = b_2^{-\frac{2}{n-2s}} - \delta_0$$
 and  $C_2 = b_1^{-\frac{2}{n-2s}} + \delta_0.$  (3.6)

For any  $x = (x_1, \ldots, x_{(m+1)^k}) \in \mathbb{R}^{(m+1)^k}$ , we denote  $||x||_0 = \max_j \{|x_j|\}$ . Let  $|\frac{x_{i_0}}{b_{i_0}}| = ||\frac{x}{b}||_0$ . From the claim above, we know  $|x_{i_0}| \ge C ||x||_0$ . Using (3.5), we have

$$\begin{aligned} |(D^2 F(b)x)_{i_0}| &= |c_2 \sum_{h \neq i_0} A_{i_0h} x_h - c_1 (\frac{2\beta}{n-2s} - 1) b_{i_0}^{\frac{2n}{n-2s} - 2} x_{i_0}| \\ &\geq c_1 (\frac{2\beta}{n-2s} - 1) b_{i_0}^{\frac{2\beta}{n-2s} - 2} |x_{i_0}| - c_2 |\sum_{h \neq i} A_{i_0h} x_h| \\ &\geq c_1 (\frac{2\beta}{n-2s} - 1) b_{i_0}^{\frac{2\beta}{n-2s} - 1} |\frac{x_{i_0}}{b_{i_0}}| - c_2 \sum_{h \neq i} A_{i_0h} b_h |\frac{x_{i_0}}{b_{i_0}}| \\ &= c_1 (\frac{2\beta}{n-2s} - 2) b_{i_0}^{\frac{2\beta}{n-2s} - 2} |x_{i_0}| \geq C_4 ||x||_0. \end{aligned}$$

From the definition of  $\|\cdot\|_0$ , we get  $\|D^2F(b)x\|_0 \ge C_4\|x\|_0$ .

Let  $\theta = (\theta_1, \dots, \theta_{(m+1)^k}) \in \mathbb{R}^{(m+1)^k}$  whose component  $\theta_i := d_i - b_i, i = 1, \dots, (m+1)^k$ . We define  $X := (X^1, \dots, X^{(m+1)^k}) \in \mathbb{R}^{n \times (m+1)^k}$ , in which  $X^i = (X^i_1, \dots, X^i_n) \in \mathbb{R}^n$  for  $i = 1, \dots, (m+1)^k$ . For any  $Y = (Y^1, \dots, Y^{(m+1)^k}) \in \mathbb{R}^{n \times (m+1)^k}$ , we use the notation  $||Y|| := \max_{i=1,\dots,(m+1)^k} \{|Y^i|\}$  to denote the maximum norm.

To simplify the equations (3.4), we need to define some vector value functions below. Let  $\Xi^{(1)} := \Xi^{(1)}(P, \Lambda, \lambda) \in \mathbb{R}^{n \times (m+1)^k}$  and  $\Theta^{(1)} := \Theta^{(1)}(P, \Lambda, \lambda) \in \mathbb{R}^{n \times (m+1)^k}$  with their exponents defined by

$$(\Xi^{(1)})_{j}^{i} = \frac{\Lambda_{i}^{\beta-2}\Xi_{j}^{i}|P^{i} - X^{i}|^{2}}{c_{3}a_{j}\|P - X\|^{2}} \quad \text{and} \quad (\Theta^{(1)})_{j}^{i} = \frac{\Lambda_{i}^{\beta-2}}{c_{3}a_{j}}\Theta_{j}^{i}, \quad i = 1, \dots, (m+1)^{k}; j = 1, \dots, n.$$

Let  $\Xi^{(2)} := \Xi^{(2)}(P, \Lambda, \lambda) \in \mathbb{R}^{(m+1)^k}, \Theta^{(2)} := \Theta^{(2)}(P, \Lambda, \lambda) \in \mathbb{R}^{(m+1)^k}$  with exponents defined by  $(\Xi^{(2)})^i = \Lambda_i^{\frac{n-2s}{2}+1} \Xi_{n+1}^i \frac{|P^i - X^i|^{\min\{2,\beta-1\}}}{\|P - X\|^{\min\{2,\beta-1\}}}$  and  $(\Theta^{(2)})^i = \Lambda_i^{\frac{n-2s}{2}+1} \Theta_{n+1}^i, \quad i = 1, \dots, (m+1)^k.$ 

Define  $\Pi(\theta) := (\Pi(\theta)^1, \dots, \Pi(\theta)^{(m+1)^k})$ , where  $\Pi(\theta)^i (i = 1, \dots, (m+1)^k)$  is defined by

$$(\Pi(\theta))^{i} = \int_{0}^{1} (D^{3}F(b+s\theta)\theta,\theta)_{i}(1-s)ds = \int_{0}^{1} c_{1}(\frac{2\beta}{n-2s}-1)(\frac{2\beta}{n-2s}-2)(b_{i}+s\theta_{i})^{\frac{2\beta}{n-2s}-3}\theta_{i}^{2}.$$

From their definition, we know there is a constant C and a constant  $C_{\lambda}$  satisfying  $C_{\lambda} \to 0$  as  $\lambda \to \infty$  such that  $\|\Xi^{(1)}\| \leq C$ ,  $\|\Xi^{(2)}\|_0 \leq C$ ;  $\|\Theta^{(1)}\| \leq C_{\lambda}$  and  $\|\Theta^{(2)}\|_0 \leq C_{\lambda}$ .

Using these notations and Taylor expansion, we can write the equations (3.4) into another form:

$$\begin{cases} P - X = \|P - X\|^2 \Xi^{(1)} + \Theta^{(1)}; \\ D^2 F(b)\theta = \|P - X\|^{\min\{2,\beta-1\}} \Xi^{(2)} + \Theta^{(2)} + \Pi(\theta), \end{cases}$$
(3.7)

Let

$$B = \left(\prod_{i=1}^{(m+1)^{k}} B_{2C_{\lambda}}(X^{i})\right) \times B_{3C_{4}^{-1}C_{\lambda}}(0) \in \mathbb{R}^{n \times (m+1)^{k}} \times \mathbb{R}^{(m+1)^{k}}.$$

Define a function

$$G: B \to B$$
  
(P, \theta)  $\mapsto (X + \Xi^{(1)} \| P - X \|^2 + \Theta^{(1)}, D^2 F(b)^{-1} (\| P - X \|^{\beta - 1} \Xi^{(2)} + \Theta^{(2)} + \Pi(\theta)))$ 

For each  $(P, \theta) \in B$ , Choose  $C_{\lambda}$  small enough, we have

$$\|\Xi^{(1)}\|P - X\|^2 + \Theta^{(1)}\| \le C(2C_{\lambda})^2 + C_{\lambda} \le 2C_{\lambda}^2,$$

and

$$\|D^2 F(b)^{-1}(\|P - X\|^{\min\{2,\beta-1\}} \Xi^{(2)} + \Pi(\theta))\|_0 \le C_4^{-1}(CC_\lambda^{\min\{2,\beta-1\}} + C_\lambda + C\theta^2) \le 3C_4^{-1}C_\lambda.$$

Since  $C_{\lambda} \to 0$  as  $\lambda \to \infty$ , so for  $\lambda$  large enough, we use the Brouwer fixed-point theorem to get a solution  $(P^1, \ldots, P^{(m+1)^k}, \theta)$  of (3.7) in *B*. It holds that

$$|P^{i} - X^{i}| \le 2C_{\lambda}$$
 and  $|\theta_{i}| = |b_{i} - \Lambda_{i}^{-\frac{n-2s}{2}}| \le 3C_{4}^{-1}C_{\lambda}.$ 

Hence we find a critical point of I of the form  $u_m := W_m + \phi_m$  with  $m < \infty$ . Next, we prove  $u_m$  is a positive function. Denote  $u_m^- = \min\{0, u_m\}$  and  $u_m^+ = u_m - u_m^-$ . Then we have

$$\int_{\mathbb{R}^n} (-\Delta)^s u_m(x) u_m^-(x) dx = \int_{\mathbb{R}^n} K(\frac{x}{\lambda}) u_m^{\frac{n+2s}{n-2s}}(x) u_m^-(x) dx.$$

From the definition of  $(-\Delta)^s$ ,

$$\begin{split} \int_{\mathbb{R}^n} (-\Delta)^s u_m u_m^- dx &= \int_{\mathbb{R}^n} (-\Delta)^s u_m^- u_m^- dx + \int_{\mathbb{R}^n} (-\Delta)^s u_m^+ u_m^- dx \\ &= \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u_m^-|^2 dx + \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{(u_m^+(x) - u_m^+(y)) u_m^-(x)}{|x - y|^{n + 2s}} dx dy \\ &\ge \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u_m^-|^2 dx. \end{split}$$

The Hardy-Littlewood-Sobolev inequality yields

$$\left( \int_{\mathbb{R}^n} |u_m^-|^{\frac{2n}{n-2s}} dx \right)^{\frac{n-2s}{n}} \leq C \int_{\mathbb{R}^n} |(-\Delta)^{\frac{s}{2}} u_m^-|^2 dx \\ \leq C \int_{\mathbb{R}^n} K(\frac{x}{\lambda}) u_m^{\frac{n+2s}{n-2s}}(x) u_m^-(x) dx \leq C \int_{\mathbb{R}^n} |u_m^-|^{\frac{2n}{n-2s}} dx.$$

Suppose  $u_m^- \neq 0$ , we have  $\int_{\mathbb{R}^n} |u_m^-|^{\frac{2n}{n-2s}} dx \ge C$ . It is easy to get  $u_m^- \le |\phi_m|$ . From this fact,

$$\begin{split} \int_{\mathbb{R}^n} |u_m^-|^{\frac{2n}{n-2s}} dx &\leq \int_{\mathbb{R}^n} |\phi_m|^{\frac{2n}{n-2s}} dx \\ &\leq \|\phi_m\|_*^{\frac{2n}{n-2s}} \int_{\mathbb{R}^n} \left(\gamma(x) \sum_{h=1}^{(m+1)^k} \frac{1}{(1+|x-X^h|)^{\frac{n-2s}{2}+\tau}}\right)^{\frac{2n}{n-2s}} dx \\ &\leq \|\phi_m\|_*^{\frac{2n}{n-2s}} C(m,k) \sum_h \int_{\mathbb{R}^n} \frac{1}{(1+|x-X^h|)^{n+\frac{2n}{n-2s}\tau}} dx. \end{split}$$

So we get  $C \leq \int_{\mathbb{R}^n} |u_m^-|^{\frac{2n}{n-2s}} dx \leq C(m,k) \|\phi_m\|_* \to 0$  as  $\lambda \to \infty$ , which is impossible. Hence  $u_m \geq 0$ . Suppose there is a point  $x_0$  such that  $u_m(x_0) = 0$ , then

$$0 = K(\frac{x_0}{\lambda})u_m^{\frac{n+2s}{n-2s}}(x_0) = (-\Delta)^s u_m(x_0) = P.V. \int_{\mathbb{R}^n} \frac{u_m(x_0) - u_m(y)}{|x_0 - y|^{n+2s}} dy = P.V. \int_{\mathbb{R}^n} \frac{-u_m(y)}{|x_0 - y|^{n+2s}} dy.$$

Then  $u_m \equiv 0$  which is impossible. Hence  $u_m > 0$ .

According to Proposition 2.5,  $u_m = W_m + \phi_m \in C^{0,\alpha}(\mathbb{R}^n) \cap \dot{H}^s(\mathbb{R}^n)$ . Using local Schauder estimate [18, Proposition 2.11] and a bootstrap argument, we know  $u_m \in C^{2,\alpha'}_{loc}(\mathbb{R}^n)$ , for some  $\alpha' \in (0,1)$ .

What is more,  $|u_m|_{L^{\infty}}(\mathbb{R}^n) \leq C$  with C independent of m. In fact, Choosing  $x \in \Omega_1$  with no loss of generality, we have

$$W_m(x) \le \sum_{i=1}^{(m+1)^k} \frac{C}{(1+|x-X^i|)^{n-2s}} \le C + \sum_{i\neq 1}^{(m+1)^k} \frac{C}{|X^i-X^1|^{n-2s}} \le C + \frac{C}{(\lambda l)^{n-2s}} \le C,$$

and

$$|\phi_m| \le \|\phi_m\|_* \sum_{h=1}^{(m+1)^k} \frac{1}{(1+|x-X^h|)^{\frac{n-2s}{2}+\tau}} \le \|\phi_m\|_* \sum_{h=1}^{\infty} \frac{1}{(1+|x-X^h|)^{\frac{n-2s}{2}+\tau}} \le C \|\phi_m\|_* \le C.$$

Since  $\phi_m$  satisfies the equation  $(-\Delta)^s \phi_m - \frac{n+2s}{n-2s} K(\frac{x}{\lambda}) W_m^{\frac{4s}{n-2s}} \phi_m = N(\phi_m) + l_m$ , then from Lemma A.8, Lemma 2.4 and Proposition 2.5, we know that for any  $x, y \in \mathbb{R}^n$  with  $x \neq y$ , there holds

$$\frac{|\phi_m(x) - \phi_m(y)|}{|x - y|^{\alpha}} \le \frac{C}{\lambda^{\tau - s}} \max\{\|\phi_m\|_*, \|N(\phi_m)\|_{**} + \|l_m\|_{**}\} \le C, \text{ where } \alpha = \min\{1, 2s\}.$$

Also from simple calculation, we get for any  $x \in \mathbb{R}^n$  and R > 0,  $||W_m||_{C^{0,\alpha}(B_{2R}(x))} \leq C(n, R)$ , where C(n, R) is a constant independent of m. Hence  $||u_m||_{C^{0,\alpha}(B_{2R}(x))} \leq C(n, R)$ . Local Schauder estimate and a bootstrip argument yields that  $||u_m||_{C^{2,\alpha'}(B_R(x))} \leq C(n, R)$ . Thanks to Azellà-Ascolli theorem, we have  $u_m$  convergent uniformly to a  $C_{loc}^{2,\alpha'}$  function  $u_{\infty} = W_{\infty} + \phi_{\infty}$  on compact sets as  $m \to \infty$ . We know  $u_{\infty}$  satisfies  $|u_{\infty}|_{L^{\infty}(\mathbb{R}^n)} \leq C$  and  $||u_{\infty}||_{C^{2,\alpha'}(B_R(x))} \leq C(n, R)$ .

We will show that  $u_{\infty}$  satisfies the equation (1.5). Let  $v_m = u_m - u_{\infty}$ . From above, we know  $v_m$  has the property  $|v_m|_{L^{\infty}(\mathbb{R}^n)} \leq C$ ;  $|v_m|_{C^2(B_1(x))} \leq C$  and  $v_m \to 0$  uniformly on compact sets.

From the definition of  $(-\Delta)^s$ , we have for any  $x \in \mathbb{R}^n$ 

$$C(n,s)^{-1}|(-\Delta)^{s}v_{m}(x)| \leq P.V \int_{\mathbb{R}^{n}} \frac{|v_{m}(x) - v_{m}(y)|}{|x - y|^{n + 2s}} dy \\ = P.V. \int_{B_{\varepsilon_{0}}(x)} \frac{|v_{m}(x) - v_{m}(y)|}{|x - y|^{n + 2s}} dy + \int_{B_{R}(x) \setminus B_{\varepsilon_{0}}(x)} \frac{|v_{m}(x) - v_{m}(y)|}{|x - y|^{n + 2s}} dy \\ + \int_{\mathbb{R}^{n} \setminus B_{R}(x)} \frac{|v_{m}(x) - v_{m}(y)|}{|x - y|^{n + 2s}} dy \\ =: T_{1} + T_{2} + T_{3}.$$

For the term  $T_1$ , we have

$$T_{1} = \frac{1}{2} P.V. \int_{B_{\varepsilon_{0}}(0)} \frac{|v_{m}(x+y) + v_{m}(x-y) - 2v_{m}(x)|}{|y|^{n+2s}}$$
  

$$\leq C|v_{m}|_{C^{2}B_{1}(x)} \int_{B_{\varepsilon_{0}}(0)} |y|^{2-2s-n} dy \leq C|v_{m}|_{C^{2}B_{1}(x)} \varepsilon_{0}^{2-2s} \to 0 \text{ as } \varepsilon_{0} \to 0.$$

For the third term,

$$T_3 \le C \int_{\mathbb{R}^n \setminus B_R(x)} \frac{1}{|x-y|^{n+2s}} = CR^{-2s} \to 0 \text{ as } R \to \infty.$$

Then we estimate the term  $T_2$ . For fixed R large enough and  $\varepsilon_0$  small enough,  $B_R(x) \setminus B_{\varepsilon_0}(x)$  is a compact set. So we have  $T_2 \to 0$  as  $m \to \infty$ . Hence  $(-\Delta)^s u_m(x) \to (-\Delta)^s u_\infty(x)$  as  $m \to \infty$ . Therefore  $u_\infty$  satisfies equation (1.5).

Proof of Corollary 1.3. Fix the constant  $m < \infty$ . Using a similar argument as in [29], we can expand  $I(W_m + \phi_m)$  as

$$I(W_m + \phi_m) = (m+1)^k \left(\frac{s}{n} \int U_{0,1}^{\frac{2n}{n-2s}} + o(1)\right), \text{ as } l \to \infty.$$

For each  $m < \infty$ ,  $I(W_m + \phi_m) \to (m+1)^k \frac{s}{n} \int U_{0,1}^{\frac{2n}{n-2s}}$  as  $l \to \infty$ . For any  $m_1, m_2 \in \mathbb{N}_+$  such that  $m_1 \neq m_2$ , we can find two solutions  $W_{m_1} + \phi_{m_1}$  and  $W_{m_2} + \phi_{m_2}$  of (1.5), such that  $I(W_{m_1} + \phi_{m_1}) \neq I(W_{m_2} + \phi_{m_2})$ . Hence we can find infinitely many solutions of (1.5).

### APPENDIX A. BASIC ESTIMATES

**Lemma A.1.** (cf. [21, 29]) For any  $x_i, x_j, y \in \mathbb{R}^n$  and constant  $\sigma \in [0, \min\{\alpha, \beta\}]$ , we have

$$\frac{1}{(1+|y-x_i|)^{\alpha}(1+|y-x_j|)^{\beta}} \le \frac{2^{\sigma}}{(1+|x_i-x_j|)^{\sigma}} \left(\frac{1}{(1+|y-x_i|)^{\alpha+\beta-\sigma}} + \frac{1}{(1+|y-x_j|)^{\alpha+\beta-\sigma}}\right) + \frac{1}{(1+|y-x_j|)^{\alpha+\beta-\sigma}} = \frac{1}{(1+|y-x_j|)^{\alpha+\beta-\sigma}} + \frac{1}{(1+|y-x_$$

**Lemma A.2.** For any  $\sigma > 0$  with  $\sigma \neq n - 2s$ , there is a constant C > 0 such that

$$\int_{\mathbb{R}^n} \frac{1}{|y-z|^{n-2s}} \frac{1}{(1+|z|)^{2s+\sigma}} dz \le \frac{C}{(1+|y|)^{\min(\sigma,n-2s)}}.$$

For  $\sigma = n - 2s$ , there is also a constant C > 0, such that

$$\int_{\mathbb{R}^n} \frac{1}{|y-z|^{n-2s}} \frac{1}{(1+|z|)^n} \le C \frac{\max(1,\log|y|)}{(1+|y|)^{n-2s}}$$

*Proof.* The proof follows from the same argument as [21, Lemma A.2]. See also [29, Lemma B.2].

Recall that  $X^i \in X_{i,m} = \{X^i\}_{i=1}^{(m+1)^k}$ ,  $B_i = B_{\lambda l}(X^i)$  and  $B_{i,m} = B_{\max\{\frac{m}{4},1\}\lambda l}(X^i)$ .

**Lemma A.3.** (cf. [21]) For any  $\theta > k$ , there exists a constant  $C(\theta, k, n) > 1$  independent of m, such that if  $y \in B_i \cap \Omega_i$ , there holds

$$\frac{1}{(1+|y-X^i|)^{\theta}} \le \sum_j \frac{1}{(1+|y-X^j|)^{\theta}} \le \frac{C}{(1+|y-X^i|)^{\theta}}.$$

If  $y \in B_i^c \cap B_{i,m} \cap \Omega_i$ , there holds

$$\frac{1}{C(1+|y-X^i|)^{\theta-k}(\lambda l)^k} \le \sum_j \frac{1}{(1+|y-X^j|)^{\theta}} \le \frac{C}{(1+|y-X^i|)^{\theta-k}(\lambda l)^k}.$$
 (A.1)

and if  $y \in B_{i,m}^c \cap \Omega_i$ , there holds

$$\frac{m^k}{C(1+|y-X^i|)^{\theta}} \le \sum_j \frac{1}{(1+|y-X^j|)^{\theta}} \le \frac{Cm^k}{(1+|y-X^i|)^{\theta}} \le \frac{C}{(1+|y-X^i|)^{\theta-k}(\lambda l)^k}.$$

**Lemma A.4.** Let n > 2s + 2 and  $0 < \tau < \frac{n+2s}{2}$ . If  $\phi$  satisfies  $\|\phi\|_* \leq \frac{C}{\lambda^{\frac{n+2s}{2}-\tau}}$ , then for any c > 0, there exists  $\lambda_0$  such that for any  $\lambda > \lambda_0$ , there holds  $|\phi| \leq cW_m$  in  $\cup_h (\Omega_h \cap B_h)$ .

*Proof.* We prove this lemma indirectly. Suppose that there exists  $c_0 > 0$ , such that for any  $\lambda_0 > 0$ , there is a  $\lambda > \lambda_0$  and  $y \in \bigcup_l (\Omega_l \cap B_l)$  such that  $|\phi(y)| \ge c_0 W_m(y)$ . Then

$$\begin{aligned} |\phi(y)| &\geq C \sum_{h=1}^{(m+1)^k} \frac{1}{(1+|y-X^h|)^{n-2s}} \\ &\geq C\gamma(y) \sum_{h=1}^{(m+1)^k} \frac{1}{(1+|y-X^h|)^{\frac{n-2s}{2}+\tau}} \frac{1}{(\lambda l)^{\frac{n-2s}{2}-\tau}} \end{aligned}$$

If  $0 < \tau < \frac{n-2s}{2}$ , we have

$$\frac{1}{\lambda^{\frac{n+2s}{2}} - \tau} \ge \|\phi\|_* \ge \frac{C}{(\lambda l)^{\frac{n-2s}{2}} - \tau},$$

which does not hold for  $\lambda$  large enough.

If  $\frac{n-2s}{2} \le \tau < \frac{n+2s}{2}$ , we can also get

$$\frac{1}{\lambda^{\frac{n+2s}{2}-\tau}} \ge \|\phi\|_* \ge C,$$

which also is a contradiction for  $\lambda$  large.

**Lemma A.5.** For n > 2s + 2 and  $1 \le k < \tau < \frac{n-2s}{2}$ , we have

$$\begin{split} &\int_{\mathbb{R}^n} \frac{1}{|x-y|^{n-2s}} W_{P,\Lambda}^{\frac{4s}{n-2s}}(y) \gamma(y) \sum_h \frac{1}{(1+|y-X^h|)^{\frac{n-2s}{2}+\tau}} dy \\ &\leq C \left( \gamma(x) \sum_h \frac{1}{(1+|x-X^h|)^{\frac{n-2s}{2}+\tau+\theta}} + \frac{1}{(\lambda l)^{\frac{4s}{n-2s}k}} \gamma(x) \sum_h \frac{1}{(1+|x-X^h|)^{\frac{n-2s}{2}+\tau}} \right), \end{split}$$

where  $\theta > 0$  is a small constant and C > 0 does not depend on m.

*Proof.* Without loss of generality, we assume  $x \in \Omega_1$ . We write

We now estimate each term  $T_i(i = 1, 2, 3)$ .

Using Lemma A.2 and Lemma A.3, we have

$$T_{1} \leq C \int_{\bigcup_{h}(\Omega_{h}\cap B_{h})} \frac{1}{|x-y|^{n-2s}} \sum_{h} \frac{1}{(1+|y-X^{h}|)^{4s+\frac{n-2s}{2}+\tau}} dy$$
  
$$\leq C \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\min\{\frac{n+2s}{2}+\tau,n-2s\}}}$$
  
$$= C \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\frac{n-2s}{2}+\tau+\theta_{1}}}, \quad \text{where } \theta_{1} = \min\{2s, \frac{n-2s}{2}-\tau\}. \quad (A.2)$$

Similarly, we also obtain

$$T_{1} \leq C \int_{\bigcup_{h}(\Omega_{h}\cap B_{h})} \frac{1}{|x-y|^{n-2s}} \frac{1}{\lambda^{\tau-s}} \sum_{h} \frac{1}{(1+|y-X^{h}|)^{\frac{n}{2}+4s}} dy$$

$$\leq \frac{C}{\lambda^{\tau-s}} \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\min\{\frac{n}{2}+2s,n-2s\}}}$$

$$\leq C \frac{(1+|x-X^{1}|)^{\tau-s}}{\lambda^{\tau-s}} \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\min\{n-3s+\tau,\frac{n}{2}+\tau+s\}}}$$

$$= C \frac{(1+|x-X^{1}|)^{\tau-s}}{\lambda^{\tau-s}} \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\frac{n-2s}{2}+\tau+\theta_{2}}}, \quad (A.3)$$

where  $\theta_2 = \min\{\frac{n-4s}{2}, 2s\}$ . Combining the estimate (A.2) and (A.3), we have

$$T_1 \le C\gamma(x) \sum_h \frac{1}{(1+|x-X^h|)^{\frac{n-2s}{2}+\tau+\theta}},$$

where  $\theta = \min\{2s, \frac{n-4s}{2}, \frac{n-2s}{2} - \tau\}.$ For the term  $T_2$ , we have

$$T_{2} \leq \frac{C}{(\lambda l)^{\frac{4s}{n-2s}k}} \int_{\mathbb{R}^{n}} \frac{1}{|x-y|^{n-2s}} \sum_{h} \frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+4s+\tau-\frac{4s}{n-2s}k}} dy$$

$$\leq \frac{C}{(\lambda l)^{\frac{4s}{n-2s}k}} \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\min\{n-2s,\frac{n-2s}{2}+\tau+2s-\frac{4s}{n-2s}k\}}}$$

$$\leq \frac{C}{(\lambda l)^{\frac{4s}{n-2s}k}} \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\frac{n-2s}{2}+\tau}},$$

and

$$T_{2} \leq \frac{C}{\lambda^{\tau-s}(\lambda l)^{\frac{4s}{n-2s}k}} \int_{\mathbb{R}^{n}} \frac{1}{|x-y|^{n-2s}} \sum_{h} \frac{1}{(1+|y-X^{h}|)^{\frac{n}{2}+4s-\frac{4s}{n-2s}k}} dy$$

$$\leq \frac{C}{\lambda^{\tau-s}(\lambda l)^{\frac{4s}{n-2s}k}} \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\min\{n-2s,\frac{n}{2}+2s-\frac{4s}{n-2s}k\}}}$$

$$\leq \frac{C}{(\lambda l)^{\frac{4s}{n-2s}k}} \frac{(1+|x-X^{1}|)^{\tau-s}}{\lambda^{\tau-s}} \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\frac{n-2s}{2}+\tau}}.$$

Thus

$$T_2 \le \frac{C}{(\lambda l)^{\frac{4s}{n-2s}k}} \gamma(x) \sum_h \frac{1}{(1+|x-X^h|)^{\frac{n-2s}{2}+\tau}}.$$

By the same procedure, we have

$$T_3 \le \frac{C}{(\lambda l)^{\frac{4s}{n-2s}k}} \gamma(x) \sum_h \frac{1}{(1+|x-X^h|)^{\frac{n-2s}{2}+\tau}}$$

Hence this lemma follows.

Remember 
$$Z_{i,j} = \frac{\partial U_{P^i,\Lambda_i}}{\partial P_j^i}$$
 for  $j = 1, 2, \dots, n$  and  $Z_{i,n+1} = \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i}$ 

**Lemma A.6.** For t = 1, 2..., n + 1, we have

$$\left| \int_{\mathbb{R}^n} K(\frac{x}{\lambda}) W_m^{\frac{4s}{n-2s}} Z_{r,t} \phi \right| \le \frac{C \|\phi\|_*}{\lambda^{\tau-s} (\lambda l)^{\frac{n}{2}}}$$

*Proof.* In the proof of this lemma, we denote  $\hat{W}_{m,r} = \sum_{h \neq r} U_{P^h,\Lambda_h}$ . It is easy to get

$$\int_{\mathbb{R}^{n}} K(\frac{x}{\lambda}) W_{m}^{\frac{4s}{n-2s}} Z_{r,t} \phi = \int_{\mathbb{R}^{n}} K(\frac{x}{\lambda}) U_{P^{r},\Lambda_{r}}^{\frac{4s}{n-2s}} Z_{r,t} \phi + O\left(\int_{\hat{W}_{m,r} > U_{P^{r},\Lambda_{r}}} \hat{W}_{m,r}^{\frac{4s}{n-2s}} Z_{r,t} \phi\right)$$

$$+ O\left(\int_{\hat{W}_{m,r} \le U_{P^{r},\Lambda_{r}}} U_{P^{r},\Lambda_{r}}^{\frac{4s}{n-2s}} \hat{W}_{m,r} \phi\right).$$
(A.4)

We need to estimate each term in the equality above.

For  $i \neq r$ , from Lemma A.3, we have

$$\begin{aligned} & \left| \int_{\Omega_i \cap B_i} \hat{W}_{m,r}^{\frac{4s}{n-2s}} Z_{r,t} \phi \right| \\ & \leq C \frac{\|\phi\|_*}{\lambda^{\tau-s}} \int_{\Omega_i \cap B_i} \left( \sum_{h \neq r} \frac{1}{(1+|x-X^h|)^{n-2s}} \right)^{\frac{4s}{n-2s}} \frac{1}{(1+|x-X^r|)^{n-2s}} \sum_h \frac{1}{(1+|x-X^h|)^{\frac{n}{2}}} dx \\ & \leq C \frac{\|\phi\|_*}{\lambda^{\tau-s}} \int_{\Omega_i \cap B_i} \frac{1}{(1+|x-X^r|)^{n-2s}} \frac{1}{(1+|x-X^i|)^{\frac{n}{2}+4s}} dx \leq C \frac{\|\phi\|_*}{\lambda^{\tau-s}|X^r-X^i|^{\frac{n}{2}}}. \end{aligned}$$

With the help of Lemma A.1 and Lemma A.3, we get

$$\begin{aligned} \left| \int_{\cup_{h}(\Omega_{h}\cap B_{h}^{c})} \hat{W}_{m,r}^{\frac{4s}{n-2s}} Z_{r,t} \phi \right| \\ &\leq C \|\phi\|_{*} \int_{\cup_{h}(\Omega_{h}\cap B_{h}^{c})} \frac{1}{(1+|x-X^{r}|)^{n-2s}} \left( \sum_{h\neq r} \frac{1}{(1+|x-X^{h}|)^{n-2s}} \right)^{\frac{4s}{n-2s}} \\ &\qquad \times \left( \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\frac{n-2s}{2}+\tau}} \right) dx \end{aligned} \tag{A.5}$$

$$&\leq C \frac{\|\phi\|_{*}}{(\lambda l)^{\frac{4s}{n-2s}k}} \int_{\cup_{h}(\Omega_{h}\cap B_{h}^{c})} \frac{1}{(1+|x-X^{r}|)^{n-2s}} \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\frac{n}{2}+3s+\tau-\frac{4s}{n-2s}k}} dx \\ &\leq C \frac{\|\phi\|_{*}}{(\lambda l)^{\frac{n}{2}+s+\tau}}. \end{aligned}$$

Since in  $\Omega_r \cap B_r$ , there holds  $\hat{W}_{m,r} \leq \sum_{j \neq r} \frac{C}{|X^j - X^r|^{n-2s}} \leq \frac{C}{(\lambda l)^{n-2s}}$ . Then if  $n \geq 6s$ , we have

$$\begin{aligned} \left| \int_{\substack{\Omega_{r} \cap B_{r} \\ \hat{W}_{m,r} > U_{Pr}, \Lambda_{r}}} \hat{W}_{m,r}^{\frac{4s}{n-2s}} Z_{r,t} \phi \right| &\leq C \int_{\Omega_{r} \cap B_{r}} \hat{W}_{m,r}^{\frac{n+2s}{2(n-2s)}} U_{Pr,\Lambda_{r}}^{\frac{n+2s}{2(n-2s)}} |\phi| \\ &\leq C \frac{\|\phi\|_{*}}{\lambda^{\tau-s}} \frac{1}{(\lambda l)^{\frac{n+2s}{2}}} \int_{\Omega_{r} \cap B_{r}} \frac{1}{(1+|x-X^{r}|)^{\frac{n+2s}{2}}} \sum_{h} \frac{1}{(1+|x-X^{h}|)^{\frac{n}{2}}} dx \\ &\leq C \frac{\|\phi\|_{*}}{\lambda^{\tau-s} (\lambda l)^{\frac{n+2s}{2}}}. \end{aligned}$$

And if n < 6s, we get  $\frac{n+2s}{2} < 4s$ . In this case

$$\begin{aligned} \left| \int_{\substack{\Omega_r \cap B_r \\ \hat{W}_{m,r} > U_{P^r,\Lambda_r}}} \hat{W}_{m,r}^{\frac{4s}{n-2s}} Z_{r,t} \phi \right| &\leq \frac{C \|\phi\|_*}{\lambda^{\tau-s} (\lambda l)^{4s}} \int_{\Omega_r \cap B_r} \frac{1}{(1+|x-X^r|)^{n-2s}} \sum_h \frac{1}{(1+|x-X^h|)^{\frac{n}{2}}} dx \\ &\leq \frac{C \|\phi\|_*}{\lambda^{\tau-s} (\lambda l)^{\frac{n+2s}{2}}}. \end{aligned}$$

From these arguments above, we arrive

$$\left| \int_{\hat{W}_{m,r} > U_{P^r,\Lambda_r}} \hat{W}_{m,r}^{\frac{4s}{n-2s}} Z_{r,t} \phi \right| \le C \frac{\|\phi\|_*}{\lambda^{\tau-s} (\lambda l)^{\frac{n}{2}}}.$$
(A.6)

By a similar procedure, we get

$$\left| \int_{\hat{W}_{m,r} \le U_{P^r,\Lambda_r}} U_{P^r,\Lambda_r}^{\frac{4s}{n-2s}} \hat{W}_{m,r} \phi \right| \le C \frac{\|\phi\|_*}{\lambda^{\tau-s} (\lambda l)^{\frac{n}{2}}}.$$
(A.7)

Now we estimate the first term on the right hand side of the equality (A.4). Since  $\phi$  satisfies the second equality in (2.10), we have

$$\left| \int_{\mathbb{R}^n} K(\frac{x}{\lambda}) U_{P^r,\Lambda_r}^{\frac{4s}{n-2s}} Z_{r,t} \phi \right| = \left| \int_{\mathbb{R}^n} \left( K(\frac{x}{\lambda}) - 1 \right) U_{P^r,\Lambda_r}^{\frac{4s}{n-2s}} Z_{r,t} \phi \right|$$
  
$$\leq \frac{\|\phi\|_*}{\lambda^{\tau-s}} \int_{\mathbb{R}^n} \left| K(\frac{x}{\lambda}) - 1 \right| U_{P^r,\Lambda_r}^{\frac{n+2s}{n-2s}} \sum_h \frac{1}{(1+|x-X^h|)^{\frac{n}{2}}} dx.$$

On one hand, Lemma A.1 implies that

$$\begin{split} \int_{\mathbb{R}^n} \left| K(\frac{x}{\lambda}) - 1 \right| U_{P^r,\Lambda_r}^{\frac{n+2s}{n-2s}} \sum_{h \neq r} \frac{1}{(1+|x-X^h|)^{\frac{n}{2}}} dx \\ &\leq C \int_{\mathbb{R}^n} \frac{1}{(1+|x-X^r|)^{n+2s}} \sum_{h \neq r} \frac{1}{(1+|x-X^h|)^{\frac{n}{2}}} dx \\ &\leq \frac{C}{(\lambda l)^{\frac{n}{2}}}. \end{split}$$

On the another hand, choose  $\delta$  to be a fixed constant small enough,

$$\int_{\mathbb{R}^{n}} \left| K(\frac{x}{\lambda}) - 1 \right| U_{P^{r},\Lambda_{r}}^{\frac{n+2s}{n-2s}} \frac{1}{(1+|x-X^{r}|)^{\frac{n}{2}}} \\ = \int_{|x-X^{r}| \le \delta\lambda} \left| K(\frac{x}{\lambda}) - 1 \right| U_{P^{r},\Lambda_{r}}^{\frac{n+2s}{n-2s}} \frac{1}{(1+|x-X^{r}|)^{\frac{n}{2}}} \\ + \int_{|x-X^{r}| > \delta\lambda} \left| K(\frac{x}{\lambda}) - 1 \right| U_{P^{r},\Lambda_{r}}^{\frac{n+2s}{n-2s}} \frac{1}{(1+|x-X^{r}|)^{\frac{n}{2}}} =: J_{1} + J_{2}.$$

From the condition  $(H_3)$ , we have

$$|J_1| \le \frac{C}{\lambda^{\beta}} \int_{|x-X^r| \le \delta\lambda} \frac{|x-X^r|^{\beta}}{(1+|x-X^r|)^{n+2s+\frac{n}{2}}} \le \begin{cases} \frac{C\log\lambda}{\lambda^{\frac{n+4s}{2}}}, & \text{if } \beta \ge \frac{n+4s}{2}, \\ \frac{C}{\lambda^{\beta}}, & \text{if } \beta < \frac{n+4s}{2}. \end{cases}$$

For the term  $J_2$ , a direct calculation yields

$$J_{2} = \int_{|x-X^{r}|>\delta\lambda} \left| K(\frac{x}{\lambda}) - 1 \right| U_{P^{r},\Lambda_{r}}^{\frac{n+2s}{n-2s}} \frac{1}{(1+|x-X^{r}|)^{\frac{n}{2}}} \\ \leq \int_{|x-X^{r}|>\delta\lambda} \frac{1}{(1+|x-X^{r}|)^{n+2s+\frac{n}{2}}} \\ \leq \frac{C}{\lambda^{\frac{n+4s}{2}}}.$$

Since  $\min\{\beta, \frac{n+4s}{2}\} > \frac{n}{2} \frac{\beta}{n-2s}$ , the definition of  $\lambda$  implies

$$\int_{\mathbb{R}^n} \left| K(\frac{x}{\lambda}) - 1 \right| U_{P^r,\Lambda_r}^{\frac{n+2s}{n-2s}} \frac{1}{(1+|x-X^r|)^{\frac{n}{2}}} \le \frac{C}{(\lambda l)^{\frac{n}{2}}}.$$

Hence we obtain

$$\left| \int_{\mathbb{R}^n} K(\frac{x}{\lambda}) U_{P^r,\Lambda_r}^{\frac{4s}{n-2s}} Z_{r,t} \phi \right| \le C \frac{\|\phi\|_*}{\lambda^{\tau-s} (\lambda l)^{\frac{n}{2}}}.$$
(A.8)

Putting (A.6), (A.7) and (A.8) into (A.4), we get this lemma.

Lemma A.7. It holds that

$$\int_{\bigcup_{h}(\Omega_{h}\cap B_{h}^{c})} |\phi|^{\frac{n+2s}{n-2s}} U_{P^{i},\Lambda_{i}} \leq C \frac{\|\phi\|_{*}^{\frac{n+2s}{n-2s}}}{(\lambda l)^{\frac{n-2s}{2}+\frac{n+2s}{n-2s}\tau}},$$
(A.9)

and

$$\left| \int_{\mathbb{R}^n} W_m^{\frac{6s-n}{n-2s}} \phi^2 U_{P^i,\Lambda_i} \right| \le \frac{C \|\phi\|_*^2}{\lambda^{2(\tau-s)}}.$$
 (A.10)

*Proof.* The estimate (A.9) follows by the same method as in (A.5). So we only prove the estimation (A.10).

Using the same trick as in (A.5), we have

$$\left| \int_{\bigcup_{h} (\Omega_{h} \cap B_{h}^{c})} W_{m}^{\frac{6s-n}{n-2s}} \phi^{2} U_{P^{i},\Lambda_{i}} \right| \leq \frac{C \|\phi\|_{*}^{2}}{\lambda^{2(\tau+s)}}.$$
(A.11)

According to Lemma A.3, we have for  $t \neq i$ ,

$$\left| \int_{\Omega_t \cap B_t} W_m^{\frac{6s-n}{n-2s}} \phi^2 U_{P^i,\Lambda_i} \right| \le \frac{C \|\phi\|_*^2}{(\lambda l)^{2(\tau-s)}} \int_{\Omega_t \cap B_t} \frac{1}{(1+|y-X^t|)^{6s}} \frac{1}{(1+|y-X^i|)^{n-2s}} dy.$$

If  $n \ge 6s$ 

$$\int_{\Omega_t \cap B_t} \frac{1}{(1+|y-X^t|)^{6s}} \frac{1}{(1+|y-X^i|)^{n-2s}} dy \le \frac{(\lambda l)^{n-6s} \log(\lambda l)}{|X^i - X^t|^{n-2s}};$$
  
< 6s, there holds

If otherwise, n < 6s, there holds

$$\int_{\Omega_t \cap B_t} \frac{1}{(1+|y-X^t|)^{6s}} \frac{1}{(1+|y-X^i|)^{n-2s}} dy \le \frac{1}{|X^i - X^t|^{n-2s}}.$$

Hence

$$\sum_{t \neq i} \left| \int_{\Omega_t \cap B_t} W_m^{\frac{6s-n}{n-2s}} \phi^2 U_{P^i,\Lambda_i} \right| \le \frac{C \|\phi\|_*^2}{(\lambda l)^{2(\tau-s)}} \frac{C \log(\lambda l)}{(\lambda l)^{\min\{n-2s,4s\}}}.$$
(A.12)

Using Lemma A.3, we also have

$$\left| \int_{\Omega_i \cap B_i} W_m^{\frac{6s-n}{n-2s}} \phi^2 U_{P^i,\Lambda_i} \right| \le \frac{C \|\phi\|_*^2}{(\lambda l)^{2(\tau-s)}} \int_{\Omega_i \cap B_i} \frac{1}{(1+|y-X^i|)^{n+4s}}.$$
 (A.13)

So we obtain the estimate (A.10) from (A.11), (A.12) and (A.13).

**Lemma A.8.** If  $\phi$  is the solution of the equation

$$(-\Delta)^{s}\phi(x) - \frac{n+2s}{n-2s}K(\frac{x}{\lambda})W_{m}^{\frac{4s}{n-2s}}(x)\phi(x) = g(x),$$
(A.14)

satisfying  $\|\phi\|_* < +\infty$ , then we have

$$\sup_{x_1 \neq x_2} \frac{|\lambda^{\tau-s}\phi(x_1) - \lambda^{\tau-s}\phi(x_2)|}{|x_1 - x_2|^{\alpha}} \le C \max\{\|\phi\|_*, \|g\|_{**}\},\$$

where  $\alpha = \min\{2s, 1\}$  and the constant C does not depend on  $\lambda$  and m.

*Proof.* Since  $|\lambda^{\tau-s}\phi(x)| \leq \|\phi\|_* \sum_h \frac{1}{(1+|x-X^h|)^{\frac{n}{2}}} \leq C\|\phi\|_*$ , we can assume  $|x_1-x_2| \leq \frac{1}{3}$  with no loss of generality. Using the Green function of  $(-\Delta)^s$  (see [5]), we can write (A.14) into the following form

$$\phi(x) = C \int_{\mathbb{R}^n} \frac{1}{|x - y|^{n - 2s}} \left( \frac{n + 2s}{n - 2s} K(\frac{y}{\lambda}) W_m^{\frac{4s}{n - 2s}}(y) \phi(y) + g(y) \right) dy,$$

Then we get

$$\begin{aligned} |\phi(x_1) - \phi(x_2)| &\leq C \left| \int_{\mathbb{R}^n} \left( \frac{1}{|x_1 - y|^{n-2s}} - \frac{1}{|x_2 - y|^{n-2s}} \right) K(\frac{y}{\lambda}) W_m^{\frac{4s}{n-2s}}(y) \phi(y) dy \right| \\ &+ C \left| \int_{\mathbb{R}^n} \left( \frac{1}{|x_1 - y|^{n-2s}} - \frac{1}{|x_2 - y|^{n-2s}} \right) g(y) dy \right| \\ &=: C(H_1 + H_2). \end{aligned}$$

Using the definition of the norm  $\|\cdot\|_*,$  there hold

$$\begin{split} |H_1| &\leq \|\phi\|_* \left| \int_{\mathbb{R}^n} \left( \frac{1}{|x_1 - x_2 - y|^{n-2s}} - \frac{1}{|y|^{n-2s}} \right) K(\frac{y + x_2}{\lambda}) W_m^{\frac{4s}{n-2s}}(y + x_2) \times \\ &\quad \times \gamma(y + x_2) \sum_h \frac{1}{(1 + |y + x_2 - X^h|)^{\frac{n-2s}{2} + \tau}} dy \right| \\ &= \|\phi\|_* \left\{ \int_{|y| \leq 3|x_1 - x_2|} \left( \frac{1}{|x_1 - x_2 - y|^{n-2s}} - \frac{1}{|y|^{n-2s}} \right) K(\frac{y + x_2}{\lambda}) W_m^{\frac{4s}{n-2s}}(y + x_2) \times \\ &\quad \times \gamma(y + x_2) \sum_h \frac{1}{(1 + |y + x_2 - X^h|)^{\frac{n-2s}{2} + \tau}} dy \\ &\quad + \int_{|y| \geq 3|x_1 - x_2|} \left( \frac{1}{|x_1 - x_2 - y|^{n-2s}} - \frac{1}{|y|^{n-2s}} \right) K(\frac{y + x_2}{\lambda}) W_m^{\frac{4s}{n-2s}}(y + x_2) \times \\ &\quad \times \gamma(y + x_2) \sum_h \frac{1}{(1 + |y + x_2 - X^h|)^{\frac{n-2s}{2} + \tau}} dy \\ &=: \|\phi\|_* (K_1 + K_2). \end{split}$$

For the term  $K_1$ , we have

$$|K_1| \le \frac{C}{\lambda^{\tau-s}} \int_{|y| \le 4|x_1 - x_2|} \frac{1}{|y|^{n-2s}} \le \frac{C}{\lambda^{\tau-s}} |x_1 - x_2|^{2s}.$$

For the term  $K_2$ , we have

$$\begin{aligned} |K_2| &\leq C|x_1 - x_2| \int_0^1 dt \int_{|y| \geq 3|x_1 - x_2|} \frac{1}{|t(x_1 - x_2) - y|^{n-2s+1}} W_m^{\frac{4s}{n-2s}}(y + x_2) \\ &\quad \times \gamma(y + x_2) \sum_h \frac{1}{(1 + |y + x_2 - X^h|)^{\frac{n-2s}{2} + \tau}} dy \\ &= C|x_1 - x_2| \int_0^1 dt \left( \int_{1 > |y| \geq 3|x_1 - x_2|} + \int_{|y| \geq 1} \right) \frac{1}{|t(x_1 - x_2) - y|^{n-2s+1}} W_m^{\frac{4s}{n-2s}}(y + x_2) \\ &\quad \times \gamma(y + x_2) \sum_h \frac{1}{(1 + |y + x_2 - X^h|)^{\frac{n-2s}{2} + \tau}} dy \\ &=: C|x_1 - x_2| (M_1 + M_2). \end{aligned}$$

Since it holds that  $|t(x_1 - x_2) - y| \in [2|x_1 - x_2|, \frac{4}{3})$  for  $1 > |y| \ge 3|x_1 - x_2|$ , we get

$$|M_1| \le \frac{1}{\lambda^{\tau-s}} \int_{2|x_1-x_2| \le |y| \le \frac{4}{3}} \frac{1}{|y|^{n-2s+1}} \le \frac{1}{\lambda^{\tau-s}} (C+C|x_1-x_2|^{2s-1}).$$

For  $|y| \ge 1$ , we have  $\frac{2}{3}|y| \le |t(x_1 - x_2) - y| \le \frac{4}{3}|y|$ . Lemma A.5 yields

$$M_{2} \leq C \int_{|y|\geq 1} \frac{1}{|y|^{n-2s+1}} W_{m}^{\frac{4s}{n-2s}}(y+x_{2})\gamma(y+x_{2}) \sum_{h} \frac{1}{(1+|y+x_{2}-X^{h}|)^{\frac{n-2s}{2}+\tau}} dy$$
  
$$\leq C \int_{\mathbb{R}^{n}} \frac{1}{|y-x_{2}|^{n-2s}} W_{m}^{\frac{4s}{n-2s}}(y)\gamma(y) \sum_{h} \frac{1}{(1+|y-X^{h}|)^{\frac{n-2s}{2}+\tau}} dy$$
  
$$\leq \frac{C}{\lambda^{\tau-s}}.$$

Hence  $|H_1| \leq \frac{C}{\lambda^{\tau-s}} \|\phi\|_* |x_1 - x_2|^{\alpha}$ . The same procedure with the help of Lemma A.2 yields that  $|H_2| \leq \frac{C}{\lambda^{\tau-s}} \|g\|_{**} |x_1 - x_2|^{\alpha}$ . Then Lemma A.8 follows.

Appendix B. Expansions of the functionals  $\frac{\partial}{\partial \Lambda_i} I(W_m)$  and  $\frac{\partial}{\partial P_i^i} I(W_m)$ 

In this section, we will expand the functionals  $\frac{\partial}{\partial \Lambda_i} I(W_m)$  and  $\frac{\partial}{\partial P_j^i} I(W_m)$ . A direct computation yields

$$\frac{\partial I}{\partial \Lambda_{i}}(W_{m}) = \int_{\mathbb{R}^{n}} W_{m}(-\Delta)^{s} \frac{\partial W_{m}}{\partial \Lambda_{i}} - \int_{\mathbb{R}^{n}} K\left(\frac{x}{\lambda}\right) W_{m}^{\frac{n+2s}{n-2s}} \frac{\partial W_{m}}{\partial \Lambda_{i}} \\
= \int_{\mathbb{R}^{n}} \sum_{h=1}^{(m+1)^{k}} U_{P^{h},\Lambda_{h}}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial \Lambda_{i}} - \int_{\mathbb{R}^{n}} K\left(\frac{x}{\lambda}\right) W_{m}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial \Lambda_{i}}, \quad (B.1)$$

and

$$\frac{\partial I}{\partial P_j^i}(W_m) = \int_{\mathbb{R}^n} W_m(-\Delta)^s \frac{\partial W_m}{\partial P_j^i} - \int_{\mathbb{R}^n} K\left(\frac{x}{\lambda}\right) W_m^{\frac{n+2s}{n-2s}} \frac{\partial W_m}{\partial P_j^i} \\
= \int_{\mathbb{R}^n} \sum_{h \neq i} U_{P^h,\Lambda_h}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial P_j^i} - \int_{\mathbb{R}^n} K\left(\frac{x}{\lambda}\right) W_m^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial P_j^i}.$$
(B.2)

In order to get the useful expansions, we need to estimate each term on the right hand side of (B.1) and (B.2) above.

# Lemma B.1. There holds

$$\begin{split} \int_{\mathbb{R}^n} K\left(\frac{x}{\lambda}\right) W_m^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} &= \int_{\mathbb{R}^n} K\left(\frac{x}{\lambda}\right) \sum_h U_{P^h,\Lambda_h}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} \\ &+ \frac{n+2s}{n-2s} \int_{\mathbb{R}^n} K\left(\frac{x}{\lambda}\right) U_{P^i,\Lambda_i}^{\frac{4s}{n-2s}} \sum_{h\neq i} U_{P^h,\Lambda_h} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} + O\left((\lambda l)^{-n}\right). \end{split}$$

*Proof.* We estimate the integration on different region. By the same method used in (A.5), we have

$$\int_{\bigcup_h (\Omega_h \cap B_h^c)} K(\frac{x}{\lambda}) W_m^{\frac{n+2s}{n-2s}} \Big| \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} \Big| = O\left( (\lambda l)^{-n} \right).$$
(B.3)

In the domain  $\Omega_j \cap B_j$ , where  $j \neq i$ , there holds  $\hat{W}_{m,j}(y) \leq \sum_{h \neq j} \frac{C}{|X^j - X^h|} \leq \frac{C}{(\lambda l)^{n-2s}} \leq CU_{P^j,\Lambda_j}$ . Taylor expansion yields

$$\int_{\Omega_{j}\cap B_{j}} K\left(\frac{x}{\lambda}\right) W_{m}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial \Lambda_{i}}$$

$$= \int_{\Omega_{j}\cap B_{j}} K\left(\frac{x}{\lambda}\right) U_{P^{j},\Lambda_{j}}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial \Lambda_{i}} + O\left(\int_{\Omega_{j}\cap B_{j}} U_{P^{j},\Lambda_{j}}^{\frac{4s}{n-2s}} \hat{W}_{m,j} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial \Lambda_{i}}\right).$$

For the error term, a direct computation yields

$$\int_{\Omega_j \cap B_j} U_{P^j, \Lambda_j}^{\frac{4s}{n-2s}} \hat{W}_{m,j} \frac{\partial U_{P^i, \Lambda_i}}{\partial \Lambda_i} = O\left(\frac{1}{(\lambda l)^{2s} |X^i - X^j|^{n-2s}}\right).$$

**Claim:** For  $j \neq i$ , there holds

$$\int_{\Omega_j \cap B_j} K\left(\frac{x}{\lambda}\right) U_{P^j,\Lambda_j}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} = \int_{\mathbb{R}^n} K\left(\frac{x}{\lambda}\right) U_{P^j,\Lambda_j}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} + O\left(\frac{1}{(\lambda l)^{2s} |X^i - X^j|^{n-2s}}\right).$$
(B.4)

From direct computation

$$\int_{\Omega_i \cap B_i} K(\frac{x}{\lambda}) U_{P^j, \Lambda_j}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i, \Lambda_i}}{\partial \Lambda_i} = O\left(\frac{(\lambda l)^{2s}}{|X^i - X^j|^{n+2s}}\right).$$
(B.5)

Using Lemma A.1, we can obtain

$$\sum_{h \neq i,j} \int_{\Omega_h \cap B_h} K(\frac{x}{\lambda}) U_{P^j,\Lambda_j}^{\frac{n+2s}{n-2s}} \left| \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} \right| \leq \sum_{h \neq i,j} \frac{C}{|X^i - X^j|^{n-2s}} \int_{\Omega_h \cap B_h} \frac{1}{(1+|x-X^i|)^{n+2s}} \\ \leq \frac{C}{|X^i - X^j|^{n-2s}} \sum_{h \neq i} \frac{(\lambda l)^n}{|X^h - X^i|^{n+2s}} \qquad (B.6)$$

$$\leq \frac{C}{(\lambda l)^{2s} |X^i - X^j|^{n-2s}},$$

and

$$\int_{\bigcup_h (\Omega_h \cap B_h^c)} K(\frac{x}{\lambda}) U_{P^j, \Lambda_j}^{\frac{n+2s}{n-2s}} \Big| \frac{\partial U_{P^i, \Lambda_i}}{\partial \Lambda_i} \Big| = O\left(\frac{1}{(\lambda l)^{2s} |X^i - X^j|^{n-2s}}\right).$$
(B.7)

From (B.5), (B.6) and (B.7), we know the Claim is true.

Hence for  $j \neq i$ ,

$$\int_{\Omega_j \cap B_j} K\left(\frac{x}{\lambda}\right) W_m^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} = \int_{\mathbb{R}^n} K\left(\frac{x}{\lambda}\right) U_{P^j,\Lambda_j}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} + O\left(\frac{1}{(\lambda l)^{2s} |X^i - X^j|^{n-2s}}\right).$$
(B.8)

Now we estimate the integration on  $\Omega_i \cap B_i$ . By Taylor expansion,

$$\int_{\Omega_{i}\cap B_{i}} K\left(\frac{x}{\lambda}\right) W_{m}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial\Lambda_{i}} \\
= \int_{\Omega_{i}\cap B_{i}} K\left(\frac{x}{\lambda}\right) U_{P^{i},\Lambda_{i}}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial\Lambda_{i}} + \frac{n+2s}{n-2s} \int_{\Omega_{i}\cap B_{i}} K\left(\frac{x}{\lambda}\right) U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}} \sum_{h\neq i} U_{P^{h},\Lambda_{h}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial\Lambda_{i}} \\
+ O\left(\int_{\Omega_{i}\cap B_{i}} K\left(\frac{x}{\lambda}\right) U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}} \hat{W}_{m,i}^{2}\right) \\
= \int_{\Omega_{i}\cap B_{i}} K\left(\frac{x}{\lambda}\right) U_{P^{i},\Lambda_{i}}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial\Lambda_{i}} + \frac{n+2s}{n-2s} \int_{\Omega_{i}\cap B_{i}} K\left(\frac{x}{\lambda}\right) U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}} \sum_{h\neq i} U_{P^{h},\Lambda_{h}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial\Lambda_{i}} \\
+ O\left((\lambda l)^{-n}\right). \tag{B.9}$$

Since in the domain  $\Omega_i^c \cup B_i^c$ , we have  $|y - X^i| \ge \min\{\lambda l, \min_{j \ne i} \frac{1}{2}|X^i - X^j|\} \ge \frac{1}{2}\lambda l$ . Then

$$\int_{\Omega_i \cap B_i} K\left(\frac{x}{\lambda}\right) U_{P^i,\Lambda_i}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} = \int_{\mathbb{R}^n} K\left(\frac{x}{\lambda}\right) U_{P^i,\Lambda_i}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} + O\left((\lambda l)^{-n}\right). \tag{B.10}$$

By a similar method used in the proof of (B.4), we get

$$\int_{\Omega_i \cap B_i} K\left(\frac{x}{\lambda}\right) U_{P^i,\Lambda_i}^{\frac{4s}{n-2s}} U_{P^j,\Lambda_j} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} = \int_{\mathbb{R}^n} K\left(\frac{x}{\lambda}\right) U_{P^i,\Lambda_i}^{\frac{4s}{n-2s}} U_{P^j,\Lambda_j} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} \qquad (B.11)$$

$$+ O\left(\frac{1}{(\lambda l)^{2s} |X^i - X^j|^{n-2s}}\right).$$

Substituting (B.10) and (B.11) into (B.9), we have

$$\int_{\Omega_{i}\cap B_{i}} K\left(\frac{x}{\lambda}\right) W_{n}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial \Lambda_{i}}$$

$$= \int_{\mathbb{R}^{n}} K\left(\frac{x}{\lambda}\right) U_{P^{i},\Lambda_{i}}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial \Lambda_{i}} + \frac{n+2s}{n-2s} \int_{\mathbb{R}^{n}} K\left(\frac{x}{\lambda}\right) U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}} \sum_{h\neq i} U_{P^{h},\Lambda_{h}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial \Lambda_{i}} \quad (B.12)$$

$$+ O\left((\lambda l)^{-n}\right).$$

Now Lemma B.1 follows from the estimate (B.3), (B.8) and (B.12).

**Lemma B.2.** For  $h \neq i$ , there holds

$$\int_{\mathbb{R}^n} K\left(\frac{x}{\lambda}\right) U_{P^h,\Lambda_h}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} = \int_{\mathbb{R}^n} U_{P^h,\Lambda_h}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} + O\left(\frac{1}{\lambda^{2s}|X^i - X^h|^{n-2s}}\right), \tag{B.13}$$

and

$$\frac{n+2s}{n-2s} \int_{\mathbb{R}^n} K\left(\frac{x}{\lambda}\right) U_{P^i,\Lambda_i}^{\frac{4s}{n-2s}} U_{P^h,\Lambda_h} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} = \int_{\mathbb{R}^n} U_{P^h,\Lambda_h}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} + O\left(\frac{1}{\lambda^{2s} |X^i - X^h|^{n-2s}}\right). \tag{B.14}$$

*Proof.* Notice the fact

$$\int_{\mathbb{R}^{n}} U_{P^{h},\Lambda_{h}}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial\Lambda_{i}} = \int_{\mathbb{R}^{n}} (-\Delta)^{s} U_{P^{h},\Lambda_{h}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial\Lambda_{i}} \\
= \int_{\mathbb{R}^{n}} U_{P^{h},\Lambda_{h}} (-\Delta)^{s} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial\Lambda_{i}} \\
= \frac{n+2s}{n-2s} \int_{\mathbb{R}^{n}} U_{P^{h},\Lambda_{h}} U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial\Lambda_{i}}.$$

So the proof of (B.14) and (B.13) are identical. We only give a proof of (B.13).

Choose  $\delta$  to be a fixed constant some enough. Since  $n > 4s > n + 2s - \beta$ , the condition  $(H_3)$  implies

$$\left| \int_{B_{\delta\lambda}(X^h)} \left| K\left(\frac{x}{\lambda}\right) - 1 \right| U_{P^h,\Lambda_h}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} \right|$$

$$\leq C \int_{B_{\delta\lambda}(X^h)} \frac{|x-X^h|^{\beta}}{\lambda^{\beta}} \frac{1}{(1+|x-P^h|)^{n+2s}} \frac{1}{(1+|x-P^i|)^{n-2s}}$$

$$\leq \frac{C}{\lambda^{\beta} |X^i - X^h|^{n-2s}} \int_{B_{\delta\lambda}(X^h)} \frac{|x-X^h|^{\beta}}{(1+|x-X^h|)^{n+2s}} \leq \frac{C}{\lambda^{2s} |X^i - X^h|^{n-2s}}.$$
(B.15)

A direct calculation yields

$$\left| \int_{B_{\delta\lambda}(X^i)} \left| K\left(\frac{x}{\lambda}\right) - 1 \right| U_{P^h,\Lambda_h}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} \right| \le \frac{C\lambda^{2s}}{|X^i - X^h|^{n+2s}} \le \frac{C}{\lambda^{2s}|X^i - X^h|^{n-2s}}.$$
 (B.16)

Using Lemma A.1, we have

$$\left| \int_{B_{\delta\lambda}^{c}(X^{h}) \cap B_{\delta\lambda}^{c}(X^{i})} \left| K\left(\frac{x}{\lambda}\right) - 1 \right| U_{P^{h},\Lambda_{h}}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial \Lambda_{i}} \right| \\
\leq C \int_{B_{\delta\lambda}^{c}(X^{h}) \cap B_{\delta\lambda}^{c}(X^{i})} \frac{1}{(1+|x-X^{h}|)^{n+2s}} \frac{1}{(1+|x-X^{i}|)^{n-2s}} \\
\leq \frac{C}{|X^{h}-X^{i}|^{n-2s}} \int_{B_{\delta\lambda}^{c}(X^{h})} \frac{1}{(1+|x-X^{h}|)^{n+2s}} \leq \frac{C}{\lambda^{2s}|X^{i}-X^{h}|^{n-2s}}. \quad (B.17)$$

Hence (B.13) follows from (B.15), (B.16) and (B.17).

Lemma B.3. We have

$$\int_{\mathbb{R}^{n}} K\left(\frac{x}{\lambda}\right) U_{P^{i},\Lambda_{i}}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial \Lambda_{i}} = -\frac{n-2s}{2n} \frac{\beta C_{0}(n,s)^{\frac{2n}{n-2s}}(\sum_{h}a_{h})}{\Lambda_{i}^{\beta+1}\lambda^{\beta}} \int_{\mathbb{R}^{n}} \frac{|x_{1}|^{\beta}}{(1+|x|^{2})^{n}} +O\left(\frac{|P^{i}-X^{i}|^{\min\{2,\beta-1\}}}{\lambda^{\beta}}\right) + o\left(\lambda^{-\beta}\right), \quad (B.18)$$

and

$$\int_{\mathbb{R}^{n}} K\left(\frac{x}{\lambda}\right) U_{P^{i},\Lambda_{i}}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial P_{j}^{i}} = \frac{(n-2s)C_{0}(n,s)^{\frac{2n}{n-2s}}\beta a_{j}}{\Lambda_{i}^{\beta-2}\lambda^{\beta}} \int_{\mathbb{R}^{n}} \frac{|x_{1}|^{\beta}}{(1+|x|^{2})^{n+1}} (P_{j}^{i}-X_{j}^{i}) + O\left(\frac{|P^{i}-X^{i}|^{2}}{\lambda^{\beta}}\right) + O\left(\lambda^{-\beta}\right),$$
(B.19)

where  $i = 1, ..., (m+1)^k$  and j = 1, ..., n.

*Proof.* The two formulas follows from some standard calculations, see [21, Lemma A.9, Lemma A.10] for details.  $\hfill \square$ 

**Lemma B.4.** For  $h \neq i$ , we have

$$\int_{\mathbb{R}^n} U_{P^h,\Lambda_h}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^i,\Lambda_i}}{\partial \Lambda_i} = c_0 \frac{\partial \varepsilon_{ih}}{\partial \Lambda_i} + \frac{1}{\Lambda_i} O(\varepsilon_{hi}^{\frac{n}{n-2s}} \log \varepsilon_{hi}),$$
  
where  $c_0 = C_0(n,s)^{\frac{2n}{n-2s}} \int_{\mathbb{R}^n} \frac{1}{(1+|y|^2)^{\frac{n+2s}{2}}}$  and  $\varepsilon_{ih} = \left(\frac{1}{\frac{\Lambda_i}{\Lambda_h} + \frac{\Lambda_h}{\Lambda_i} + \Lambda_i\Lambda_h |P^i - P^h|^2}\right)^{\frac{n-2s}{2}}$ 

*Proof.* The proof of this lemma is rather standard. We refer to [3] and [9] for ideas.

### **Proposition B.5.** It holds that

$$\begin{aligned} \frac{\partial I}{\partial \Lambda_i}(W_m) &= -\frac{c_1}{\Lambda_i^{\beta+1}\lambda^{\beta}} + \sum_{h \neq i} \frac{c_2}{\Lambda_i(\Lambda_i\Lambda_h)^{\frac{n-2s}{2}} |X^i - X^h|^{n-2s}} + O\left(\frac{|P^i - X^i|^{\min\{2,\beta-1\}}}{\lambda^{\beta}}\right) + o(\lambda^{-\beta}), \\ where \ c_1 &= \frac{(n-2s)\beta C_0(n,s)^{\frac{2n}{n-2s}}(-\sum_h a_h)}{2n} \int_{\mathbb{R}^n} \frac{|x_1|^{\beta}}{(1+|x|^2)^n} > 0 \ and \ c_2 &= \frac{n-2s}{2} C_0(n,s)^{\frac{2n}{n-2s}} \int_{\mathbb{R}^n} \frac{1}{(1+|y|^2)^{\frac{n+2s}{2}}} \\ \end{bmatrix}$$

*Proof.* This proposition is a consequence of Lemma B.1, (B.13), (B.14), (B.18), Lemma B.4 and the definition of  $\lambda$ . We need to remind that

$$\frac{\partial \varepsilon_{ih}}{\partial \Lambda_i} = -\frac{n-2s}{2\Lambda_i (\Lambda_i \Lambda_h)^{\frac{n-2s}{2}} |X^i - X^h|^{n-2s}} + O(\frac{1}{|X^i - X^h|^{n-2s+1}}),$$

which is directly from  $P_h \in B_{\frac{1}{2}}(X^h)$  and the definition of  $\{X^h\}_{h=1}^{(m+1)^k}$ .

Proposition B.6. We have

where  $c_3 = (r$ 

$$\frac{\partial I}{\partial P_j^i}(W_m) = -\frac{c_3 a_j}{\Lambda_i^{\beta-2} \lambda^{\beta}} (P_j^i - X_j^i) + O\left(\frac{|P^i - X^i|^2}{\lambda^{\beta}}\right) + o(\lambda^{-\beta}),$$
  
$$(n-2s)C_0(n,s)^{\frac{2n}{n-2s}} \beta \int \frac{|x_1|^{\beta}}{(1+|x|^2)^{n+1}}.$$

*Proof.* We need to estimate each term on the right hand side of the equality (B.2). By simple calculation, we have

$$\left| W_m^{\frac{n+2s}{n-2s}} - U_{P^i,\Lambda_i}^{\frac{n+2s}{n-2s}} - \frac{n+2s}{n-2s} U_{P^i,\Lambda_i}^{\frac{4s}{n-2s}} \sum_{h \neq i} U_{P^h,\Lambda_h} \right| \leq \begin{cases} \left( \sum_{h \neq i} U_{P^h,\Lambda_h} \right)^{\frac{n+2s}{n-2s}}, & \text{if } U_{P^i,\Lambda_i} \leq \sum_{h \neq i} U_{P^h,\Lambda_h}, \\ U_{P^i,\Lambda_i}^{\frac{6s-n}{n-2s}} \left( \sum_{h \neq i} U_{P^h,\Lambda_h} \right)^2, & \text{otherwise.} \end{cases}$$

Then we get

$$\int_{\mathbb{R}^{n}} K\left(\frac{x}{\lambda}\right) W_{m}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial P_{j}^{i}} = \int_{\mathbb{R}^{n}} K\left(\frac{x}{\lambda}\right) U_{P^{i},\Lambda_{i}}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial P_{j}^{i}} + \frac{n+2s}{n-2s} \int_{\mathbb{R}^{n}} K\left(\frac{x}{\lambda}\right) U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}} \sum_{h\neq i} U_{P^{h},\Lambda_{h}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial P_{j}^{i}} \qquad (B.20) + O\left(\int_{\mathbb{R}^{n}} (\sum_{h\neq i} U_{P^{h},\Lambda_{h}})^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial P_{j}^{i}}\right) + O\left(\int_{U_{P^{i},\Lambda_{i}} \geq \sum_{h\neq i} U_{P^{h},\Lambda_{h}}} U_{P^{i},\Lambda_{i}}^{\frac{6s-n}{n-2s}} (\sum_{h\neq i} U_{P^{h},\Lambda_{h}})^{2} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial P_{j}^{i}}\right).$$

We first estimate the error terms above. Since n > 2s + 2 > 4s, we get  $(n - s)\frac{n-2s}{n+2s} > \frac{n-2s}{2} > k$ . From Lemma A.1 we have

$$\int_{\mathbb{R}^{n}} (\sum_{h \neq i} U_{P^{h}, \Lambda_{h}})^{\frac{n+2s}{n-2s}} \left| \frac{\partial U_{P^{i}, \Lambda_{i}}}{\partial P_{j}^{i}} \right| \leq C \int_{\mathbb{R}^{n}} \left( \sum_{h \neq i} \frac{1}{(1+|y-X^{h}|)^{n-2s}} \right)^{\frac{n+2s}{n-2s}} \frac{1}{(1+|y-X^{i}|)^{n-2s+1}} \\
\leq C \int_{\mathbb{R}^{n}} \left( \sum_{h \neq i} \frac{1}{(1+|y-X^{h}|)^{n-2s}} \frac{1}{(1+|y-X^{i}|)^{(n-s)\frac{n-2s}{n+2s}}} \right)^{\frac{n+2s}{n-2s}} \\
\leq \left( \sum_{h \neq i} \frac{1}{|X^{h}-X^{i}|^{(n-s)\frac{n-2s}{n+2s}}} \right)^{\frac{n+2s}{n-2s}} \int_{\mathbb{R}^{n}} \frac{1}{(1+|y-X^{i}|)^{n+2s}} \\
\leq C(\lambda l)^{-(n-s)}. \qquad (B.21)$$

The similar argument yields

$$\int_{U_{P^{i},\Lambda_{i}} > \sum_{h \neq i} U_{P^{h},\Lambda_{h}}} U_{P^{i},\Lambda_{i}}^{\frac{6s-n}{n-2s}} (\sum_{h \neq i} U_{P^{h},\Lambda_{h}})^{2} \left| \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial P_{j}^{i}} \right| \leq \int_{\mathbb{R}^{n}} (\sum_{h \neq i} U_{P^{h},\Lambda_{h}})^{\frac{n}{n-2s}} U_{P^{i},\Lambda_{i}}^{\frac{2s}{n-2s}} \left| \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial P_{j}^{i}} \right| \leq C(\lambda l)^{-n}.$$
(B.22)

For  $h \neq i$ , we see that

$$\frac{n+2s}{n-2s} \int_{\mathbb{R}^{n}} K\left(\frac{x}{\lambda}\right) U_{P^{i},\Lambda_{i}}^{\frac{4s}{n-2s}} U_{P^{h},\Lambda_{h}} \frac{\partial U_{P^{i},\Lambda_{i}}}{\partial P_{j}^{i}}$$

$$= \frac{\partial}{\partial P_{j}^{i}} \int_{\mathbb{R}^{n}} K\left(\frac{x}{\lambda}\right) U_{P^{i},\Lambda_{i}}^{\frac{n+2s}{n-2s}} U_{P^{h},\Lambda_{h}}$$

$$= \frac{1}{\lambda} \int_{\mathbb{R}^{n}} \frac{\partial K}{\partial t_{j}} \left(\frac{x+P^{i}}{\lambda}\right) U_{0,\Lambda_{i}}^{\frac{n+2s}{n-2s}} U_{P^{h}-P^{i},\Lambda_{h}} - \int_{\mathbb{R}^{n}} K\left(\frac{x+P^{i}}{\lambda}\right) U_{0,\Lambda_{i}}^{\frac{n+2s}{n-2s}} \frac{\partial U_{P^{h}-P^{i},\Lambda_{h}}}{\partial P_{j}^{i}}$$

$$= O(\lambda^{-1} \frac{1}{|X^{i}-X^{h}|^{n-2s}}) + O(\frac{1}{|X^{i}-X^{h}|^{n-s}}).$$
(B.23)

The first part of (B.2) can be estimated as

$$\int_{\mathbb{R}^{n}} \sum_{h \neq i} U_{P^{h}, \Lambda_{h}}^{\frac{n+2s}{n-2s}} \left| \frac{\partial U_{P^{i}, \Lambda_{i}}}{\partial P_{j}^{i}} \right| \leq \sum_{h \neq i} \int_{\mathbb{R}^{n}} \frac{C}{(1+|x-X^{h}|)^{n+2s}} \frac{1}{(1+|x-X^{i}|)^{n-2s+1}} \\ \leq \sum_{h \neq i} \frac{C}{|X^{h}-X^{i}|^{n-s}} \int_{\mathbb{R}^{n}} \frac{1}{(1+|y-X^{h}|)^{n+s+1}} \\ \leq C(\lambda l)^{-(n-s)}. \tag{B.24}$$

Then the expansion of  $\frac{\partial I}{\partial P_j^i}(W_m)$  follows from (B.19), (B.20), (B.21), (B.22), (B.23) and (B.24).

#### References

- Wael Abdelhedi and Hichem Chtioui, On a Nirenberg-type problem involving the square root of the Laplacian, J. Funct. Anal. 265 (2013), no. 11, 2937–2955.
- Wael Abdelhedi, Hichem Chtioui, and Hichem Hajaiej, A complete study of the lack of compactness and existence results of a fractional Nirenberg equation via a flatness hypothesis, I, Anal. PDE 9 (2016), no. 6, 1285–1315.
- Abbas Bahri, Critical points at infinity in some variational problems, Pitman Research Notes in Mathematics Series, vol. 182, Longman Scientific & Technical, Harlow UK, 1989.
- Thoams P. Branson, Sharp inequalities, the functional determinant, and the complementary series, Trans. Amer. Math. Soc. 347 (1995), no. 10, 3671–3742.
- Luis Caffarelli and Luis Silvestre, An extension problem related to the fractional Laplacian, Comm. Partial Differential Equations 32 (2007), no. 7-9, 1245–1260.
- Sun-Yung Alice Chang and María del Mar González, Fractional Laplacian in conformal geometry, Adv. Math. 226 (2011), no. 2, 1410–1432.
- 7. Guoyuan Chen and Youquan Zheng, A perturbation result for the  $Q_{\gamma}$  curvature problem on  $\mathbb{S}^n$ , Nonlinear Anal. 97 (2014), 4–14.
- Wenxiong Chen, Congming Li, and Biao Ou, Classification of solutions for an integral equation, Comm. Pure Appl. Math. 59 (2006), no. 3, 330–343.
- Yan-Hong Chen, Chungen Liu, and Youquan Zheng, Existence results for the fractional Nirenberg problem, J. Funct. Anal. 270 (2016), no. 11, 4043–4086.
- Yan-Hong Chen and Youquan Zheng, Peak solutions for the fractional Nirenberg problem, Nonlinear Anal. 122 (2015), 100–124.
- Hichem Chtioui and Wael Abdelhedi, On a fractional Nirenberg problem on n-dimensional spheres: existence and multiplicity results, Bull. Sci. Math. 140 (2016), no. 6, 617–628.
- Juan Dávila, Manuel del Pino, Serena Dipierro, and Enrico Valdinoci, Concentration phenomena for the nonlocal Schrödinger equation with Dirichlet datum, Anal. PDE 8 (2015), no. 5, 1165–1235.
- Juan Dávila, Manuel del Pino, and Yannick Sire, Nondegeneracy of the bubble in the critical case for nonlocal equations, Proc. Amer. Math. Soc. 141 (2013), no. 11, 3865–3870.
- Manuel del Pino, Patricio Felmer, and Monica Musso, Two-bubble solutions in the super-critical Bahri-Coron's problem, Calc. Var. Partial Differential Equations 16 (2003), no. 2, 113–145.
- Eleonora Di Nezza, Giampiero Palatucci, and Enrico Valdinoci, Hitchhiker's guide to the fractional Sobolev spaces, Bull. Sci. Math. 136 (2012), no. 5, 521–573.
- C. Robin Graham, Ralph Jenne, Lionel J. Mason, and George A. J. Sparling, Conformally invariant powers of the Laplacian. I. Existence, J. London Math. Soc. (2) 46 (1992), no. 3, 557–565.
- C. Robin Graham and Maciej Zworski, Scattering matrix in conformal geometry, Invent. Math. 152 (2003), no. 1, 89–118.
- Tianling Jin, Yanyan Li, and Jingang Xiong, On a fractional Nirenberg problem, part I: blow up analysis and compactness of solutions, J. Eur. Math. Soc. 16 (2014), no. 6, 1111–1171.
- <u>On a fractional Nirenberg problem</u>, Part II: Existence of solutions, Int. Math. Res. Not. IMRN 2015 (2015), no. 6, 1555–1589.

- Yanyan Li, Remark on some conformally invariant integral equations: the method of moving spheres, J. Eur. Math. Soc. 6 (2004), no. 2, 153–180.
- 21. Yanyan Li, Juncheng Wei, and Haoyuan Xu, Multi-bump solutions of  $-\Delta u = K(x)u^{\frac{n+2}{n-2}}$  on lattice in  $\mathbb{R}^n$ , to appear in J. Reine Angew. Math., arXiv:1305.4698v2.
- Yanyan Li and Meijun Zhu, Uniqueness theorems through the method of moving spheres, Duke Math. J. 80 (1995), no. 2, 383–417.
- Elliott Lieb, Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities, Ann. of Math.(2) 118 (1983), no. 2, 349–374.
- 24. Chungen Liu and Qiang Ren, Infinitely many non-radial solutions for fractional Nirenberg problem, preprint, 2015.
- Zhongyuan Liu, Concentration of solutions for the fractional Nirenberg problem, Commun. Pure Appl. Anal. 15 (2016), no. 2, 563–576.
- 26. Wei Long and Jing Yang, Positive or sign-changing solutions for a critical semilinear nonlocal equation, Z. Angew. Math. Phys. 67 (2016), no. 3, Art. 45, 30 pp.
- 27. Stephen Mark Paneitz, A quartic conformally covariant differential operator for arbitrary pseudo-Riemannian manifolds, SIGMA Symmetry Integrability Geom. Methods Appl. 4 (2008), Paper 036, 3 pp.
- Lawrence J. Peterson, Conformally covariant pseudo-differential operators, Differential Geom. Appl. 13 (2000), no. 2, 197–211.
- Juncheng Wei and Shusen Yan, Infinitely many solutions for the prescribed scalar curvature problem on S<sup>N</sup>, J. Funct. Anal. 258 (2010), 3048–3081.

(Chungen Liu) School of Mathematical Sciences and LPMC, Nankai University, Tianjin 300071, P.R. China

*E-mail address*: liucg@nankai.edu.cn

(Qiang Ren) School of Mathematical Sciences and LPMC, Nankai University, Tianjin 300071, P.R. China

E-mail address: tjftp@mail.nankai.edu.cn