Local Energy Conservation Law for Spatially-Discretized Hamiltonian Vlasov-Maxwell System

Jianyuan Xiao,^{1,2} Hong Qin,^{1,3,*} Jian Liu,^{1,2} and Ruili Zhang^{1,2}

¹School of Nuclear Science and Technology and Department of Modern Physics,

University of Science and Technology of China, Hefei, Anhui 230026, China

²Key Laboratory of Geospace Environment, CAS, Hefei, Anhui 230026, China

³Plasma Physics Laboratory, Princeton University, Princeton, NJ 08543, U.S.

Abstract

Structure-preserving geometric algorithm for the Vlasov-Maxwell (VM) equations is currently an active research topic. We show that spatially-discretized Hamiltonian systems for the VM equations admit a local energy conservation law in space-time. This is accomplished by proving that for a general spatially-discretized system, a global conservation law always implies a discrete local conservation law in space-time when the algorithm is local. This general result demonstrates that Hamiltonian discretizations can preserve local conservation laws, in addition to the symplectic structure, both of which are the intrinsic physical properties of infinite dimensional Hamiltonian systems in physics.

PACS numbers: 02.60.Jh, 03.50.-z, 52.65.Rr, 52.25.Dg

^{*} corresponding author: hongqin@ustc.edu.cn

The dynamics of a collection of charged particles and electromagnetic fields is governed by the well-known Vlasov-Maxwell (VM) equations

$$\frac{\partial f_s}{\partial t} + \mathbf{v} \cdot \nabla f_s + \frac{q_s}{m_s} \left(\mathbf{E} + \mathbf{v} \times \mathbf{B} \right) \cdot \frac{\partial f_s}{\partial \mathbf{v}} = 0, \qquad (1)$$

$$\frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E} \,, \tag{2}$$

$$\frac{\partial \mathbf{E}}{\partial t} = \nabla \times \mathbf{B} - \sum_{s} \int \mathrm{d}\mathbf{v} q_{s} f_{s}\left(\mathbf{x}, \mathbf{v}, t\right) \mathbf{v}, \qquad (3)$$

where **E** and **B** are electromagnetic fields, f_s , m_s and q_s are the number density distribution function, mass and charge of the *s*'th particle, respectively. Here the permittivity and permeability are set to 1 for simple notation. This set of equations is also a Hamiltonian Partial Differential Equation (PDE) [1–3], which means that solutions of the equations conserve the symplectic structure [4, 5] and various types of invariants. As one of these invariants, the Hamiltonian itself is conserved, i.e.,

$$\frac{\partial}{\partial t} \int d\mathbf{x} \left(\frac{E^2 + B^2}{2} + \sum_s \int d\mathbf{v} \left(\frac{1}{2} m_s v^2 f_s \left(\mathbf{x}, \mathbf{v}; t \right) \right) \right) = 0 .$$
(4)

For the Vlasov-Maxwell equations and many other Hamiltonian PDE systems, there are also local conservation laws, which can be written in the following form

$$\frac{\partial p}{\partial t} + \nabla \cdot \mathbf{u} = 0 , \qquad (5)$$

where p is a scalar field or a component of a tensor or vector field, and **u** is the flux corresponding to p. As an important example, the local energy conservation law for the Vlasov-Maxwell system reads

$$\frac{\partial}{\partial t} \left(\frac{E^2 + B^2}{2} + \sum_s \int d\mathbf{v} \left(\frac{1}{2} m_s v^2 f_s \left(\mathbf{x}, \mathbf{v}; t \right) \right) \right) + \nabla \cdot \left(\mathbf{E} \times \mathbf{B} + \sum_s \int d\mathbf{v} \left(\frac{1}{2} m_s v^2 \mathbf{v} f_s \left(\mathbf{x}, \mathbf{v}; t \right) \right) \right) = 0 .$$
(6)

It can be verified directly by using the Vlasov-Maxwell equations (1)-(3). The conservation law can also be obtained by using Noether's theorem and the weak Euler-Lagrangian equations [6]. Local conservation laws are more fundamental and practical than global conservation laws, because we often consider systems without global conservations, for example, systems with open boundaries and particle sources. Nowadays, Particle-In-Cell (PIC) simulations [7, 8] are commonly used in the investigation of Vlasov-Maxwell systems [9–16], and advanced structure-preserving geometric algorithms based on variational or Hamiltonian discretization have been developed recently [17–32]. Some of these structure-preserving PIC schemes are able to bound the global energy errors for all simulation time-steps and are effective for solving multi-scale problems [19, 24, 32]. For these algorithms, the corresponding spatially-discretized systems conserve global energy as a consequence of time symmetry admitted by the system. However, as an essential physical property, the discrete local energy conservation has not yet been discussed. If a discrete system has a local conservation law, it implies that at every grid point there is a conservation law, which is much stronger than just one global conservation law for the entire system. There have been some investigations on the discrete local energy conservation law associated with Yee's FDTD schemes for Maxwell's equations [33, 34]. However, the local conservation law for discrete particle-field systems is still an unexplored topic. In this paper, we prove a discrete local energy conservation law for the spatially-discretized Vlasov-Maxwell system described in Ref. [24]. This is accomplished by proving in the Appendix a theorem stating that for a general spatially-discretized system, a global conservation law always implies a discrete local conservation law in space-time when the algorithm is local. Here, an algorithm is called local if the time-advance of a field at a grid point or a particle involving only its neighboring grid points and particles. With this theorem, for any geometric spatial-discretizations with local algorithms, we only need to search for global conservation laws and then the corresponding local conservation laws are automatically satisfied. This general result demonstrates that Hamiltonian discretizations can preserve local conservation laws in space-time, in addition to the symplectic structure, both of which are the intrinsic physical properties of many important infinite dimensional Hamiltonian systems in physics.

The idea of structure-preserving spatial discretization of the Vlasov-Maxwell equations can be traced back to Lewis [35] who proposed a spatial-discretized Lagrangian for Vlasov plasmas. Today, modern geometric discretizations for constructing PIC schemes are based on Discrete Exterior Calculus (DEC) [36, 37], interpolation forms [24, 38, 39] or Finite Element Exterior Calculus (FEEC) [40–42] to ensure the conservation of charge, the gauge invariance, and the symplectic structure [17, 18, 24, 28, 31, 32]. We start from the Lagrangian of the spatially-discretized Vlasov-Maxwell system in Ref. [24],

$$L_{sd} = \frac{1}{2} \left(\sum_{J} \left(-\dot{\mathbf{A}}_{J} - \sum_{I} \nabla_{\mathrm{d}JI} \phi_{I} \right)^{2} - \sum_{K} \left(\sum_{J} \mathrm{curl}_{\mathrm{d}KJ} \mathbf{A}_{J} \right)^{2} \right) \Delta V + \sum_{s} \left(\frac{1}{2} m_{s} \dot{\mathbf{x}}_{s}^{2} + q_{s} \left(\dot{\mathbf{x}}_{s} \cdot \sum_{J} W_{\sigma_{1J}} \left(\mathbf{x}_{s} \right) \mathbf{A}_{J} - \sum_{I} W_{\sigma_{0I}} \left(\mathbf{x}_{s} \right) \phi_{I} \right) \right) .$$
(7)

Here, integers I, J and K are indices of grid points in a cubic-mesh, and ∇_d , curl_d and div_d are the discrete gradient, curl and divergence operators defined as follows,

$$(\nabla_{d}\phi)_{i,j,k} = [\phi_{i+1,j,k} - \phi_{i,j,k}, \phi_{i,j+1,k} - \phi_{i,j,k}, \phi_{i,j,k+1} - \phi_{i,j,k}], \qquad (8)$$

$$(\operatorname{curl}_{d} \mathbf{A})_{i,j,k} = \begin{bmatrix} (A_{zi,j+1,k} - A_{zi,j,k}) - (A_{y_{i,j,k+1}} - A_{y_{i,j,k}}) \\ (A_{xi,j,k+1} - A_{xi,j,k}) - (A_{zi+1,j,k} - A_{zi,j,k}) \\ (A_{y_{i+1,j,k}} - A_{y_{i,j,k}}) - (A_{xi,j+1,k} - A_{xi,j,k}) \end{bmatrix} , \qquad (9)$$
$$(\operatorname{div}_{d} \mathbf{B})_{i,j,k} = (B_{xi+1,j,k} - B_{xi,j,k}) + (B_{y_{i,j+1,k}} - B_{y_{i,j,k}}) + (B_{zi,j,k+1} - B_{zi,j,k}) . \qquad (10)$$

These operators are local linear operators on the discrete fields ϕ_I , \mathbf{A}_J and \mathbf{B}_K . Functions $W_{\sigma_{0J}}$, $W_{\sigma_{1I}}$ and $W_{\sigma_{2K}}$ are interpolation functions (Whitney forms) for 0-forms, e.g., scalar potential, 1-forms, e.g., vector potential, and 2-forms, e.g., magnetic fields, respectively. These discrete operators and interpolation functions satisfy the following properties [24, 36, 38, 39],

$$\nabla \sum_{I} W_{\sigma_{0I}} \left(\mathbf{x} \right) \phi_{I} = \sum_{I,J} W_{\sigma_{1J}} \left(\mathbf{x} \right) \nabla_{\mathrm{d}JI} \phi_{I} , \qquad (11)$$

$$\nabla \times \sum_{J} W_{\sigma_{1J}} \left(\mathbf{x} \right) \mathbf{A}_{J} = \sum_{J,K} W_{\sigma_{2K}} \left(\mathbf{x} \right) \operatorname{curl}_{\mathrm{d}KJ} \mathbf{A}_{J} , \qquad (12)$$

$$\nabla \cdot \sum_{K} W_{\sigma_{2K}} \left(\mathbf{x} \right) \mathbf{B}_{K} = \sum_{K,L} W_{\sigma_{3L}} \left(\mathbf{x} \right) \operatorname{div}_{\mathrm{d}LK} \mathbf{B}_{K} .$$
(13)

Periodic boundary in all three directions are adopted to simplify the discussion. Because time t does not explicitly appear in the Lagrangian, the total energy

$$H_{sd} = \frac{\partial L_{sd}}{\partial \dot{q}} \dot{q}^T - L_{sd} \tag{14}$$

$$= \frac{1}{2}\Delta V \left(\sum_{J} \mathbf{E}_{J}^{2} + \sum_{K} \mathbf{B}_{K}\right) + \sum_{s} \frac{1}{2} m_{s} \dot{\mathbf{x}}_{s}^{2}$$
(15)

is conserved, i.e., $\dot{H}_{sd} = 0$. Here, q is the generalized coordinates, i.e., $q = [\mathbf{A}_J, \phi_I, \mathbf{x}_s]$, and

 \mathbf{E}_J and \mathbf{B}_K are discrete electromagnetic fields defied as

$$\mathbf{E}_J = -\dot{\mathbf{A}}_J - \sum_J \nabla_{\mathrm{d}JI} \phi_I \;, \tag{16}$$

$$\mathbf{B}_K = \sum_J \operatorname{curl}_{\mathrm{d}KJ} \mathbf{A}_J \ . \tag{17}$$

The corresponding Poisson bracket for this system is [24]

$$\{F,G\} = \frac{1}{\Delta V} \sum_{J} \left(\frac{\partial F}{\partial \mathbf{E}_{J}} \cdot \sum_{K} \frac{\partial G}{\partial \mathbf{B}_{K}} \operatorname{curl}_{\mathrm{d}KJ} - \sum_{K} \frac{\partial F}{\partial \mathbf{B}_{K}} \operatorname{curl}_{\mathrm{d}KJ} \cdot \frac{\partial G}{\partial \mathbf{E}_{J}} \right) + \sum_{s} \frac{1}{m_{s}} \left(\frac{\partial F}{\partial \mathbf{x}_{s}} \cdot \frac{\partial G}{\partial \dot{\mathbf{x}}_{s}} - \frac{\partial F}{\partial \dot{\mathbf{x}}_{s}} \cdot \frac{\partial G}{\partial \mathbf{x}_{s}} \right) + \sum_{s} \frac{q_{s}}{m_{s}\Delta V} \left(\frac{\partial F}{\partial \dot{\mathbf{x}}_{s}} \cdot \sum_{J} W_{\sigma_{1J}} \left(\mathbf{x}_{s} \right) \frac{\partial G}{\partial \mathbf{E}_{J}} - \frac{\partial G}{\partial \dot{\mathbf{x}}_{s}} \cdot \sum_{J} W_{\sigma_{1J}} \left(\mathbf{x}_{s} \right) \frac{\partial F}{\partial \mathbf{E}_{J}} \right) + \sum_{s} \frac{q_{s}}{m_{s}^{2}} \frac{\partial F}{\partial \dot{\mathbf{x}}_{s}} \cdot \sum_{J} W_{\sigma_{1J}} \left(\mathbf{x}_{s} \right) \frac{\partial G}{\partial \mathbf{E}_{J}} - \frac{\partial G}{\partial \dot{\mathbf{x}}_{s}} \cdot \sum_{J} W_{\sigma_{1J}} \left(\mathbf{x}_{s} \right) \frac{\partial F}{\partial \mathbf{E}_{J}} \right) + \sum_{s} \frac{q_{s}}{m_{s}^{2}} \frac{\partial F}{\partial \dot{\mathbf{x}}_{s}} \cdot \left[\sum_{K} W_{\sigma_{2K}} \left(\mathbf{x}_{s} \right) \mathbf{B}_{K} \right] \times \frac{\partial G}{\partial \dot{\mathbf{x}}_{s}} . \quad (18)$$

With this Poisson bracket, the time evolution of the system is

$$\dot{g} = \{g, H_{sd}\}$$
, (19)

where $g = [\mathbf{E}_J, \mathbf{B}_K, \mathbf{x}_s, \dot{\mathbf{x}}_s]$. Now we introduce the discrete local energy ϵ_I at the *I*'th grid,

$$\varepsilon_I = \sum_s \frac{1}{2} m_s \dot{\mathbf{x}}_s^2 W_{\sigma_0 I} \left(\mathbf{x}_s \right) + \mathbf{E}_I^2 + \mathbf{B}_I^2.$$
⁽²⁰⁾

The evolution of ε_I is

$$\dot{\varepsilon}_I = \{\varepsilon_I, H_{sd}\} \quad , \tag{21}$$

or more specifically,

$$\dot{\varepsilon}_{I} = \sum_{s} q_{s} \left(\dot{\mathbf{x}}_{s} \cdot \sum_{J'} \mathbf{E}_{J'} W_{\sigma_{1} J'} \left(\mathbf{x}_{s} \right) \right) W_{\sigma_{0} I} \left(\mathbf{x}_{s} \right) + \sum_{s} \frac{1}{2} m_{s} \dot{\mathbf{x}}_{s}^{2} \left(\dot{\mathbf{x}}_{s} \cdot \nabla W_{\sigma_{0} I} \left(\mathbf{x}_{s} \right) \right) + \mathbf{E}_{I} \cdot \sum_{K} \operatorname{curl}_{\mathrm{d} K I} \mathbf{B}_{K} - \mathbf{E}_{I} \cdot \sum_{s} q_{s} \dot{\mathbf{x}}_{s} W_{\sigma_{1} I} \left(\mathbf{x}_{s} \right) - \mathbf{B}_{I} \cdot \sum_{J} \operatorname{curl}_{\mathrm{d} I J} \mathbf{E}_{J} .$$

$$(22)$$

We will see that the right hand side of Eq. (22) can be written as a discrete divergence of a discrete vector field, which means that Eq. (22) is a discrete energy conservation law.

Let us divide the RHS of Eq. (22) into three terms,

$$\dot{\varepsilon}_I = T_1 + T_2 + T_3 , \qquad (23)$$

where

$$T_1 = \mathbf{E}_I \cdot \sum_K \operatorname{curl}_{\mathrm{d}KI} \mathbf{B}_K - \mathbf{B}_I \cdot \sum_J \operatorname{curl}_{\mathrm{d}IJ} \mathbf{E}_J , \qquad (24)$$

$$T_2 = \sum_s \frac{1}{2} m_s \dot{\mathbf{x}}_s^2 \left(\dot{\mathbf{x}}_s \cdot \nabla W_{\sigma_0 I} \left(\mathbf{x}_s \right) \right) , \qquad (25)$$

$$T_{3} = \sum_{s} q_{s} \left(\dot{\mathbf{x}}_{s} \cdot \sum_{J'} \mathbf{E}_{J'} W_{\sigma_{1} J'} \left(\mathbf{x}_{s} \right) \right) W_{\sigma_{0} I} \left(\mathbf{x}_{s} \right) - \mathbf{E}_{I} \cdot \sum_{s} q_{s} \dot{\mathbf{x}}_{s} W_{\sigma_{1} I} \left(\mathbf{x}_{s} \right) .$$
(26)

Firstly, for T_1 , we can check that this term can be written as a discrete divergence

$$T_1 = -\sum_K \operatorname{div}_{\mathrm{d}IK} (\mathbf{E} \times *\mathbf{B})_K , \qquad (27)$$

where $\mathbf{E}\times \ast \mathbf{B}$ is defined as

$$(\mathbf{E} \times *\mathbf{B})_{i,j,k} = \begin{bmatrix} E_{y_{i,j,k}} B_{z_{i-1,j,k}} - E_{z_{i,j,k}} B_{y_{i-1,j,k}} \\ E_{z_{i,j,k}} B_{x_{i,j-1,k}} - E_{x_{i,j,k}} B_{z_{i,j-1,k}} \\ E_{x_{i,j,k}} B_{y_{i,j,k-1}} - E_{y_{i,j,k}} B_{x_{i,j,k-1}} \end{bmatrix}^{T}.$$
(28)

Next, we investigate ${\cal T}_2$ which represents the energy flow of particles,

$$T_{2} = \sum_{s} \frac{1}{2} m_{s} \dot{\mathbf{x}}_{s}^{2} \left(\dot{\mathbf{x}}_{s} \cdot \nabla W_{\sigma_{0}I} \left(\mathbf{x}_{s} \right) \right)$$
$$= \sum_{s} \frac{1}{2} m_{s} \dot{\mathbf{x}}_{s}^{2} \left(\dot{\mathbf{x}}_{s} \cdot \sum_{J} W_{\sigma_{1}J} \left(\mathbf{x}_{s} \right) \nabla_{dJI} \right)$$
$$= \sum_{J} \nabla_{dJI} \sum_{s} \frac{1}{2} m_{s} \dot{\mathbf{x}}_{s}^{2} \dot{\mathbf{x}}_{s} \cdot W_{\sigma_{1}J} \left(\mathbf{x}_{s} \right)$$
$$= \sum_{J} \nabla_{dJI} \mathbf{S}_{J} = -\sum_{K} \operatorname{div}_{dIK} \left(* \mathbf{S} \right)_{K} , \qquad (29)$$

where

$$\mathbf{S}_{J} = \sum_{s} \frac{1}{2} m_{s} \dot{\mathbf{x}}_{s}^{2} \dot{\mathbf{x}}_{s} \cdot W_{\sigma_{1}J} \left(\mathbf{x}_{s} \right) , \qquad (30)$$

$$(*\mathbf{S})_{K} = [S_{xi-1,j,k}, S_{y_{i,j-1,k}}, S_{zi,j,k-1}] .$$
(31)

Thus this term is also a discrete divergence. Finally, let us look at T_3 , which appears only in the discrete particle-field system. It is

$$T_{3} = \sum_{s} q_{s} \left(\dot{\mathbf{x}}_{s} \cdot \sum_{J} \mathbf{E}_{J} W_{\sigma_{1}J} \left(\mathbf{x}_{s} \right) \right) W_{\sigma_{0}I} \left(\mathbf{x}_{s} \right) - \mathbf{E}_{I} \cdot \sum_{s} q_{s} \dot{\mathbf{x}}_{s} W_{\sigma_{1}I} \left(\mathbf{x}_{s} \right)$$
(32)

$$=\sum_{s} q_{s} \left(\sum_{J'} \dot{\mathbf{x}}_{s} \cdot \mathbf{E}_{J} W_{\sigma_{1}J} \left(\mathbf{x}_{s} \right) W_{\sigma_{0}I} \left(\mathbf{x}_{s} \right) - \mathbf{E}_{I} \cdot \dot{\mathbf{x}}_{s} W_{\sigma_{1}I} \left(\mathbf{x}_{s} \right) \right)$$
(33)

$$=\sum_{s} q_{s} F\left(s\right) \left(W_{\sigma_{0}I}\left(\mathbf{x}_{s}\right) - \frac{p\left(s,I\right)}{F\left(s\right)} \right) , \qquad (34)$$

where

$$F(s) = \sum_{I} p(s, I) , \qquad (35)$$

$$p(s, I) = \mathbf{E}_{I} \cdot \dot{\mathbf{x}}_{s} W_{\sigma_{1}I}(\mathbf{x}_{s}) \quad .$$
(36)

From the definition of $W_{\sigma_0 I}(\mathbf{x})$, we have $\sum_I W_{\sigma_0 I}(\mathbf{x}_s) = 1$. Therefore,

$$\sum_{I} \left(W_{\sigma_0 I} \left(\mathbf{x}_s \right) - \frac{p\left(s, I \right)}{F\left(s \right)} \right) = 0 , \qquad (37)$$

which means that the sum of T_3 over all spatial grid points vanishes. Now we invoke the two theorems approved in the Appendix, which states that a discrete sum-free field must be a discrete divergence field and that if the sum-free field is local, then the divergence field is local. Therefore, for each particle, T_3 can be expressed as discrete divergence of a discrete vector field \mathbf{G}_s ,

$$T_{3} = \sum_{s} q_{s} \sum_{J'} \operatorname{div}_{\mathrm{d}IJ'} \mathbf{G}_{sJ'} F\left(s\right) .$$
(38)

For a particular s, $W_{\sigma_0 I}(\mathbf{x}_s)$, $W_{\sigma_1 J}(\mathbf{x}_s)$ are local near \mathbf{x}_s , so is $W_{\sigma_0 I}(\mathbf{x}_s) - p(s, I) / F(s)$. Thus \mathbf{G}_s is also local near \mathbf{x}_s .

Finally, we obtain the local energy conservation law for the spatial-discretized Vlasov-Maxwell system.

$$\frac{\partial}{\partial t} \left(\sum_{s} \frac{1}{2} m_{s} \dot{\mathbf{x}}_{s}^{2} W_{\sigma_{0}I} \left(\mathbf{x}_{s} \right) + \mathbf{E}_{I}^{2} + \mathbf{B}_{I}^{2} \right) + \sum_{K} \operatorname{div}_{\mathrm{d}IK} \left((\mathbf{E} \times * \mathbf{B})_{K} + (*\mathbf{S})_{K} - \sum_{s} q_{s} \mathbf{G}_{sK} F\left(s\right) \right) = 0 .$$
(39)

In conclusion, we started from the Hamiltonian theory of the spatially-discretized Vlasov-Maxwell system, used the property of Whitney forms, and derived a discrete local energy conservation law. In practice the spatially-discretized Vlasov-Maxwell system also need a temporal discretize to become a numerical scheme, after which the total energy will not be an exact invariant. However if we apply a symplectic integrator to perform the temporal discretize, then the total energy error can be bounded within a small value for all simulation time-steps [43–46]. Investigation on discrete conservation laws for such systems are planned for future work.

Appendix A: a discrete sum-free field is a discrete divergence field

In this appendix, we prove the following two theorems.

Theorem 1. If a discrete scalar field R is sum-free, i.e.,

$$\sum_{I} R_{I} = 0 , \qquad (A1)$$

then it can be expressed as a discrete divergence of a discrete vector field G, i.e.,

$$R_I = \sum_J \operatorname{div}_{\mathrm{d}IJ} \mathbf{G}_J \ . \tag{A2}$$

The discrete field R in Theorem 1 can also be a function of the continuous spatial coordinate \mathbf{x} , such as the Whitney form $W_{\sigma_{0J}}(\mathbf{x})$. A discrete field or a field component $F(\mathbf{x})$ is called local, if for every \mathbf{x}_0 , there exists a positive constant C such that

$$F_I(\mathbf{x}_0) = 0, \forall I \in \{J | | |\mathbf{x}_J - \mathbf{x}_0| > C\}.$$
(A3)

Theorem 2. If a discrete sum-free scalar field R is local, then there exists a local discrete vector field \mathbf{G} such that Eq. (A2) holds.

These two theorems also imply that a (local) discrete sum-free vector field or tensor field can be expressed as a discrete divergence of a (local) tensor or high order tensor, because for each component of the vector field or tensor field, Theorems 1 and 2 apply. To prove these two theorems, we first prove the following three lemmas.

Lemma 1. The discrete scalar field $R_x(i', j', k')$ defied as

$$R_{xI}(i',j',k') = \begin{cases} 1, & \text{if } I = [i'+1,j',k'] ,\\ -1, & \text{if } I = [i',j',k'] ,\\ 0, & \text{otherwise }, \end{cases}$$
(A4)

can be expressed as a composition of div_d and a discrete vector field $\mathbf{G}_x(i',j',k')$,

$$R_{xI}(i',j',k') = \sum_{J} \operatorname{div}_{\mathrm{d}IJ} \mathbf{G}_{xJ}(i',j',k') .$$
(A5)

Proof. Let $\mathbf{G}_{x}(i',j',k')$ be

$$\mathbf{G}_{xJ}(i',j',k') = \begin{cases} [-1,0,0], \text{ if } J = [i'+1,j',k'], \\ [0,0,0], & \text{otherwise}. \end{cases}$$
(A6)

Then it is straightforward to verify that

$$R_{xI}(i',j',k') = \sum_{J} \operatorname{div}_{\mathrm{d}IJ} \mathbf{G}_{xJ}(i',j',k') .$$
(A7)

Using the similar technique, we can construct $\mathbf{G}_{y}(i', j', k')$ and $\mathbf{G}_{z}(i', j', k')$ for scalar fields $R_{y}(i', j', k')$ and $R_{z}(i', j', k')$ as well, i.e.,

$$R_{yI}(i',j',k') = \begin{cases} 1, & \text{if } I = [i',j'+1,k'], \\ -1, & \text{if } I = [i',j',k'], \\ 0, & \text{otherwise}, \end{cases}$$
(A8)
$$R_{zI}(i',j',k') = \begin{cases} 1, & \text{if } I = [i',j',k'+1], \\ -1, & \text{if } I = [i',j',k'], \\ 0, & \text{otherwise}, \end{cases}$$
(A9)

$$\mathbf{G}_{yJ}(i',j',k') = \begin{cases} [0,-1,0], \text{ if } J = [i',j'+1,k'], \\ [0,0,0], & \text{otherwise}, \end{cases}$$
(A10)

$$\mathbf{G}_{zJ}(i',j',k') = \begin{cases} [0,0,-1], \text{ if } J = [i',j',k'+1], \\ [0,0,0], & \text{otherwise}. \end{cases}$$
(A11)

Lemma 2. The discrete scalar field $R^{\dagger}(I_1, I_2)$ defined as

$$R_{I}^{\dagger}(I_{1}, I_{2}) = \begin{cases} -1, & \text{if } I = I_{1} ,\\ 1, & \text{if } I = I_{2} ,\\ 0, & \text{otherwise} \end{cases}$$
(A12)

can be written in the following form

$$R_{I}^{\dagger}(I_{1}, I_{2}) = \sum_{I' \in Y_{x}} a_{x} R_{xI}(I') + \sum_{I' \in Y_{y}} a_{y} R_{yI}(I') + \sum_{I' \in Y_{z}} a_{z} R_{zI}(I') , \qquad (A13)$$

where Y_x , Y_y and Y_z are some indices sets, and a_x , a_y and a_z are some integers.

Proof. Choose Y_x , Y_y and Y_z as

$$Y_x = \{ [i, j, k] | \min(i_1, i_2) \le i < \max(i_1, i_2), j = j_2, k = k_2 \} ,$$
 (A14)

$$Y_y = \{ [i, j, k] | i = i_1, \min(j_1, j_2) \le j < \max(j_1, j_2), k = k_2 \} ,$$
 (A15)

$$Y_z = \{ [i, j, k] | i = i_1, j = j_1, \min(k_1, k_2) \le k < \max(k_1, k_2) \} .$$
(A16)

Choose a_x , a_y and a_z as

$$a_x = \begin{cases} 1, & \text{if } i_1 < i_2 ,\\ -1, & \text{otherwise} , \end{cases}$$
(A17)

$$a_y = \begin{cases} 1, & \text{if } j_1 < j_2 ,\\ -1, & \text{otherwise }, \end{cases}$$
(A18)

$$a_z = \begin{cases} 1, & \text{if } k_1 < k_2 ,\\ -1, & \text{otherwise} . \end{cases}$$
(A19)

It is straightforward to verify that Eq. (A13) holds.

Lemma 3. A sum-free scalar field R can be written as

$$R_{I} = \sum_{[I',I'']\in Z} b_{[I',I'']} R_{I}^{\dagger} (I',I'') \quad , \tag{A20}$$

where Z is some index-pair set.

Proof. If $\forall I$ such that $R_I = 0$, then Z can be chosen as \emptyset . Otherwise let $Y_d = \{I_1, I_2, I_3, \dots\}$ is the index set such that if $I \in Y_d$, $R_I \neq 0$ and $I \notin Y_d$, $R_I = 0$. We can choose the index-pair set Z as

$$Z = \{ [I', I''] | I' = I_1, I'' \in Y_d \text{ and } I'' \neq I_1 \},$$
(A21)

where I_1 is one arbitrarily chosen element in Y_d , and the corresponding $b_{[I',I'']}$ is

$$b_{[I',I'']} = R_{I''}.$$
 (A22)

Using the fact that

$$R_{I_1} + \sum_{I'' \in Y_d \text{ and } I'' \neq I_1} R_{I''} = \sum_I R_I = 0 , \qquad (A23)$$

we have

$$R_{I_1} = -\sum_{I'' \in Y_d \text{ and } I'' \neq I_1} R_{I''} .$$
(A24)

Then it is straightforward to verify that Eq. (A20) holds.

Composing these three lemmas, we can see that the Theorems 1 and 2 are proved.

ACKNOWLEDGMENTS

This research is supported by National Magnetic Confinement Fusion Energy Research Project (2015GB111003, 2014GB124005), National Natural Science Foundation of China (NSFC-11575185, 11575186, 11305171), JSPS-NRF-NSFC A3 Foresight Program (NSFC-11261140328), Chinese Scholar Council (201506340103), Key Research Program of Frontier Sciences CAS (QYZDB-SSW-SYS004), and the GeoAlgorithmic Plasma Simulator (GAPS) Project.

- [1] P. J. Morrison, Physics Letters A 80, 383 (1980).
- [2] A. Weinstein and P. J. Morrison, Physics Letters A 86, 235 (1981).
- [3] J. E. Marsden and A. Weinstein, Physica D: Nonlinear Phenomena 4, 394 (1982).
- [4] J. E. Marsden, G. W. Patrick, and S. Shkoller, Communications in Mathematical Physics 199, 351 (1998).
- [5] J. E. Marsden and T. Ratiu, Introduction to mechanics and symmetry: a basic exposition of classical mechanical systems, Vol. 17 (Springer Science & Business Media, 2013).
- [6] H. Qin, J. W. Burby, and R. C. Davidson, Physical Review E 90, 043102 (2014).
- [7] C. K. Birdsall and A. B. Langdon, *Plasma Physics via Computer Simulation* (IOP Publishing, 1991) p. 293.
- [8] R. W. Hockney and J. W. Eastwood, *Computer Simulation Using Particles* (CRC Press, 1988).
- [9] J. Cary and I. Doxas, Journal of Computational Physics 107, 98 (1993).
- [10] C. Nieter and J. R. Cary, Journal of Computational Physics 196, 448 (2004).
- [11] N. Xiang and J. R. Cary, Phys. Rev. Lett. **100**, 085002 (2008).
- [12] K. Germaschewski, W. Fox, S. Abbott, N. Ahmadi, K. Maynard, L. Wang, H. Ruhl, and A. Bhattacharjee, Journal of Computational Physics **318**, 305 (2016).
- [13] C.-K. Huang, Y. Zeng, Y. Wang, M. Meyers, S. Yi, and B. Albright, Computer Physics Communications 207, 123 (2016).
- [14] J. Qiang, arXiv:1610.04763 (2016).
- [15] A. Oeftiger, in *Proceedings of HB2016* (Malmo, Sweden, 2016) p. 126.
- [16] T. Planche and P. M. Jung, arXiv:1603.02976 (2016).

- [17] J. Squire, H. Qin, and W. M. Tang, Geometric Integration Of The Vlasov-Maxwell System With A Variational Particle-in-cell Scheme, Tech. Rep. PPPL-4748 (Princeton Plasma Physics Laboratory, 2012).
- [18] J. Squire, H. Qin, and W. M. Tang, Physics of Plasmas (1994-present) 19, 084501 (2012).
- [19] J. Xiao, J. Liu, H. Qin, and Z. Yu, Phys. Plasmas 20, 102517 (2013).
- [20] M. Kraus, arXiv:1307.5665 (2013).
- [21] E. Evstatiev and B. Shadwick, Journal of Computational Physics 245, 376 (2013).
- [22] B. A. Shadwick, A. B. Stamm, and E. G. Evstatiev, Physics of Plasmas 21, 055708 (2014).
- [23] J. Xiao, J. Liu, H. Qin, Z. Yu, and N. Xiang, Physics of Plasmas (1994-present) 22, 092305 (2015).
- [24] J. Xiao, H. Qin, J. Liu, Y. He, R. Zhang, and Y. Sun, Physics of Plasmas (1994-present) 22, 112504 (2015).
- [25] N. Crouseilles, L. Einkemmer, and E. Faou, Journal of Computational Physics 283, 224 (2015).
- [26] H. Qin, Y. He, R. Zhang, J. Liu, J. Xiao, and Y. Wang, Journal of Computational Physics 297, 721 (2015).
- [27] Y. He, H. Qin, Y. Sun, J. Xiao, R. Zhang, and J. Liu, Physics of Plasmas (1994-present) 22, 124503 (2015).
- [28] Y. He, Y. Sun, H. Qin, and J. Liu, Physics of Plasmas (1994-present) 23, 092108 (2016).
- [29] H. Qin, J. Liu, J. Xiao, R. Zhang, Y. He, Y. Wang, Y. Sun, J. W. Burby, L. Ellison, and Y. Zhou, Nuclear Fusion 56, 014001 (2016).
- [30] S. D. Webb, Plasma Physics and Controlled Fusion 58, 034007 (2016).
- [31] M. Kraus, K. Kormann, P. J. Morrison, and E. Sonnendrücker, arXiv:1609.03053 (2016).
- [32] J. Xiao, H. Qin, P. J. Morrison, J. Liu, Z. Yu, R. Zhang, and Y. He, Physics of Plasmas (1994-present) 23, 112107 (2016).
- [33] W. Chew, Journal of Applied Physics **75**, 4843 (1994).
- [34] J. De Moerloose and D. De Zutter, Microwave and Optical Technology Letters 8, 257 (1995).
- [35] H. R. Lewis, Journal of Computational Physics 6, 136 (1970).
- [36] A. N. Hirani, Discrete Exterior Calculus, Ph.D. thesis, California Institute of Technology (2003).
- [37] M. Desbrun, A. N. Hirani, M. Leok, and J. E. Marsden, arXiv:math/0508341 (2005).

- [38] H. Whitney, Geometric Integration Theory (Princeton University Press, 1957).
- [39] M. Desbrun, E. Kanso, and Y. Tong, in *Discrete Differential Geometry* (Springer, 2008) pp. 287–324.
- [40] D. N. Arnold, R. S. Falk, and R. Winther, Acta numerica 15, 1 (2006).
- [41] D. Arnold, R. Falk, and R. Winther, Bulletin of the American mathematical society 47, 281 (2010).
- [42] P. Monk, *Finite element methods for Maxwell's equations* (Oxford University Press, 2003).
- [43] R. D. Ruth, IEEE Trans. Nucl. Sci **30**, 2669 (1983).
- [44] K. Feng, in the Proceedings of 1984 Beijing Symposium on Differential Geometry and Differential Equations, edited by K. Feng (Science Press, 1985) pp. 42–58.
- [45] K. Feng, J. Comput. Maths. 4, 279 (1986).
- [46] E. Hairer, C. Lubich, and G. Wanner, Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations (Springer, New York, 2002).