

ON WELL-POSEDNESS OF ERICKSEN-LESLIE'S PARABOLIC-HYPERBOLIC LIQUID CRYSTAL MODEL

NING JIANG AND YILONG LUO

ABSTRACT. We establish a local well-posedness of Ericksen-Leslie's parabolic-hyperbolic liquid crystal model under an assumption on the smallness of initial data and with constraints of the dissipation coefficients $\mu_4 > 2(\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6) > 0$. In particular, if the dissipation coefficients $\mu_1 = \mu_2 = \mu_3 = \mu_5 = \mu_6 = 0$, we construct a local well-posedness of Ericksen-Leslie's hyperbolic liquid crystal model with large initial data. With a further constrain on the coefficients of dissipation terms ($\mu_2 < \mu_3$), we prove a global classical solution with small initial data.

1. INTRODUCTION

In this paper, we consider the so-called Ericksen-Leslie's parabolic-hyperbolic system that models the hydrodynamics of nematic liquid crystals in dimension n ($n=2, 3$):

$$\begin{cases} \partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v} - \frac{1}{2} \mu_4 \Delta \mathbf{v} + \nabla p = -\operatorname{div}(\nabla \mathbf{d} \odot \nabla \mathbf{d}) + \mathcal{L}(\mathbf{d}, \mathbf{v}), \\ \operatorname{div} \mathbf{v} = 0, |\mathbf{d}| = 1, \\ \sigma \ddot{\mathbf{d}} = \Delta \mathbf{d} + \gamma \mathbf{d} + \lambda_1 (\dot{\mathbf{d}} - B \mathbf{d}) + \lambda_2 A \mathbf{d}, \end{cases} \quad (1.1)$$

on $\Omega \times \mathbb{R}^+ \subset \mathbb{R}^n \times \mathbb{R}^+$ ($n = 2, 3$), where

$$\begin{aligned} \mathcal{L}(\mathbf{d}, \mathbf{v})_i = \partial_j & \left[\mu_1 \mathbf{d}_k \mathbf{d}_p A_{kp} \mathbf{d}_i \mathbf{d}_j + \mu_2 \mathbf{d}_j (\dot{\mathbf{d}}_i + B_{ki} \mathbf{d}_k) \right. \\ & \left. + \mu_3 \mathbf{d}_i (\dot{\mathbf{d}}_j + B_{kj} \mathbf{d}_k) + \mu_5 \mathbf{d}_j \mathbf{d}_k A_{ki} + \mu_6 \mathbf{d}_i \mathbf{d}_k A_{kj} \right] \end{aligned}$$

for $1 \leq i, j \leq n$, and the Lagrangian multiplier γ is

$$\gamma = -|\dot{\mathbf{d}}|^2 + |\nabla \mathbf{d}|^2 + \lambda_1 \mathbf{d}^\top B \mathbf{d} - \lambda_2 \mathbf{d}^\top A \mathbf{d},$$

and $A_{ij} = \frac{1}{2}(\partial_i \mathbf{v}_j + \partial_j \mathbf{v}_i)$, $B_{ij} = \frac{1}{2}(\partial_j \mathbf{v}_i - \partial_i \mathbf{v}_j)$, and the coefficients $\mu_4 > 0$, $\mu_i \geq 0$ ($1 \leq i \leq 6, i \neq 4$), and $\lambda_1 = \mu_2 - \mu_3$, $\lambda_2 = \mu_5 - \mu_6$, and $\sigma \geq 0$.

Here the material derivatives $\dot{\mathbf{d}} = \partial_t \mathbf{d} + \mathbf{v} \cdot \nabla \mathbf{d}$ and $(\nabla \mathbf{d} \odot \nabla \mathbf{d})_{ij} = \partial_i \mathbf{d}_k \partial_j \mathbf{d}_k$, $\Omega = \mathbb{R}^n$ or Ω is a bounded domain in \mathbb{R}^n . The vector $\mathbf{v}(x, t) \in \mathbb{R}^n$ is the bulk velocity and $\mathbf{d}(x, t) \in \mathbb{R}^n$ is the direction field of the liquid molecule with $|\mathbf{d}| = 1$.

A simple consideration is that the dissipative term $\mathcal{L}(\mathbf{d}, \mathbf{v})$ vanishes. Namely, the coefficients $\mu_1, \mu_2, \mu_3, \mu_5, \mu_6$ of $\mathcal{L}(\mathbf{d}, \mathbf{v})$ are chosen as 0, which immediately implies $\lambda_1 = \lambda_2 = 0$. Consequently, the system (1.1) reduces to a model which is Navier-Stokes equations coupled with a wave map

$$\begin{cases} \partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v} + \nabla p = \frac{1}{2} \mu_4 \Delta \mathbf{v} - \operatorname{div}(\nabla \mathbf{d} \odot \nabla \mathbf{d}), \\ \operatorname{div} \mathbf{v} = 0, \\ \sigma \ddot{\mathbf{d}} = \Delta \mathbf{d} + (-|\dot{\mathbf{d}}|^2 + |\nabla \mathbf{d}|^2) \mathbf{d}. \end{cases} \quad (1.2)$$

If Ω is a bounded domain in \mathbb{R}^n , the boundary conditions are given as

$$\mathbf{v}(x, t) = 0, \text{ and } \frac{\partial \mathbf{d}}{\partial \nu}(x, t) = 0 \quad (1.3)$$

on $(x, t) \in \partial\Omega \times [0, +\infty)$, where ν is the external normal vector of the boundary $\partial\Omega$. We impose the initial data in Ω as following:

$$\mathbf{v}|_{t=0} = \mathbf{v}^{in}, \dot{\mathbf{d}}|_{t=0} = \dot{\mathbf{d}}^{in}, \mathbf{d}|_{t=0} = \mathbf{d}^{in}, \quad (1.4)$$

which satisfy the compatibilities $\mathbf{v}^{in}(x)|_{\partial\Omega} = 0$ and $\frac{\partial \mathbf{d}^{in}}{\partial \nu}(x)|_{\partial\Omega} = 0$ if Ω is a bounded domain.

1.1. $\sigma = 0, \lambda_1 = -1$, parabolic model. When the coefficients $\sigma = 0$ and $\lambda_1 = -1$ in the third equation of (1.1), the system reduces to the parabolic type equation, which is also called Ericksen-Leslie's system in the literatures. The static analogue of the parabolic Ericksen-Leslie's system is the so-called Oseen-Frank model, whose mathematical study was initiated from Hardt-Kinderlehrer-Lin [8]. Since then there have been many works in this direction. In particular, the existence and regularity or partial regularity of the approximation (usually Ginzburg-Landau approximation as in [16]) dynamical Ericksen-Leslie's system was started by the work of Lin and Liu in [16], [17] and [18].

For the simplest system preserving the basic energy law

$$\begin{cases} \partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v} + \nabla p = \Delta \mathbf{v} - \operatorname{div}(\nabla \mathbf{d} \odot \nabla \mathbf{d}), \\ \operatorname{div} \mathbf{v} = 0, \\ \partial_t \mathbf{d} + \mathbf{v} \cdot \nabla \mathbf{d} = \Delta \mathbf{d} + |\nabla \mathbf{d}|^2 \mathbf{d}, \quad |\mathbf{d}| = 1, \end{cases} \quad (1.5)$$

which can be obtained by neglecting the Leslie stress and specifying some elastic constants. In 2-D case, global weak solutions with at most a finite number of singular times was proved by Lin-Lin-Wang [15]. The uniqueness of weak solutions was later on justified by Lin-Wang [20] and Xu-Zhang [27]. Recently, Lin and Wang proved global existence of weak solution for 3-D case in [21].

For the more general parabolic Ericksen-Leslie's system, local well-posedness is proved by Wang-Zhang-Zhang in [26], and in [10] regularity and existence of global solutions in \mathbb{R}^2 was established by Huang-Lin-Wang. The existence and uniqueness of weak solutions, also in \mathbb{R}^2 was proved by Hong-Xin and Li-Titi-Xin in [9] [14] respectively. Similar result was also obtained by Wang-Wang in [24]. For more complete review of the works for the parabolic Ericksen-Leslie's system, please see the reference listed above.

1.2. $\sigma = 1$, parabolic-hyperbolic model. If $\sigma = 1$, (1.1) is a parabolic-hyperbolic system for which there is very few works comparing the corresponding parabolic model. The only notable exception might be for the most simplified model, say, in (1.2), taking $\mathbf{v} = \mathbf{0}$, the spacial dimension is 1. For this case, the system (1.2) can be reduced to a so-called nonlinear variational wave equation. Zhang and Zheng (later on with Bressan and others) studied systematically the dissipative and energy conservative solutions in series work starting from late 90's [28, 29, 30, 31, 32, 33, 34, 35, 1, 36, 37, 2].

For the multidimensional case, to our best acknowledgement, there was no mathematical work on the original parabolic-hyperbolic Ericksen-Leslie's system (1.1). Very recently, De Anna and Zarnescu [3] considered the inertial Qian-Sheng model of liquid crystals which couples a hyperbolic type equation involving a second order derivative with a forced incompressible Navier-Stokes equations. It is a system describing the hydrodynamics of nematic liquid crystals in the Q-tensor framework. They proved global well-posedness and twist-wave solutions. Furthermore, for the inviscid version of the Qian-Sheng model, in [7], Feireisl-Rocca-Schimperna-Zarnescu proved a global existence of the *dissipative solution* which is inspired from that of incompressible Euler equation defined by P-L. Lions [22].

1.3. Ericksen-Leslie's parabolic-hyperbolic liquid crystal model. From now on, we always take $\sigma = 1$ in this paper. Leslie [12, 13] obtained a general parabolic-hyperbolic liquid crystal system (see also Section 5.1 of [19]). This system can be written in the following form:

$$\begin{cases} \partial_t \mathbf{v}_i + \mathbf{v}_k \partial_k \mathbf{v}_i = \partial_j (-p \delta_{ij} + \sigma_{ji}) + \mathbf{F}_i, \\ \partial_i \mathbf{v}_i = 0, \\ \ddot{\mathbf{d}}_i = \gamma \mathbf{d}_i + \mathbf{g}_i + \partial_j \pi_{ji} + \mathbf{G}_i. \end{cases} \quad (1.6)$$

Here $\mathbf{v} = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ is the bulk velocity, $\mathbf{d} = (\mathbf{d}_1, \dots, \mathbf{d}_n)$ is the direction field of the liquid molecules with the constraint $|\mathbf{d}| = 1$, p is the pressure, \mathbf{F}_i , \mathbf{G}_i are external forces and γ is

a Lagrange multiplier due to the constraint $|\mathbf{d}| = 1$. The extra stress σ_{ji} , the extra director stress π_{ji} and the extra intrinsic director body force \mathbf{g}_i read as

$$\begin{cases} \sigma_{ji} = -\frac{\partial W}{\partial(\partial_j \mathbf{d}_k)} \partial_i \mathbf{d}_k + \hat{\sigma}'_{ji}, \\ \pi_{ji} = \frac{\partial W}{\partial(\partial_j \mathbf{d}_i)}, \\ \mathbf{g}_i = -\frac{\partial W}{\partial \mathbf{d}_i} + \hat{\mathbf{g}}'_i, \end{cases} \quad (1.7)$$

respectively. Here the Oseen-Frank energy W can be written as

$$2W = k_1(\operatorname{div} \mathbf{d})^2 + k_2|\mathbf{d} \cdot (\nabla \times \mathbf{d})|^2 + k_3|\mathbf{d} \times (\nabla \times \mathbf{d})|^2 + (k_2 + k_4)[\operatorname{tr}(\nabla \mathbf{d})^2 - (\operatorname{div} \mathbf{d})^2], \quad (1.8)$$

where the coefficients k_1 , k_2 , k_3 , and k_4 are the measure of viscosity, depending on the material and the temperature. The dissipative term $\hat{\sigma}_{ji}$ is

$$\hat{\sigma}'_{ji} = \mu_1 \mathbf{d}_k \mathbf{d}_p A_{kp} \mathbf{d}_i \mathbf{d}_j + \mu_2 \mathbf{d}_j N_i + \mu_3 \mathbf{d}_i N_j + \mu_4 A_{ij} + \mu_5 \mathbf{d}_j \mathbf{d}_k A_{ki} + \mu_6 \mathbf{d}_i \mathbf{d}_k A_{kj}, \quad (1.9)$$

where $N_i = \dot{\mathbf{d}}_i + B_{ki} \mathbf{d}_k$ and the coefficients $\mu_i (1 \leq i \leq 6)$ may depend upon the temperature. The term $\hat{\mathbf{g}}'_i$ can be represented as

$$\hat{\mathbf{g}}'_i = \lambda_1 N_i + \lambda_2 \mathbf{d}_j A_{ij} \quad (1.10)$$

with the relations $\lambda_1 = \mu_2 - \mu_3$, $\lambda_2 = \mu_5 - \mu_6$.

The system (1.6) is derived from the conservation law of the form (proposed by Ericksen [4, 5, 6])

$$\frac{d}{dt} \int_{\Omega} (|\mathbf{v}|^2 + W + |\dot{\mathbf{d}}|^2) dx = - \int_{\Omega} \Delta \mathbf{d} + \text{boundary terms and harmless terms}.$$

Specifically, let $k_1 = k_2 = k_3 = 1$, $k_4 = 0$ in the equality (1.8) and \mathbf{F}_i , $\mathbf{G}_i = 0$ in (1.6), then

$$2W = |\mathbf{d} \cdot (\nabla \times \mathbf{d})|^2 + |\mathbf{d} \times (\nabla \times \mathbf{d})|^2 + \operatorname{tr}(\nabla \mathbf{d})^2.$$

Since $|\mathbf{d}| = 1$, one directly calculates that $2W = |\nabla \mathbf{d}|^2$, which implies

$$\frac{\partial W}{\partial \mathbf{d}_i} = 0, \quad \frac{\partial W}{\partial(\partial_j \mathbf{d}_i)} = \partial_j \mathbf{d}_i. \quad (1.11)$$

Then the equality (1.11) reduces to

$$\partial_j \pi_{ji} = \partial_j \left(\frac{\partial W}{\partial(\partial_j \mathbf{d}_i)} \right) = \partial_j (\partial_j \mathbf{d}_i) = \Delta \mathbf{d}_i, \quad (1.12)$$

and the relations (1.7), (1.10) and (1.11) imply that

$$\begin{aligned} \mathbf{g}_i &= \hat{\mathbf{g}}'_i \\ &= \lambda_1 N_i + \lambda_2 \mathbf{d}_j A_{ij} \\ &= \lambda_1 (\dot{\mathbf{d}} - B_{ik} \mathbf{d}_k) + \lambda_2 \mathbf{d}_j A_{ij} \\ &= \lambda_1 (\dot{\mathbf{d}} - (B \mathbf{d})_i) + \lambda_2 (A \mathbf{d})_i. \end{aligned} \quad (1.13)$$

Then it is derived from (1.12), (1.13) and the third equation of (1.6) with $\mathbf{F}_i = 0$ that

$$\sigma \ddot{\mathbf{d}} = \Delta \mathbf{d} + \gamma \mathbf{d} + \lambda_1 (\dot{\mathbf{d}} - B \mathbf{d}) + \lambda_2 A \mathbf{d}. \quad (1.14)$$

Since $|\mathbf{d}| = 1$, it is derived from multiplying \mathbf{d} in (1.14) that

$$\gamma = -|\dot{\mathbf{d}}|^2 + |\nabla \mathbf{d}|^2 + \lambda_1 \mathbf{d}^\top B \mathbf{d} - \lambda_2 \mathbf{d}^\top A \mathbf{d}. \quad (1.15)$$

Combining the first equality of (1.7) and (1.11), one can obtain that

$$\begin{aligned} \partial_j \sigma_{ji} &= -\partial_j (\partial_i \mathbf{d}_k \partial_j \mathbf{d}_k) + \partial_j \hat{\sigma}'_{ji} \\ &= -\operatorname{div}(\nabla \mathbf{d} \odot \nabla \mathbf{d})_i + \partial_j \hat{\sigma}'_{ji}, \end{aligned}$$

which means that the first equation of (1.6) can be rewritten as

$$\partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v} + \nabla p = -\operatorname{div}(\nabla \mathbf{d} \odot \nabla \mathbf{d}) + \operatorname{div} \hat{\sigma}'. \quad (1.16)$$

On the other hand, it can yield that by (1.9)

$$\operatorname{div} \hat{\sigma}' = \frac{1}{2} \mu_4 \Delta \mathbf{v} + \mathcal{L}(\mathbf{d}, \mathbf{v}),$$

hence, we get Leslie's parabolic-hyperbolic liquid crystal model (1.1).

1.4. Main results. In this paper, we first establish the local well-posedness of the Ericksen-Leslie's parabolic-hyperbolic liquid crystal system (1.1)-(1.3)-(1.4) with an assumption on smallness of the initial data \mathbf{d}^{in} and \mathbf{v}^{in} . However, if letting $\mu_1 = \mu_2 = \mu_3 = \mu_5 = \mu_6 = 0$, we can show the local well-posedness of the system (1.2)-(1.3)-(1.4) with large initial data \mathbf{v}^{in} and \mathbf{d}^{in} . Furthermore, if we give an additional assumption $\mu_2 < \mu_3$ (i.e. $\lambda_1 < 0$) on the model (1.1), we can prove the existence of a unique global smooth solution to the parabolic-hyperbolic liquid crystal system (1.1)-(1.3)-(1.4) with small initial data \mathbf{v}^{in} and \mathbf{d}^{in} .

Theorem 1.1. *For any integer $s > \frac{n}{2} + 2$ ($n = 2, 3$) and for given initial data \mathbf{v}^{in} , $\dot{\mathbf{d}}^{in} \in H^s(\Omega)$, $\mathbf{d}^{in} \in H^{s+1}(\Omega)$, the following statements are valid:*

(I) *If $\mu_4 > 2(\mu_2 + \mu_3 + \mu_5 + \mu_6)$, $\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6 > 0$ and the initial data satisfy $E^{in} \equiv |\mathbf{v}^{in}|_{H^s}^2 + |\dot{\mathbf{d}}^{in}|_{H^s}^2 + |\nabla \mathbf{d}^{in}|_{H^s}^2 \leq C_2 \equiv \frac{\mu_4}{4C(1+\mu_1+\mu_2+\mu_3+\mu_5+\mu_6)}$ for some positive constant $C = C(n, s) > 0$, then there exists a unique local solution $\mathbf{v} \in L^\infty(0, T; H^s(\Omega)) \cap L^2(0, T; H^{s+1}(\Omega))$, $\nabla \mathbf{d} \in L^\infty(0, T; H^s(\Omega))$ and $\dot{\mathbf{d}} \in L^\infty(0, T; H^s(\Omega))$ to the Ericksen-Leslie's parabolic-hyperbolic system (1.1)-(1.3)-(1.4), where T satisfies*

$$0 < T < \frac{1}{2|\lambda_1| + 1} \min \left\{ \ln \left(1 + \frac{2|\lambda_1| + 1}{C_1 E^{in}} \right), \ln \frac{\frac{4}{3} C_2 (C_1 E^{in} + 2|\lambda_1| + 1)}{(2|\lambda_1| + 1 + \frac{4}{3} C_1 C_2) E^{in}} \right\},$$

where $C_1 = C(1 + |\lambda_1| + |\lambda_2|) + \frac{C^2(1+|\lambda_1|+|\lambda_2|+\mu_2+\mu_3)^2}{2\mu_4} > 0$. Moreover, the solution (\mathbf{v}, \mathbf{d}) satisfies

$$(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^2}^2 + |\nabla \mathbf{d}|_{H^s}^2)(t) + \frac{1}{6} \mu_4 \int_0^t |\nabla \mathbf{v}|_{H^s}^2(s) ds \leq C^*(E^{in}, T)$$

for all $0 \leq t \leq T$, where

$$C^*(E^{in}, T) = E^{in} + [2|\lambda_1| C_3(E^{in}, T) + C_1(C_3(E^{in}, T))^2]T > 0$$

and

$$C_3(E^{in}, T) \equiv \frac{E^{in}(2|\lambda_1| + 1)e^{(2|\lambda_1|+1)T}}{C_1 E^{in} + 2|\lambda_1| + 1 - C_1 E^{in} e^{(2|\lambda_1|+1)T}} > 0.$$

(II) *If $\mu_1 = \mu_2 = \mu_3 = \mu_5 = \mu_6 = 0$ and the initial data satisfy $E^{in} < \infty$, then for some positive $C = C(n, s) > 0$ there exists a time number $0 < T < \frac{1}{C_4 E^{in}}$ such that there is a unique local solution $\mathbf{v} \in L^\infty(0, T; H^s(\Omega)) \cap L^2(0, T; H^{s+1}(\Omega))$, $\nabla \mathbf{d} \in L^\infty(0, T; H^s(\Omega))$ and $\dot{\mathbf{d}} \in L^\infty(0, T; H^s(\Omega))$ to the hyperbolic-type system (1.2)-(1.3)-(1.4). Moreover, the solution (\mathbf{v}, \mathbf{d}) satisfies*

$$(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2)(t) + \frac{1}{2} \mu_4 \int_0^t |\nabla \mathbf{v}|_{H^s}^2(s) ds \leq E^{in} + \frac{C_4(E^{in})^2 T}{(1 - C_4 E^{in} T)^2}$$

for all $0 \leq t \leq T$, where $C_4 = C(\frac{C}{2\mu_4} + 1) > 0$.

(III) *If $\mu_4 > 2(\mu_2 + \mu_3 + \mu_5 + \mu_6)$, $\mu_2 < \mu_3$ (i.e. $\lambda_1 < 0$) and the initial data satisfy $\mathcal{E}^{in} \equiv |\mathbf{v}^{in}|_{H^s}^2 + |\mathbf{d}^{in}|_{H^s}^2 + |\dot{\mathbf{d}}^{in}|_{H^s}^2 + |\nabla \mathbf{d}^{in}|_{H^s}^2 \leq \frac{\delta(\delta|\lambda_1|+1-\delta)}{4C_5[2\delta(\delta|\lambda_1|+1-\delta)+\delta|\lambda_1|+1]}$, then there exists a unique global solution $\mathbf{v} \in L^\infty(0, \infty; H^s(\Omega)) \cap L^2(0, \infty; H^{s+1}(\Omega))$, $\nabla \mathbf{d} \in L^\infty(0, \infty; H^s(\Omega))$*

and $\dot{\mathbf{d}} \in L^\infty(0, \infty; H^s(\Omega))$ to the parabolic-hyperbolic system (1.1)-(1.3)-(1.4). Moreover, the solution (\mathbf{v}, \mathbf{d}) satisfies

$$\sup_{0 \leq t \leq T} \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) (t) + \int_0^T |\nabla \mathbf{v}|_{H^s}^2 dt \leq \frac{1}{4C_5 \min \left\{ \delta, \frac{4-3\delta}{16} \mu_4 \right\}}$$

for any fixed $T \in (0, +\infty)$, where

$$\begin{aligned} C_5 = & C \left(\frac{8}{4-3\delta} + \frac{1}{2(|\lambda_1|-\delta)} + \frac{1}{2\delta} \right) \left(\frac{1}{\delta|\lambda_1|+1-\delta} + \frac{1}{1-\delta} + 1 \right) \left[\delta \left(\frac{1+|\lambda_1|^2+|\lambda_2|^2}{\mu_4} + 1 + |\lambda_1| + |\lambda_2| \right) \right. \\ & \left. + \frac{C(1+|\lambda_1|+|\lambda_2|+\mu_2+\mu_3)^2}{2\mu_4} + 1 + |\lambda_1| + |\lambda_2| + \mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6 \right] > 0, \end{aligned}$$

and $\delta = \frac{1}{2} \min\{|\lambda_1|, 1\} \in (0, \frac{1}{2}]$ and the positive constant $C = C(n, s) > 0$.

2. APRIORI ESTIMATES

Lemma 2.1. Assume that (\mathbf{v}, \mathbf{d}) is a sufficiently smooth solution to Ericksen-Leslie's parabolic-hyperbolic model (1.1). Then for any integer $s > \frac{n}{2} + 2$ ($n = 2, 3$) and for some $C = C(n, s) > 0$ we have

$$\begin{aligned} & \frac{d}{dt} \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) + \mu_4 |\nabla \mathbf{v}|_{H^s}^2 - 2\lambda_1 |\dot{\mathbf{d}}|_{H^s}^2 + 2\mu_1 \sum_{k=0}^s |\mathbf{d}^\top (\nabla^{k+1} \mathbf{v}) \mathbf{d}|_{L^2}^2 \\ & \leq C(1 + |\lambda_1| + |\lambda_2| + \mu_2 + \mu_3) \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) |\nabla \mathbf{v}|_{H^s} \\ & + C(1 + |\lambda_1| + |\lambda_2|) \left(|\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right)^2 + C(\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6) |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2, \end{aligned} \quad (2.1)$$

where $\dot{H}^s = H^s - L^2$.

Remark 2.1. (1) If $\mu_1 = \mu_2 = \mu_3 = \mu_5 = \mu_6 = 0$, which immediately means that $\lambda_1 = \mu_2 - \mu_3 = 0$ and $\lambda_2 = \mu_5 - \mu_6 = 0$, then the apriori estimate (2.1) can be represented as

$$\begin{aligned} & \frac{d}{dt} \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) + \mu_4 |\nabla \mathbf{v}|_{H^s}^2 \\ & \leq C \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) |\nabla \mathbf{v}|_{H^s} + C \left(|\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right)^2, \end{aligned} \quad (2.2)$$

where the constant $C = C(n, s) > 0$ is mentioned in Lemma 2.1.

(2) If $\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6 > 0$, the last term $C(\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6) |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2$ in (2.1) must be cancelled by the left term $\mu_4 |\nabla \mathbf{v}|_{H^s}^2$ under the assumption that the last term $C(\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6) |\nabla \mathbf{d}|_{H^s}^2 < \mu_4$. In order to guarantee this assumption, we have to impose on the smallness of the initial data E^{in} .

Proof. L^2 -Estimates. Multiplying $\dot{\mathbf{d}}$ in the third equation of (1.1) and integrating on B by parts, on may obtain

$$\int_\Omega \ddot{\mathbf{d}} \cdot \dot{\mathbf{d}} dx = \int_\Omega [\Delta \mathbf{d} \dot{\mathbf{d}} + \gamma \mathbf{d} \cdot \dot{\mathbf{d}} + \lambda_1 (\dot{\mathbf{d}} - B \mathbf{d}) \cdot \dot{\mathbf{d}} + \lambda_2 A \mathbf{d} \cdot \dot{\mathbf{d}}] dx,$$

where

$$\begin{aligned} \text{Left} &= \int_\Omega (\partial_t \dot{\mathbf{d}} + \mathbf{v} \cdot \nabla \dot{\mathbf{d}}) \cdot \dot{\mathbf{d}} dx = \frac{1}{2} \frac{d}{dt} \int_\Omega |\dot{\mathbf{d}}|^2 dx + \int_\Omega (\mathbf{v} \cdot \nabla \dot{\mathbf{d}}) \cdot \dot{\mathbf{d}} dx \\ &= \frac{1}{2} \frac{d}{dt} |\dot{\mathbf{d}}|_{L^2}^2, \end{aligned}$$

and the facts $\operatorname{div} \mathbf{v} = 0$ and $|\mathbf{d}| = 1$ imply that

$$\begin{aligned} \text{Right} &= \int_\Omega \Delta \mathbf{d} (\partial_t \mathbf{d} + \mathbf{v} \cdot \nabla \mathbf{d}) dx + \int_\Omega \gamma \left(\frac{1}{2} (\partial_t + v \cdot \nabla) |\mathbf{d}|^2 \right) dx \\ &+ \lambda_1 |\dot{\mathbf{d}}|_{L^2}^2 - \lambda_1 \int_\Omega \mathbf{d}^\top B \mathbf{d} dx + \lambda_2 \int_\Omega \mathbf{d}^\top A \mathbf{d} dx \end{aligned}$$

$$\begin{aligned}
&= - \int_{\Omega} \nabla \mathbf{d} : \nabla (\partial_t \mathbf{d} + \mathbf{v} \cdot \nabla \mathbf{d}) dx + \lambda_1 |\dot{\mathbf{d}}|_{L^2}^2 - \lambda_1 \int_{\Omega} \mathbf{d}^\top B \mathbf{d} dx + \lambda_2 \int_{\Omega} \mathbf{d}^\top A \mathbf{d} dx \\
&= - \frac{1}{2} \frac{d}{dt} |\nabla \mathbf{d}|_{L^2}^2 + \int_{\Omega} \mathbf{v} \cdot \nabla \mathbf{d} \Delta \mathbf{d} dx + \lambda_1 |\dot{\mathbf{d}}|_{L^2}^2 - \lambda_1 \int_{\Omega} \mathbf{d}^\top B \mathbf{d} dx + \lambda_2 \int_{\Omega} \mathbf{d}^\top A \mathbf{d} dx \\
&\leq - \frac{1}{2} \frac{d}{dt} |\nabla \mathbf{d}|_{L^2}^2 + \int_{\Omega} \mathbf{v} \cdot \nabla \mathbf{d} \Delta \mathbf{d} dx + \lambda_1 |\dot{\mathbf{d}}|_{L^2}^2 + (|\lambda_1| + |\lambda_2|) |\mathbf{d}|_{L^\infty} |\dot{\mathbf{d}}|_{L^2} |\nabla \mathbf{v}|_{L^2}.
\end{aligned}$$

Hence the following relation holds:

$$\frac{1}{2} \frac{d}{dt} \left(|\dot{\mathbf{d}}|_{L^2}^2 + |\nabla \mathbf{d}|_{L^2}^2 \right) - \lambda_1 |\dot{\mathbf{d}}|_{L^2}^2 \leq \int_{\Omega} \mathbf{v} \cdot \nabla \mathbf{d} \Delta \mathbf{d} dx + (|\lambda_1| + |\lambda_2|) |\mathbf{d}|_{L^\infty} |\dot{\mathbf{d}}|_{L^2} |\nabla \mathbf{v}|_{L^2}. \quad (2.3)$$

Multiplying by \mathbf{v} in the first equation of (1.1) and integrating on B by parts, one can derive by the fact $\operatorname{div} \mathbf{v} = 0$

$$\begin{aligned}
\frac{1}{2} \frac{d}{dt} |\mathbf{v}|_{L^2}^2 + \mu |\nabla \mathbf{v}|_{L^2}^2 &= - \int_{\Omega} \mathbf{v}_i \partial_j (\partial_i \mathbf{d}_k \partial_j \mathbf{d}_k) dx + \int_{\Omega} \mathcal{L}(\mathbf{d}, \mathbf{v}) \cdot \mathbf{v} dx \\
&= - \int_{\Omega} \mathbf{v}_i \partial_j \partial_i \mathbf{d}_k \partial_j \mathbf{d}_k dx - \int_{\Omega} \mathbf{v}_i \partial_i \mathbf{d}_k \partial_j \partial_j \mathbf{d}_k dx + \int_{\Omega} \mathcal{L}(\mathbf{d}, \mathbf{v}) \cdot \mathbf{v} dx \quad (2.4) \\
&= - \int_{\Omega} \mathbf{v}_i \partial_i \left(\frac{1}{2} |\nabla \mathbf{d}|^2 \right) dx - \int_{\Omega} \mathbf{v} \cdot \nabla \mathbf{d} \cdot \Delta \mathbf{d} dx + \int_{\Omega} \mathcal{L}(\mathbf{d}, \mathbf{v}) \cdot \mathbf{v} dx \\
&= - \int_{\Omega} \mathbf{v} \cdot \nabla \mathbf{d} \cdot \Delta \mathbf{d} dx + \int_{\Omega} \mathcal{L}(\mathbf{d}, \mathbf{v}) \cdot \mathbf{v} dx.
\end{aligned}$$

Next we are going to estimate the term $\int_{\Omega} \mathcal{L}(\mathbf{d}, \mathbf{v}) \cdot \mathbf{v} dx$. It is calculated that by integrating by parts, Hölder inequality and the fact $\mathbf{d}^\top B \mathbf{d} = 0$

$$\mu_1 \int_{\Omega} \partial_j ((\mathbf{d}^\top A \mathbf{d}) \mathbf{d}_i \mathbf{d}_j) \mathbf{v}_i dx = -\mu_1 |\mathbf{d}^\top \nabla \mathbf{v} \mathbf{d}|_{L^2}^2,$$

and

$$\begin{aligned}
\mu_2 \int_{\Omega} \partial_j [\mathbf{d}_j (\dot{\mathbf{d}}_i + B_{ki} \mathbf{d}_k)] \mathbf{v}_i dx &= -\mu_2 \int_{\Omega} \mathbf{d}_j (\dot{\mathbf{d}}_i + B_{ki} \mathbf{d}_k) \partial_j \mathbf{v}_i dx \\
&\leq \mu_2 |\mathbf{d}|_{L^\infty} |\dot{\mathbf{d}}|_{L^2} |\nabla \mathbf{v}|_{L^2} + \mu_2 |\mathbf{d}|_{L^\infty}^2 |\nabla \mathbf{v}|_{L^2}^2,
\end{aligned}$$

and similarly

$$\mu_3 \int_{\Omega} \partial_j [\mathbf{d}_i (\dot{\mathbf{d}}_j + B_{kj} \mathbf{d}_k)] \mathbf{v}_i dx \leq \mu_3 |\mathbf{d}|_{L^\infty} |\dot{\mathbf{d}}|_{L^2} |\nabla \mathbf{v}|_{L^2} + \mu_3 |\mathbf{d}|_{L^\infty}^2 |\nabla \mathbf{v}|_{L^2}^2,$$

and

$$\begin{aligned}
\mu_5 \int_{\Omega} \partial_j (\mathbf{d}_j \mathbf{d}_k A_{ki}) \mathbf{v}_i dx &= -\mu_5 \int_{\Omega} \mathbf{d}_j \mathbf{d}_k A_{ki} \partial_j \mathbf{v}_i dx \\
&\leq \mu_5 |\mathbf{d}|_{L^\infty}^2 |\nabla \mathbf{v}|_{L^2}^2,
\end{aligned}$$

and similarly

$$\mu_6 \int_{\Omega} \partial_j (\mathbf{d}_i \mathbf{d}_k A_{kj}) \mathbf{v}_i dx \leq \mu_6 |\mathbf{d}|_{L^\infty}^2 |\nabla \mathbf{v}|_{L^2}^2.$$

Hence the estimate of the term $\int_{\Omega} \mathcal{L}(\mathbf{d}, \mathbf{v}) \cdot \mathbf{v} dx$ is

$$\begin{aligned}
\int_{\Omega} \mathcal{L}(\mathbf{d}, \mathbf{v}) \cdot \mathbf{v} dx &\leq -\mu_1 |\mathbf{d}^\top \nabla \mathbf{v} \mathbf{d}|_{L^2}^2 + (\mu_2 + \mu_3) |\mathbf{d}|_{L^\infty} |\dot{\mathbf{d}}|_{L^2} |\nabla \mathbf{v}|_{L^2} \\
&\quad + (\mu_2 + \mu_3 + \mu_5 + \mu_6) |\mathbf{d}|_{L^\infty}^2 |\nabla \mathbf{v}|_{L^2}^2. \quad (2.5)
\end{aligned}$$

It is implied that by the inequalities (2.3), (2.4) and (2.5)

$$\begin{aligned}
&\frac{1}{2} \frac{d}{dt} \left(|\mathbf{v}|_{L^2}^2 + |\dot{\mathbf{d}}|_{L^2}^2 + |\nabla \mathbf{d}|_{L^2}^2 \right) + \frac{1}{2} \mu_4 |\nabla \mathbf{v}|_{L^2}^2 - \lambda_1 |\dot{\mathbf{d}}|_{L^2}^2 + \mu_1 |\mathbf{d}^\top A \mathbf{d}|_{L^2}^2 \\
&\leq (|\lambda_1| + |\lambda_2| + \mu_2 + \mu_3) |\mathbf{d}|_{L^\infty} |\dot{\mathbf{d}}|_{L^2} |\nabla \mathbf{v}|_{L^2} + (\mu_2 + \mu_3 + \mu_5 + \mu_6) |\mathbf{d}|_{L^\infty}^2 |\nabla \mathbf{v}|_{L^2}^2. \quad (2.6)
\end{aligned}$$

H^s -Estimates. For any integer $1 \leq k \leq s$, we take ∇^k in the fist equation of (1.1), multiply by $\nabla^k \mathbf{v}$ and integrate by parts, then we have

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} |\nabla^k \mathbf{v}|_{L^2}^2 + \mu |\nabla^{k+1} \mathbf{v}|_{L^2}^2 \\ &= - \left\langle \nabla^k (\mathbf{v} \cdot \nabla \mathbf{v}), \nabla^k \mathbf{v} \right\rangle - \left\langle \nabla^k \operatorname{div}(\nabla \mathbf{d} \odot \nabla \mathbf{d}), \nabla^k \mathbf{v} \right\rangle + \left\langle \nabla^k \mathcal{L}(\mathbf{d}, \mathbf{v}), \nabla^k \mathbf{v} \right\rangle. \end{aligned} \quad (2.7)$$

It is derived that by Hölder inequality, Sobolev embedding and the fact $\operatorname{div} \mathbf{v} = 0$

$$\begin{aligned} - \left\langle \nabla^k (\mathbf{v} \cdot \nabla \mathbf{v}), \nabla^k \mathbf{v} \right\rangle &= - \sum_{\substack{a+b=k \\ a \geq 1}} \left\langle \nabla^a \mathbf{v} \nabla^{b+1} \mathbf{v}, \nabla^k \mathbf{v} \right\rangle \\ &= - 2 \left\langle \nabla \mathbf{v} \nabla^k \mathbf{v}, \nabla^k \mathbf{v} \right\rangle - \sum_{\substack{a+b=k \\ a \geq 2, b \geq 1}} \left\langle \nabla^a \mathbf{v} \nabla^{b+1} \mathbf{v}, \nabla^k \mathbf{v} \right\rangle \\ &\leq C |\nabla \mathbf{v}|_{L^\infty} |\nabla^k \mathbf{v}|_{L^2}^2 + C \sum_{\substack{a+b=k \\ a \geq 2, b \geq 1}} |\nabla^a \mathbf{v}|_{L^4} |\nabla^{b+1} \mathbf{v}|_{L^4} |\nabla^k \mathbf{v}|_{L^2} \\ &\leq C |\nabla^3 \mathbf{v}|_{L^2} |\nabla^k \mathbf{v}|_{L^2}^2 + C |\nabla^k \mathbf{v}|_{L^2}^3 \\ &\leq C |\nabla \mathbf{v}|_{H^s} |\mathbf{v}|_{\dot{H}^s}^2, \end{aligned} \quad (2.8)$$

where $\dot{H}^s = H^s - L^2$ and

$$\begin{aligned} & - \left\langle \nabla^k \operatorname{div}(\nabla \mathbf{d} \odot \nabla \mathbf{d}), \nabla^k \mathbf{v} \right\rangle \\ &= \left\langle \nabla^k (\nabla \mathbf{d} \odot \nabla \mathbf{d}), \nabla^{k+1} \mathbf{v} \right\rangle \\ &= \sum_{a+b=k} \left\langle \nabla^{a+1} \mathbf{d} \odot \nabla^{b+1} \mathbf{d}, \nabla^{k+1} \mathbf{v} \right\rangle \\ &\leq 2 |\nabla \mathbf{d}|_{L^\infty} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2} + \sum_{\substack{a+b=k \\ a, b \geq 1}} \left\langle \nabla^{a+1} \mathbf{d} \odot \nabla^{b+1} \mathbf{d}, \nabla^{k+1} \mathbf{v} \right\rangle \\ &\leq C |\nabla^3 \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2} + \sum_{\substack{a+b=k \\ a, b \geq 1}} |\nabla^{a+1} \mathbf{d}|_{L^4} |\nabla^{b+1} \mathbf{d}|_{L^4} |\nabla^{k+1} \mathbf{v}|_{L^2} \\ &\leq C |\nabla^3 \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2} + C |\nabla^{k+1} \mathbf{d}|_{L^2}^2 |\nabla^{k+1} \mathbf{v}|_{L^2} \\ &\leq C |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}. \end{aligned} \quad (2.9)$$

The term $\langle \nabla^k \mathcal{L}(\mathbf{d}, \mathbf{v}), \nabla^k \mathbf{v} \rangle$, which can be divided into five parts $(\mu_1, \mu_2, \mu_3, \mu_5, \mu_6)$, remains to be estimated. First, we estimate the μ_1 -part: $\langle \mu_1 \nabla^k \partial_j(\mathbf{d}^\top A \mathbf{d}_i \mathbf{d}_j), \nabla^k \mathbf{v}_i \rangle$.

$$\begin{aligned} & \left\langle \mu_1 \nabla^k \partial_j(\mathbf{d}^\top A \mathbf{d}_i \mathbf{d}_j), \nabla^k \mathbf{v}_i \right\rangle \\ &= -\mu_1 \left\langle \nabla^k (\mathbf{d}^\top A \mathbf{d}_i \mathbf{d}_j), \nabla^k \partial_j \mathbf{v}_i \right\rangle \\ &= -\mu_1 \left\langle \mathbf{d}^\top \nabla^{k+1} \mathbf{v} \mathbf{d}, \mathbf{d}^\top \nabla^{k+1} \mathbf{v} \mathbf{d} \right\rangle - \mu_1 \sum_{\substack{a+b=k \\ b \geq 1}} \left\langle \nabla^a (\mathbf{d}_p \mathbf{d}_q) \nabla^b \partial_p \mathbf{v}_q \mathbf{d}_i \mathbf{d}_j, \nabla^k \partial_j \mathbf{v}_i \right\rangle \\ &\quad - \mu_1 \sum_{\substack{a+b=k \\ b \geq 1}} \left\langle \mathbf{d}_p \mathbf{d}_q \nabla^a \partial_p \mathbf{v}_q \nabla^b (\mathbf{d}_i \mathbf{d}_j), \nabla^k \partial_j \mathbf{v}_i \right\rangle - \mu_1 \sum_{\substack{a_1+a_2+b=k \\ a_1, b \geq 1}} \left\langle \nabla^{a_1} (\mathbf{d}_p \mathbf{d}_q) \nabla^{a_2} \partial_p \mathbf{v}_q \nabla^b (\mathbf{d}_i \mathbf{d}_j), \nabla^k \partial_j \mathbf{v}_i \right\rangle \\ &\equiv -\mu_1 |\mathbf{d}^\top \nabla^{k+1} \mathbf{v} \mathbf{d}|_{L^2}^2 + I_1 + I_2 + I_3. \end{aligned}$$

According to Hölder inequality, Sobolev embedding and the fact $|\mathbf{d}| = 1$, we calculate that

$$\begin{aligned}
I_1 &\leq \mu_1 |\mathbf{d}|_{L^\infty}^2 |\nabla^{k+1} \mathbf{v}|_{L^2}^2 + \mu_1 \sum_{\substack{a_1+a_2+b=k \\ a_1+a_2 \geq 1, b \geq 1}} \left\langle |\nabla^{a_1} \mathbf{d}_p \nabla^{a_2} \mathbf{d}_q \nabla^b \partial_p \mathbf{v}_q|, |\nabla^{k+1} \mathbf{v}| \right\rangle \\
&\leq \mu_1 |\mathbf{d}|_{L^\infty}^2 |\nabla^{k+1} \mathbf{v}|_{L^2}^2 + \mu_1 \sum_{\substack{a_1+a_2+b=k \\ a_1+a_2 \geq 1, b \geq 1}} |\nabla^{a_1} \mathbf{d}|_{L^6} |\nabla^{a_2} \mathbf{d}|_{L^6} |\nabla^{b+1} \mathbf{v}|_{L^6} |\nabla^{k+1} \mathbf{v}|_{L^2} \\
&\leq \mu_1 |\mathbf{d}|_{L^\infty}^2 |\nabla^{k+1} \mathbf{v}|_{L^2}^2 + C\mu_1 \sum_{\substack{a_1+a_2+b=k \\ a_1+a_2 \geq 1, b \geq 1}} |\nabla^{a_1+1} \mathbf{d}|_{L^2} |\nabla^{a_2+1} \mathbf{d}|_{L^2} |\nabla^{b+2} \mathbf{v}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2} \\
&\leq C\mu_1 |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2,
\end{aligned}$$

and

$$\begin{aligned}
I_2 &\leq \mu_1 |\mathbf{d}|_{L^\infty}^2 \sum_{\substack{a+b_1+b_2=k \\ b_1+b_2 \geq 1}} \left\langle |\nabla^a \partial_p \mathbf{v}_q| |\nabla^{b_1} \mathbf{d}_i| |\nabla^{b_2} \mathbf{d}_j|, \nabla^k \partial_j \mathbf{v}_i \right\rangle \\
&\leq 2\mu_1 |\mathbf{d}|_{L^\infty}^3 \sum_{\substack{a+b_1=k \\ b_1 \geq 1}} \left\langle |\nabla^a \partial_p \mathbf{v}_q| |\nabla^{b_1} \mathbf{d}|, |\nabla^{k+1} \mathbf{v}| \right\rangle \\
&\quad + \mu_1 |\mathbf{d}|_{L^\infty}^2 \sum_{\substack{a+b_1+b_2=k \\ b_1, b_2 \geq 1}} \left\langle |\nabla^a \partial_p \mathbf{v}_q| |\nabla^{b_1} \mathbf{d}| |\nabla^{b_2} \mathbf{d}|, |\nabla^{k+1} \mathbf{v}| \right\rangle \\
&\leq 2\mu_1 |\mathbf{d}|_{L^\infty}^3 \sum_{\substack{a+b_1=k \\ b_1 \geq 1}} |\nabla^{a+1} \mathbf{v}|_{L^4} |\nabla^{b_1} \mathbf{d}|_{L^4} |\nabla^{k+1} \mathbf{v}|_{L^2} \\
&\quad + \mu_1 |\mathbf{d}|_{L^\infty}^2 \sum_{\substack{a+b_1+b_2=k \\ b_1, b_2 \geq 1}} |\nabla^{a+1} \mathbf{v}|_{L^6} |\nabla^{b_1} \mathbf{d}|_{L^6} |\nabla^{b_2} \mathbf{d}|_{L^6} |\nabla^{k+1} \mathbf{v}|_{L^2} \\
&\leq C\mu_1 |\mathbf{d}|_{L^\infty}^3 |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2}^2 + C\mu_1 |\mathbf{d}|_{L^\infty}^2 |\nabla^{k+1} \mathbf{d}|_{L^2}^2 |\nabla^{k+1} \mathbf{v}|_{L^2}^2 \\
&\leq C\mu_1 |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2,
\end{aligned}$$

and

$$\begin{aligned}
I_3 &\leq \mu_1 \sum_{\substack{a_1+a_2+b=k \\ b \geq 1}} \left\langle |\nabla^{a_1} (\mathbf{d}_p \mathbf{d}_q)| |\nabla^{a_2} \partial_p \mathbf{v}_q| |\nabla^b (\mathbf{d}_i \mathbf{d}_j)|, |\nabla^k \partial_j \mathbf{v}_i| \right\rangle \\
&\leq \mu_1 \sum_{\substack{a_1+a_2+b=k \\ b \geq 1}} |\nabla^{a_1} (\mathbf{d}_p \mathbf{d}_q)|_{L^6} |\nabla^{a_2+1} \mathbf{v}|_{L^6} |\nabla^b (\mathbf{d}_i \mathbf{d}_j)|_{L^6} |\nabla^{k+1} \mathbf{v}|_{L^2} \\
&\leq C\mu_1 \sum_{a_1+b \leq k} |\nabla^{a_1} (\mathbf{d}_p \mathbf{d}_q)|_{L^6} |\nabla^b (\mathbf{d}_i \mathbf{d}_j)|_{L^6} |\nabla^{k+1} \mathbf{v}|_{L^2}^2 \\
&\leq C\mu_1 \sum_{a_1+b \leq k} |\mathbf{d}|_{L^\infty}^2 |\nabla^{a_1} \mathbf{d}|_{L^6} |\nabla^b \mathbf{d}|_{L^6} |\nabla^{k+1} \mathbf{v}|_{L^2}^2 \\
&\leq C\mu_1 |\mathbf{d}|_{L^\infty}^2 |\nabla^{k+1} \mathbf{d}|_{L^2}^2 |\nabla^{k+1} \mathbf{v}|_{L^2}^2 \\
&\leq C\mu_1 |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2.
\end{aligned}$$

So we get the estimate of the μ_1 -part

$$\left\langle \mu_1 \nabla^k \partial_j (\mathbf{d}^\top A \mathbf{d} \mathbf{d}^\top \mathbf{d}_j), \nabla^k \mathbf{v}_i \right\rangle \leq -\mu_1 |\mathbf{d}^\top \nabla^{k+1} \mathbf{v} \mathbf{d}|_{L^2}^2 + C\mu_1 |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2. \quad (2.10)$$

Second, we estimate the μ_2 -part: $\langle \mu_2 \nabla^k \partial_j [\mathbf{d}_j(\dot{\mathbf{d}}_i + B_{ki}\mathbf{d}_k)], \nabla^k \mathbf{v}_i \rangle$.

$$\begin{aligned}
& \left\langle \mu_2 \nabla^k \partial_j [\mathbf{d}_j(\dot{\mathbf{d}}_i + B_{ki}\mathbf{d}_k)], \nabla^k \mathbf{v}_i \right\rangle \\
&= -\mu_2 \left\langle \nabla^k [\mathbf{d}_j(\dot{\mathbf{d}}_i + B_{ki}\mathbf{d}_k)], \nabla^k \partial_j \mathbf{v}_i \right\rangle \\
&\leq \mu_2 |\nabla^k (\dot{\mathbf{d}}_i \mathbf{d}_j)|_{L^2} |\nabla^k \partial_j \mathbf{v}_i|_{L^2} + \mu_2 |\nabla^k (\mathbf{d}_j B_{ki} \mathbf{d}_k)|_{L^2} |\nabla^k \partial_j \mathbf{v}_i|_{L^2} \\
&\leq C\mu_2 (|\nabla^k \dot{\mathbf{d}}|_{L^2} |\mathbf{d}|_{L^\infty} + |\nabla^k \mathbf{d}|_{L^2} |\dot{\mathbf{d}}|_{L^\infty}) |\nabla^{k+1} \mathbf{v}|_{L^2} \\
&\quad + C\mu_2 (|\mathbf{d}|_{L^\infty} |\nabla^k (B_{ki} \mathbf{d}_k)|_{L^2} + |\nabla^k \mathbf{d}_j|_{L^2} |B_{ki} \mathbf{d}_k|_{L^\infty}) |\nabla^k \partial_j \mathbf{v}_i|_{L^2} \\
&\leq C\mu_2 |\nabla^k \dot{\mathbf{d}}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2} + C\mu_2 |\nabla^{k+1} \mathbf{d}|_{L^2} |\mathbf{d}|_{L^\infty} |\nabla \mathbf{v}|_{L^\infty} |\nabla^{k+1} \mathbf{v}|_{L^2} \\
&\quad + C\mu_2 |\nabla^{k+1} \mathbf{d}|_{L^2} (|\nabla^{k+1} \mathbf{v}|_{L^2} |\mathbf{d}|_{L^\infty} + |\nabla^k \mathbf{d}|_{L^2} |\nabla \mathbf{v}|_{L^\infty}) |\nabla^{k+1} \mathbf{v}|_{L^2} \\
&\leq C\mu_2 |\nabla^k \dot{\mathbf{d}}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2} + C\mu_2 |\nabla^{k+1} \mathbf{d}|_{L^2}^2 |\nabla^{k+1} \mathbf{v}|_{L^2}^2 \\
&\leq C\mu_2 |\dot{\mathbf{d}}|_{H^s} |\nabla \mathbf{d}|_{H^s} |\nabla \mathbf{v}|_{H^s} + C\mu_2 |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2. \tag{2.11}
\end{aligned}$$

Third, by similar arguments on estimating the μ_2 -parts, we get the estimate of the μ_3 -part:

$$\langle \mu_3 \nabla^k \partial_j [\mathbf{d}_i(\dot{\mathbf{d}}_j + B_{kj}\mathbf{d}_k)], \nabla^k \mathbf{v}_i \rangle \leq C\mu_3 |\dot{\mathbf{d}}|_{H^s} |\nabla \mathbf{d}|_{H^s} |\nabla \mathbf{v}|_{H^s} + C\mu_3 |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2. \tag{2.12}$$

Fourth, we estimate the μ_5 -part: $\langle \mu_5 \nabla^k \partial_j (\mathbf{d}_j \mathbf{d}_p A_{pi}), \nabla^k \mathbf{v}_i \rangle$.

$$\begin{aligned}
& \left\langle \mu_5 \nabla^k \partial_j (\mathbf{d}_j \mathbf{d}_p A_{pi}), \nabla^k \mathbf{v}_i \right\rangle \\
&= -\mu_5 \left\langle \nabla^k (\mathbf{d}_j \mathbf{d}_p A_{pi}), \nabla^k \partial_j \mathbf{v}_i \right\rangle \\
&\leq C\mu_5 (|\mathbf{d}_j \mathbf{d}_p|_{L^\infty} |\nabla^k A_{pi}|_{L^2} + |A_{pi}|_{L^\infty} |\nabla^k (\mathbf{d}_j \mathbf{d}_p)|_{L^2}) |\nabla^k \partial_j \mathbf{v}_i|_{L^2} \\
&\leq C\mu_5 |\mathbf{d}|_{L^\infty}^2 |\nabla^{k+1} \mathbf{v}|_{L^2}^2 + C\mu_5 |\mathbf{d}|_{L^\infty} |\nabla^k \mathbf{d}|_{L^2} |\nabla \mathbf{v}|_{L^\infty} |\nabla^{k+1} \mathbf{v}|_{L^2} \\
&\leq C\mu_5 |\nabla^{k+1} \mathbf{d}|_{L^2}^2 |\nabla^{k+1} \mathbf{v}|_{L^2}^2 \\
&\leq C\mu_5 |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2. \tag{2.13}
\end{aligned}$$

Finally, the same arguments on the estimate of the μ_5 -part implies the bound of the μ_6 -part:

$$\langle \mu_6 \nabla^k \partial_j (\mathbf{d}_i \mathbf{d}_p A_{pj}), \nabla^k \mathbf{v}_i \rangle \leq C\mu_6 |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2. \tag{2.14}$$

Substituting the inequalities (2.8)-(2.14) into (2.7), one has

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} |\nabla^k \mathbf{v}|_{L^2}^2 + \frac{1}{2} \mu_4 |\nabla^{k+1} \mathbf{v}|_{L^2}^2 + \mu_1 |\mathbf{d}^\top (\nabla^{k+1} \mathbf{v}) \mathbf{d}|_{L^2}^2 \\
&\leq C(1 + \mu_2 + \mu_3) \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) |\nabla \mathbf{v}|_{H^s} \\
&\quad + C(\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6) |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2. \tag{2.15}
\end{aligned}$$

Taking ∇^k in the third equation of (1.1), multiplying by $\nabla^k \dot{\mathbf{d}}$, integrating by parts, one easily obtains

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \left(|\nabla^k \dot{\mathbf{d}}|_{L^2}^2 + |\nabla^{k+1} \mathbf{d}|_{L^2}^2 \right) - \lambda_1 |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 \\
&= - \sum_{\substack{a+b=k \\ a \geq 1}} \left\langle \nabla^a \mathbf{v} \nabla^{b+1} \dot{\mathbf{d}}, \nabla^k \dot{\mathbf{d}} \right\rangle - \sum_{\substack{a+b=k+1 \\ a \geq 1}} \left\langle \nabla^a \mathbf{v} \nabla^{b+1} \mathbf{d}, \nabla^{k+1} \mathbf{d} \right\rangle \\
&\quad - \sum_{a+b+c=k} \left\langle \nabla^a \dot{\mathbf{d}} \nabla^b \dot{\mathbf{d}} \nabla^c \mathbf{d}, \nabla^k \dot{\mathbf{d}} \right\rangle + \sum_{a+b+c=k} \left\langle \nabla^{a+1} \mathbf{d} \nabla^{b+1} \mathbf{d} \nabla^c \mathbf{d}, \nabla^k \dot{\mathbf{d}} \right\rangle \\
&\quad + \lambda_1 \sum_{a+b=k} \left\langle \nabla^a (\mathbf{d}^\top B \mathbf{d}) \nabla^b, \nabla^k \dot{\mathbf{d}} \right\rangle - \lambda_2 \sum_{a+b=k} \left\langle \nabla^a (\mathbf{d}^\top A \mathbf{d}) \nabla^b \mathbf{d}, \nabla^k \dot{\mathbf{d}} \right\rangle \tag{2.16}
\end{aligned}$$

$$\begin{aligned}
& -\lambda_1 \sum_{a+b=k} \left\langle \nabla^a B \nabla^b \mathbf{d}, \nabla^k \dot{\mathbf{d}} \right\rangle + \lambda_2 \sum_{a+b=k} \left\langle \nabla^a A \nabla^b \mathbf{d}, \nabla^k \dot{\mathbf{d}} \right\rangle \\
& \equiv I_1 + I_2 + I_3 + I_4 + I_5 + I_6 + I_7 + I_8.
\end{aligned}$$

By using Hölder inequality, Sobolev embedding, the facts $\operatorname{div} \mathbf{v} = 0$ and $|\mathbf{d}| = 1$, one can directly calculate the following estimates:

$$\begin{aligned}
I_1 &= - \sum_{\substack{a+b=k \\ a \geq 1}} \left\langle \nabla^a \mathbf{v} \nabla^{b+1} \dot{\mathbf{d}}, \nabla^k \dot{\mathbf{d}} \right\rangle \\
&\leq |\nabla \mathbf{v}|_{L^\infty} |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 + \sum_{\substack{a+b=k \\ a \geq 2}} |\nabla^a \mathbf{v}|_{L^4} |\nabla^{b+1} \dot{\mathbf{d}}|_{L^4} |\nabla^k \dot{\mathbf{d}}|_{L^2} \\
&\leq C |\nabla^3 \mathbf{v}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 + C |\nabla^{k+1} \mathbf{v}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 \\
&\leq C |\nabla \mathbf{v}|_{H^s} |\dot{\mathbf{d}}|_{H^s}^2,
\end{aligned} \tag{2.17}$$

and

$$\begin{aligned}
I_2 &= - \sum_{\substack{a+b=k+1 \\ a \geq 1}} \left\langle \nabla^a \mathbf{v} \nabla^{b+1} \mathbf{d}, \nabla^{k+1} \mathbf{d} \right\rangle \\
&\leq |\nabla \mathbf{v}|_{L^\infty} |\nabla^{k+1} \mathbf{d}|_{L^2}^2 + |\nabla \mathbf{d}|_{L^\infty} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2} + \sum_{\substack{a+b=k+1 \\ a \geq 2, b \geq 2}} |\nabla^a \mathbf{v}|_{L^4} |\nabla^{b+1} \mathbf{d}|_{L^4} |\nabla^{k+1} \mathbf{d}|_{L^2} \\
&\leq C |\nabla^3 \mathbf{v}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2}^2 + C |\nabla^3 \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2} + C |\nabla^{k+1} \mathbf{v}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2}^2 \\
&\leq C |\nabla \mathbf{v}|_{H^s} |\nabla \mathbf{d}|_{H^s}^2,
\end{aligned} \tag{2.18}$$

and

$$\begin{aligned}
I_3 &= - \sum_{a+b+c=k} \left\langle \nabla^a \dot{\mathbf{d}} \nabla^b \dot{\mathbf{d}} \nabla^c \mathbf{d}, \nabla^k \dot{\mathbf{d}} \right\rangle \\
&\leq \sum_{a+b=k} |\mathbf{d}|_{L^\infty} |\nabla^a \dot{\mathbf{d}} \nabla^b \dot{\mathbf{d}}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} - \sum_{\substack{a+b+c=k \\ c \geq 1}} \left\langle \nabla^a \dot{\mathbf{d}} \nabla^b \dot{\mathbf{d}} \nabla^c \mathbf{d}, \nabla^k \dot{\mathbf{d}} \right\rangle \\
&\leq C |\nabla^2 \mathbf{d}|_{L^2} |\dot{\mathbf{d}}|_{L^\infty} |\nabla^k \dot{\mathbf{d}}|_{L^2} + C \sum_{\substack{a+b=k \\ a, b \geq 1}} |\nabla^2 \mathbf{d}|_{L^2} |\nabla^a \dot{\mathbf{d}}|_{L^4} |\nabla^b \dot{\mathbf{d}}|_{L^4} |\nabla^k \dot{\mathbf{d}}|_{L^2} \\
&\quad + |\dot{\mathbf{d}}|_{L^\infty}^2 |\nabla^k \mathbf{d}|_{L^2} |\nabla \dot{\mathbf{d}}|_{L^2} + \sum_{\substack{a+b+c=k \\ 1 \leq c \leq k-1}} |\nabla^c \mathbf{d}|_{L^\infty} |\nabla^a \dot{\mathbf{d}}|_{L^4} |\nabla^b \dot{\mathbf{d}}|_{L^4} |\nabla^k \dot{\mathbf{d}}|_{L^2} \\
&\leq C |\nabla^2 \mathbf{d}|_{L^2} |\nabla^2 \dot{\mathbf{d}}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 + C |\nabla^2 \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2}^3 \\
&\quad + C |\nabla^2 \dot{\mathbf{d}}|_{L^2}^2 |\nabla^k \dot{\mathbf{d}}|_{L^2} |\nabla^k \mathbf{d}|_{L^2} + C |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2}^3 \\
&\leq C |\dot{\mathbf{d}}|_{H^s}^3 |\nabla \mathbf{d}|_{H^s},
\end{aligned} \tag{2.19}$$

and

$$\begin{aligned}
I_4 &= \sum_{a+b+c=k} \left\langle \nabla^{a+1} \mathbf{d} \nabla^{b+1} \mathbf{d} \nabla^c \mathbf{d}, \nabla^k \dot{\mathbf{d}} \right\rangle \\
&\leq \sum_{a+b=k} |\mathbf{d}|_{L^\infty} |\nabla^{a+1} \mathbf{d} \nabla^{b+1} \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} + \sum_{\substack{a+b+c=k \\ c \geq 1}} \left\langle \nabla^{a+1} \mathbf{d} \nabla^{b+1} \mathbf{d} \nabla^c \mathbf{d}, \nabla^k \dot{\mathbf{d}} \right\rangle \\
&\leq C |\nabla^2 \mathbf{d}|_{L^2} |\nabla \mathbf{d}|_{L^\infty} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} + C \sum_{\substack{a+b=k \\ a, b \geq 1}} |\nabla^2 \mathbf{d}|_{L^2} |\nabla^{a+1} \mathbf{d}|_{L^4} |\nabla^{b+1} \mathbf{d}|_{L^4} |\nabla^k \dot{\mathbf{d}}|_{L^2}
\end{aligned}$$

$$\begin{aligned}
& + |\nabla \mathbf{d}|_{L^\infty}^2 |\nabla^k \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} + \sum_{\substack{a+b+c=k \\ 1 \leq c \leq k-1}} |\nabla^c \mathbf{d}|_{L^\infty} |\nabla^{a+1} \mathbf{d}|_{L^4} |\nabla^{b+1} \mathbf{d}|_{L^4} |\nabla^k \dot{\mathbf{d}}|_{L^2} \\
& \leq C |\nabla^2 \mathbf{d}|_{L^2} |\nabla^3 \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} + C |\nabla^2 \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2}^2 |\nabla^k \dot{\mathbf{d}}|_{L^2} \\
& \quad + C |\nabla^3 \mathbf{d}|_{L^2}^2 |\nabla^k \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} + C |\nabla^{k+1} \mathbf{d}|_{L^2}^3 |\nabla^k \dot{\mathbf{d}}|_{L^2} \\
& \leq C |\nabla \mathbf{d}|_{H^s}^3 |\dot{\mathbf{d}}|_{H^s},
\end{aligned} \tag{2.20}$$

and

$$\begin{aligned}
I_5 &= \lambda_1 \sum_{a+b=k} \left\langle \nabla^a (\mathbf{d}^\top B \mathbf{d}) \nabla^b \mathbf{d}, \nabla^k \dot{\mathbf{d}} \right\rangle \\
&\leq |\lambda_1| \sum_{\substack{a+b=k \\ b \leq k-1}} |\nabla^b \mathbf{d}|_{L^\infty} |\nabla^a (\mathbf{d}^\top B \mathbf{d})|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} + |\lambda_1| |\mathbf{d}^\top B \mathbf{d}|_{L^\infty} |\nabla^k \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} \\
&\leq C |\lambda_1| |\nabla^{k+1} \mathbf{d}|_{L^2} \sum_{1 \leq a \leq k} |\nabla^a (\mathbf{d}^\top B \mathbf{d})|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} + C |\lambda_1| |\mathbf{d}|_{L^\infty}^2 |\nabla \mathbf{v}|_{L^\infty} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} \\
&\leq C |\lambda_1| \sum_{1 \leq a \leq k} (|\mathbf{d}_i \mathbf{d}_j|_{L^\infty} |\nabla^a B_{ij}|_{L^2} + |B_{ij}|_{L^\infty} |\nabla^a (\mathbf{d}_i \mathbf{d}_j)|_{L^2}) |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} \\
&\quad + C |\lambda_1| |\mathbf{d}|_{L^\infty}^2 |\nabla \mathbf{v}|_{L^\infty} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2} \\
&\leq C |\lambda_1| (|\mathbf{d}|_{L^\infty}^2 |\nabla^{k+1} \mathbf{v}|_{L^2} + |\nabla \mathbf{v}|_{L^\infty} |\mathbf{d}|_{L^\infty} |\nabla^{k+1} \mathbf{d}|_{L^2}) |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2} \\
&\quad + C |\lambda_1| |\mathbf{d}|_{L^\infty}^2 |\nabla \mathbf{v}|_{L^\infty} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2} \\
&\leq C |\lambda_1| |\nabla^{k+1} \mathbf{d}|_{L^2}^2 |\nabla^{k+1} \mathbf{v}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} \\
&\leq C |\lambda_1| |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s} |\dot{\mathbf{d}}|_{H^s},
\end{aligned} \tag{2.21}$$

and by similar arguments on estimating the term I_5

$$I_6 \leq C |\lambda_2| |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s} |\dot{\mathbf{d}}|_{H^s}, \tag{2.22}$$

and

$$\begin{aligned}
I_7 &= -\lambda_1 \sum_{a+b=k} \left\langle \nabla^a B \nabla^b \mathbf{d}, \nabla^k \dot{\mathbf{d}} \right\rangle \\
&\leq |\lambda_1| \sum_{\substack{a+b=k \\ 0 \leq b \leq k-1}} |\nabla^b \mathbf{d}|_{L^\infty} |\nabla^a B|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} + |\lambda_1| |B|_{L^\infty} |\nabla^k \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} \\
&\leq C |\lambda_1| \sum_{1 \leq a \leq k} |\nabla^{a+1} \mathbf{v}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} + C |\lambda_1| |\nabla \mathbf{v}|_{L^\infty} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} \\
&\leq C |\lambda_1| |\nabla^{k+1} \mathbf{v}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} \\
&\leq C |\lambda_1| |\nabla \mathbf{v}|_{H^s} |\nabla \mathbf{d}|_{H^s} |\dot{\mathbf{d}}|_{H^s},
\end{aligned} \tag{2.23}$$

and by similarly estimating on the term I_7

$$I_8 \leq C |\lambda_2| |\nabla \mathbf{v}|_{H^s} |\nabla \mathbf{d}|_{H^s} |\dot{\mathbf{d}}|_{H^s}. \tag{2.24}$$

Combining the inequalities (2.17), (2.18), (2.19), (2.20), (2.21), (2.22), (2.23) and (2.24), it can be derived from (2.16) that the following estimate is valid:

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \left(|\nabla^k \dot{\mathbf{d}}|_{L^2}^2 + |\nabla^{k+1} \mathbf{d}|_{L^2}^2 \right) - \lambda_1 |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 \\
& \leq C (|\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2) |\dot{\mathbf{d}}|_{H^s} |\nabla \mathbf{d}|_{H^s} + C (|\lambda_1| + |\lambda_2|) |\nabla \mathbf{d}|_{H^s}^2 |\dot{\mathbf{d}}|_{H^s} |\nabla \mathbf{v}|_{H^s} \\
& \quad + C (|\lambda_1| + |\lambda_2|) |\nabla \mathbf{d}|_{H^s} |\dot{\mathbf{d}}|_{H^s} |\nabla \mathbf{v}|_{H^s}.
\end{aligned} \tag{2.25}$$

Then the inequalities (2.15) and (2.25) imply that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(|\nabla^k \mathbf{v}|_{L^2}^2 + |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 + |\nabla^{k+1} \mathbf{d}|_{L^2}^2 \right) + \frac{1}{2} \mu_4 |\nabla^{k+1} \mathbf{v}|_{L^2}^2 - \lambda_1 |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 + \mu_1 |\mathbf{d}^\top (\nabla^{k+1} \mathbf{v}) \mathbf{d}|_{L^2}^2 \\ & \leq C(1 + |\lambda_1| + |\lambda_2| + \mu_2 + \mu_3)(|\mathbf{v}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2) |\nabla \mathbf{v}|_{H^s} + C \left(|\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right)^2 \\ & \quad + C(\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6) |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2 + C(|\lambda_1| + |\lambda_2|) |\nabla \mathbf{d}|_{H^s}^2 |\dot{\mathbf{d}}|_{H^s} |\nabla \mathbf{v}|_{H^s}. \end{aligned} \quad (2.26)$$

for all $1 \leq k \leq s$, which combining the inequality (2.6) and Schwartz inequality gives us

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) + \frac{1}{2} \mu_4 |\nabla \mathbf{v}|_{H^s}^2 - \lambda_1 |\dot{\mathbf{d}}|_{H^s}^2 + \mu_1 \sum_{k=0}^s |\mathbf{d}^\top (\nabla^{k+1} \mathbf{v}) \mathbf{d}|_{L^2}^2 \\ & \leq C(1 + |\lambda_1| + |\lambda_2| + \mu_2 + \mu_3) \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) |\nabla \mathbf{v}|_{H^s} \\ & \leq C(1 + |\lambda_1| + |\lambda_2|) \left(|\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right)^2 + C(\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6) |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2, \end{aligned}$$

and then the proof of Lemma 2.1 is completed. \square

3. THE APPROXIMATED SYSTEMS OF ERICKSEN-LESLIE'S PARABOLIC-HYPERBOLIC MODEL

In order to obtain the local existence of Ericksen-Leslie's parabolic-hyperbolic liquid crystal model (1.1), the approximated system of (1.1) is constructed as follows:

$$\begin{cases} \partial_t \mathbf{v}^\epsilon + \mathcal{P} J_\epsilon [J_\epsilon \mathbf{v}^\epsilon \cdot \nabla J_\epsilon \mathbf{v}^\epsilon] = \frac{1}{2} \mu_4 J_\epsilon (\Delta J_\epsilon \mathbf{v}^\epsilon) - \mathcal{P} J_\epsilon \operatorname{div}(J_\epsilon \nabla J_\epsilon \mathbf{d}^\epsilon \odot J_\epsilon \nabla J_\epsilon \mathbf{d}^\epsilon) \\ \quad + \mathcal{P} J_\epsilon \mathcal{L}(J_\epsilon \mathbf{d}^\epsilon, J_\epsilon \mathbf{v}^\epsilon), \\ \operatorname{div} \mathbf{v}^\epsilon = 0, \\ \partial_t \dot{\mathbf{d}}^\epsilon + J_\epsilon [\mathbf{v}^\epsilon \cdot \nabla J_\epsilon \dot{\mathbf{d}}^\epsilon] = J_\epsilon \Delta J_\epsilon \mathbf{d}^\epsilon + \gamma_\epsilon \mathbf{d}^\epsilon + \lambda_1 (J_\epsilon \dot{\mathbf{d}}^\epsilon - J_\epsilon B_\epsilon J_\epsilon \mathbf{d}^\epsilon) + \lambda_1 J_\epsilon A_\epsilon J_\epsilon \mathbf{d}^\epsilon, \\ \dot{\mathbf{d}}^\epsilon = \partial_t \mathbf{d}^\epsilon + J_\epsilon [\mathbf{v}^\epsilon \cdot \nabla J_\epsilon \mathbf{d}^\epsilon], \end{cases} \quad (3.1)$$

where J_ϵ is mollifier operators, \mathcal{P} is the Leray projection, $B_\epsilon = \frac{1}{2} (\nabla J_\epsilon \mathbf{v}^\epsilon - \nabla J_\epsilon \mathbf{v}^{\epsilon \top})$, $A_\epsilon = \frac{1}{2} (\nabla J_\epsilon \mathbf{v}^\epsilon + \nabla J_\epsilon \mathbf{v}^{\epsilon \top})$, and

$$\begin{aligned} \mathcal{L}(J_\epsilon \mathbf{d}^\epsilon, J_\epsilon \mathbf{v}^\epsilon) = & \partial_j \left[\mu_1 J_\epsilon \mathbf{d}_k^\epsilon J_\epsilon \mathbf{d}_p^\epsilon J_\epsilon (A_{kp})_\epsilon J_\epsilon \mathbf{d}_i^\epsilon J_\epsilon \mathbf{d}_j^\epsilon + \mu_2 J_\epsilon \mathbf{d}_j^\epsilon (J_\epsilon \dot{\mathbf{d}}_i^\epsilon + J_\epsilon (B_{ki})_\epsilon J_\epsilon \mathbf{d}_k^\epsilon) \right. \\ & \left. + \mu_3 J_\epsilon \mathbf{d}_i^\epsilon (J_\epsilon \dot{\mathbf{d}}_j^\epsilon + J_\epsilon (B_{kj})_\epsilon J_\epsilon \mathbf{d}_k^\epsilon) + \mu_5 J_\epsilon \mathbf{d}_j^\epsilon J_\epsilon \mathbf{d}_k^\epsilon J_\epsilon (A_{ki})_\epsilon + \mu_6 J_\epsilon \mathbf{d}_i^\epsilon J_\epsilon \mathbf{d}_k^\epsilon J_\epsilon (A_{kj})_\epsilon \right] \end{aligned}$$

and

$$\gamma_\epsilon = -|J_\epsilon \dot{\mathbf{d}}^\epsilon|^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|^2 + \lambda_1 (J_\epsilon \mathbf{d}^\epsilon)^\top J_\epsilon B_\epsilon J_\epsilon \mathbf{d}^\epsilon - \lambda_2 (J_\epsilon \mathbf{d}^\epsilon)^\top J_\epsilon A_\epsilon \mathbf{d}^\epsilon.$$

Furthermore, the approximated system (3.1) is imposed on the same initial data (1.4) with the system (1.1).

One can obviously rewrite the approximated system (3.1) as the form:

$$\begin{cases} \frac{d}{dt} \mathbf{v}^\epsilon = F_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon), \\ \frac{d}{dt} \dot{\mathbf{d}}^\epsilon = G_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon), \\ \frac{d}{dt} \mathbf{d}^\epsilon = H_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon), \end{cases} \quad (3.2)$$

where

$$\begin{cases} F_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon) = \frac{1}{2} \mu_4 J_\epsilon^2 \Delta \mathbf{v}^\epsilon - \mathcal{P} J_\epsilon [J_\epsilon \mathbf{v}^\epsilon \cdot \nabla J_\epsilon \mathbf{v}^\epsilon] \\ \quad - \mathcal{P} J_\epsilon \operatorname{div}(J_\epsilon \nabla J_\epsilon \mathbf{d}^\epsilon \odot J_\epsilon \nabla J_\epsilon \mathbf{d}^\epsilon) + \mathcal{P} J_\epsilon \mathcal{L}(J_\epsilon \mathbf{d}^\epsilon, J_\epsilon \mathbf{v}^\epsilon), \\ G_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon) = J_\epsilon^2 \Delta \mathbf{d}^\epsilon - J_\epsilon [\mathbf{v}^\epsilon \cdot \nabla J_\epsilon \dot{\mathbf{d}}^\epsilon] + \gamma_\epsilon \mathbf{d}^\epsilon + \lambda_1 (J_\epsilon \dot{\mathbf{d}}^\epsilon - J_\epsilon B_\epsilon J_\epsilon \mathbf{d}^\epsilon) + \lambda_1 J_\epsilon A_\epsilon J_\epsilon \mathbf{d}^\epsilon, \\ H_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon) = \dot{\mathbf{d}}^\epsilon - J_\epsilon [\mathbf{v}^\epsilon \cdot \nabla J_\epsilon \mathbf{d}^\epsilon]. \end{cases}$$

Denoting by $\mathcal{F}_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon) = (F_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon), G_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon), H_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon))^\top$, one can represent the system (3.2) as a simpler form

$$\begin{cases} \frac{d}{dt} \begin{pmatrix} \mathbf{v}^\epsilon \\ \dot{\mathbf{d}}^\epsilon \\ \mathbf{d}^\epsilon \end{pmatrix} = \mathcal{F}_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon), \\ \left. \begin{pmatrix} \mathbf{v}^\epsilon \\ \dot{\mathbf{d}}^\epsilon \\ \mathbf{d}^\epsilon \end{pmatrix} \right|_{t=0} = \begin{pmatrix} \mathbf{v}^{in} \\ \dot{\mathbf{d}}^{in} \\ \mathbf{d}^{in} \end{pmatrix} \end{cases} \quad (3.3)$$

Our goal is to show the global existence of the strong classical solution to the approximated equations (3.1) by the basic ODE theory. After doing that, we can construct some useful results of compactness, so that we could imply the local existence of the strong solution to the system (1.1) by passing to the limit from the approximated system (3.1). In this section, we first prove the local existence of the classical solutions to the approximated equations (3.1). Then the constructed local solution can be extended to the global time interval $[0, \infty)$.

3.1. The local existence of the approximated system (3.1). The following basic properties will be used to estimate the Lipschitz continuity of the functions $\mathcal{F}_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon)$ in some proper spaces.

Lemma 3.1. (1) *The following calculus inequality*

$$|\mathbf{u}\mathbf{v}|_{H^s} \leq |\mathbf{u}|_{L^\infty} |\mathbf{v}|_{H^s} + |\mathbf{v}|_{L^\infty} |\mathbf{u}|_{H^s}$$

holds for all $\mathbf{u}, \mathbf{v} \in H^s \cap L^\infty$.

(2) For any $\mathbf{v} \in H^s$, as $\epsilon \rightarrow 0$, the convergence $|J_\epsilon \mathbf{v} - \mathbf{v}|_{H^s} \rightarrow 0$ is valid, and the inequality $|J_\epsilon \mathbf{v} - \mathbf{v}|_{H^{s-1}} \leq C\epsilon |\mathbf{v}|_{H^s}$ holds for some positive constant C .

(3) For any $\mathbf{v} \in H^s$ and $k \in \mathbb{Z}^+ \cup \{0\}$, the inequalities

$$|J_\epsilon \mathbf{v}|_{H^{s+k}} \leq \frac{C(s, k)}{\epsilon^k} |\mathbf{v}|_{H^s}$$

and

$$|J_\epsilon \nabla^k \mathbf{v}|_{L^\infty} \leq \frac{C(k)}{\epsilon^{\frac{n}{2}+k}} |\mathbf{v}|_{L^2}$$

are valid.

Proof. These properties are fundamental facts and we omit the details of the proof. The details can be seen in [23] for instance. \square

Proposition 3.1. Let $s > \frac{n}{2} + 2(n = 2, 3)$. For given initial data $\mathbf{v}^{in}, \dot{\mathbf{d}}^{in}, \mathbf{d}^{in} \in H^s$ there exists a unique solution $(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon)^\top \in C([0, T_\epsilon); H^s \times H^s \times H^s)$ to the equations (3.1) for some $T_\epsilon > 0$.

Proof. Our goal is going to verify the Lipschitz continuity of the functions $\mathcal{F}_\epsilon(\cdot, \cdot, \cdot)$ in the product space $H^s \times H^s \times H^s$. By the way, we define the norms on a space $X \times X \times X$ as

$$\left| \begin{pmatrix} a \\ b \\ c \end{pmatrix} \right|_{X \times X \times X} \equiv |a|_X + |b|_X + |c|_X.$$

It is calculated by Sobolev embedding and Lemma 3.1 that for any $\mathbf{v}^1, \mathbf{v}^2, \dot{\mathbf{d}}^1, \dot{\mathbf{d}}^2, \mathbf{d}^1, \mathbf{d}^2 \in H^s$

$$\begin{aligned} & |J_\epsilon \operatorname{div}(J_\epsilon \nabla J_\epsilon \mathbf{d}^1 \odot J_\epsilon \nabla J_\epsilon \mathbf{d}^1) - J_\epsilon \operatorname{div}(J_\epsilon \nabla J_\epsilon \mathbf{d}^2 \odot J_\epsilon \nabla J_\epsilon \mathbf{d}^2)|_{H^s} \\ & \leq |\Delta J_\epsilon^2(\mathbf{d}^1 - \mathbf{d}^2) \odot J_\epsilon \nabla J_\epsilon \mathbf{d}^1|_{H^s} + |J_\epsilon \nabla J_\epsilon(\mathbf{d}^1 - \mathbf{d}^2) \odot \Delta J_\epsilon^2 \mathbf{d}^1|_{H^s} \\ & \quad + |\Delta J_\epsilon^2 \mathbf{d}^2 \odot J_\epsilon \nabla J_\epsilon(\mathbf{d}^1 - \mathbf{d}^2)|_{H^s} + |J_\epsilon \nabla J_\epsilon \mathbf{d}^2 \odot \Delta J_\epsilon^2(\mathbf{d}^1 - \mathbf{d}^2)|_{H^s} \\ & \lesssim |\nabla J_\epsilon \mathbf{d}^1|_{L^\infty} |J_\epsilon \Delta(\mathbf{d}^1 - \mathbf{d}^2)|_{H^s} + |\Delta J_\epsilon(\mathbf{d}^1 - \mathbf{d}^2)|_{L^\infty} |\nabla J_\epsilon \mathbf{d}^1|_{H^s} \end{aligned}$$

$$\begin{aligned}
& + |\Delta J_\epsilon \mathbf{d}^1|_{L^\infty} |\nabla J_\epsilon(\mathbf{d}^1 - \mathbf{d}^2)|_{H^s} + |\nabla J_\epsilon(\mathbf{d}^1 - \mathbf{d}^2)|_{L^\infty} |\Delta J_\epsilon \mathbf{d}^1|_{H^s} \\
& + |\Delta J_\epsilon \mathbf{d}^2|_{L^\infty} |\nabla J_\epsilon(\mathbf{d}^1 - \mathbf{d}^2)|_{H^s} + |\nabla J_\epsilon(\mathbf{d}^1 - \mathbf{d}^2)|_{L^\infty} |\Delta J_\epsilon \mathbf{d}^2|_{H^s} \\
& + |\nabla J_\epsilon \mathbf{d}^2|_{L^\infty} |J_\epsilon \Delta(\mathbf{d}^1 - \mathbf{d}^2)|_{H^s} + |\Delta J_\epsilon(\mathbf{d}^1 - \mathbf{d}^2)|_{L^\infty} |\nabla J_\epsilon \mathbf{d}^2|_{H^s} \\
& \lesssim \frac{1}{\epsilon^{\frac{n}{2}}} |\nabla \mathbf{d}^1|_{L^2} \cdot \frac{1}{\epsilon^2} |\mathbf{d}^1 - \mathbf{d}^2|_{H^s} + \frac{1}{\epsilon^s} |\nabla \mathbf{d}^1|_{L^2} \cdot \frac{1}{\epsilon^{\frac{n}{2}+2}} |\mathbf{d}^1 - \mathbf{d}^2|_{L^2} \\
& \quad + \frac{1}{\epsilon^{\frac{n}{2}+1}} |\nabla \mathbf{d}^1|_{L^2} \cdot \frac{1}{\epsilon} |\mathbf{d}^1 - \mathbf{d}^2|_{H^s} + \frac{1}{\epsilon^{s+1}} |\nabla \mathbf{d}^1|_{L^2} \cdot \frac{1}{\epsilon^{\frac{n}{2}+1}} |\mathbf{d}^1 - \mathbf{d}^2|_{L^2} \\
& \quad + \frac{1}{\epsilon^{\frac{n}{2}+2}} |\nabla \mathbf{d}^2|_{L^2} |\mathbf{d}^1 - \mathbf{d}^2|_{H^s} + \frac{1}{\epsilon^{\frac{n}{2}+s+2}} |\nabla \mathbf{d}^2|_{L^2} |\mathbf{d}^1 - \mathbf{d}^2|_{L^2} \\
& \leq \frac{C}{\epsilon^{\frac{n}{2}+s+2}} (|\nabla \mathbf{d}^1|_{L^2} + |\nabla \mathbf{d}^2|_{L^2}) |\mathbf{d}^1 - \mathbf{d}^2|_{H^s},
\end{aligned}$$

and

$$\begin{aligned}
& |\mu_1 J_\epsilon \partial_j [J_\epsilon \mathbf{d}_k^1 J_\epsilon \mathbf{d}_p^1 J_\epsilon (A_{kp}^1)_\epsilon J_\epsilon \mathbf{d}_i^1 J_\epsilon \mathbf{d}_j^1] - \mu_1 J_\epsilon \partial_j [J_\epsilon \mathbf{d}_k^2 J_\epsilon \mathbf{d}_p^2 J_\epsilon (A_{kp}^2)_\epsilon J_\epsilon \mathbf{d}_i^2 J_\epsilon \mathbf{d}_j^2]|_{H^s} \\
& \leq \mu_1 |J_\epsilon \partial_j [J_\epsilon (\mathbf{d}_k^1 - \mathbf{d}_k^2) J_\epsilon \mathbf{d}_p^1 J_\epsilon (A_{kp}^1)_\epsilon J_\epsilon \mathbf{d}_i^1 J_\epsilon \mathbf{d}_j^1]|_{H^s} \\
& \quad + \mu_1 |J_\epsilon \partial_j [J_\epsilon \mathbf{d}_k^1 J_\epsilon (\mathbf{d}_p^1 - \mathbf{d}_p^2) J_\epsilon (A_{kp}^1)_\epsilon J_\epsilon \mathbf{d}_i^1 J_\epsilon \mathbf{d}_j^1]|_{H^s} \\
& \quad + \mu_1 |J_\epsilon \partial_j [J_\epsilon \mathbf{d}_k^1 J_\epsilon \mathbf{d}_p^1 J_\epsilon (A_{kp}^1 - A_{kp}^2)_\epsilon J_\epsilon \mathbf{d}_i^1 J_\epsilon \mathbf{d}_j^1]|_{H^s} \\
& \quad + \mu_1 |J_\epsilon \partial_j [J_\epsilon \mathbf{d}_k^1 J_\epsilon \mathbf{d}_p^1 J_\epsilon (A_{kp}^1)_\epsilon J_\epsilon (\mathbf{d}_i^1 - \mathbf{d}_i^2) J_\epsilon \mathbf{d}_j^1]|_{H^s} \\
& \quad + \mu_1 |J_\epsilon \partial_j [J_\epsilon \mathbf{d}_k^1 J_\epsilon \mathbf{d}_p^1 J_\epsilon (A_{kp}^1)_\epsilon J_\epsilon \mathbf{d}_i^1 J_\epsilon (\mathbf{d}_j^1 - \mathbf{d}_j^2)]|_{H^s} \\
& \equiv I_1 + I_2 + I_3 + I_4 + I_5,
\end{aligned}$$

where

$$\begin{aligned}
I_1 & \leq \frac{C(s)\mu_1}{\epsilon} |J_\epsilon(\mathbf{d}_k^1 - \mathbf{d}_k^2) J_\epsilon \mathbf{d}_p^1 J_\epsilon (A_{kp}^1)_\epsilon J_\epsilon \mathbf{d}_i^1 J_\epsilon \mathbf{d}_j^1|_{H^s} \\
& \leq \frac{C(s)\mu_1}{\epsilon} |J_\epsilon(\mathbf{d}_k^1 - \mathbf{d}_k^2)|_{H^s} |J_\epsilon \mathbf{d}_p^1 J_\epsilon (A_{kp}^1)_\epsilon J_\epsilon \mathbf{d}_i^1 J_\epsilon \mathbf{d}_j^1|_{L^\infty} \\
& \quad + \frac{C(s)\mu_1}{\epsilon} |J_\epsilon(\mathbf{d}_k^1 - \mathbf{d}_k^2)|_{L^\infty} |J_\epsilon \mathbf{d}_p^1 J_\epsilon (A_{kp}^1)_\epsilon J_\epsilon \mathbf{d}_i^1 J_\epsilon \mathbf{d}_j^1|_{H^s} \\
& \leq \frac{C(s)\mu_1}{\epsilon} |\mathbf{d}^1 - \mathbf{d}^2|_{H^s} \cdot C |\nabla^2 J_\epsilon \mathbf{d}^1|_{L^2}^3 |J_\epsilon \nabla \mathbf{v}^1|_{L^\infty} \\
& \quad + \frac{C(s)\mu_1}{\epsilon^{\frac{n}{2}+1}} |\mathbf{d}^1 - \mathbf{d}^2|_{L^2} \left(|J_\epsilon \mathbf{d}_p^1 J_\epsilon \mathbf{d}_i^1 J_\epsilon \mathbf{d}_j^1|_{L^\infty} |J_\epsilon (A_{kp}^1)_\epsilon|_{H^s} + |J_\epsilon (A_{kp}^1)_\epsilon|_{L^\infty} |J_\epsilon \mathbf{d}_p^1 J_\epsilon \mathbf{d}_i^1 J_\epsilon \mathbf{d}_j^1|_{H^s} \right) \\
& \leq \frac{C(s)\mu_1}{\epsilon} |\mathbf{d}^1 - \mathbf{d}^2|_{H^s} |\nabla^2 J_\epsilon \mathbf{d}^1|_{L^2}^3 |J_\epsilon \nabla \mathbf{v}^1|_{L^\infty} \\
& \quad + \frac{C(s)\mu_1}{\epsilon^{\frac{n}{2}+1}} |\mathbf{d}^1 - \mathbf{d}^2|_{L^2} \left(|\nabla^2 J_\epsilon \mathbf{d}^1|_{L^2}^3 |\nabla J_\epsilon \mathbf{v}^1|_{H^s} + |J_\epsilon \nabla \mathbf{v}^1|_{L^\infty} |J_\epsilon \mathbf{d}^1|_{L^\infty}^2 |J_\epsilon \mathbf{d}^1|_{H^s} \right) \\
& \leq \frac{C(s)\mu_1}{\epsilon^{\frac{n}{2}+5}} |\mathbf{d}^1 - \mathbf{d}^2|_{H^s} |\nabla \mathbf{d}^1|_{L^2} |\mathbf{v}^1|_{L^2} + \frac{C(s)\mu_1}{\epsilon^{\frac{n}{2}+s+5}} |\mathbf{d}^1 - \mathbf{d}^2|_{L^2} |\nabla \mathbf{d}^1|_{L^2}^3 |\mathbf{v}^1|_{L^2} \\
& \quad + \frac{C(s)\mu_1}{\epsilon^{n+s+3}} |\mathbf{d}^1 - \mathbf{d}^2|_{L^2} |\mathbf{v}^1|_{L^2} |\nabla \mathbf{d}^1|_{L^2}^2 \\
& \leq C(s, \mu_1, \frac{1}{\epsilon}) \left(|\nabla \mathbf{d}^1|_{L^2} |\mathbf{v}^1|_{L^2} + |\nabla \mathbf{d}^1|_{L^2}^3 |\mathbf{v}^1|_{L^2} + |\nabla \mathbf{d}^1|_{L^2}^2 |\mathbf{v}^1|_{L^2} \right) |\mathbf{d}^1 - \mathbf{d}^2|_{H^s},
\end{aligned}$$

and by similar estimating

$$\begin{aligned}
I_2 & \leq \frac{C(s)\mu_1}{\epsilon^{\frac{n}{2}+s+5}} |\nabla \mathbf{d}^1|_{L^2}^2 |\nabla \mathbf{d}^2|_{L^2} |\mathbf{v}^1|_{L^2} |\mathbf{d}^1 - \mathbf{d}^2|_{H^s}, \\
I_3 & \leq \frac{C(s)\mu_1}{\epsilon^{\frac{n}{2}+s+5}} |\nabla \mathbf{d}^1|_{L^2}^2 |\nabla \mathbf{d}^2|_{L^2}^2 |\mathbf{v}^1 - \mathbf{v}^2|_{H^s},
\end{aligned}$$

$$\begin{aligned} I_4 &\leq \frac{C(s)\mu_1}{\epsilon^{\frac{n}{2}+s+5}} |\nabla \mathbf{d}^1|_{L^2} |\nabla \mathbf{d}^2|_{L^2}^2 |\mathbf{v}^2|_{L^2} |\mathbf{d}^1 - \mathbf{d}^2|_{H^s}, \\ I_5 &\leq \frac{C(s)\mu_1}{\epsilon^{\frac{n}{2}+s+5}} |\nabla \mathbf{d}^2|_{L^2}^3 |\mathbf{v}^2|_{L^2} |\mathbf{d}^1 - \mathbf{d}^2|_{H^s}. \end{aligned}$$

Furthermore, similar arguments and Lemma 3.1 induce to the estimates for all $\mathbf{v}^1, \mathbf{v}^2, \dot{\mathbf{d}}^1, \mathbf{d}^1, \dot{\mathbf{d}}^2, \mathbf{d}^2 \in H^s$

$$\begin{aligned} &\left| F_\epsilon(\mathbf{v}^1, \dot{\mathbf{d}}^1, \mathbf{d}^1) - F_\epsilon(\mathbf{v}^2, \dot{\mathbf{d}}^2, \mathbf{d}^2) \right|_{H^s} \\ &\leq C\left(\frac{1}{\epsilon}\right) f(|\mathbf{v}^1|_{L^2} + |\mathbf{v}^2|_{L^2}, |\nabla \mathbf{d}^1|_{L^2} + |\nabla \mathbf{d}^2|_{L^2}, |\dot{\mathbf{d}}^1|_{L^2} + |\dot{\mathbf{d}}^2|_{L^2}) \\ &\quad \times (|\mathbf{v}^1 - \mathbf{v}^2|_{H^s} + |\mathbf{d}^1 - \mathbf{d}^2|_{H^s}), \end{aligned} \quad (3.4)$$

and

$$\begin{aligned} &\left| G_\epsilon(\mathbf{v}^1, \dot{\mathbf{d}}^1, \mathbf{d}^1) - G_\epsilon(\mathbf{v}^2, \dot{\mathbf{d}}^2, \mathbf{d}^2) \right|_{H^s} \\ &\leq C\left(\frac{1}{\epsilon}\right) g(|\mathbf{v}^1|_{L^2} + |\mathbf{v}^2|_{L^2}, |\dot{\mathbf{d}}^1|_{L^2} + |\dot{\mathbf{d}}^2|_{L^2}, |\nabla \mathbf{d}^1|_{L^2} + |\nabla \mathbf{d}^2|_{L^2}) \\ &\quad \times (|\mathbf{v}^1 - \mathbf{v}^2|_{H^s} + |\dot{\mathbf{d}}^1 - \dot{\mathbf{d}}^2|_{H^s} + |\mathbf{d}^1 - \mathbf{d}^2|_{H^s}) \end{aligned} \quad (3.5)$$

for some positive increasing functions $f(\cdot, \cdot, \cdot)$ and $g(\cdot, \cdot, \cdot)$ on their variables, and

$$\begin{aligned} &\left| H_\epsilon(\mathbf{v}^1, \dot{\mathbf{d}}^1, \mathbf{d}^1) - H_\epsilon(\mathbf{v}^2, \dot{\mathbf{d}}^2, \mathbf{d}^2) \right|_{H^s} \\ &\leq \left| \dot{\mathbf{d}}^1 - \dot{\mathbf{d}}^2 \right|_{H^s} + \frac{C}{\epsilon^{\frac{n}{2}+s+1}} (|\mathbf{v}^1|_{L^2} + |\mathbf{v}^2|_{L^2} + |\nabla \mathbf{d}^1|_{L^2} + |\nabla \mathbf{d}^2|_{L^2}) (|\mathbf{v}^1 - \mathbf{v}^2|_{H^s} + |\mathbf{d}^1 - \mathbf{d}^2|_{H^s}). \end{aligned} \quad (3.6)$$

Then combining the estimates (3.4), (3.5) and (3.6) implies that for all $\mathbf{v}^i, \dot{\mathbf{d}}^i, \mathbf{d}^i \in H^s$, $i = 1, 2$

$$\begin{aligned} &\left| \mathcal{F}_\epsilon(\mathbf{v}^1, \dot{\mathbf{d}}^1, \mathbf{d}^1) - \mathcal{F}_\epsilon(\mathbf{v}^2, \dot{\mathbf{d}}^2, \mathbf{d}^2) \right|_{H^s \times H^s \times H^s} \\ &\leq C\left(\frac{1}{\epsilon}\right) \mathfrak{f}(|\mathbf{v}^1|_{L^2} + |\mathbf{v}^2|_{L^2}, |\dot{\mathbf{d}}^1|_{L^2} + |\dot{\mathbf{d}}^2|_{L^2}, |\nabla \mathbf{d}^1|_{L^2} + |\nabla \mathbf{d}^2|_{L^2}) \\ &\quad \times (|\mathbf{v}^1 - \mathbf{v}^2|_{H^s} + |\dot{\mathbf{d}}^1 - \dot{\mathbf{d}}^2|_{H^s} + |\mathbf{d}^1 - \mathbf{d}^2|_{H^s}), \end{aligned} \quad (3.7)$$

where $\mathfrak{f}(\cdot, \cdot, \cdot)$ is a positive increasing function on its variables. Hence \mathcal{F}_ϵ is locally Lipschitz on any open subset $\mathcal{O}^M \subset H^s \times H^s \times H^s$, where

$$\mathcal{O}^M = \left\{ (\mathbf{v}, \dot{\mathbf{d}}, \mathbf{d})^\top \in H^s \times H^s \times H^s \mid |\mathbf{v}|_{H^s} + |\dot{\mathbf{d}}|_{H^s} + |\mathbf{d}|_{H^s} < M, \operatorname{div} \mathbf{v} = 0 \right\}$$

for any fixed number $M > 0$. Then the ODE theory implies that for any given $\mathbf{v}^{in}, \dot{\mathbf{d}}^{in}, \mathbf{d}^{in} \in H^s$ there exists a unique solution $(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon)^\top \in C([0, T_\epsilon]; H^s \times H^s \times H^s \cap \mathcal{O}^M)$ to the system (3.3) and then Proposition 3.1 holds. \square

Remark 3.1. This proposition is valid for general case $\mu_4 > 0, \mu_1, \mu_2, \mu_3, \mu_5, \mu_6 \geq 0$.

3.2. The global existence of the approximated system (3.1). Next we are going to extend globally the solutions constructed in Proposition 3.1 to the time interval $[0, \infty)$ under the assumption $\mu_4 > 2(\mu_2 + \mu_3 + \mu_5 + \mu_6) \geq 0$.

Proposition 3.2. *If $\mu_4 > 2(\mu_2 + \mu_3 + \mu_5 + \mu_6) \geq 0$, then solutions constructed in Proposition 3.1 can be extended to the time existed interval $[0, \infty)$.*

Proof. Case 1: $\mu_1 = \mu_2 = \mu_3 = \mu_5 = \mu_6 = 0, \mu_4 > 0$. Multiplying by \mathbf{v}^ϵ in the first equation of (3.1), integrating by parts, one derives that

$$\frac{1}{2} \frac{d}{dt} |\mathbf{v}^\epsilon|_{L^2}^2 + \mu |\nabla J_\epsilon \mathbf{v}^\epsilon|_{L^2}^2 = \langle J_\epsilon \nabla J_\epsilon \mathbf{d}^\epsilon \odot J_\epsilon \nabla J_\epsilon \mathbf{d}^\epsilon, \nabla J_\epsilon \mathbf{v}^\epsilon \rangle.$$

We multiply by $\dot{\mathbf{d}}^\epsilon$ in the third equation of (3.1), integrate by parts and then we have

$$\frac{1}{2} \frac{d}{dt} \left(|\dot{\mathbf{d}}^\epsilon|_{L^2}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{L^2}^2 \right) = \langle J_\epsilon \Delta J_\epsilon \mathbf{d}^\epsilon, J_\epsilon [J_\epsilon \mathbf{v}^\epsilon \cdot \nabla J_\epsilon \mathbf{d}^\epsilon] \rangle.$$

Noticing that the cancellation

$$\langle J_\epsilon \nabla J_\epsilon \mathbf{d}^\epsilon \odot J_\epsilon \nabla J_\epsilon \mathbf{d}^\epsilon, \nabla J_\epsilon \mathbf{v}^\epsilon \rangle + \langle J_\epsilon \Delta J_\epsilon \mathbf{d}^\epsilon, J_\epsilon [J_\epsilon \mathbf{v}^\epsilon \cdot \nabla J_\epsilon \mathbf{d}^\epsilon] \rangle = 0,$$

one knows that

$$\frac{1}{2} \frac{d}{dt} \left(|\mathbf{v}^\epsilon|_{L^2}^2 + |\dot{\mathbf{d}}^\epsilon|_{L^2}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{L^2}^2 \right) + \frac{1}{2} \mu_4 |\nabla J_\epsilon \mathbf{v}^\epsilon|_{L^2}^2 = 0,$$

which immediately implies that the solution $(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon)^\top$ to the system (3.1) exists on any interval $[0, T]$, and

$$\sup_{0 \leq t \leq T} \left(|\mathbf{v}^\epsilon|_{L^2}^2 + |\dot{\mathbf{d}}^\epsilon|_{L^2}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{L^2}^2 \right) \leq 2 \left(|\mathbf{v}^{in}|_{L^2} + |\dot{\mathbf{d}}^{in}|_{L^2} + |\nabla \mathbf{d}^{in}|_{L^2} \right). \quad (3.8)$$

Letting $(\mathbf{v}^1, \dot{\mathbf{d}}^1, \mathbf{d}^1)^\top = (\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon)^\top$ be the solution to the system (3.1) obtained in Proposition 3.1 on the interval $[0, T_\epsilon]$ and $(\mathbf{v}^2, \dot{\mathbf{d}}^2, \mathbf{d}^2)^\top = (0, 0, 0)^\top$ in the relation (3.7), one has

$$\left| \mathcal{F}_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon) \right|_{H^s \times H^s \times H^s} \leq C(\frac{1}{\epsilon}) f(|\mathbf{v}^\epsilon|_{L^2}, |\dot{\mathbf{d}}^\epsilon|_{L^2}, |\nabla \mathbf{d}^\epsilon|_{L^2}) \left(|\mathbf{v}^\epsilon|_{H^s} + |\dot{\mathbf{d}}^\epsilon|_{H^s} + |\mathbf{d}^\epsilon|_{H^s} \right). \quad (3.9)$$

Recalling that

$$\frac{d}{dt} \begin{pmatrix} \mathbf{v}^\epsilon \\ \dot{\mathbf{d}}^\epsilon \\ \mathbf{d}^\epsilon \end{pmatrix} = \mathcal{F}_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon),$$

which implies that by multiplying by $(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon)^\top$ and combining the bounds (3.8) and (3.9)

$$\frac{d}{dt} \left| \begin{pmatrix} \mathbf{v}^\epsilon \\ \dot{\mathbf{d}}^\epsilon \\ \mathbf{d}^\epsilon \end{pmatrix} \right|_{H^s \times H^s \times H^s}^2 \leq C(\epsilon, |\mathbf{v}^{in}|_{L^2}, |\dot{\mathbf{d}}^{in}|_{L^2}, |\nabla \mathbf{d}^{in}|_{L^2}) \left(|\mathbf{v}^\epsilon|_{H^s}^2 + |\dot{\mathbf{d}}^\epsilon|_{H^s}^2 + |\mathbf{d}^\epsilon|_{H^s}^2 \right).$$

Then Gronwall inequality deduces to

$$\sup_{0 \leq t \leq T} \left(|\mathbf{v}^\epsilon|_{H^s}^2 + |\dot{\mathbf{d}}^\epsilon|_{H^s}^2 + |\mathbf{d}^\epsilon|_{H^s}^2 \right) \leq \left(|\mathbf{v}^{in}|_{H^s}^2 + |\dot{\mathbf{d}}^{in}|_{H^s}^2 + |\mathbf{d}^{in}|_{H^s}^2 \right) e^{CT}$$

for any $T > 0$. By standard extension methods, the solutions to the system (3.1) constructed in Proposition 3.1 can be globally extended to $[0, \infty)$.

Case 2: $\mu_4 > 2(\mu_2 + \mu_3 + \mu_5 + \mu_6) \geq 0$, $\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6 > 0$. As shown in Section 2, we can get the L^2 -estimate of the approximated system (3.1)

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(|\mathbf{v}^\epsilon|_{L^2}^2 + |\dot{\mathbf{d}}^\epsilon|_{L^2}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{L^2}^2 \right) + \frac{1}{2} \mu_4 |\nabla J_\epsilon \mathbf{v}^\epsilon|_{L^2}^2 - \lambda_1 |\dot{\mathbf{d}}^\epsilon|_{L^2}^2 + \mu_1 |J_\epsilon \mathbf{d}^{\epsilon\top} J_\epsilon A_\epsilon J_\epsilon \mathbf{d}^\epsilon|_{L^2}^2 \\ & \leq (|\lambda_1| + |\lambda_2| + \mu_2 + \mu_3) |\mathbf{d}^\epsilon|_{L^\infty} |\dot{\mathbf{d}}^\epsilon|_{L^2} |\nabla J_\epsilon \mathbf{v}^\epsilon|_{L^2} + (\mu_2 + \mu_3 + \mu_5 + \mu_6) |\mathbf{d}^\epsilon|_{L^\infty}^2 |\nabla J_\epsilon \mathbf{v}^\epsilon|_{L^2}^2. \end{aligned} \quad (3.10)$$

Since $\mu_4 > 2(\mu_2 + \mu_3 + \mu_5 + \mu_6) \geq 0$ and $|\mathbf{d}^\epsilon|_{L^\infty} \leq |\mathbf{d}|_{L^\infty} = 1$, it is derived from (3.10) that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(|\mathbf{v}^\epsilon|_{L^2}^2 + |\dot{\mathbf{d}}^\epsilon|_{L^2}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{L^2}^2 \right) + \mu_1 |J_\epsilon \mathbf{d}^{\epsilon\top} J_\epsilon A_\epsilon J_\epsilon \mathbf{d}^\epsilon|_{L^2}^2 \\ & + \left(\frac{1}{2} \mu_4 - (\mu_2 + \mu_3 + \mu_5 + \mu_6) \right) |\nabla J_\epsilon \mathbf{v}^\epsilon|_{L^2}^2 \\ & \leq |\lambda_1| |\dot{\mathbf{d}}^\epsilon|_{L^2}^2 + \frac{|\lambda_1| + |\lambda_2| + \mu_2 + \mu_3}{\mu_4 - 2(\mu_2 + \mu_3 + \mu_5 + \mu_6)} |\dot{\mathbf{d}}^\epsilon|_{L^2}^2 \\ & + \frac{1}{2} \left(\frac{1}{2} \mu_4 - (\mu_2 + \mu_3 + \mu_5 + \mu_6) \right) |\nabla J_\epsilon \mathbf{v}^\epsilon|_{L^2}^2, \end{aligned}$$

which immediately means that

$$\frac{1}{2} \frac{d}{dt} \left(|\mathbf{v}^\epsilon|_{L^2}^2 + |\dot{\mathbf{d}}^\epsilon|_{L^2}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{L^2}^2 \right) + \mu_1 |J_\epsilon \mathbf{d}^{\epsilon\top} J_\epsilon A_\epsilon J_\epsilon \mathbf{d}^\epsilon|_{L^2}^2$$

$$\begin{aligned}
& + \frac{1}{2} \left(\frac{1}{2} \mu_4 - (\mu_2 + \mu_3 + \mu_5 + \mu_6) \right) |\nabla J_\epsilon \mathbf{v}^\epsilon|_{L^2}^2 \\
& \leq \left(|\lambda_1| + \frac{|\lambda_1| + |\lambda_2| + \mu_2 + \mu_3}{\mu_4 - 2(\mu_2 + \mu_3 + \mu_5 + \mu_6)} \right) |\dot{\mathbf{d}}^\epsilon|_{L^2}^2.
\end{aligned} \tag{3.11}$$

This implies by Gronwall inequality that the solution $(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon)^\top$ to the system (3.1) exists on any interval $[0, T]$, and

$$\sup_{0 \leq t \leq T} \left(|\mathbf{v}^\epsilon|_{L^2}^2 + |\dot{\mathbf{d}}^\epsilon|_{L^2}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{L^2}^2 \right) \leq \left(|\mathbf{v}^{in}|_{L^2} + |\dot{\mathbf{d}}^{in}|_{L^2} + |\nabla \mathbf{d}^{in}|_{L^2} \right) e^{CT}, \tag{3.12}$$

where $C = 2 \left(|\lambda_1| + \frac{|\lambda_1| + |\lambda_2| + \mu_2 + \mu_3}{\mu_4 - 2(\mu_2 + \mu_3 + \mu_5 + \mu_6)} \right) > 0$. Then by the same remained arguments in Case 1 we can finish the proof of this proposition. \square

4. THE PROOF OF THEOREM 1.1.

In this section, we are going to show the main theorem by analyzing some results of compactness and then passing to the limit on the approximated system (3.1). Recall that we have

defined the norms on a product space $X \times X \times X$ as $\left| \begin{pmatrix} a \\ b \\ c \end{pmatrix} \right|_{X \times X \times X} \equiv |a|_X + |b|_X + |c|_X$.

4.1. The proof of Part (I) of Theorem 1.1. By the apriori estimate (2.1), we know that for some constant $C = C(n, s) > 0$ and for all $\epsilon > 0$

$$\begin{aligned}
& \frac{d}{dt} \left(|\mathbf{v}^\epsilon|_{H^s}^2 + |\dot{\mathbf{d}}^\epsilon|_{H^s}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s}^2 \right) + \mu_4 |\nabla J_\epsilon \mathbf{v}^\epsilon|_{H^s}^2 \\
& - 2\lambda_1 |\dot{\mathbf{d}}^\epsilon|_{H^s}^2 + 2\mu_1 \sum_{k=0}^s |J_\epsilon \mathbf{d}^{\epsilon \top} (\nabla^{k+1} J_\epsilon \mathbf{v}^\epsilon) J_\epsilon \mathbf{d}^\epsilon|_{L^2}^2 \\
& \leq C(1 + |\lambda_1| + |\lambda_2| + \mu_2 + \mu_3) \left(|\mathbf{v}^\epsilon|_{H^s}^2 + |\dot{\mathbf{d}}^\epsilon|_{H^s}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s}^2 \right) |\nabla J_\epsilon \mathbf{v}^\epsilon|_{H^s} \\
& + C(1 + |\lambda_1| + |\lambda_2|) \left(|\dot{\mathbf{d}}^\epsilon|_{H^s}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s}^2 \right)^2 \\
& + C(\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6) |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s}^2 |\nabla J_\epsilon \mathbf{v}^\epsilon|_{H^s}^2.
\end{aligned} \tag{4.1}$$

Denoting by $E_\epsilon(t) \equiv |\mathbf{v}^\epsilon|_{H^s}^2(t) + |\dot{\mathbf{d}}^\epsilon|_{H^s}^2(t) + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s}^2(t)$ and $F_\epsilon(t) \equiv |\nabla J_\epsilon \mathbf{v}^\epsilon|_{H^s}^2$, the inequalities (4.1) can be rewritten as

$$\begin{aligned}
\frac{d}{dt} E_\epsilon(t) + \mu_4 F_\epsilon(t) & \leq C(1 + |\lambda_1| + |\lambda_2| + \mu_2 + \mu_3) E_\epsilon(t) F_\epsilon^{\frac{1}{2}}(t) + 2|\lambda_1| E_\epsilon(t) \\
& + C(1 + |\lambda_1| + |\lambda_2|) E_\epsilon^2(t) + C(\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6) E_\epsilon(t) F_\epsilon(t),
\end{aligned}$$

and then by Cauchy inequality

$$\frac{d}{dt} E_\epsilon(t) + \frac{1}{2} \mu_4 F_\epsilon(t) \leq 2|\lambda_1| E_\epsilon(t) + C_1 E_\epsilon^2(t) + C(\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6) E_\epsilon(t) F_\epsilon(t), \tag{4.2}$$

where the constant $C_1 = C(1 + |\lambda_1| + |\lambda_2|) + \frac{C^2(1 + |\lambda_1| + |\lambda_2| + \mu_2 + \mu_3)^2}{2\mu_4} > 0$. Recalling that $E_\epsilon(0) = E^{in} \leq C_2 \equiv \frac{\mu_4}{4C(1 + \mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6)}$, we define $T > 0$ such that $E_\epsilon(t) \leq \frac{4}{3} C_2$ for all $t \in [0, T]$. Then the inequalities (4.2) imply that

$$\frac{d}{dt} E_\epsilon(t) + \frac{1}{6} \mu_4 F_\epsilon(t) \leq 2|\lambda_1| E_\epsilon(t) + C_1 E_\epsilon^2(t),$$

which immediately imply that by Gronwall inequality

$$E_\epsilon(t) \leq C_3(E^{in}, T) \equiv \frac{E^{in}(2|\lambda_1| + 1)e^{(2|\lambda_1|+1)T}}{C_1 E^{in} + 2|\lambda_1| + 1 - C_1 E^{in} e^{(2|\lambda_1|+1)T}}$$

for all $t \in [0, T]$. Here the relation $C_1 E^{in} + 2|\lambda_1| + 1 - C_1 E^{in} e^{(2|\lambda_1|+1)T} > 0$, i.e., $T < \frac{1}{2|\lambda_1|+1} \ln \left(1 + \frac{2|\lambda_1|+1}{C_1 E^{in}} \right)$ is required. Also, by the definition of $T > 0$, we need

$$E_\epsilon(t) \leq \frac{E^{in}(2|\lambda_1|+1)e^{(2|\lambda_1|+1)T}}{C_1 E^{in} + 2|\lambda_1| + 1 - C_1 E^{in} e^{(2|\lambda_1|+1)T}} \leq \frac{4}{3} C_2,$$

hence

$$T \leq \frac{1}{2|\lambda_1|+1} \ln \frac{\frac{4}{3} C_2 (C_1 E^{in} + 2|\lambda_1| + 1)}{(2|\lambda_1| + 1 + \frac{4}{3} C_1 C_2) E^{in}}.$$

So, we can take $T > 0$ such that

$$0 < T < \frac{1}{2|\lambda_1|+1} \min \left\{ \ln \left(1 + \frac{2|\lambda_1|+1}{C_1 E^{in}} \right), \ln \frac{\frac{4}{3} C_2 (C_1 E^{in} + 2|\lambda_1| + 1)}{(2|\lambda_1| + 1 + \frac{4}{3} C_1 C_2) E^{in}} \right\},$$

and furthermore for all $t \in [0, T]$

$$|\mathbf{v}^\epsilon|_{H^s}^2(t) + |\dot{\mathbf{d}}^\epsilon|_{H^s}^2(t) + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s}^2(t) + \frac{1}{6} \mu_4 \int_0^t |\nabla J_\epsilon \mathbf{v}^\epsilon|_{H^s}^2(s) ds \leq C^*(E^{in}, T), \quad (4.3)$$

where $C^*(E^{in}, T) = E^{in} + [2|\lambda_1|C_3(E^{in}, T) + C_1(C_3(E^{in}, T))^2]T > 0$. Thus we know that there exist functions $\mathbf{v} \in L^\infty(0, T; H^s) \cap L^2(0, T; H^{s+1})$, $\mathbf{d} \in L^\infty(0, T; H^{s+1})$ and $\mathbf{w} \in L^\infty(0, T; H^s)$ such that (extracting subsequence if necessary)

$$\begin{cases} \mathbf{v}^\epsilon \rightharpoonup \mathbf{v} \text{ weakly* in } L^\infty(0, T; H^s), \\ \mathbf{v}^\epsilon \rightharpoonup \mathbf{v} \text{ weakly in } L^2(0, T; H^{s+1}), \\ \mathbf{d}^\epsilon \rightharpoonup \mathbf{d} \text{ weakly* in } L^\infty(0, T; H^{s+1}), \\ \dot{\mathbf{d}}^\epsilon \rightharpoonup \mathbf{w} \text{ weakly* in } L^\infty(0, T; H^s). \end{cases} \quad (4.4)$$

Noticing that $s > \frac{n}{2} + 2$, one can estimate by Lemma 3.1, the equations (3.1), the bounds (4.3) and Sobolev embedding theory that

$$\begin{aligned} \sup_{0 \leq t \leq T} |\partial_t \mathbf{d}^\epsilon|_{H^s} &\leq \sup_{0 \leq t \leq T} |\dot{\mathbf{d}}^\epsilon|_{H^s} + \sup_{0 \leq t \leq T} |J_\epsilon \mathbf{v}^\epsilon \cdot \nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s} \\ &\leq \sup_{0 \leq t \leq T} |\dot{\mathbf{d}}^\epsilon|_{H^s} + C \sup_{0 \leq t \leq T} (|J_\epsilon \mathbf{v}^\epsilon|_{L^\infty} |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s} + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{L^\infty} |J_\epsilon \mathbf{v}^\epsilon|_{H^s}) \\ &\leq \sup_{0 \leq t \leq T} |\dot{\mathbf{d}}^\epsilon|_{H^s} + C \sup_{0 \leq t \leq T} |\mathbf{v}^\epsilon|_{H^s} \sup_{0 \leq t \leq T} |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s} \\ &\leq C(E^{in}, T), \end{aligned}$$

and

$$\begin{aligned} |\partial_t \mathbf{v}^\epsilon|_{H^{s-2}} &\leq |\mathcal{P} J_\epsilon [J_\epsilon \mathbf{v}^\epsilon \cdot \nabla J_\epsilon \mathbf{v}^\epsilon]|_{H^{s-2}} + \frac{1}{2} \mu_4 |J_\epsilon (\Delta J_\epsilon \mathbf{v}^\epsilon)|_{H^{s-2}} \\ &\quad + |\mathcal{P} J_\epsilon \operatorname{div}(J_\epsilon \nabla J_\epsilon \mathbf{d}^\epsilon \odot J_\epsilon \nabla J_\epsilon \mathbf{d}^\epsilon)|_{H^{s-2}} + |\mathcal{P} J_\epsilon \mathcal{L}(J_\epsilon \mathbf{d}^\epsilon, J_\epsilon \mathbf{v}^\epsilon)|_{H^{s-2}} \\ &\leq |J_\epsilon \mathbf{v}^\epsilon \cdot \nabla J_\epsilon \mathbf{v}^\epsilon|_{H^{s-2}} + \frac{1}{4} \mu_4 |\mathbf{v}^\epsilon|_{H^s} + |\operatorname{div}(J_\epsilon \nabla J_\epsilon \mathbf{d}^\epsilon \odot J_\epsilon \nabla J_\epsilon \mathbf{d}^\epsilon)|_{H^{s-2}} \\ &\quad + |\mathcal{P} J_\epsilon \mathcal{L}(J_\epsilon \mathbf{d}^\epsilon, J_\epsilon \mathbf{v}^\epsilon)|_{H^{s-2}} \\ &\leq C |\mathbf{v}^\epsilon|_{H^s}^2 + \frac{1}{2} \mu_4 |\mathbf{v}^\epsilon|_{H^s} + C |\nabla \mathbf{d}^\epsilon|_{H^s}^2 + |\mathcal{P} J_\epsilon \mathcal{L}(J_\epsilon \mathbf{d}^\epsilon, J_\epsilon \mathbf{v}^\epsilon)|_{H^{s-2}} \\ &\leq C(E^{in}, T) + |\mathcal{P} J_\epsilon \mathcal{L}(J_\epsilon \mathbf{d}^\epsilon, J_\epsilon \mathbf{v}^\epsilon)|_{H^{s-2}}, \end{aligned}$$

where

$$\begin{aligned} &|\mathcal{P} J_\epsilon \mathcal{L}(J_\epsilon \mathbf{d}^\epsilon, J_\epsilon \mathbf{v}^\epsilon)|_{H^{s-2}} \\ &\leq \mu_1 |(J_\epsilon \mathbf{d}^\epsilon \otimes J_\epsilon \mathbf{d}^\epsilon : J_\epsilon A_\epsilon) J_\epsilon \mathbf{d}^\epsilon \otimes J_\epsilon \mathbf{d}^\epsilon|_{H^{s-1}} + (\mu_2 + \mu_3) \left| J_\epsilon \mathbf{d}^\epsilon \otimes J_\epsilon \dot{\mathbf{d}}^\epsilon \right|_{H^{s-1}} \\ &\quad + (\mu_2 + \mu_3) \left| J_\epsilon \mathbf{d}^\epsilon \otimes J_\epsilon \dot{\mathbf{d}}^\epsilon : J_\epsilon B_\epsilon \right|_{H^{s-1}} + (\mu_5 + \mu_6) |J_\epsilon \mathbf{d}^\epsilon \otimes J_\epsilon \mathbf{d}^\epsilon : J_\epsilon A_\epsilon|_{H^{s-1}} \end{aligned}$$

$$\begin{aligned}
&\leq C\mu_1 \left(|J_\epsilon \mathbf{d}\epsilon|_{L^\infty}^4 |J_\epsilon A_\epsilon|_{H^{s-1}} + |J_\epsilon \mathbf{d}\epsilon|_{L^\infty}^3 |J_\epsilon \dot{\mathbf{d}}\epsilon|_{H^{s-1}} |J_\epsilon A_\epsilon|_{L^\infty} \right) \\
&\quad + C(\mu_2 + \mu_3) \left(|J_\epsilon \mathbf{d}\epsilon|_{L^\infty} |J_\epsilon \dot{\mathbf{d}}\epsilon|_{H^{s-1}} + |J_\epsilon \mathbf{d}\epsilon|_{H^{s-1}} |J_\epsilon \dot{\mathbf{d}}\epsilon|_{L^\infty} \right) \\
&\quad + C(\mu_2 + \mu_3) \left(|J_\epsilon \mathbf{d}\epsilon|_{L^\infty} |J_\epsilon \dot{\mathbf{d}}\epsilon|_{L^\infty} |J_\epsilon B_\epsilon|_{H^{s-1}} + |J_\epsilon \mathbf{d}\epsilon|_{L^\infty} |J_\epsilon \dot{\mathbf{d}}\epsilon|_{H^{s-1}} |J_\epsilon B_\epsilon|_{L^\infty} \right. \\
&\quad \quad \quad \left. + |J_\epsilon \mathbf{d}\epsilon|_{H^{s-1}} |J_\epsilon \dot{\mathbf{d}}\epsilon|_{L^\infty} |J_\epsilon B_\epsilon|_{L^\infty} \right) \\
&\quad + C(\mu_5 + \mu_6) \left(|J_\epsilon \mathbf{d}\epsilon|_{L^\infty}^2 |J_\epsilon A_\epsilon|_{H^{s-1}} + |J_\epsilon \mathbf{d}\epsilon|_{L^\infty} |J_\epsilon \mathbf{d}\epsilon|_{H^{s-1}} |J_\epsilon A_\epsilon|_{L^\infty} \right) \\
&\leq C\mu_1 |\mathbf{d}\epsilon|_{H^{s+1}}^4 |\mathbf{v}^\epsilon|_{H^s} + C(\mu_5 + \mu_6) |\mathbf{d}\epsilon|_{H^{s+1}}^2 |\mathbf{v}^\epsilon|_{H^s} \\
&\quad + C(\mu_2 + \mu_3) \left(|\mathbf{d}\epsilon|_{H^{s+1}} |\dot{\mathbf{d}}\epsilon|_{H^s} + |\mathbf{d}\epsilon|_{H^{s+1}} |\dot{\mathbf{d}}\epsilon|_{H^s} |\mathbf{v}^\epsilon|_{H^s} \right) \\
&\leq C(E^{in}, T)(\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6).
\end{aligned}$$

Namely, we have obtained the bounds

$$\sup_{0 \leq t \leq T} |\partial_t \mathbf{d}\epsilon|_{H^s} \leq C(E^{in}, T), \quad (4.5)$$

and

$$\sup_{0 \leq t \leq T} |\partial_t \mathbf{v}^\epsilon|_{H^{s-2}} \leq C(E^{in}, T)(1 + \mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6). \quad (4.6)$$

Consequently, combining the construction of the approximated equations (3.1) we know that $\partial_t \mathbf{d}\epsilon \rightharpoonup \partial_t \mathbf{d}$ weakly in $L^\infty(0, T; H^s)$ and $\partial_t \mathbf{v}^\epsilon \rightharpoonup \partial_t \mathbf{v}$ weakly in $L^\infty(0, T; H^{s-2})$, where the limits are considered in distributional sense.

We also assert that the sequences $\{\mathbf{v}^\epsilon\}$, $\{\dot{\mathbf{d}}\epsilon\}$ and $\{\mathbf{d}\epsilon\}$ form Cauchy sequences in $C([0, T]; L^2(\Omega))$. Namely, there exists a constant $C = C(E^{in}, T) > 0$ such that for any $\epsilon, \epsilon' > 0$

$$\sup_{0 \leq t \leq T} \left| \begin{pmatrix} \mathbf{v}^\epsilon - \mathbf{v}^{\epsilon'} \\ \dot{\mathbf{d}}\epsilon - \dot{\mathbf{d}}^{\epsilon'} \\ \mathbf{d}\epsilon - \mathbf{d}^{\epsilon'} \end{pmatrix} \right|_{L^2 \times L^2 \times L^2} \leq C \max(\epsilon, \epsilon'). \quad (4.7)$$

Indeed, the equations (3.3) reduce to

$$\frac{d}{dt} \begin{pmatrix} \mathbf{v}^\epsilon - \mathbf{v}^{\epsilon'} \\ \dot{\mathbf{d}}\epsilon - \dot{\mathbf{d}}^{\epsilon'} \\ \mathbf{d}\epsilon - \mathbf{d}^{\epsilon'} \end{pmatrix} = \mathcal{F}_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}\epsilon, \mathbf{d}\epsilon) - \mathcal{F}_{\epsilon'}(\mathbf{v}^{\epsilon'}, \dot{\mathbf{d}}^{\epsilon'}, \mathbf{d}^{\epsilon'}).$$

Multiplying by $(\mathbf{v}^\epsilon - \mathbf{v}^{\epsilon'}, \dot{\mathbf{d}}\epsilon - \dot{\mathbf{d}}^{\epsilon'}, \mathbf{d}\epsilon - \mathbf{d}^{\epsilon'})^\top$ in the above equations yields that

$$\frac{d}{dt} \left| \begin{pmatrix} \mathbf{v}^\epsilon - \mathbf{v}^{\epsilon'} \\ \dot{\mathbf{d}}\epsilon - \dot{\mathbf{d}}^{\epsilon'} \\ \mathbf{d}\epsilon - \mathbf{d}^{\epsilon'} \end{pmatrix} \right|_{L^2 \times L^2 \times L^2} \leq \left| \mathcal{F}_\epsilon(\mathbf{v}^\epsilon, \dot{\mathbf{d}}\epsilon, \mathbf{d}\epsilon) - \mathcal{F}_{\epsilon'}(\mathbf{v}^{\epsilon'}, \dot{\mathbf{d}}^{\epsilon'}, \mathbf{d}^{\epsilon'}) \right|_{L^2 \times L^2 \times L^2}.$$

It can be estimated by Lemma 3.1 and similar calculation in Section 3.1 that

$$\frac{d}{dt} \left| \begin{pmatrix} \mathbf{v}^\epsilon - \mathbf{v}^{\epsilon'} \\ \dot{\mathbf{d}}\epsilon - \dot{\mathbf{d}}^{\epsilon'} \\ \mathbf{d}\epsilon - \mathbf{d}^{\epsilon'} \end{pmatrix} \right|_{L^2 \times L^2 \times L^2} \leq C(|\mathbf{v}^\epsilon|_{H^s}, |\dot{\mathbf{d}}\epsilon|_{H^s}, |\mathbf{d}\epsilon|_{H^s}) \left(\max(\epsilon, \epsilon') + \left| \begin{pmatrix} \mathbf{v}^\epsilon - \mathbf{v}^{\epsilon'} \\ \dot{\mathbf{d}}\epsilon - \dot{\mathbf{d}}^{\epsilon'} \\ \mathbf{d}\epsilon - \mathbf{d}^{\epsilon'} \end{pmatrix} \right|_{L^2 \times L^2 \times L^2} \right),$$

which implies that by the inequality (4.3) and Gronwall inequality

$$\sup_{0 \leq t \leq T} \left| \begin{pmatrix} \mathbf{v}^\epsilon - \mathbf{v}^{\epsilon'} \\ \dot{\mathbf{d}}\epsilon - \dot{\mathbf{d}}^{\epsilon'} \\ \mathbf{d}\epsilon - \mathbf{d}^{\epsilon'} \end{pmatrix} \right|_{L^2 \times L^2 \times L^2} \leq e^{C(E^{in}T)} \max(\epsilon, \epsilon'),$$

and the assertion (4.7) is valid.

So the bound (4.7) tells us that $\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon$ and \mathbf{d}^ϵ strongly converge to \mathbf{v}, \mathbf{w} and \mathbf{d} in $C([0, T]; L^2(\Omega))$ respectively and

$$\sup_{0 \leq t \leq T} \left(|\mathbf{v}^\epsilon - \mathbf{v}|_{L^2} + |\dot{\mathbf{d}}^\epsilon - \mathbf{w}|_{L^2} + |\mathbf{d}^\epsilon - \mathbf{d}|_{L^2} \right) \leq C\epsilon.$$

Then the interpolation inequality in Sobolev space implies that for all $0 < s' < s$

$$\begin{aligned} & \sup_{0 \leq t \leq T} \left(|\mathbf{v}^\epsilon - \mathbf{v}|_{H^{s'}} + |\dot{\mathbf{d}}^\epsilon - \mathbf{w}|_{H^{s'}} + |\mathbf{d}^\epsilon - \mathbf{d}|_{H^{s'}} \right) \\ & \leq C \sup_{0 \leq t \leq T} |\mathbf{v}^\epsilon - \mathbf{v}|_{L^2}^{1-\frac{s'}{s}} \sup_{0 \leq t \leq T} |\mathbf{v}^\epsilon - \mathbf{v}|_{H^s}^{\frac{s'}{s}} + C \sup_{0 \leq t \leq T} |\dot{\mathbf{d}}^\epsilon - \mathbf{w}|_{L^2}^{1-\frac{s'}{s}} \sup_{0 \leq t \leq T} |\dot{\mathbf{d}}^\epsilon - \mathbf{w}|_{H^s}^{\frac{s'}{s}} \\ & \quad + C \sup_{0 \leq t \leq T} |\mathbf{d}^\epsilon - \mathbf{d}|_{L^2}^{1-\frac{s'}{s}} \sup_{0 \leq t \leq T} |\mathbf{d}^\epsilon - \mathbf{d}|_{H^s}^{\frac{s'}{s}} \\ & \leq C\epsilon^{1-\frac{s'}{s}}, \end{aligned}$$

which immediately means that $\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon$ and \mathbf{d}^ϵ converge strongly to \mathbf{v}, \mathbf{w} and \mathbf{d} in $L^\infty(0, T; H^{s'})$ respectively for all $0 < s' < s$. Consequently, one also easily know that $J_\epsilon[J_\epsilon \mathbf{v}^\epsilon \cdot \nabla J_\epsilon \mathbf{d}^\epsilon]$ strongly converges to $\mathbf{v} \cdot \nabla \mathbf{d}$ in $L^\infty(0, T; H^{s'})$ for all $0 < s' < s$. Then we have that, in $L^\infty(0, T; H^{s'})$, $\mathbf{w} = \lim_{\epsilon \rightarrow 0} \dot{\mathbf{d}}^\epsilon = \lim_{\epsilon \rightarrow 0} \partial_t \mathbf{d}^\epsilon + \mathbf{v} \cdot \nabla \mathbf{d}$, that is, $\partial_t \mathbf{d}^\epsilon$ converges strongly to $\mathbf{w} - \mathbf{v} \cdot \nabla \mathbf{d}$ in $L^\infty(0, T; H^{s'})$ for all $0 < s' < s$. Recalling that $\partial_t \mathbf{d}^\epsilon \rightharpoonup \partial_t \mathbf{d}$ weakly in $L^\infty(0, T; H^s)$ in the sense of distribution, we know that $\partial_t \mathbf{d} = \mathbf{w} - \mathbf{v} \cdot \nabla \mathbf{d} \in L^\infty(0, T; H^{s'})$, namely, $\mathbf{w} = \dot{\mathbf{d}}$.

Since $s > s' \geq \frac{n}{2} + 2$, the space $H^{s'}$ embeds into C^2 , which means that $(\mathbf{v}^\epsilon, \dot{\mathbf{d}}^\epsilon, \mathbf{d}^\epsilon)$ converge strongly to $(\mathbf{v}, \dot{\mathbf{d}}, \mathbf{d})$ in $C([0, T]; C^2(\Omega))$. Therefore by passing to the limit as $\epsilon \rightarrow 0$ in the approximated system (3.1), we know that (\mathbf{v}, \mathbf{d}) is a solution of the equations (1.1). Furthermore, by Fatou lemma, the bound (4.3) implies that

$$|\mathbf{v}|_{H^s}^2(t) + |\dot{\mathbf{d}}|_{H^s}^2(t) + |\nabla \mathbf{d}|_{H^s}^2(t) + \frac{1}{6}\mu_4 \int_0^t |\nabla \mathbf{v}|_{H^s}^2(s) ds \leq C^*(E^{in}, T)$$

for all $t \in [0, T]$. The proof of Part (I) of Theorem 1.1 is finished.

4.2. The proof of Part (II) of Theorem 1.1. Since $\mu_1 = \mu_2 = \mu_3 = \mu_5 = \mu_6 = 0$, by the apriori estimates in Remark 2.2.(1) in Section 2, we get the relation for some constant $C = C(n, s) > 0$

$$\begin{aligned} & \frac{d}{dt} \left(|\mathbf{v}^\epsilon|_{H^s}^2 + |\dot{\mathbf{d}}^\epsilon|_{H^s}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s}^2 \right) + \mu_4 |\nabla J_\epsilon \mathbf{v}^\epsilon|_{H^s}^2 \\ & \leq C \left(|\mathbf{v}^\epsilon|_{H^s}^2 + |\dot{\mathbf{d}}^\epsilon|_{H^s}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s}^2 \right) |\nabla J_\epsilon \mathbf{v}^\epsilon|_{H^s} + C \left(|\dot{\mathbf{d}}^\epsilon|_{H^s}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s}^2 \right)^2 \\ & \leq C_4 \left(|\mathbf{v}^\epsilon|_{H^s}^2 + |\dot{\mathbf{d}}^\epsilon|_{H^s}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s}^2 \right)^2 + \frac{1}{2}\mu_4 |\nabla J_\epsilon \mathbf{v}^\epsilon|_{H^s}^2, \end{aligned}$$

where the latter inequality is deduced by Cauchy inequality and the constant $C_4 = C(\frac{C}{2\mu_4} + 1) > 0$. Namely, the following inequality is valid:

$$\frac{d}{dt} \left(|\mathbf{v}^\epsilon|_{H^s}^2 + |\dot{\mathbf{d}}^\epsilon|_{H^s}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s}^2 \right) + \frac{1}{2}\mu_4 |\nabla J_\epsilon \mathbf{v}^\epsilon|_{H^s}^2 \leq C_4 \left(|\mathbf{v}^\epsilon|_{H^s}^2 + |\dot{\mathbf{d}}^\epsilon|_{H^s}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon|_{H^s}^2 \right)^2. \quad (4.8)$$

Thus provided the initial data $E^{in} \equiv |\mathbf{v}^{in}|_{H^s}^2 + |\dot{\mathbf{d}}^{in}|_{H^s}^2 + |\nabla \mathbf{d}^{in}|_{H^s}^2 < \infty$, it is derived from Gronwall inequality that for any $T \in (0, \frac{1}{C_4 E^{in}})$ and for all $0 \leq t \leq T$

$$|\mathbf{v}^\epsilon(t)|_{H^s}^2 + |\dot{\mathbf{d}}^\epsilon(t)|_{H^s}^2 + |\nabla J_\epsilon \mathbf{d}^\epsilon(t)|_{H^s}^2 + \frac{1}{2}\mu_4 \int_0^t |\nabla J_\epsilon \mathbf{v}^\epsilon|_{H^s}^2 ds \leq E^{in} + \frac{C_4(E^{in})^2 T}{(1 - C_4 E^{in} T)^2}, \quad (4.9)$$

which means that there exist functions $\mathbf{v} \in L^\infty(0, T; H^s) \cap L^2(0, T; H^{s+1})$, $\mathbf{d} \in L^\infty(0, T; H^{s+1})$ and $\mathbf{w} \in L^\infty(0, T; H^s)$ such that (extracting subsequence if necessary)

$$\begin{cases} \mathbf{v}^\epsilon \rightharpoonup \mathbf{v} \text{ weakly* in } L^\infty(0, T; H^s), \\ \mathbf{v}^\epsilon \rightharpoonup \mathbf{v} \text{ weakly in } L^2(0, T; H^{s+1}), \\ \mathbf{d}^\epsilon \rightharpoonup \mathbf{d} \text{ weakly* in } L^\infty(0, T; H^{s+1}), \\ \dot{\mathbf{d}}^\epsilon \rightharpoonup \mathbf{w} \text{ weakly* in } L^\infty(0, T; H^s). \end{cases}$$

Using the same arguments in the proof of Part (I) of Theorem 1.1 in Section 4.1, we show that (\mathbf{v}, \mathbf{d}) is a strong solution to the hyperbolic-type system (1.2). Moreover, Fatou lemma and the bound (4.9) yield that the solution (\mathbf{v}, \mathbf{d}) satisfies the inequality

$$|\mathbf{v}(t)|_{H^s}^2 + |\dot{\mathbf{d}}(t)|_{H^s}^2 + |\nabla \mathbf{d}(t)|_{H^s}^2 + \frac{1}{2}\mu_4 \int_0^t |\nabla \mathbf{v}|_{H^s}^2 ds \leq E^{in} + \frac{C_4(E^{in})^2 T}{(1 - C_4 E^{in} T)^2}$$

for all $t \in [0, T]$, and then we complete the proof of Part (II) of Theorem 1.1.

4.3. The proof of Part (III) of Theorem 1.1. In order to prove the existence of the unique global classical solution, we must construct a stronger energy estimate. We have shown the apriori estimate in Lemma 2.1:

$$\begin{aligned} & \frac{d}{dt} \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) + \mu_4 |\nabla \mathbf{v}|_{H^s}^2 - 2\lambda_1 |\dot{\mathbf{d}}|_{H^s}^2 + 2\mu_1 \sum_{k=0}^s |\mathbf{d}^\top (\nabla^{k+1} \mathbf{v}) \mathbf{d}|_{L^2}^2 \\ & \leq C(1 + |\lambda_1| + |\lambda_2| + \mu_2 + \mu_3) \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) |\nabla \mathbf{v}|_{H^s} \\ & \quad + C(1 + |\lambda_1| + |\lambda_2|) \left(|\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right)^2 + C(\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6) |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2. \end{aligned}$$

Since $\mu_2 < \mu_3$, $-\lambda_1 = |\lambda_1| = \mu_3 - \mu_2 > 0$. Then Cauchy inequality reduce to

$$\begin{aligned} & \frac{d}{dt} \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) + \frac{1}{2}\mu_4 |\nabla \mathbf{v}|_{H^s}^2 + 2|\lambda_1| |\dot{\mathbf{d}}|_{H^s}^2 + 2\mu_1 \sum_{k=0}^s |\mathbf{d}^\top (\nabla^{k+1} \mathbf{v}) \mathbf{d}|_{L^2}^2 \\ & \leq C \left(\frac{C(1+|\lambda_1|+|\lambda_2|+\mu_2+\mu_3)^2}{2\mu_4} + 1 + |\lambda_1| + |\lambda_2| \right) \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right)^2 \\ & \quad + C(\mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6) |\nabla \mathbf{d}|_{H^s}^2 |\nabla \mathbf{v}|_{H^s}^2 \\ & \leq C \left(\frac{C(1+|\lambda_1|+|\lambda_2|+\mu_2+\mu_3)^2}{2\mu_4} + 1 + |\lambda_1| + |\lambda_2| + \mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6 \right) \\ & \quad \times \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) \cdot \left(|\nabla \mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right). \end{aligned} \tag{4.10}$$

Next we are going to construct a more apriori estimate. Multiplying by \mathbf{d} in the third direction equation of (1.1), integrating by parts and controlling by Hölder inequality, we have

$$\begin{aligned} \langle \ddot{\mathbf{d}}, \mathbf{d} \rangle + |\nabla \mathbf{d}|_{L^2}^2 &= \langle \gamma \mathbf{d}, \mathbf{d} \rangle + \lambda_1 \langle \dot{\mathbf{d}} - B \mathbf{d}, \mathbf{d} \rangle + \lambda_2 \langle A \mathbf{d}, \mathbf{d} \rangle \\ &\leq |\mathbf{d}|_{L^\infty}^2 \left(|\dot{\mathbf{d}}|_{L^2}^2 + |\nabla \mathbf{d}|_{L^2}^2 \right) + |\lambda_1| |\mathbf{d}|_{L^\infty}^3 |\mathbf{d}|_{L^2} |\nabla \mathbf{v}|_{L^2} + |\lambda_2| |\mathbf{d}|_{L^\infty}^3 |\mathbf{d}|_{L^2} |\nabla \mathbf{v}|_{L^2} \\ &\leq |\mathbf{d}|_{L^\infty}^2 \left(|\dot{\mathbf{d}}|_{L^2}^2 + |\nabla \mathbf{d}|_{L^2}^2 \right) + (|\lambda_1| + |\lambda_2|) |\mathbf{d}|_{L^\infty} |\mathbf{d}|_{L^2} |\nabla \mathbf{v}|_{L^2}, \end{aligned}$$

where the latter inequality is implied by the fact $|\mathbf{d}| = 1$. Notice that by the fact $\operatorname{div} \mathbf{v} = 0$

$$\begin{aligned} \langle \ddot{\mathbf{d}}, \mathbf{d} \rangle &= \langle \partial_t \dot{\mathbf{d}} + \mathbf{v} \cdot \nabla \dot{\mathbf{d}}, \mathbf{d} \rangle \\ &= \frac{d}{dt} \langle \dot{\mathbf{d}}, \mathbf{d} \rangle - \langle \dot{\mathbf{d}}, \partial_t \mathbf{d} \rangle - \langle \dot{\mathbf{d}}, \mathbf{v} \cdot \nabla \mathbf{d} \rangle \\ &= \frac{d}{dt} \langle \dot{\mathbf{d}}, \mathbf{d} \rangle - |\dot{\mathbf{d}}|_{L^2}^2 \\ &= \frac{1}{2} \frac{d}{dt} \left(|\mathbf{d} + \dot{\mathbf{d}}|_{L^2}^2 - |\dot{\mathbf{d}}|_{L^2}^2 - |\mathbf{d}|_{L^2}^2 \right) - |\dot{\mathbf{d}}|_{L^2}^2. \end{aligned}$$

So we know that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(|\mathbf{d} + \dot{\mathbf{d}}|_{L^2}^2 - |\dot{\mathbf{d}}|_{L^2}^2 - |\mathbf{d}|_{L^2}^2 \right) - |\dot{\mathbf{d}}|_{L^2}^2 \\ & \leq |\mathbf{d}|_{L^\infty}^2 \left(|\dot{\mathbf{d}}|_{L^2}^2 + |\nabla \mathbf{d}|_{L^2}^2 \right) + (|\lambda_1| + |\lambda_2|) |\mathbf{d}|_{L^\infty} |\mathbf{d}|_{L^2} |\nabla \mathbf{v}|_{L^2} \\ & \leq C \left(\frac{|\lambda_1|^2 + |\lambda_2|^2}{\mu_4} + 1 \right) |\nabla \mathbf{d}|_{H^s}^2 \left(|\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) + \frac{1}{16} \mu_4 |\nabla \mathbf{v}|_{H^s}^2. \end{aligned} \quad (4.11)$$

For all $1 \leq k \leq s$ ($s > \frac{n}{2} + 2$), we take ∇^k in the third direction equation of the parabolic-hyperbolic model (1.1), multiply by $\nabla^k \mathbf{d}$, integrate by parts and then we get

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(|\nabla^k (\dot{\mathbf{d}} + \mathbf{d})|_{L^2}^2 + (|\lambda_1| - 1) |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 - |\nabla^k \mathbf{d}|_{L^2}^2 \right) - |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 + |\nabla^{k+1} \mathbf{d}|_{L^2}^2 \\ & = - \sum_{\substack{a+b=k \\ a \geq 1}} \langle \nabla^k \dot{\mathbf{d}}, \nabla^a \mathbf{v} \nabla^{b+1} \mathbf{d} \rangle - \sum_{\substack{a+b=k \\ a \geq 1}} \langle \nabla^a \mathbf{v} \nabla^{b+1} \dot{\mathbf{d}}, \nabla^k \mathbf{d} \rangle \\ & \quad + \langle \nabla^k (\gamma \mathbf{d}), \nabla^k \mathbf{d} \rangle + \lambda_1 \langle \nabla^k (\mathbf{v} \cdot \nabla \mathbf{d} - B \mathbf{d}), \nabla^k \mathbf{d} \rangle + \lambda_2 \langle \nabla^k (A \mathbf{d}), \nabla^k \mathbf{d} \rangle \\ & \equiv M_1 + M_2 + M_3 + M_4 + M_5, \end{aligned} \quad (4.12)$$

where $\gamma = -|\dot{\mathbf{d}}|^2 + |\nabla \mathbf{d}|^2 + \lambda_1 \mathbf{d}^\top B \mathbf{d} - \lambda_2 \mathbf{d}^\top A \mathbf{d}$. By using the Sobolev embedding $W^{1,2} \hookrightarrow L^4$ and $W^{2,2} \hookrightarrow L^\infty$ for $n = 2, 3$ and Hölder inequality, we have

$$\begin{aligned} M_1 + M_2 & \leq \sum_{\substack{a+b=k \\ a \geq 1}} |\nabla^k \dot{\mathbf{d}}|_{L^2} |\nabla^a \mathbf{v}|_{L^4} |\nabla^{b+1} \mathbf{d}|_{L^4} \\ & \quad + \sum_{\substack{a+b=k \\ a \geq 2}} |\nabla^k \mathbf{d}|_{L^2} |\nabla^a \mathbf{v}|_{L^4} |\nabla^{b+1} \dot{\mathbf{d}}|_{L^4} + |\nabla \mathbf{v}|_{L^\infty} |\nabla^k \dot{\mathbf{d}}|_{L^2} |\nabla^k \mathbf{d}|_{L^2} \\ & \leq C |\nabla^k \dot{\mathbf{d}}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2} + C |\nabla^k \dot{\mathbf{d}}|_{L^2} |\nabla^{k+1} \mathbf{v}|_{L^2} |\nabla^k \mathbf{d}|_{L^2} \\ & \quad + C |\nabla^3 \mathbf{v}|_{L^2} |\nabla^k \dot{\mathbf{d}}|_{L^2} |\nabla^k \mathbf{d}|_{L^2} \\ & \leq C |\dot{\mathbf{d}}|_{H^s} |\nabla \mathbf{d}|_{H^s} |\nabla \mathbf{v}|_{H^s} \\ & \leq \frac{C}{\mu_4} (|\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2)^2 + \frac{1}{16} \mu_4 |\nabla \mathbf{v}|_{H^s}^2, \end{aligned} \quad (4.13)$$

and

$$\begin{aligned} M_3 & = - \sum_{a+b+c=k} \langle \nabla^a \dot{\mathbf{d}} \nabla^b \dot{\mathbf{d}} \nabla^c \mathbf{d}, \nabla^k \mathbf{d} \rangle - \sum_{a+b+c=k} \langle \nabla^{a+1} \dot{\mathbf{d}} \nabla^{b+1} \mathbf{d} \nabla^c \mathbf{d}, \nabla^k \mathbf{d} \rangle \\ & \quad + \lambda_1 \langle \nabla^k [(\mathbf{d}^\top B \mathbf{d}) \mathbf{d}], \nabla^k \mathbf{d} \rangle - \lambda_2 \langle \nabla^k [(\mathbf{d}^\top A \mathbf{d}) \mathbf{d}], \nabla^k \mathbf{d} \rangle \\ & \leq 2 |\dot{\mathbf{d}}|_{L^4} |\nabla^k \dot{\mathbf{d}}|_{L^2} |\nabla^k \mathbf{d}|_{L^4} |\mathbf{d}|_{L^\infty} + \sum_{\substack{a+b+c=k \\ a,b \leq k-1}} |\nabla^a \dot{\mathbf{d}}|_{L^4} |\nabla^b \dot{\mathbf{d}}|_{L^4} |\nabla^c \mathbf{d}|_{L^4} |\nabla^k \mathbf{d}|_{L^4} \\ & \quad + 2 |\nabla \mathbf{d}|_{L^4} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^k \mathbf{d}|_{L^4} |\mathbf{d}|_{L^\infty} + \sum_{\substack{a+b+c=k \\ a,b \leq k-1}} |\nabla^{a+1} \mathbf{d}|_{L^4} |\nabla^{b+1} \mathbf{d}|_{L^4} |\nabla^c \mathbf{d}|_{L^4} |\nabla^k \mathbf{d}|_{L^4} \\ & \quad + C(|\lambda_1| + |\lambda_2|) |\nabla^k \mathbf{d}|_{L^2} \left(|\mathbf{d}|_{L^\infty}^3 |\nabla^{k+1} \mathbf{v}|_{L^2} + |\nabla \mathbf{v}|_{L^\infty} |\mathbf{d}|_{L^\infty}^2 |\nabla^k \mathbf{d}|_{L^2} \right) \\ & \leq C |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 |\nabla^{k+1} \mathbf{d}|_{L^2}^2 + C |\nabla^{k+1} \mathbf{d}|_{L^2}^4 + C(|\lambda_1| + |\lambda_2|) |\nabla^{k+1} \mathbf{d}|_{L^2}^3 |\nabla^{k+1} \mathbf{v}|_{L^2} \\ & \leq C(1 + |\lambda_1| + |\lambda_2|) |\nabla^{k+1} \mathbf{d}|_{L^2}^2 \left(|\nabla^{k+1} \mathbf{v}|_{L^2}^2 + |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 + |\nabla^{k+1} \mathbf{d}|_{L^2}^2 \right) \\ & \leq C(1 + |\lambda_1| + |\lambda_2|) |\nabla \mathbf{d}|_{H^s}^2 \left(|\nabla \mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right), \end{aligned} \quad (4.14)$$

and

$$\begin{aligned}
M_4 + M_5 &\leq |\lambda_1| \sum_{a+b=k} \left\langle |\nabla^a \mathbf{v}| |\nabla^{b+1} \mathbf{d}|, |\nabla^k \mathbf{d}| \right\rangle + (|\lambda_1| + |\lambda_2|) \sum_{a+b=k} \left\langle |\nabla^{a+1} \mathbf{v}| |\nabla^b \mathbf{d}|, |\nabla^k \mathbf{d}| \right\rangle \\
&\leq C |\lambda_1| |\nabla^k \mathbf{v}|_{L^4} |\nabla^{k+1} \mathbf{d}|_{L^2} |\nabla^k \mathbf{d}|_{L^4} + C(|\lambda_1| + |\lambda_2|) |\mathbf{d}|_{L^\infty} |\nabla^{k+1} \mathbf{v}|_{L^2} + |\nabla^k \mathbf{d}|_{L^2} \\
&\quad + C(|\lambda_1| + |\lambda_2|) \sum_{\substack{a+b=k \\ a \leq k-1}} |\nabla^{a+1} \mathbf{v}|_{L^4} |\nabla^b \mathbf{d}|_{L^4} |\nabla^k \mathbf{d}|_{L^2} \\
&\leq C |\lambda_1| |\nabla^{k+1} \mathbf{v}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2}^2 + C(|\lambda_1| + |\lambda_2|) |\nabla^{k+1} \mathbf{v}|_{L^2} |\nabla^{k+1} \mathbf{d}|_{L^2}^2 \\
&\leq C \frac{|\lambda_1|^2 + |\lambda_2|^2}{\mu_4} |\nabla^{k+1} \mathbf{d}|_{L^2}^4 + \frac{1}{16} \mu_4 |\nabla^{k+1} \mathbf{v}|_{L^2}^2 \\
&\leq C \frac{|\lambda_1|^2 + |\lambda_2|^2}{\mu_4} |\nabla \mathbf{d}|_{H^s}^4 + \frac{1}{16} \mu_4 |\nabla \mathbf{v}|_{H^s}^2.
\end{aligned} \tag{4.15}$$

It is immediately implied that by substituting (4.13), (4.14) and (4.15) into (4.12)

$$\begin{aligned}
&\frac{1}{2} \frac{d}{dt} \left(|\nabla^k (\dot{\mathbf{d}} + \mathbf{d})|_{L^2}^2 + (|\lambda_1| - 1) |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 - |\nabla^k \mathbf{d}|_{L^2}^2 \right) - |\nabla^k \dot{\mathbf{d}}|_{L^2}^2 + |\nabla^{k+1} \mathbf{d}|_{L^2}^2 \\
&\leq C \left(\frac{1+|\lambda_1|^2+|\lambda_2|^2}{\mu_4} + 1 + |\lambda_1| + |\lambda_2| \right) \left(|\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) \left(|\nabla \mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) \\
&\quad + \frac{1}{8} \mu_4 |\nabla \mathbf{v}|_{H^s}^2
\end{aligned}$$

for all $1 \leq k \leq s$. Then, by combining the inequality (4.11), we sum up the above inequalities for $1 \leq k \leq s$ and have that

$$\begin{aligned}
&\frac{d}{dt} \left(|\dot{\mathbf{d}} + \mathbf{d}|_{H^s}^2 + (|\lambda_1| - 1) |\dot{\mathbf{d}}|_{H^s}^2 - |\mathbf{d}|_{H^s}^2 \right) - 2 |\dot{\mathbf{d}}|_{H^s}^2 + 2 |\nabla \mathbf{d}|_{H^s}^2 - \frac{3}{8} \mu_4 |\nabla \mathbf{v}|_{H^s}^2 \\
&\leq C \left(\frac{1+|\lambda_1|^2+|\lambda_2|^2}{\mu_4} + 1 + |\lambda_1| + |\lambda_2| \right) \left(|\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) \left(|\nabla \mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right).
\end{aligned} \tag{4.16}$$

Taking a positive constant $\delta = \frac{1}{2} \min\{|\lambda_1|, 1\} \in (0, \frac{1}{2}]$ and noticing that $\frac{d}{dt} |\mathbf{d}|_{L^2}^2 = 0$, we multiply by δ in the inequality (4.16) and then add it to the inequality (4.10), so that we know that

$$\begin{aligned}
&\frac{d}{dt} \left(|\mathbf{v}|_{H^s}^2 + \delta |\dot{\mathbf{d}} + \mathbf{d}|_{H^s}^2 + (\delta |\lambda_1| + 1 - \delta) |\dot{\mathbf{d}}|_{H^s}^2 + (1 - \delta) |\nabla \mathbf{d}|_{H^s}^2 + \delta |\nabla^{s+1} \mathbf{d}|_{L^2}^2 \right) \\
&\quad + 2(|\lambda_1| - \delta) |\dot{\mathbf{d}}|_{H^s}^2 + (\frac{1}{2} - \frac{3}{8} \delta) \mu_4 |\nabla \mathbf{v}|_{H^s}^2 + 2\delta |\nabla \mathbf{d}|_{H^s}^2 + 2\mu_1 \sum_{k=1}^s |\mathbf{d}^\top (\nabla^{k+1} \mathbf{v}) \mathbf{d}|_{L^2}^2 \\
&\leq C \delta \left(\frac{1+|\lambda_1|^2+|\lambda_2|^2}{\mu_4} + 1 + |\lambda_1| + |\lambda_2| \right) \left(|\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) \left(|\nabla \mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) \\
&\quad + C \left(\frac{C(1+|\lambda_1|+|\lambda_2|+\mu_2+\mu_3)^2}{2\mu_4} + 1 + |\lambda_1| + |\lambda_2| + \mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6 \right) \\
&\quad \times \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) \cdot \left(|\nabla \mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right).
\end{aligned} \tag{4.17}$$

Denoted by

$$\mathcal{E}(t) \equiv |\mathbf{v}|_{H^s}^2 + \delta |\dot{\mathbf{d}} + \mathbf{d}|_{H^s}^2 + (\delta |\lambda_1| + 1 - \delta) |\dot{\mathbf{d}}|_{H^s}^2 + (1 - \delta) |\nabla \mathbf{d}|_{H^s}^2 + \delta |\nabla^{s+1} \mathbf{d}|_{L^2}^2$$

and

$$\mathcal{D}(t) \equiv 2(|\lambda_1| - \delta) |\dot{\mathbf{d}}|_{H^s}^2 + (\frac{1}{2} - \frac{3}{8} \delta) \mu_4 |\nabla \mathbf{v}|_{H^s}^2 + 2\delta |\nabla \mathbf{d}|_{H^s}^2 + 2\mu_1 \sum_{k=1}^s |\mathbf{d}^\top (\nabla^{k+1} \mathbf{v}) \mathbf{d}|_{L^2}^2,$$

the inequality (4.17) reduces to

$$\frac{d}{dt} \mathcal{E}(t) + \mathcal{D}(t) \leq C_5 \mathcal{E}(t) \mathcal{D}(t), \tag{4.18}$$

where the positive constant C_5 is

$$\begin{aligned} C_5 = & C \left(\frac{8}{4-3\delta} + \frac{1}{2(|\lambda_1|-\delta)} + \frac{1}{2\delta} \right) \left(\frac{1}{\delta|\lambda_1|+1-\delta} + \frac{1}{1-\delta} + 1 \right) \left[\delta \left(\frac{1+|\lambda_1|^2+|\lambda_2|^2}{\mu_4} + 1 + |\lambda_1| + |\lambda_2| \right) \right. \\ & \left. + \frac{C(1+|\lambda_1|+|\lambda_2|+\mu_2+\mu_3)^2}{2\mu_4} + 1 + |\lambda_1| + |\lambda_2| + \mu_1 + \mu_2 + \mu_3 + \mu_5 + \mu_6 \right] > 0. \end{aligned}$$

Recalling that $\mathcal{E}^{in} \leq \frac{\delta(\delta|\lambda_1|+1-\delta)}{4C_5[2\delta(\delta|\lambda_1|+1-\delta)+\delta|\lambda_1|+1]}$, we have

$$\begin{aligned} \mathcal{E}(0) = & |\mathbf{v}^{in}|_{H^s}^2 + \delta|\dot{\mathbf{d}}^{in}|_{H^s}^2 + |\mathbf{d}^{in}|_{H^s}^2 + (\delta|\lambda_1| + 1 - \delta)|\dot{\mathbf{d}}^{in}|_{H^s}^2 + (1 - \delta)|\nabla \mathbf{d}^{in}|_{H^s}^2 + \delta|\nabla^{s+1} \mathbf{d}|_{L^2}^2 \\ \leq & \left(2 + \frac{1}{\delta} + \frac{1}{\delta|\lambda_1|+1-\delta} \right) \mathcal{E}^{in} \\ \leq & \frac{1}{4C_5}. \end{aligned}$$

We define $T^* = \sup \left\{ t > 0; \mathcal{E}(t) < \frac{1}{2C_5} \right\}$. Then for all $t \in [0, T^*]$

$$\frac{d}{dt} \mathcal{E}(t) + \frac{1}{2} \mathcal{D}(t) \leq 0,$$

which implies that for all $t \in [0, T^*]$

$$\mathcal{E}(t) \leq \mathcal{E}(0) \leq \frac{1}{4C_5} < \frac{1}{2C_5}.$$

This yields that $T^* = +\infty$ and for any fixed $T \in (0, +\infty)$

$$\sup_{0 \leq t \leq T} \mathcal{E}(t) + \frac{1}{2} \int_0^T \mathcal{D}(t) dt \leq \mathcal{E}(0) \leq \frac{1}{4C_5}.$$

Consequently, under the assumption $\mathcal{E}^{in} \leq \frac{\delta(\delta|\lambda_1|+1-\delta)}{4C_5[2\delta(\delta|\lambda_1|+1-\delta)+\delta|\lambda_1|+1]}$ the following global apriori bound is valid for any fixed $T \in (0, +\infty)$:

$$\sup_{0 \leq t \leq T} \left(|\mathbf{v}|_{H^s}^2 + |\dot{\mathbf{d}}|_{H^s}^2 + |\nabla \mathbf{d}|_{H^s}^2 \right) (t) + \int_0^T |\nabla \mathbf{v}|_{H^s}^2 dt \leq \frac{1}{4C_5 \min \left\{ \delta, \frac{4-3\delta}{16} \mu_4 \right\}}. \quad (4.19)$$

Finally, by using the global apriori bound (4.19) we follow the arguments in Section 4.1 and complete the proof of the existence of the global classical solution.

ACKNOWLEDGEMENT

We take this opportunity to appreciate Prof. Fanghua Lin, who suggested the Ericksen-Leslie's parabolic-hyperbolic liquid crystal model to us when we visited the NYU-ECNU Institute of Mathematical Sciences at NYU Shanghai during the spring semester of 2015. We also thanks the hospitality of host institute.

REFERENCES

- [1] A. Bressan, P. Zhang and Y-X. Zheng, Asymptotic variational wave equations. *Arch. Ration. Mech. Anal.* **183** (2007), no. 1, 163-185.
- [2] G. Chen, P. Zhang and Y-X. Zheng, Energy conservative solutions to a nonlinear wave system of nematic liquid crystals. *Commun. Pure Appl. Anal.* **12** (2013), no. 3, 1445-1468.
- [3] F. De Anna and A. Zarnescu, Global well-posedness and twist-wave solutions for the inertial Qian-Sheng model of liquid crystals. arXiv:1608.08872
- [4] J. L. Ericksen, Hydrostatic theory of liquid crystals. *Arch. Rational Mech. Anal.* **9** 1962, 371-378.
- [5] J. L. Ericksen, Conservation laws for liquid crystals. *Trans. Soc. Rheology* **5** 1961, 23-34.
- [6] J. L. Ericksen, On equations of motion for liquid crystals. *Q. J. Mechanics Appl. Math.* (1976) **29** (2), 203-208.
- [7] E. Feireisl, E. Rocca, G. Schimperna and A. Zarnescu, On a hyperbolic system arising in liquid crystals modeling. arXiv:1610.07828
- [8] R. Hardt, D. Kinderlehrer and F-H. Lin, Existence and partial regularity of static liquid crystal configurations. *Comm. Math. Phys.* **105** (1986), no. 4, 547-570.

- [9] M-C. Hong and Z-P. Xin, Global existence of solutions of the liquid crystal flow for the Oseen-Frank model in \mathbb{R}^2 . *Adv. Math.* **231** (2012), no. 3-4, 1364-1400.
- [10] J-R. Huang, F-H. Lin and C-Y. Wang, Regularity and existence of global solutions to the Ericksen-Leslie system in \mathbb{R}^2 . *Comm. Math. Phys.* **331** (2014), no. 2, 805-850.
- [11] T. Huang, F-H. Lin, C. Liu and C-Y. Wang, Finite time singularity of the nematic liquid crystal flow in dimension three. *Arch. Ration. Mech. Anal.* **221** (2016), no. 3, 1223-1254.
- [12] F. M. Leslie, Some thermal effects in cholesteric liquid crystals. *Proc. Roy. Soc. A.*, **307** (1968), 359-372.
- [13] F. M. Leslie, Some constitutive equations for liquid crystals. *Arch. Rational Mech. Anal.* **28** (1968), no. 4, 265-283.
- [14] J-K. Li, E. Titi and Z-P. Xin, On the uniqueness of weak solutions to the Ericksen-Leslie liquid crystal model in \mathbb{R}^2 . *Math. Models Methods Appl. Sci.* **26** (2016), no. 4, 803-822.
- [15] F-H. Lin, J-Y. Lin and C-Y. Wang, Liquid crystal flows in two dimensions. *Arch. Ration. Mech. Anal.* **197** (2010), no. 1, 297-336.
- [16] F-H. Lin and C. Liu, Nonparabolic dissipative systems modeling the flow of liquid crystals. *Comm. Pure Appl. Math.* **48** (1995), no. 5, 501-537.
- [17] F-H. Lin and C. Liu, Partial regularity of the dynamic system modeling the flow of liquid crystals. *Discrete Contin. Dynam. Systems* **2** (1996), no. 1, 1-22.
- [18] F-H. Lin and C. Liu, Existence of solutions for the Ericksen-Leslie system. *Arch. Ration. Mech. Anal.* **154** (2000), no. 2, 135-156.
- [19] F-H. Lin and C. Liu, Static and dynamic theories of liquid crystal. *J. Partial Differential Equations*, **14** (2001), no. 4, 289-330.
- [20] F-H. Lin and C-Y. Wang, On the uniqueness of heat flow of harmonic maps and hydrodynamic flow of nematic liquid crystals. *Chin. Ann. Math. Ser. B* **31** (2010), no. 6, 921-938.
- [21] F-H. Lin and C-Y. Wang, Global existence of weak solutions of the nematic liquid crystal flow in dimension three. *Comm. Pure Appl. Math.* **69** (2016), no. 8, 1532-1571.
- [22] P-L. Lions. *Mathematical Topics in Fluid Mechanics, Vol. 1 : Incompressible Models*. Clarendon Press. Oxford, 1996.
- [23] A. J. Majda and A. L. Bertozzi, *Vorticity and incompressible flow*, Cambridge University Press, 2002.
- [24] M. Wang and W-D. Wang, Global existence of weak solution for the 2-D Ericksen-Leslie system. *Calc. Var. Partial Differential Equations* **51** (2014), no. 3-4, 915-962.
- [25] M. Wang, W-D. Wang and Z-F. Zhang, On the uniqueness of weak solution for the 2-D Ericksen-Leslie system. *Discrete Contin. Dyn. Syst. Ser. B* **21** (2016), no. 3, 919-941.
- [26] W. Wang, P-W. Zhang and Z-F. Zhang, Well-posedness of the Ericksen-Leslie system. *Arch. Ration. Mech. Anal.* **210** (2013), no. 3, 837-855.
- [27] X. Xu and Z-F. Zhang, Global regularity and uniqueness of weak solution for the 2-D liquid crystal flows. *J. Differential Equations* **252** (2012), no. 2, 1169-1181.
- [28] P. Zhang and Y-X. Zheng, On oscillations of an asymptotic equation of a nonlinear variational wave equation. *Asymptot. Anal.* **18** (1998), no. 3-4, 307-327.
- [29] P. Zhang and Y-X. Zheng, On the existence and uniqueness of solutions to an asymptotic equation of a variational wave equation. *Acta Math. Sin. (Engl. Ser.)* **15** (1999), no. 1, 115-130.
- [30] P. Zhang and Y-X. Zheng, Existence and uniqueness of solutions of an asymptotic equation arising from a variational wave equation with general data. *Arch. Ration. Mech. Anal.* **155** (2000), no. 1, 49-83.
- [31] P. Zhang and Y-X. Zheng, Singular and rarefactive solutions to a nonlinear variational wave equation. *Chinese Ann. Math. Ser. B* **22** (2001), no. 2, 159-170.
- [32] P. Zhang and Y-X. Zheng, Rarefactive solutions to a nonlinear variational wave equation of liquid crystals. *Comm. Partial Differential Equations* **26** (2001), no. 3-4, 381-419.
- [33] P. Zhang and Y-X. Zheng, On the second-order asymptotic equation of a variational wave equation. *Proc. Roy. Soc. Edinburgh Sect. A* **132** (2002), no. 2, 483-509.
- [34] P. Zhang and Y-X. Zheng, Weak solutions to a nonlinear variational wave equation. *Arch. Ration. Mech. Anal.* **166** (2003), no. 4, 303-319.
- [35] P. Zhang and Y-X. Zheng, Weak solutions to a nonlinear variational wave equation with general data. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **22** (2005), no. 2, 207-226.
- [36] P. Zhang and Y-X. Zheng, Conservative solutions to a system of variational wave equations of nematic liquid crystals. *Arch. Ration. Mech. Anal.* **195** (2010), no. 3, 701-727.
- [37] P. Zhang and Y-X. Zheng, Energy conservative solutions to a one-dimensional full variational wave system. *Comm. Pure Appl. Math.* **65** (2012), no. 5, 683-726.

(Ning Jiang)

SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN, 430072, P. R. CHINA

E-mail address: njiang@whu.edu.cn

(Yi-Long Luo)

INSTITUTE OF MATHEMATICS, ACADEMY OF MATHEMATICS AND SYSTEM SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING, 100190, P. R. CHINA

E-mail address: yl-luo@amss.ac.cn