

# Maximum Likelihood Estimation in Markov Regime-Switching Models with Covariate-Dependent Transition Probabilities\*

Demian Pouzo,<sup>†</sup>Zacharias Psaradakis<sup>‡</sup>and Martin Sola<sup>§</sup>

December 7, 2021

## Abstract

This paper considers maximum likelihood (ML) estimation in a large class of models with hidden Markov regimes. We investigate consistency of the ML estimator and local asymptotic normality for the models under general conditions which allow for autoregressive dynamics in the observable process, Markov regime sequences with covariate-dependent transition matrices, and possible model misspecification. A Monte Carlo study examines the finite-sample properties of the ML estimator in correctly specified and misspecified models. An empirical application is also discussed.

*Key words and phrases:* Autoregressive model; consistency; covariate-dependent transition probabilities; hidden Markov model; Markov-switching model; maximum likelihood; local asymptotic normality; misspecified models.

---

\*We would like to thank Nouredine El Karoui, Adityanand Guntuboyina, Michael Jansson, Ulrich Müller, and three anonymous referees for useful comments and suggestions. All errors are our own.

<sup>†</sup>e-mail: dpouzo@berkeley.edu (corresponding author). Address: UC Berkeley, Dept. of Economics; 530 Evans Hall, Berkeley CA 94619, USA.

<sup>‡</sup>e-mail: z.psaradakis@bbk.ac.uk. Address: Birkbeck, University of London, Dept. of Economics, Mathematics & Statistics; Malet Street, London WC1E 7HX, UK

<sup>§</sup>e-mail: msola@utdt.edu. Address: Universidad Torcuato Di Tella, Dept. of Economics; Figueroa Alcorta 7350, C1428 Buenos Aires, Argentina.

# 1 Introduction

Stochastic models with parameters that are subject to changes driven by an unobservable Markov chain (the regime or state sequence) have attracted considerable attention in many different areas, the influential work by [Hamilton \[1989\]](#) being a prominent example from the econometrics literature. An important subclass of such models, the so-called hidden Markov models, in which observations are conditionally independent given the regime sequence, are also widely used in a variety of disciplines. A common assumption in these models is that the unobservable Markov chain is temporally homogeneous.

In this paper, we focus on a larger class of models in which the hidden regime process and the observation process (conditional on the regimes) are both temporally inhomogeneous Markov chains. This is a useful generalization of models with a time-invariant transition mechanism that has found numerous applications, especially in economics and finance.<sup>1</sup> Statistical inference in this class of models is predominantly likelihood based, even though very little is known about the asymptotic properties of the relevant inferential procedures. In a typical application, inference is conducted on the implicit assumption that the maximum likelihood (ML) estimator of unknown parameters has its familiar properties of consistency, asymptotic normality, and asymptotic efficiency, and associated confidence sets and hypotheses tests are constructed in the usual manner. It hardly needs noting that, unless these properties of likelihood-based inferential procedures known from regular parametric estimation problems hold for Markov regime-switching models with covariate-dependent transition probabilities, inferences drawn from them cannot be justified in any meaningful way and should be interpreted very cautiously. Arguing, for instance, as is common in applied work, that an economic variable is a useful leading indicator for business-cycle phases because it appears to have a ‘statistically significant’ coefficient in the transition functions of a Markov-switching model for output growth is problematic when little is known about the properties of the relevant estimators and related tests.

The main contribution of this paper is to provide consistency and asymptotic normality results for a large class of models that are relevant in applications. Our approach allows for autoregressive dynamics in the observable process, covariate-dependence in the transition functions of the hidden regime

---

<sup>1</sup>Examples include, among many others, applications to the analysis of business-cycle fluctuations (e.g., [Filardo \[1994\]](#), [Filardo and Gordon \[1998\]](#), [Ravn and Sola \[1999\]](#), [Simpson et al. \[2001\]](#), [Gadea Rivas and Perez-Quiros \[2015\]](#)), interest rates and yields (e.g., [Gray \[1996\]](#), [Bekaert et al. \[2001\]](#), [Ang and Bekaert \[2002a\]](#), [Ang and Bekaert \[2002b\]](#), [Psaradakis and Sola \[2021\]](#)), consumption growth ([Whitelaw \[2000\]](#)), currency crises (e.g., [Martinez Peria \[2002\]](#), [Mouratidis \[2008\]](#)), and cryptocurrency returns ([Tan et al. \[2021\]](#)).

process, and potential model misspecification. To the best of our knowledge, the only asymptotic results available on ML estimation in Markov regime-switching models with covariate-dependent transition probabilities are those of [Ailliot and Pène \[2015\]](#), who investigate consistency of the ML estimator in a correctly specified model (i.e., a model which contains the data-generating process). Our results include both consistency of the ML estimator and local asymptotic normality (LAN) for the model, from which asymptotic normality of the ML estimator can be inferred. Unlike [Ailliot and Pène \[2015\]](#), who allow for a general hidden state space, we require the latter to be finite, but do not restrict the model to be correctly specified. In doing so, we also extend some results of [White \[1982\]](#) for independent, identically distributed (i.i.d.) data to the case of dependent observations and for classes of parametric distributions associated with dynamic models with hidden Markov regimes. As such stochastic specifications are typically highly parametric, it is important to understand the properties of likelihood-based inferential procedures in situations where the true probability structure of the data does not necessarily lie within the parametric family of distributions specified by the model. We show that the ML estimator in our setting converges to the true parameter value if the model is correctly specified and to a pseudo-true parameter set if the model is misspecified.<sup>2</sup> We also show that the sample log-likelihood satisfies the LAN property, establish an asymptotic linear representation for the ML estimator, obtain the asymptotic distribution of the estimator, and present results relating to consistent estimation of its asymptotic covariance matrix. These are the most general results available for Markov regime-switching models with autoregressive dynamics and covariate-dependent transition probabilities.

In related earlier work, [Mevel and Finesso \[2004\]](#) examine consistency and asymptotic normality of the ML estimator in misspecified hidden Markov models with a finite state space, while [Douc and Moulines \[2012\]](#) consider consistency under general state spaces. [Bickel and Ritov \[1996\]](#), [Bickel et al. \[1998\]](#), [Jensen and Petersen \[1999\]](#), [Douc and Matias \[2001\]](#), and [Douc et al. \[2011\]](#) investigate consistency and/or asymptotic normality in correctly specified hidden Markov models with regime sequences defined on either a finite or a general state space. [Francq and Roussignol \[1998\]](#) and [Krishnamurthy and Rydén \[1998\]](#) consider consistency in correctly specified autoregressive models with Markov regimes defined on a finite state space. [Douc et al. \[2004b\]](#) and [Kasahara and Shimotsu \[2019\]](#) investigate consistency and asymptotic normality in a similar autoregressive setup, but allow the regime sequence to take values in a space that is not necessarily countable. In all of these papers, the hidden regime sequence is

---

<sup>2</sup>We refer to estimators for the various models discussed throughout as ML estimators even though they may be obtained from a pseudo-likelihood based on a misspecified model.

assumed to be a temporally homogeneous Markov chain.

In the sequel, we follow [Bickel et al. \[1998\]](#) and [Douc et al. \[2004b\]](#) fairly closely in terms of the technical tools and the arguments used to establish our results, but our setup is more general in certain respects. Like [Bickel et al. \[1998\]](#), we consider models with a finite hidden state space, but allow for autoregressive dynamics in the observation sequence, covariate-dependence in the transition probabilities of the regime sequence, and potential model misspecification. In [Douc et al. \[2004b\]](#), the hidden Markov chain is allowed to take values in a compact topological space, but is restricted to be temporally homogeneous and the model is assumed to be correctly specified. The cornerstone of the methods used in these papers for establishing the asymptotic properties of the ML estimator are mixing-type results for the unobservable regime sequence conditional on the observation sequence (see also [Bickel and Ritov \[1996\]](#)). This is also true for our approach, although the aforementioned results cannot be invoked directly because they are established under the assumption of temporal homogeneity of the hidden Markov chain. We, therefore, extend these results to allow for more general Markov regime sequences; in particular, we establish mixing-type results for the unobservable regime sequence given the observed data, allowing for a particular form of covariate-dependence in the transition kernels. This last result is, to our knowledge, novel and may be of interest in its own right.

The remainder of the paper is organized as follows. [Section 2](#) defines the class of models under consideration, gives sufficient conditions for stationarity and ergodicity of the observation process, and describes the estimation problem. [Section 3](#) investigates consistency of the ML estimator in a general setting. [Section 4](#) contains results on the LAN property of the model and the asymptotic normality of the ML estimator. [Section 5](#) presents simulation results on the finite-sample properties of estimators based on well-specified and misspecified likelihoods. [Section 6](#) presents an illustration using real-world data. Proofs of the main results are gathered in an Appendix.

The following notational conventions are used throughout the paper. For an infinite sequence  $(V_j)_j$ ,  $V_a^b = (V_a, \dots, V_b)$  for any  $a \leq b$ ;  $\mathcal{P}(\mathbb{V})$  denotes the set of Borel probability measures on a Polish space  $\mathbb{V}$ ; for a probability measure  $P$ ,  $E_P(\cdot)$  denotes expectation with respect to  $P$ ,  $o_P(\cdot)$  and  $O_P(\cdot)$  indicate order in probability under  $P$ ,  $\Rightarrow_P$  signifies weak convergence under  $P$ , and  $L^r(P)$ ,  $1 \leq r < \infty$ , denotes the class of measurable functions integrable to order  $r$  with respect to  $P$ ;  $\nabla_{\vartheta}$  and  $\nabla_{\vartheta}^2$  are the gradient and Hessian operators, respectively, with respect to  $\vartheta$ ;  $\|\cdot\|$  denotes the Euclidean norm of a vector or matrix;  $1\{\cdot\}$  denotes the indicator function;  $\mathbb{N}$  denotes the set of positive integers. Unless stated otherwise, limits are taken as the sample size,  $T$ , diverges to infinity.

## 2 Model and Estimation

### 2.1 Statistical Model

Let  $(X_t, S_t)_{t=0}^\infty$  be a discrete-time stochastic process such that, for each  $t \in \mathbb{N}$ ,  $S_t \in \mathbb{S} \equiv \{s_1, \dots, s_{|\mathbb{S}|}\} \subset \mathbb{R}$  is the unobservable state and  $X_t \in \mathbb{X} \subseteq \mathbb{R}^h$ , for some  $h \in \mathbb{N}$ , is the observable state. Moreover, for each  $t \in \mathbb{N}$ , the conditional distribution of  $X_t$ , given  $X_0^{t-1}$  and  $S_0^t$ , depends only on  $X_{t-1}$  and  $S_t$ , and the conditional distribution of  $S_t$ , given  $X_0^{t-1}$  and  $S_0^{t-1}$ , depends only on  $X_{t-1}$  and  $S_{t-1}$ , so that

$$\begin{aligned} X_t \mid (X_0^{t-1}, S_0^t) &\sim P_*(X_{t-1}, S_t, \cdot), \\ S_t \mid (X_0^{t-1}, S_0^{t-1}) &\sim Q_*(X_{t-1}, S_{t-1}, \cdot), \end{aligned}$$

with  $(x, s) \mapsto P_*(x, s, \cdot) \in \mathcal{P}(\mathbb{X})$  and  $(x, s) \mapsto Q_*(x, s, \cdot) \in \mathcal{P}(\mathbb{S})$  denoting the true transition probabilities. It is further assumed that, for each  $(x, s) \in \mathbb{X} \times \mathbb{S}$ ,  $P_*(x, s, \cdot)$  admits a density  $p_*(x, s, \cdot)$  with respect to some  $\sigma$ -finite measure on  $\mathbb{X}$ . Our framework imposes no additional restrictions on this measure; for instance, it can be the Lebesgue measure (i.e., allow for continuous  $X_t$ ) or a counting measure (i.e., allow for discrete  $X_t$ ).

The researcher's model is given by a family of transition probabilities  $(x, s) \mapsto P_\theta(x, s, \cdot) \in \mathcal{P}(\mathbb{X})$  and  $(x, s) \mapsto Q_\theta(x, s, \cdot) \in \mathcal{P}(\mathbb{S})$  indexed by an (unknown) parameter  $\theta \in \Theta \subseteq \mathbb{R}^q$ , for some  $q \in \mathbb{N}$ , such that, for each  $\theta \in \Theta$ ,

$$\begin{aligned} X_t \mid (X_0^{t-1}, S_0^t) &\sim P_\theta(X_{t-1}, S_t, \cdot), \\ S_t \mid (X_0^{t-1}, S_0^{t-1}) &\sim Q_\theta(X_{t-1}, S_{t-1}, \cdot), \end{aligned}$$

and, for each  $(x, s) \in \mathbb{X} \times \mathbb{S}$ ,  $P_\theta(x, s, \cdot)$  admits a density  $p_\theta(x, s, \cdot)$  with respect to the same measure used to define  $p_*(x, s, \cdot)$ .

A few remarks about this setup are worth making. First, and perhaps most importantly, the unobservable states (regimes) are a Markov chain whose transition kernel can depend on the lagged value of the observable state. This is the main departure from prior literature which, with the exception of [Ailliot and Pène \[2015\]](#), has focused on the case where the conditional distribution of  $S_t$ , given  $X_0^{t-1}$  and  $S_0^{t-1}$ , depends only on  $S_{t-1}$ . Second, the model  $\{(P_\theta, Q_\theta) : \theta \in \Theta\}$  is allowed to be misspecified in the sense that  $(P_*, Q_*) \notin \{(P_\theta, Q_\theta) : \theta \in \Theta\}$ . This setup encompasses a rich family of models that arise in econometric and statistical applications, some examples of which are given below. Note that, although the family  $\{Q_\theta : \theta \in \Theta\}$  may or may not contain  $Q_*$ , it is defined on the same finite state space  $\mathbb{S}$  as  $Q_*$ , so misspecification of the number of unobservable regimes is ruled out.<sup>3</sup> Finally, even though the conditional distribution

<sup>3</sup>Such misspecification would introduce additional complexities (e.g., locally non-quadratic likelihood surfaces, non-identifiable parameters) which are beyond the scope of this paper.

of  $S_t$ , given  $X_0^{t-1}$  and  $S_0^{t-1}$ , is assumed to depend only on  $X_{t-1}$  and  $S_{t-1}$ , it is straightforward to extend all our results to situations where this distribution is dependent on additional higher-order lags of  $X_t$  and/or  $S_t$ .

**Example 1** (Hidden Markov Model with Covariate-Dependent Transition Probabilities). Let  $x = (y, z) \in \mathbb{X} = \mathbb{R}^2$  and  $\mathbb{S} = \{0, 1\}$ . Let  $P_*$  be determined by the equations

$$\begin{aligned} Y_t &= \mu^*(S_t) + \sigma^*(S_t)U_{1,t}, \\ Z_t &= \mu_2^* + \psi^*Z_{t-1} + \sigma_2^*U_{2,t}, \end{aligned}$$

where  $(U_{1,t}, U_{2,t})_t$  are i.i.d. (independent of  $(S_t)_t$ ) with zero mean and covariance matrix indexed by a parameter  $\rho^*$ , e.g.,  $\begin{bmatrix} 1 & \rho^* \\ \rho^* & 1 \end{bmatrix}$ . The transition probabilities of  $(S_t)_t$  are allowed to depend on  $Z_{t-1}$ ; for instance,  $(s, z) \mapsto Q_*(z, s, s) \equiv \Pr(S_t = s \mid Z_{t-1} = z, S_{t-1} = s) = [1 + \exp(-\alpha_s^* - \beta_s^*z)]^{-1}$  for  $s \in \mathbb{S}$ . The homogeneous specification with  $\beta_0^* = \beta_1^* = 0$  has been used to model regime shifts in a variety of economic and financial time series, including output growth (Albert and Chib [1993], Gadea Rivas and Perez-Quiros [2015]), foreign exchange rates (Engel and Hamilton [1990], Bollen et al. [2008]), and equity returns (Rydén et al. [1998], Ang and Bekaert [2002a]). The homogeneity restriction is relaxed in Diebold et al. [1994], Engel and Hakio [1996], and Ang and Bekaert [2002a], among others, to allow the transition probabilities to depend on  $Z_{t-1}$ . The results obtained here establish the asymptotic properties of the ML estimator in this class of models.  $\triangle$

**Example 2** (Markov-Switching Autoregressive Model with Covariate-Dependent Transition Probabilities). A useful generalization of the previous example is one where the outcome equation is extended to

$$Y_t = \mu^*(S_t) + \phi^*Y_{t-1} + \sigma^*(S_t)U_{1,t}.$$

Variations of the model with  $\beta_0^* = \beta_1^* = 0$  have found widespread application in economics (e.g., Hansen [1992], McCulloch and Tsay [1994], Ruge-Murcia [1995], Ang et al. [2008]) and beyond. Generalizations of the model without the restriction of time-invariant transition probabilities are also very popular and have been used, for example, in the modeling of output growth (Gadea Rivas and Perez-Quiros [2015]), interest rates (Ang and Bekaert [2002b]), consumption growth (Whitelaw [2000]), and bond spreads (Psaradakis and Sola [2021]). The results obtained here establish the asymptotic properties of the ML estimator in this class of models.  $\triangle$

**Example 3** (Mixture Autoregressive Model). Let  $x \in \mathbb{X} = \mathbb{R}$ ,  $\mathbb{S} = \{0, 1\}$ , and for each  $t \in \mathbb{N}$ ,  $\Pr(S_t = 0 \mid X_{t-1}) = G_*(X_{t-1})$  for some  $x \mapsto G_*(x) \in [0, 1]$  and

$X_t \sim P_*(X_{t-1}, S_t, \cdot)$ . This specification implies a conditional density for  $X_t$ , given  $X_{t-1}$ , that is a mixture of the type

$$x \mapsto p_*(x \mid x_{t-1}) = G_*(x_{t-1})p_*(x_{t-1}, 0, x) + (1 - G_*(x_{t-1}))p_*(x_{t-1}, 1, x).$$

The covariate-dependence of the transition functions is reflected by the fact that  $G_*$  depends on  $X_{t-1}$ . Models which belong to the general class of mixture autoregressive models (e.g., Dueker et al. [2007], Tadjuidje et al. [2009], Dueker et al. [2011], Kalliovirta et al. [2015]) are covered by this framework.  $\triangle$

The examples above, as well as those that follow, illustrate that in many areas of application the stochastic process  $(X_t, S_t)_{t=0}^\infty$  is typically highly complex and it is natural/desirable to allow for feedback from past realizations of the observable process  $(X_t)_{t=0}^\infty$  to the law of the unobservable regime sequence  $(S_t)_{t=0}^\infty$ ; a tractable way for modeling such feedback is to allow the transition kernel  $Q_\theta$  to depend on  $X_{t-1}$ . This feature adds an additional level of complexity and with it sources of potential misspecification. For instance, a common assumption in most applications is that the transition probabilities of hidden regimes are time-invariant, an assumption which may result in misspecification of the transitions functions. In applications that allow for covariate-dependent transition functions, a somewhat more subtle and often overlooked source of misspecification, associated with endogeneity of the transition-driving covariates, may come into play. Specifically, in the context of models such as those in Examples 1 and 2, contemporaneous correlation between  $U_{1,t}$  and  $U_{2,t}$  is typically ignored and inference is based on the likelihood implied by the outcome equation alone; we consider this case in more detail in Sections 5.2 and 6.

Before discussing estimation of  $\theta$ , we give a result regarding the mixing and ergodicity properties of  $(X_t)_{t=0}^\infty$ . To do so, let  $\bar{P}_*^\kappa$  denote the true distribution over  $(X_t)_{t=0}^\infty$  when the distribution of  $(X_0, S_0)$  is  $\kappa$ . Under the following assumptions, Lemma 1 below ensures that there exists a Borel probability measure on  $\mathbb{X} \times \mathbb{S}$ , denoted henceforth by  $\nu$ , for which  $(X_t)_{t=0}^\infty$  is stationary and ergodic.

**Assumption 1.** *There exists a continuous function  $\underline{q} : \mathbb{X} \rightarrow \mathbb{R}_+ \setminus \{0\}$  such that, for all  $Q \in \{Q_\theta : \theta \in \Theta\} \cup Q_*$ ,  $Q(x, s, s') \geq \underline{q}(x)$  for all  $(s', s, x) \in \mathbb{S}^2 \times \mathbb{X}$ .*

**Assumption 2.** *There exist constants  $\lambda' \in (0, 1)$ ,  $\gamma \in (0, 1)$ ,  $b' > 0$  and  $R > 2b'/(1 - \gamma)$ , a lower semi-continuous function  $\mathcal{U} : \mathbb{X} \rightarrow [1, \infty)$ , and a measure  $\varpi \in \mathcal{P}(\mathbb{X})$  such that, for all  $s \in \mathbb{S}$ : (i)  $\int_{\mathbb{X}} \mathcal{U}(x') P_*(x, s, dx') \leq \gamma \mathcal{U}(x) + b' 1\{x \in A\}$ , with  $A \equiv \{x \in \mathbb{X} : \mathcal{U}(x) \leq R\}$ ; (ii)  $A$  is bounded and  $\varpi(A) > 0$ ; (iii)  $\inf_{x \in A} P_*(x, s, C) \geq \lambda' \varpi(C)$  for any Borel set  $C \subseteq \mathbb{X}$ .*

The following lemma establishes stationarity, ergodicity, and  $\beta$ -mixing of  $(X_t)_{t=0}^\infty$ .

**Lemma 1.** *Suppose Assumptions 1 and 2 hold. Then, there exists a  $\nu \in \mathcal{P}(\mathbb{X} \times \mathbb{S})$  such that, under  $\bar{P}_*^\nu$ ,  $(X_t)_{t=0}^\infty$  is stationary, ergodic, and  $\beta$ -mixing with mixing coefficients  $\beta_n = O(\gamma^n)$ ,  $n \in \mathbb{N}$ .*

*Proof.* See Supplemental Material SM.1. □

The result follows in a standard manner by using Assumptions 1 and 2 to establish that the implied transition kernel of the joint process  $(X_t, S_t)_{t=0}^\infty$  has a unique invariant distribution and also that it is Harris recurrent and aperiodic. This fact, in turn, is used to show that  $(X_t)_{t=0}^\infty$  is stationary, ergodic, and  $\beta$ -mixing at a geometric rate.

**Remark 1** (Discussion of Assumptions 1 and 2). Assumption 1 is an extension of a common assumption in the literature (cf. Douc et al. [2004b], Ailliot and Pène [2015]) to the case where the transition kernel of  $(S_t)_{t=0}^\infty$  depends on  $X_{t-1}$ . Allowing the lower bound  $\underline{q}$  to depend on  $x$  is especially relevant when the support of  $X_t$  is unbounded because, while  $\underline{q}(x) > 0$ , it is allowed to converge to zero as  $\|x\| \rightarrow \infty$ . Although this assumption is not innocuous, we view it as mild because it accommodates the typical specifications used in the literature, where  $Q$  is parameterized by a standard Gaussian or logistic cumulative distribution function and a single index  $x^\top \beta$ , with  $\beta$  restricted to take values in a bounded subset of a finite-dimensional Euclidean space.

Assumption 2(iii) is an analogous condition for the transition kernel  $P_*$ . By inspection of the proof of Lemma 1, it is easy to see that it suffices to obtain a minorization condition for the “joint” kernel, i.e.,  $\inf_{x \in A} P_*(x, s', C)Q(x, s, s') \geq \lambda \tilde{\omega}(C, s')$  for any Borel set  $C \subseteq \mathbb{X}$  and for some  $\tilde{\omega} \in \mathcal{P}(\mathbb{X} \times \mathbb{S})$  and  $\lambda \in (0, 1)$ . Thus, Assumptions 1(i) and 2(i) could be relaxed; e.g., the former could be relaxed to  $Q(x, s, s') \geq \underline{q}(x)\varrho(s')$ , where  $\varrho \in \mathcal{P}(\mathbb{S})$ , or the latter could be relaxed to  $\inf_{x \in A} P_*(x, s', C) \geq \lambda' \tilde{\omega}(C, s')$ , where  $\tilde{\omega} \in \mathcal{P}(\mathbb{X} \times \mathbb{S})$ .

Assumption 2(i),(ii) is a so-called Foster–Lyapunov drift condition; see Meyn and Tweedie [1993] and references therein for a discussion of the assumption. △

In view of Lemma 1, under  $\nu$ , the process  $(X_t)_{t=0}^\infty$  can be extended to a two-sided sequence  $(X_t)_{t=-\infty}^\infty$ . With a slight abuse of notation, we still use  $\bar{P}_*^\nu$  to denote the true probability distribution over  $(X_t, S_t)_{t=-\infty}^\infty$ ;  $\bar{P}_\theta^\nu$  is defined analogously for the model  $(Q_\theta, p_\theta, \nu)$ .<sup>4</sup>

---

<sup>4</sup>Throughout the text, we use  $\bar{P}_\theta^\nu$  to denote any marginal or conditional probabilities associated with  $\bar{P}_\theta^\nu$ ; the same holds for  $\bar{P}_*^\nu$ .



## 2.2 Parameter Estimation

For any  $T \in \mathbb{N}$ , let  $\ell_T^\nu : \mathbb{X}^{T+1} \times \Theta \rightarrow \mathbb{R}$  be the sample criterion function given by

$$\ell_T^\nu(X_0^T, \theta) = T^{-1} \sum_{t=1}^T \log p_t^\nu(X_t | X_0^{t-1}, \theta), \quad (1)$$

where  $p_t^\nu(X_t | X_0^{t-1}, \theta)$  denotes the conditional density of  $X_t$  given  $X_0^{t-1}$  for any  $\theta \in \Theta$ ; the latter is defined recursively as follows: for any  $t \geq 1$ ,

$$p_t^\nu(X_t | X_0^{t-1}, \theta) = \sum_{s' \in \mathbb{S}} \sum_{s \in \mathbb{S}} p_\theta(X_{t-1}, s', X_t) Q_\theta(X_{t-1}, s, s') \delta_t^{\theta, \nu}(s),$$

and  $s \mapsto \delta_t^{\theta, \nu}(s) \equiv \bar{P}_\theta^\nu(S_{t-1} = s | X_0^{t-1})$ . For each  $t \geq 2$  and any  $s \in \mathbb{S}$ ,  $s \mapsto \delta_t^{\theta, \nu}(s)$  satisfies the recursion

$$\delta_t^{\theta, \nu}(s) = \sum_{\tilde{s} \in \mathbb{S}} \frac{Q_\theta(X_{t-1}, \tilde{s}, s) p_\theta(X_{t-2}, \tilde{s}, X_{t-1}) \delta_{t-1}^{\theta, \nu}(\tilde{s})}{\sum_{s' \in \mathbb{S}} p_\theta(X_{t-2}, s', X_{t-1}) \delta_{t-1}^{\theta, \nu}(s')},$$

with  $s \mapsto \delta_1^{\theta, \nu}(s) = \sum_{\tilde{s} \in \mathbb{S}} Q_\theta(X_0, \tilde{s}, s) \nu(\tilde{s} | X_0)$ , where  $\nu(\cdot | \cdot)$  is the conditional density corresponding to  $\nu$ .

For a given initial distribution  $\kappa \in \mathcal{P}(\mathbb{X} \times \mathbb{S})$  over  $(X_0, S_0)$ , we define our estimator as  $\hat{\theta}_{\kappa, T}$ , where

$$\ell_T^\kappa(X_0^T, \hat{\theta}_{\kappa, T}) \geq \sup_{\theta \in \Theta} \ell_T^\kappa(X_0^T, \theta) - \eta_T, \quad (2)$$

for some  $\eta_T \geq 0$  and  $\eta_T = o(1)$ .

## 3 Consistency

The main result of this section establishes convergence of the estimator  $\hat{\theta}_{\nu, T}$  to the set of points in  $\Theta$  that are closest to the true model under the Kullback–Leibler information criterion.

Let  $H^* : \Theta \rightarrow \mathbb{R}_+ \cup \{\infty\}$  be the Kullback–Leibler information criterion  $\theta \mapsto H^*(\theta)$ , which is given by

$$H^*(\theta) = E_{\bar{P}_*^\nu} \left[ \log \frac{p_*^\nu(X_0 | X_{-\infty}^{-1})}{p^\nu(X_0 | X_{-\infty}^{-1}, \theta)} \right],$$

where, for any  $\theta \in \Theta$ ,  $p^\nu(X_0 | X_{-\infty}^{-1}, \theta)$  denotes the conditional density of  $X_0$  given  $X_{-\infty}^{-1}$  induced by  $(P_\theta, Q_\theta, \nu)$ , and  $p_*^\nu(X_0 | X_{-\infty}^{-1})$  is its counterpart

induced by the true transition kernels  $(P_*, Q_*, \nu)$ ; we refer the reader to the Supplemental Material [SM.3](#) for details about the construction of these objects and their properties.

Our results allow for misspecified models, and thus  $p_*^\nu \notin \{p^\nu(\cdot | \cdot, \theta) : \theta \in \Theta\}$ . Hence, as in [White \[1982\]](#), the relevant limiting set for our estimator is

$$\Theta_* = \arg \min_{\theta \in \Theta} H^*(\theta),$$

which is the *pseudo-true parameter (set)* that minimizes the Kullback–Leibler information criterion. Under Assumption [3](#) below,  $\Theta_*$  is non-empty and compact by the Weierstrass Theorem.

**Assumption 3.** (i)  $\Theta$  is compact; (ii)  $H^*$  exists and is lower semi-continuous.

The following additional assumptions are used to establish the main result. To state these, for any  $\delta > 0$  and any  $\dot{\theta} \in \Theta$ , let  $B(\delta, \dot{\theta}) \equiv \{\theta \in \Theta : \|\dot{\theta} - \theta\| < \delta\}$ .

**Assumption 4.** (i) For any  $\epsilon > 0$ , there exists some  $\delta > 0$  such that

$$\max_{\dot{\theta} \in \Theta} E_{\bar{P}_*^\nu} \left[ \sup_{\theta \in B(\delta, \dot{\theta})} \frac{p^\nu(X_0 | X_{-\infty}^{-1}, \theta)}{p^\nu(X_0 | X_{-\infty}^{-1}, \dot{\theta})} \right] \leq 1 + \epsilon;$$

(ii) there exists a function  $(x, x') \mapsto C(x, x') \in \mathbb{R}_+$  such that  $\sup_{\theta \in \Theta} \frac{\max_{s \in \mathbb{S}} p_\theta(X, s, X')}{\min_{s \in \mathbb{S}} p_\theta(X, s, X')} \leq C(X, X')$  and  $\frac{\max_{s \in \mathbb{S}} p_*(X, s, X')}{\min_{s \in \mathbb{S}} p_*(X, s, X')} \leq C(X, X')$  a.s.- $\bar{P}_*^\nu$ .

**Assumption 5.**  $T^{-1} \sum_{t=1}^T \max\{1, C(X_{t-1}, X_t)\} \prod_{i=0}^{t-1} (1 - \underline{q}(X_i)) = o_{\bar{P}_*^\nu}(1)$ .

**Remark 2** (Discussion of Assumptions [3](#), [4](#) and [5](#)). Assumption [3](#)(i) is standard. Assumption [3](#)(ii) is high-level but can be obtained from lower-level conditions (e.g., by following the reasoning in Proposition 11 of [Douc and Moulines \[2012\]](#)).

Assumption [4](#)(i) is a high-level condition used for establishing uniform law of large numbers results (see Lemma [3](#) in Section [A.1](#)). Assumption [4](#)(ii) is akin to Assumption A4 in [Bickel et al. \[1998\]](#); it essentially restricts the support of  $p_\theta$  and  $p_*$  for different values of the state variable.<sup>5</sup>

Assumption [5](#) essentially requires that  $\underline{q}$  is not “too close” to zero on average and that  $C(X_{t-1}, X_t)$  is finite a.s.- $\bar{P}_*^\nu$ . For instance, if  $\underline{q}(x) \geq c$  for some  $c > 0$ , then Assumption [5](#) is automatically satisfied provided  $E_{\bar{P}_*^\nu}[C(X_0, X_1)] < \infty$ . Moreover, by exploiting the fact that  $(X_t)_t$  is  $\beta$ -mixing (see Lemma [1](#)), Lemma [13](#) in the Supplemental Material [SM.4](#) provides sufficient conditions of the form  $E_{\bar{P}_*^\nu}[\underline{q}(X_1)] > 0$  and  $E_{\bar{P}_*^\nu}[C(X_1, X_0)^l] < \infty$  for some  $l > 1$ .<sup>6</sup>  $\triangle$

<sup>5</sup>The second part of Assumption [4](#)(ii) is used to show that  $p_*^\nu(\cdot | X_{-\infty}^{-1})$  integrates to one.

<sup>6</sup>We thank a referee and the editor for suggestions on how to weaken Assumption [4](#), which lead to these sufficient conditions for Assumption [5](#).

We now establish consistency of the estimator defined by (2).

**Theorem 1.** *Suppose Assumptions 1–4 hold. Then,  $d_\Theta(\hat{\theta}_{\nu,T}, \Theta_*) = o_{\bar{P}_*^\nu}(1)$ .<sup>7</sup>*

*Proof.* See Appendix A.1.  $\square$

This result is analogous to Theorem 2 in Douc and Moulines [2012] but for a somewhat different setup; specifically, we allow for autoregressive dynamics and covariate-dependent transition probabilities, but restrict  $\mathbb{S}$  to be finite.<sup>8</sup>

Clearly, if the model is correctly specified and point identified, i.e., there exists a  $\theta \in \Theta$  such that  $(P_*, Q_*) = (P_\theta, Q_\theta)$ , then  $\Theta_* = \{\theta\}$  and our estimator converges in  $\bar{P}_*^\nu$ -probability to this point. If, however, the model is misspecified, our estimator converges to the set of parameters that is closest to the true set, when closeness is measured by means of the Kullback–Leibler information criterion (cf. White [1982], Douc and Moulines [2012]).

To prove Theorem 1, we first show that  $T^{-1} \sum_{t=1}^T \log p_t^\nu(X_t \mid X_0^{t-1}, \theta)$  is well-approximated by  $T^{-1} \sum_{t=1}^T \log p^\nu(X_t \mid X_{-\infty}^{t-1}, \theta)$  (see Lemma 2 in Appendix A.1). Second, relying on ergodicity (Lemma 1) and Assumption 4, we establish a *uniform* law of large numbers for the latter quantity (see Lemma 3 in Appendix A.1). The proof of consistency then follows the standard Wald approach.

The approximation result in the first step relies on “mixing” results for the process  $(S_t)_{t=-\infty}^\infty$ , given  $(X_t)_{t=-\infty}^\infty$ . The following theorem, which might be of independent interest, establishes such a “mixing” result in our setting.

**Theorem 2.** *Take any  $(j, m) \in \mathbb{N}^2$ . Suppose that, for any  $\theta \in \Theta$ , there exist mappings  $x \mapsto \varrho(x, \cdot) \in \mathcal{P}(\mathbb{S})$  and  $q : \mathbb{X} \rightarrow \mathbb{R}_+$  such that, for all  $(s, s') \in \mathbb{S}^2$ ,*

$$Q_\theta(X, s, s') \geq \underline{q}(X) \varrho(X, s') \quad \text{a.s.}-\bar{P}_*^\nu. \quad (3)$$

*Then,*<sup>9</sup>

$$\max_{(b,c) \in \mathbb{S}^2} \left\| \bar{P}_\theta^\nu(S_{j+1} = \cdot \mid S_{-m} = b, X_{-m}^j) - \bar{P}_\theta^\nu(S_{j+1} = \cdot \mid S_{-m} = c, X_{-m}^j) \right\|_1 \leq \prod_{n=-m}^j (1 - \underline{q}(X_n)).$$

*Proof.* See Appendix A.2.  $\square$

<sup>7</sup>For any set  $A \subseteq \Theta$ ,  $d_\Theta(\theta, A) \equiv \inf_{\hat{\theta} \in A} \|\theta - \hat{\theta}\|$ .

<sup>8</sup>Finiteness of  $\mathbb{S}$  simplifies the proofs as we do not have to be concerned with uniformity issues of certain quantities as functions of the hidden state. By combining techniques available in the literature (e.g., Douc and Moulines [2012]) – which essentially amount to imposing requirements like compactness of  $\mathbb{S}$  and continuity – with ours, we conjecture that our results can be extended to the case where  $\mathbb{S}$  is a compact set.

<sup>9</sup>For any  $P, Q \in \mathcal{P}(\mathbb{S})$ ,  $\|P - Q\|_1 \equiv \sum_{s \in \mathbb{S}} |P(s) - Q(s)|$ .

This result is analogous to results in Douc et al. [2004b] (e.g., their Lemma 1 and Corollary 1) but for a more general transition function  $Q_\theta$  (albeit under the requirement that  $\mathbb{S}$  is finite). Specifically, we allow  $Q_\theta$  to depend on  $X$ , and do not restrict the lower bound in (3) to be uniform in  $s'$ ; these features are, to our knowledge, novel.

The proof of Theorem 2 relies on bounding the Dobrushin coefficient of the transition kernel  $\bar{P}_\theta(S_{l+1} = \cdot | S_l = \cdot, X_{-m}^j)$  by  $1 - \underline{q}(X_l)$ , for each  $l \in \{-m, \dots, j\}$ . The case where  $\varrho(x, \cdot)$ , in condition (3), is uniformly bounded from below has been studied in the literature, and such a bound can be obtained by elementary calculations. For the general case, however, the standard technique cannot be used, and we develop a novel, to our knowledge, approach based on coupling techniques; see Appendix A.2 for details.

**Remark 3.** Remark 1 and the fact that Theorem 2 is established under condition (3), imply that, in regards to consistency, Assumption 1 could be replaced by the weaker condition (3). This remark, however, does not extend to the LAN results of Section 4, since we do not know whether Assumption 1 could be weakened to condition (3) in this case.  $\triangle$

We discuss next a canonical example that encompasses many commonly used specifications, such as those in Examples 1 and 2. The purpose of this example is to verify our assumptions under primitive and low-level conditions.

**Example 4** (Canonical Example). We verify the regularity conditions for a model with  $\mathbb{S} = \{0, 1\}$  and

$$\begin{aligned} X_t &= \mu(S_t) + \Phi^\top X_{t-1} + \Sigma^{1/2}(S_t) \varepsilon_t, \\ S_t &\sim Q_{\bar{\vartheta}}(X_{t-1}, S_{t-1}, \cdot), \end{aligned}$$

where  $(\varepsilon_t)_t \sim \text{i.i.d. } \mathcal{N}(0, I)$ ,  $I$  being the identity matrix, and  $x \mapsto Q_{\bar{\vartheta}}(x, s, s) \equiv \Pr(S_t = s | X_{t-1} = x, S_{t-1} = s) = \Psi(x^\top \bar{\vartheta}_s)$  for  $s \in \mathbb{S}$ , where  $\Psi$  is a full-support, continuous cumulative distribution function (e.g., logistic or normal). It is assumed that the parameter set  $\Theta$  is compact and that any  $\theta \equiv ((\mu(s), \Sigma(s))_{s \in \mathbb{S}}, \Phi, \bar{\vartheta})$  in it is such that  $\Sigma(\cdot)$  and  $\Phi\Phi^\top$  have eigenvalues uniformly bounded away from zero and infinity. For simplicity, we assume that the true model is indexed by  $((\mu_*(s), \Sigma_*(s))_{s \in \mathbb{S}}, \Phi_*, \bar{\vartheta}_*)$ , for which analogous conditions hold.

Let  $q(x) \equiv \inf_b \Psi(x^\top b)$  and note that  $q(x) > 0$  for each finite  $x$  as  $\Psi$  has full support. Thus, Assumption 1 holds and  $E_{\bar{P}_*}[q(X)] > 0$ . The conditions of Assumption 2 follow by the results in Douc et al. [2004a]; see Lemma 14 in the Supplemental Material SM.5 for the formal argument. In Lemma 15 in the Supplemental Material SM.5, it is shown that

$$C^{-1} \underline{p}(x, y) \leq f_{\mathcal{N}}(\{y - \Phi^\top x - \mu(s)\} \Sigma^{-1/2}(s)) \leq C \bar{p}(x, y),$$

for some  $C \geq 1$ , where  $f_{\mathcal{N}}$  is the  $\mathcal{N}(0, I)$  probability density function,

$$\begin{aligned}\underline{p}(x, y) &\equiv \exp\{-0.5(a_1 \|y\|^2 + a_2 \|x\|^2 + 2a_3 \|x\| \|y\|) - a_4 \|y\| - a_5 \|x\|\}, \\ \bar{p}(x, y) &\equiv \exp\{-0.5(b_1 \|y\|^2 + b_2 \|x\|^2 - 2b_3 \|x\| \|y\|) + b_4 \|y\| + b_5 \|x\|\},\end{aligned}$$

and  $a_i, b_i$  ( $i = 1, \dots, 5$ ) are positive constants that are functions of uniform bounds of  $\Sigma(\cdot)$ ,  $\mu(\cdot)$  and  $\Phi$  over  $\Theta$ ; they are defined in Lemmas 15 and 16 in the Supplemental Material SM.4. Therefore, by Lemma 12 in the Supplemental Material SM.3, it follows that Assumptions 3 and 4 hold with  $(x, x') \mapsto C(x, x') \equiv \bar{p}(x, x')/\underline{p}(x, x')$ , provided that  $E_{\bar{P}_\nu}[\exp\{-0.5l((b_1 - a_1) \|Y\|^2 + (b_2 - a_2) \|X\|^2 - 2(b_3 + a_3) \|X\| \|Y\|) + l(b_4 + a_4) \|y\| + l(b_5 + a_5) \|X\|\}] < \infty$  for some  $l \geq 1$ . Lemma 16 in the Supplemental Material SM.5 shows that the latter condition holds under some restrictions on the eigenvalues of  $\Sigma(\cdot)$ ,  $\Sigma_*(\cdot)$ ,  $\Phi\Phi^\top$ , and  $\Phi_*\Phi_*^\top$  (see the aforementioned lemma and its subsequent remark for the exact formulation and more details). Finally, under these conditions, Lemma 13 in the Supplemental Material SM.4 shows that Assumption 5 holds.  $\triangle$

Lastly, we note that further characterization and interpretation of the pseudo-true parameter set  $\Theta_*$  is not straightforward in our general setup, and we believe that progress in this direction should be made on a case-by-case basis. While pursuing this program is outside the scope of the present paper, in Section 5 we present numerical results that attempt to shed some light on the characterization of this set. In addition, Example 6 in the Supplemental Material SM.6 provides analytical results on the characterization of  $\Theta_*$  for a subclass of the models in Example 2, namely hidden Markov models with covariate-dependent transition probabilities, when the (misspecified) model is a simple mixture.

## 4 Asymptotic Distribution Theory

In this section, we establish a LAN property ([Ibragimov and Has'minskii, 1981, Ch. II], Le Cam [1986]) for our model and an asymptotic linear representation for our estimator, from which asymptotic normality of the estimator is inferred. For this, we require the following assumptions.

**Assumption 6.** (i)  $\Theta_* = \{\theta_*\} \subset \text{int}(\Theta)$ ; (ii)  $\theta \mapsto p_\theta(X, s, X')$  and  $\theta \mapsto Q_\theta(X, s, s')$  are twice continuously differentiable a.s.- $\bar{P}_\nu^*$  for all  $(s, s') \in \mathbb{S}^2$ .

**Assumption 7.** For some  $\delta > 0$  and  $a \geq 1$ , and for all  $(s', s) \in \mathbb{S}^2$ : (i)

$$E_{\bar{P}_\nu^*} \left[ \sup_{\theta \in B(\delta, \theta_*)} \|\nabla_\theta \log p_\theta(X, s, X')\|^{2a} \right] < \infty \text{ and } E_{\bar{P}_\nu^*} \left[ \sup_{\theta \in B(\delta, \theta_*)} \|\nabla_\theta \log Q_\theta(X, s, s')\|^{2a} \right] < \infty;$$

(ii)

$$E_{\bar{P}^\nu_*} \left[ \sup_{\theta \in B(\delta, \theta_*)} \|\nabla_\theta^2 \log p_\theta(X, s, X')\|^{2a} \right] < \infty \text{ and } E_{\bar{P}^\nu_*} \left[ \sup_{\theta \in B(\delta, \theta_*)} \|\nabla_\theta^2 \log Q_\theta(X, s, s')\|^{2a} \right] < \infty.$$

**Assumption 8.**  $\sum_{j=0}^\infty \left( E_{\bar{P}^\nu_*} \left[ \prod_{i=0}^j (1 - \underline{q}(X_i))^{\frac{2a}{1-a}} \right] \right)^{p(\frac{1-a}{2a})} < \infty$  for some  $p \in (0, 2/3)$  (and the same  $a \geq 1$  that appears in Assumption 7).

**Remark 4** (Discussion of Assumptions 6, 7 and 8). Part (i) of Assumption 6 is standard in the literature. The restriction that  $\Theta_*$  is a singleton could be relaxed using the ideas of Liu and Shao [2003] for non-identified ML estimators. This extension, albeit interesting, would present nuances that are beyond the scope of the present paper. Part (ii) of Assumption 6 is also standard, and so is Assumption 7 (see Bickel et al. [1998] for a discussion). Finally, Assumption 8 is a strengthening of Assumption 5, and is required in order to establish the existence of a random sequence  $(\Delta_t(\theta_*))_t$  which approximates the “score” function well (in the sense of Lemma 18 in Appendix A.3). As was the case with Assumption 5, Lemma 13 in the Supplemental Material SM.4 provides sufficient conditions of the form  $E_{\bar{P}^\nu_*} [(1 - \underline{q}(X_1))^{2a/(1-a)}] < 1$ .  $\triangle$

The next theorem establishes a LAN-type property for the log-likelihood criterion function defined in (1).

**Theorem 3.** Suppose Assumptions 1, 2, 6, 7 and 8 hold. Then, there exists a stationary and ergodic process  $(\Delta_t(\theta_*))_t$  in  $L^2(\bar{P}^\nu_*)$ , a sequence of negative definite matrices  $(\xi_t(\theta_*))_t$ , and a compact set  $K \subseteq \Theta$  that includes 0, such that, for any  $v \in K$ ,

$$\begin{aligned} \ell_T^\nu(X_0^T, \theta_* + v) - \ell_T^\nu(X_0^T, \theta_*) &= v^\top \left( T^{-1} \sum_{t=0}^T \Delta_t(\theta_*) + o_{\bar{P}^\nu_*}(T^{-1/2}) \right) \\ &\quad + (1/2)v^\top \left( T^{-1} \sum_{t=0}^T \xi_t(\theta_*) + o_{\bar{P}^\nu_*}(1) \right) v + R_T(v), \end{aligned}$$

where  $v \mapsto R_T(v) \in \mathbb{R}$  is such that  $\lim_{\delta \rightarrow 0} \bar{P}^\nu_* \left( \sup_{v \in B(\delta, 0)} \|v\|^{-2} R_T(v) \geq \epsilon \right) = 0$  for any  $\epsilon > 0$  and any  $T \in \mathbb{N}$ .

*Proof.* See Appendix A.3.  $\square$

Theorem 3 extends the results in Bickel and Ritov [1996] and Bickel et al. [1998] (see their remark on p. 1620) to a more general setup which allows for covariate-dependent transition probabilities, autoregressive dynamics, and misspecified models. The proof develops along the same lines as theirs. The main

difference relates to the way in which one establishes that the “score”  $\nabla_{\theta} \ell_T^{\nu}(\cdot, \theta_*)$  and the Hessian  $\nabla_{\theta}^2 \ell_T^{\nu}(\cdot, \theta_*)$  can be approximated by  $T^{-1} \sum_{t=0}^T \Delta_t(\theta_*)$  and  $T^{-1} \sum_{t=0}^T \zeta_t(\theta_*)$ , respectively (see Lemmas 7 and 8 in Appendix A.3). As mentioned earlier, these approximations rely on “mixing” properties of the *temporally inhomogeneous* hidden Markov chain; see Lemma 22 in the Supplemental Material SM.8.

Theorem 3 may be used to establish the following asymptotic linear representation for our estimator in terms of  $(\Delta_t(\theta_*))_{t=0}^{\infty}$  and  $(\xi_t(\theta_*))_{t=0}^{\infty}$ .

**Theorem 4.** *Suppose Assumptions 1–8 hold and  $\eta_T = o(T^{-1})$ . Then,*

$$\frac{\sqrt{T}(\hat{\theta}_{\nu,T} - \theta_*)}{\sqrt{\text{tr}\{\Sigma_T(\theta_*)\}}} = -\{E_{\bar{P}_*^{\nu}}[\xi_1(\theta_*)] + o_{\bar{P}_*^{\nu}}(1)\}^{-1} T^{-1/2} \sum_{t=0}^T \frac{\Delta_t(\theta_*)}{\sqrt{\text{tr}\{\Sigma_T(\theta_*)\}}} + o_{\bar{P}_*^{\nu}}(1),$$

where  $\Sigma_T(\theta_*) \equiv T^{-1} E_{\bar{P}_*^{\nu}}[\{\sum_{t=0}^T \Delta_t(\theta_*)\} \{\sum_{t=0}^T \Delta_t(\theta_*)^{\top}\}]$ .

*Proof.* See Appendix A.3. □

Theorem 4 readily implies that, if  $T^{-1/2} \Sigma_T(\theta_*)^{-1/2} \sum_{t=0}^T \Delta_t(\theta_*) \Rightarrow_{\bar{P}_*^{\nu}} \mathcal{N}(0, I)$ , then

$$\sqrt{T} \Sigma_T(\theta_*)^{-1/2} E_{\bar{P}_*^{\nu}}[\xi_1(\theta_*)](\hat{\theta}_{\nu,T} - \theta_*) \Rightarrow_{\bar{P}_*^{\nu}} \mathcal{N}(0, I). \quad (4)$$

(If  $\Sigma_T(\theta_*) \rightarrow \Sigma(\theta_*)$ ,  $\Sigma_T(\theta_*)$  may be replaced by  $\Sigma(\theta_*)$  in these statements). This result is akin to results in White [1982] and shares the same features, i.e., the asymptotic covariance matrix  $\Omega_T(\theta_*) \equiv (E_{\bar{P}_*^{\nu}}[\xi_1(\theta_*)])^{-1} \Sigma_T(\theta_*) (E_{\bar{P}_*^{\nu}}[\xi_1(\theta_*)])^{-1}$  has a “sandwich” form and the familiar Fisher information equality does not necessarily hold (see also [White, 1994, Ch. 6]).

The next theorem complements Theorem 4 by presenting results on consistent estimation of the asymptotic covariance matrix of  $\hat{\theta}_{\nu,T}$ , vis.,  $\Omega_T(\theta_*)$ , in both correctly specified and potentially misspecified models. In the latter case, the proposed estimator is of the “heteroskedasticity and autocorrelation consistent” type. As in many other results for such estimators (e.g., Newey and West [1987], [White, 1994, Ch. 8]), consistency requires some conditions over the score covariance process  $(\Delta_{t+\tau}(\cdot) \Delta_t(\cdot)^{\top})_{t=0}^T$ ,  $\tau \in \mathbb{N}$ , namely continuity and law of large numbers results (see Lemma 25 in the Supplemental Material SM.9). In what follows, the modulus of continuity is defined as

$$\ddot{\omega}(\delta') \equiv \max_t \left\| \sup_{\|\theta - \theta_*\| < \delta'} \|\Delta_t(\theta) \Delta_0(\theta)^{\top} - \Delta_t(\theta_*) \Delta_0(\theta_*)^{\top}\| \right\|_{L^1(\bar{P}_*^{\nu})},$$

for any  $\delta' \in (0, \delta]$ , where  $\delta$  is as in Assumption 7.<sup>10</sup>

<sup>10</sup>Lemma 25 in the Supplementary Material SM.9 shows that the modulus of continuity converges to zero as  $\delta$  vanishes.

**Theorem 5.** Suppose the assumptions of Theorem 4 hold.

- (a) If  $\theta_*$  is such that, for any  $t \geq 0$  and  $T \geq 1$ ,  $p_t^\nu(\cdot | X_{t-T}^{t-1}; \theta^*) = p_*^\nu(\cdot | X_{t-T}^{t-1})$ , then

$$\|\Omega_T(\theta_*) - \{-H_T^{-1}(\hat{\theta}_{\nu,T})\}\| = o_{\bar{P}_*^\nu}(1),$$

where

$$H_T(\theta) \equiv T^{-1} \sum_{t=1}^T \nabla_\theta^2 \log p_t^\nu(X_t | X_0^{t-1}, \theta).$$

- (b) If, for any  $l \geq 0$ ,  $\|E_{\bar{P}_*^\nu}[\Delta_l(\theta_*)\Delta_0(\theta_*)^\top]\| \leq \bar{v}(l)$  for some integrable function  $l \mapsto \bar{v}(l) \in \mathbb{R}_+$ , then

$$\|\Omega_T(\theta_*) - H_T^{-1}(\hat{\theta}_{\nu,T})J_T(\hat{\theta}_{\nu,T})H_T^{-1}(\hat{\theta}_{\nu,T})\| = o_{\bar{P}_*^\nu}(1),$$

where

$$\begin{aligned} J_T(\theta) &\equiv T^{-1} \sum_{t=1}^T \nabla_\theta \log p_t^\nu(X_t | X_0^{t-1}, \theta) \nabla_\theta \log p_t^\nu(X_t | X_0^{t-1}, \theta)^\top \\ &\quad + \sum_{\tau=1}^{L_T} \frac{\omega(\tau, L_T)}{T - \tau} \sum_{t=1}^{T-\tau} \nabla_\theta \log p_{t+\tau}^\nu(X_{t+\tau} | X_0^{t+\tau-1}, \theta) \nabla_\theta \log p_t^\nu(X_t | X_0^{t-1}, \theta)^\top \\ &\quad + \sum_{\tau=1}^{L_T} \frac{\omega(\tau, L_T)}{T - \tau} \sum_{t=1}^{T-\tau} \nabla_\theta \log p_t^\nu(X_t | X_0^{t-1}, \theta) \nabla_\theta \log p_{t+\tau}^\nu(X_{t+\tau} | X_0^{t+\tau-1}, \theta)^\top, \end{aligned}$$

$\omega(\cdot, \cdot)$  are bounded real weights with  $\lim_{T \rightarrow \infty} \omega(\tau, L_T) = 1$  for all  $\tau \geq 1$ , and  $(L_T)_{T=1}^\infty \subseteq \mathbb{N}$  is such that  $L_T(\ddot{w}(T^{-1/2} \log \log T) \log \log T + r_T + T^{-1/2}) = o(1)$ , with  $(r_T)_T$  being a positive sequence converging to zero.

*Proof.* See Supplemental Material SM.9.  $\square$

Part (a) of Theorem 5 deals with correctly specified models and makes use of the Fisher information equality. Part (b) provides a consistent covariance estimator in the general case of potentially misspecified models. The conditions for the weights  $\omega(\tau, L_T)$  and the tuning parameters  $(L_T)_T$  are standard for estimators of this type.<sup>11</sup> The terms  $\ddot{w}(T^{-1/2} \log \log T) \log \log T$  and  $r_T$  in the growth condition for  $L_T$  are analogous to those appearing in Newey and West [1987] and arise as “costs” of working with  $\nabla_\theta \log p_t^\nu(X_t | X_0^{t-1}, \hat{\theta}_{\nu,T})$  as opposed

<sup>11</sup>The additional condition that  $\omega(\tau, L_T) = \sum_{j=1+\tau}^{L_T} c(j, L_T)c(j - \tau, L_T)$  for some constants  $c(1, L_T), \dots, c(L_T, L_T)$  guarantees that  $J_T(\hat{\theta}_{\nu,T})$  is positive semidefinite. Weights obtained from the commonly used Bartlett, Parzen and quadratic-spectral kernels satisfy this condition.



to  $\nabla_\theta \log p_t^\nu(X_t | X_0^{t-1}, \theta_*)$ , and of working with sample averages as opposed to population means, respectively.<sup>12</sup> On the other hand, the  $T^{-1/2}$  term does not appear in Newey and West [1987] and arises because we need to approximate  $\Delta_t(\theta_*)$ , that depends on  $X_{-\infty}^t$ , with the score, that depends on  $X_0^t$ .

Theorems 4 and 5, together with an asymptotic normality result such as (4), provide the means for constructing asymptotically correct confidence sets and hypotheses tests for  $\theta_*$ . In correctly specified models,  $(\Delta_t(\theta_*))_{t=0}^\infty$  is a martingale difference sequence, and thus result (4) can be obtained by invoking a martingale central limit theorem. In potentially misspecified models,  $(\Delta_t(\theta_*))_{t=0}^\infty$  will not, in general, be a martingale difference sequence, so one should use a different approach; in some situations, a central limit theorem for  $\beta$ -mixing sequences can be used instead. The following corollary formalizes this discussion.

**Corollary 1.** *Suppose the assumptions of Theorem 4 hold.*

- (a) *If  $\theta_*$  is such that, for any  $t \geq 0$  and  $T \geq 1$ ,  $p_t^\nu(\cdot | X_{t-T}^{t-1}; \theta^*) = p_*^\nu(\cdot | X_{t-T}^{t-1})$ , then*

$$\sqrt{T}\{-H_T(\hat{\theta}_{\nu,T})\}^{1/2}(\hat{\theta}_{\nu,T} - \theta_*) \Rightarrow_{\bar{P}_*^\nu} \mathcal{N}(0, I).$$

- (b) *If there exists  $\bar{L} > 0$  such that, for any  $k, T > \bar{L}$ ,  $p_k^\nu(X_k | X_{k-T}^{k-1}; \theta^*) = p_k^\nu(X_k | X_{k-\bar{L}}^{k-1}; \theta^*)$ ,  $\liminf_{T \rightarrow \infty} e_{\min}(\Sigma_T(\theta_*)) > 0$ ,<sup>13</sup> and  $E_{\bar{P}_*^\nu}[\|\Delta_1(\theta_*)\|^{4+4\delta}] < \infty$  for some  $\delta > 0$ , then*

$$\sqrt{T}\hat{\Omega}_T(\hat{\theta}_{\nu,T})^{-1/2}(\hat{\theta}_{\nu,T} - \theta_*) \Rightarrow_{\bar{P}_*^\nu} \mathcal{N}(0, I),$$

$$\text{where } \hat{\Omega}_T(\hat{\theta}_{\nu,T}) \equiv H_T^{-1}(\hat{\theta}_{\nu,T})J_T(\hat{\theta}_{\nu,T})H_T^{-1}(\hat{\theta}_{\nu,T}).$$

*Proof.* See Appendix A.3. □

Regarding the asymptotic normality result, both parts of the corollary rely on the structure of the process defined by the random variables  $\nabla_\theta \log p_k^\nu(X_k | X_{-\infty}^{k-1}; \theta^*)$ ; this is a martingale difference sequence in part (a) and a geometrically  $\beta$ -mixing sequence in part (b). The latter result is a consequence of the  $\beta$ -mixing structure of  $(X_t)_{t=-\infty}^\infty$  and of the fact that the conditional density at  $\theta_*$  depends on a finite number of lags. Examples of models for which such properties hold true are mixture models (cf. Example 6 in the Supplemental Material SM.6) and mixture autoregressive models (cf. Example 3).<sup>14</sup>

<sup>12</sup>The  $\log \log T$  factor is used in order to avoid working with constants.

<sup>13</sup>For any real symmetric matrix  $A$ ,  $e_{\min}(A)$  denotes its minimum eigenvalue.

<sup>14</sup>At this level of generality, we cannot establish an asymptotic normality result in the general case where the model is misspecified and  $\nabla_\theta \log p_k^\nu(X_k | X_{-\infty}^{k-1}; \theta^*)$  depends on the entire  $X_{-\infty}^{k-1}$ . The reason is that, although the process  $X_{-\infty}^\infty$  is  $\beta$ -mixing, there is no guarantee that  $\nabla_\theta \log p_k^\nu(X_k | X_{-\infty}^{k-1}; \theta^*)$  inherits these mixing properties, or that it is a martingale difference sequence.

Regarding the covariance estimators used in parts (a) and (b) of the corollary, these rely on the corresponding parts of Theorem 5. The result in part (b) is established by exploiting the  $\beta$ -mixing structure of the score process  $(\Delta_k(\theta_*))_{k=-\infty}^{\infty}$  and the finiteness of its  $4 + 4\delta$  moments (under  $\bar{P}_*^\nu$ ) for some  $\delta > 0$ . Lemma 9 in Appendix A.3 shows that  $r_T = L_T T^{-1/2}$ , and, since  $\theta \mapsto \log p_\theta$  and  $\theta \mapsto \log Q_\theta$  are smooth (see Assumptions 6 and 7), it follows that  $\delta' \mapsto \ddot{w}(\delta') = C\delta'$  for some finite constant  $C$ . Hence, the growth condition on  $(L_T)_T$  translates into  $L_T T^{-1/4} \log \log T = o(1)$ , which is analogous to that in Newey and West [1987] (apart from the  $\log \log T$  factor, which was introduced in Theorem 5 for convenience).

The next example verifies the assumptions in the context of the models considered in Example 4.

**Example 5.** In view of the results in Example 4, we only need to verify Assumptions 6–8. Part (i) of Assumption 6 is standard and is directly imposed, while part (ii) follows from the setup of the example. Assumption 7 follows by the continuity of the derivatives. Finally, Lemma 13 in the Supplemental Material SM.2 implies that Assumption 8 holds.

Thus, Theorem 4 holds for the class of models considered in Example 4. In particular, in the correctly specified case, Corollary 1(a) guarantees asymptotic normality of the studentized ML estimator of  $\theta_*$ , thereby providing the basis for inference. These results are, to our knowledge, new in the context of Markov-switching autoregressive models with covariate-dependent transition probabilities.  $\triangle$

## 5 Monte Carlo Simulations

The objective in this section is twofold. First, to assess the quality of approximations provided by our asymptotic results by examining the finite-sample properties of the ML estimator and related statistics in a correctly specified Markov-switching autoregressive model with covariate-dependent transition probabilities. Second, to explore the effects of a type of empirically relevant misspecification which involves the use of an incomplete approximation to the likelihood function that ignores potential contemporaneous correlation between the observation variable ( $Y_t$ ) and the variable ( $Z_t$ ) upon the lagged value of which the transition probabilities depend.

Monte Carlo experiments are based on artificial data  $(X_t = (Y_t, Z_t))_t$  generated according to the equations

$$Y_t = \mu_0(1 - S_t) + \mu_1 S_t + \phi Y_{t-1} + [\sigma_0(1 - S_t) + \sigma_1 S_t] U_{1,t}, \quad (5)$$

$$Z_t = \mu_2 + \psi Z_{t-1} + \sigma_2 U_{2,t}, \quad (6)$$

for  $t \in \mathbb{N}$ , with  $X_0 = (0.5, 0.4)$ ,  $\mu_0 = -\mu_1 = 1$ ,  $\phi = 0.9$ ,  $\sigma_0 = \sigma_1 = 1$ ,  $\mu_2 = 0.2$ ,  $\psi = 0.8$ ,  $\sigma_2 = 0.6$ , and  $(U_{1,t}, U_{2,t})_t \sim \text{i.i.d. } \mathcal{N}\left(\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix}\right)$  with  $\rho \in \{0, 0.8\}$ . The regimes  $(S_t)_t$  are a Markov chain on  $\{0, 1\}$ , independent of  $(U_{1,t}, U_{2,t})_t$ , with transition probabilities

$$Q_\theta(z, s, s) \equiv \Pr(S_t = s \mid Z_{t-1} = z, S_{t-1} = s) = [1 + \exp(-\alpha_s - \beta_s z)]^{-1}, \quad (7)$$

for  $s \in \{0, 1\}$ , where  $\alpha_0 = \alpha_1 = 2$  and  $\beta_0 = -\beta_1 = -0.5$ . The model defined by (5)–(7) is a prototypical Markov-switching autoregressive model with covariate-dependent transition probabilities (cf. Example 2). In each of 1000 independent Monte Carlo replications,  $100 + T$  data points for  $(X_t)_t$  are generated, with  $T \in \{200, 800, 1600, 3200\}$ , and the last  $T$  points are used to compute estimates of the parameters of interest. In order to conserve space, only a selection of the results are reported (the full set of results is available upon request).

## 5.1 Correct Specification

In the first set of experiments, we consider estimation of the parameters of the model in (5)–(7) using the likelihood function based on the conditional distribution of  $X_t$  given  $X_0^{t-1}$ . Table 1 reports the deviation of the mean of the finite-sample distributions of the ML estimators of the elements of  $\vartheta = (\mu_0, \mu_1, \phi, \sigma_0, \sigma_1, \alpha_0, \beta_0, \alpha_1, \beta_1)$  from the corresponding true parameter values (bias) when  $\rho = 0.8$ . We also report the ratio of the sampling standard deviation of the estimators to the estimated standard errors (averaged across replications); the latter are computed using the Hessian estimator (cf. Theorem 5(a)).<sup>15</sup> Although the estimators of  $\beta_0$  and  $\beta_1$  are somewhat biased when  $T = 200$ , bias is insignificant in the rest of the cases. Estimated standard errors are somewhat downwards biased in most cases but, unless  $T$  is small, the bias is not generally substantial and decreases as  $T$  increases.

We also examine conventional hypothesis tests for  $\vartheta$ . Table 2 reports the rejection frequencies of: (i) a  $t$ -type test of  $\mathcal{H}_0 : \vartheta_j = \vartheta_j^*$  versus  $\mathcal{H}_1 : \vartheta_j \neq \vartheta_j^*$ , where  $\vartheta_j$  is the  $j$ -th element of  $\vartheta$  and  $\vartheta_j^*$  is its true value; (ii) a  $t$ -type test of  $\mathcal{H}_0 : \vartheta_j = 0$  versus  $\mathcal{H}_1 : \vartheta_j \neq 0$ . These rejection frequencies are referred to as “size” and “power”, respectively, and are computed using the 0.975 standard-normal quantile as critical value.<sup>16</sup> Tests tend to have Type I error probabilities which are generally close to the nominal 0.05 level, especially for  $T > 200$ . Tests are also powerful enough to reject the hypothesis of a zero parameter

<sup>15</sup>Results for  $\rho = 0$  are not reported since they are very similar to those for  $\rho = 0.8$ .

<sup>16</sup>Results should be interpreted with caution in the case of  $\mathcal{H}_0 : \sigma_i = 0$ ,  $i \in \{0, 1\}$ , because the null value of  $\sigma_i$  is on the boundary of the maintained hypothesis. Our asymptotic theory does not allow for parameters that may lie on the boundary of the parameter space.

Table 1: Bias and Standard Deviation of ML Estimators ( $\rho = 0.8$ )

$T$	$\mu_0$	$\mu_1$	$\alpha_1$	$\beta_1$	$\alpha_0$	$\beta_0$	$\sigma_0$	$\sigma_1$	$\phi$
Bias									
200	-0.007	0.023	0.004	0.126	0.043	0.209	-0.011	-0.005	-0.005
800	-0.002	0.004	-0.002	0.023	0.002	0.026	-0.001	0.000	-0.001
1600	-0.001	0.002	-0.002	0.019	0.007	0.019	0.001	0.000	-0.001
3200	0.001	0.002	0.002	-0.005	0.002	0.011	0.001	0.000	0.000
Ratio of sampling standard deviation to estimated standard error									
200	1.085	1.016	1.084	1.078	1.132	1.178	1.066	1.043	1.072
800	0.983	0.991	0.980	1.024	1.044	1.018	1.012	1.010	1.045
1600	1.021	1.002	0.964	0.955	1.033	1.042	1.000	0.999	0.997
3200	1.026	0.979	1.010	1.017	0.990	0.955	1.012	1.020	1.020

Table 2: Size and Power of  $t$ -Type Tests ( $\rho = 0.8$ )

$T$	$\mu_0$	$\mu_1$	$\alpha_1$	$\beta_1$	$\alpha_0$	$\beta_0$	$\sigma_0$	$\sigma_1$	$\phi$
Size									
200	0.063	0.076	0.069	0.069	0.081	0.043	0.084	0.088	0.108
800	0.053	0.063	0.061	0.051	0.074	0.049	0.065	0.065	0.073
1600	0.065	0.061	0.047	0.043	0.060	0.048	0.052	0.047	0.068
3200	0.049	0.058	0.053	0.058	0.054	0.042	0.051	0.061	0.053
Power									
200	0.999	1.000	0.990	0.261	0.934	0.207	1.000	1.000	1.000
800	1.000	1.000	1.000	0.812	0.999	0.732	1.000	1.000	1.000
1600	1.000	1.000	1.000	0.988	1.000	0.966	1.000	1.000	1.000
3200	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000

value, except in the case of  $\beta_0$  and  $\beta_1$  with  $T = 200$ . The distributions of studentized statistics associated with the elements of the ML estimator of  $\vartheta$  (ratio of estimation error to corresponding estimated standard error) generally tend to have mean and variance (not shown) that do not differ substantially from zero and one, respectively, and Gaussianity is never rejected for  $T > 200$ .<sup>17</sup>

## 5.2 Misspecification

In the second set of experiments, we consider estimation of the parameter  $\vartheta = (\mu_0, \mu_1, \phi, \sigma_0, \sigma_1, \alpha_0, \beta_0, \alpha_1, \beta_1)$  using the partial likelihood function based on

<sup>17</sup>Statistics associated with  $\phi$ ,  $\sigma_0$  and  $\sigma_1$  appear to fare somewhat worse than others when  $T = 200$ , a finding similar to that reported in Psaradakis and Sola [1998] for models with a time-invariant transition mechanism.

the conditional distribution of  $Y_t$  given  $(Y_{t-1}, S_t)$ . Inference in models like (5)–(7) is predominantly based on such a partial likelihood that ignores the equation for  $Z_t$  (see, e.g., Diebold et al. [1994], Filardo [1994]). Formally, the misspecified model may be viewed as defined by equations (5)–(7), with the additional assumption that  $\rho = 0$ . Under this assumption, estimation of  $\vartheta$  is based on (5) alone, the (potentially incorrect) rationale behind this approach being that, since the conditional distribution of  $Y_t$  given  $(Y_{t-1}, S_t)$  and the transition probabilities of  $(S_t)_t$  depend only on  $Z_{t-1}$ , (5) may be analyzed, without loss of relevant information, independently of  $(Z_t)_t$ . Even though such an approach may be appealing because of its relative simplicity, it is far from obvious that it provides a valid way for conducting inference on  $\vartheta$ ; it is unclear, for example, what the limit point of the ML estimator based on the partial likelihood might be when  $\rho \neq 0$ . In earlier sections, we considered a theoretical framework that acknowledges this source of misspecification (among others) and provided tools for asymptotically valid inference. We now quantify the implications of this misspecification in finite samples. For brevity, we refer to the maximizer of the partial likelihood function associated with the conditional distribution of  $Y_t$  given  $(Y_{t-1}, S_t)$  as the ‘partial ML’ estimator to distinguish it from the ‘joint ML’ estimator based on the joint model for the conditional distribution of  $X_t$  given  $X_0^{t-1}$  (cf. Section 5.1).

Table 3 shows the estimated bias of the partial ML estimators of the elements of  $\vartheta$  and the ratio of the sampling standard deviation of the estimators to the estimated standard errors (averaged across replications) when  $\rho = 0.8$ . To reflect what is common practice in applied research, standard errors are computed using the Hessian estimator (which relies on the assumption of a correctly specified likelihood) instead of a “sandwich” estimator (which allows for misspecification). It is immediately apparent that the partial ML estimator of most of the parameters is considerably more biased than the joint ML estimator. The differences between the two estimators are more pronounced for parameters associated with the transition probabilities  $(\alpha_0, \beta_0, \alpha_1, \beta_1)$ , the partial ML estimators of which are significantly biased even for  $T = 3200$ . This suggests that the bias of the partial ML estimator when  $\rho \neq 0$  is not associated only with small samples, a finding that is consistent with our asymptotic results. Regarding the accuracy of estimated standard errors, the latter are downwards biased in most cases, the bias being somewhat larger than it is for joint ML estimators. However, unless  $T$  is small, this bias is not generally substantial and declines as  $T$  increases, despite the fact that standard errors are obtained from the Hessian.<sup>18</sup>

---

<sup>18</sup>Results for  $\rho = 0$  (not shown) are not substantially different from those obtained from the joint ML procedure. This is not surprising since the joint and partial ML estimators are both consistent for the true parameter value when  $\rho = 0$ .

Table 3: Bias and Standard Deviation of Partial ML Estimators ( $\rho = 0.8$ )

$T$	$\mu_0$	$\mu_1$	$\alpha_1$	$\beta_1$	$\alpha_0$	$\beta_0$	$\sigma_0$	$\sigma_1$	$\phi$
Bias									
200	0.019	0.046	-0.172	0.898	0.397	1.363	-0.056	-0.039	-0.001
800	0.017	0.017	-0.156	0.485	0.216	0.838	-0.025	-0.024	0.002
1600	0.008	0.011	-0.150	0.446	0.211	0.824	-0.018	-0.021	0.003
3200	0.012	0.008	-0.149	0.444	0.196	0.773	-0.020	-0.019	0.003
Ratio of sampling standard deviation to estimated standard error									
200	1.244	1.135	1.395	1.585	1.349	1.406	1.161	1.062	1.250
800	1.027	1.041	1.153	1.170	1.067	1.119	0.988	1.020	1.080
1600	0.991	1.014	1.070	1.075	1.072	1.161	1.006	0.959	1.054
3200	1.047	1.027	1.050	1.063	1.034	1.088	1.021	1.004	1.056

We note that hypothesis tests based on studentized statistics analogous to those considered in Section 5.1 (not shown) are unreliable when partial ML estimates are used. This is especially true in the case of parameters associated with the transition probabilities, the corresponding tests being either excessively conservative or excessively liberal. Although tests of this type are extensively used in applied work, they should be interpreted with caution since the statistics on which they are based have an asymptotically normal null distribution only when  $\rho = 0$ . We also note that using the “sandwich” estimator of Theorem 5(b) instead of the estimator based on the observed information matrix is not without difficulty when  $\rho \neq 0$ . As Freedman [2006] points out, the use of such an estimator for inference is unlikely to produce results that are any less misleading under misspecification since the problem of bias/inconsistency of the ML estimator for the true parameter value remains. It is indeed clear from the results in Table 3 that the bias of the partial ML estimator presents a much more serious problem in our setting than the inaccuracy of conventionally computed standard errors.

## 6 Empirical Illustration

In this section, we present an empirical illustration based on a regime-switching model of a type that is commonly used in economics.<sup>19</sup> Specifically, we investigate the potential contribution of the interest rate spread and the growth in tax revenues in predicting regime changes in U.S. real output growth. The model

<sup>19</sup>An additional empirical example, which examines the predictive ability of an index of leading indicators for regime changes in U.S. output growth, is discussed in the Supplemental Material SM.10.

is a variant of the specification used in the simulations and is given by

$$Y_t = \mu_0(1 - S_t) + \mu_1 S_t + \sum_{i=1}^{h_1} \phi_i Y_{t-i} + \sigma_1 U_{1,t}, \quad (8)$$

$$Z_t = \mu_2 + \sum_{i=1}^{h_2} \psi_i Z_{t-i} + \sigma_2 U_{2,t}, \quad (9)$$

for some  $h_1, h_2 \in \mathbb{N}$ , with the hidden, two-state Markov chain  $(S_t)_t$  being governed by the transition probabilities

$$Q_\theta(z, s, s) \equiv \Pr(S_t = s \mid Z_{t-1} = z, S_{t-1} = s) = [1 + \exp(-\alpha_s - \beta_s z)]^{-1}, \quad (10)$$

with  $s \in \{0, 1\}$ , and  $(U_{1,t}, U_{2,t})_t$  postulated to be i.i.d.  $\mathcal{N}\left(\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix}\right)$  and independent of  $(S_t)_t$ . In (8)–(10),  $Y_t$  stands for the growth rate of real gross domestic product and  $Z_t$  is either the spread between the 10-year Treasury note rate and the 3-month Treasury bill rate or the growth rate of real government receipts of direct and indirect taxes.<sup>20</sup> The data are quarterly and span the period 1954:3–2009:2.<sup>21</sup>

Since the aim is not only to assess the predictive ability of the interest rate spread and tax revenues for regime changes in output growth but also to examine whether treating these variables as exogenous yields results which are different from those obtained from a joint model, we compute two sets of estimates: partial ML estimates based on (8) alone and joint ML estimates based on the system (8)–(9). We note that in econometric models of the business cycle such as (8)–(10), it is common to rely on partial ML estimation (see, e.g., [Filardo \[1994\]](#)). Parameter estimates are reported in Tables 4 and 5, with estimated standard errors given in parentheses; the latter are obtained from the “sandwich” estimator of Theorem 5(b).<sup>22</sup> On the basis of  $t$ -type tests based on

<sup>20</sup>The model could be generalized to allow for Markov changes in all the parameters. However, since  $Z_t$  is thought of here as a potential leading indicator for business-cycle phases, it does not seem sensible to allow the parameters in both (8) and (9) to be subject to changes driven by  $(S_t)_t$ . Modeling regime changes in  $(Y_t)_t$  and  $(Z_t)_t$  as being driven by two independent Markov processes is more attractive, but we choose to abstract from this as it is not directly related to the main problem under study.

<sup>21</sup>Interest rate data are taken from the FRED database; output and tax data are taken from [Auerbach and Gorodnichenko \[2012\]](#). The likelihood ratio test of [Hansen \[1992\]](#) rejects the hypothesis that  $\mu_0 = \mu_1$  in (8).

<sup>22</sup>The weights  $\omega(\cdot, L_T)$  are obtained from the Parzen kernel and  $L_T$  is determined using the automatic plug-in method of [Andrews \[1991\]](#).

Table 4: ML Estimates (Output Growth, Interest Rate Spread)

Partial ML		Joint ML			
$\mu_0$	0.0091 (0.0012)	$\mu_0$	0.0092 (0.0012)		
$\mu_1$	0.0001 (0.0013)	$\mu_1$	0.0004 (0.0013)	$\mu_2$	0.2956 (0.0984)
$\phi_1$	0.1543 (0.0761)	$\phi_1$	0.1449 (0.0763)	$\psi_1$	0.8098 (0.0012)
$\phi_2$	0.0620 (0.0855)	$\phi_2$	0.0595 (0.0852)	$\psi_2$	-0.1065 (0.1251)
$\phi_3$	-0.0468 (0.0867)	$\phi_3$	-0.0435 (0.0759)	$\psi_3$	0.3368 (0.1747)
$\phi_4$	-0.0136 (0.0929)	$\phi_4$	-0.0183 (0.0914)	$\psi_4$	-0.2495 (0.0859)
$\alpha_0$	-1.3367 (3.8866)	$\alpha_0$	-1.5210 (2.6935)	$\sigma_2$	0.7053 (0.1091)
$\beta_0$	8.8363 (9.4778)	$\beta_0$	9.1169 (6.4769)	$\rho$	-0.0849 (0.1198)
$\alpha_1$	3.1927 (0.7646)	$\alpha_1$	3.1218 (0.7688)		
$\beta_1$	-1.0779 (0.4304)	$\beta_1$	-1.0338 (0.4293)		
$\sigma_1$	0.0077 (0.0006)	$\sigma_1$	0.0078 (0.0006)		

joint ML estimates, at least one of the parameters  $(\beta_0, \beta_1)$  is significantly different from zero, indicating that the spread and tax revenues contain significant information about the probability of switching between the two regimes.

Regarding the implications of treating  $Z_t$  as exogenous, the differences between partial and joint ML estimates are substantial in the model with tax revenues (especially for autoregressive coefficients and the parameters associated with the transition probabilities) but much less so in the model with the interest rate spread. This is not entirely unexpected in view of the fact that the estimated value of the conditional correlation  $\rho$  is relatively large (0.6034) in the former model but much smaller ( $-0.0849$ ), and insignificantly different from zero, in the latter. Such findings are in line with the analytical and simulation results presented in previous sections. The relatively large estimate of  $\rho$  in Table 5 also suggests that inference based on the partial ML estimator is potentially misleading because of the likely bias of the estimator.



Table 5: ML Estimates (Output Growth, Growth in Taxes)

Partial ML		Joint ML			
$\mu_0$	0.0071 (0.0014)	$\mu_0$	0.0081 (0.0014)		
$\mu_1$	-0.0119 (0.0034)	$\mu_1$	-0.0100 (0.0075)	$\mu_2$	0.5455 (0.2633)
$\phi_1$	0.2076 (0.0679)	$\phi_1$	0.0987 (0.0665)	$\psi_1$	0.1723 (0.1220)
$\phi_2$	0.0709 (0.0953)	$\phi_2$	0.0754 (0.0833)	$\sigma_2$	3.0458 (0.3082)
$\phi_3$	-0.0530 (0.0754)	$\phi_3$	-0.1168 (0.0610)	$\rho$	0.6034 (0.0673)
$\phi_4$	-0.0291 (0.0885)	$\phi_4$	-0.0345 (0.0721)		
$\alpha_0$	3.4588 (0.6379)	$\alpha_0$	3.8835 (1.3467)		
$\beta_0$	0.2754 (0.1237)	$\beta_0$	0.4061 (0.1379)		
$\alpha_1$	0.3852 (0.8704)	$\alpha_1$	-2.5349 (4.7906)		
$\beta_1$	0.2558 (0.1053)	$\beta_1$	0.0579 (0.1713)		
$\sigma_1$	0.0076 (0.0006)	$\sigma_1$	0.0084 (0.0009)		

## A Appendix: Proofs

### A.1 Consistency

In order to prove Theorem 1, we need two lemmas (the proofs of which are relegated to the Supplemental Material SM.2). The first lemma shows that the log-likelihood function  $\ell_T^\nu(X_0^T, \cdot)$  can be approximated by the sample average of  $(\log p^\nu(X_t | X_{-\infty}^{t-1}, \cdot))_{t \in \mathbb{N}}$ ; this function is used to construct the function  $H^*$  that defines the pseudo-true parameter set. The result relies on “mixing” properties established in Theorem 2 (see Lemma 11 in the Supplemental Material SM.2).

**Lemma 2.** *Suppose Assumptions 1, 4(ii) and 5 hold. Then,*

$$\sup_{\theta \in \Theta} \left| T^{-1} \sum_{t=1}^T (\log p_t^\nu(X_t | X_0^{t-1}, \theta) - \log p^\nu(X_t | X_{-\infty}^{t-1}, \theta)) \right| = o_{\bar{P}_*^\nu}(1).$$

The second lemma essentially establishes a uniform law of large numbers for the sample average of  $(\log p^\nu(X_t | X_{-\infty}^{t-1}, \cdot))_{t \in \mathbb{N}}$ .

**Lemma 3.** *Suppose Assumptions 1, 2, 3 and 4(i) hold. Then: (i) For any compact  $K \subseteq \Theta$  and any  $\epsilon > 0$ , there exists  $T(\epsilon) \in \mathbb{N}$  such that*

$$\bar{P}_*^\nu \left( \sup_{\theta \in K} T^{-1} \sum_{t=1}^T (\log p^\nu(X_t | X_{-\infty}^{t-1}, \theta) - E_{\bar{P}_*^\nu} [\log p^\nu(X_t | X_{-\infty}^{t-1}, \theta)]) > \epsilon \right) \leq \epsilon, \quad (11)$$

for all  $T \geq T(\epsilon)$ .

(ii) For any  $\theta_* \in \Theta_*$ ,

$$\left| T^{-1} \sum_{t=1}^T \left( \log p^\nu(X_t | X_{-\infty}^{t-1}, \theta_0) - E_{\bar{P}^\nu} [\log p^\nu(X_t | X_{-\infty}^{t-1}, \theta_0)] \right) \right| = o_{\bar{P}^\nu}(1).$$

*Proof of Theorem 1.* For simplicity, we set  $\eta_T = 0$  throughout the proof. Formally, we wish to establish that, for all  $\epsilon > 0$ , there exists  $T(\epsilon) \in \mathbb{N}$  such that

$$\bar{P}^\nu \left( d_\Theta(\hat{\theta}_{\nu,T}, \Theta_*) \geq \epsilon \right) < \epsilon,$$

for all  $t \geq T(\epsilon)$ . For this, it suffices to show that there exists a  $\theta_0 \in \Theta_*$  such that, for any  $\epsilon > 0$ , there exists a  $T(\theta_0, \epsilon)$  such that

$$\bar{P}^\nu \left( \sup_{\theta \in \Theta \setminus \Theta_*^\epsilon} \ell_T^\nu(X_0^T, \theta) \geq \ell_T^\nu(X_0^T, \theta_0) \right) < \epsilon,$$

for all  $T \geq T(\theta_0, \epsilon)$ , where  $\Theta_*^\epsilon = \{\theta \in \Theta : d_\Theta(\theta, \Theta_*) < \epsilon\}$ . Since, by Lemma 2,  $\ell_T^\nu(X_0^T, \cdot)$  is well approximated by  $\ell_T^\nu(X_{-\infty}^T, \cdot) \equiv T^{-1} \sum_{t=1}^T \log p^\nu(X_t | X_{-\infty}^{t-1}, \cdot)$ , it suffices to work with the latter function.

Let  $A_T(\delta) = \left\{ X_{-\infty}^\infty : \sup_{\theta \in \Theta \setminus \Theta_*^\epsilon} T^{-1} \sum_{t=1}^T \left( \log p^\nu(X_t | X_{-\infty}^{t-1}, \theta) - E_{\bar{P}^\nu} [\log p^\nu(X_t | X_{-\infty}^{t-1}, \theta)] \right) \leq \delta \right\}$  and  $B_T(\delta) = \left\{ X_{-\infty}^\infty : \left| T^{-1} \sum_{t=1}^T \left( -\log p^\nu(X_t | X_{-\infty}^{t-1}, \theta_0) + E_{\bar{P}^\nu} [\log p^\nu(X_t | X_{-\infty}^{t-1}, \theta_0)] \right) \right| \leq \delta \right\}$ , for any  $\delta > 0$  and any  $\theta_0 \in \Theta_*$ . Observe that

$$\begin{aligned} & \bar{P}^\nu \left( \sup_{\theta \in \Theta \setminus \Theta_*^\epsilon} \ell_T^\nu(X_{-\infty}^T, \theta) \geq \ell_T^\nu(X_{-\infty}^T, \theta_0) \right) \\ & \leq \bar{P}^\nu \left( \sup_{\theta \in \Theta \setminus \Theta_*^\epsilon} \ell_T^\nu(X_{-\infty}^T, \theta) \geq \ell_T^\nu(X_{-\infty}^T, \theta_0) \cap A_T(\delta) \cap B_T(\delta) \right) + \bar{P}^\nu(A_T(\delta)^c) + \bar{P}^\nu(B_T(\delta)^c) \\ & \leq \bar{P}^\nu \left( \sup_{\theta \in \Theta \setminus \Theta_*^\epsilon} T^{-1} \sum_{t=1}^T E_{\bar{P}^\nu} \left[ \log \frac{p^\nu(X_t | X_{-\infty}^{t-1}, \theta)}{p^\nu(X_t | X_{-\infty}^{t-1})} \right] \geq T^{-1} \sum_{t=1}^T E_{\bar{P}^\nu} \left[ \log \frac{p^\nu(X_t | X_{-\infty}^{t-1}, \theta_0)}{p^\nu(X_t | X_{-\infty}^{t-1})} \right] - 2\delta \right) \\ & \quad + \bar{P}^\nu(A_T(\delta)^c) + \bar{P}^\nu(B_T(\delta)^c) \\ & \leq \bar{P}^\nu \left( \inf_{\theta \in \Theta \setminus \Theta_*^\epsilon} H^*(\theta) \leq H^*(\theta_0) + 2\delta \right) + \bar{P}^\nu(A_T(\delta)^c) + \bar{P}^\nu(B_T(\delta)^c), \end{aligned}$$

where the last line follows from the stationarity of  $X_{-\infty}^\infty$  and the definition of  $H^*$ . By Assumption 3 and the fact that, for any  $\theta \in \Theta \setminus \Theta_*^\epsilon$ ,  $H^*(\theta) > H^*(\theta_0)$  (otherwise,  $\theta$  would belong to  $\Theta_*$ ), it follows that  $\inf_{\theta \in \Theta \setminus \Theta_*^\epsilon} H^*(\theta) - H^*(\theta_0) \equiv \Delta > 0$ . Hence, choosing  $\delta < 0.5\Delta$ , the first term of the right-hand side (RHS) vanishes. By Assumption 3(i),  $\Theta \setminus \Theta_*^\epsilon$  is compact; hence, by Lemma 3, there exists a  $T'$  (which may depend on  $\epsilon$  and  $\theta_0$ ) such that  $\bar{P}^\nu(A_T(\delta)^c) + \bar{P}^\nu(B_T(\delta)^c) \leq \epsilon$  for any  $\delta \leq 0.5\epsilon$  and all  $T \geq T'$ , and thus the desired result follows.  $\square$

## A.2 Mixing Results

Throughout, fix  $m$  and  $j$  as in the statement of Theorem 2. For any  $n, n'$  such that  $-m \leq n, n' \leq j+1$ , we denote the Dobrushin coefficient of  $\bar{P}_\theta^\nu(S_{n'} = \cdot \mid S_n = \cdot, X_{-m}^j)$  as

$$\alpha_{\theta, n', n}(X_{-m}^j) \equiv \frac{1}{2} \max_{(a, b) \in \mathbb{S}^2} \left\| \bar{P}_\theta^\nu(S_{n'} = \cdot \mid S_n = a, X_{-m}^j) - \bar{P}_\theta^\nu(S_{n'} = \cdot \mid S_n = b, X_{-m}^j) \right\|_1. \quad (12)$$

Since  $\alpha_{\theta, j+1, -m}(X_{-m}^j) \leq \prod_{n=-m}^j \alpha_{\theta, n+1, n}(X_{-m}^j)$  (e.g., Dobrushin [1956], Sethuraman and Varadhan [2005]), to prove Theorem 2, it suffices to show the following.

**Lemma 4.** *For any  $l \in \{-m, \dots, j\}$  and any  $\theta \in \Theta$ ,  $\alpha_{\theta, l+1, l}(X_{-m}^j) \leq 1 - \underline{q}(X_l)$  a.s.- $\bar{P}_*^\nu$ .*

Lemma 4 follows immediately from Lemmas 5 and 6 below. To state these lemmas, we construct the following processes that will be used for coupling. For any  $i \in \{1, 2\}$  and any  $\theta \in \Theta$ , let  $(X_{i,t}, \eta_{i,t}, v_{i,t})_{t=-m}^\infty$ , with  $(X_{i,t}, \eta_{i,t}, v_{i,t}) \in \mathbb{X} \times \mathbb{S} \times \{0, 1\}$ , be defined as follows:  $(X_{i,-m}, \eta_{i,-m}) \sim \nu$ ; given  $(X_{i,t})_{t=-m}^\infty, (v_{i,t})_{t=-m}^\infty$  is i.i.d. with  $\Pr(v_{i,t} = 1 \mid X_{i,-m}^\infty) = \Pr(v_{i,t} = 1 \mid X_{i,-m}^t) \equiv \underline{q}(X_{i,t})$ ; for each  $t \geq -m$ ,  $\eta_{i,t+1} \sim \varrho(X_{i,t}, \cdot)$  if  $v_{i,t} = 1$ , and  $\eta_{i,t+1} \sim \frac{Q_\theta(X_{i,t}, \eta_{i,t}, \cdot) - \underline{q}(X_{i,t})\varrho(X_{i,t}, \cdot)}{1 - \underline{q}(X_{i,t})}$  if  $v_{i,t} = 0$  (the last quotient expression is a valid transition kernel under condition (3) in Theorem 2); finally,  $X_{i,t+1} \sim p_\theta(X_{i,t}, \eta_{i,t+1}, \cdot)$ .

This construction implies that the transition kernel of  $(\eta_{i,t})_t$  is given by

$$\begin{aligned} \Pr_\theta(\eta_{i,t+1} = \cdot \mid \eta_{i,t}, X_{i,t}) &= \underline{q}(X_{i,t})\varrho(X_{i,t}, \cdot) + (1 - \underline{q}(X_{i,t})) \frac{Q_\theta(X_{i,t}, \eta_{i,t}, \cdot) - \underline{q}(X_{i,t})\varrho(X_{i,t}, \cdot)}{1 - \underline{q}(X_{i,t})} \\ &= Q_\theta(X_{i,t}, \eta_{i,t}, \cdot), \end{aligned}$$

and since the transition for  $X_{i,t+1}$  given  $(X_{i,t}, \eta_{i,t+1})$  is governed by  $p_\theta$ , the following result holds (its proof is relegated to the Supplemental Material SM.7).

**Lemma 5.** *For any  $l \in \{-m, \dots, j\}$  and any  $\theta \in \Theta$ ,*

$$\bar{P}_\theta^\nu(S_{l+1} = \cdot \mid S_l = \cdot, X_{-m}^j) = \Pr_\theta(\eta_{i,l+1} = \cdot \mid \eta_{i,l} = \cdot, X_{-m}^j), \quad \forall i \in \{1, 2\},$$

a.s.- $\bar{P}_*^\nu$ .

Furthermore, since  $\eta_{i,t+1}$  becomes independent of its own past whenever  $v_{i,t} = 1$ , the following result can be established (its proof is relegated to the Supplemental Material SM.7).

**Lemma 6.** For any  $l \in \{-m, \dots, j\}$  and any  $\theta \in \Theta$ ,

$$\frac{1}{2} \max_{(a,b) \in \mathbb{S}^2} \left\| \Pr_{\theta}(\eta_{1,l+1} = \cdot \mid \eta_{1,l} = a, X_{-m}^j) - \Pr_{\theta}(\eta_{2,l+1} = \cdot \mid \eta_{2,l} = b, X_{-m}^j) \right\|_1 \leq 1 - \underline{q}(X_l),$$

*a.s.*- $\bar{P}_*^{\nu}$ .

It is easy to see that Lemma 4 (and thus Theorem 2) follows from the last two lemmas.

### A.3 Asymptotic Distribution Theory

The next two lemmas are used to prove Theorems 3 and 4 (their proofs are relegated to the Supplemental Material SM.8.1). In what follows, for vector/matrix-valued functions  $X \mapsto f(X)$ ,  $\|f\|_{L^r(P)}$  is short-hand notation for the  $L^r(P)$ -norm of  $x \mapsto \|f(x)\|$ , where  $\|\cdot\|$  denotes the Euclidean/dual norm of  $f$ .

**Lemma 7.** Suppose Assumptions 1, 2, 6, 7(i) and 8 hold. Then, there exists a stationary and ergodic (under  $\bar{P}_*^{\nu}$ ) process  $(\Delta_t(\theta_*))_{t=-\infty}^{\infty}$  in  $L^2(\bar{P}_*^{\nu})$  such that

$$\lim_{T \rightarrow \infty} \left\| T^{-1/2} \sum_{t=0}^T \{ \nabla_{\theta} \log p_t^{\nu}(\cdot \mid \cdot, \theta_*) - \Delta_t(\theta_*) \} \right\|_{L^2(\bar{P}_*^{\nu})} = 0.$$

**Lemma 8.** Suppose Assumptions 1, 2, 6, 7 and 8 hold. Then, there exists a sequence of  $\mathbb{R}^{q \times q}$ -valued continuous functions  $(\theta \mapsto \xi_t(\theta))_t$  such that  $\xi_t(\theta)$  is negative definite for all  $t$  and

$$\lim_{T \rightarrow \infty} \left\| \sup_{\theta \in B(\delta, \theta_*)} \left\| T^{-1} \sum_{t=0}^T \{ \nabla_{\theta}^2 \log p_t^{\nu}(\cdot \mid \cdot, \theta) - \xi_t(\theta) \} \right\| \right\|_{L^1(\bar{P}_*^{\nu})} = 0,$$

where  $\delta > 0$  is the same as in Assumption 7.

*Proof of Theorem 3.* Choose  $K$  compact such that, for any  $v \in K$ ,  $\|v\| \leq \delta$  for  $\delta > 0$  as in Lemma 8. For any  $v \in K$ , by Assumption 6,

$$\ell_T^{\nu}(X_0^T, \theta_* + v) - \ell_T^{\nu}(X_0^T, \theta_*) = v^{\top} \nabla_{\theta} \ell_T^{\nu}(X_0^T, \theta_*) + 0.5 v^{\top} \left( \int_0^1 \nabla_{\theta}^2 \ell_T^{\nu}(X_0^T, \theta_* + wv) dw \right) v.$$

By Lemmas 7 and 8, and the fact that  $(\theta_* + wv) \in B(v, \theta_*)$ ,

$$\begin{aligned} \ell_T^{\nu}(X_0^T, \theta_* + v) - \ell_T^{\nu}(X_0^T, \theta_*) &= v^{\top} \left( T^{-1} \sum_{t=0}^T \Delta_t(\theta_*) + o_{\bar{P}_*^{\nu}}(T^{-1/2}) \right) \\ &\quad + 0.5 v^{\top} \left( \int_0^1 T^{-1} \sum_{t=0}^T \xi_t(\theta_* + wv) dw + o_{\bar{P}_*^{\nu}}(1) \right) v. \end{aligned}$$

Now, let  $R_T(v) \equiv v^\top \left( T^{-1} \sum_{t=0}^T \int_0^1 \{ \xi_t(\theta_* + wv) - \xi_t(\theta_*) \} dw \right) v$ . Observe that  $\|v\|^{-2} |R_T(v)| \leq \int_0^1 \left\| T^{-1} \sum_{t=0}^T \{ \xi_t(\theta_* + wv) - \xi_t(\theta_*) \} \right\| dw$ , so, for any  $\delta > 0$ ,

$$\begin{aligned} \bar{P}_*^\nu \left( \sup_{v \in B(\delta, 0)} \frac{|R_T(v)|}{\|v\|^2} \geq \epsilon \right) &\leq \bar{P}_*^\nu \left( \sup_{v \in B(\delta, 0)} \int_0^1 \left\| T^{-1} \sum_{t=0}^T \{ \xi_t(\theta_* + wv) - \xi_t(\theta_*) \} \right\| dw \geq \epsilon \right) \\ &\leq \epsilon^{-1} E_{\bar{P}_*^\nu} \left[ \sup_{v \in B(\delta, 0)} \int_0^1 \| \xi_1(\theta_* + wv) - \xi_1(\theta_*) \| dw \right], \end{aligned}$$

where the second line follows by the Markov inequality and stationarity. The desired result then follows by the continuity of  $\xi_1$  (see Lemma 8) and the same arguments as in [Bickel et al., 1998, p. 1634].  $\square$

*Proof of Theorem 4.* Henceforth, let  $\bar{\Delta}_T \equiv T^{-1} \sum_{t=0}^T \Delta_t(\theta_*) + o_{\bar{P}_*^\nu}(T^{-1/2})$ . By Theorem 1,  $\hat{\theta}_{\nu, T} - \theta_*$  converges to zero with probability approaching one (w.p.a.1). Thus,  $R_T(v_T) = o_{\bar{P}_*^\nu}(\|v_T\|^2)$  and, by Theorem 3,

$$\begin{aligned} \ell_T^\nu(X_0^T, \hat{\theta}_{\nu, T}) - \ell_T^\nu(X_0^T, \theta_*) &= (\hat{\theta}_{\nu, T} - \theta_*)^\top \bar{\Delta}_T \\ &\quad + 0.5(\hat{\theta}_{\nu, T} - \theta_*)^\top \left( T^{-1} \sum_{t=0}^T \xi_t(\theta_*) + o_{\bar{P}_*^\nu}(1) \right) (\hat{\theta}_{\nu, T} - \theta_*). \end{aligned}$$

Ergodicity of  $X_\infty^\infty$  (Lemma 1) implies ergodicity of  $(\xi_t(\theta_*))_{t=-\infty}^\infty$ ; therefore, by Lemma 8 and Birkhoff's ergodic theorem,

$$\begin{aligned} \ell_T^\nu(X_0^T, \hat{\theta}_{\nu, T}) - \ell_T^\nu(X_0^T, \theta_*) &= (\hat{\theta}_{\nu, T} - \theta_*)^\top \bar{\Delta}_T \\ &\quad + 0.5(\hat{\theta}_{\nu, T} - \theta_*)^\top \left( E_{\bar{P}_*^\nu}[\xi_1(\theta_*)] + o_{\bar{P}_*^\nu}(1) \right) (\hat{\theta}_{\nu, T} - \theta_*), \end{aligned} \tag{13}$$

and  $E_{\bar{P}_*^\nu}[\xi_1(\theta_*)]$  is non-singular. The rest of the proof proceeds in two steps.

STEP 1. Let  $r_T \equiv \min\{o_{\bar{P}_*^\nu}(1), o(T^{-1}) + E_{\bar{P}_*^\nu}[(\bar{\Delta}_T)^\top(\bar{\Delta}_T)]\}$ . We first establish that  $\|\hat{\theta}_{\nu, T} - \theta_*\| = O_{\bar{P}_*^\nu}(\sqrt{r_T})$ ; by Theorem 1, the  $o_{\bar{P}_*^\nu}(1)$  part of  $r_T$  has been established.

By (13) and the fact that  $\hat{\theta}_{\nu, T}$  is an (approximate) maximizer of the likelihood function,

$$-2\eta_T \leq 2(\hat{\theta}_{\nu, T} - \theta_*)^\top \bar{\Delta}_T - (\hat{\theta}_{\nu, T} - \theta_*)^\top A(\theta_*)(\hat{\theta}_{\nu, T} - \theta_*),$$

with  $A(\theta_*) \equiv \left( -E_{\bar{P}_*^\nu}[\xi_1(\theta_*)] + o_{\bar{P}_*^\nu}(1) \right)$ . Simple algebra yields

$$-2\eta_T \leq - \left\| (\hat{\theta}_{\nu, T} - \theta_*)^\top A(\theta_*)^{1/2} - \bar{\Delta}_T^\top A(\theta_*)^{-1/2} \right\|^2 + \bar{\Delta}_T^\top A(\theta_*)^{-1} \bar{\Delta}_T.$$

Moreover, by simple algebra and the Markov inequality,

$$\left\| (\hat{\theta}_{\nu,T} - \theta_*)^\top A(\theta_*)^{1/2} \right\| = O_{\bar{P}_*^\nu} \left( \sqrt{\eta_T} + \sqrt{E_{\bar{P}_*^\nu} \left[ (\bar{\Delta}_T)^\top (E_{\bar{P}_*^\nu}[\xi_1(\theta_*)])^{-1} (\bar{\Delta}_T) \right]} \right).$$

This expression, the fact that  $E_{\bar{P}_*^\nu}[\xi_1(\theta_*)]$  is non-singular, and  $\eta_T = o(T^{-1})$ , imply the desired result.

STEP 2. We now show that, for any  $\epsilon > 0$ ,

$$\bar{P}_*^\nu \left( r_T^{-1/2} \left\| (\hat{\theta}_{\nu,T} - \theta_*) - (-E_{\bar{P}_*^\nu}[\xi_1(\theta_*)] + o_{\bar{P}_*^\nu}(1))^{-1} \bar{\Delta}_T \right\| \geq \epsilon \right) \rightarrow 0.$$

Since, by Step 1,  $\left\| \hat{\theta}_{\nu,T} - \theta_* \right\| = O_{\bar{P}_*^\nu}(\sqrt{r_T})$ , it suffices to show that

$$\bar{P}_*^\nu \left( \left\{ r_T^{-1/2} \left\| (\hat{\theta}_{\nu,T} - \theta_*) - (-E_{\bar{P}_*^\nu}[\xi_1(\theta_*)] + o_{\bar{P}_*^\nu}(1))^{-1} \bar{\Delta}_T \right\| \geq \epsilon \right\} \cap \left\{ \left\| \hat{\theta}_{\nu,T} - \theta_* \right\| \leq \sqrt{r_T} M \right\} \right) \rightarrow 0, \quad (14)$$

where  $M > 0$ . To this end, note that, by Theorem 3 and the fact that  $T^{-1} \sum_{t=1}^T \xi_t(\theta_*) = E_{\bar{P}_*^\nu}[\xi_1(\theta_*)] + o_{\bar{P}_*^\nu}(1)$ , it follows that

$$\ell_T^\nu(X_0^T, \theta_* + v) - \ell_T^\nu(X_0^T, \theta_*) = (\bar{\Delta}_T)^\top v - 0.5v^\top (-E_{\bar{P}_*^\nu}[\xi_1(\theta_*)] + o_{\bar{P}_*^\nu}(1))v + R_T(v),$$

for any  $v \in K$ . Letting  $\Lambda_T(v) \equiv \ell_T^\nu(X_0^T, \theta_* + v) - \ell_T^\nu(X_0^T, \theta_*)$  and  $Q_T(v) \equiv (\bar{\Delta}_T)^\top v - 0.5v^\top (-E_{\bar{P}_*^\nu}[\xi_1(\theta_*)] + o_{\bar{P}_*^\nu}(1))v$ , we show that  $\sup_{v \in \{v: \|v\| \leq \sqrt{r_T} M\}} r_T^{-1} |\Lambda_T(v) - Q_T(v)| = o_{\bar{P}_*^\nu}(1)$ . To do so, it suffices to prove that  $\sup_{v \in \{v: \|v\| \leq \sqrt{r_T} M\}} |R_T(v)| = o_{\bar{P}_*^\nu}(r_T)$ . But this follows from Theorem 3 and the fact that  $\sqrt{r_T} = o_{\bar{P}_*^\nu}(1)$ . Since  $(\hat{\theta}_{\nu,T} - \theta_*) \in \{v: \|v\| \leq \sqrt{r_T} M\}$  and maximizes  $\Lambda_T(\cdot)$  (within a  $\eta_T$  margin), the previous result implies that

$$\begin{aligned} \hat{\theta}_{\nu,T} - \theta_* &= \arg \max_{v \in \{v: \|v\| \leq T^{-1/2} M\}} Q_T(v) + o_{\bar{P}_*^\nu}(\sqrt{r_T}) + \eta_T \\ &= (-E_{\bar{P}_*^\nu}[\xi_1(\theta_*)] + o_{\bar{P}_*^\nu}(1))^{-1} \bar{\Delta}_T + o_{\bar{P}_*^\nu}(\sqrt{r_T}), \end{aligned}$$

and thus (14) follows.  $\square$

The proof of Corollary 1 uses the following lemma (whose proof is relegated to the Supplemental Material SM.8.2).

**Lemma 9.** *Suppose there exists  $\bar{L} \in \mathbb{N}$  such that  $\nabla_\theta \log p_t^\nu(X_t | X_0^{t-1}, \theta) = \nabla_\theta \log p_t^\nu(X_t | X_{t-\bar{L}}^{t-1}, \theta)$  for all  $t \geq 0$ . Then:*

(a) *For all  $t \geq 0$ ,*

$$\Delta_t(\theta_*) \equiv \Delta_{t,-\infty}(\theta_*)(X_{-\infty}^t) = \nabla_\theta \log p_t^\nu(X_t | X_{t-\bar{L}}^{t-1}, \theta_*).$$

(b) If, in addition, there exists  $\delta > 0$  such that  $E_{\bar{P}_*^\nu} [|\nabla_\theta \log p_1^\nu(X_1 | X_{1-\bar{L}}^0, \theta_*)|^{4+4\delta}] < \infty$ , then, for any  $L \in \mathbb{N}$ ,

$$\max_{j \in \{0, \dots, L\}} \|T^{-1} \sum_{t=1}^T \Delta_{t+j, -\infty}(\theta_*) \Delta_{t, -\infty}(\theta_*)^\top - E_{\bar{P}_*^\nu} [\Delta_{j, -\infty}(\theta_*) \Delta_{0, -\infty}(\theta_*)^\top]\| = O_{\bar{P}_*^\nu} \left( \frac{L}{\sqrt{T}} \right).$$

*Proof of Corollary 1.* Throughout the proof, we use  $C$  to denote a universal constant that can take different values in different places. Also, for any  $k, T \geq 0$  and any  $\theta \in \Theta$ , we write  $\Delta_{k, k-T}(\theta) \equiv \Delta_{k, k-T}(\theta)(X_{k-T}^k) = \nabla_\theta \log p_k^\nu(X_k | X_{k-T}^{k-1}; \theta)$  (for the last equality, see Lemma 17 in the Supplemental Material SM.8.1) and  $\Delta_k(\theta) \equiv \Delta_{k, -\infty}(\theta)$ .

We first show that, under the conditions of part (a),  $(\Delta_t(\theta_*))_t$  is a martingale difference sequence (MDS) with respect to the natural filtration of  $(X_t)_t$ . To establish this, observe that

$$\begin{aligned} E_{\bar{P}_*^\nu} [\Delta_{k, k-T}(\theta_*) | X_{-\infty}^{k-1}] &= E_{\bar{P}_*^\nu} [\Delta_{k, k-T}(\theta_*) | X_{k-T}^{k-1}] \\ &= \int \frac{\nabla_\theta p_k^\nu(x_k | X_{k-T}^{k-1}; \theta_*)}{p_k^\nu(x_k | X_{k-T}^{k-1}; \theta_*)} p_k^\nu(x_k | X_{k-T}^{k-1}) dx_k. \end{aligned}$$

By assumption,  $p_k^\nu(x_k | X_{k-T}^{k-1}; \theta_*) = p_k^*(x_k | X_{k-T}^{k-1})$ . Moreover, by using the representation of  $\Delta_{k, k-T}(\cdot)$  in the Supplemental Material SM.8.1 and Assumption 7, it can be shown that  $\nabla_\theta \log p_k^\nu(\cdot | X_{k-T}^{k-1}; \theta)$  is uniformly bounded for all  $\theta \in B(\delta, \theta_*)$  (where  $\delta > 0$  is as in Assumption 7). This result allow us to use standard ‘‘Fisher information equality’’ calculations and thus deduce that  $E_{\bar{P}_*^\nu} [\Delta_{k, k-T}(\theta_*) | X_{-\infty}^{k-1}] = 0$ .

Next, we show that, for any  $k \geq 0$ ,  $A_k \equiv E_{\bar{P}_*^\nu} [\Delta_{k, -\infty}(\theta_*) | X_{-\infty}^{k-1}] = 0$ . To do so, observe that, for any  $k, T \geq 0$ ,

$$\begin{aligned} \|A_k\|_{L^2(\bar{P}_*^\nu)}^2 &= E_{\bar{P}_*^\nu} \left[ \left( E_{\bar{P}_*^\nu} [\Delta_{k, -\infty}(\theta_*) | X_{-\infty}^{k-1}] - E_{\bar{P}_*^\nu} [\Delta_{k, k-T}(\theta_*) | X_{-\infty}^{k-1}] \right)^2 \right] \\ &\leq \|\Delta_{k, -\infty}(\theta_*) - \Delta_{k, k-T}(\theta_*)\|_{L^2(\bar{P}_*^\nu)}^2 \\ &= \|\Delta_{0, -\infty}(\theta_*) - \Delta_{0, -T}(\theta_*)\|_{L^2(\bar{P}_*^\nu)}^2 \\ &\leq C \left( \max \left\{ \sum_{j=[-T/2]}^{-1} \varrho(j, -T), \sum_{j=-T}^{[-T/2]-1} \varrho(-1, j) \right\} \right), \end{aligned}$$

where the second line follows from the Jensen inequality, the third line follows from stationarity, and the fourth line follows from Lemma 18(i). Now, recall

that, for any  $j \geq k$ ,  $\varrho(j, k) \equiv \left( E_{\bar{P}_*^\nu} \left[ \prod_{i=k}^j (1 - \underline{q}(X_i))^{\frac{2a}{1-a}} \right] \right)^{\frac{1-a}{2a}} = \left( E_{\bar{P}_*^\nu} \left[ \prod_{i=0}^{j-k} (1 - \underline{q}(X_i))^{\frac{2a}{1-a}} \right] \right)^{\frac{1-a}{2a}} =$

$\varrho(j-k, 0)$ . Moreover, by Assumption 8,  $(\varrho(j, 0))_j$  is  $p$ -summable with  $p < 2/3$ , and thus  $\lim_{j \rightarrow \infty} \varrho(j, 0)^p j = 0$  (if not, then  $\varrho(j, 0) > c/j^{1/p}$  for some  $c > 0$  and all  $j$  above certain point, and this violates the assumption). Hence, for sufficiently large  $T$ ,

$$\begin{aligned} \|A_k\|_{L^2(\bar{P}_*)}^2 &\leq C \left( \max \left\{ \sum_{j=-T/2}^{-1} \varrho(j+T, 0), \sum_{j=-T}^{[-T/2]-1} \varrho(-1-j, 0) \right\} \right) \\ &= C \left( \max \left\{ \sum_{l=[T/2]}^{T-1} \varrho(l, 0), \sum_{l=[T/2]}^{T-1} \varrho(l, 0) \right\} \right) \\ &\leq C \sum_{l=[T/2]}^{T-1} l^{-(1/p)}. \end{aligned}$$

As  $1/p > 1$ , it follows that, as  $T$  diverges, the RHS converges to zero, and thus  $\|A_k\|_{L^2(\bar{P}_*)}^2 = 0$ . Since  $\Delta_k(\theta_*) = \Delta_{k,-\infty}(\theta_*)$  for any  $k$ , the desired result follows.

By Theorem 4 and the central limit theorem for MDS, we have, therefore,

$$\sqrt{T}(\hat{\theta}_{\nu,T} - \theta_*) \Rightarrow_{\bar{P}_*} \mathcal{N}(0, (E_{\bar{P}_*}[\xi_1(\theta_*)])^{-1} \Sigma(\theta_*) (E_{\bar{P}_*}[\xi_1(\theta_*)])^{-1}),$$

where  $\Sigma(\theta_*) \equiv \lim_{T \rightarrow \infty} \Sigma_T(\theta_*)$ , with  $\Sigma(\theta_*) = E_{\bar{P}_*}[(\Delta_1(\theta_*))(\Delta_1(\theta_*))^\top]$  since  $(\Delta_t(\theta_*))_t$  is a stationary MDS. Moreover, it can be shown that  $E_{\bar{P}_*}[\xi_1(\theta_*)] = -E_{\bar{P}_*}[(\Delta_1(\theta_*))(\Delta_1(\theta_*))^\top]$ . This follows by standard ‘‘Fisher information equality’’ calculations and derivations analogous to those above (so they are omitted). Hence, the proof of part (a) is complete once we show that  $-H_T^{-1}(\hat{\theta}_{\nu,T})$  converges in  $\bar{P}_*$ -probability to  $(E_{\bar{P}_*}[\xi_1(\theta_*)])^{-1} \Sigma(\theta_*) (E_{\bar{P}_*}[\xi_1(\theta_*)])^{-1}$ . But this follows from Theorem 5(a).

Under the conditions of part (b), and in view of Lemma 9(a),  $\Delta_k(\theta_*) = \nabla_{\theta} \log p_k^\nu(X_k | X_{k-L}^{k-1}; \theta_*)$  for all  $k \geq 0$ . This result and the fact that  $(X_k)_{k=-\infty}^\infty$  is  $\beta$ -mixing with mixing coefficients  $\beta_n = O(\gamma^n)$  (see Lemma 1) imply that  $(\Delta_k(\theta_*))_{k=-\infty}^\infty$  is also  $\beta$ -mixing with mixing coefficients of the same order. Hence, to establish that  $T^{-1/2} \sum_{t=0}^T \Delta_t(\theta_*) \Rightarrow_{\bar{P}_*} \mathcal{N}(0, \Sigma(\theta_*))$ , it is enough to verify that, for some  $\delta > 0$ ,  $E_{\bar{P}_*}[|\Delta_1(\theta_*)|^{2+\delta}] < \infty$  and  $\sum_{n=1}^\infty \alpha_n^{\delta/(2+\delta)} < \infty$ , where  $(\alpha_n)_n$  are the  $\alpha$ -mixing coefficients of  $(\Delta_k(\theta_*))_{k=-\infty}^\infty$  (see, e.g., [Doukhan et al. \[1994\]](#)). But, the summability condition on the  $\alpha$ -mixing coefficients is satisfied, because the  $\beta$ -mixing coefficients of the process decay at rate  $O(\gamma^n)$ , and the moment condition is directly assumed.

The proof of part (b) is completed by showing that  $\hat{\Omega}_T(\hat{\theta}_{\nu,T})$  converges in  $\bar{P}_*$ -probability to  $(E_{\bar{P}_*}[\xi_1(\theta_*)])^{-1} \Sigma(\theta_*) (E_{\bar{P}_*}[\xi_1(\theta_*)])^{-1}$ , which is non-singular by assumption. We do so by invoking Theorem 5(b) and verifying its conditions.



As stated above,  $\alpha_n = O(\gamma^n)$ . Moreover, by Corollary 6.17 in White [2001], there exists  $C < \infty$  such that, for any  $l > 0$ ,

$$\|E_{\bar{P}_*^\nu}[\Delta_l(\theta_*)\Delta_0(\theta_*)^\top]\| \leq C(\alpha_l)^{\frac{2}{2+2\delta}} \sqrt{E_{\bar{P}_*^\nu}[\|\Delta_l(\theta_*)\|^2]}(E_{\bar{P}_*^\nu}[\|\Delta_l(\theta_*)\|^{2+2\delta}])^{\frac{1}{2+2\delta}}.$$

Under our assumptions, the RHS equals  $C(\alpha_l)^{\frac{2}{2+2\delta}}$  for some  $C < \infty$ . Hence, one can set  $\bar{v}(l) \equiv C(\gamma^{\frac{2}{2+2\delta}})^l$ . Since  $\gamma < 1$ , the function  $l \mapsto \bar{v}(l)$  is integrable. Also, by Lemma 9(b), it follows that  $r_T = L_T T^{-1/2}$ .

Next, we show that  $\ddot{\omega}(\delta') = C\delta'$  for some  $C < \infty$  and any  $\delta' \leq \delta$ , where  $\delta > 0$  is as in Assumption 7. To do so, we note that for any  $\theta$  and any  $t \in \mathbb{N}$ ,

$$\begin{aligned} \|\Delta_t(\theta)\Delta_0(\theta)^\top - \Delta_t(\theta_*)\Delta_0(\theta_*)^\top\| &\leq \|\Delta_0(\theta)\| \times \|\Delta_t(\theta) - \Delta_t(\theta_*)\| \\ &\quad + \|\Delta_t(\theta_*)\| \times \|\Delta_0(\theta) - \Delta_0(\theta_*)\|. \end{aligned}$$

By the calculations in the proof of Lemma 18 in the Supplemental Material SM.8,  $\|\Delta_t(\theta)\|$  is a linear combination of terms involving  $\Gamma(\cdot, \theta)$  and  $\Lambda(\cdot, \theta)$ . Since  $\Delta_t(\theta)$  only depends on  $X_{t-\bar{L}}^t$  for some finite  $\bar{L}$ , there are only finitely many terms in this linear combination. Moreover, by Assumption 7, there exists  $C < \infty$  such that  $E_{\bar{P}_*^\nu}[\sup_{\theta \in B(\delta, \theta_*)} \|\Delta_t(\theta)\|^2] \leq C$ , where  $\delta > 0$  is as in Assumption 7. This bound is also uniform over  $t$ . Hence, on account of this result and stationarity, there exists  $C < \infty$  such that

$$E_{\bar{P}_*^\nu} \left[ \sup_{\|\theta - \theta_*\| \leq \delta'} \|\Delta_t(\theta)\Delta_0(\theta)^\top - \Delta_t(\theta_*)\Delta_0(\theta_*)^\top\| \right] \leq C \left( E_{\bar{P}_*^\nu} \left[ \sup_{\|\theta - \theta_*\| \leq \delta'} \|\Delta_0(\theta) - \Delta_0(\theta_*)\|^2 \right] \right)^{1/2},$$

for any  $\delta' \leq \delta$  and any  $t \in \mathbb{N}$ . It remains to show that the RHS is of the form  $C\delta'$  for some finite constant  $C$ . By the mean value theorem,  $\|\Delta_0(\theta) - \Delta_0(\theta_*)\| \leq \sup_{\theta \in B(\delta, \theta_*)} \|\nabla_\theta \Delta_0(\theta)\| \times \|\theta - \theta_*\|$ , so it suffices to show that  $E_{\bar{P}_*^\nu}[\sup_{\theta \in B(\delta, \theta_*)} \|\nabla_\theta \Delta_0(\theta)\|^2] \leq C$  for some  $C < \infty$ . Since  $\Delta_0(\theta) = \nabla_\theta \log p_0^\nu(X_0 | X_{-\bar{L}}^{-1})$ , this condition is equivalent to  $E_{\bar{P}_*^\nu}[\sup_{\theta \in B(\delta, \theta_*)} \|\nabla_\theta^2 \log p_0^\nu(X_0 | X_{-\bar{L}}^{-1}, \theta)\|^2] \leq C$ . By the calculations in [Bickel et al., 1998, pp. 1627–1628],  $\nabla_\theta^2 \log p_0^\nu(X_0 | X_{-\bar{L}}^0, \theta)$  is a linear combination of finitely many terms involving  $\Gamma(\cdot, \theta)$ ,  $\Lambda(\cdot, \theta)$ , and their derivatives. By Assumption 7 and calculations analogous to those in the proof of Lemma 18, it then follows that  $E_{\bar{P}_*^\nu}[\sup_{\theta \in B(\delta, \theta_*)} \|\nabla_\theta^2 \log p_0^\nu(X_0 | X_{-\bar{L}}^{-1}, \theta)\|^2] \leq C$ . Thus, there exists  $C < \infty$  such that  $\ddot{\omega}(\delta') \leq C\delta'$  for any  $\delta' \leq \delta$ . Therefore, all the conditions of Theorem 5(b) are satisfied.  $\square$

## References

- P. Ailliot and F. Pène. Consistency of the maximum likelihood estimate for non-homogeneous Markov-switching models. *ESAIM: Probability and Statistics*, 19:268–292, 2015.
- J. H. Albert and S. Chib. Bayes inference via gibbs sampling of autoregressive time series subject to Markov mean and variance shifts. *Journal of Business and Economic Statistics*, 11:1–15, 1993.
- D. W. K. Andrews. Heteroskedasticity and autocorrelation consistent covariance matrix estimation. *Econometrica*, 59:817–858, 1991.
- A. Ang and G. Bekaert. International asset allocation with regime shifts. *Review of Financial Studies*, 15:1137–1187, 2002a.
- A. Ang and G. Bekaert. Regime switches in interest rates. *Journal of Business and Economic Statistics*, 20:163–182, 2002b.
- A. Ang, G. Bekaert, and M. Wei. The term structure of real rates and expected inflation. *Journal of Finance*, 63:797–848, 2008.
- A. J. Auerbach and Y. Gorodnichenko. Measuring the output responses to fiscal policy. *American Economic Journal: Economic Policy*, 4:1–27, 2012.
- G. Bekaert, R. J. Hodrick, and D. A. Marshall. Peso problem explanations for term structure anomalies. *Journal of Monetary Economics*, 48:241–270, 2001.
- P. J. Bickel and Y. Ritov. Inference in hidden Markov models i: Local asymptotic normality in the stationary case. *Bernoulli*, 2:199–228, 1996.
- P. J. Bickel, Y. Ritov, and T. Rydén. Asymptotic normality of the maximum-likelihood estimator for general hidden Markov models. *Annals of Statistics*, 26:1614–1635, 1998.
- N. P. B. Bollen, S. F. Gray, and R. E. Whaley. Regime switching in foreign exchange rates: Evidence from currency option prices. *Journal of Econometrics*, 94:239–276, 2008.
- F. X. Diebold, J.-H. Lee, and G. C. Weinbach. Regime switching with time-varying transition probabilities. In C. P. Hargreaves, editor, *Nonstationary Time Series Analysis and Cointegration*, pages 283–302. Oxford University Press, Oxford, 1994.
- R. L. Dobrushin. Central limit theorem for nonstationary Markov chains. II. *Theory of Probability and Its Applications*, 1:329–383, 1956.
- R. Douc and C. Matias. Asymptotics of the maximum likelihood estimator for general hidden Markov models. *Bernoulli*, 7:381–420, 2001.
- R. Douc and E. Moulines. Asymptotic properties of the maximum likelihood estimation in misspecified hidden Markov models. *Annals of Statistics*, 40:2697–2732, 2012.

- R. Douc, G. Fort, E. Moulines, and P. Soulier. Practical drift conditions for subgeometric rates of convergence. *Annals of Applied Probability*, 14:1353–1377, 2004a.
- R. Douc, É. Moulines, and T. Rydén. Asymptotic properties of the maximum likelihood estimator in autoregressive models with Markov regime. *Annals of Statistics*, 32:2254–2304, 2004b.
- R. Douc, E. Moulines, J. Olsson, and R. van Handel. Consistency of the maximum likelihood estimator for general hidden Markov models. *Annals of Statistics*, 39:474–513, 2011.
- P. Doukhan, P. Massart, and E. Rio. The functional central limit theorem for strongly mixing processes. *Annales de l’Institut Henri Poincaré, Section B*, 30:63–82, 1994.
- M. J. Dueker, M. Sola, and F. Spagnolo. Contemporaneous threshold autoregressive models: estimation, testing and forecasting. *Journal of Econometrics*, 141:517–547, 2007.
- M. J. Dueker, Z. Psaradakis, M. Sola, and F. Spagnolo. Multivariate contemporaneous threshold autoregressive models. *Journal of Econometrics*, 160:311–325, 2011.
- C. Engel and C. S. Hakkio. The distribution of the exchange rate in the EMS. *International Journal of Finance and Economics*, 1:55–67, 1996.
- C. Engel and J. D. Hamilton. Long swings in the Dollar: Are they in the data and do markets know it? *American Economic Review*, 80:689–713, 1990.
- A. J. Filardo. Business-cycle phases and their transitional dynamics. *Journal of Business and Economic Statistics*, 12:299–308, 1994.
- A. J. Filardo and S. F. Gordon. Business cycle durations. *Journal of Econometrics*, 85:99–123, 1998.
- C. Francq and M. Roussignol. Ergodicity of autoregressive processes with Markov-switching and consistency of the maximum-likelihood estimator. *Statistics*, 32:151–173, 1998.
- D. A. Freedman. On the so-called “Huber sandwich estimator” and “robust standard errors”. *American Statistician*, 60:299–302, 2006.
- M. D. Gadea Rivas and G. Perez-Quiros. The failure to predict the Great Recession—A view through the role of credit. *Journal of the European Economic Association*, 13:534–559, 2015.
- S. F. Gray. Modeling the conditional distribution of interest rates as a regime-switching process. *Journal of Financial Economics*, 42:27–62, 1996.
- J. D. Hamilton. A new approach to the economic analysis of nonstationary time series and the business cycle. *Econometrica*, 57:357–384, 1989.

- B. E. Hansen. The likelihood ratio test under nonstandard conditions: Testing the Markov switching model of GNP. *Journal of Applied Econometrics*, 7:S61–S82 (Erratum: 11, 195–198), 1992.
- I. A. Ibragimov and R. Z. Has'minskii. *Statistical Estimation: Asymptotic Theory*. Springer-Verlag, New York, 1981.
- J. L. Jensen and N. V. Petersen. Asymptotic normality of the maximum likelihood estimator in state space models. *Annals of Statistics*, 27:514–535, 1999.
- L. Kalliovirta, M. Meitz, and P. Saikkonen. A Gaussian mixture autoregressive model for univariate time series. *Journal of Time Series Analysis*, 36:247–266, 2015.
- H. Kasahara and K. Shimotsu. Asymptotic properties of the maximum likelihood estimator in regime switching econometric models. *Journal of Econometrics*, 208: 442–467, 2019.
- V. Krishnamurthy and T. Rydén. Consistent estimation of linear and non-linear autoregressive models with Markov regime. *Journal of Time Series Analysis*, 19:291–307, 1998.
- L. Le Cam. *Asymptotic Methods in Statistical Decision Theory*. Springer-Verlag, New York, 1986.
- X. Liu and Y. Shao. Asymptotics for likelihood ratio tests under loss of identifiability. *Annals of Statistics*, 31:807–832, 2003.
- M. S. Martinez Peria. A regime-switching approach to the study of speculative attacks: A focus on the EMS crisis. *Empirical Economics*, 27:299–334, 2002.
- R. E. McCulloch and R. S. Tsay. Statistical analysis of economic time series via Markov switching models. *Journal of Time Series Analysis*, 15:523–539, 1994.
- L. Mevel and L. Finesso. Asymptotical statistics of misspecified hidden Markov models. *IEEE Transactions on Automatic Control*, 49:1123–1132, 2004.
- S. P. Meyn and R. L. Tweedie. *Markov Chains and Stochastic Stability*. Springer-Verlag, London, 1993.
- K. Mouratidis. Evaluating currency crises: A Bayesian Markov switching approach. *Journal of Macroeconomics*, 30:1688–1711, 2008.
- W. K. Newey and K. D. West. A simple, positive-definite, heteroskedasticity and autocorrelation consistent covariance matrix. *Econometrica*, 55:703–708, 1987.
- Z. Psaradakis and M. Sola. Finite-sample properties of the maximum likelihood estimator in autoregressive models with Markov switching. *Journal of Econometrics*, 86:369–386, 1998.

- Z. Psaradakis and M. Sola. Markov-switching models with state-dependent time-varying transition probabilities. *Econometrics and Statistics*, 2021. in press (<https://doi.org/10.1016/j.ecosta.2021.04.007>).
- M. O. Ravn and M. Sola. Business cycle dynamics: Predicting transitions with macrovariables. In P. Rothman, editor, *Nonlinear Time Series Analysis of Economic and Financial Data*, pages 231–265. Kluwer Academic Publishers, Dordrecht, 1999.
- F. J. Ruge-Murcia. Credibility and changes in policy regime. *Journal of Political Economy*, 103:176–208, 1995.
- T. Rydén, T. Teräsvirta, and S. Åsbrink. Stylized facts of daily returns series and the hidden Markov model. *Journal of Applied Econometrics*, 13:217–244, 1998.
- S. Sethuraman and S. R. S. Varadhan. A martingale proof of Dobrushin’s theorem for non-homogeneous Markov chains. *Electronic Journal of Probability*, 10:1221–1235, 2005.
- P. W. Simpson, D. R. Osborn, and M. Sensier. Modelling business cycle movements in the uk economy. *Economica*, 68:243–267, 2001.
- K. J. Tadjuidje, H. Ombao, and R. A. Davis. Autoregressive processes with data-driven regime switching. *Journal of Time Series Analysis*, 30:505–533, 2009.
- C.-Y. Tan, Y.-B. Koh, K.-H. Ng, and K.-H. Ng. Dynamic volatility modelling of Bitcoin using time-varying transition probability Markov-switching GARCH model. *North American Journal of Economics and Finance*, 56:Article 101377, 2021.
- H. White. Maximum likelihood estimation of misspecified models. *Econometrica*, 50:1–25, 1982.
- H. White. *Estimation, Inference and Specification Analysis*. Cambridge University Press, Cambridge, 1994.
- H. White. *Asymptotic Theory for Econometricians*. Revised Edition, Academic Press, London, 2001.
- R. F. Whitelaw. Stock market risk and return: An equilibrium approach. *Review of Financial Studies*, 13:521–547, 2000.

# Online Supplemental Material

For any measure  $P$ , we use  $L^r(P)$ ,  $1 \leq r < \infty$ , to denote the class of measurable functions integrable to order  $r$  with respect to  $P$ ;  $\|\cdot\|_{L^r(P)}$  denotes the usual  $r$ -norm in  $L^r(P)$ . For a vector/matrix-valued functions  $X \mapsto f(X)$ ,  $\|f\|_{L^r(P)}$  is short-hand notation for the  $L^r(P)$ -norm of  $x \mapsto \|f(x)\|$ , where  $\|\cdot\|$  denotes the Euclidean/dual norm of  $f$ . For any two sequences of random variables  $(X_n)_n$  and  $(Y_n)_n$ ,  $X_n \preceq Y_n$ , implies that  $X_n \leq CY_n$  for some universal positive finite constant  $C$ .

## SM.1 Ergodicity and Stationarity

Let  $(\zeta_t)_{t=-\infty}^\infty$  be a Markov chain with transition kernel  $\zeta \mapsto \mathbb{P}(\zeta, \cdot) \in \mathcal{P}(\mathbb{Z})$  and  $\zeta_t \in \mathbb{Z} \subseteq \mathbb{R}^d$  for some  $d > 0$ . Also, for any probability measure  $P$  over  $\mathbb{Z}$  and any  $f : \mathbb{Z} \rightarrow \mathbb{R}$ , let  $P[f] \equiv \int f(z)P(dz)$  and  $P[f](z) \equiv \int f(u)P(z, du)$  (if it exists).

**Assumption 9.** *There exist constants  $\gamma \in (0, 1)$ ,  $\lambda \in (0, 1)$ ,  $b > 0$  and  $R > 2b/(1 - \gamma)$ , a function  $\mathbf{V} : \mathbb{Z} \rightarrow [1, \infty)$ , and a probability measure  $\varrho$  such that:*  
*(i)  $\mathbb{P}[\mathbf{V}](\zeta) \leq \gamma \mathbf{V}(\zeta) + b1\{\zeta \in \mathcal{C}\}$  for all  $\zeta \in \mathbb{Z}$  with  $\mathcal{C} \equiv \{\zeta \in \mathbb{Z} : \mathbf{V}(\zeta) \leq R\}$  ;*  
*(ii)  $\inf_{\zeta \in \mathcal{C}} \mathbb{P}(\zeta, \cdot) \geq \lambda \varrho(\cdot)$ , with  $\varrho(\mathcal{C}) > 0$ .*

The next result is used for the proof of Lemma 1; it contains well-known results that are stated and proved here for convenience. In particular, the first part of Lemma 10 is a re-statement of Theorem 1.2 in Hairer and Mattingly [2011]. The second part of Lemma 10 and Assumption 9(ii) imply that  $\mathbb{P}$  is Harris recurrent (see [Athreya and Lahiri, 2006, Ch. 14]) and aperiodic (see [Thierney, 1996, p. 65]). The proof follows from standard arguments.

Let  $v \mapsto \|v\|_{\mathbf{V}} \equiv \sup_{\zeta} \frac{|v(\zeta)|}{1 + \mathbf{V}(\zeta)}$ . Also, for any  $A \subseteq \mathbb{Z}$ , let  $T_A = \inf\{t \geq 0 : \zeta_t \in A\}$ .

**Lemma 10.** *If Assumption 9 holds, then:*

*(i)  $\mathbb{P}$  admits a unique invariant measure  $\nu^*$ , and there exist constants  $\gamma \in (0, 1)$  and  $C > 0$  such that*

$$\|\mathbb{P}^n[v] - \nu^*[v]\|_{\mathbf{V}} \leq C\gamma^n \|v - \nu^*[v]\|_{\mathbf{V}}$$

*for every measurable function  $v$  such that  $\|v\|_{\mathbf{V}} < \infty$ , where  $\nu^*[v] \equiv \int v(\zeta)\nu^*(d\zeta)$ .*

*(ii)  $\mathbb{P}(\zeta, \{T_{\mathcal{C}} < \infty\}) = 1$  for all  $\zeta \in \mathbb{Z}$ , and  $\mathbb{P}(\zeta_0, \mathcal{C}) > 0$  for all  $\zeta_0 \in \mathcal{C}$ .*

*Proof of Lemma 10.* Part (i) is Theorem 1.2 in Hairer and Mattingly [2011]. Assumption 9(i) implies their Assumption 1 with  $K = b$  and Assumption 9(ii) implies their Assumption 2.

For part (ii), we first establish that  $\mathbb{P}(\zeta_0, \mathcal{C}) = \mathbb{P}(\zeta_0, \{\zeta_1 \in \mathcal{C}\}) > 0$  for all  $\zeta_0 \in \mathcal{C}$ . For this, note that  $\mathbb{P}(\zeta_0, \mathcal{C}) \geq \inf_{\zeta \in \mathcal{C}} \mathbb{P}(\zeta, \mathcal{C}) \geq \lambda \varrho(\mathcal{C}) > 0$  by Assumption 9(ii).

We now show that  $\mathbb{P}(\zeta, \{T_{\mathcal{C}} < \infty\}) = 1$  for all  $\zeta \in \mathbb{Z}$ . It suffices to show that  $\mathbb{P}[T_{\mathcal{C}}](\zeta) < \infty$  for all  $\zeta \notin \mathcal{C}$ . Under Assumption 9(i),  $\mathbf{V} \geq 1$ , so

$$\mathbb{P}[T_{\mathcal{C}}](\zeta) \leq \mathbb{P}\left[\sum_{j=0}^{T_{\mathcal{C}}-1} \mathbf{V}(\zeta_j)\right](\zeta) = \sum_{T=0}^{\infty} \sum_{j=0}^T E_{\mathbb{P}}[\mathbf{V}(\zeta_j) \mid T_{\mathcal{C}} = T+1, \zeta] \Pr(\zeta, \{T_{\mathcal{C}} = T+1\})$$

for any  $\zeta \in \mathbb{Z} \setminus \mathcal{C}$ . To establish the desired result, it is sufficient to show that  $\sup_T \sum_{j=0}^T E_{\mathbb{P}}[\mathbf{V}(\zeta_j) \mid T_{\mathcal{C}} = T+1, \zeta] < \infty$ .

Take any  $T \geq 0$  and any  $j \leq T$ , and note that

$$\begin{aligned} E_{\mathbb{P}}[\mathbb{P}[\mathbf{V}](\zeta_j) \mid \zeta_l \notin \mathcal{C}, \forall l \leq T+1] &= E_{\mathbb{P}}\left[\int_{\zeta \notin \mathcal{C}} \mathbb{P}[\mathbf{V}](\zeta) \mathbb{P}(\zeta_{j-1}, d\zeta) \mid \zeta_l \notin \mathcal{C}, \forall l \leq j-1\right] \\ &\leq \gamma E_{\mathbb{P}}\left[\int_{\zeta \notin \mathcal{C}} \mathbf{V}(\zeta) \mathbb{P}(\zeta_{j-1}, d\zeta) \mid \zeta_l \notin \mathcal{C}, \forall l \leq j-1\right] \\ &\leq \gamma E_{\mathbb{P}}[\mathbb{P}[\mathbf{V}](\zeta_{j-1}) \mid \zeta_l \notin \mathcal{C}, \forall l \leq j-1] \\ &\leq \gamma^j \mathbf{V}(\zeta_0), \end{aligned}$$

where the second line follows from Assumption 9(i) and the fact that  $\zeta \notin \mathcal{C}$ , the third line follows from the fact that  $\mathbf{V} > 0$ , and the last line follows from repeated iteration of the first lines. Note that  $T_{\mathcal{C}} = T+1$  is equivalent to  $\zeta_j \notin \mathcal{C}, \forall j \leq T$  and  $\zeta_{T+1} \in \mathcal{C}$ . Thus, the previous display implies that

$$E_{\mathbb{P}}[\mathbf{V}(\zeta_j) \mid T_{\mathcal{C}} = T+1] \leq \gamma^j \mathbf{V}(\zeta_0)$$

for any  $T \geq 0$  and any  $j \leq T$ . Consequently,  $\sum_{j=0}^T E_{\mathbb{P}}[\mathbf{V}(\zeta_j) \mid T_{\mathcal{C}} = T+1, \zeta] \leq \mathbf{V}(\zeta) \sum_{j=0}^T \gamma^j \leq \frac{\mathbf{V}(\zeta)}{1-\gamma}$ , and thus the result follows.  $\square$

*Proof of Lemma 1.* Let  $(\zeta_t)_{t=-\infty}^{\infty}$  be the stochastic process given by  $\zeta_t \equiv (X_t, S_t)$ . This process is a Markov chain with transition kernel  $\mathbb{X} \times \mathbb{S} \ni \zeta \mapsto \mathbb{P}(\zeta, \cdot) \in \mathcal{P}(\mathbb{X} \times \mathbb{S})$  given by

$$\mathbb{P}((x, s), \{\zeta_{t+1} \in A_1 \times A_2\}) = \sum_{s' \in A_2} Q_*(x, s, s') P_*(x, s', A_1),$$

for any Borel sets  $A_1 \subseteq \mathbb{X}$  and  $A_2 \subseteq \mathbb{S}$ .

By Lemma 10, there exists a unique invariant measure  $\nu$ , provided that the conditions of Assumption 9 are met. In order to verify the first part of

Assumption 9, consider  $\mathbf{V}(\zeta) = \mathcal{U}(x)$ , and  $\mathcal{C} \equiv \mathcal{C}_1 \times \mathbb{S}$  with  $\mathcal{C}_1 \equiv \{x \in \mathbb{X} : \mathcal{U}(x) \leq R\}$ . By Assumption 2(i),

$$\mathbb{P}[\mathbf{V}](\zeta) = \int_{\mathbb{X}} \mathcal{U}(x') \left\{ \sum_{s' \in \mathbb{S}} Q_*(x, s, s') P_*(x, s', dx') \right\} \leq \gamma \mathcal{U}(x) + 2b' 1\{x \in \mathcal{C}_1\}.$$

Thus,  $b \equiv 2b'$ . Regarding Assumption 9(ii), observe that, by Assumption 1(i), for  $C$  and any  $s \in \mathbb{S}$ ,

$$\mathbb{P}((x, s), C \times \{s'\}) \geq \underline{q}(x) P_*(x, s', C),$$

and, by Assumption 2(iii),  $P_*(x, s', C) \geq \lambda' \varpi(C)$  and  $\lambda' \in (0, 1)$ . Also note that, by Assumption 1,  $\underline{q}$  is continuous and  $\underline{q}(x) > 0$  for all  $x \in \mathbb{X}$ . Furthermore, by Assumption 2(ii),  $\mathcal{U}$  is lower semi-compact, because  $\{x \in \mathbb{X} : \mathcal{U}(x) \leq R\}$  is closed ( $x \mapsto \mathcal{U}(x)$  is lower semi-continuous), and is also bounded. Therefore,  $\inf_{x: \mathcal{U}(x) \leq R} \underline{q}(x) = \min_{x: \mathcal{U}(x) \leq R} \underline{q}(x) \geq c > 0$  (because it is a minimization of a continuous function on compact set). Therefore,

$$\mathbb{P}(\zeta, C \times \{s'\}) \geq c\lambda' \varpi(C) \frac{1}{|\mathbb{S}|},$$

and, by putting  $\varrho = \varpi(\cdot) \frac{1}{|\mathbb{S}|}$  and  $\lambda \equiv c\lambda'$ , Assumption 9(ii) follows since  $\varpi(\mathcal{C}_1) > 0$ . Since  $\nu$  is unique, it is trivially ergodic. Therefore, the process with initial probability measure  $\nu$  is stationary. Ergodicity of  $(\zeta_t)_t$  follows from Theorem 14.2.11 in [Athreya and Lahiri \[2006\]](#) (recall that  $\mathbb{P}$  is Harris recurrent and aperiodic). Since  $X_t$  is a deterministic function of  $\zeta_t$ ,  $X_0^\infty$  is also stationary and ergodic. Finally, observe that

$$\int \sup_{0 \leq f \leq 1} |\mathbb{P}^n[f](\zeta) - \nu[f]| \nu(d\zeta) \lesssim \gamma^n \int |1 + \mathcal{U}(x)| \nu(d\zeta).$$

Since  $\mathcal{U}$  satisfies Assumption 9(i), it follows that  $\int \mathbb{P}[\mathcal{U}](\zeta) \nu(d\zeta) \leq \gamma \nu[\mathcal{U}] + K$ . Since  $\nu$  is the invariant measure of  $\mathbb{P}$  and  $\gamma \in (0, 1)$ , this implies that  $\nu(d\zeta) \leq K/(1 - \gamma)$ . Therefore,

$$\int \sup_{0 \leq f \leq 1} |\mathbb{P}^n[f](\zeta) - \nu[f]| \nu(d\zeta) \lesssim \gamma^n,$$

thereby implying that  $(\zeta_t)_t$  is  $\beta$ -mixing with rate  $\beta_n = O(\gamma^n)$  (see [Davydov \[1973\]](#)). Since  $X_t$  is a deterministic function of  $\zeta_t$ , the same holds for  $X_0^\infty$ .  $\square$



## SM.2 Proofs of Supplementary Lemmas in Appendix A.1

To prove Lemmas 2 and 3 we use the following result.

**Lemma 11.** *Suppose Assumptions 1 and 4(ii) hold. Then, for all  $t \in \mathbb{N}$  and  $-n \leq -m \leq t-1$ ,*

$$\sup_{\theta \in \Theta} |\log p_t^\nu(X_t | X_{-m}^{t-1}, \theta) - \log p_t^\nu(X_t | X_{-n}^{t-1}, \theta)| \leq C(X_{t-1}, X_t) \prod_{i=-m}^{t-1} (1 - \underline{q}(X_i))$$

a.s.- $\bar{P}_*^\nu$ .

*Proof of Lemma 11.* Observe that, for any  $n \in \mathbb{N}$ ,

$$\log p_t^\nu(X_t | X_{-n}^{t-1}, \theta) = \log \sum_{s \in \mathbb{S}} p_\theta(X_{t-1}, s, X_t) \bar{P}_\theta^\nu(s | X_{-n}^{t-1}),$$

and since  $\log x - \log y \leq x/y - 1$ , it suffices to study  $\frac{\sum_{s \in \mathbb{S}} p_\theta(X_{t-1}, s, X_t) (\bar{P}_\theta^\nu(S_t = s | X_{-m}^{t-1}) - \bar{P}_\theta^\nu(S_t = s | X_{-n}^{t-1}))}{\sum_{s \in \mathbb{S}} p_\theta(X_{t-1}, s, X_t) \bar{P}_\theta^\nu(s | X_{-n}^{t-1})}$ .

This expression can be bounded above by

$$\frac{\max_{s \in \mathbb{S}} p_\theta(X_{t-1}, s, X_t)}{\min_{s \in \mathbb{S}} p_\theta(X_{t-1}, s, X_t)} \|\bar{P}_\theta^\nu(S_t = \cdot | X_{-m}^{t-1}) - \bar{P}_\theta^\nu(S_t = \cdot | X_{-n}^{t-1})\|_1.$$

By Assumption 4(ii),  $\sup_{\theta \in \Theta} \frac{\max_{s \in \mathbb{S}} p_\theta(X_{t-1}, s, X_t)}{\min_{s \in \mathbb{S}} p_\theta(X_{t-1}, s, X_t)} \leq C(X_{t-1}, X_t)$  a.s.- $\bar{P}_*^\nu$ . So it suffices to bound  $\|\bar{P}_\theta^\nu(S_t = \cdot | X_{-m}^{t-1}) - \bar{P}_\theta^\nu(S_t = \cdot | X_{-n}^{t-1})\|_1$ .

By Lemma B.2.2 in Stachurski [2009],

$$\begin{aligned} & \|\bar{P}_\theta^\nu(S_t = \cdot | X_{-m}^{t-1}) - \bar{P}_\theta^\nu(S_t = \cdot | X_{-n}^{t-1})\|_1 \\ & \leq \frac{1}{2} \sup_{b, c \in \mathbb{S}^2} \|\bar{P}_\theta^\nu(S_t = \cdot | S_{-m} = b, X_{-m}^{t-1}) - \bar{P}_\theta^\nu(S_t = \cdot | S_{-m} = c, X_{-n}^{t-1})\|_1 \\ & = \frac{1}{2} \sup_{b, c \in \mathbb{S}^2} \|\bar{P}_\theta^\nu(S_t = \cdot | S_{-m} = b, X_{-m}^{t-1}) - \bar{P}_\theta^\nu(S_t = \cdot | S_{-m} = c, X_{-m}^{t-1})\|_1, \end{aligned}$$

where the last line follows from the fact that, given  $S_{-m}$ , it is the same to condition on  $X_{-m}^{t-1}$  and on  $X_{-n}^{t-1}$ . Hence,

$$\sup_{\theta \in \Theta} |\log p_t^\nu(X_t | X_{-m}^{t-1}, \theta) - \log p_t^\nu(X_t | X_{-n}^{t-1}, \theta)| \leq C' \alpha_{\theta, t, -m}(X_{-m}^{t-1}),$$

where  $\alpha_{\theta, t, -m}(X_{-m}^{t-1})$  is defined in expression (12). By Applying Lemmas 6 and 5 and the fact that  $\alpha_{\theta, t, -m}(X_{-m}^{t-1}) \leq \prod_{j=-m}^{t-1} \alpha_{\theta, j, j+1}(X_{-m}^{t-1})$ , it follows that

$$\sup_{\theta \in \Theta} |\log p_t^\nu(X_t | X_{-m}^{t-1}, \theta) - \log p_t^\nu(X_t | X_{-n}^{t-1}, \theta)| \leq C(X_{t-1}, X_t) \prod_{i=-m}^{t-1} (1 - \underline{q}(X_i)),$$

a.s.- $\bar{P}_*^\nu$ . □

We now prove Lemmas 2 and 3.

*Proof of Lemma 2.* Fix any  $\epsilon > 0$ . By Lemma 11, with  $m = 0$  and  $n = M$ ,

$$\begin{aligned} & \sup_{\theta \in \Theta} |\log p_t^\nu(X_t | X_0^{t-1}, \theta) - \log p^\nu(X_t | X_{-\infty}^{t-1}, \theta)| \\ & \leq \limsup_{M \rightarrow \infty} \sup_{\theta \in \Theta} |\log p_t^\nu(X_t | X_0^{t-1}, \theta) - \log p_t^\nu(X_t | X_{-M}^{t-1}, \theta)| \\ & \leq C(X_{t-1}, X_t) \prod_{i=0}^{t-1} (1 - \underline{q}(X_i)). \end{aligned}$$

Thus, it suffices to show that there exists a  $T$  such that for all  $t \geq T$ ,  $\bar{P}_\nu^* \left( T^{-1} \sum_{t=1}^T C(X_{t-1}, X_t) \prod_{i=0}^{t-1} (1 - \underline{q}(X_i)) \geq \epsilon \right) < \epsilon$ . This follows from Assumption 5.  $\square$

*Proof of Lemma 3.* Recall that, by Lemma 1, the process  $X_{-\infty}^\infty$  is ergodic and stationary under  $\bar{P}_*^\nu$ .

**Part (i).** Consider a  $\delta > 0$  and an open cover  $\{B(\theta, \delta) : \theta \in \Theta\}$  where  $B(\theta, \delta)$  is an open ball centered around  $\theta$  with radius  $\delta > 0$ . Since  $\Theta$  is compact (Assumption 3), there exists a finite sub-cover  $B_j \equiv B(\theta_j, \delta)$  with  $j = 1, \dots, J$ . Also note that, pointwise in  $\theta \in \Theta$ ,  $\ell_T^\nu(X_{-\infty}^T, \theta) - E_{\bar{P}_*^\nu}[\ell_T^\nu(X_{-\infty}^T, \theta)] \rightarrow 0$  a.s.- $\bar{P}_*^\nu$  by the ergodic theorem and the fact that  $X_{-\infty}^\infty \mapsto \ell_T^\nu(X_{-\infty}^T, \theta) \in L^1(\bar{P}_{\theta_*}^\nu)$ . Thus, it suffices to show that there exists a  $T(j, \epsilon)$  such that, for all  $t \geq T(j, \epsilon)$ ,

$$\bar{P}_*^\nu \left( \sup_{\theta \in B_j} T^{-1} \sum_{t=1}^T \left( l_t(X_{-\infty}^t, \theta) - E_{\bar{P}_*^\nu}[l_t(X_{-\infty}^t, \theta)] \right) > \epsilon \right) \leq \epsilon,$$

where  $l_t(X_{-\infty}^t, \theta) \equiv \log \frac{p^\nu(X_t | X_{-\infty}^{t-1}, \theta)}{p^\nu(X_t | X_{-\infty}^{t-1}, \theta_j)}$ . Observe that, for any  $j$ ,

$$\begin{aligned} \sup_{\theta \in B_j} \sum_{t=1}^T \left( l_t(X_{-\infty}^t, \theta) - E_{\bar{P}_*^\nu}[l_t(X_{-\infty}^t, \theta)] \right) & \leq \sum_{t=1}^T \sup_{\theta \in B_j} \left( l_t(X_{-\infty}^t, \theta) - E_{\bar{P}_*^\nu}[l_t(X_{-\infty}^t, \theta)] \right) \\ & \equiv \sum_{t=1}^T \bar{l}_t(X_{-\infty}^t). \end{aligned}$$

Moreover, observe that

$$\sup_{\theta \in B_j} \log \frac{p^\nu(X_t | X_{-\infty}^{t-1}, \theta)}{p^\nu(X_t | X_{-\infty}^{t-1}, \theta_j)} \leq \sup_{\theta \in B_j} \frac{p^\nu(X_t | X_{-\infty}^{t-1}, \theta)}{p^\nu(X_t | X_{-\infty}^{t-1}, \theta_j)} - 1.$$

By Assumption 4(i), for any  $\epsilon > 0$  there exists a  $\delta > 0$  such that  $E_{\bar{P}_*^\nu} \left[ \sup_{\theta \in B_j} \frac{p^\nu(X_0 | X_{-\infty}^{-1}, \theta)}{p^\nu(X_0 | X_{-\infty}^{-1}, \theta_j)} \right] \leq 1 + \epsilon$  for any  $j \in \{1, \dots, J\}$  and any  $t$ . Therefore, we can choose a  $\delta > 0$  such that

$$E_{\bar{P}_*^\nu} \left[ \sup_{\theta \in B_j} \log \frac{p^\nu(X_0 | X_{-\infty}^{-1}, \theta)}{p^\nu(X_0 | X_{-\infty}^{-1}, \theta_j)} \right] \leq \epsilon/4.$$

This in turn implies that  $E_{\bar{P}_*^\nu}[\bar{l}_t(X_{-\infty}^t)] \leq \epsilon/2$ . This result and the ergodic theorem establish that  $\lim_{T \rightarrow \infty} T^{-1} \sum_{t=1}^T \bar{l}_t(X_{-\infty}^t) \leq \epsilon/2$  a.s.- $\bar{P}_*^\nu$ . This implies the result in (11).

**Part (ii).** Follows directly from the ergodic theorem and the fact that  $X_{-\infty}^\infty \mapsto \log p^\nu(X_t | X_{-\infty}^{t-1}, \theta_*)$  is in  $L^1(\bar{P}_*^\nu)$ .  $\square$

### SM.3 Properties of $p_\theta(X_1 | X_{-\infty}^0)$

For any  $t \in \mathbb{N}_0 \equiv \mathbb{N} \cup \{0\}$ , any  $X_{-\infty}^t$  and any  $\theta \in \Theta$ ,  $p^\nu(X_t | X_{-\infty}^{t-1}, \theta)$  is defined as  $\liminf_{M \rightarrow \infty} p_\theta^\nu(X_t | X_{-M}^{t-1})$ ;  $p_*(X_t | X_{-\infty}^{t-1})$  is defined analogously.

**Lemma 12.** *Suppose Assumptions 4(ii) and 5 hold. Then:*

(1) *For any  $t \in \mathbb{N}_0$ ,  $x \mapsto p^\nu(\cdot | X_{-\infty}^{t-1}, \theta)$  and  $x \mapsto p_*(\cdot | X_{-\infty}^{t-1})$  are densities, a.s.- $\bar{P}_*^\nu$ .*

(2) *Suppose  $\Theta$  is compact and that for each  $n \in \mathbb{N}_0$ ,  $\theta \mapsto p_\theta^\nu(X_1 | X_{-n}^0)$  is uniformly continuous a.s.- $\bar{P}_*^\nu$ . Then, for any  $\theta_0 \in \Theta$  and  $\epsilon > 0$ , there exists a  $\delta > 0$  such that*

$$\bar{P}_*^\nu \left( \sup_{\theta_0 \in \Theta} \sup_{\theta \in B(\theta_0, \delta)} |p^\nu(X_1 | X_{-\infty}^0, \theta) - p^\nu(X_1 | X_{-\infty}^0, \theta_0)| > \epsilon \right) < \epsilon.$$

(3) *Suppose  $\Theta$  is compact and that for each  $n \in \mathbb{N}_0$ ,  $\theta \mapsto p_\theta^\nu(X_1 | X_{-n}^0)$  is uniformly continuous a.s.- $\bar{P}_*^\nu$ . Suppose also that there exists functions  $(x_1, x_0) \mapsto (\bar{p}(x_0, x_1), \underline{p}(x_0, x_1))$  such that for any  $p \in \{p_\theta : \theta \in \Theta\} \cup p_*$ ,  $\underline{p}(x_0, x_1) \leq p(x_0, s, x_1) \leq \bar{p}(x_0, x_1)$  for all  $s \in \mathbb{S}$ , and  $E_{\bar{P}_*^\nu}[\bar{p}(X_0, X_1)/\underline{p}(X_0, X_1)] < \infty$ . Then, Assumptions 3 and 4 hold.*

*Proof.* (1) We need to show that the functions integrate to 1. By analogous

steps to those in the proof of Lemma 11, it follows that

$$\begin{aligned}
& \left| \int \{p^\nu(x|X_{-\infty}^{t-1}, \theta) - p_\theta^\nu(x|X_{-n}^{t-1})\} dx \right| \\
& \leq \int \sum_s p_\theta(X_{t-1}, s, x) dx \limsup_{M \rightarrow \infty} \|\bar{P}_\theta^\nu(S_t = \cdot | X_{-M}^{t-1}) - \bar{P}_\theta^\nu(S_t = \cdot | X_{-n}^{t-1})\|_1 \\
& \leq \int \sum_s p_\theta(X_{t-1}, s, x) dx \limsup_{M \rightarrow \infty} \prod_{i=-n}^{t-1} (1 - \underline{q}(X_i)) \\
& = \prod_{i=-n}^{t-1} (1 - \underline{q}(X_i)) |\mathbb{S}|.
\end{aligned}$$

Since this holds for any  $n$  such that  $-n \leq t-1$ , we can take averages and obtain, for any  $\epsilon > 0$ , that

$$\begin{aligned}
\bar{P}_*^\nu \left( \frac{1}{M+1} \sum_{n=0}^M \left| \int p^\nu(x|X_{-\infty}^{t-1}, \theta) dx - 1 \right| > \epsilon \right) & \leq \bar{P}_*^\nu \left( \frac{1}{M+1} \sum_{n=0}^M \prod_{i=-n}^{t-1} (1 - \underline{q}(X_i)) |\mathbb{S}| > \epsilon \right) \\
& \leq \bar{P}_*^\nu \left( \frac{1}{M+1} \sum_{n=0}^M \prod_{i=-n}^0 (1 - \underline{q}(X_i)) |\mathbb{S}| > \epsilon \right).
\end{aligned}$$

Since this holds for any  $M$ , by taking  $M \rightarrow \infty$ , stationarity and Assumption 5 imply that the RHS vanishes. Thus, it follows that  $\bar{P}_*^\nu(|\int p^\nu(x|X_{-\infty}^{t-1}, \theta) dx - 1| \geq \epsilon) = 0$ . As the  $\epsilon > 0$  is arbitrary, this implies that  $\int p^\nu(x|X_{-\infty}^{t-1}, \theta) dx = 1$ , a.s.- $\bar{P}_*^\nu$ . Following the same logic, an analogous result can be shown for  $p_*^\nu(\cdot|X_{-\infty}^{t-1})$ . (2)

Similarly, for any  $n$  such that  $-n \leq t-1$ ,

$$\sup_{\theta \in \Theta} |p^\nu(X_t|X_{-\infty}^{t-1}, \theta) - p_\theta^\nu(X_t|X_{-n}^{t-1})| \leq \sum_{s \in \mathbb{S}} p_\theta(X_{t-1}, s, X_t) \prod_{i=-n}^{t-1} (1 - \underline{q}(X_i)).$$

Hence, for any  $\theta_0$  in  $\Theta$ ,

$$\begin{aligned}
& \sup_{\theta \in B(\theta_0, \delta)} |p^\nu(X_t|X_{-\infty}^{t-1}, \theta) - p^\nu(X_t|X_{-\infty}^{t-1}, \theta_0)| \\
& \leq \sup_{\theta \in B(\theta_0, \delta)} \sum_{s \in \mathbb{S}} p_\theta(X_{t-1}, s, X_t) \frac{1}{1+M} \sum_{n=0}^M \prod_{i=-n}^{t-1} (1 - \underline{q}(X_i)) \\
& \quad + \frac{1}{1+M} \sum_{n=0}^M \sup_{\theta \in B(\theta_0, \delta)} |p_\theta^\nu(X_t|X_{-n}^{t-1}, \theta) - p_{\theta_0}^\nu(X_t|X_{-n}^{t-1})|.
\end{aligned}$$

For any  $\gamma > 0$ , choose  $M$  such that  $\bar{P}_*^\nu \left( \frac{1}{1+M} \sum_{n=0}^M \prod_{i=-n}^{t-1} (1 - \underline{q}(X_i)) \geq \gamma \right) < \gamma$ ; such  $M$  exists by Assumption 5. Given such  $M$ , for any  $\epsilon > 0$  and any  $\delta > 0$ , it follows that

$$\begin{aligned} & \bar{P}_*^\nu \left( \sup_{\theta_0 \in \Theta} \sup_{\theta \in B(\theta_0, \delta)} |p^\nu(X_t | X_{-\infty}^{t-1}, \theta) - p^\nu(X_t | X_{-\infty}^{t-1}, \theta_0)| > \epsilon \right) \\ & \leq \bar{P}_*^\nu \left( \sup_{\theta \in \Theta} \sum_{s \in \mathbb{S}} p_\theta(X_{t-1}, s, X_t) \geq 0.5\epsilon/\gamma \right) + \gamma/3 \\ & \quad + \bar{P}_*^\nu \left( \frac{1}{1+M} \sum_{n=0}^M \sup_{\theta_0 \in \Theta} \sup_{\theta \in B(\theta_0, \delta)} |p_\theta^\nu(X_t | X_{-n}^{t-1}) - p_{\theta_0}^\nu(X_t | X_{-n}^{t-1})| \geq 0.5\epsilon \right). \end{aligned}$$

Let  $\delta > 0$  be such that for each  $n \in \{0, \dots, M\}$ ,  $\sup_{\theta_0 \in \Theta} \sup_{\theta \in B(\theta_0, \delta)} |p_\theta^\nu(X_t | X_{-n}^{t-1}) - p_{\theta_0}^\nu(X_t | X_{-n}^{t-1})| < 0.5\epsilon$  a.s.- $\bar{P}_*^\nu$ ; such  $\delta > 0$  exists by our conditions. So the third term in the RHS is 0. Also, under our conditions, it follows that  $\sup_{\theta \in \Theta} p_\theta(X_{t-1}, s, X_t) = O_{\bar{P}_*^\nu}(1)$  and thus the first term in the RHS can be made smaller than  $\epsilon/3$  by a suitably chosen  $\gamma \in (0, \epsilon)$ . Thus, the desired result holds. (3) By the definition

of  $p^\nu(\cdot, \cdot, \theta)$ , it follows that

$$p^\nu(X_1 | X_{-\infty}^0, \theta) \leq \max_{s \in \mathbb{S}} p_\theta(X_0, s, X_1),$$

and, for some  $n \in \mathbb{N}_0$ ,

$$\begin{aligned} p^\nu(X_1 | X_{-\infty}^0, \theta) & \geq 0.5 \sum_{s \in \mathbb{S}} p_\theta(X_0, s, X_1) Pr_\theta(S_1 = s | X_{-n}^0) \\ & \geq 0.5 \min_{s \in \mathbb{S}} p_\theta(X_0, s, X_1). \end{aligned}$$

Thus, by our conditions, for any  $\delta > 0$  and  $\theta_0 \in \Theta$ ,

$$\sup_{\theta \in B(\delta, \theta_0)} \frac{p^\nu(X_1 | X_{-\infty}^0, \theta)}{p^\nu(X_1 | X_{-\infty}^0, \theta_0)} \leq \frac{\bar{p}(X_0, X_1)}{\underline{p}(X_0, X_1)},$$

and the RHS is in  $L^1(\bar{P}_*^\nu)$ . Thus, by part (2) and the dominated convergence theorem, for any  $\epsilon > 0$ , there exists a  $\delta > 0$  such that

$$E_{\bar{P}_*^\nu} \left[ \sup_{\theta \in B(\delta, \theta_0)} \frac{p^\nu(X_1 | X_{-\infty}^0, \theta)}{p^\nu(X_1 | X_{-\infty}^0, \theta_0)} \right] \leq 1 + \epsilon,$$

for any  $\theta_0 \in \Theta$ . This readily implies Assumption 4(i). Part (ii) also follows with  $(x_0, x_1) \mapsto C(x_0, x_1) = \frac{\bar{p}(x_0, x_1)}{\underline{p}(x_0, x_1)}$ . Similarly, by noting that

$$1 - \frac{p^\nu(X_1 | X_{-\infty}^0, \theta)}{p_*^\nu(X_1 | X_{-\infty}^0)} \leq \log \frac{p^\nu(X_1 | X_{-\infty}^0, \theta)}{p_*^\nu(X_1 | X_{-\infty}^0)} \leq \frac{p^\nu(X_1 | X_{-\infty}^0, \theta)}{p_*^\nu(X_1 | X_{-\infty}^0)} - 1,$$

it follows that, for any  $\theta \in \Theta$ ,

$$\left| \log \frac{p^\nu_*(X_1|X_{-\infty}^0)}{p^\nu(X_1|X_{-\infty}^0, \theta)} \right| \leq 1 + \frac{\bar{p}(X_0, X_1)}{\underline{p}(X_0, X_1)}.$$

Since the RHS is in  $L^1(\bar{P}_*)$ , the results in part (2) and the dominated convergence theorem imply that  $\theta \mapsto H(\theta)$  is continuous.  $\square$

## SM.4 Sufficient Conditions for Assumptions 5 and 8

By exploiting the fact that  $(X_t)_{t=-\infty}^\infty$  is  $\beta$ -mixing and stationary, the following lemma provides sufficient conditions for Assumption 5.

**Lemma 13.** *Suppose Assumptions 1 and 2 hold.*

(1) *Suppose further there exists  $l \geq 1$  such that*

$$E_{\bar{P}_\nu^*} \left[ (1 - \underline{q}(X_1))^{l'} \right] < 1 \text{ and } E_{\bar{P}_\nu^*} \left[ C(X_1, X_0)^l \right] < \infty,$$

where  $1/l' + 1/l = 1$ . Then,

$$\lim_{T \rightarrow \infty} E_{\bar{P}_\nu^*} \left[ T^{-1} \sum_{t=1}^T C(X_{t-1}, X_t) \prod_{i=0}^{t-1} (1 - \underline{q}(X_i)) \right] = 0$$

(2) *Suppose further  $E_{\bar{P}_\nu^*} \left[ (1 - \underline{q}(X_1))^{\frac{2a}{1-a}} \right] < 1$ . Then, Assumption 8 holds.*

Clearly, by the Markov inequality, Part (1) implies that

$$\lim_{T \rightarrow \infty} \bar{P}_\nu^* \left( T^{-1} \sum_{t=1}^T C(X_{t-1}, X_t) \prod_{i=0}^{t-1} (1 - \underline{q}(X_i)) \geq \epsilon \right) = 0.$$

*Proof.* (1) We use well-known coupling results for  $\beta$ -mixing sequences (see Yu [1994]). For any  $q \in \mathbb{N}$ , let  $(\tilde{X}_t)_{t=-\infty}^\infty$  be such that: (a) for any  $i \in \mathbb{N}_0$ ,  $\bar{U}_i \equiv (\tilde{X}_{iq+1}, \dots, \tilde{X}_{iq+q})$  has the same distribution as  $U_i \equiv (X_{iq+1}, \dots, X_{iq+q})$ ; (b) the sequence  $(\bar{U}_i)_{i \text{ even}}$  is i.i.d. and so is  $(\bar{U}_i)_{i \text{ odd}}$ ; (c) for any  $i \in \mathbb{N}_0$ ,  $\mathbb{P}(\bar{U}_i \neq U_i) \leq \beta_q$ , where  $\mathbb{P}$  is the product measure of the random processes  $(X_t)_{t=-\infty}^\infty$  and  $(\tilde{X}_t)_{t=-\infty}^\infty$ .

By stationarity, for any  $t \in \mathbb{N}$ ,

$$\begin{aligned} E_{\bar{P}_\nu^*} \left[ C(X_{t-1}, X_t) \prod_{i=0}^{t-1} (1 - \underline{q}(X_i)) \right] &= E_{\bar{P}_\nu^*} [C(X_{t-1}, X_t)(1 - \underline{q}(X_t)) \dots (1 - \underline{q}(X_0))] \\ &= E_{\bar{P}_\nu^*} [C(X_{-1}, X_0)(1 - \underline{q}(X_{-1})) \dots (1 - \underline{q}(X_{-t}))] \\ &= E_{\bar{P}_\nu^*} \left[ C(X_{-1}, X_0) \prod_{i=1}^t (1 - \underline{q}(X_{-i})) \right]. \end{aligned}$$

By the conditions of the lemma,  $E[C(X_{-1}, X_0)^l] < \infty$ , so by Jensen's inequality (and the fact that  $1/l' \leq 1$ , where  $1/l + 1/l' = 1$ ), it suffices to show that

$$\lim_{T \rightarrow \infty} T^{-1} \sum_{t=1}^T E_{\bar{P}_\nu^*} \left[ \prod_{i=0}^{t-1} (1 - \underline{q}(X_i))^{l'} \right] = 0.$$

Note that, for any  $1 \leq m \leq T$ ,

$$T^{-1} \sum_{t=1}^T E_{\bar{P}_\nu^*} \left[ \prod_{i=0}^{t-1} (1 - \underline{q}(X_i))^{l'} \right] \leq \frac{m}{T} + T^{-1} \sum_{t=m+1}^T E_{\bar{P}_\nu^*} \left[ \prod_{i=0}^{t-1} (1 - \underline{q}(X_i))^{l'} \right].$$

We now examine the term  $E_{\bar{P}_\nu^*} \left[ \prod_{i=0}^{t-1} (1 - \underline{q}(X_i))^{l'} \right]$  for any  $t \equiv qk$  and some positive integers  $q$  and  $k$ ; if this does not hold,  $t$  in the product is simply replaced with  $qk(t)$ , where  $k(t)$  is the largest integer such that  $t \geq qk$ . Letting  $M_j \equiv \{jq + 1, \dots, (j+1)q\}$ , it follows that  $\{1, \dots, t\} = \cup_{j=0}^{k-1} M_j$ . Thus,

$$\begin{aligned} E_{\bar{P}_\nu^*} \left[ \prod_{i=0}^{t-1} (1 - \underline{q}(X_i))^{l'} \right] &= E_{\bar{P}_\nu^*} \left[ \prod_{j=0}^{k-1} \prod_{i \in M_j} (1 - \underline{q}(X_i))^{l'} \right] \\ &\leq E_{\bar{P}_\nu^*} \left[ \prod_{j \in E_k} \prod_{i \in M_j} (1 - \underline{q}(X_i))^{l'} \right], \end{aligned}$$

where  $E_k \equiv \{0 \leq j \leq k-1 : j \text{ is even}\}$ . Observe that

$$\begin{aligned} E_{\bar{P}_\nu^*} \left[ \prod_{j \in E_k} \prod_{i \in M_j} (1 - \underline{q}(X_i))^{l'} \right] &= E_{\mathbb{P}} \left[ \left| \prod_{j \in E_k} \prod_{i \in M_j} (1 - \underline{q}(X_i))^{l'} - \prod_{j \in E_k} \prod_{i \in M_j} (1 - \underline{q}(\bar{X}_i))^{l'} \right| \right] \\ &\quad + E \left[ \prod_{j \in E_k} \prod_{i \in M_j} (1 - \underline{q}(\bar{X}_i))^{l'} \right] \\ &\leq \mathbb{P}(\exists j \in E_k : \bar{U}_j \neq U_j) + E \left[ \prod_{j \in E_k} \prod_{i \in M_j} (1 - \underline{q}(\bar{X}_i))^{l'} \right] \\ &\leq \beta_q k + E \left[ \prod_{j \in E_k} \prod_{i \in M_j} (1 - \underline{q}(\bar{X}_i))^{l'} \right]. \end{aligned}$$

Because of the properties of the process  $(\bar{U}_j)_{j \text{ even}}$ ,

$$\begin{aligned} E \left[ \prod_{j \in E_k} \prod_{i \in M_j} (1 - \underline{q}(\bar{X}_i))^{l'} \right] &\leq \prod_{j \in E_k} E \left[ (1 - \underline{q}(\bar{X}_{jq+1}))^{l'} \right] \\ &= \prod_{j \in E_k} E_{\bar{P}_\nu^*} \left[ (1 - \underline{q}(X_{jq+1}))^{l'} \right] \\ &= \left( E_{\bar{P}_\nu^*} \left[ (1 - \underline{q}(X_1))^{l'} \right] \right)^{(k-1)/2}. \end{aligned}$$

Noting that  $t = qk$ , these results imply that

$$E_{\bar{P}_\nu^*} \left[ \prod_{i=0}^{t-1} (1 - \underline{q}(X_i))^{l'} \right] \leq \beta_q t/q + \left( E_{\bar{P}_\nu^*} \left[ (1 - \underline{q}(X_1))^{l'} \right] \right)^{(t/q-1)/2}.$$

Therefore, for any  $1 \leq m \leq T$ ,

$$T^{-1} \sum_{t=1}^T E_{\bar{P}_\nu^*} \left[ \prod_{i=0}^{t-1} (1 - \underline{q}(X_i))^{l'} \right] \leq \frac{m}{T} + T^{-1} \sum_{t=m+1}^T \left\{ \beta_q t/q + \left( E_{\bar{P}_\nu^*} \left[ (1 - \underline{q}(X_1))^{l'} \right] \right)^{(t/q-1)/2} \right\}.$$

By Lemma 1,  $\beta_q = \exp\{q \log \gamma\}$ . This fact and the condition  $E_{\bar{P}_\nu^*} \left[ (1 - \underline{q}(X_1))^{l'} \right] < 1$  imply that we can take, for instance,  $q \equiv t^{1/2}$  and  $m = \sqrt{T}$ , so that the RHS vanishes as  $T$  diverges.

(2) We wish to show that

$$\sum_{t=1}^{\infty} \left( E_{\bar{P}_\nu^*} \left[ \prod_{i=0}^{t-1} (1 - \underline{q}(X_i))^{\frac{2a}{1-a}} \right] \right)^{p \frac{1-a}{2a}} < \infty.$$

By our previous calculations,

$$E_{\bar{P}_\nu^*} \left[ \prod_{i=0}^{t-1} (1 - \underline{q}(X_i))^{\frac{2a}{1-a}} \right] \leq \beta_q t/q + \left( E_{\bar{P}_\nu^*} \left[ (1 - \underline{q}(X_1))^{\frac{2a}{1-a}} \right] \right)^{(t/q-1)/2};$$

thus, it suffices to show that there exists a choice  $t \mapsto q(t)$  such that  $\sum_{t=1}^{\infty} \left\{ \beta_{q(t)} t/q(t) + \left( E_{\bar{P}_\nu^*} \left[ (1 - \underline{q}(X_1))^{\frac{2a}{1-a}} \right] \right)^{(t/q(t)-1)/2} \right\} < \infty$ . Let  $q(t) = \sqrt{t}$ ; then as  $E_{\bar{P}_\nu^*} \left[ (1 - \underline{q}(X_1))^{\frac{2a}{1-a}} \right] < 1$  by assumption, it follows that the second term in the sum is finite. Since  $t \mapsto \beta_{q(t)} t/q(t) = \exp\{\sqrt{t} \log \gamma\} \sqrt{t}$  is summable (note that  $\log \gamma < 0$ ), the same holds true for the first term.

□



## SM.5 Proofs and Results for Example 4

In what follows  $e_{\max}(M)$  and  $e_{\min}(M)$  denote the maximal and minimal eigenvalues of a matrix  $M$ .

**Lemma 14.** *Assumption 2 holds.*

*Proof.* For each  $s \in \mathbb{S}$ , we apply Theorem 3.3 in Douc et al. [2004a]. To do so, we first verify their Assumptions 3.3 and 3.4. In our case,  $\epsilon \sim N(0, \Sigma(s))$ , so their Assumption 3.3 is satisfied for any  $z_0$  and  $\gamma_0 = 1$ . In their notation,  $g(x) \equiv \mu(s) + \Phi^\top x$ . Observe that  $\|g(x)\| \leq \|\mu(s)\| + e_{\max}(\Phi\Phi^\top)\|x\|$ . By assumption,  $e_{\max}(\Phi\Phi^\top) \equiv \gamma < 1$  and

$$\|\mu(s)\| \leq (1 - e_{\max}(\Phi\Phi^\top))\|x\|(1 - \|x\|^{-0.5})$$

for all  $x$  such that  $\|x\| \geq R_0$ . Such an  $R_0$  exists because  $\|\mu(s)\|$  is bounded and  $e_{\max}(\Phi\Phi^\top) < 1$ . This choice ensures that  $\|g(x)\| \leq \|x\|(1 - \|x\|^{0.5})$ , which, in turn, ensures the validity of their Assumption 3.4 with  $r = 1$  and  $\rho = 0.5$ . By their Theorem 3.3, Assumption 2(i) holds.

Assumption 2(ii),(iii) is satisfied because  $\inf_{x \in A} P_*(s, x, C) \geq \int_C \inf_{x \in A} p(x, a) da$  and, since  $A$  is bounded, it follows that  $\inf_{x \in A} p(x, a) \geq \exp\{\tilde{D} + (x')^T \tilde{F}a + \tilde{G}a\}$ , so the RHS plays the role of the measure  $\varpi$ , which clearly is such that  $\varpi(A) > 0$ .  $\square$

Let  $\bar{\kappa} \equiv \max_{s \in \mathbb{S}} \sup_{\Sigma(s) \in \Theta} e_{\max}(\Sigma(s))$  and  $\underline{\kappa} \equiv \min_{s \in \mathbb{S}} \inf_{\Sigma(s) \in \Theta} e_{\min}(\Sigma(s))$ ;  $\bar{m} = \max_{s \in \mathbb{S}} \sup_{\mu(s) \in \Theta} \|\mu(s)\|$ ;  $\bar{M} = \sup_{\Phi \in \Theta} e_{\max}(\Phi\Phi^\top)$  and  $\underline{M} = \min_{\Phi \in \Theta} e_{\min}(\Phi\Phi^\top)$ ;  $\bar{\kappa}_* \equiv \max_{s \in \mathbb{S}} e_{\max}(\Sigma_*(s))$  and  $\underline{\kappa}_* \equiv \min_{s \in \mathbb{S}} e_{\min}(\Sigma_*(s))$ ;  $\bar{m}_* = \max_{s \in \mathbb{S}} \|\mu_*(s)\|$ ;  $\bar{M}_* = e_{\max}(\Phi_*\Phi_*^\top)$  and  $\underline{M}_* = e_{\min}(\Phi_*\Phi_*^\top)$ . By the assumptions in the text,  $\bar{\kappa}, \bar{\kappa}_*, \underline{\kappa}, \underline{\kappa}_*, \underline{M}, \bar{M}, \underline{M}_*, \bar{M}_*$  are all in  $(0, \infty)$ .

**Lemma 15.** *There exists a  $C \in [1, \infty)$  such that, for any  $(x, y)$  and any  $s$ ,*

$$\begin{aligned} & f_N((y - \Phi^\top x - \mu(s))\Sigma^{-1/2}(s)) \\ & \leq C \exp\{-0.5\underline{\kappa}(\|y\|^2 + \underline{M}\|x\|^2 - 2\sqrt{\bar{M}}\|x\|\|y\|) + \underline{\kappa}(\|y\| + \sqrt{\bar{M}}\|x\|)\bar{m}\} \end{aligned}$$

and

$$\begin{aligned} & f_N((y - \Phi^\top x - \mu(s))\Sigma^{-1/2}(s)) \\ & \geq C^{-1} \exp\{-0.5\bar{\kappa}(\|y\|^2 + \bar{M}\|x\|^2 + 2\sqrt{\bar{M}}\|x\|\|y\|) - \bar{\kappa}(\|y\| + \sqrt{\bar{M}}\|x\|)\bar{m}\}. \end{aligned}$$

*An analogous bound holds for  $f_N((y - \Phi_*^\top x - \mu_*(s))\Sigma_*^{-1/2}(s))$ .*

*Proof.* We have  $(a, s, b) \mapsto p_\theta(a, s, b) \asymp \phi((b - \mu(s) - \Phi^\top a)\Sigma^{-1/2}(s))$ . Thus, there exists a  $C \in [1, \infty)$  such that, for any  $u \in \mathbb{R}^p$ ,

$$\begin{aligned} f_{\mathcal{N}}((u - \mu(s))\Sigma^{-1/2}(s)) &\leq C \exp\{-0.5\underline{\kappa}\|u - \mu(s)\|^2\}, \\ f_{\mathcal{N}}((u - \mu(s))\Sigma^{-1/2}(s)) &\geq C^{-1} \exp\{-0.5\bar{\kappa}\|u - \mu(s)\|^2\}, \end{aligned}$$

where  $\bar{\kappa} \in (0, \infty)$  and  $\underline{\kappa} \in (0, \infty)$  are the largest and smallest eigenvalues within the collection  $(\Sigma(s))_{s \in \mathbb{S}}$  in  $\Theta$ .

Observe that

$$\|x - y\|^2 \leq \|x\|^2 + \|y\|^2 + 2\|x\|\|y\|,$$

and, if  $\|x\| > \|y\|$ ,

$$\|x - y\|^2 \geq (\|x\| - \|y\|)^2 = \|x\|^2 + \|y\|^2 - 2\|x\|\|y\|,$$

and the same for  $\|x\| < \|y\|$ .

Hence,

$$\begin{aligned} f_{\mathcal{N}}((u - \mu(s))\Sigma^{-1/2}(s)) &\leq C \exp\{-0.5\underline{\kappa}(\|u\|^2 + \|\mu(s)\|^2 - 2\|u\|\|\mu(s)\|)\} \\ &\leq C \exp\{-0.5\underline{\kappa}\|u\|^2 + \underline{\kappa}\|u\|\bar{m}\} \end{aligned}$$

and

$$\begin{aligned} f_{\mathcal{N}}((u - \mu(s))\Sigma^{-1/2}(s)) &\geq C^{-1} \exp\{-0.5\bar{\kappa}(\|u\|^2 + \|\mu(s)\|^2 + 2\|u\|\|\mu(s)\|)\} \\ &\geq C^{-1} \exp\{-0.5\bar{\kappa}\|u\|^2 - 0.5\bar{\kappa}m^2 - \bar{\kappa}\|u\|m\} \\ &\equiv C^{-1} \exp\{-0.5\bar{\kappa}\|u\|^2 - \bar{\kappa}\|u\|m\}, \end{aligned}$$

where  $\bar{m} = \sup_{\mu(\cdot) \in \Theta} \max_{s \in \mathbb{S}} \|\mu(s)\|$ .

Observe that

$$\begin{aligned} \|y - \Phi^\top x\|^2 &\leq \|y\|^2 + \|\Phi^\top x\|^2 + 2\|\Phi^\top x\|\|y\| \\ &\leq \|y\|^2 + \bar{M}\|x\|^2 + 2\sqrt{\bar{M}}\|x\|\|y\| \end{aligned}$$

and

$$\begin{aligned} \|y - \Phi^\top x\|^2 &\geq \|y\|^2 + \|\Phi^\top x\|^2 - 2\|\Phi^\top x\|\|y\| \\ &\geq \|y\|^2 + \underline{M}\|x\|^2 - 2\sqrt{\underline{M}}\|x\|\|y\|, \end{aligned}$$

where  $\bar{M} = \sup_{\Phi \in \Theta} e_{\max}(\Phi\Phi^\top)$  and  $\underline{M} = \min_{\Phi \in \Theta} e_{\min}(\Phi\Phi^\top)$ , which, by our conditions, are finite and positive. Hence,

$$\begin{aligned} f_{\mathcal{N}}((y - \Phi^\top x - \mu(s))\Sigma^{-1/2}(s)) \\ \leq C \exp\{-0.5\underline{\kappa}(\|y\|^2 + \underline{M}\|x\|^2 - 2\sqrt{\underline{M}}\|x\|\|y\|) + \underline{\kappa}(\|y\| + \sqrt{\underline{M}}\|x\|)\bar{m}\} \end{aligned}$$

and

$$\begin{aligned} & f_{\mathcal{N}}((y - \Phi^\top x - \mu(s))\Sigma^{-1/2}(s)) \\ & \geq C^{-1} \exp\{-0.5\bar{\kappa}(\|y\|^2 + \bar{M}\|x\|^2 + 2\sqrt{\bar{M}}\|x\|\|y\|) - \bar{\kappa}(\|y\| + \sqrt{\bar{M}}\|x\|)\bar{m}\}. \end{aligned}$$

□

**Lemma 16.** *Suppose there exists a  $l \geq 1$  such that  $l\bar{\kappa} < l\underline{\kappa} + \underline{\kappa}_*$  and  $l\bar{\kappa}\bar{M} < l\underline{\kappa}\bar{M} + \underline{\kappa}_*$ . Then,*

$$E_{\bar{P}_\nu^*}[\exp\{-0.5l((b_1 - a_1)\|Y\|^2 + (b_2 - a_2)\|X\|^2 - 2(b_3 + a_3)\|X\|\|Y\|) + l(b_4 + a_4)\|Y\| + l(b_5 + a_5)\|X\|\}] < \infty,$$

where

$$\begin{aligned} a_1 &= \bar{\kappa}, \quad b_1 = \underline{\kappa}, \\ a_2 &= \bar{\kappa}\bar{M}, \quad b_2 = \underline{\kappa}\bar{M}, \\ a_3 &= \bar{\kappa}\sqrt{\bar{M}}, \quad b_3 = \underline{\kappa}\sqrt{\bar{M}}, \\ a_4 &= \underline{\kappa}, \quad b_4 = \bar{\kappa}, \\ a_5 &= \underline{\kappa}\sqrt{\bar{M}}, \quad b_5 = \bar{\kappa}\sqrt{\bar{M}}. \end{aligned}$$

**Remark 5.** Before going to the proof, we discuss the conditions in the lemma. They basically require that the “spread” of the eigenvalues of the matrices  $\Sigma(\cdot)$  and  $\Phi\Phi^*$  is not too large relative to the eigenvalues in  $\Sigma_*(\cdot)$ . This condition comes naturally since we are essentially requiring that the ratio of two exponential functions is integrable with respect to a Gaussian measure. For instance, if  $\Sigma(\cdot)$ ,  $\Sigma_*(\cdot)$  and  $\Phi\Phi^\top$ ,  $\Phi_*\Phi_*^\top$  are matrices with eigenvalues bounded between  $0 < a$  and  $a + \Delta$ , then sufficient conditions are given by  $l\Delta < a$  and  $l(2a\Delta + (\Delta)^2) < a$ , which is equivalent to  $\frac{l\Delta^2}{1-2l\Delta} < a$ .  $\triangle$

*Proof.* It is enough to show that

$$\begin{aligned} E_{\bar{P}_\nu^*}[T_1(Y)] &\equiv E_{\bar{P}_\nu^*}[\exp\{-0.5l(b_1 - a_1)\|Y\|^2 + l(b_4 + a_4)\|Y\|\}] < \infty, \\ E_{\bar{P}_\nu^*}[T_2(X)] &\equiv E_{\bar{P}_\nu^*}[\exp\{-0.5l(b_2 - a_2)\|X\|^2 + l(b_5 + a_5)\|X\|\}] < \infty, \\ E_{\bar{P}_\nu^*}[T_3(X, Y)] &\equiv E_{\bar{P}_\nu^*}[\exp\{l(b_3 + a_3)\|X\|\|Y\|\}] < \infty. \end{aligned}$$

For any  $d \in \{1, 2\}$ , suppose there exists  $\varphi$  such that  $\int T_d(b)p_*(x, s, b)db \leq \varphi(x)T_d(x)$  for any  $x$ , and, for any  $\gamma > 0$ ,  $\{x : \varphi(x) \geq \gamma\}$  is either empty or

compact. Then, for any  $\gamma > 0$ ,

$$\begin{aligned} \int T_d(x) \nu(dx) &= \int 1\{\varphi(x) \leq \gamma\} T_d(x) \nu(dx) + \int 1\{\varphi(x) > \gamma\} T_d(x) \nu(dx) \\ &= \int \int 1\{\varphi(x) \leq \gamma\} T_d(b) p_*(x, s, b) db \nu(dx, ds) + \int 1\{\varphi(x) > \gamma\} T_d(x) \nu(dx) \\ &\leq \gamma \int T_d(x) \nu(dx) + \sup_{x: \varphi(x) \geq \gamma} T_d(x), \end{aligned}$$

where second line follows because  $\nu$  is the invariant probability distribution. Since  $\{x : \varphi(x) \geq \gamma\}$  is bounded and compact (if it is non-empty),  $\sup_{x: \varphi(x) \geq \gamma} T_d(x) \leq M < \infty$ . Choosing  $\gamma < 1$ , it follows that  $\int T_d(x) \nu(dx) \leq \frac{M}{1-\gamma} < \infty$ , as desired.

We now show that  $\int T_1(b) p_*(x, s, b) db \leq \varphi(x) T_1(x)$  for any  $x$ . To do this, note that

$$\int T_1(b) p_*(x, s, b) db \leq \int \exp\{0.5l(\bar{\kappa} - \underline{\kappa})\|y\|^2 + l(\bar{\kappa} + \underline{\kappa})\|y\|\} p_*(x, s, y) dy,$$

and, by Lemma 15,

$$\begin{aligned} f_{\mathcal{N}}((y - \Phi_*^\top x - \mu_*(s)) \Sigma_*^{-1/2}(s)) &\leq \exp\{-0.5\underline{\kappa}_*(\|y\|^2 - 2\sqrt{\bar{M}_*}\|x\|\|y\|) + \underline{\kappa}_*\|y\|\bar{m}_*\} \\ &\quad \times B_*(x), \end{aligned}$$

with  $B_*(x) \equiv C_* \exp\{-0.5\underline{\kappa}_*\bar{M}_*\|x\|^2 + \underline{\kappa}_*\sqrt{\bar{M}_*}\|x\|\bar{m}_*\}$ . Therefore,

$$\begin{aligned} &\frac{\int T_1(b) p_*(x, s, y) dy}{B_*(x)} \\ &\leq \int \exp\{0.5(\bar{\kappa} - \underline{\kappa})\|y\|^2 + l(\bar{\kappa} + \underline{\kappa})\|y\|\} \exp\{-0.5\underline{\kappa}_*(\|y\|^2 - 2\sqrt{\bar{M}_*}\|x\|\|y\|) + \underline{\kappa}_*\bar{m}_*\|y\|\} dy \\ &= \int \exp\{0.5(l(\bar{\kappa} - \underline{\kappa}) - \underline{\kappa}_*)\|y\|^2 + \underline{\kappa}_*\sqrt{\bar{M}_*}\|x\|\|y\| + (\underline{\kappa}_*\bar{m}_* + l(\bar{\kappa} + \underline{\kappa}))\|y\|\} dy. \end{aligned}$$

By our conditions,  $l(\bar{\kappa} - \underline{\kappa}) - \underline{\kappa}_* < 0$ . Hence, the expression above is an integral of an exponential function of a quadratic form with negative leading coefficient, and is thus finite. Moreover, after some algebra, there exists a finite constant  $C$ , such that  $\int T_1(b) p_*(x, s, y) dy \leq C B_*(x) \exp\{\underline{\kappa}_*\sqrt{\bar{M}_*}\|x\|/(-(l(\bar{\kappa} - \underline{\kappa}) - \underline{\kappa}_*))\}$ ; we re-define the RHS as  $C B_*(x) \exp\{D \underline{\kappa}_*\sqrt{\bar{M}_*}\|x\|\}$  with  $D > 0$ . Therefore, the result holds with

$$x \mapsto \varphi(x) \equiv \exp\{-0.5(\underline{\kappa}_*\bar{M}_* + l(\bar{\kappa} - \underline{\kappa}))\|x\|^2 + (\underline{\kappa}_*\sqrt{\bar{M}_*}(D + \bar{m}_*) + l(b_4 + a_4))\|x\|\}.$$

As the coefficient  $\|x\|^2$  is  $-0.5(\underline{\kappa}_* \underline{M}_* + l(\bar{\kappa} - \underline{\kappa}))$ , which is negative, the function satisfies the required conditions.

The case for  $d = 2$  is analogous and is thus omitted; for this case, we use the restriction that  $l\bar{\kappa}\bar{M} < l\underline{\kappa}\underline{M} + \underline{\kappa}_*$ , instead of  $l(\bar{\kappa} - \underline{\kappa}) - \underline{\kappa}_* < 0$ .

Finally, observe that, if  $\int T_3(x, y)p_*(x, s, y)\nu(dx, ds)dy \leq C \int \exp\{c_1\|x\|\}\nu(dx)$  for some  $c_1 < \infty$ , then we can follow the same approach as before to show that  $\int T_3(x, y)p_*(x, s, y)\nu(dx, ds)dy < \infty$ . Observe that, by the same calculations as before,

$$\begin{aligned} & \int T_3(x, y)p_*(x, s, y)dy \\ & \leq B_*(x) \int \exp\{l(a_3 + b_3)\|x\|\|y\| - 0.5\underline{\kappa}(\|y\|^2 - 2\sqrt{\bar{M}}\|x\|\|y\|) + \underline{\kappa}\|y\|\bar{m}\}dy \\ & \leq CB_*(x) \exp\{(l(a_3 + b_3) + \underline{\kappa}\sqrt{\bar{M}})\|x\|\}. \end{aligned}$$

The RHS is of the form  $C \exp\{c_1\|x\|\}$ , as desired.  $\square$

## SM.6 Pseudo-True Parameter Set

In this Section, we present an example in which it is possible to characterize the pseudo-true parameter set  $\Theta_*$  of Theorem 1. The example shows that for a subclass of the models in Example 2, namely hidden Markov models with covariate-dependent transition probabilities,  $\Theta_*$  can be characterized when the (misspecified) model is a simple mixture. Specifically, we show that, even though the mixture model misspecifies the dependence structure of the hidden state – and thus of the overall system – parameters related to the outcome equation are consistently estimated.

**Example 6.** Suppose the true model is a version of the hidden Markov model of Example 1 with

$$\begin{aligned} Y_t &= \mu^*(S_t) + \sigma^*(S_t)U_{1,t}, \\ Z_t &= \mu_2^* + \psi^*Z_{t-1} + \sigma_2^*U_{2,t}, \end{aligned}$$

where  $((U_{1,t}, U_{2,t}))_t$  are i.i.d. with zero mean and covariance matrix  $\Sigma = \begin{bmatrix} 1 & \rho^* \\ \rho^* & 1 \end{bmatrix}$ , and  $S_t \sim Q_*(Z_{t-1}, S_{t-1}, \cdot)$ . The only additional requirement is that the process  $(S_t, Z_t)_t$  is stationary with invariant distribution  $\nu_{ZS}$ . Since the focus of the example is misspecification of the Markov transition mechanism, we also assume that the process for  $(Z_t)_t$  is correctly specified. The researcher,

however, postulates a mixture model (with i.i.d. regimes) of the form

$$\begin{aligned} Y_t &= \mu(S_t) + \sigma(S_t)\varepsilon_t, \\ S_t &\sim Q_{\bar{\vartheta}}, \end{aligned}$$

where  $Q_{\bar{\vartheta}}$  is parameterized as  $Q_{\bar{\vartheta}}(s) = \bar{\vartheta}_s$  for  $s \in \{0, 1\}$ , with  $\bar{\vartheta}_s \in (0, 1)$  (see, e.g., [McLachlan and Peel \[2000\]](#)).<sup>23</sup>

The following proposition shows that, although the mixture model is misspecified, ML estimates correctly the parameters of the outcome equation.

**Proposition 1.** *The choice  $\mu = \mu_*$ ,  $\sigma = \sigma_*$ , and  $\bar{\vartheta}$  such that  $Q_{\bar{\vartheta}} = E_{\nu_{ZS}}[Q_*(Z, S, \cdot)]$  is a pseudo-true parameter, i.e., minimizes  $\theta \mapsto E_{\bar{P}_*^\nu} \left[ \log \sum_{s \in \mathbb{S}} Q_{\bar{\vartheta}}(s) \sigma^{-1}(s) f\left(\frac{Y_1 - \mu(s) - \phi Y_0}{\sigma(s)}\right) \right]$ , where  $f$  is the probability density function of  $U_{1,1}$ .*

*Proof.* Throughout the proof, we omit the sub-index "1" from the parameters in the outcome equation to ease the notational burden.

Observe that  $H^*$  is proportional to

$$\theta \mapsto - \int \log \frac{\sum_{s \in \mathbb{S}} Q_{\bar{\vartheta}}(s) \sigma(s)^{-1} f((y - \mu(s))/\sigma(s))}{f_*(y)} f_*(y) dy,$$

where

$$y \mapsto f_*(y) = \sum_{s \in \mathbb{S}} Pr_*(S_1 = s) \sigma_*(s)^{-1} f((y - \mu_*(s))/\sigma_*(s)),$$

and  $Pr_*$  is the true probability over the hidden state, given by

$$s \mapsto Pr_*(S_1 = s) = \int \sum_{s_0 \in \mathbb{S}} Q_*(z, s_0, s) \nu_{ZS}(dz, ds_0).$$

It is well-known that the minimizers of this function are all  $\theta$  such that

$$\sum_{s \in \mathbb{S}} Q_{\bar{\vartheta}}(s) \sigma(s)^{-1} f((\cdot - \mu(s))/\sigma(s)) = f_*(\cdot).$$

It is straightforward to check that by choosing  $\bar{\vartheta}$  such that  $Q_{\bar{\vartheta}}(\cdot) = Pr_*(S_1 = \cdot)$  and  $\mu = \mu_*$  and  $\sigma = \sigma_*$ , the equality holds.  $\square$

<sup>23</sup>For our results to go through, it suffices that a weaker condition holds, namely there exists a  $\bar{\vartheta}$  such that  $s \mapsto Q_{\bar{\vartheta}}(s) = E_{\nu_{ZS}}[Q_*(Z, S, s)]$ . The idea is that the parametrization is "rich enough" to mimic the average behavior of the true transitions.

**Remark 6.** This result may be of practical interest because it indicates that dependence of the regimes in a hidden Markov model may be ignored if the parameters of interest are only those of the outcome equation. Note, however, that the result does not hold in cases where the right-hand side of the outcome equation contains lagged values of the dependent variable. In this case, the misspecification of the transition function will affect estimation of all parameters. [Carter and Steigerwald \[2012\]](#) make a similar point in the context of a simple data-generating process with time-invariant transition probabilities.  $\triangle$

## SM.7 Proofs of Supplemental Lemmas in Appendix A.2

*Proof of Lemma 5.* For any  $a, b$  in  $\mathbb{S}$ ,

$$\begin{aligned}\bar{P}_\theta^\nu \left( S_{l+1} = b \mid S_l = a, X_{-m}^j \right) &= \frac{\bar{P}_\theta^\nu \left( S_{l+1} = b, S_l = a, X_{-m}^j \right)}{\sum_{c \in \mathbb{S}} \bar{P}_\theta^\nu \left( S_{l+1} = c, S_l = a, X_{-m}^j \right)} \\ &= \frac{\bar{P}_\theta^\nu \left( X_{l+1}^j \mid S_{l+1} = b, S_l = a, X_{-m}^l \right) Q_\theta(X_l, a, b)}{\sum_{c \in \mathbb{S}} \bar{P}_\theta^\nu \left( X_{l+1}^j \mid S_{l+1} = c, S_l = a, X_{-m}^l \right) Q_\theta(X_l, a, c)}.\end{aligned}$$

The expression  $\bar{P}_\theta^\nu \left( X_{l+1}^j \mid S_{l+1} = b, S_l = a, X_{-m}^l \right)$  equals  $\bar{P}_\theta^\nu \left( X_{l+1}^j \mid S_{l+1} = b, X_l \right)$  because of the Markov property. The latter probability depends on the transitions of  $X_{t+1}$  given  $(X_t, S_{t+1})$  and  $S_{t+1}$  given  $(X_t, S_t)$  for each  $t \geq l+1$ . Since these are the same for the process with  $i=1$  and  $i=2$  and the "original" process"  $(S_t, X_t)_{t=-m}^\infty$ , it follows that the last line of the previous display equals  $Pr_\theta \left( \eta_{1,l+1} = b \mid \eta_{1,l} = a, X_{-m}^j \right) = Pr_\theta \left( \eta_{2,l+1} = b \mid \eta_{2,l} = a, X_{-m}^j \right)$ , as desired.  $\square$

*Proof of Lemma 6.* Throughout this proof, we omit the dependence on  $\theta$  in the probability terms and on other quantities. For any  $a, c$  in  $\mathbb{S}$ ,

$$\begin{aligned}& \left\| \Pr \left( \eta_{1,l+1} = \cdot \mid \eta_{1,l} = a, X_{-m}^j \right) - \Pr \left( \eta_{2,l+1} = \cdot \mid \eta_{1,l} = c, X_{-m}^j \right) \right\|_1 \\ & \leq \left\| \Pr \left( \eta_{1,l+1} = \cdot, v_{1,l} = 0 \mid \eta_{1,l} = a, X_{-m}^j \right) - \Pr \left( \eta_{2,l+1} = \cdot, v_{2,l} = 0 \mid \eta_{2,l} = c, X_{-m}^j \right) \right\|_1 \\ & \quad + \left\| \Pr \left( \eta_{1,l+1} = \cdot, v_{1,l} = 1 \mid \eta_{1,l} = a, X_{-m}^j \right) - \Pr \left( \eta_{2,l+1} = \cdot, v_{2,l} = 1 \mid \eta_{2,l} = c, X_{-m}^j \right) \right\|_1 \\ & \equiv Term_1 + Term_2\end{aligned}$$

To bound the second term, note that

$$\begin{aligned} \Pr\left(\eta_{1,l+1} = \cdot, v_{1,l} = 1 \mid \eta_{1,l} = a, X_{-m}^j\right) &= \Pr\left(\eta_{1,l+1} = \cdot \mid v_{1,l} = 1, \eta_{1,l} = a, X_{-m}^j\right) \\ &\quad \times \Pr\left(v_{1,l} = 1 \mid \eta_{1,l} = a, X_{-m}^j\right). \end{aligned}$$

It follows that  $\Pr\left(v_{1,l} = 1 \mid \eta_{1,l} = a, X_{-m}^j\right) = \underline{q}(X_l)$  because, given  $X_{-m}^j$ ,  $v_{1,l}$  is drawn independently according to a probability function that only depends on  $X_l$  (in particular, it does not depend on  $\eta_{1,l}$ ), and is given by  $\underline{q}(X_l)$ . By some algebra, the Markov property, and the fact that, given  $v_{1,l} = 1$  and  $X_{-m}^j$ , the random variable  $\eta_{1,l+1}$  is independent of its past, it follows that  $a \mapsto \Pr\left(\eta_{1,l+1} = \cdot \mid v_{1,l} = 1, \eta_{1,l} = a, X_{-m}^j\right)$  is constant (i.e., does not depend on  $\eta_{1,l} = a$ ). Thus,  $a \mapsto \Pr\left(\eta_{1,l+1} = \cdot, v_{1,l} = 1 \mid \eta_{1,l} = a, X_{-m}^j\right)$  is constant (i.e., does not depend on the value of  $a$ ); since one can obtain the exact result for  $c \mapsto \Pr\left(\eta_{2,l+1} = \cdot, v_{2,l} = 1 \mid \eta_{2,l} = c, X_{-m}^j\right)$  and, moreover, the laws for  $i = 1$  and  $i = 2$  coincide (see the proof of Lemma 5), it follows that  $Term_2 = 0$ .

To bound  $Term_1$ , it follows that from the previous arguments that

$$\begin{aligned} Term_1 &= \sum_{s \in \mathbb{S}} \left| \Pr\left(\eta_{1,l+1} = s \mid v_{1,l} = 0, \eta_{1,l} = a, X_{-m}^j\right) - \Pr\left(\eta_{2,l+1} = s \mid v_{2,l} = 0, \eta_{2,l} = c, X_{-m}^j\right) \right| \\ &\quad \times (1 - \underline{q}(X_l)) \\ &\leq 2(1 - \underline{q}(X_l)), \end{aligned}$$

and thus the desired result follows.  $\square$

## SM.8 Proofs of Supplementary Lemmas in Appendix A.3

### SM.8.1 Proofs of Lemmas 7 and 8

In this section, we provide the proofs of Lemmas 7 and 8. To do this, we use a series of lemmas which we state below (their proofs are relegated to the end of this section).

Henceforth, for any  $j \geq m$ , let

$$\varrho(j, m) \equiv \left( E_{\bar{P}^\nu_*} \left[ \prod_{i=m}^j (1 - \underline{q}(X_i))^{\frac{2a}{1-a}} \right] \right)^{\frac{1-a}{2a}}, \quad (15)$$

where the constant  $a$  is the same as in Assumption 7. We also introduce the following notation: for any  $\theta \in \Theta$ ,  $(x', x, s) \mapsto \Gamma(x'|x, s; \theta) \equiv \nabla_\theta \log p_\theta(x, s, x')$



and  $(s, x, s) \mapsto \Lambda(s'|s, x; \theta) \equiv \nabla_\theta \log Q_\theta(x, s, s')$ . Furthermore, for any  $k \geq n$  and any  $l \geq m$ , let

$$\Phi_\theta(k, n, l, m) \equiv E_{\bar{P}_\theta^\nu} \left[ \sum_{j=n}^k \Gamma(X_j | X_{j-1}, S_j; \theta) \mid X_m^l \right]$$

$$\text{and } \Psi_\theta(k, n, l, m) \equiv E_{\bar{P}_\theta^\nu} \left[ \sum_{j=n}^k \Lambda(S_j | S_{j-1}, X_{j-1}; \theta) \mid X_m^l \right].$$

To state the first lemma, for any  $k, T$  and  $X_{k-T}^k$  and any  $\theta$ , let

$$\begin{aligned} \Delta_{k, k-T}(\theta)(X_{k-T}^k) &\equiv \Phi_\theta(k-1, k-T, k, k-T) + \Psi_\theta(k-1, k-T-1, k, k-T) \\ &\quad - \Phi_\theta(k-1, k-T, k-1, k-T) - \Psi_\theta(k-1, k-T-1, k-1, k-T) \\ &\quad + \Phi_\theta(k, k, k, k-T) + \Psi_\theta(k, k, k, k-T) \\ &= \sum_{j=k-T}^{k-1} E_{\bar{P}_\theta^\nu} \left[ \Gamma(X_j | X_{j-1}, S_j; \theta) \mid X_{k-T}^k \right] - E_{\bar{P}_\theta^\nu} \left[ \Gamma(X_j | X_{j-1}, S_j; \theta) \mid X_{k-T}^{k-1} \right] \\ &\quad + \sum_{j=k-T}^{k-1} E_{\bar{P}_\theta^\nu} \left[ \Lambda(S_l | S_{l-1}, X_{l-1}; \theta) \mid X_{k-T}^k \right] - E_{\bar{P}_\theta^\nu} \left[ \Lambda(S_l | S_{l-1}, X_{l-1}; \theta) \mid X_{k-T}^{k-1} \right] \\ &\quad + E_{\bar{P}_\theta^\nu} \left[ \Gamma(X_k | X_{k-1}, S_k; \theta) \mid X_{k-T}^k \right] + E_{\bar{P}_\theta^\nu} \left[ \Lambda(S_k | S_{k-1}, X_{k-1}; \theta) \mid X_{k-T}^k \right], \end{aligned} \tag{16}$$

The next lemma is analogous to the results in [Douc et al. \[2004b\]](#) and [Bickel et al. \[1998\]](#), and uses ideas of missing data models.

**Lemma 17.** *Suppose Assumption 6 holds. Then, for any  $k, T \geq 0$  and for any  $\theta \in \Theta$ ,*

$$\nabla_\theta \log p_k^\nu(X_k | X_{k-T}^{k-1}; \theta) = \Delta_{k, k-T}(\theta)(X_{k-T}^k)$$

*a.s.- $\bar{P}_*^\nu$ .*<sup>24</sup>

This Lemma characterizes the asymptotic behavior of the score functions; in particular, it shows that they are well-approximated by  $(\Delta_{t, -\infty}(\theta_*))_t$ , which is to be defined below, but at this stage is worth pointing out that it is stationary and ergodic; this last fact is established in Lemma 19 below.

**Lemma 18.** *Suppose Assumptions 1, 2, 6, 7(i) and 8 hold. Then:*

---

<sup>24</sup>When there is no risk of confusion, we will omit the dependence of  $\Delta_{k, k-T}$  on the data  $X_{k-T}^k$ .

(i) There exists a finite constant  $C > 0$  such that for any  $k$  and  $T \geq 0$ ,

$$\|\Delta_{k,k-T}(\theta_*) - \Delta_{k,-\infty}(\theta_*)\|_{L^2(\bar{P}_*^\nu)} \leq C \left( \max\left\{ \sum_{j=[k-T/2]}^{k-1} \varrho(j, k-T), \sum_{j=k-T}^{[k-T/2]-1} \varrho(k-1, j) \right\} \right);$$

(ii)

$$\lim_{T \rightarrow \infty} \left\| T^{-1/2} \sum_{t=0}^T \{\Delta_{t,-\infty}(\theta_*) - \nabla_\theta \log p_t^\nu(\cdot | \cdot, \theta_*)\} \right\|_{L^2(\bar{P}_*^\nu)} = 0.$$

**Lemma 19.** Suppose Assumptions 1, 2 and 5 hold. Then,  $(\Delta_{t,-\infty}(\theta_*))_{t=-\infty}^\infty$  is a stationary and ergodic  $L^2(\bar{P}_*^\nu)$  process (under  $\bar{P}_*^\nu$ ).

**Lemma 20.** Suppose Assumption 1 holds. Then, there exists a finite constant  $L > 0$  such that:

(i) For  $-m \leq j < k$  and any  $\theta \in \Theta$ ,

$$\left\| \bar{P}_\theta^\nu(S_j = \cdot | X_{-m}^k) - \bar{P}_\theta^\nu(S_j = \cdot | X_{-m}^{k-1}) \right\|_1 \leq L \prod_{i=j}^{k-1} (1 - \underline{q}(X_i)),$$

a.s.- $\bar{P}_*^\nu$ .

(ii) For  $-n \leq -m \leq j < k$  and any  $\theta \in \Theta$ ,

$$\left\| \bar{P}_\theta^\nu(S_j = \cdot | X_{-m}^k) - \bar{P}_\theta^\nu(S_j = \cdot | X_{-n}^{k-1}) \right\|_1 \leq L \prod_{i=-m}^j (1 - \underline{q}(X_i)),$$

a.s.- $\bar{P}_*^\nu$ .

*Proof of Lemma 7.* Follows directly from Lemmas 18 and 19.  $\square$

*Proof of Lemma 8.* Lemma 8 is analogous to Lemma 10 in Bickel et al. [1998]. The proof follows by their Lemma 9, which in turn holds by analogous steps to theirs and by invoking Lemma 20 (which is analogous to their Lemma 7).  $\square$

### SM.8.1.1 Proofs of Lemmas

Throughout this section, in cases where the expectations are taken with respect to  $\bar{P}_*^\nu$ , we omit the probability from the notation and simply use  $E[\cdot]$ .

*Proof of Lemma 17.* By [Louis, 1982, p. 227],

$$\begin{aligned}\nabla_\theta \log p_k^\nu(X_k | X_{k-T}^{k-1}; \theta) &= \nabla_\theta \log p_k^\nu(X_{k-T}^k; \theta) - \nabla_\theta \log p_{k-1}^\nu(X_{k-T}^{k-1}; \theta) \\ &= E_{\bar{P}_\theta^\nu}[\nabla_\theta \log p_k^\nu(X_{k-T}^k, S_{k-T}^k; \theta) | X_{k-T}^k] \\ &\quad - E_{\bar{P}_\theta^\nu}[\nabla_\theta \log p_{k-1}^\nu(X_{k-T}^{k-1}, S_{k-T}^{k-1}; \theta) | X_{k-T}^{k-1}].\end{aligned}$$

(Note that the expectation is with respect to  $S_{k-T}^k$ , which takes finitely many values; thus interchanging differentiation and integration is allowed).

Since  $p_k^\nu(X_{k-T}^k, S_{k-T}^k; \theta) = p_\theta(X_{k-1}, S_k, X_k) Q_\theta(X_{k-1}, S_{k-1}, S_k) \times p_k^\nu(X_{k-T}^{k-1}, S_{k-T}^{k-1}; \theta)$  (and an analogous result holds for  $p_k^\nu(X_{k-T}^{k-1}, S_{k-T}^{k-1}; \theta)$ ), it follows that

$$\begin{aligned}&\nabla_\theta \log p_k^\nu(X_k | X_{k-T}^{k-1}; \theta) \\ &= E_{\bar{P}_\theta^\nu} \left[ \sum_{j=k-T}^k \nabla_\theta \log p_\theta(X_{j-1}, S_j, X_j) | X_{k-T}^k \right] + E_{\bar{P}_\theta^\nu} \left[ \sum_{j=k-T}^k \nabla_\theta \log Q_\theta(X_{j-1}, S_{j-1}, S_j) | X_{k-T}^k \right] \\ &\quad - E_{\bar{P}_\theta^\nu} \left[ \sum_{j=k-T}^{k-1} \nabla_\theta \log p_\theta(X_{j-1}, S_j, X_j) | X_{k-T}^{k-1} \right] - E_{\bar{P}_\theta^\nu} \left[ \sum_{j=k-T}^{k-1} \nabla_\theta \log Q_\theta(X_{j-1}, S_{j-1}, S_j) | X_{k-T}^{k-1} \right] \\ &= E_{\bar{P}_\theta^\nu} \left[ \sum_{j=k-T}^{k-1} \Gamma(X_j | X_{j-1}, S_j; \theta) | X_{k-T}^k \right] + E_{\bar{P}_\theta^\nu} \left[ \sum_{l=k-T-1}^{k-1} \Lambda(S_l | S_{l-1}, X_{l-1}; \theta) | X_{k-T}^k \right] \\ &\quad - E_{\bar{P}_\theta^\nu} \left[ \sum_{j=k-T}^{k-1} \Gamma(X_j | X_{j-1}, S_j; \theta) | X_{k-T}^{k-1} \right] - E_{\bar{P}_\theta^\nu} \left[ \sum_{l=k-T-1}^{k-1} \Lambda(S_l | S_{l-1}, X_{l-1}; \theta) | X_{k-T}^{k-1} \right] \\ &\quad + E_{\bar{P}_\theta^\nu} \left[ \Gamma(X_k | X_{k-1}, S_k; \theta) | X_{k-T}^k \right] + E_{\bar{P}_\theta^\nu} \left[ \Lambda(S_k | S_{k-1}, X_{k-1}; \theta) | X_{k-T}^k \right].\end{aligned}$$

□

The proof of Lemma 18 requires the following lemma.

**Lemma 21.** *Suppose that Assumptions 1 and 7(i) hold. Then, there exists a finite constant  $C > 0$  such that:*

(i) *for any  $-n \leq -m \leq -m' \leq l \leq k$ ,*

$$\|\Phi_{\theta_*}(l, -m', k, -m) - \Phi_{\theta_*}(l, -m', k, -n)\|_{L^2(\bar{P}_*^\nu)} \leq C \left( \sum_{j=-m'}^l \varrho(j, -m) \right);$$

(ii) *for any  $-m \leq -m' < l \leq k-1$ ,*

$$\|\Phi_{\theta_*}(l, -m', k, -m) - \Phi_{\theta_*}(l, -m', k-1, -m)\|_{L^2(\bar{P}_*^\nu)} \leq C \left( \sum_{j=-m'}^l \varrho(k-1, j) \right);$$

(iii) for any  $-n \leq -m \leq -m' < l \leq k$ ,

$$\|\Psi_{\theta_*}(l, -m', k, -m) - \Psi_{\theta_*}(l, -m', k, -n)\|_{L^2(\bar{P}_*^\nu)} \leq C \left( \sum_{j=-m'}^l \varrho(j-1, -m) \right);$$

(iv) for any  $-m \leq -m' < l \leq k-1$ ,

$$\|\Psi_{\theta_*}(l, -m', k, -m) - \Psi_{\theta_*}(l, -m', k-1, -m)\|_{L^2(\bar{P}_*^\nu)} \leq C \left( \sum_{j=-m'}^l \varrho(k-1, j) \right).$$

*Proof of Lemma 21.* Throughout the proof we omit the dependence of  $E[\cdot]$  on  $\bar{P}_{\theta_*}^\nu$ . Also, let  $L$  denote the constant in Lemma 22.

**Part (i).** Observe that, for any  $j \leq k$ ,

$$\begin{aligned} & \left\| E \left[ \Gamma(X_j | X_{j-1}, S_j; \theta_*) \mid X_{-m}^k \right] - E \left[ \Gamma(X_j | X_{j-1}, S_j; \theta_*) \mid X_{-n}^k \right] \right\| \\ &= \left\| \sum_{a \in \mathbb{S}} \Gamma(X_j | X_{j-1}, a; \theta_*) \{ \bar{P}_{\theta_*}^\nu(S_j = a \mid X_{-m}^k) - \bar{P}_{\theta_*}^\nu(S_j = a \mid X_{-n}^k) \} \right\| \\ &\leq \max_{a \in \mathbb{S}} \|\Gamma(X_j | X_{j-1}, a; \theta_*)\| \left\| \bar{P}_{\theta_*}^\nu(S_j = \cdot \mid X_{-m}^k) - \bar{P}_{\theta_*}^\nu(S_j = \cdot \mid X_{-n}^k) \right\|_1. \end{aligned}$$

By Lemma 20(ii),

$$\begin{aligned} & \left\| E \left[ \Gamma(X_j | X_{j-1}, S_j; \theta_*) \mid X_{-m}^k \right] - E \left[ \Gamma(X_j | X_{j-1}, S_j; \theta_*) \mid X_{-n}^k \right] \right\| \\ &\leq L \max_{a \in \mathbb{S}} \|\Gamma(X_j | X_{j-1}, a; \theta_*)\| \prod_{i=-m}^j (1 - \underline{q}(X_i)), \end{aligned}$$

Therefore, by the Hölder inequality, it follows that, for  $a^{-1} + b^{-1} = 1$  (with  $a$  as in Assumption 7),

$$\begin{aligned} & \left\| \sum_{j=-m'}^l \left\{ E \left[ \Gamma(X_j | X_{j-1}, S_j; \theta_*) \mid X_{-m}^k \right] - E \left[ \Gamma(X_j | X_{j-1}, S_j; \theta_*) \mid X_{-n}^k \right] \right\} \right\|_{L^2(\bar{P}_*^\nu)} \\ &\leq \sum_{j=-m'}^l \left\| E \left[ \Gamma(X_j | X_{j-1}, S_j; \theta_*) \mid X_{-m}^k \right] - E \left[ \Gamma(X_j | X_{j-1}, S_j; \theta_*) \mid X_{-n}^k \right] \right\|_{L^2(\bar{P}_*^\nu)} \\ &\leq L \left( \sum_{a \in \mathbb{S}} E_{\bar{P}_*^\nu} [\|\Gamma(X_1 | X_0, a; \theta_*)\|^{2a}] \right)^{1/(2a)} \sum_{j=-m'}^l \left( E_{\bar{P}_*^\nu} \left[ \prod_{i=-m}^j (1 - \underline{q}(X_i))^{2b} \right] \right)^{1/(2b)}, \end{aligned}$$

where the second line follows from the triangle inequality and the third follows from stationarity (Lemma 1). The fact that  $\Gamma(X_1|X_0, a; \theta_*) = \nabla_{\theta} \log p_{\theta_*}(X_0, a, X_1)$ , Assumption 7(i) and definition of  $\varrho$  imply the desired result.

**Part (ii).** Follows from analogous calculations to those in part (i) and Lemma 20(i).

**Parts (iii) and (iv).** We only work out part (iii) since (iv) is analogous. Observe that

$$\begin{aligned}
& \left\| E \left[ \Lambda(S_j | S_{j-1}, X_{j-1}; \theta_*) \mid X_{-m}^k \right] - E \left[ \Lambda(S_j | S_{j-1}, X_{j-1}; \theta_*) \mid X_{-n}^k \right] \right\| \\
&= \left\| \sum_{(a,b) \in \mathbb{S}^2} \Lambda(b|a, X_{j-1}; \theta_*) \{ \bar{P}_{\theta_*}^{\nu}(S_j = b, S_{j-1} = a \mid X_{-m}^k) - \bar{P}_{\theta_*}^{\nu}(S_j = b, S_{j-1} = a \mid X_{-n}^k) \} \right\| \\
&\leq \sum_{(a,b) \in \mathbb{S}^2} \|\Lambda(b|a, X_{j-1}; \theta_*)\| \left| \bar{P}_{\theta_*}^{\nu}(S_j = b, S_{j-1} = a \mid X_{-m}^k) - \bar{P}_{\theta_*}^{\nu}(S_j = b, S_{j-1} = a \mid X_{-n}^k) \right| \\
&= \sum_{(a,b) \in \mathbb{S}^2} \|\Lambda(b|a, X_{j-1}; \theta_*)\| \bar{P}_{\theta_*}^{\nu}(S_j = b \mid S_{j-1} = a, X_{j-1}^k) \left| \bar{P}_{\theta_*}^{\nu}(S_{j-1} = a \mid X_{-m}^k) - \bar{P}_{\theta_*}^{\nu}(S_{j-1} = a \mid X_{-n}^k) \right|
\end{aligned}$$

where the last line follows from the Markov property of the model and the fact that  $-m \leq j$ . Thus

$$\begin{aligned}
& \left\| E \left[ \Lambda(S_j | S_{j-1}, X_{j-1}; \theta_*) \mid X_{-m}^k \right] - E \left[ \Lambda(S_j | S_{j-1}, X_{j-1}; \theta_*) \mid X_{-n}^k \right] \right\| \\
&\leq \max_{a,b} \|\Lambda(b|a, X_{j-1}; \theta_*)\| \left\| \bar{P}_{\theta_*}^{\nu}(S_{j-1} = \cdot \mid X_{-m}^k) - \bar{P}_{\theta_*}^{\nu}(S_{j-1} = \cdot \mid X_{-n}^k) \right\|_1 \\
&\leq L \max_{a,b} \|\Lambda(b|a, X_{j-1}; \theta_*)\| \prod_{i=-m}^{j-1} (1 - \underline{q}(X_i))
\end{aligned}$$

where the last line follows from Lemma 20(ii). Hence

$$\begin{aligned}
& \left\| \sum_{j=-m'}^l \{ E \left[ \Lambda(S_j | S_{j-1}, X_{j-1}; \theta_*) \mid X_{-m}^k \right] - E \left[ \Lambda(S_j | S_{j-1}, X_{j-1}; \theta_*) \mid X_{-n}^k \right] \} \right\|_{L^2(\bar{P}_*^{\nu})} \\
&\leq L \left( \sum_{(a,b) \in \mathbb{S}^2} E_{\bar{P}_*^{\nu}} \left[ \|\Lambda(b|a, X_0; \theta_*)\|^{2a} \right] \right)^{1/(2a)} \sum_{j=-m'}^l \left( E_{P_*^{\nu}} \left[ \prod_{i=-m}^{j-1} (1 - \underline{q}(X_i))^{2b} \right] \right)^{1/(2b)},
\end{aligned}$$

so the result follows from Assumption 7(i).  $\square$

*Proof of Lemma 18.* Throughout the proof we denote  $\|\cdot\|_{L^2(\bar{P}_*^\nu)}$  as  $\|\cdot\|_{L^2}$ . Also, we use  $\Phi$  and  $\Psi$  to denote  $\Phi_{\theta_*}$  and  $\Psi_{\theta_*}$ , resp.

**Part (i):** Observe that  $\Phi(k-1, k-T, l, k-T) = \Phi(k-1, [k-T/2], l, k-T) + \Phi([k-T/2]-1, k-T, l, k-T)$  and an analogous result holds for  $\Psi$ . Therefore, by the definition of  $\Delta_{k,k-T}$  and analogous calculations to those in [Bickel et al., 1998, pp. 1624–1626],

$$\begin{aligned}
& \|\Delta_{k,k-T}(\theta_*) - \Delta_{k,-\infty}(\theta_*)\|_{L^2} \\
& \leq \|\Phi(k-1, [k-T/2], k, k-T) - \Phi(k-1, [k-T/2], k, -\infty)\|_{L^2} \\
& \quad + \|\Phi(k-1, [k-T/2], k-1, k-T) - \Phi(k-1, [k-T/2], k-1, -\infty)\|_{L^2} \\
& \quad + \|\Phi([k-T/2]-1, k-T, k, k-T) - \Phi([k-T/2]-1, k-T, k-1, k-T)\|_{L^2} \\
& \quad + \|\Psi(k-1, [k-T/2], k, k-T) - \Psi(k-1, [k-T/2], k, -\infty)\|_{L^2} \\
& \quad + \|\Psi(k-1, [k-T/2], k-1, k-T) - \Psi(k-1, [k-T/2], k-1, -\infty)\|_{L^2} \\
& \quad + \|\Psi([k-T/2]-1, k-T-1, k, k-T) - \Psi([k-T/2]-1, k-T-1, k-1, k-T)\|_{L^2} \\
& \quad + \|\Phi(k, k, k, k-T) - \Phi(k, k, k, -\infty)\|_{L^2} + \|\Psi(k, k, k, k-T) - \Psi(k, k, k, -\infty)\|_{L^2} \\
& \equiv \sum_{i=1}^8 Term_i.
\end{aligned}$$

Terms 1 and 2 are analogous of the form

$$\begin{aligned}
& \|\Phi(k-1, [k-T/2], k, k-T) - \Phi(k-1, [k-T/2], k, -\infty)\|_{L^2} \\
& = \left\| \sum_{j=[k-T/2]}^{k-1} E_{\bar{P}_{\theta_*}^\nu} [\Gamma(X_j | X_{j-1}, S_j; \theta_*) | X_{k-T}^m] - \sum_{j=[k-T/2]}^{k-1} E_{\bar{P}_{\theta_*}^\nu} [\Gamma(X_j | X_{j-1}, S_j; \theta_*) | X_{-\infty}^m] \right\|_{L^2}
\end{aligned}$$

for  $m \in \{k, k-1\}$ . By Lemma 21(i), for  $i \in \{1, 2\}$ ,  $Term_i \lesssim \left( \sum_{j=[k-T/2]}^{k-1} \varrho(j, k-T) \right)$ .  $Term_7$  has the same bound by analogous calculations.

The  $Term_3$  is of the form

$$\left\| \sum_{j=k-T}^{[k-T/2]-1} E_{\bar{P}_{\theta_*}^\nu} [\Gamma(X_j | X_{j-1}, S_j; \theta_*) | X_{k-T}^k] - \sum_{j=k-T}^{[k-T/2]-1} E_{\bar{P}_{\theta_*}^\nu} [\Gamma(X_j | X_{j-1}, S_j; \theta_*) | X_{k-T}^{k-1}] \right\|_{L^2}.$$

By Lemma 21(ii),  $Term_3 \lesssim \left( \sum_{j=k-T}^{[k-T/2]-1} \varrho(k-1, j) \right)$ . The term  $Term_8$  has the same bound by analogous calculations.

The terms  $Term_4$  and  $Term_5$  are of the form

$$\left\| \sum_{j=[k-T/2]}^{k-1} E_{\bar{P}_{\theta_*}^\nu} [\Lambda(S_j | S_{j-1}, X_{j-1}; \theta_*) | X_{k-T}^m] - \sum_{j=[k-T/2]}^{k-1} E_{\bar{P}_{\theta_*}^\nu} [\Lambda(S_j | S_{j-1}, X_{j-1}; \theta_*) | X_{-\infty}^m] \right\|_{L^2}$$

and by Lemma 21(iii) are bounded by  $C \left( \sum_{j=[k-T/2]}^{k-1} \varrho(j-1, k-T) \right)$  for some universal constant  $C > 0$ .

Finally, by analogous calculations to those for  $Term_3$ ,  $Term_6$  is bounded by  $C \left( \sum_{j=k-T}^{[k-T/2]-1} \varrho(k-1, j) \right)$  by Lemma 21(iv).

**Part (ii).** By part (i) and Lemma 17,

$$\begin{aligned} \left\| T^{-1/2} \sum_{t=1}^T \{ \Delta_{t,-\infty}(\theta_*) - \nabla_{\theta} \log p_t^{\nu}(\cdot | \cdot; \theta_*) \} \right\|_{L^2} &\leq T^{-1/2} \sum_{t=1}^T \| \{ \Delta_{t,-\infty}(\theta_*) - \Delta_{t,0}(\theta_*) \} \|_{L^2} \\ &\lesssim \left( T^{-1/2} \sum_{t=1}^T \sum_{j=[t/2]}^{t-1} \varrho(j, 0) + T^{-1/2} \sum_{t=1}^T \sum_{j=0}^{[t/2]-1} \varrho(t, j) \right). \end{aligned}$$

By Kronecker's lemma, it suffices to show that

$$\sum_{t=1}^T t^{-1/2} \sum_{j=[t/2]}^{t-1} \varrho(j, 0) \quad \text{and} \quad \sum_{t=1}^T t^{-1/2} \sum_{j=0}^{[t/2]-1} \varrho(t, j) \quad (17)$$

are bounded uniformly in  $T$ , where, recall,  $\varrho(j, k) \equiv \left( E \left[ \prod_{i=k}^j (1 - \underline{q}(X_i))^{\frac{2a}{1-a}} \right] \right)^{\frac{1-a}{2a}}$ .

Moreover,  $j \mapsto \varrho(j, k)$  is non-increasing and  $k \mapsto \varrho(j, k)$  is non-decreasing since  $1 - \underline{q}(\cdot) \leq 1$ . By Assumption 8,  $(\varrho(j, 0))_j$  is  $p$ -summable with  $p < 2/3$ , thus  $\lim_{j \rightarrow \infty} \varrho(j, 0)^p j = 0$  (if not, then  $\varrho(j, 0) > c/j^{1/p}$  for some  $c > 0$  and all  $j$  above certain point and this violates the assumption). Hence,

$$\sum_{j=[t/2]}^{t-1} \varrho(j, 0) < \sum_{j=[t/2]}^{t-1} \frac{1}{j^{1/p}} \leq \int_{[t/2]+1}^t x^{-1/p} dx \leq \frac{p}{1-p} (t/2)^{1-1/p},$$

for all  $t \geq \tau$  and some  $\tau > 0$ , and this implies that, for some constant  $const > 0$ ,

$$\sum_{t=1}^T t^{-1/2} \sum_{j=[t/2]}^{t-1} \varrho(j, 0) \leq C(\tau) + const \times \sum_{t=\tau+1}^T \frac{p}{1-p} t^{1-1/p-1/2} \leq C < \infty,$$

because  $1 - 1/p - 1/2 < -1 \Leftrightarrow p < 2/3$  ( $C$  is a finite constant, which may depend on  $\tau$ ).

By stationarity of  $X_{-\infty}^{\infty}$  (Lemma 1),

$$\begin{aligned}
\sum_{j=0}^{[t/2]-1} \varrho(t, j) &= \left( E \left[ \prod_{i=0}^t (1 - \underline{q}(X_i))^{\frac{2a}{1-a}} \right] \right)^{\frac{1-a}{2a}} + \left( E \left[ \prod_{i=1}^t (1 - \underline{q}(X_i))^{\frac{2a}{1-a}} \right] \right)^{\frac{1-a}{2a}} \\
&\quad + \dots + \left( E \left[ \prod_{i=[t/2]-1}^t (1 - \underline{q}(X_i))^{\frac{2a}{1-a}} \right] \right)^{\frac{1-a}{2a}} \\
&= \left( E \left[ \prod_{i=0}^t (1 - \underline{q}(X_i))^{\frac{2a}{1-a}} \right] \right)^{\frac{1-a}{2a}} + \left( E \left[ \prod_{i=0}^{t-1} (1 - \underline{q}(X_i))^{\frac{2a}{1-a}} \right] \right)^{\frac{1-a}{2a}} \\
&\quad + \dots + \left( E \left[ \prod_{i=0}^{[t/2]-1} (1 - \underline{q}(X_i))^{\frac{2a}{1-a}} \right] \right)^{\frac{1-a}{2a}} \\
&= \sum_{j=0}^{[t/2]+1} \varrho(t-j, 0).
\end{aligned}$$

Thus  $\sum_{j=0}^{[t/2]+1} \varrho(t-j, 0) \leq \sum_{j=0}^{[t/2]+1} \frac{1}{(t-j)^{1/p}} \leq \int_{[t/2]-1}^t \frac{1}{u^{1/p}} du$  and by our previous calculation the result follows. Thus, the terms in (17) are uniformly bounded.  $\square$

*Proof of Lemma 19.* It is easy to see that  $\Delta_{t,-\infty}(\theta_*)$  is adapted to the filtration associated with the  $\sigma$ -algebra generated by  $X_{-\infty}^t$ . Since  $X_{-\infty}^{\infty}$  is stationary and ergodic (by Lemma 1), so is  $(\Delta_{t,-\infty}(\theta_*))_{t=-\infty}^{\infty}$ .  $\square$

To prove Lemma 20, we need the following results.

**Lemma 22.** *Suppose Assumption 1 holds. Then, there exists a finite constant  $L > 0$ , such that, for any  $-m \leq j < n \leq k$  and any  $\theta \in \Theta$ ,*

$$\max_{b,c} \left\| \bar{P}_{\theta}^{\nu} \left( S_j = \cdot | S_n = b, X_{-m}^k \right) - \bar{P}_{\theta}^{\nu} \left( S_j = \cdot | S_n = c, X_{-m}^k \right) \right\|_1 \leq L \prod_{i=j}^n (1 - q(X_i))$$

*a.s.- $\bar{P}_{*}^{\nu}$ .*

**Lemma 23.** *For any  $-m < i < l \leq r \leq n$ , let  $\mathbf{S}_l^r \equiv (S_l, \dots, S_r)$ . Then, for any  $\theta \in \Theta$ ,*

$$\bar{P}_{\theta}^{\nu} (S_i | \mathbf{S}_l^r, X_{-m+1}^n) = \bar{P}_{\theta}^{\nu} (S_i | S_l, X_{-m+1}^n),$$

*i.e., the Markov property holds backward in time.*



*Proof of Lemma 22.* Observe that, for any  $b, c \in \mathbb{S}^2$ ,

$$\begin{aligned} & \left\| \bar{P}_\theta^\nu \left( S_j = \cdot | S_n = b, X_{-m}^k \right) - \bar{P}_\theta^\nu \left( S_j = \cdot | S_n = c, X_{-m}^k \right) \right\|_1 \\ &= \left\| \sum_{s \in \mathbb{S}} \bar{P}_\theta^\nu \left( S_j = \cdot | S_{j+1} = s, X_{-m}^{j+1} \right) \left( \bar{P}_\theta^\nu \left( S_{j+1} = s | S_n = b, X_{-m}^k \right) - \bar{P}_\theta^\nu \left( S_{j+1} = s | S_n = c, X_{-m}^k \right) \right) \right\|_1 \\ &\leq \alpha_{\theta, j+1, j}(X_{-m}^k) \left\| \bar{P}_\theta^\nu \left( S_{j+1} = \cdot | S_n = b, X_{-m}^k \right) - \bar{P}_\theta^\nu \left( S_{j+1} = \cdot | S_n = c, X_{-m}^k \right) \right\|_1 \end{aligned}$$

where the second line follows from Lemma 23 with  $i = j$ ,  $r = l = j + 1$  and  $n = k$ , and the third follows from Lemma B.2.1 in Stachurski [2009] and the definition of  $\alpha_{\theta, j+1, j}(X_{-m}^k)$  in expression (12). Iterating in this fashion it follows that

$$\left\| \bar{P}_\theta^\nu \left( S_j = \cdot | S_n = b, X_{-m}^k \right) - \bar{P}_\theta^\nu \left( S_j = \cdot | S_n = c, X_{-m}^k \right) \right\|_1 \leq 2 \prod_{l=j}^n \alpha_{\theta, l+1, l}(X_{-m}^k).$$

Thus, it suffices to show that  $\alpha_{\theta, l+1, l}(X_{-m}^k) \leq 1 - \underline{q}(X_l)$ . Since

$$\alpha_{\theta, l+1, l}(X_{-m}^k) = 1 - \min_{a, b} \sum_{s' \in \mathbb{S}} \min \{ \bar{P}_\theta^\nu \left( S_l = s' | S_{l+1} = a, X_{-m}^k \right), \bar{P}_\theta^\nu \left( S_l = s' | S_{l+1} = b, X_{-m}^k \right) \}$$

(see [Stachurski, 2009, p. 344]), it suffices to show that, for any  $(a, b) \in \mathbb{S}^2$ ,

$$\bar{P}_\theta^\nu \left( S_l = a | S_{l+1} = b, X_{-m}^k \right) \geq \underline{q}(X_l) \varpi(X_{-m+1}^l, a),$$

where  $\varpi(X_{-m+1}^{k-1}, \cdot) \in \mathcal{P}(\mathbb{S})$ .

To do this, first note that  $\bar{P}_\theta^\nu \left( S_l = a | S_{l+1} = b, X_{-m}^k \right) = \bar{P}_\theta^\nu \left( S_l = a | S_{l+1} = b, X_{-m}^{l+1} \right)$ , by Lemma 23, and

$$\begin{aligned} \bar{P}_\theta^\nu \left( S_l = a | S_{l+1} = b, X_{-m+1}^{l+1} \right) &= \frac{p_\theta(X_l, b, X_{l+1}) Q_\theta(X_l, a, b) \bar{P}_\theta^\nu(X_{-m+1}^l, S_l = a)}{\sum_{s \in \mathbb{S}} p_\theta(X_l, b, X_{l+1}) Q_\theta(X_l, s, b) \bar{P}_\theta^\nu(X_{-m+1}^l, S_l = s)} \\ &\geq \underline{q}(X_l) \frac{\bar{P}_\theta^\nu(X_{-m+1}^l, S_l = a)}{\sum_{s \in \mathbb{S}} \bar{P}_\theta^\nu(X_{-m+1}^l, S_l = s)}, \end{aligned}$$

where the last line follows from Assumption 1. Letting  $\varpi(\cdot, X_{-m+1}^l) \equiv \frac{\bar{P}_\theta^\nu(S_l = \cdot | X_{-m+1}^l)}{\sum_{s \in \mathbb{S}} \bar{P}_\theta^\nu(S_l = s | X_{-m+1}^l)}$ , the desired result is obtained.  $\square$

*Proof of Lemma 23.* Throughout the proof, we omit  $\theta$  from the notation. Let  $\mathbf{S}_{i:l}^r \equiv (S_i, S_l, S_{l+1}, \dots, S_{r-1}, S_r)$  and note that

$$\bar{P}_\theta^\nu \left( S_i | \mathbf{S}_l^r, X_{-m+1}^n \right) = \frac{\bar{P}_\theta^\nu \left( X_r^n | \mathbf{S}_{i:l}^r, X_{-m+1}^{r-1} \right) \bar{P}_\theta^\nu \left( \mathbf{S}_{i:l}^r, X_{-m+1}^{r-1} \right)}{\bar{P}_\theta^\nu \left( X_r^n | \mathbf{S}_l^r, X_{-m+1}^{r-1} \right) \bar{P}_\theta^\nu \left( \mathbf{S}_l^r, X_{-m+1}^{r-1} \right)},$$

by Bayes' rule. By the Markov property, it follows that  $\bar{P}_\theta^\nu(X_r^n \mid \mathbf{S}_{i:l}^r, X_{-m+1}^{r-1}) = \bar{P}_\theta^\nu(X_r^n \mid X_{r-1}, S_r)$ , so

$$\bar{P}_\theta^\nu(S_i \mid \mathbf{S}_l^r, X_{-m+1}^n) = \frac{\bar{P}_\theta^\nu(\mathbf{S}_{i:l}^r, X_{-m+1}^{r-1})}{\bar{P}_\theta^\nu(\mathbf{S}_l^r, X_{-m+1}^{r-1})} = \frac{\bar{P}_\theta^\nu(S_r \mid \mathbf{S}_{i:l}^{r-1}, X_{-m+1}^{r-1})}{\bar{P}_\theta^\nu(S_r \mid \mathbf{S}_l^{r-1}, X_{-m+1}^{r-1})} \frac{\bar{P}_\theta^\nu(\mathbf{S}_{i:l}^{r-1}, X_{-m+1}^{r-1})}{\bar{P}_\theta^\nu(\mathbf{S}_l^{r-1}, X_{-m+1}^{r-1})}.$$

Observe that  $\bar{P}_\theta^\nu(S_r \mid \mathbf{S}_l^{r-1}, X_{-m+1}^{r-1}) = Q_\theta(X_{r-1}, S_{r-1}, S_r)$ , and thus  $\bar{P}_\theta^\nu(S_i \mid X_{-m+1}^r, \mathbf{S}_l^r) = \frac{\bar{P}_\theta^\nu(\mathbf{S}_{i:l}^{r-1}, X_{-m+1}^{r-1})}{\bar{P}_\theta^\nu(\mathbf{S}_l^{r-1}, X_{-m+1}^{r-1})}$  and, by iterating, it follows that

$$\bar{P}_\theta^\nu(S_i \mid \mathbf{S}_l^r, X_{-m+1}^n) = \frac{\bar{P}_\theta^\nu(S_i, S_l, X_{-m+1}^l)}{\bar{P}_\theta^\nu(\mathbf{S}_l^r, X_{-m+1}^l)} = \bar{P}_\theta^\nu(S_i \mid S_l, X_{-m+1}^l),$$

as desired.  $\square$

*Proof of Lemma 20. Part (i).* By Lemma 23 with  $l = r = k - 1$ ,  $n = k$ , it follows that

$$\begin{aligned} \bar{P}_\theta^\nu(S_i \mid X_{-m+1}^k) &= \sum_{s \in \mathbb{S}} \bar{P}_\theta^\nu(S_i \mid S_{k-1} = s, X_{-m+1}^k) \bar{P}_\theta^\nu(S_{k-1} = s \mid X_{-m+1}^k) \\ &= \sum_{s \in \mathbb{S}} \bar{P}_\theta^\nu(S_i \mid S_{k-1} = s, X_{-m+1}^{k-1}) \bar{P}_\theta^\nu(S_{k-1} = s \mid X_{-m+1}^k), \end{aligned}$$

and similarly,

$$\bar{P}_\theta^\nu(S_i \mid X_{-m+1}^{k-1}) = \sum_{s \in \mathbb{S}} \bar{P}_\theta^\nu(S_i \mid S_{k-1} = s, X_{-m+1}^{k-1}) \bar{P}_\theta^\nu(S_{k-1} = s \mid X_{-m+1}^{k-1}).$$

Thus, by Lemma B.2.2 in Stachurski [2009],

$$\begin{aligned} &\left\| \bar{P}_\theta^\nu(S_j = \cdot \mid X_{-m+1}^k) - \bar{P}_\theta^\nu(S_j = \cdot \mid X_{-m+1}^{k-1}) \right\|_1 \\ &\leq \max_{a,b} \left\| \bar{P}_\theta^\nu(S_j = \cdot \mid S_{k-1} = a, X_{-m+1}^{k-1}) - \bar{P}_\theta^\nu(S_j = \cdot \mid S_{k-1} = b, X_{-m+1}^{k-1}) \right\|_1 \\ &\leq L \prod_{l=j}^{k-1} (1 - q(X_l)), \end{aligned}$$

where the second line follows by Lemma 22 with  $n = k - 1$ . Thus, the desired result follows.

**Part (ii).** The proof is analogous to that of Lemma 5 (third part) in Bickel

et al. [1998]. By analogous calculations to those in part (i),

$$\begin{aligned}
& \left\| \bar{P}_\theta^\nu(S_j = \cdot \mid X_{-m}^k) - \bar{P}_\theta^\nu(S_j = \cdot \mid X_{-n}^k) \right\|_1 \\
& \leq \max_{b,b'} \left\| \bar{P}_\theta^\nu(S_j = \cdot \mid S_{-m} = b, X_{-m}^k) - \bar{P}_\theta^\nu(S_j = \cdot \mid S_{-m} = b', X_{-n}^k) \right\|_1 \\
& = \max_{b,b'} \left\| \bar{P}_\theta^\nu(S_j = \cdot \mid S_{-m} = b, X_{-m}^k) - \bar{P}_\theta^\nu(S_j = \cdot \mid S_{-m} = b', X_{-m}^k) \right\|_1,
\end{aligned}$$

where the last line follows from the fact that, given  $S_{-m}$ , it is the same to condition on  $X_{-m}^k$  and on  $X_{-n}^k$ . The results thus follows from following the same steps as those in the proof of Theorem 2.  $\square$

### SM.8.2 Proof of Lemma 9

*Proof of Lemma 9.* To simplify the exposition, we present the proof for the case where  $\Delta_{t,-\infty}(\theta_*)(X_{-\infty}^t)$  is a scalar; since the dimension of this quantity is finite, the vector case follows readily from the results below.

Part (a) follows easily from Lemma 17 in the Supplemental Material SM.8.1.

For part (b), it follows from part (a) that  $\Delta_{t+j,-\infty}(\theta_*)(X_{-\infty}^{t+j})\Delta_{t,-\infty}(\theta_*)(X_{-\infty}^t)$  depends only on  $(X_{t-\bar{L}}, \dots, X_{t+j})$ . By Lemma 1,  $(X_t)_{t=-\infty}^\infty$  is  $\beta$ -mixing. Since, for any fixed  $j$ , the  $\sigma$ -algebra generated by  $(\Delta_{s+j,-\infty}(\theta_*)\Delta_{s,-\infty}(\theta_*))_{s \leq t}$  is contained in the  $\sigma$ -algebra generated by  $(X_{s+j})_{s \leq t}$ , and the  $\sigma$ -algebra generated by  $(\Delta_{s+j,-\infty}(\theta_*)\Delta_{s,-\infty}(\theta_*))_{s \geq t}$  is contained in the  $\sigma$ -algebra generated by  $(X_s)_{s \geq t-\bar{L}}$ , it follows that, for each  $j$ ,  $(\Delta_{t+j,-\infty}(\theta_*)\Delta_{t,-\infty}(\theta_*))_{t=-\infty}^\infty$  is also  $\beta$ -mixing with mixing coefficients that decay at rate  $O(\gamma^{n-2\bar{L}})$  as  $n$  diverges through the positive integers; as  $\bar{L}$  is taken to be fixed, the decay rate is  $O(\gamma^n)$ . As is well known, this result implies that the corresponding  $\alpha$ -mixing coefficients  $(\alpha_n)_n$  decay at the same rate; i.e.,  $\alpha_n = O(\gamma^n)$  for all  $n$ .

Henceforth, let  $\Omega_{t+j,t,-\infty}(\theta_*) \equiv \Delta_{t+j,-\infty}(\theta_*)(X_{-\infty}^{t+j})\Delta_{t,-\infty}(\theta_*)(X_{-\infty}^t)$  and  $\bar{\Omega}_{t+j,t,-\infty}(\theta_*) \equiv \Omega_{t+j,t,-\infty}(\theta_*) - E_{\bar{P}_*^\nu}[\Omega_{t+j,t,-\infty}(\theta_*)]$ , for any  $t, j$ . Observe that, for any  $j$ ,

$$\begin{aligned}
E_{\bar{P}_*^\nu} \left[ \left( T^{-1} \sum_{t=1}^T \bar{\Omega}_{t+j,t,-\infty}(\theta_*) \right)^2 \right] &= T^{-1} \sum_{t=1}^T E_{\bar{P}_*^\nu} \left[ (\bar{\Omega}_{t+j,t,-\infty}(\theta_*))^2 \right] \\
&\quad + 2T^{-2} \sum_{t=1}^T \sum_{\tau=0}^{t-1} E_{\bar{P}_*^\nu} \left[ (\bar{\Omega}_{t+j,t,-\infty}(\theta_*)) (\bar{\Omega}_{s+j,\tau,-\infty}(\theta_*)) \right] \\
&= T^{-1} E_{\bar{P}_*^\nu} \left[ (\bar{\Omega}_{j,0,-\infty}(\theta_*))^2 \right] \\
&\quad + 2T^{-1} \sum_{t=0}^{T-1} (1 - t/T) E_{\bar{P}_*^\nu} \left[ (\bar{\Omega}_{t+j,t,-\infty}(\theta_*)) (\bar{\Omega}_{j,0,-\infty}(\theta_*)) \right],
\end{aligned}$$

where the last equality follows by stationarity. By Corollary 6.17 in [White \[2001\]](#), for any  $m \in \mathbb{N}$ ,

$$|E_{\bar{P}_*^\nu}[\bar{\Omega}_{j,0,-\infty}(\theta_*)\bar{\Omega}_{j+m,m,-\infty}(\theta_*)]| \lesssim (\alpha_m)^{\frac{2}{2+2\delta}} \sqrt{E_{\bar{P}_*^\nu}[(\bar{\Omega}_{j,0,-\infty}(\theta_*))^2]} (E_{\bar{P}_*^\nu}[|\bar{\Omega}_{j+m,m,-\infty}(\theta_*)|^{2+2\delta}])^{\frac{1}{2+2\delta}},$$

for any  $j$  (the implicit constant in the display does not depend on  $j$ ). Thus, for any  $j$ ,

$$\begin{aligned} E_{\bar{P}_*^\nu} \left[ \left( T^{-1} \sum_{t=1}^T \bar{\Omega}_{t+j,t,-\infty}(\theta_*) \right)^2 \right] &\lesssim T^{-1} E_{\bar{P}_*^\nu} [(\bar{\Omega}_{j,0,-\infty}(\theta_*))^2] \\ &+ 2T^{-1} \sum_{t=0}^{T-1} (1-t/T) (\alpha_t)^{2/(2+2\delta)} \sqrt{E_{\bar{P}_*^\nu}[(\bar{\Omega}_{j,0,-\infty}(\theta_*))^2]} (E_{\bar{P}_*^\nu}[|\bar{\Omega}_{j+m,m,-\infty}(\theta_*)|^{2+2\delta}])^{1/(2+2\delta)}. \end{aligned}$$

By the Cauchy-Schwarz inequality and stationarity, for any  $j$ ,

$$E_{\bar{P}_*^\nu} [(\bar{\Omega}_{j,0,-\infty}(\theta_*))^2] \lesssim E_{\bar{P}_*^\nu} [\Delta_{0,-\infty}(\theta_*) (X_{-\infty}^0)^4],$$

which is bounded by assumption. In addition, by similar calculations, it follows that, for any  $j$ ,

$$E_{\bar{P}_*^\nu} [(\bar{\Omega}_{j,0,-\infty}(\theta_*))^{2+2\delta}] \lesssim E_{\bar{P}_*^\nu} [\Delta_{0,-\infty}(\theta_*) (X_{-\infty}^0)^{4+4\delta}],$$

which is bounded by assumption. Therefore, there exists a finite constant  $C$  (which does not depend on  $j$ ) such that

$$E_{\bar{P}_*^\nu} \left[ \left( T^{-1} \sum_{t=1}^T \bar{\Omega}_{t+j,t,-\infty}(\theta_*) \right)^2 \right] \leq C(T^{-1} + 2T^{-1} \sum_{t=0}^{T-1} (1-t/T) (\alpha_t)^{2/(2+2\delta)}).$$

As  $\alpha_t = O(\gamma^t)$ , it follows that  $\sum_{t=0}^{T-1} (1-t/T) (\alpha_t)^{2/(2+2\delta)} = O\left(\sum_{t=0}^{T-1} (1-t/T) (\gamma^{2/(2+2\delta)})^t\right)$ .

Since  $\gamma < 1$ , it follows that  $\sum_{t=0}^{T-1} (1-t/T) (\alpha_t)^{2/(2+2\delta)} = O(1)$ , which in turn implies that  $E_{\bar{P}_*^\nu} \left[ \left( T^{-1} \sum_{t=1}^T \bar{\Omega}_{t+j,t,-\infty}(\theta_*) \right)^2 \right] \leq CT^{-1}$ . Hence, by the Markov inequality, for any  $a > 0$ ,

$$\begin{aligned} &\bar{P}_*^\nu \left( \max_{j \in \{0, \dots, L\}} \left\| T^{-1} \sum_{t=1}^T \Delta_{t+j,-\infty}(\theta_*) (X_{-\infty}^{t+j}) \Delta_{t,-\infty}(\theta_*) (X_{-\infty}^t)^\top - E_{\bar{P}_*^\nu} [\Delta_{j,-\infty}(\theta_*) (X_{-\infty}^j) \Delta_{0,-\infty}(\theta_*) (X_{-\infty}^0)^\top] \right\| \geq a \right) \\ &\leq \sum_{j=0}^L \bar{P}_*^\nu \left( \left\| T^{-1} \sum_{t=1}^T \Delta_{t+j,-\infty}(\theta_*) (X_{-\infty}^{t+j}) \Delta_{t,-\infty}(\theta_*) (X_{-\infty}^t)^\top - E_{\bar{P}_*^\nu} [\Delta_{j,-\infty}(\theta_*) (X_{-\infty}^j) \Delta_{0,-\infty}(\theta_*) (X_{-\infty}^0)^\top] \right\| \geq a \right) \\ &\leq Ca^{-2} LT^{-1}, \end{aligned}$$

which implies the desired result.  $\square$

## SM.9 Proof of Theorem 5

To prove Theorem 5, we use the following results, whose proofs are relegated to the end of this section. The following lemma shows that we can “quantify” convergence in probability.

**Lemma 24.** *Suppose a random sequence  $(X_T)_{T=0}^\infty$  converges to zero in probability. Then, there exists a deterministic positive sequence  $(r_T)_{T=0}^\infty$  such that  $r_T = o(1)$  and, for any  $\epsilon > 0$ , there exists  $T_\epsilon$  such that*

$$\Pr(|X_T| \geq r_T) \leq \epsilon,$$

for all  $T \geq T_\epsilon$ . In particular,  $|X_T| = O_{\Pr}(r_T)$ .

The next lemma presents some useful properties for the “score process”.

**Lemma 25.** *Under the Assumptions of Theorem 5, the following are true:*

1.  $\|\sup_{\theta \in B(\delta, \theta_*)} \|\Delta_{0,-\infty}(\theta)\|\|_{L^2(\bar{P}_*)} < \infty$  ( $\delta > 0$  is as in Assumption 7).
2.  $\Delta_{t,-\infty}$  and  $\Delta_{t,-\infty}\Delta_{0,-\infty}^\top$  are continuous in  $L^1(\bar{P}_*)$ -norm, i.e., for any  $\epsilon > 0$ , there exists  $\delta > 0$  such that

$$\max_t \left\| \sup_{\|\theta - \theta_0\| < \delta} \|\Delta_{t,-\infty}(\theta)\Delta_{t,-\infty}(\theta)^\top - \Delta_{t,-\infty}(\theta_0)\Delta_{t,-\infty}(\theta_0)^\top\| \right\|_{L^1(\bar{P}_*)} \leq \epsilon$$

and

$$\ddot{\omega}(\delta) \equiv \max_t \left\| \sup_{\|\theta - \theta_0\| < \delta} \|\Delta_{t,-\infty}(\theta)\Delta_{0,-\infty}(\theta)^\top - \Delta_{t,-\infty}(\theta_0)\Delta_{0,-\infty}(\theta_0)^\top\| \right\|_{L^1(\bar{P}_*)} \leq \epsilon.$$

3. There exists a constant  $C < \infty$  such that, for any  $t$  and  $M$ ,

$$\left\| \sup_{\theta \in B(\delta, \theta_*)} \Delta_{t,-\infty}(\theta) - \Delta_{t,t-M}(\theta) \right\|_{L^2(\bar{P}_*)} \leq CM^{1-1/p}.$$

Moreover, by Assumption 8,  $p \in (0, 2/3)$  and thus the RHS vanishes as  $M$  diverges.

*Proof of Theorem 5.* The proof has several parts and steps.

PART (A). We show that

$$\|T^{-1} \sum_{t=1}^T \nabla_\theta^2 \log p_t^\nu(X_t | X_0^{t-1}, \hat{\theta}_{\nu,T}) - E_{\bar{P}_*}[\xi_1(\theta_*)]\| = o_{\bar{P}_*}(1).$$

We do this by using the triangle inequality and showing that the following expressions hold:

$$\lim_{T \rightarrow \infty} \left\| T^{-1} \sum_{t=1}^T \{ \nabla_{\theta}^2 \log p_t^{\nu}(\cdot | \cdot, \hat{\theta}_{\nu, T}) - \xi_t(\hat{\theta}_{\nu, T}) \} \right\|_{L^1(\bar{P}_{\nu}^*)} = 0$$

(which implies convergence in probability),

$$\| T^{-1} \sum_{t=1}^T \{ \xi_t(\hat{\theta}_{\nu, T}) - \xi_t(\theta_*) \} \| = o_{\bar{P}_{\nu}^*}(1),$$

and

$$\| T^{-1} \sum_{t=1}^T \xi_t(\theta_*) - E_{\bar{P}_{\nu}^*}[\xi_1(\theta_*)] \| = o_{\bar{P}_{\nu}^*}(1).$$

The first expression holds true because, by Theorem 1, for any  $\delta' \leq \delta$ ,  $\hat{\theta}_{\nu, T} \in B(\delta', \theta_*)$  w.p.a.1 and, hence, by Lemma 8, the desired result follows.

Regarding the second expression, again by Theorem 1,  $\hat{\theta}_{\nu, T} \in B(\delta', \theta_*)$  w.p.a.1. Thus, it follows that, for any  $\epsilon > 0$ , there exists  $T(\epsilon)$  such that, for any  $t \geq T(\epsilon)$ ,

$$\begin{aligned} \bar{P}_{\nu}^* \left( \left\| T^{-1} \sum_{t=1}^T \{ \xi_t(\hat{\theta}_{\nu, T}) - \xi_t(\theta_*) \} \right\| \geq \epsilon \right) &\leq \bar{P}_{\nu}^* \left( \left\| T^{-1} \sum_{t=1}^T \{ \xi_t(\hat{\theta}_{\nu, T}) - \xi_t(\theta_*) \} \right\| \geq \epsilon, \hat{\theta}_{\nu, T} \in B(\delta', \theta_*) \right) + \epsilon \\ &\leq \epsilon^{-1} E_{\bar{P}_{\nu}^*} \left[ \sup_{\theta \in B(\delta', \theta_*)} \| \xi_1(\theta) - \xi_1(\theta_*) \| \right] + 0.5\epsilon, \end{aligned}$$

where the second line follows from the Markov inequality and stationarity. By Lemma 8,  $\xi_1$  is continuous – and thus uniformly continuous over compact sets. Since  $\delta' > 0$  can be chosen to be any number less than  $\delta$  ( $\delta$  as in Assumption 7), we can choose it so that the first term in the RHS is less than  $0.5\epsilon$ . Hence, the desired follows.

Finally, ergodicity of  $X_{-\infty}^{\infty}$  (Lemma 1) implies ergodicity of  $(\xi_t(\theta_*))_{t=-\infty}^{\infty}$ ; therefore, by Lemma 8 and Birkhoff's ergodic theorem,  $\| T^{-1} \sum_{t=1}^T \xi_t(\theta_*) - E_{\bar{P}_{\nu}^*}[\xi_1(\theta_*)] \| = o_{\bar{P}_{\nu}^*}(1)$ . Hence,

$$\| T^{-1} \sum_{t=1}^T \nabla_{\theta}^2 \log p_t^{\nu}(X_t | X_0^{t-1}, \hat{\theta}_{\nu, T}) - E_{\bar{P}_{\nu}^*}[\xi_1(\theta_*)] \| = o_{\bar{P}_{\nu}^*}(1).$$

With this result and the Fisher information equality (established in the proof of Corollary 1), the result of part (a) of the theorem follows.

PART (B). STEP 1 To prove part (b), it suffices to show that

$$\|E_{\bar{P}_*}[\xi_1(\theta_*)] - H_T(\hat{\theta}_{\nu,T})\| = o_{\bar{P}_*}(1)$$

and

$$\|\Sigma_T(\theta_*) - J_T(\hat{\theta}_{\nu,T})\| = o_{\bar{P}_*}(1).$$

The first expression was established in Part (A). Regarding the second expression, we introduce some notation. For any  $t, l, M \in \mathbb{N}$  and any  $\theta \in \Theta$ , let

$$\Omega_{t,l,M}(\theta) \equiv \Delta_{t,M}(\theta)(X_M^t)\Delta_{l,M}(\theta)(X_M^l)^\top,$$

where it is left implicit that this quantity depends on  $X_{-M}^{\max\{t,l\}}$ . Also, for any  $\tau \in \{1, \dots, L_T\}$ ,

$$\theta \mapsto \hat{\gamma}_{T,\tau,0}(\theta) \equiv (T - \tau)^{-1} \sum_{t=1}^{T-\tau} \Omega_{t+\tau,t,0}(\theta).$$

Recall that  $\Delta_{t,0}(\theta)(X_0^t) = \nabla_{\theta} p_t^\nu(X_t | X_0^{t-1}, \theta)$ , so  $\hat{\gamma}_{T,\tau,0}(\theta)$  is the sample covariance of  $\Omega_{t+\tau,t,0}(\theta)$ .

Given this notation, observe that

$$\Sigma_T(\theta_*) = T^{-1} \sum_{t=1}^T E_{\bar{P}_*}[\Omega_{t,t,-\infty}(\theta_*)] + T^{-1} \sum_{t=1}^T \sum_{l=0}^{t-1} \{E_{\bar{P}_*}[\Omega_{t,l,-\infty}(\theta_*)] + E_{\bar{P}_*}[\Omega_{t,l,-\infty}(\theta_*)^\top]\}.$$

The aim is to show that each of the summands above is well-approximated by its counterpart in  $J_T$ . For the first summand, we show in Step 2 below that

$$\|T^{-1} \sum_{t=1}^T \{\Omega_{t,t,0}(\hat{\theta}_{\nu,T}) - E_{\bar{P}_*}[\Omega_{t,t,-\infty}(\theta_*)]\| = o_{\bar{P}_*}(1).$$

Regarding the second summand, we observe that, for any  $t \geq l$ ,

$$E_{\bar{P}_*}[\Omega_{t,l,-\infty}(\theta_*)] = E_{\bar{P}_*}[\Omega_{t-l,0,-\infty}(\theta_*)] \equiv \gamma_{t-l}(\theta_*)$$

(the first equality, which follows from stationarity, can be established by analogous arguments to those presented at the beginning of Step 2). Hence,

$$\begin{aligned} T^{-1} \sum_{t=1}^T \sum_{l=0}^{t-1} E_{\bar{P}_*}[\Omega_{t,l,-\infty}(\theta_*)] &= T^{-1} (\gamma_1(\theta_*) + (\gamma_2(\theta_*) + \gamma_1(\theta_*)) + \dots + (\gamma_T(\theta_*) + \dots + \gamma_1(\theta_*))) \\ &= \sum_{j=0}^{T-1} (1 - j/T) \gamma_{j+1}(\theta_*). \end{aligned}$$

Thus, it suffices to show that

$$\left\| \sum_{j=0}^{T-1} (1 - j/T) \gamma_{j+1}(\theta_*) - \sum_{j=0}^{L_T-1} \omega(j, L) \hat{\gamma}_{T,j+1,0}(\hat{\theta}_{T,\nu}) \right\| = o_{\bar{P}_*^\nu}(1),$$

the proof of which is in Step 3 below.

STEP 2 We now show that

$$\left\| T^{-1} \sum_{t=1}^T \{ \Delta_{t,0}(\hat{\theta}_{\nu,T})(X_0^t) \Delta_{t,0}(\hat{\theta}_{\nu,T})(X^t)^\top - E_{\bar{P}_*^\nu} [\Delta_{t,-\infty}(\theta_*)(X_{-\infty}^t) \Delta_{t,-\infty}(\theta_*)(X_{-\infty}^t)^\top] \} \right\| = o_{\bar{P}_*^\nu}(1),$$

where, by the definition of  $\Delta_{t,M}$ ,  $\Delta_{t,0}(\theta)(X^t) = \nabla_\theta \log p_t(X_t | X_0^{t-1}, \theta)$ .

First, observe that

$$E_{\bar{P}_*^\nu} [\Delta_{t,-\infty}(\theta_*)(X_{-\infty}^t) \Delta_{t,-\infty}(\theta_*)(X_{-\infty}^t)^\top] = E_{\bar{P}_*^\nu} [\Delta_{0,-\infty}(\theta_*)(X_{-\infty}^0) \Delta_{0,-\infty}(\theta_*)(X_{-\infty}^0)^\top].$$

This follows from stationarity (see Lemma 1) and the fact that  $\Delta_{t,-\infty}(\theta_*)$  can be approximated (uniformly in  $t$ ) by  $\Delta_{t,t-M}(\theta_*)$  (see Lemma 18). Hence, it suffices to show that

$$\left\| T^{-1} \sum_{t=1}^T \Delta_{t,0}(\hat{\theta}_{\nu,T})(X^t) \Delta_{t,0}(\hat{\theta}_{\nu,T})(X^t)^\top - E_{\bar{P}_*^\nu} [\Delta_{0,-\infty}(\theta_*)(X_{-\infty}^0) \Delta_{0,-\infty}(\theta_*)(X_{-\infty}^0)^\top] \right\| = o_{\bar{P}_*^\nu}(1).$$

By Lemma 1, ergodicity of  $\Delta_{t,t-M}(\theta_*)$  for any  $M$  follows. This, Birkhoff's ergodic theorem, and Lemma 18 imply that

$$\left\| T^{-1} \sum_{t=1}^T \bar{\Delta}_\infty(\theta_*)(X_{-\infty}^t) - E_{\bar{P}_*^\nu} [\bar{\Delta}_\infty(\theta_*)(X_{-\infty}^0)] \right\| = o_{\bar{P}_*^\nu}(1),$$

where, for any  $M \in \mathbb{Z} \cup \{\infty\}$ ,  $\bar{\Delta}_M(\theta)(X_{t-M}^t) \equiv \Delta_{0,-M}(\theta)(X_{t-M}^t) \Delta_{0,-M}(\theta)(X_{t-M}^t)^\top$ .

Hence, in order to obtain the desired result it suffices to show that

$$\left\| T^{-1} \sum_{t=1}^T \{ \bar{\Delta}_t(\hat{\theta}_{\nu,T})(X_0^t) - \bar{\Delta}_\infty(\theta_*)(X_{-\infty}^t) \} \right\| = o_{\bar{P}_*^\nu}(1).$$

In order to do so, by the triangle inequality, it is sufficient to show that

$$\left\| T^{-1} \sum_{t=1}^T \{ \bar{\Delta}_\infty(\theta_*)(X_{-\infty}^t) - \bar{\Delta}_\infty(\hat{\theta}_{\nu,T})(X_{-\infty}^t) \} \right\| = o_{\bar{P}_*^\nu}(1) \quad (18)$$



and

$$\|T^{-1} \sum_{t=1}^T \{\bar{\Delta}_t(\hat{\theta}_{\nu,T})(X_0^t) - \bar{\Delta}_\infty(\hat{\theta}_{\nu,T})(X_{-\infty}^t)\}\| = o_{\bar{P}_*^\nu}(1). \quad (19)$$

Expression (18) holds by Lemma 25, the fact that, for any  $\delta > 0$ ,  $\hat{\theta}_{T,\nu} \in B(\delta, \theta_*)$  w.p.a.1- $\bar{P}_*^\nu$  (by Theorem 1) and the Markov inequality. Regarding expression (19), by the Markov inequality and the fact that  $\hat{\theta}_{T,\nu} \in B(\delta, \theta_*)$  w.p.a.1- $\bar{P}_*^\nu$  (by Theorem 1), it is sufficient to show that

$$T^{-1} \sum_{t=1}^T E_{\bar{P}_*^\nu} \left[ \sup_{\theta \in B(\delta, \theta_*)} \|\bar{\Delta}_t(\theta)(X_0^t) - \bar{\Delta}_\infty(\theta_*)(X_{-\infty}^t)\| \right] = o(1).$$

In turn, the LHS is bounded by

$$\begin{aligned} & T^{-1} \sum_{t=1}^T \left\| \sup_{\theta \in B(\delta, \theta_*)} \|\Delta_{t,0}(\theta) - \Delta_{t,-\infty}(\theta_*)\| \right\|_{L^2(\bar{P}_*^\nu)} \\ & \times \left( \left\| \sup_{\theta \in B(\delta, \theta_*)} \|\Delta_{t,0}(\theta)\| \right\|_{L^2(\bar{P}_*^\nu)} + \left\| \sup_{\theta \in B(\delta, \theta_*)} \|\Delta_{t,-\infty}(\theta)\| \right\|_{L^2(\bar{P}_*^\nu)} \right). \end{aligned}$$

By Lemma 25(3), the first term in the RHS is bounded (up to constants) by  $T^{-1} \sum_{t=1}^T t^{1-1/p}$ . The second term in the RHS is bounded by Lemma 25(1). Thus, under Assumption 8, the whole expression converges to zero and the desired result follows.

STEP 3. We next show that, for any  $L \equiv L_T$  such that  $\lim_{T \rightarrow \infty} L_T = \infty$  and  $L_T(\ddot{w}(T^{-1/2} \log \log T) \log \log T + r_T + T^{-1/2}) = o(1)$ ,

$$\left\| \sum_{j=0}^{T-1} (1 - j/T) \gamma_{j+1}(\theta_*) - \sum_{j=0}^{L_T-1} \omega(j, L) \hat{\gamma}_{T,j+1,0}(\hat{\theta}_{T,\nu}) \right\| = o_{\bar{P}_*^\nu}(1),$$

where, for any  $\tau \in \{1, \dots, L_T\}$  and any  $M \leq 1$ ,

$$\theta \mapsto \hat{\gamma}_{T,\tau,M}(\theta) \equiv T^{-1} \sum_{t=1}^{T-\tau} \Delta_{t+\tau,M}(\theta)(X_M^{t+\tau}) \Delta_{t,M}(\theta)(X_M^t)^\top$$

(recall that  $\Delta_{t,0}(\theta)(X_0^t) = \nabla_{\theta} p_t^\nu(X_t | X_0^{t-1}, \theta)$ ).

Putting  $\hat{\gamma}_{T,\tau} \equiv \hat{\gamma}_{T,\tau,-\infty}$ , we have, by the triangle inequality,

$$\begin{aligned}
& \left\| \sum_{j=0}^{T-1} (1-j/T) \gamma_{j+1}(\theta_*) - \sum_{j=0}^{L_T-1} \omega(j, L) \hat{\gamma}_{T,j+1,0}(\hat{\theta}_{T,\nu}) \right\| \\
& \leq \left\| \sum_{j=0}^{T-1} (1-j/T) \gamma_{j+1}(\theta_*) - \sum_{j=0}^{L_T-1} \omega(j, L) \gamma_{j+1}(\theta_*) \right\| \\
& \quad + \left\| \sum_{j=0}^{L_T-1} \omega(j, L) \{ \gamma_{j+1}(\theta_*) - \hat{\gamma}_{T,j+1}(\theta_*) \} \right\| \\
& \quad + \left\| \sum_{j=0}^{L_T-1} \omega(j, L) \{ \hat{\gamma}_{T,j+1}(\theta_*) - \hat{\gamma}_{T,j+1}(\hat{\theta}_{T,\nu}) \} \right\| \\
& \quad + \left\| \sum_{j=0}^{L_T-1} \omega(j, L) \{ \hat{\gamma}_{T,j+1,0}(\hat{\theta}_{T,\nu}) - \hat{\gamma}_{T,j+1}(\hat{\theta}_{T,\nu}) \} \right\|.
\end{aligned}$$

We now bound each term in the RHS individually.

By assumption, for any  $l \geq 0$ ,  $\|\gamma_l(\theta_*)\| \leq v(l)$  and thus, for any  $L$ ,

$$\begin{aligned}
\left\| \sum_{j=0}^{T-1} (1-j/T) \gamma_{j+1}(\theta_*) - \sum_{j=0}^{L-1} \omega(j, L) \gamma_{j+1}(\theta_*) \right\| & \leq \left\| \sum_{j=L}^{T-1} \gamma_{j+1}(\theta_*) \right\| \\
& \quad + \left\| \sum_{j=0}^{L-1} \{ (1-j/T) - \omega(j, L) \} \gamma_{j+1}(\theta_*) \right\| \\
& \leq \sum_{j=L}^{\infty} v(j) \\
& \quad + \left\| \sum_{j=0}^{L-1} \{ (1-j/T) - \omega(j, L) \} \gamma_{j+1}(\theta_*) \right\|.
\end{aligned}$$

Since,  $v$  is integrable, the first term in the RHS converges to zero as  $L$  diverges. Furthermore, since  $\omega(\cdot, \cdot)$  is bounded,  $\|\gamma_{j+1}(\theta_*)\| \leq v(j+1)$ , which is integrable, and  $(1-j/T) - \omega(j, L)$  converges to zero pointwise in  $j$  as  $T$  (and thus  $L = L_T$ ) diverges so, by the dominated convergence theorem, the second term also converges to zero as  $T$  (and thus  $L = L_T$ ) diverges. Therefore, for any  $\epsilon > 0$ , there exists  $T_\epsilon$  such that, for all  $T \geq T_\epsilon$ ,

$$\left\| \sum_{j=0}^{T-1} (1-j/T) \gamma_{j+1}(\theta_*) - \sum_{j=0}^{L_T-1} \omega(j, L_T) \gamma_{j+1}(\theta_*) \right\| \leq \epsilon.$$

We now consider

$$\bar{P}_*^\nu \left( \left\| \sum_{j=0}^{L-1} \omega(j, L) \{ T^{-1} \sum_{t=1}^{T-j} \Delta_{t+j, -\infty}(\theta_*) (X_{-\infty}^{t+j}) \Delta_{t, -\infty}(\theta_*) (X_{-\infty}^t)^\top - \gamma_j(\theta_*) \} \right\| \geq \delta \right)$$

for any  $\delta > 0$ . Since  $\sum_{j=0}^{L-1} \omega(j, L) \leq L$ , it follows that this expression is bounded above by

$$\bar{P}_*^\nu \left( \max_{j \in \{0, \dots, L\}} \left\| T^{-1} \sum_{t=1}^{T-j} \Delta_{t+j, -\infty}(\theta_*) (X_{-\infty}^{t+j}) \Delta_{t, -\infty}(\theta_*) (X_{-\infty}^t)^\top - \gamma_j(\theta_*) \right\| \geq \delta/L \right).$$

By similar arguments to those presented in Step 2 and Birkhoff's ergodic theorem, it follows that, for each  $L$ ,

$$\max_{j \in \{0, \dots, L\}} \left\| T^{-1} \sum_{t=1}^{T-j} \Delta_{t+j, -\infty}(\theta_*) (X_{-\infty}^{t+j}) \Delta_{t, -\infty}(\theta_*) (X_{-\infty}^t)^\top - \gamma_j(\theta_*) \right\| = o_{\bar{P}_*^\nu}(1).$$

By Lemma 24, for each  $L$ , there exists a positive sequence  $(r_T)_T$  such that  $r_T = o(1)$  and  $\bar{P}_*^\nu \left( \max_{j \in \{0, \dots, L\}} \left\| T^{-1} \sum_{t=1}^T \Delta_{t+j, -\infty}(\theta_*) (X_{-\infty}^{t+j}) \Delta_{t, -\infty}(\theta_*) (X_{-\infty}^t)^\top - \gamma_j(\theta_*) \right\| \geq r_T \right) = o(1)$ . Thus, by setting  $\delta = 2r_T L$ , it follows that, for any  $\epsilon > 0$ , there exists  $T_\epsilon$  such that, for all  $T \geq T_\epsilon$ ,

$$\bar{P}_*^\nu \left( \left\| \sum_{j=0}^{L-1} \omega(j, L) \{ T^{-1} \sum_{t=1}^{T-j} \Delta_{t+j, -\infty}(\theta_*) (X_{-\infty}^{t+j}) \Delta_{t, -\infty}(\theta_*) (X_{-\infty}^t)^\top - \gamma_j(\theta_*) \} \right\| \geq 2r_T L \right) \leq \epsilon.$$

By Theorem 4,  $\hat{\theta}_{\nu, T} \in B(T^{-1/2} \log \log T, \theta_*)$  w.p.a.1- $\bar{P}_*^\nu$ . Hence, for any  $\epsilon > 0$  there exists  $T_\epsilon$  such that for, all  $T \geq T_\epsilon$ ,

$$\begin{aligned} & \bar{P}_*^\nu \left( \left\| \sum_{j=0}^{L-1} \omega(j, L) \{ \hat{\gamma}_{T, j+1}(\theta_*) - \hat{\gamma}_{T, j+1}(\hat{\theta}_{\nu, T}) \} \right\| \geq L \ddot{\omega}(T^{-1/2} \log \log T) \log \log T \right) \\ & \leq \bar{P}_*^\nu \left( \sup_{\theta \in B(T^{-1/2} \log \log T, \theta_*)} \left\| \sum_{j=0}^{L-1} \omega(j, L) \{ \hat{\gamma}_{T, j+1}(\theta_*) - \hat{\gamma}_{T, j+1}(\theta) \} \right\| \geq L \ddot{\omega}(T^{-1/2} \log \log T) \log \log T \right) \\ & \quad + \bar{P}_*^\nu(\hat{\theta}_{\nu, T} \notin B(T^{-1/2} \log \log T, \theta_*)) \\ & \leq \bar{P}_*^\nu \left( \sup_{\theta \in B(T^{-1/2} \log \log T, \theta_*)} \sum_{j=0}^{L-1} \omega(j, L) \left\| \hat{\gamma}_{T, j+1}(\theta_*) - \hat{\gamma}_{T, j+1}(\theta) \right\| \geq L \ddot{\omega}(T^{-1/2} \log \log T) \log \log T \right) + \epsilon. \end{aligned}$$

Moreover, by the Markov inequality,

$$\begin{aligned}
& \bar{P}_*^\nu \left( \sup_{\theta \in B(T^{-1/2} \log \log T, \theta_*)} \sum_{j=0}^{L-1} \omega(j, L) \|\hat{\gamma}_{T,j+1}(\theta_*) - \hat{\gamma}_{T,j+1}(\theta)\| \geq L\ddot{w}(T^{-1/2} \log \log T) \log \log T \right) \\
& \leq \frac{1}{L\ddot{w}(T^{-1/2} \log \log T) \log \log T} E_{\bar{P}_*^\nu} \left[ \sup_{\theta \in B(T^{-1/2} \log \log T, \theta_*)} \sum_{j=0}^{L-1} \omega(j, L) \|\hat{\gamma}_{T,j}(\theta_*) - \hat{\gamma}_{T,j}(\theta)\| \right] \\
& \leq \frac{1}{L\ddot{w}(T^{-1/2} \log \log T) \log \log T} \sum_{j=0}^{L-1} \omega(j, L) E_{\bar{P}_*^\nu} \left[ \sup_{\theta \in B(T^{-1/2} \log \log T, \theta_*)} \|\tilde{\Delta}_{j,-\infty}(\theta_*) - \tilde{\Delta}_{j,-\infty}(\theta)\| \right] \\
& \leq \frac{\sum_{j=0}^{L-1} \omega(j, L) \ddot{w}(T^{-1/2} \log \log T)}{L\ddot{w}(T^{-1/2} \log \log T) \log \log T},
\end{aligned}$$

where the last line follows from Lemma 25 and  $\tilde{\Delta}_{t,-M}(\theta) \equiv \Delta_{t,-M}(\theta) \Delta_{0,-M}(\theta)^\top$  for any  $t \in \{0, \dots, T, \dots\}$  and  $M \in \mathbb{N} \cup \{\infty\}$ . Since  $\sum_{j=0}^{L-1} \omega(j, L)/L \leq 1$ , the last expression is less than  $\epsilon$  for sufficiently large  $T$ . Thus,

$$\bar{P}_*^\nu \left( \left\| \sum_{j=0}^{L-1} \omega(j, L) \{\hat{\gamma}_{T,j+1}(\theta_*) - \hat{\gamma}_{T,j+1}(\hat{\theta}_{\nu,T})\} \right\| \geq L\ddot{w}(T^{-1/2} \log \log T) \log \log T \right) \leq \epsilon.$$

Finally, since, for any  $\tau \in \{1, \dots, L\}$ ,  $\theta \mapsto \hat{\gamma}_{T,\tau,0}(\theta) \equiv T^{-1} \sum_{t=1}^{T-\tau} \Delta_{t+\tau,0}(\theta) (X_0^{t+\tau}) \Delta_{t,0}(\theta) (X_0^t)^\top$ , by Lemma 25(3) (with  $M = t$ ),

$$\left\| \sum_{j=0}^{L-1} \omega(j, L) \{\hat{\gamma}_{T,j+1}(\hat{\theta}_{\nu,T}) - \hat{\gamma}_{T,j+1,0}(\hat{\theta}_{\nu,T})\} \right\|_{L^2(\bar{P}_*^\nu)} \lesssim \sum_{j=0}^{L-1} \omega(j, L) T^{-1} \sum_{t=1}^{T-j} t^{1-1/p}.$$

By the proof of Lemma 18,  $T^{-1/2} \sum_{t=1}^T t^{1-1/p}$  vanishes; thus, the RHS is of order  $o(LT^{-1/2})$ .

Therefore, we have shown that, for any  $\epsilon > 0$ , there exists  $T_\epsilon$  such that, for all  $T \geq T_\epsilon$ ,

$$\bar{P}_*^\nu \left( \left\| \sum_{j=0}^{T-1} (1-j/T) \gamma_{j+1}(\theta_*) - \sum_{j=0}^{L_T-1} \omega(j, L) \hat{\gamma}_{T,j+1,0}(\hat{\theta}_{\nu,T}) \right\| \geq \epsilon + L_T (\ddot{w}(T^{-1/2} \log \log T) \log \log T + 2r_T + T^{-1/2}) \right) \leq \epsilon,$$

where the  $\epsilon$  inside the probability arises from bounding  $\left\| \sum_{j=0}^{T-1} (1-j/T) \gamma_{j+1}(\theta_*) - \sum_{j=0}^{L_T-1} \omega(j, L_T) \gamma_{j+1}(\theta_*) \right\|$  and requires  $L_T$  to diverge. Therefore, by taking  $L \equiv L_T$  such that  $\lim_{T \rightarrow \infty} L_T = \infty$  and  $L_T (\ddot{w}(T^{-1/2} \log \log T) \log \log T + r_T + T^{-1/2}) = o(1)$ , we establish the desired result.  $\square$

### SM.9.1 Proofs of Supplementary Lemmas

*Proof of Lemma 24.* Since  $(X_T)_{T=0}^\infty$  converges to zero in probability, for any  $\epsilon > 0$ , there exists  $T_\epsilon$  such that

$$\Pr(|X_T| \geq \epsilon) \leq \epsilon,$$

for all  $T \geq T_\epsilon$ . Now consider the sequence  $\epsilon_l = 1/2^l$  for  $l \in \{0, 1, \dots\}$ . For each  $l$ , let  $T_l$  be the smallest  $T \geq T_{l-1}$  ( $T_{-1}$  is set to 0) for which

$$\Pr(|X_T| \geq 1/2^l) \leq 1/2^l.$$

Such  $T$  always exists by the definition of convergence in probability.

Next, we construct  $(r_T)_{T=0}^\infty$ . For any  $T \in \{0, 1, \dots\}$ , there exists an  $l$  for which  $T_l \leq T \leq T_{l+1}$ , so set  $r_T = 1/2^l$ . First, observe that for any  $\delta > 0$ , there exists a sufficiently large  $T_\delta$  for which the corresponding  $l$  is such that  $1/2^l < \delta$ . Thus,  $r_T$  converges to 0 as  $T$  diverges. Second, take any  $\epsilon > 0$  and choose  $l_\epsilon$  such that  $1/2^{l_\epsilon} \leq \epsilon$ , and finally, take  $T_\epsilon \equiv T_{l_\epsilon}$ . For any  $T \geq T_\epsilon$ , it follows that

$$\Pr(|X_T| \geq r_T) = \Pr(|X_T| \geq 1/2^{l_\epsilon}) \leq 1/2^{l_\epsilon} \leq \epsilon,$$

as desired. □

*Proof of Lemma 25.* We show that  $\|\sup_{\theta \in B(\delta, \theta_*)} \|\Delta_{0,-\infty}(\theta)\|\|_{L^2(\bar{P}_*^\nu)}$  is bounded ( $\delta$  is as in Assumption 7) and that  $\Delta_{t,-\infty}$  and  $\Delta_{t,-\infty}\Delta_{0,-\infty}$  are continuous in  $L^1(\bar{P}_*^\nu)$  norm, i.e., for any  $\epsilon > 0$ , there exists  $\delta > 0$  such that

$$\omega(\delta) \equiv \max_t \left\| \sup_{\|\theta - \theta_0\| < \delta} \{\Delta_{t,-\infty}(\theta)\Delta_{t,-\infty}(\theta)^\top - \Delta_{t,-\infty}(\theta_0)\Delta_{t,-\infty}(\theta_0)^\top\} \right\|_{L^1(\bar{P}_*^\nu)} \leq \epsilon \quad (20)$$

and

$$\ddot{\omega}(\delta) \equiv \max_t \left\| \sup_{\|\theta - \theta_0\| < \delta} \{\Delta_{t,-\infty}(\theta)\Delta_{0,-\infty}(\theta)^\top - \Delta_{t,-\infty}(\theta_0)\Delta_{0,-\infty}(\theta_0)^\top\} \right\|_{L^1(\bar{P}_*^\nu)} \leq \epsilon. \quad (21)$$

We also show that there exist a constant  $C < \infty$  such that, for any  $t$  and  $M$ ,

$$\left\| \sup_{\theta \in B(\delta, \theta_*)} \Delta_{t,-\infty}(\theta) - \Delta_{t,t-M}(\theta) \right\|_{L^2(\bar{P}_*^\nu)} \leq CM^{1-1/p}. \quad (22)$$

Moreover, by Assumption 8,  $p \in (0, 2/3)$  and thus the RHS vanishes as  $M$  diverges.

We first establish (22). To do so, observe that, by inspection of the proof of Lemma 18, it follows that the conclusion of that lemma holds uniformly in  $\theta$  (and also in  $t$ ), i.e.,

$$\| \sup_{\theta \in B(\delta, \theta_*)} \Delta_{t, -\infty}(\theta) - \Delta_{t, t-M}(\theta) \|_{L^2(\bar{P}_*^\nu)} \lesssim \sum_{j=[t-M/2]}^{t-1} \varrho(j, t-M) + \sum_{j=[t-M]}^{[t-M/2]-1} \varrho(t-1, j). \quad (23)$$

By the definition of  $\varrho$  and stationarity, we have that, for any  $j \geq k$ ,  $\varrho(j, k) = \varrho(j-k, 0)$  and thus  $\sum_{j=[t-M/2]}^{t-1} \varrho(j, t-M) \leq \sum_{j=[t-M/2]}^{t-1} 1/(j-(t-M))^{1/p} \leq \int_{[M/2]+1}^M 1/(x)^{1/p} dx \leq \frac{p}{1-p} (M/2)^{1-1/p}$ , and  $\sum_{j=[t-M]}^{[t-M/2]-1} \varrho(t-1-j, 0) \leq \sum_{j=[M/2]-1}^{M-1} 1/(x)^{1/p} \leq \frac{p}{1-p} (M/2)^{1-1/p}$ . Thus,

$$\| \sup_{\theta \in B(\delta, \theta_*)} \Delta_{t, -\infty}(\theta) - \Delta_{t, t-M}(\theta) \|_{L^2(\bar{P}_*^\nu)} \lesssim M^{1-1/p},$$

as desired.

Since, under Assumption 7,  $\| \sup_{\theta \in B(\delta, \theta_*)} \| \Delta_{0, -M}(\theta) \| \| \|_{L^2(\bar{P}_*^\nu)} < \infty$  for any finite  $M$ , (22) implies that  $\| \sup_{\theta \in B(\delta, \theta_*)} \| \Delta_{0, -\infty}(\theta) \| \| \|_{L^2(\bar{P}_*^\nu)}$  is bounded.

We show next that (21) holds (the proof of (20) is completely analogous and is, therefore, omitted). To this end, observe that

$$\begin{aligned} \| \sup_{\|\theta - \theta_0\| < \delta} \tilde{\Delta}_{t, -\infty}(\theta) - \tilde{\Delta}_{t, -\infty}(\theta_0) \|_{L^1(\bar{P}_*^\nu)} &\leq \| \sup_{\theta \in B(\delta, \theta_*)} \tilde{\Delta}_{t, -\infty}(\theta) - \tilde{\Delta}_{t, t-M}(\theta) \|_{L^1(\bar{P}_*^\nu)} \\ &\quad + \| \sup_{\{\|\theta - \theta_0\| < \delta\} \cap B(\delta, \theta_*)} \tilde{\Delta}_{t, t-M}(\theta) - \tilde{\Delta}_{t, t-M}(\theta_0) \|_{L^1(\bar{P}_*^\nu)} \\ &\quad + \| \sup_{\theta_0 \in B(\delta, \theta_*)} \tilde{\Delta}_{t, -\infty}(\theta_0) - \tilde{\Delta}_{t, t-M}(\theta_0) \|_{L^1(\bar{P}_*^\nu)} \\ &\equiv Term_{1,t,M} + Term_{2,t,M} + Term_{3,t,M}, \end{aligned}$$

where, for any  $t \in \{0, \dots, T, \dots\}$  and  $M \in \mathbb{N} \cup \{\infty\}$ ,  $\tilde{\Delta}_{t, -M}(\theta) \equiv \Delta_{t, -M}(\theta) \Delta_{0, -M}(\theta)^\top$ . We now bound each of these terms to obtain the desired result.

Regarding terms 1 and 3, by simple algebra and the fact that  $\| \sup_{\theta \in B(\delta, \theta_*)} \| \Delta_{0, -M}(\theta) \| \| \|_{L^2(\bar{P}_*^\nu)} < \infty$  for any  $M \in \mathbb{N} \cup \{\infty\}$ , it follows that, for some finite  $C < \infty$ ,

$$Term_{1,t,M} + Term_{3,t,M} \leq C \| \sup_{\theta \in B(\delta, \theta_*)} \| \Delta_{t, -\infty}(\theta) - \Delta_{t, t-M}(\theta) \| \| \|_{L^2(\bar{P}_*^\nu)},$$

and, by (22), the RHS is bounded by  $O(M^{1-1/p})$ . Therefore, under Assumption 8, for any  $\epsilon > 0$ , there exists an  $M$  such that, uniformly over  $t$ ,  $Term_{1,t,M} + Term_{3,t,M} \leq \epsilon$ . Henceforth, fix this  $M$ .

Regarding term 2, observe that  $M < \infty$  and that  $\tilde{\Delta}_{t,t-M}$  is the product of two functions that are comprised of  $M$ -term-sums of products of  $\theta \mapsto \log p_\theta(x, s, x')$ ,  $\theta \mapsto \log Q_\theta(x, s, x')$  and their derivatives, all of which are continuous functions by Assumption 6. Thus, it can be shown that  $\Delta_{t,t-M}$  is continuous, thereby implying that, for any  $\epsilon > 0$ , there exists some  $\delta_{M,\epsilon}$  (which could always be chosen to be smaller than  $\delta > 0$ ) such that  $Term_{2,t,M} < \epsilon$ . This completes the proof.  $\square$

## SM.10 Additional Empirical Example

In this empirical example, we examine the predictive ability of an index of leading indicators for regime changes in U.S. output growth. The model, based on Filardo [1994], allows for shifts in the mean output growth, and is given by

$$Y_t = \mu_0 + \mu_1 S_t + \sum_{i=1}^4 \phi_i (Y_{t-i} - \mu_0 - \mu_1 S_{t-i}) + \sigma_1 U_{1,t}, \quad (24)$$

$$Z_t = \mu_2 + \sum_{i=1}^4 \psi_i Z_{t-i} + \sigma_2 U_{2,t}, \quad (25)$$

with the hidden, two-state Markov chain  $(S_t)_t$  being governed by the transition probabilities

$$\Pr(S_t = s \mid Z_{t-1} = z, S_{t-1} = s) = [1 + \exp(-\alpha_s - \beta_s z)]^{-1}, \quad s \in \{0, 1\}, \quad (26)$$

and  $(U_{1,t}, U_{2,t})_t$  postulated to be i.i.d.  $\mathcal{N}\left(\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix}\right)$  and independent of  $(S_t)_t$ . Here,  $Y_t$  stands for the growth rate of total industrial production and  $Z_t$  stands for the growth rate in the Composite Index of Eleven Leading Indicators (CLI). The data, taken from [Kim and Nelson, 1999, Ch. 4], are monthly, span the period 1948:1–1991:4, and are transformed in the way detailed in Filardo [1994]. Note that the analysis in Filardo [1994] is based on a slightly longer sample period ending in 1992:8. The likelihood ratio test of Hansen [1992] rejects the hypothesis that  $\mu_0 = \mu_1$  in (24).

We compute two sets of ML estimates: partial ML estimates based on (24) and (26) (as in Filardo [1994]) and joint ML estimates based on the system (24)–(26). Results are presented in Table 6, with estimated standard errors given in parentheses. The latter are obtained from the “sandwich” estimator defined in Theorem 5(b); the weights  $\omega(\cdot, L_T)$  are generated from the Parzen kernel and the truncation parameter  $L_T$  is determined using the automatic plug-in method

Table 6: ML Estimates (Real Output, Index of Leading Indicators)


Partial ML		Joint ML			
$\mu_0$	-0.8567 (0.3257 )	$\mu_0$	-0.7892 (0.2786)		
$\mu_1$	1.3762 (0.3257)	$\mu_1$	1.3099 (0.2179)	$\mu_2$	0.0090 (0.0678)
$\phi_1$	0.1949 (0.0784)	$\phi_1$	0.1952 (0.0694)	$\psi_1$	0.3769 (0.0490)
$\phi_2$	0.0802 (0.0540)	$\phi_2$	0.1278 (0.0493)	$\psi_2$	0.1891 (0.0538 )
$\phi_3$	0.1094 (0.0602)	$\phi_3$	0.1161 (0.0453)	$\psi_3$	0.0049 (0.0486 )
$\phi_4$	0.1229 (0.0611)	$\phi_4$	0.1169 (0.0484)	$\psi_4$	0.0570 (0.0515 )
$\alpha_0$	1.6311 (0.5351)	$\alpha_0$	1.7158 (0.6145)	$\sigma_2$	0.7729 (0.0437)
$\beta_0$	-1.0428 (0.7048)	$\beta_0$	-0.8408 (0.4131)	$\rho$	0.4498 (0.0345)
$\alpha_1$	4.3743 (0.6891 )	$\alpha_1$	4.1188 (0.7618)		
$\beta_1$	1.7669 (0.4805)	$\beta_1$	1.6020 (0.8209)		
$\sigma_1$	0.6969 (0.0386)	$\sigma_1$	0.6928 (0.0393)		

of Andrews [1991]. The differences between partial and joint ML estimates are not very substantial, a finding which is not perhaps surprising in view of the fact that the estimated conditional correlation  $\rho$  is moderate (0.4498). Conventional  $t$ -type tests based on joint ML estimates reject the hypotheses that  $\beta_0$  and  $\beta_1$  are zero, suggesting that the lagged CLI has significant information about the probability of switching between the two Markov regimes. Needless to say, inference based on the partial ML estimator must be viewed with caution unless  $\rho = 0$ .

It is worth noting that, although partial ML and joint ML estimates of the parameters are not substantially different, the inferred probabilities that  $S_t = 1$ , conditional on sample information available at time  $t$ , based on these estimates are not always close. This can be seen in Figure 1, which plots the difference between the inferred probabilities based on the two sets of estimates. The estimated covariate-dependent transition probabilities based on the two sets of ML estimates (not shown) are, on the other hand, fairly similar.



Figure 1: Difference between the inferred probability of  $S_t = 1$  based on partial ML and joint ML estimates.



## Supplemental Material References

- D. W. K. Andrews. Heteroskedasticity and autocorrelation consistent covariance matrix estimation. *Econometrica*, 59:817–858, 1991.
- K. B. Athreya and S. N. Lahiri. *Measure Theory and Probability Theory*. Springer, New York, 2006.
- P. J. Bickel, Y. Ritov, and T. Rydén. Asymptotic normality of the maximum-likelihood estimator for general hidden Markov models. *Annals of Statistics*, 26:1614–1635, 1998.
- A. V. Carter and D. G. Steigerwald. Testing for regime switching: A comment. *Econometrica*, 80:1809–1812, 2012.
- Y. A. Davydov. Mixing conditions for Markov chains. *Theory of Probability and Its Applications*, 18:312–328, 1973.
- R. Douc, G. Fort, E. Moulines, and P. Soulier. Practical drift conditions for subgeometric rates of convergence. *Annals of Applied Probability*, 14:1353–1377, 2004a.
- R. Douc, É. Moulines, and T. Rydén. Asymptotic properties of the maximum likelihood estimator in autoregressive models with Markov regime. *Annals of Statistics*, 32: 2254–2304, 2004b.
- A. J. Filardo. Business-cycle phases and their transitional dynamics. *Journal of Business and Economic Statistics*, 12:299–308, 1994.

- M. Hairer and J. C. Mattingly. Yet another look at Harris' ergodic theorem for Markov chains. In R. Dalang, M. Dozzi, and F. Russo, editors, *Seminar on Stochastic Analysis, Random Fields and Applications VI*, pages 109–117. Springer, Basel, 2011.
- B. E. Hansen. The likelihood ratio test under nonstandard conditions: Testing the Markov switching model of GNP. *Journal of Applied Econometrics*, 7:S61–S82 (Erratum: 11, 195–198), 1992.
- C.-J. Kim and C. R. Nelson. *State-Space Models with Regime Switching*. MIT Press, Cambridge, Massachusetts, 1999.
- T. A. Louis. Finding the observed information matrix when using the EM algorithm. *Journal of the Royal Statistical Society, Series B*, 44:226–233, 1982.
- G. J. McLachlan and D. Peel. *Finite Mixture Models*. Wiley, New York, 2000.
- J. Stachurski. *Economic Dynamics: Theory and Computation*. MIT Press, Cambridge, Massachusetts, 2009.
- L. Tierney. Introduction to general state-space Markov chain theory. In W. R. Gilks, S. Richardson, and D. J. Spiegelhalter, editors, *Markov Chain Monte Carlo in Practice*, pages 59–74. Chapman & Hall/CRC, London, 1996.
- H. White. *Asymptotic Theory for Econometricians*. Revised Edition, Academic Press, London, 2001.
- B. Yu. Rates of convergence for empirical processes of stationary mixing sequences. *Annals of Probability*, 22:94–116, 1994.