L^{∞} RESOLVENT BOUNDS FOR STEADY BOLTZMANN'S EQUATION

KEVIN ZUMBRUN

ABSTRACT. We derive lower bounds on the resolvent operator for the linearized steady Boltzmann equation over weighted L^{∞} Banach spaces in velocity, comparable to those derived by Pogan&Zumbrun in an analogous weighted L^2 Hilbert space setting. These show in particular that the operator norm of the resolvent kernel is unbounded in $L^p(\mathbb{R})$ for all 1 , resolvingan apparent discrepancy in behavior between the two settings suggested by previous work.

1. INTRODUCTION

In this note, we derive an L^{∞} lower bound on the resolvent operator for the linearized steady Boltzmann equation, in the process resolving a discrepancy between results of [2] and [4]. Let $L_{r,\xi}^{\infty}$ denote the space of functions $h(\xi)$, $\xi \in \mathbb{R}^3$, with finite norm $\|h\|_{L^{\infty}_{r,\xi}} := \sup_{\xi \in \mathbb{R}^3} (1+|\xi|)^r |h(\xi)|$. A key element in the study [2] of existence of invariant manifolds for the steady Boltzmann equation with hard sphere potential is the resolvent estimate [2, Thm. 11, (108)]

(1.1)
$$\|h\|_{L^{\infty}_{3,\xi}}(x) \le C \int_{\mathbb{R}} e^{-\beta |x-y|} \|g\|_{L^{\infty}_{2,\xi}}(y) dy, \quad \beta > 0$$

for solutions of the linearized inhomogeneous equation with "microscopic" data,

(1.2)
$$\xi_1 \partial_x h - Lh = g, \qquad g \in \mathrm{Ker} L^\perp$$

where $L := \overline{M}^{-1/2} \mathscr{L} \overline{M}^{1/2}$, with \mathscr{L} the linearized collision operator about a reference Maxwellian where $L := M + \mathcal{L} M + \mathcal{L}$, with \mathcal{L} the interface constant operator about a reference maxwellian $\overline{M}(\xi) = ce^{-|\xi-v|^2/d}$, and $h := (f - \overline{M})/\overline{M}^{1/2}$, with $f(x,\xi,t)$ the standard Boltzmann variable denoting density of particles of velocity $\xi \in \mathbb{R}^3$ at point $(x,t) \in \mathbb{R}^2$ in space and time [1, 2].¹ Estimate (1.1) may be expressed equivalently as a bound $|R(x,\cdot,\cdot)|_{L^{\infty}_{2,\xi} \to L^{\infty}_{3,\xi}} \leq Ce^{-\beta|x|}, \beta > 0$ on

the resolvent kernel $R(x,\xi,\xi_*)$ for (1.2), that is, the kernel of resolvent operator

$$\mathscr{R} = (\xi_1 \partial_x - L)^{-1}|_{\operatorname{Ker} L^{\perp}}, \qquad (\mathscr{R}g)(x,\xi) = \int_{\mathbb{R}} \int_{\mathbb{R}^3} R(x-y,\xi,\xi_*)g(y,\xi_*)d\xi_*dy.$$

(See [2] for a construction of R.) Either implies by standard convolution bounds uniform estimates

(1.3)
$$||h||_{L^{\infty}(\mathbb{R}, L^{\infty}_{3,\xi})} \le C ||g||_{L^{q}(\mathbb{R}, L^{\infty}_{2,\xi})}, \quad 1 \le q \le \infty.$$

Taking $q = \infty$ in (1.3), and noting [1, 3] that the associated bilinear collision operator Q(h, h) of the full, nonlinear equation is bounded from $L_{3,\xi}^{\infty} \times L_{3,\xi}^{\infty} \to L_{2,\xi}^{\infty}$, yields $L^{\infty}(\mathbb{R}, L_{3,\xi}^{\infty})$ -contractivity of the map $h \to \mathscr{R}Q(h,h)$, giving the basis for the fixed-point iteration schemes defined in [2].

By comparison, in a nearby Hilbert-space setting,² it has been shown [4] that the operator norm of the resolvent kernel is unbounded in all $L^p(\mathbb{R})$, by explicit computation using self-adjointness of L to diagonalize by unitary transformation. This is in sharp contrast with (1.1), especially since finite-dimensional intuition would suggest that optimal bounds be found in coordinates for which L is unitarily diagonalizable. Here, we resolve this apparent paradox by (i) establishing the following result contradicting (1.3), hence (1.1), and (ii) identifying a corresponding error in [2].

Research of K.Z. was partially supported under NSF grant no. DMS-0300487.

¹ Here, $(\cdot)^{\perp}$ denotes orthogonal complement with respect to the usual, unweighted space $L^2 \supset L_{2,\xi}^{\infty}$ in ξ . ² Precisely, the L^2 norm in ξ , after the rescaling $h \to \langle \xi \rangle^{1/2} h$, $g \to \langle \xi \rangle^{-1/2} g$, where $\langle \xi \rangle := (1 + |\xi|^2)^{1/2}$; see [3, 4].

Proposition 1.1. The solution operator for (1.2) does not satisfy $||h||_{L^{\infty}(\mathbb{R},L^{\infty}_{3,\xi})} \leq C||g||_{L^{q}(\mathbb{R},L^{\infty}_{2,\xi})}$ for any $q < \infty$, C = C(q). Likewise, $|R(x,\cdot,\cdot)|_{L^{\infty}_{2,\xi} \to L^{\infty}_{3,\xi}}$ is unbounded in $L^{p}(\mathbb{R})$, p > 1.

Proof. Recall [1] that the linearized collision operator L appearing in (1.2) may be decomposed as $L = -\nu(\xi) + K$, where $\nu(\xi) \sim \langle \xi \rangle := (1 + |\xi|^2)^{1/2}$ and $(Kh)(\xi) = \int_{\mathbb{R}^3} k(\xi, \xi_*)h(\xi_*)d\xi_*$, with kernel $|k(\xi, \xi_*)| \leq C|\xi - \xi_*|^{-1}e^{-c|\xi-\xi_*|^2}$. By $||\xi|^{-1}e^{-c|\xi|^2}||_{L^1} < \infty$ and standard convolution bounds, K is bounded on $L^{\infty}(\xi)$. Similarly, using the inequality $\langle \xi \rangle / \langle \xi - \xi_* \rangle \leq C \langle \xi_* \rangle$ and $||\xi|^{r-1}e^{-c|\xi|^2}||_{L^1} < \infty$, we obtain the standard result $|K|_{L^{\infty}_{r,\xi}} < +\infty$ [1, 2]. Recall, further, that KerL is finite-dimensional, tangent to the 5-dimensional manifold of Maxwellians [1]. Defining $\mathscr{S} = (\xi_1 \partial_x + \nu(\xi))^{-1}$, we have the explicit solution formula $(\mathscr{S}g)(x,\xi) = \int_{\mathbb{R}} S_{\xi}(x-y)g(y,\xi)dy$, with $S_{\xi}(\theta) = \xi_1^{-1}e^{-\nu(\xi)\xi_1^{-1}|\theta|}$ for $\theta\xi_1 > 0$, 0 otherwise. Computing $||S_{\xi}(\cdot)||_{L^1(\mathbb{R})} = 1/\nu(\xi) \leq C/\langle \xi \rangle$, we have by standard convolution bounds is $|\mathscr{S}|_{L^{\infty}(\mathbb{R}, L^{\infty}_{2,\xi}) \to L^{\infty}(\mathbb{R}, L^{\infty}_{3,\xi})} < +\infty$.

Writing (1.2) as $(\xi_1 \partial_x + \nu(\xi))h = Kh + g$, applying \mathscr{S} , and rearranging, gives $\mathscr{S}g = h - \mathscr{S}Kh$, hence $|\mathscr{S}g|_{L^{\infty}(\mathbb{R}, L^{\infty}_{3,\xi})} \leq C|h|_{L^{\infty}(3, L^{\infty}_{r,\xi})}$ by boundedness of $|K|_{L^{\infty}_{3,\xi}}$, $|\mathscr{S}|_{L^{\infty}(\mathbb{R}, L^{\infty}_{3,\xi})}$. Thus, (1.3) would imply a comparable bound $|\mathscr{S}g|_{L^{\infty}(\mathbb{R}, L^{\infty}_{3,\xi})} \leq C|g|_{L^{q}(\mathbb{R}, L^{\infty}_{2,\xi})}$ on \mathscr{S} . This is easily seen to be false for $g \in L^{\infty}_{2,\xi}$, using test functions $g_{\alpha}(x,\xi) = \langle \xi \rangle^{-2} \alpha^{-1/q} e^{-\alpha^{-1}|x|}$, $||g_{\alpha}||_{L^{q}(\mathbb{R}, L^{\infty}_{2,\xi})} \equiv 1, \alpha \to 0$, and computing $\mathscr{S}g_{\alpha}(0, (\alpha, 1, 0)) = (1 + \alpha^2)^{-1} \int_0^{+\infty} \alpha^{-1} e^{-(1 + \alpha^2)^{1/2} \alpha^{-1} y} \alpha^{-1/q} e^{-\alpha^{-1} y} dy \sim \alpha^{-1/q} \to +\infty$. Using linear combinations of members g_{α_j} of the infinite-dimensional family $\{g_{\alpha}\}$, we may obtain a contradiction also for g ranging on any finite-codimension subspace, in particular on Ker L^{\perp} . \Box

The argument of [2]. In [2], the authors study the steady Boltzmann equation by a timeregularization scheme based on detailed pointwise bounds on the time-evolutionary Green function $G(x, t, \xi, \xi_*)$, through the relation $R = \int_{\mathbb{R}^+} G dt$. However, this analysis contains a key error. By [2, (67)]: $S_x b = \int_{\mathbb{R}^+} \int_{\mathbb{R}^3} G(x, t, \cdot, \xi_*) \xi_{1,*} b(\xi_*) d\xi_* dt$ and the symmetric [2, (68)], [2, (52)] is equivalent to $\|\int_{\mathbb{R}^+} \int_{\mathbb{R}^3} G(x, t, \cdot, \xi_*) \xi_{1,*} b(\xi_*) d\xi_* dt \|_{L^{\infty}_{3,\xi}} \leq C e^{-\beta |x|} \|b\|_{L^{\infty}_{3,\xi}}$. In [2, (102)], this is used to estimate

(1.4)
$$\|\int_{\mathbb{R}^+} \int_{\mathbb{R}^3} G(x,t,\cdot,\xi_*) \tilde{b}(\xi_*) \, d\xi_* dt \|_{L^{\infty}_{3,\xi}} \le C e^{-\beta|x|} \|\tilde{b}\|_{L^{\infty}_{2,\xi}}.$$

in effect asserting $\|\xi_1^{-1}\tilde{b}\|_{L^{\infty}_{3,\xi}} \lesssim \|\tilde{b}\|_{L^{\infty}_{2,\xi}}$, or $|\xi_1|^{-1} \lesssim (1+|\xi|)^{-1}$ - evidently false for ξ_1 small. But, (1.4) is the basis for (1.1), the resolvent estimate underlying the fixed-point constructions of [2].

Discussion and open problems. Proposition 1.1 shows that [2, Thm. 11] is incorrect, invalidating the conclusions of [2] on existence of invariant manifolds. Our proof of Proposition 1.1 depends on the property that the principle part \mathscr{S} of the resolvent \mathscr{R} is unbounded from $L^q(x, L^{\infty}_{2,\xi}) \to L^{\infty}(x, L^{\infty}_{3,\xi})$ for $q < \infty$, which is much stronger than $|S(x, \cdot, \cdot)|_{L^{\infty}_{2,\xi} \to L^{\infty}_{3,\xi}} \notin L^p$ for 1 = 1/p + 1/q. Estimate $|S(x, \cdot, \cdot)|_{L^{\infty}_{2,\xi} \to L^{\infty}_{3,\xi}} \sim |x|^{-1}$ as $x \to 0$ shows that the latter holds also at the boundary p = 1; we conjecture that it holds for R as well. This, and the determination of upper bounds (1.3), $q = \infty$ on \mathscr{R} we regard as very interesting open problems.

References

- H. Grad, Asymptotic theory of the Boltzmann equation. II, 1963 Rarefied Gas Dynamics (Proc. 3rd Internat. Sympos., Palais de l'UNESCO, Paris, 1962), Vol. I pp. 26–59 Academic Press, New York.
- T.-P. Liu and S.-H. Yu, Invariant Manifolds for Steady Boltzmann Flows and Applications, Arch. Rational Mech. Anal. 209 (2013) 869–997.
- [3] G. Métivier and K. Zumbrun, Existence and sharp localization in velocity of small-amplitude Boltzmann shocks, Kinet. Relat. Models 2 (2009), no. 4, 667–705.
- [4] A. Pogan and K. Zumbrun, Stable manifolds for a class of degenerate evolution equations and exponential decay of kinetic shocks, preprint, 39 pp.; arXiv:1607.03028.

INDIANA UNIVERSITY, BLOOMINGTON, IN 47405 $E\text{-mail}\ address:$ kzumbrun@indiana.edu