

FREE BOUNDARY MINIMAL SURFACES OF UNBOUNDED GENUS

DANIEL KETOVER

ABSTRACT. For each integer $g \geq 1$ we use variational methods to construct in the unit 3-ball B a free boundary minimal surface Σ_g of symmetry group \mathbb{D}_{g+1} . For g large, Σ_g has three boundary components and genus g . As $g \rightarrow \infty$ the surfaces Σ_g converge as varifolds to the union of the disk and critical catenoid. These examples are the first with genus greater than 1 and were conjectured to exist by Fraser-Schoen. We also construct several new free boundary minimal surfaces in B with the symmetry groups of the cube, tetrahedron and dodecahedron. Finally, we prove that free boundary minimal surfaces isotopic to those of Fraser-Schoen can be constructed variationally using an equivariant min-max procedure. We also prove an ϵ -regularity theorem for free boundary minimal surfaces in B .

1. INTRODUCTION

Denote by U an open domain in \mathbb{R}^{n+1} . A hypersurface $\Sigma^n \subset U$ with $\partial\Sigma \subset \partial U$ is called a *free boundary minimal surface* if it is minimal in U and

$$(1.1) \quad \Sigma \perp \partial U.$$

Free boundary minimal surfaces arise variationally as critical points to the volume functional for hypersurfaces with boundary in ∂U where one permits variations that move $\partial\Sigma$ within ∂U . They have been studied already since the 1940s by Courant [C]. Recently Fraser-Schoen [FS] have found some connections between free boundary minimal surfaces and extremal metrics for Steklov eigenvalues.

As for existence theory, Grüter-Jost ([GJ],[GJ2]) used the min-max method pioneered by Almgren-Pitts [Pi] and Simon-Smith [SS] to produce a free boundary minimal disk in three-dimensional convex bodies. Dropping the convexity assumption, Li [Li] recently obtained a related existence result. Applying White's degree theory [W], Maximo-Nunes-Smith [MNS] showed the existence of a free boundary minimal annulus

The author was partially supported by NSF-PRF DMS-1401996 as well as ERC-2011-StG-278940.

in convex bodies. De Lellis-Ramic [DR] have produced free boundary surfaces in higher dimensions (see also [LZ]). See [FGM] for explicit higher dimensional examples.

In this paper, we will be concerned with free boundary minimal surfaces in the standard three-ball B in \mathbb{R}^3 . Very few explicit examples are known. The simplest examples are the flat disks through the origin. Secondly, there is the “critical catenoid” which is a free boundary annulus obtained by rescaling the catenoid in \mathbb{R}^3 so that it intersects the boundary of the unit ball orthogonally. Throughout this paper, D will denote the unique free boundary minimal disk in the xy plane, and C the unique critical catenoid that is rotationally symmetric about the z axis.

For each $k \geq 2$, Fraser-Schoen [FS] constructed a free boundary minimal surface F_k resembling a “doubling” of the flat disk D in the sense that $F_k \rightarrow 2D$ as $k \rightarrow \infty$. The surfaces F_k consist of two disks joined by many half-necks at the boundary ∂B and so have genus 0 and k ends. Later Folha-Pacard-Zolotareva [FPZ] using gluing methods gave another construction of F_k when k is large. They also constructed related genus 1 examples G_k (when k is large) by adding a catenoidal neck at the center of the F_k examples joining the two layers.

It has been an open question whether one can construct higher genus examples. Fraser-Schoen conjectured (Section 1 in [FS2]) that there should be a sequence of free boundary minimal surfaces with genus approaching infinity converging to the union of the critical catenoid and the disk. In this paper, we confirm their conjecture and prove:

Theorem 1.1. *For each integer $g \geq 1$, there exists a free boundary minimal surface Σ_g in the unit 3-ball B with dihedral symmetry \mathbb{D}_{g+1} that is not the flat disk D . When g is large, Σ_g has three boundary components and genus g . Moreover,*

$$(1.2) \quad \Sigma_g \rightarrow D \cup C \text{ in the varifold sense as } g \rightarrow \infty.$$

Furthermore, $|\Sigma_g| < |D| + |C|$ for all $g \geq 1$ (where $|\Sigma|$ denotes the 2-dimensional Hausdorff measure of Σ).

The proof of Theorem 1.1 was inspired by a sketch of Pitts-Rubinstein [PR] for a variational construction of the minimal surfaces of Costa-Hoffman-Meeks (see also [HM]). The minimal surfaces obtained by Theorem 1.1 can be interpreted as free boundary analogs to these surfaces. Surprisingly, in the proof of Theorem 1.1 we need the sharp isoperimetric inequality for the ball B to prove that the sweepouts we consider are nontrivial.

When g is small, it is possible that the surface Σ_g produced by Theorem 1.1 has one boundary component in ∂B rather than three. One could rule this out if one could show that the only free boundary minimal surface in B with one boundary component is a flat disk. This would imply the existence of a free boundary minimal surface of each genus g .

We also apply the catenoid estimate [KMN] to show:

Theorem 1.2. *For each $k \geq 2$, a free boundary minimal surface isotopic to F_k can be constructed variationally through a one-parameter equivariant min-max procedure.*

Finally we produce several new genus zero examples associated with the Platonic solids:

Theorem 1.3. *There exists a free boundary minimal surface in B with octahedral symmetry of genus 0 and 6 ends, an example with tetrahedral symmetry of genus 0 and 4 ends, and an example of genus 0 and 12 ends of dodecahedral symmetry.*

Let us describe the genus 0 surface with 6 ends produced by Theorem 1.3. Consider the graph \mathcal{G} in B consisting of the union of the x , y and z axes in B . The free boundary minimal surface we construct is isotopic to the boundary of a tubular neighborhood of \mathcal{G} , resembling a three-dimensional “cross.”

Since the methods involved in proving Theorem 1.1 arise from a global variational principle, they are quite versatile and apply in other ambient geometries. They can be used to give a min-max construction of the self-shrinkers discovered by Kapouleas-Kleene-Møller [KKM] and independently Nguyen [Ng]:

Theorem 1.4. *(Kapouleas-Kleene-Møller [KKM], Nguyen [Ng]) For g large enough, there exist a self-shrinker N_g with dihedral symmetry $\mathbb{D}_{2(g+1)}$ (acting by rotations about the z -axis) with one end and genus g . As $g \rightarrow \infty$, N_g converge as varifolds to the union of the self-shrinking xy -plane and self-shrinking sphere.*

While our existence theorem could produce examples of low genus with the symmetries of those of Theorem 1.4 when g is small, we would need to take g large to rule out that the min-max limit we obtain has many ends.

In gluing constructions, one always needs to take the genus along the desingularizing curves to be large. One advantage of the variational approach is that one can produce minimal surfaces with low genus as in Theorem 1.1. However, one still has to rule out various compressions

(“neck-pinches”) that occur as the min-max sequence converges to its limiting minimal surface. Thus uniqueness or classification theorems are still needed when the genus is low to rule out unwanted behavior such as the surfaces Σ_g having genus greater than zero but one end. When the genus is large, typically one can take a limit in the underlying symmetry group to see that the limiting stationary varifold one produces has enough symmetries to be easily classifiable (see for instance the proof of Theorem 1.1 or Theorem 1.7 in [KMN]).

On a related note, using higher parameter families, Marques-Neves [MN] proved that manifolds with positive Ricci curvature contain infinitely many embedded minimal surfaces. Li-Zhou [LZ] have adapted their argument to the free boundary setting. Of course, manifolds with many symmetries such as the 3-ball trivially contain infinitely many minimal free boundary surfaces by considering all rotations of the disk or critical catenoid. Thus it is not so clear how to use Marques-Neves’ method to obtain new minimal surfaces. Recently Aiex [Ai] has shown that the set of min-max surfaces produced by the Marques-Neves procedure is non-compact, which should rule out that the infinitely many surfaces produced are simply rotations of the basic ones. Still, these methods would not allow any precise control on the topological type.

Let us explain the geometric idea behind the existence of the surfaces Σ_g purported by Theorem 1.1. They arise via an equivariant min-max procedure. The sweepouts we consider are somewhat unusual in that they do not start and end at surfaces with zero area as one usually has in min-max theory. Instead, they begin and end at the unit disk D (thus one can alternatively think of them as being parameterized by \mathbb{S}^1 instead of $[0, 1]$). Let us describe these sweepouts. There’s an optimal sweepout C_t of B by annuli orthogonal to D with the critical catenoid sitting in the middle of the foliation. We consider the family $\Sigma_t = C_t \cup D$, desingularized by adding in genus g in a \mathbb{D}_{g+1} -equivariant manner along the circle of intersection between D and C_t . By the \mathbb{D}_{g+1} -equivariance, each surface in this family divides B into two components of equal volume. By Almgren’s solution [Al] of the isoperimetric problem for B , it follows that each surface in Σ_t has area at least π . The key observation is that the sweepout Σ_t (and any sweepout in the equivariant saturation of Σ_t) *interchanges* the two hemispheres of B determined by D .

One then considers the width W for the min-max problem for sweepouts in the equivariant saturation of Σ_t . In order to show that this sweepout is non-trivial, we must show that W is greater than the area of the disk, π . If $W = \pi$, then by definition of width, one could find sweepouts all of whose slices have area very close to π . By Almgren’s

result again [Al], it follows that all of these slices are very close as varifolds to the disk D as D is the only equivariant surface solving the isoperimetric problem. But one can show that such sweepouts cannot swap the two hemispheres.

While generalizing the min-max theory to the free boundary setting has already been carried out ([DR], [LZ], [LZ2]), from a min-max perspective, there are some new ingredients that are needed in Theorem 1.1. In min-max theory, the regularity of the surface produced is proved by showing that the min-max sequence is approximated locally by stable surfaces and using the compactness theorem for such surfaces due to Schoen [S].

In [Ke2], this theory was extended to considering sweepouts satisfying a symmetry. There one restricts to sweepouts of G -equivariant surfaces for some group of isometries G so that each surface in the sweepout intersects the singular set of the group action transversally. One shows there (Section 4 in [Ke2]) that one can produce G -stable replacements (i.e. minimal surfaces that are stable among G -equivariant deformations) and moreover that G -stability is equivalent to stability. Thus Schoen's compactness result applies.

In the setting of Theorem 1.1, there is a key difference. The min-max sequence contains segments of the singular set of the action of \mathbb{D}_{g+1} on B . In balls about such segments one can find approximating \mathbb{Z}_2 -stable surfaces. But because the surface contains the axis, \mathbb{Z}_2 -stability *does not* seem to imply stability. Thus we need to replace Schoen's estimates in the regularity theory with the fact that minimal surfaces of bounded area and genus have a convergent subsequence [CS]. The convergence may be non-smooth over finitely many points, but this does not impede the regularity theory or genus bounds. The arguments thus are inherently two-dimensional and only apply to the Simon-Smith theory [SS].

The organization of this paper is as follows. In Section 2 we prove an ϵ -regularity type theorem expressing the rigidity of the free boundary disk. In Section 3 we state the equivariant min-max theorem we need in this paper in the free boundary setting. In Section 4 we prove Theorem 1.1. In Section 5 we construct the examples of Fraser-Schoen, and in Section 6 the Platonic examples of genus 0. Finally in Section 7 we prove the equivariant min-max theorem.

It has been brought to my attention that N. Kapouleas and M. Li have applied gluing methods to obtain an analog of Theorem 1.1 when g is large.

Acknowledgements: I thank Brian White for a conversation and Otis Chodosh for bringing the work of Volkman [V] to my attention. I

thank Fernando Marques and André Neves for discussions and encouragement.

2. RIGIDITY OF FREE BOUNDARY MINIMAL DISK

We need the following strong rigidity statement for the free boundary minimal disk which may be of independent interest:

Proposition 2.1. *There exists $\epsilon > 0$ so that if Σ is an embedded free boundary minimal surface in B and*

$$(2.1) \quad |\Sigma| < \pi + \epsilon,$$

then Σ is a flat free boundary disk D .

One can think of Proposition 2.1 as strong form of Allard's regularity theorem [A] for free boundary minimal surfaces that holds *up the boundary*. Indeed, Allard's theorem loosely speaking gives that a minimal surface in the ball with area close enough to the flat disk is a graph over this flat disk in a sub-ball. Adding the free boundary condition to these assumptions, Proposition 2.1 gives a stronger conclusion as one obtains graphicality up to the boundary and moreover, one gets a unique graph, i.e., the flat disk itself. This improvement is one of the special features of free boundary minimal surfaces. For related (sharp) gap theorems for the free boundary minimal disk, see [AN].

The proof of Proposition 2.1 uses a monotonicity formula for free boundary minimal surfaces that was discovered in various guises independently by Fraser-Schoen (Theorem 5.4 in [FS]), Ros-Vergasta [RV], Brendle [B], and obtained in most general form by Volkman [V].

To state the consequence of Volkman's formula that we need, let Σ be a free boundary minimal surface in B . For any x_0 in the support of Σ , there holds (by plugging $H = 0$ into Theorem 3.4.3 and using equation (4) in [V]):

$$(2.2) \quad \int_{\Sigma} \frac{|(x - x_0)^{\perp}|^2}{|x - x_0|^4} \leq |\Sigma| - |D|.$$

The vector $(x - x_0)^{\perp}$ is the component of the vector $x - x_0$ that is orthogonal to the tangent plane $T_{x_0}\Sigma$. Volkman's monotonicity formula compares a quantity at two scales – letting one scale approach 0, and the other approach ∞ in his formula one obtains (2.2).

One easy consequence of (2.2) is the fact that D has the smallest area among free boundary minimal surfaces, and is the unique surface with this property (cf. [B]). The integrated term in (2.2)

$$(2.3) \quad E(x_0, \Sigma) := \int_{\Sigma} \frac{|(x - x_0)^{\perp}|^2}{|x - x_0|^4}$$

is a kind of “tilt-excess,” that controls the deviation of Σ from a plane. A convenient feature is that the point x_0 in (2.2) is arbitrary, while the right hand side of (2.2) does not depend on x_0 . A key point is that $E(x_0, \Sigma)$ is *scale invariant* if Σ is rescaled about x_0 . In other words

$$(2.4) \quad E(x_0, \Sigma) = E(0, \tau(\Sigma - x_0)) \text{ for any } \tau > 0.$$

Proof of Proposition 2.1

We first establish the following claim:

claim: There exists $\epsilon > 0$ and $C > 0$ so that whenever Σ is an embedded free boundary minimal surface with $|\Sigma| \leq \pi + \epsilon$ there hold the following curvature bounds

$$(2.5) \quad \sup_{x \in \Sigma} |A|^2(x) \leq C.$$

This claim easily implies Proposition 2.1. Indeed, suppose the claim is true. If Proposition 2.1 failed, it means there is a sequence of free boundary surfaces that are different from flat disks, with areas approaching π and by the claim, uniformly bounded curvature. Thus by Theorem 5.1 in [FL], Σ_i converge smoothly (up to the boundary) to a disk (see also Section 5 of [LZ2]). Since the convergence is smooth, it follows that Σ_i are all disks for i large enough. But by a theorem of Nitsche [N], the only free boundary minimal disks are the flat ones. This is a contradiction.

It remains to prove the claim. We will use a blowup argument (cf. [W2]). Suppose the claim is false. Thus there is a sequence Σ_i of free boundary minimal surfaces in B where

$$(2.6) \quad |\Sigma_i| - \pi \rightarrow 0$$

while

$$(2.7) \quad A_i := \sup_{x \in \Sigma_i} |A|^2(x) \rightarrow \infty.$$

For each i , choose a point $x_i \in \Sigma_i$ where the sup A_i is attained in (2.7) (of course x_i may be in the boundary of Σ_i). Then consider the sequence of surfaces

$$(2.8) \quad \tilde{\Sigma}_i := A_i(\Sigma_i - x_i),$$

which satisfy

$$(2.9) \quad \sup_{x \in \tilde{\Sigma}_i} |A|^2 \leq 1$$

and

$$(2.10) \quad |A|_{\tilde{\Sigma}_i}^2(0) = 1.$$

Since dilations preserve angles, the surface $\tilde{\Sigma}_i$ still satisfies the free boundary condition for the new domain $A_i(B - x_i)$.

By the curvature bounds (2.9) it follows that $\tilde{\Sigma}_i \rightarrow \Sigma_\infty$ smoothly. Note that Σ_∞ is either complete without boundary or (after rotating Σ_j potentially) is contained in a half-space and its boundary is contained in a plane. The convergence in this latter case is smooth up to the boundary by the boundary Schauder estimates of Agmon-Douglis-Nirenberg (Theorem 9.1 in [Ag]) because the free boundary condition implies that the surfaces locally near the boundary satisfy an elliptic equation with homogeneous boundary conditions. By the smooth convergence $\Sigma_j \rightarrow \Sigma_\infty$ (up to the boundary if it exists) and (2.10) we still have:

$$(2.11) \quad |A|_{\Sigma_\infty}^2(0) = 1.$$

In light of (2.6) and (2.2), we obtain

$$(2.12) \quad E(x_0, \Sigma) \rightarrow 0.$$

By the scale invariance (2.4), we obtain

$$(2.13) \quad E(0, \tilde{\Sigma}_i) \rightarrow 0.$$

Since the convergence $\tilde{\Sigma}_j \rightarrow \Sigma_\infty$ is smooth, we have

$$(2.14) \quad 0 \leq E(0, \Sigma_\infty) \leq \liminf E(0, \tilde{\Sigma}_i) = 0.$$

Since $E(0, \Sigma_\infty) = 0$, it follows that Σ_∞ is a plane or half-plane. But this violates (2.11). Thus the claim is established. \square

3. EQUIVARIANT MIN-MAX THEORY

In [Ke2], an equivariant min-max theory was developed to produce min-max minimal surfaces with symmetries. Throughout this paper G will denote either \mathbb{Z}_n , \mathbb{D}_n or one of the three groups associated to the Platonic solids T_{12} , O_{24} , or I_{60} acting standardly on B .

For any $x \in B$ we first define the *isotropy subgroup* G_x at x as:

$$G_x = \{g \in G \mid gx = x\}$$

We then define the *singular locus* of the group action as points with nontrivial isotropy subgroup:

$$\mathcal{S} = \{x \in B \mid G_x \neq \{e\}\}$$

The set \mathcal{S} consists of straight line segments emanating from the origin. In [Ke2], one considers sweepouts of B that intersect \mathcal{S} transversally. But the sweepouts we consider in Section 4 to prove Theorem 1.1 are not transverse to the singular set – they *contain* line segments which are part of \mathcal{S} . Thus the theory developed in [Ke2] needs to be

extended to that setting. We make the following more general definition:

A (*genus g*) G -sweepout of B is a family of closed sets $\{\Sigma_t\}_{t=0}^1$, continuously varying in the Hausdorff topology such that:

- i. Σ_t is a smooth embedded surface of genus g for $0 < t < 1$ varying smoothly
- ii. Σ_0 and Σ_1 are the union of a smooth surface together with a collection of arcs
- iii. Each Σ_t is G -equivariant, i.e. $g(\Sigma_t) = \Sigma_t$ for $0 \leq t \leq 1$ and all $g \in G$

Remark 3.1. *Note that if Σ_t contains an arc $\mathcal{A} \subset \mathcal{S}$ for some t , then \mathcal{A} is in the support of Σ_t for all t . This follows from the fact the surfaces vary smoothly and one cannot smoothly “push off the axis.” See Lemma 3.6 in [Ke2].*

Given such a G -sweepout $\{\Sigma_t\}_{t=0}^1$ we may define the G -equivariant saturation $\Pi = \Pi_{\{\Sigma_t\}}$ identically as in [Ke2]. We can then define the min-max width:

$$(3.1) \quad W_{\Pi}^G = \inf_{\{\Lambda_t\} \in \Pi} \sup_{t \in [0,1]} |\Lambda_t|.$$

We can then consider a sequence of sweepouts $\{\Sigma_t\}^i$ the area of whose maximal slice converges to W_{Π}^G . From $\{\Sigma_t\}^i$ we may then choose a sequence of slices $\Sigma_i := \Sigma_{t_i}^i$ with area converging to W_{Π}^G . Such a sequence of surfaces we will call a *min-max sequence*.

With this notation we have the following Min-Max theorem (where the relevant terms in ii) are explained in the remarks following the theorem):

Theorem 3.2. *If*

$$(3.2) \quad W_{\Pi}^G > \max(|\Sigma_0|, |\Sigma_1|)$$

then there exists a min-max sequence Σ_j converging as varifolds to $n\Gamma$, where Γ is a smooth embedded connected free boundary minimal surface in B and n is a positive integer. Moreover, the following statements hold:

- i. $W_{\Pi}^G = n|\Gamma|$
- ii. *For j large enough, after performing finitely many surgeries on Σ_j and discarding some components, each remaining component of Σ_j is isotopic to Γ and there are n such components. Thus*

$$(3.3) \quad n(\text{genus}(\Gamma)) \leq g.$$

- iii. If the surfaces Σ_t all contain a segment of isotropy \mathbb{Z}_2 , then Γ contains this segment as well and n is odd.
- iv. If the surfaces Σ_t are orthogonal to or are disjoint from an arc $\mathcal{A} \subset \mathcal{S}$ of isotropy \mathbb{Z}_2 , then Γ either intersects \mathcal{A} orthogonally (if at all) or else contains \mathcal{A} and n is even.
- v. Γ intersects arcs of isotropy \mathbb{Z}_n (for $n \neq 2$) orthogonally (if at all).

Remark 3.3. The connectedness of Γ follows from the Frankel-type property of embedded free boundary minimal surfaces in convex bodies: any two must intersect (Lemma 2.4 in [FL]).

Remark 3.4. By “surgeries” in ii) is meant either “ G -equivariant neck-pinch” (cf. Remark 1.6 in [Ke2]) or “collapse of topology” which we will define below. A “ G -equivariant neckpinch,” is obtained in three possible ways: 1) by removing a cylinder and gluing in two disks so that the cylinder and disks as well as the ball they bound is disjoint from \mathcal{S} , 2) removing a cylinder centered around \mathcal{S} and adding in two disks, where each disk intersects \mathcal{S} once, or as a “half neckpinch” at the boundary. This third type of surgery is needed because of the free boundary and is defined as follows. Let Σ be a surface in B with $\partial\Sigma \subset \partial B$. Consider a disk $D \subset \Sigma$ with $\partial D = \cup_{i=1}^4 \gamma_i$ where γ_i are smooth arcs and the four arcs are concatenated in increasing order to give ∂D . The arc γ_1 is contained in ∂B , γ_2 is contained in the interior of B , γ_3 contained in ∂B and γ_4 is contained in the interior of B . The arcs γ_2 and γ_4 bound the disks $D_1 \subset \overline{B}$ and $D_3 \subset \overline{B}$ where $\partial D_1 = \gamma_2 \cup \alpha_1$ where α_1 is an arc contained in ∂B . Likewise $\partial D_3 = \gamma_4 \cup \alpha_3$ where α_3 is an arc contained in ∂B . The surgery on Σ is the removal of D from Σ and addition of D_1 and D_2 . This third type of surgery is loosely speaking a “half neck-pinch” at the boundary ∂B .

Finally, in the case where the sweepout surfaces Σ_t contain an arc \mathcal{A}' of isotropy \mathbb{Z}_2 , we include “collapse of topology” in the admissible surgeries. This consists of the following. Fix $x \in \overline{\mathcal{A}'}$ so that G_x is either \mathbb{D}_k or \mathbb{Z}_2 . Fix a small G_x -invariant ball B_x about x and let Σ be a G_x -invariant surface in B_x with $\partial\Sigma \subset \partial B_x$. We say Σ' arises from Σ by collapse of topology if Σ' is the varifold limit of G -equivariant isotopies supported in B_x and if there is a sequence of neck-pinches supported in B_x (not necessarily G -equivariant), so that Σ' arises topologically from Σ after performing these neck-pinches.

Remark 3.5. The reason we include the “collapse of topology” in the admissible surgeries in ii) is the following. In the case where the sweepout surfaces contain an axis of the singular set, we only prove that the \mathbb{Z}_2 -stable replacements V_j around the axis converge smoothly

to their smooth limit V away from finitely many points, and near such points it may not be possible to perform G -equivariant neck-pinches to obtain V .

As an example of this phenomenon, consider the Costa-Hoffman-Meeks minimal surface C_g of genus g in \mathbb{R}^3 , that has dihedral symmetry and contains the \mathbb{Z}_2 isotropy axes of the symmetry group. One can consider the family of rescalings $\{\lambda_i C_g\}$ for $\lambda_i \rightarrow 0$ that converges to the plane with multiplicity 3. The convergence is smooth away from the origin, where the genus is collapsing. There is *no way* to perform equivariant neck-pinches on this family to obtain a surface isotopic to three disjoint sheets as any neck-pinch will break the dihedral symmetry. On the other hand, the configuration consisting of three planes is indeed in the limit of G -equivariant isotopies of C_g and it is achievable through surgeries (though not equivariant ones).

Remark 3.6. While we only state Theorem 3.2 in the case where the ambient manifold is a three-ball, since all considerations are local, it is clear that it holds for a general closed three-manifold.

We defer the proof of Theorem 3.2 til Section 7.

4. PROOF OF THEOREM 1.1

4.1. Sweepouts. Let us first construct the sweepouts we will need for the surfaces constructed in Theorem 1.1. In this section, denote by D the unit disk in the xy plane.

Fix $g \geq 1$. Consider the group \mathbb{D}_{g+1} acting on B by rotations and of $2\pi/(g+1)$ about the z axis, and also rotations of π about the $g+1$ line segments $\{L_i\}_{i=0}^g$ (setting $\theta_i = i\pi/(g+1)$):

$$(4.1) \quad L_i := \{(r \cos(\theta_i), r \sin(\theta_i), 0) \in \mathbb{R}^3 \mid -1 \leq r \leq 1\}$$

Denote by C the unique critical catenoid with symmetry group \mathbb{D}_{g+1} encircling the z -axis. There exists a sweepout $\{C_t\}_{t=-1}^1$ of B with dihedral symmetry \mathbb{D}_{g+1} with the following properties:

- i. $C_{-1} = \{z\text{-axis}\} \cap B$
- ii. $C_1 = \partial D$
- iii. C_t is a smooth annulus for $-1 < t < 1$.
- iv. C_0 is the critical catenoid C
- v. $|C_t| < |C| - At^2$ for some $A > 0$
- vi. $C_t \cap D$ is a round circle R_t for all $-1 < t \leq 1$ and R_t sweep-out D .
- vii. C_t is orthogonal to D for $-1 < t < 1$

Note that the only singular slices in the foliation C_t occur when $t = 1$ and $t = -1$ when C_t consists of one dimensional graphs.

In order to construct C_t , we start with the critical catenoid C_0 . It is stable if one considers deformations that vanish on ∂B . On the other hand, if one considers more general deformations, then its Morse index has recently been computed to be 4 ([SZ], [T], [D]).

Let ϕ be a smooth function defined on C_0 and let n be a choice of unit normal on C_0 . Then the formula for the second derivative of area is (c.f. Section 5.1 [D]):

$$(4.2) \quad \left. \frac{d^2}{dt^2} \right|_{t=0} |C_0 + t\phi| = \int_{\Sigma} |\nabla \phi|^2 - |A|^2 \phi^2 - \int_{\partial \Sigma} \phi^2.$$

Setting $\phi_1 = 1$ in (4.2) we obtain

$$(4.3) \quad \left. \frac{d^2}{dt^2} \right|_{t=0} |C_0 + t\phi_1| = - \int_{\Sigma} |A|^2 d\mu - \mathcal{H}^1(\partial \Sigma) < 0.$$

Thus ϕ_1 , while not an eigenfunction of the stability operator, still gives a rotationally symmetric direction for decreasing the area of the critical catenoid.

Let us now give construction of the optimal sweepout of annuli C_t . Let R denote the region of $B \setminus C_0$ that is disjoint from the z -axis. We can consider dilations of the critical catenoid λC_0 for $\lambda \geq 1$. Such dilations preserve the region R in that $\lambda R \subset R$. For some value λ_0 , $\{\lambda C_0\}_{\lambda=1}^{\lambda=\lambda_0}$ gives a foliation of R interpolating between C_0 and ∂D . Since dilating by λ takes the ball of radius $B_{1/\lambda}(0)$ to the ball B , it follows from the monotonicity formula that the area of $B \cap \lambda C_0$ is a decreasing function in λ . This gives the required foliation of R . To fill out $R' = B \setminus R$ we can argue as follows. First use ϕ_1 defined above to extend C_0 into R' to C_t for $-\epsilon \leq t \leq 0$ by

$$(4.4) \quad C_t := C_0 + t\phi_1(x)n(x).$$

In light of (4.3), $|C_t| \leq |C_0| - At^2$ for some $A > 0$.

Note that the region $R' \setminus \cup_{t \in [-\epsilon, 0]} C_t$ is mean convex and the boundary circles of $C_{-\epsilon}$ no longer bound a minimal annulus since the circles $\partial C_{-\epsilon}$ are contained in the region in ∂B where no catenoids can penetrate. Thus there is a path of rotationally symmetric surfaces beginning at $C_{-\epsilon}$ and ending at the two flat disks bounded by $\partial C_{-\epsilon}$ that increases area an arbitrarily small amount along the way. It is then easy to use parallel disks to $\partial C_{-\epsilon}$ joined by a tiny tube about the z -axis to fill out the rest of R' . This gives the required sweepout.

Let us define the singular family of sets for $-1 \leq t \leq 1$:

$$(4.5) \quad \Sigma_t = C_t \cup D.$$

We will now amend each surface Σ_t in the sweepout in a very small neighborhood of $D \cap C_t$. The $g+1$ lines L_i intersect $C_t \cap D$ in $2(g+1)$ equally spaced points and divide $C_t \cap D$ into $2(g+1)$ consecutive arcs $\{A_i\}_{i=1}^{2(g+1)}$. Along A_1 are being joined 4 pieces of smooth surfaces in an “X”: two from C_t and two from D . Desingularize this intersection along A_1 by pairing one of the pieces of C_t with one from D , and the other C_t piece with the other D piece. Then proceed to pair off the pieces in the opposite way along A_2 and continue in this alternating fashion along ∂D to arrive at a new surface $\tilde{\Sigma}_t$. Note that to preserve \mathbb{D}_{g+1} symmetry, once the desingularization has been performed along A_1 , it extends in a unique way to the rest of ∂D . Thus rotating 180° about any of the lines L_i is still a symmetry of $\tilde{\Sigma}_t$.

It is clear that we can perform this desingularization so that all changes are supported in $T_{\epsilon(t)}(C_t \cap D)$ for any suitably small continuous positive function $\epsilon(t)$ and we can moreover choose $\epsilon(t)$ to approach 0 as t approaches -1 or 1 . This desingularization is an area-decreasing procedure since desingularizing amounts to “rounding corners” and thus lowers area. We can think that each arc A_i is labelled alternatively $+$ or $-$, and combining two consecutive such arcs gives a period of the rotational symmetry of the resulting surface. Note that $\tilde{\Sigma}_t$ still contains the lines $\{L_i\}_{i=0}^g$ since the surfaces are only being adjusted in the interior of each arc A_i .

This desingularization has the effect that $\tilde{\Sigma}_t$ is now a surface of genus g with \mathbb{D}_{g+1} symmetry: rotations of angle $2\pi/(g+1)$ together with such rotations composed with a “flip” about any of the lines L_i . In summary, one obtains a sweepout $\tilde{\Sigma}_t$ satisfying the following properties (reparameterizing the sweepout by $[0, 1]$ instead of $[-1, 1]$):

- i. $|\tilde{\Sigma}_t| < |C| + |D|$ for all $0 \leq t \leq 1$
- ii. $\tilde{\Sigma}_0 = D \cup (\{z\text{-axis}\} \cap B)$
- iii. $\tilde{\Sigma}_1 = \partial D$
- iv. For each $0 < t < 1$ the genus of $\tilde{\Sigma}_t$ is g
- v. $\tilde{\Sigma}_t$ has dihedral symmetry \mathbb{D}_{g+1}
- vi. For all t , $\tilde{\Sigma}_t$ contains the lines $\{L_i\}_{i=0}^g$ and (in particular) the origin

The sweepout $\{\tilde{\Sigma}_t\}$ satisfies the definition of \mathbb{D}_{g+1} -sweepout in Section 3. Let us denote by Π the equivariant saturation of sweepouts containing $\{\tilde{\Sigma}_t\}$.

Remark 4.1. *Note that unlike in 2-dimensions, there are several different types of dihedral symmetry in 3-dimensions. The Costa-Hoffman-Meeks surfaces have an extra symmetry coming from reflections in certain planes, and their symmetry group is $\mathbb{D}_{2(g+1)}$. Our group is only half as large. We do this so that the resulting orbifold B/G has no boundary aside from $\partial B/G$ which is the setting in which equivariant min-max theory was developed in [Ke2].*

The set $B \setminus D$ has two components. Denote by C_1 the component in the northern hemisphere, and C_2 the component in the southern. The sweepout $\tilde{\Sigma}_t$ has the following key property.

- (1) For each t , $\tilde{\Sigma}_t$ divides B into two components $A(t)$ and $B(t)$ of equal volume
- (2) $A(0) = C_1$, $B(0) = C_2$
- (3) $A(1) = C_2$, $B(1) = C_1$.

In other words, in the course of the sweepout, the two hemispheres of B are swapped. One can see (1) because rotating 180° through any of the lines L_i acts by isometry to preserve the surface $\tilde{\Sigma}_t$ while interchanging the components $A(t)$ and $B(t)$.

Moreover, any $\{\Lambda_t\} \in \Pi$ satisfies (1)-(3) as well. As a consequence, for any such $\{\Lambda_t\}$ (denoting by $A(t)$ and $B(t)$ the components of $B \setminus \Lambda_t$) there is a $t \in [0, 1]$ with

$$(4.6) \quad \text{vol}(A(t) \cap C_1) = \frac{1}{2} \text{vol}(C_1) = \frac{2\pi}{3}.$$

4.2. Non-triviality of the sweepout. In this section we prove

Proposition 4.2. *For each $g \geq 1$ the width satisfies:*

$$(4.7) \quad W_g > \pi = |D|.$$

The equation (4.7) expresses the non-triviality of the sweepout and is what permits the min-max method to work.

We need the following fundamental theorem about isoperimetric surfaces in the unit ball:

Theorem 4.3. *(Almgren [Al], Bokowski-Sperner [BS], Ros [R]) The isoperimetric surfaces in B are hyperplanes through the origin or spherical caps meeting ∂B orthogonally. Thus if a surface Σ in B with $\partial \Sigma \subset \partial B$ divides B into two components with equal volume, then $|\Sigma| \geq \pi$, and equality holds if and only if Σ is the intersection of B with a hyperplane through the origin.*

Proof of Proposition 4.2:

Assume toward a contradiction that $W_g = |D| = \pi$. Thus by the

definition of width, there is a sequence of sweepouts $\{\Sigma_t\}^i$ in the saturation Π so that

$$(4.8) \quad \sup_{t \in [0,1]} |\Sigma_t^j| \leq \pi + \epsilon_j,$$

for some sequence $\epsilon_j \rightarrow 0$.

Since each surface $\{\Sigma_t\}^j$ divides B into two components of equal volume, it follows that for each fixed t , $\{\Sigma_t\}^j$ is a minimizing sequence for the isoperimetric problem in B , and thus for large j by Theorem 4.3 each surface $\{\Sigma_t\}^j$ must be close as varifolds to some disk. Since the sweepouts are \mathbb{D}_{g+1} equivariant, it follows that this disk is precisely D . In other words for some $\delta_i \rightarrow 0$, and j large enough, we have

$$(4.9) \quad \sup_{t \in [0,1]} \mathbb{F}(\Sigma_t^j, D) \leq \delta_j,$$

where \mathbb{F} denotes the \mathbb{F} -metric on varifolds.

On the other hand, for any j , the sweepout $\{\Sigma_t\}^j$ must interchange the two components C_1 and C_2 of $B \setminus D$, which readily violates (4.9).

Let us give more details. By (4.6), since each sweepout $\{\Sigma_t\}^j$ interchanges C_1 and C_2 , for each j , there is some $t_j \in [0, 1]$ so that $\Sigma_{t_j}^i$ bounds two regions R_1 , and R_2 , where

$$(4.10) \quad \text{vol}(R_1 \cap C_1) = \frac{1}{2} \text{vol}(C_1) = \frac{2}{3} \pi$$

and

$$(4.11) \quad \text{vol}(R_2 \cap C_1) = \frac{1}{2} \text{vol}(C_1) = \frac{2}{3} \pi.$$

Choose $\epsilon > 0$ and consider the cap

$$(4.12) \quad S_\epsilon := (C_1 \setminus T_\epsilon(D)) \cap B_{1-\epsilon}(0),$$

where $B_{1-\epsilon}(0)$ denotes the ball of radius $1 - \epsilon$ about the origin, and $T_\epsilon(D)$ is the ϵ -tubular neighborhood about D in \mathbb{R}^3 . The boundary of S_ϵ is contained in the sets

$$(4.13) \quad \partial S_\epsilon^1 = \{(x, y, z) \in \mathbb{R}^3 \mid z = \epsilon\} \cap B$$

together with

$$(4.14) \quad \partial S_\epsilon^2 = \{(x, y, z) \in \mathbb{R}^3 \mid z \geq \epsilon\} \cap \partial B_{1-\epsilon}(0).$$

In light of (4.9), for t_i chosen above and j large enough and any $\epsilon > 0$ and any $\eta > 0$

$$(4.15) \quad |\Sigma_{t_i}^j \cap S_\epsilon| < \eta.$$

Thus by the coarea formula

$$(4.16) \quad \int_{\epsilon}^{2\epsilon} \mathcal{H}^1(\Sigma_{t_i}^j \cap \partial S_{\sigma}^1) d\sigma \leq C |\Sigma_{t_i}^j \cap S_{\epsilon}| < C\eta.$$

By (4.16) and Sard's lemma we can find $\sigma \in [\epsilon, 2\epsilon]$ so that

$$(4.17) \quad \mathcal{H}^1(\Sigma_{t_i}^j \cap \partial S_{\sigma}^1) \leq \frac{2C\eta}{\epsilon}.$$

and

$$(4.18) \quad \Sigma_{t_i}^i \cap \partial S_{\sigma}^1$$

consists of several closed circles or half-circles. By the coarea formula again we can choose σ so that in addition ∂S_{σ}^2 consists of several closed circles or half-circles and

$$(4.19) \quad \mathcal{H}^1(\Sigma_{t_j}^j \cap \partial S_{\sigma}^2) \leq \frac{2C\eta}{\epsilon}.$$

By the isoperimetric inequality for the plane and $\partial B_{1-\sigma}$, (4.17) and (4.19) the disks D_i in ∂S_{σ} bounded by the collection of circles $\Sigma_{t_j}^j \cap \partial S_{\sigma}$ have total area:

$$(4.20) \quad \sum_i |D_i| \leq C'(\eta/\epsilon)^2.$$

Note the decomposition

$$(4.21) \quad S_{\sigma} = (\cup A_i) \cup X.$$

In (4.21), A_i are the components of the interior of the closed surface obtained by capping off a component of $\Sigma_{t_j}^j$ in ∂S_{σ} with one of the disks D_i (i.e. "filigree"). The set X is defined to be the complement of the A_i . Note that the sets A_i are not necessarily pairwise disjoint as they may be nested. Nonetheless, by the isoperimetric inequality in \mathbb{R}^3 , (4.15) and (4.20) it follows that

$$(4.22) \quad \sum_i \text{vol}(A_i) \leq C((\eta/\epsilon)^2 + C\eta)^{3/2}.$$

Since $\sigma < \epsilon$ we can estimate the volume of the shell region

$$(4.23) \quad \text{vol}(C_1 \setminus S_{\sigma}) < C\epsilon.$$

The set X is connected and so is contained in either R_1 or R_2 . Suppose without loss of generality that $X \subset R_1$. It follows that $R_2 \cap C_1$ is composed of some of the filigree sets A_i together with some parts of $C_1 \setminus S_{\sigma}$. Thus combining (4.23) with (4.22) we obtain:

$$(4.24) \quad \text{vol}(R_2 \cap C_1) \leq C((\eta/\epsilon)^2 + C\eta)^{3/2} + C\epsilon.$$

If first ϵ and then η are chosen small enough, (4.24) contradicts (4.11).

□

Remark 4.4. *As an alternative approach to proving Proposition 4.2 one could try to argue as follows. If $W_g = \pi$, then after a pull-tight procedure, each slice in a tightened sweepout Γ_t (with maximal area very close to π) is close to a stationary varifold with area π . Since each surface passes through the origin, by the monotonicity formula one obtains that all surfaces in Γ_t are close to a disk. The argument then proceeds as in Proposition 4.2. This argument is less elementary since to use the monotonicity formula it seems one needs the integrality of limits of min-max sequences, which already uses the almost minimizing property.*

4.3. Completion of the proof of Theorem 1.1. Since $W_g > \pi$ by Proposition 4.2, (3.2) is satisfied, and thus we may apply Theorem 3.2 to obtain a \mathbb{D}_{g+1} -equivariant free boundary minimal surface Σ_g in B . It remains to determine the topological type of Σ_g .

In the following, we will first enumerate the possible neck-pinches and collapse of topology that may occur and the topological type of possible min-max limits. With more work one can show that the neck-pinches we enumerate are the only possible ones. The only fact we shall use in the proof of Theorem 1.1 is that the genus of the min-max limit obtained is either g or 0 , which follows just from the equivariance. The reader may thus wish to skip to (4.25) as the intervening paragraphs below are not logically necessarily.

Recall from Remark 3.4 that there are three possible types of surgeries: \mathbb{Z}_{g+1} neckpinches occurring by removing an annulus centered around an arc of the singular set, surgeries removing a “half-annulus” at the boundary ∂B , and *ordinary* neckpinches performed in the interior of B that occur in regions disjoint from \mathcal{S} . In the following, we will describe the possible compressions.

Let us describe the potential \mathbb{Z}_{g+1} -neckpinch. We can describe this compression on the initial sweepout surfaces $\tilde{\Sigma}_t$. In the northern hemisphere, $\tilde{\Sigma}_t$ coincides with C_t . Consider the circle γ on C_t that is homotopically nontrivial obtained by intersecting C_t with a plane of constant positive z -value. The circle γ encircles the z -axis (which has isotropy \mathbb{Z}_{g+1}) and it is possible to compress along this circle. By equivariance, if this circle is compressed, so must be a corresponding circle in the southern hemisphere. After this surgery on $\tilde{\Sigma}_t$, one obtains two disks (one in each hemisphere) as well as one surface of genus g which has one boundary circle. Only this latter surface with genus g can contribute to the min-max limit since the min-max limit contains the origin in

its support. Thus if this \mathbb{Z}_{g+1} neckpinch occurs, one obtains a surface with one boundary component and genus g .

We now will argue that there are no possible \mathbb{Z}_2 -compressions or ordinary neckpinches that can bring down the genus of $\tilde{\Sigma}_t$. To see this, consider any simple closed curve γ in the interior of B representing a non-trivial homology class of $\tilde{\Sigma}_t$. Such a curve is contained in the tiny tubular neighborhood of $C_t \cap D$ on which C_t and D are desingularized to produce $\tilde{\Sigma}_t$. But such a curve has non-zero intersection number with one of the lines L_i . It follows that no such compression can occur since \mathbb{Z}_2 -neckpinches as well as ordinary neckpinches by definition are obtained by removing an annulus that is disjoint from the singular set of the group action. One way to understand this phenomenon is to consider Scherk's singly periodic surface in \mathbb{R}^3 that is the desingularization of two orthogonal planes and has dihedral symmetry. There are two ways to "snap the necks" but neither of the resulting configurations still has dihedral symmetry.

One can still collapse the genus of $\tilde{\Sigma}_g$ by "collapse of topology" as in Remark 3.5. In this case, one can use an isotopy to press all of the genus of $\tilde{\Sigma}_t$ to the origin in a \mathbb{D}_{g+1} -equivariant fashion, in which case Γ is a free boundary disk with odd multiplicity (by Theorem 3.2iii). Since $W_g < |D| + |C| < 3|D|$, it follows that the multiplicity is 1. By a theorem of Nitsche [N], this disk must be flat, violating Proposition 4.2 as $W_g > |D|$. Thus the throwing away of topology cannot occur.

Let us now consider the compressions that occur along "half-annuli" at the boundary ∂B . Let us describe the arcs along which the compression can occur in the model $\tilde{\Sigma}_t$ surface. Recall that in producing $\tilde{\Sigma}_t$, along the arc A_1 of $C_t \cap D$, the part of C_t contained in the northern hemisphere is being connected with the component of the disk $D \setminus (C_t \cap D)$ that touches ∂B . Thus there is an arc α contained on $\tilde{\Sigma}_t$ beginning at the boundary $\partial B \cap C_t$ and ending at $\partial B \cap D$. It is clear α bounds a "half-disk" and such a surgery is admissible. By the equivariance, if such a surgery is performed, there are $2(g+1)$ copies of it that must be performed concurrently. The effect of this surgery is to bring the genus of $\tilde{\Sigma}_t$ down to zero while the number of ends becomes one (though this end is rather checkered). After performing this surgery, one can see that no further ones are possible.

In total, we have shown either i) Σ_g has genus g and three ends, or else ii) Σ_g has genus g and one end, or else iii) genus zero and one end. Let us now show that when g is large, the last two cases cannot occur.

To achieve this, we prove (1.2), i.e., that

$$(4.25) \quad \Sigma_g \rightarrow D \cup C,$$

from which it follows that when g is large, Σ_g has genus g and three ends.

Consider any subsequence (not relabelled) of Σ_g . We will show that Σ_g has a subsequence converging to $D \cup C$, which implies (4.25). Let Σ_∞ denote the (free boundary) stationary integral varifold that is a subsequential limit of Σ_g as $g \rightarrow \infty$. Since Σ_g is invariant under \mathbb{D}_{g+1} , it follows that the support of Σ_∞ is invariant under rotations about the z axis.

Since Σ_g contains the lines $\{L_i\}_{i=0}^g$ that become denser and denser as $g \rightarrow \infty$, it follows that the disk D is in the support of Σ_∞ . By the monotonicity formula, it also follows that the support of Σ_∞ is connected.

Let us first consider the blowup set \mathcal{B} for the curvature of Σ_g :

$$(4.26) \quad \mathcal{B} = \{x \in B \mid \inf_{r>0} \liminf_{g \rightarrow \infty} \int_{B_r(x)} |A|_{\Sigma_g}^2 d\mu \geq \epsilon_0\}$$

The constant $\epsilon_0 > 0$ is chosen so that by the ϵ -regularity theorem of Choi-Schoen [CS], if $x \in B \setminus \mathcal{B}$, then some subsequence of Σ_g satisfies uniform curvature estimates in a neighborhood of x .

We claim:

The set $\mathcal{B} \cap \text{int}(B)$ consists of a single circle S centered about 0 in D of some radius $r \in (0, 1)$.

Let us first prove this claim. By the equivariance of Σ_g which increases as $g \rightarrow \infty$ it follows that \mathcal{B} is the union of (potentially infinitely many) round circles about the z -axis.

Let us first show that $\mathcal{B} \cap \text{int}(B)$ contains a circle in the disk D . Suppose not. Then for any sub-disk D' of D (passing to a subsequence in g) there exists a $r > 0$ so that

$$(4.27) \quad \sup_{x \in T_r(D')} |A|_{\Sigma_g}^2 \leq C$$

where $T_r(D')$ denotes the r -tubular neighborhood about D' . By (4.27), it follows that $\Sigma_g \rightarrow \Sigma_\infty$ smoothly on compact subsets of D . By the free boundary condition and the unique continuation property for minimal surfaces (Lemma 5 in [MY]) and since Σ_∞ contains D , it follows that Σ_∞ restricted to $T_r(D)$ consists of kD where k is a positive integer. By the equivariance, it follows that k is odd. Since $|\Sigma_g| < |D| + |C| < 3|D|$, it follows that $k = 1$. As Σ_∞ restricted to $T_r(D)$ is precisely D , the connectness of Σ_∞ implies that $\Sigma_\infty = D$.

To rule this situation out, observe that if indeed $\Sigma_g \rightarrow D$, then for g large enough, the area of Σ_g approaches that of D . By Proposition 2.1, this implies Σ_g is itself D , which contradicts the fact that $|\Sigma_g| > |D|$. Thus we have a contradiction, and it follows that \mathcal{B} contains a circle S in the interior of the disk D . Note that circle may be trivial, i.e. have zero radius, and we will rule this situation out later.

We claim that for some $r_i(g) \rightarrow 0$, the genus of Σ_g restricted to $T_{r_i}(S)$ is equal to g . To see this, suppose that instead $T_r(S)$ contains no genus for g large (by the equivariance, the genus g is either all contained in $T_r(S)$ or zero in $T_r(S)$). Then by Ilmanen's integrated Gauss-Bonnet argument (Lemma 1 in Lecture 3 in [I]), one obtains for some $C < \infty$

$$(4.28) \quad \sup_{g \in \mathbb{N}^+} \int_{\Sigma_g \cap T_{r/2}(S)} |A|^2 \leq C.$$

However, since $S \subset \mathcal{B}$, then by definition of \mathcal{B} and the rotational invariance we obtain

$$(4.29) \quad \sup_{g \in \mathbb{N}^+} \int_{\Sigma_g \cap T_{r/2}(S)} |A|^2 = \infty.$$

This is a contradiction.

Since we have just proved that the genus is collapsing into S , it follows that

$$(4.30) \quad \mathcal{B} \cap \text{int}(B) = S.$$

Indeed, let S_2 be another circle contained in $(\mathcal{B} \cap \text{int}(B)) \setminus S$. Since the genus of Σ_g in a neighborhood of S_2 is 0 (as the genus is contained in smaller and smaller neighborhoods about S), we can again apply Ilmanen's integrated Gauss-Bonnet argument [I], to obtain for some $\epsilon > 0$ sufficiently small and $C_\epsilon < \infty$

$$(4.31) \quad \sup_{g \in \mathbb{N}^+} \int_{\Sigma_g \cap T_\epsilon(S_2)} |A|^2 \leq C_\epsilon.$$

But since we have assumed $S_2 \subset \mathcal{B}$, then by definition of \mathcal{B} and the rotational invariance we obtain

$$(4.32) \quad \sup_{g \in \mathbb{N}^+} \int_{\Sigma_g \cap T_\epsilon(S_2)} |A|^2 = \infty.$$

The identity (4.32) contradicts (4.31) and thus in fact (4.30) holds.

To prove the claim, it now suffices to show the circle S is non-trivial, i.e. its radius r is not zero. If instead $r = 0$, then Σ_g converge to a multiple of the disk D . One can see this as follows. In this case, the curvature of Σ_g is bounded away from the origin, so that $\Sigma_\infty \setminus \{0\}$ is a smooth free boundary minimal surface (maybe with multiplicity).

Since Σ_∞ contains the disk, Σ_∞ is either D or some multiple kD . Any other component C_∞ would be a smooth rotationally symmetric disk containing the origin in the northern hemisphere, together with its mirror image in the south. But this violates the maximum principle unless both of these components are D . So in fact $\Sigma_\infty = kD$ for some integer k . We have already ruled out the case $k = 1$. From the equivariance of Σ_∞ , it follows that k is odd. But because $|\Sigma_g| < |C| + |D| < 3|D|$ we obtain $|\Sigma_\infty| < 3|D|$ and thus $k < 3$. Thus the case $r = 0$ is ruled out and r is instead some value in $(0, 1)$.

In sum we obtain that Σ_∞ is a rotationally symmetric stationary varifold in B containing D that is singular only at the circle $A(r)$ for some $0 < r < 1$. Since $\Sigma_\infty \neq D$, it follows that Σ_∞ contains in the northern hemisphere (and by symmetry in the southern too) an additional smooth rotationally symmetric minimal component Σ'_∞ with boundary $A(r)$. This component either hits the boundary ∂B or stays in the interior of B . If it stays in the interior, then by the convex hull property of minimal surfaces, Σ'_∞ is contained in D , and thus Σ_∞ has differing integer multiplicities in the disk D' bounded by $A(r)$ and its complement $D \setminus D'$. This violates the Constancy Theorem which states that a stationary varifold supported on a smooth surface is an integer multiple of the surface. If instead Σ'_∞ reaches ∂B , then because the critical catenoid is the unique rotationally symmetric free boundary surface aside from the disk, it follows that $\Sigma_\infty = D \cup C$.

This completes the proof of Theorem 1.1. □

Remark 4.5. *Assuming one knew that Σ_g had three boundary components, then instead of using Proposition 2.1 to rule out the situation where $\Sigma_g \rightarrow D$ one could alternatively apply the following argument. By Theorem 5.4 in Fraser-Schoen [FS3], since Σ_g is a free boundary minimal surface, one has*

$$(4.33) \quad |\partial\Sigma_g| = 2|\Sigma_g|.$$

Thus if

$$(4.34) \quad |\Sigma_g| \rightarrow \pi,$$

then by (4.33)

$$(4.35) \quad |\partial\Sigma_g| \rightarrow 2\pi.$$

By assumption $\partial\Sigma_g$ consists of three circles. If $\Sigma_g \rightarrow D$, then either the middle circle converges to ∂D and the top and bottom ones vanish at the north and south poles or else the three circles converge to ∂D with multiplicity 3. The first case is ruled out by applying the monotonicity

formula to Σ_g at points near the north or south pole. The second case is ruled out by (4.35) as $|\partial\Sigma_g|$ would be converging to 6π in this case.

5. VARIATIONAL CONSTRUCTION OF THE FRASER-SCHOEN EXAMPLES

In this section, we prove Theorem 1.2 which we restate

Theorem 5.1. *For each $k \geq 2$, a free boundary minimal surface isotopic to F_k can be constructed variationally through a one-parameter equivariant min-max procedure.*

Proof. Let $G = \mathbb{D}_k$. We construct a G -invariant sweepout of B as follows. For $t \in (-1, 1)$, denote

$$(5.1) \quad D_t = B \cap \{(x, y, z) \in \mathbb{R}^3 \mid z = t\}.$$

Note that

$$(5.2) \quad |D_t| = \pi(1 - t^2).$$

For $t \in [0, 1]$ consider the surfaces $\Sigma_t = D_t \cup D_{-t}$ (so that Σ_0 is the disk $D_0 = D$ with multiplicity 2). Fix k evenly spaced lines L_i of longitude joining the north pole of ∂B to the south pole. Let $L_i(\epsilon)$ be the ϵ -tubular neighborhood (in \mathbb{R}^3) of L_i intersected with B and let $P_i(\epsilon)$ denote the part of the boundary of $L_i(\epsilon)$ contained in the interior of B .

Now for some function $f(t) : [0, 1] \rightarrow [0, \delta]$ consider the surfaces

$$(5.3) \quad \tilde{\Sigma}_t = \Sigma_t \cup \bigcup_i P_i(f(t)) \setminus \left(\bigcup_i \Sigma_t \cap L_i(f(t)) \right).$$

In other words, we add to Σ_t the half-tubes P_i and remove the two half-disks in Σ_t abutting the boundary of B . By construction $\tilde{\Sigma}_t$ are \mathbb{D}_k -equivariant, have genus 0 and k ends as long as $f(t) > 0$.

Choose $f(t)$ so that $f(t) \rightarrow 0$ as $t \rightarrow 1$. In other words, as $t \rightarrow 1$, the necks disappear and $\tilde{\Sigma}_t$ converges to the graph \mathcal{G}_1 consisting of the collection of arcs L_i . Fix $\epsilon > 0$ small and enforce for f that $f(\epsilon) = 0$ and also $f(x) > 0$ for $x \in [\epsilon, 1]$.

Thus we have in light of (5.2) and the choice of f

$$(5.4) \quad |\tilde{\Sigma}_\epsilon| \leq 2|D| - 2\pi\epsilon^2$$

and in fact

$$(5.5) \quad \sup_{t \in [\epsilon, 1]} |\tilde{\Sigma}_t| \leq 2|D| - C,$$

for some $C > 0$.

For each i , let p_i denote the point $L_i \cap D$. Note that there is a retraction R_t from $D \setminus \cup_i p_i$ onto the graph \mathcal{G}_2 consisting of k equally spaced lines passing through the origin in D . Thus applying the Catenoid Estimate [KMN] and using (5.5) one can adjust the sweepout $\tilde{\Sigma}_t$ in the region $t \in [0, \epsilon]$ so that

$$(5.6) \quad \sup_{t \in [0,1]} |\tilde{\Sigma}_t| < 2|D|.$$

and $\tilde{\Sigma}_t \rightarrow \mathcal{G}_2$ as $t \rightarrow 0$.

Applying Theorem 3.2 to the family $\tilde{\Sigma}_t$ and the group G , we obtain a free boundary minimal surface F_k . The only possible neckpinches would result in two disks, which would imply $F_k = 2D$, contradicting (5.6). Thus there are no neckpinches and one obtains free boundary minimal surfaces isotopic to those of [FS]. \square

6. NEW FREE BOUNDARY MINIMAL SURFACES ASSOCIATED TO THE PLATONIC SOLIDS

In this section, we prove Theorem 1.3, which we restate:

Theorem 6.1. *There exists a free boundary minimal surface in B with octahedral symmetry of genus 0 and 6 ends, an example with tetrahedral symmetry of genus 0 and 4 ends, and an example of genus 0 and 12 ends of dodecahedral symmetry.*

Proof. Let us produce the surface of genus 0 and 6 ends, as the others follow analogously. Consider O_{24} (the symmetry group of a cube) acting on B . We can construct a O_{24} -invariant sweepout Σ_t of B as follows. Let Σ_0 be the graph \mathcal{G} consisting of the x -axis, y -axis, and z -axis restricted to B . For t small, set $\Sigma_t = \partial T_\epsilon(\mathcal{G})$. One can extend this sweepout O_{24} -equivariantly to the rest of B so that Σ_1 consists of the tessellation of ∂B by six squares.

Applying Theorem 3.2 to the sweepout Σ_t with $G = O_{24}$ one obtains an embedded connected free boundary minimal surface Γ . It is easy to see that the only possible degeneration is into several disjoint disks. But no disk is invariant under O_{24} . It follows that Γ is a free boundary minimal surface with six ends and genus zero, resembling a three-dimensional “cross.” \square

7. PROOF OF THE MIN-MAX THEOREM 3.2

We will assume the reader is familiar with the min-max construction of Simon-Smith [SS] and only focus on the changes necessary from the standard arguments. For an exposition of the theory see for instance Colding-De Lellis [CD], or [Ke] for the control on the genus of the

limiting minimal surface. In Section 2 of [Ke2] is a detailed account of the changes needed in the equivariant setting. De-Lellis-Ramic provide a detailed account of the min-max theory in the free boundary setting (the “unconstrained problem”) which includes a proof of the regularity of free boundary minimal surfaces at their free boundary.

Note that the “boundary” considered here is not entirely “free.” Consider for instance the situation of Theorem 1.1 where the sweepout surfaces contain an axis of the singular set \mathbb{Z}_2 and so cannot move off these lines. Thus the boundary is constrained to contain the points that are the intersections of these lines with ∂B . But this is not really a problem because of the Schwarz reflection principle: if a varifold is stationary with respect to variations preserving a line in the surface, it is stationary with respect to all variations (cf. Lemma 3.8 in [Ke2]).

7.1. Existence of a free boundary equivariant stationary varifold \mathcal{V} . Given a vector field χ defined on \overline{B} , denote by ϕ_χ the family of isotopies obtained by integrating χ . Let us consider the family of vector fields χ so that the corresponding isotopy $\phi_\chi(t) : \overline{B} \rightarrow \overline{B}$ satisfies

- (1) For all $t \in [0, 1]$, ϕ_t preserves ∂B .
- (2) $g\phi_t(x) = \phi_t(gx)$ for all $t, g \in G$ and $x \in \overline{B}$.

Denote by Is_{FB}^G the set of vector fields χ so that ϕ_χ satisfies items (1) and (2). The set Is_{FB}^G consists of the *free boundary equivariant vector fields*. A varifold $\mathcal{W} \subset B$ is called equivariant if $g_\#(\mathcal{W}) = \mathcal{W}$ for all $g \in G$.

An equivariant varifold \mathcal{W} is called a *free boundary equivariantly stationary varifold* if $\delta_V(\mathcal{W}) = 0$ for all vector fields $V \in Is_{FB}^G$. Note that such V are tangent to ∂B by definition.

As observed in [Ke2], the set of vector fields Is_{FB}^G is a convex vector space. Thus the “pull-tight” procedure (Proposition 3.2 in [DR]) applies to produce a free boundary equivariantly stationary varifold \mathcal{V} . By Lemma 3.8 in [Ke2], a free boundary equivariantly stationary varifold is in fact stationary. Thus we produce an equivariant, free boundary stationary varifold \mathcal{V} with mass equal to W_Π^G .

7.2. Regularity of \mathcal{V} . As in Section 4.2 in [Ke2] and Proposition 4.3 in [DR], one can find a min-max sequence Σ_j that is almost minimizing in annuli small enough. Precisely, for each $x \in \overline{B}$, there is a radius $r(x) > 0$ so that for any annulus An about x of outer radius at most $r(x)$, Σ_j is $1/j$ G -almost minimizing in An . In other words, Σ_j is almost minimizing but only among variations through G -equivariant deformations.

We must construct smooth replacements for \mathcal{V} in such annuli. There are several cases:

- (1) $x \notin \mathcal{S}$
- (2) $x \in \mathcal{S}$ and Σ_t do not contain the arc \mathcal{A} containing x
- (3) $x \in \mathcal{S}$ and Σ_t does contain the arc \mathcal{A} containing x

We can further subdivide case (1) (2) and (3) into subcase a) if x is in the interior of B and b) if x is in the boundary of B .

Case (1ab) and (2a) follow from previous work on free boundary problems (see for instance [GJ]) and [Ke]. Let us first consider (3ab).

Let us first recall the following definitions (Section 4 in [Ke2]):

Definition 7.1. Let Σ be a smooth G -equivariant surface contained in a G -ball. Choose a normal vector field n on Σ . Let us call a smooth function ϕ defined on Σ an *equivariant deformation* if for all t small enough, the following set is G -equivariant:

$$(7.1) \quad \Sigma_{t\phi} = \{\exp_p(n(p)t\phi(p)) \mid p \in \Sigma\}.$$

In other words, ϕ is an equivariant deformation if moving normally to Σ according to ϕ gives rise to G -equivariant surfaces.

Definition 7.2. A G -equivariant surface Σ is G -stable if it is stable among equivariant deformations, i.e.,

$$(7.2) \quad \left. \frac{d^2}{dt^2} \right|_{t=0} \mathcal{H}^2(\Sigma + nt\phi) \geq 0,$$

for all equivariant deformations ϕ .

Given a point p in case (3a), one can minimize among G -equivariant $1/j$ -isotopies in a fixed annulus An based about p to produce a \mathbb{Z}_2 -stable surface V_j which is a replacement for Σ_j in An . Assume for the moment that V_j is smooth.

If V_j intersects \mathcal{S} transversally (as in case (2)), it was proved in Section 4 in [Ke2] that G -stability implies stability. In case (3a) however, V_j contains an arc $\mathcal{A} \subset \mathcal{S}$. In this setting, \mathbb{Z}_2 -stability may not be equivalent to stability. It is easy to see that V_j is stable among odd deformations, but it may not be stable among even deformations. Given any fundamental domain \mathcal{F} of the \mathbb{Z}_2 action, any compactly supported variation χ supported in \mathcal{F} extends by $\chi(\tau(x)) := -\chi(x)$ to give an odd variation of An . Thus V_j is stable in any such \mathcal{F} . By curvature estimates for stable surface [S], this implies that

$$(7.3) \quad |A|^2(x) \leq C \text{dist}(x, (\mathcal{A} \cap An) \cup \partial An)^{-2}.$$

In other words, the curvature of V_j is bounded in the interior of An but may blow up as one approaches the axis \mathcal{A} .

On the other hand, since Σ_j has bounded genus and area, it follows that V_j have bounded genus and area (see for instance Proposition 4.7 in [Ke]). By the integrated Gauss-Bonnet argument of Ilmanen (Lemma 1 in Lecture 3 in [I]), in any proper subannulus $An' \subset An$, there holds

$$(7.4) \quad \sup_j \int_{V_j \cap An'} |A|_{V_j}^2 \leq C(An'),$$

where $C(An')$ is a constant depending on An' which blows up as An' approaches An . Note that one cannot expect any curvature bound up to the boundary of An . Consider for instance the Fraser-Schoen free boundary minimal surfaces F_k in B . They have genus 0 and bounded areas, but do not converge smoothly to $2D$ over ∂B .

Using (7.4), in any such An' , V_j has a convergent subsequence by classical results due to Choi-Schoen [CS]. Taking a sequence of annuli $An' \rightarrow An$ and a diagonal argument, one can produce a subsequence of V_j converging to a smooth \mathbb{Z}_2 -stable minimal surface V_∞ in compact subsets of An . Note that the convergence may not be smooth in any given subannulus, An' . There may be finitely many points where the convergence fails to be smooth. Note also that the convergence $V_j \rightarrow V_\infty$ is smooth in the interior of An away from the axis \mathcal{A} by (7.3).

Thus we have replaced \mathcal{V} in An by a smooth minimal surface V_∞ . Moreover, the replacement is stationary over $\partial(An)$ as in Proposition 7.5 in [CD] and thus gives a replacement for \mathcal{V} .

In Section 6 of [CD], the fact that V has local smooth replacements is used to prove the regularity of \mathcal{V} . One can peruse the proof of Theorem 6.3 in [CD] to see that the curvature estimates of Schoen's are not needed at all. Colding-De Lellis explicitly point this out in Section 2.5: "In fact what we will use is not the actual curvature estimate, rather it is the following consequence of it: a sequence of stable minimal surfaces has a convergent subsequence." We replace this compactness theorem of Schoen [S] with the fact that \mathbb{Z}_2 -stable surfaces with bounded area and genus subconverge smoothly away from finitely many points concentrating along the axis \mathcal{A} . This then implies regularity of \mathcal{V} .

It remains to prove that one can minimize among G -equivariant $1/j$ -isotopies in An to produce the smooth \mathbb{Z}_2 -stable replacements V_j .

As observed in Section 4.3 in [Ke2], by a Squeezing Lemma, to prove regularity of the replacements V_j it is enough to prove the following Proposition 7.3, which states that one can minimize area restricting to equivariant isotopies. The only difference here from Section 4.3 in [Ke2] is that one of the curves in the boundary is not acted upon freely by \mathbb{Z}_2 . Some of the proof is identical to Proposition 4.14 in [Ke2]. The

added difficulty is that here it does not follow in the same way as in [Ke2] that area minimizing disks are equivariant (though I do not know an explicit example where this fails).

Proposition 7.3. *Suppose \mathbb{Z}_2 acts on a 3-ball B as a rotation of 180° about the line \mathcal{S} . Let $\{\gamma_i\}_{i=1}^k$ be a collection of Jordan curves in ∂B bounding a \mathbb{Z}_2 -equivariant surface $\Sigma \subset B$ of genus 0 and so that \mathbb{Z}_2 acts freely on the curves $\{\gamma_i\}_{i=2}^k$ but non-freely on γ_1 (and thus γ_1 contains the two points of $\mathcal{S} \cap \partial B$). Consider a minimizing sequence Σ_i for area among surfaces that are contained in B and \mathbb{Z}_2 -isotopic to Σ . Then after passing to a subsequence (not relabeled) Σ_i converges with multiplicity 1 to a smooth embedded \mathbb{Z}_2 -equivariant minimal surface V with boundary $\{\gamma_i\}_{i=1}^k$ and genus 0.*

Proof. Regularity of V away from the axis \mathcal{S} follows from the replacement theory [Pi] and [MSY]. It remains to prove that V extends smoothly over \mathcal{S} . To that end, we can fix a small ball N centered about \mathcal{S} . As in Proposition 4.14 in [Ke2], Steps 1) and 2), we can perform finitely many neckpinches so that the surgered sequence Σ_i still converges to V and moreover, we can replace Σ_i so that it consists of disks $\cup_k D_k^j$ inside N . Let $D_{k'}^j$ denote the unique disk that in N with containing the singular axis. We then can replace each disk in the collection of disks $(\cup_k D_k^j) \setminus D_{k'}^j$ with the area minimizing disk $\cup_{k \neq k'} A_k^j$ in N with the same boundary in ∂N . By the Meeks-Yau cut-and-paste argument [MY] the disks in the collection $\cup_{k \neq k'} A_k^j$ are pairwise disjoint (since their boundaries are) and are moreover minimizers among all \mathbb{Z}_2 isotopies. The disk $D_{k'}^j$ is problematic since it is not clear that the area-minimizer is \mathbb{Z}_2 -equivariant. If it is, then we can replace $D_{k'}^j$ with the area-minimizer, which again by Meeks-Yau is disjoint from the disks $\cup_{k \neq k'} A_k^j$. In that case, taking $j \rightarrow \infty$ we produce a replacement for the varifold V in N , which implies regularity as in [Pi] [CD].

Suppose instead the area minimizer S for disks with boundary γ_1 in N is not \mathbb{Z}_2 -equivariant. Then consider $\tau(S)$, where τ generates the \mathbb{Z}_2 action. By the Meeks-Yau argument, $\tau(S)$ and S are disjoint except for their boundaries which coincide. Since S and $\tau(S)$ are area minimizing, it follows by Meeks-Yau again that $\cup_{k \neq k'} A_k^j$ is disjoint from $\tau(S)$ and S and thus avoids entirely the 3-ball R in N bounded between $\tau(S)$ and S . By Lemma 7.4 below we can minimize area for $D_{k'}^j$ in the region R among \mathbb{Z}_2 -equivariant isotopies to produce a \mathbb{Z}_2 -equivariant minimal disk $A_{k'}^j$ in R (though potentially not stable among all variations). By Schoen-Simon [SchS] it follows that simply connected minimal surfaces satisfy curvature estimates. Thus $A_{k'}^j$ still

has a convergent subsequence as $j \rightarrow \infty$. The limit in j of this disk together with the limits of $\cup_{k \neq k'} A_k^j$ gives a smooth replacement for V in N , and thus establishes regularity. \square

Lemma 7.4. *Let N be a domain in \mathbb{R}^3 with boundary consisting of two minimal disks D_1, D_2 . Assume N is invariant under a rotation τ of 180° that interchanges D_1 and D_2 . Let \mathcal{S} denote the singular set of the \mathbb{Z}_2 action which intersects $\partial D_1 = \partial D_2$ in two points and is contained in N . Let γ be the \mathbb{Z}_2 -equivariant closed curve given by $\partial D_1 = \partial D_2$ bounding a disk Σ in N that contains \mathcal{S} . Then one can minimize area for Σ restricting to \mathbb{Z}_2 -equivariant isotopies in N to produce a \mathbb{Z}_2 -stable minimal disk Σ_∞ containing \mathcal{S} with boundary γ .*

Proof. It follows as before that the minimizing sequence among \mathbb{Z}_2 isotopies converges to a smooth minimal disk away from \mathcal{S} . Since each half of Σ_∞ contains the line \mathcal{S} , by Schwarz reflection, Σ_∞ is a smooth surface, embedded away from \mathcal{S} , but potentially containing points of self-intersection at \mathcal{S} . Thus we need only show that any tangent cone of Σ_∞ at \mathcal{S} consists of a plane with some multiplicity.

The proof of this fact is essentially contained in Theorem 3 in [AS] and Section 8 in [DP] so we merely sketch the argument. Consider a sequence of dilations $\lambda_j \rightarrow \infty$ and the sequence of rescaled surfaces $\tilde{\Sigma}_j := \lambda_j(\Sigma_j - x)$ approaching a tangent cone C at x . Let us restrict attention to the sequence $\tilde{\Sigma}_j$ contained in the unit ball \mathcal{B}_1 in \mathbb{R}^3 . The cone C in \mathcal{B}_1 consists of several half disks $\cup_k P_k$ meeting along the z -axis. This follows as the surface Σ_∞ has bounded genus and is smooth away from the axis \mathcal{S} and thus a tangent cone at \mathcal{S} cannot acquire any singular points beyond \mathcal{S} itself. By a desingularization procedure (Section 8.3 in [DP]), one can assume $\tilde{\Sigma}_j \cap \mathcal{B}_1$ consists of several disks $\cup_i D_i^j$, and the area of each disk is very close to a minimal area disk with the same boundary values. One of these disks D_1^j is the one containing $\mathcal{S} \cap \mathcal{B}_1$, and by the \mathbb{Z}_2 -equivariance this disk converges to a union of two half-disks, say $P_1 \cup P_2$, where P_2 is the half disk obtained by rotating P_1 by 180° . The other disks $\cup_{i \geq 2} D_i^j$ either have no limit in C , or secondly converge to the flat disk $P_1 \cup P_2$ or thirdly can converge to the union of several of the half-disks P_i that are contained in one of the hemispheres $\mathcal{B} \setminus (P_1 \cup P_2)$. This third possibility cannot occur because such disks would be “folding” along a line and thus be very far in area from the infimal area of disks with their boundary values. See Section 8.5 in [DP] for a demonstration of this, or the proof of Theorem 3 in [AS]. \square

As for cases (2b), and (3b), the changes necessary are minor. Fix $x \in \partial B$ and an annulus An centered about x for case (2b) or (3b). Then one can minimize appropriate $1/j$ -isotopies (allowing the boundary of Σ_j in ∂B to move) as in Section 9 in [DR] to obtain a smooth free boundary minimal surface V_j in An . In case (2b) one obtains a \mathbb{Z}_n -stable free boundary minimal surface V_j in An . The surface produced is in fact stable among all variations for the free boundary problem by Proposition 4.6 in [Ke2]. Thus one can apply the curvature estimates for such surfaces (Theorem 7.3 in [DR]) to obtain a smooth replacement in An . In case (3b) one obtains a \mathbb{Z}_2 -stable free boundary minimal surface. Since \mathbb{Z}_2 -stability may not imply stability in this setting, one can appeal to the arguments of case (3a) to obtain nevertheless a convergent subsequence away from finitely many points in An .

7.3. Completion of proof of Theorem 3.2. Since the regularity of \mathcal{V} has been established, the other claims Theorem 3.2ii, iii, iv and v. follow easily. The proofs of iv. and v. are identical to the arguments in Section 5 of [Ke2]. Let us show iii.

In this case, the min-max sequence Σ_i contains the singular axis \mathcal{A} with \mathbb{Z}_2 isotropy. Let us first show that Γ also contains \mathcal{A} . Fix a point $x \in \mathcal{A}$ and a ball B about x so that $B \cap \mathcal{S} = \mathcal{A}'$ and Σ_i intersects ∂B transversally. For each i , there is a distinguished circle C_i contained in ∂B that contains the two points N and S of intersection of \mathcal{A}' with ∂B . The circle C_i is comprised of two arcs A_i and B_i , each starting at N and ending at S and being interchanged by the \mathbb{Z}_2 action τ . Consider the piece P_i of the surface Σ_i in B bounded by the closed piecewise smooth curve $\mathcal{A}' \cup A_i$. Since this curve is evidently bounded away from zero in the flat topology as $i \rightarrow \infty$, it cannot happen that the area of P_i tends to zero as $i \rightarrow \infty$. Thus the varifold limit of P_i is contained in the min-max minimal surface Γ . Since each P_i contains \mathcal{A} , so does Γ .

Finally let us also show the further statement in iii) that when Σ_t contains \mathcal{A} , the multiplicity n of Γ is an odd integer. Fix again a ball B centered about a point $p \in \mathcal{A}$. Taking replacement for Σ_i in B , one obtains a \mathbb{Z}_2 -stable surface V_j . We have seen that V_j converges to \mathcal{V} smoothly away from finitely many points. Choose a point $p \in B$ centered around the axis \mathcal{A} and a ball B_1 centered around it that avoid these finitely many points. Then in B_1 , V_j consists of m graphs f_1^j, \dots, f_m^j each converging smoothly to \mathcal{V} (as $j \rightarrow \infty$). Precisely one of the graphs f_s^j among the f_1^j, \dots, f_m^j contains the axis \mathcal{A} in B_1 . The graph f_s^j is preserved by the involution τ generating \mathbb{Z}_2 . The other

graphs, being disjoint from the singular axis, must be swapped one with another. It follows that m is odd. Thus iii. is established.

The argument for ii) follows with straightforward modifications of the proof of the Improved Lifting Lemma from the arguments of [Ke], [Ke2] and we omit it. □

REFERENCES

- [Ai] N. Aiex, *Non-compactness of the space of minimal hypersurfaces*, available at arXiv:1601.01049v2.
- [A] W.K. Allard, *On the first variation of a varifold*, Ann. of Math.(2) 95 (1972) no. 3, 417–491.
- [Al] F.J. Almgren. *Existence and regularity almost everywhere of solutions to elliptic variational problems with constraints*, Mem.AMS 165 (1976).
- [AS] F.J. Almgren and L. Simon. *Existence of embedded solutions of Plateau’s problem*. Ann. Scuola Norm. Sup. Pisa 6 (1979) 447-495.
- [Ag] S. Agmon, A. Douglis, and L. Nirenberg. *Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I*, Comm. Pure Appl. Math. 12 (1959), 623–727.
- [AN] L. Ambrozio and I. Nunes. *A gap theorem for free boundary minimal surfaces in the three-ball*, preprint available at arXiv:1608.05689v1.
- [BS] J. Bokowski and E. Sperner. *Zerlegung konvexer Körper durch minimale Trennflächen*. J. Reine Angew. Math. 311-312 (1979), 80-100.
- [B] S. Brendle. *A sharp bound for the area of minimal surfaces in the unit ball*, Geom. Funct. Anal. vol 22 (2012) 621–626.
- [C] R. Courant, *The existence of minimal surfaces of given topological structure under prescribed boundary conditions*, Acta Math. 72 (1940), 51-98.
- [CD] T. H. Colding and C. De Lellis. *The min-max construction of minimal surfaces*, Surveys in Differential Geometry, vol 8 (2003) 75-107.
- [CS] H.I. Choi and R. Schoen. *The space of minimal embeddings of a surface into a 3-manifold with positive Ricci curvature*. Invent. Math., 81, 1985, 357–394.
- [DP] C. De Lellis and F. Pellandini. *Genus bounds for minimal surfaces arising from the min-max construction* Jour. Reine. Angewdte Math. (2010).
- [DR] C. De Lellis and J. Ramic, *Min-max theory for minimal hypersurfaces with boundary*, preprint available at arXiv:1611.00926.
- [D] B. Devyver. *Index of the critical catenoid*, preprint available at 1609.02315v1.pdf.
- [FGM] B. Freidin, M. Gulian and P. McGrath, *Free boundary minimal surfaces in the unit ball with low cohomogeneity*, preprint, available at arXiv:1601.07588.
- [FPZ] A. Folha, F. Pacard, and T. Zolotareva. *Free boundary minimal surfaces in the unit 3-ball*, preprint, available at abs/1502.06812.
- [FL] A. Fraser and M. Li. *Compactness of the space of embedded minimal surfaces with free boundary in three-manifolds with nonnegative Ricci curvature and convex boundary*, J. Differential Geom. vol. 96, no. 2 (2014), 183–200.

- [FS] A. Fraser and R. Schoen. *Sharp eigenvalue bounds and minimal surfaces in the ball*, Invent. math., vol 203, Issue 3, pp 823–890.
- [FS2] A. Fraser and R. Schoen. *Minimal surfaces and eigenvalue problems*, Contemp. Math. 599, Amer. Math. Soc., Providence, RI, 105–121, 2013.
- [FS3] A. Fraser and R. Schoen. *The first Steklov eigenvalue, conformal geometry, and free boundary minimal surfaces*, Adv. Math. 226 (2011), no. 5, 4011–4030.
- [HM] D. Hoffman and W. H. Meeks. *A variational approach to the existence of complete embedded minimal surfaces*, Duke Math. J. , 57(3) (1988) 877–893.
- [GJ] M. Grüter and J. Jost, *On embedded minimal disks in convex bodies*, Ann. Inst. H. Poincaré Anal. No Linaire 3 (1986), no. 5, 345–390
- [GJ2] M. Grüter and J. Jost. *Allard type regularity results for varifolds with free boundaries*, Ann. Scuola Nor Sup. Pisa Cl. Sci. (4) 13 (1986), no. 1, 129–169.
- [I] T. Ilmanen. *Lectures on Mean Curvature Flow and Related Equations* in Conference on Partial Differential Equations and Applications to Geometry, 21 August - 1 September, 1995, ICTP, Trieste.
- [KKM] N. Kapouleas, S. Kleene, and N. Møller, *Mean curvature self-shrinkers of high genus: non-compact examples*, preprint available at abs/1106.5454.
- [Ke] D. Ketover. *Genus bounds for min-max minimal surfaces*, preprint, available at abs/1312.2666.
- [Ke2] D. Ketover. *Equivariant min-max theory*, preprint.
- [KMN] D. Ketover, F.C. Marques and A. Neves. *Catenoid estimate and its geometric applications*, preprint, available at abs/1601.04514.
- [KPS] H. Karcher, U. Pinkall and I. Sterling. *New minimal surfaces in \mathbb{S}^3* , Jour. Diff. Geom. (1988) no. 2 169-185.
- [Li] M. Li, *A general existence theorem for embedded minimal surfaces with free boundary*, Comm. Pure Appl. Math. 68 (2015), no. 2, 286-331.
- [MN] F.C. Marques and A. Neves. *Existence of infinitely many minimal hypersurfaces in positive Ricci curvature*. preprint available at abs/1311.6501.
- [MNS] D. Maximo, I. Nunes and G. Smith. *Free boundary minimal annuli in convex three-manifolds*, J. Differential Geom, to appear.
- [MSY] W. Meeks, L. Simon, and S.T. Yau. *Embedded minimal surfaces, exotic spheres, and manifolds with positive Ricci curvature*. Ann. of Math. 2-116 (1982) 621-659.
- [MY] W. H. Meeks, III and S. T. Yau. *The classical Plateau problem and the topology of three-dimensional manifolds*. Topology , 21(4): (1982) 409–442.
- [Ng] X.H. Nguyen, *Construction of complete embedded self-similar surfaces under mean curvature flow, part I*, ProQuest LLC, Ann Arbor, MI, 2006. Thesis (Ph.D.), The University of Wisconsin - Madison. MR2709389.
- [N] J. C. C. Nitsche, *Stationary partitioning of convex bodies*, Arch. Rational Mech. Anal. 89 (1985), no. 1, 119.
- [LZ] M. Li and X. Zhou. *Min-max theory for free boundary minimal hypersurfaces I - regularity theory*, preprint available at abs/1611.02612.
- [LZ2] M. Li and X. Zhou. *Curvature estimates for stable free boundary minimal hypersurfaces*, preprint available at 1611.02605v2.

- [Pi] J. Pitts. *Existence and regularity of minimal surfaces on Riemannian manifolds*. volume 27 of Mathematical Notes. Princeton University Press, Princeton, N.J., 1981.
- [PR] J. Pitts and J.H. Rubinstein. *Applications of minimax to minimal surfaces and the topology of 3-manifolds*, Miniconference on geometry and partial differential equations, 2, Canberra, (1986), 137–170; Proc. Centre Math. Anal. Austral. Nat. Univ.
- [PR2] J. Pitts and J.H. Rubinstein. *Equivariant minimax and minimal surfaces in geometric three-manifolds*, Bull. Amer. Math. Soc. (N.S.) vol. 19, no 1 (1988), 303-309.
- [RV] A. Ros and E. Vergasta. *Stability for hypersurfaces of constant mean curvature with free boundary*, Geom. Dedicata, 56, (1995), 19–33.
- [R] A. Ros. *The isoperimetric problem*, In Proceedings of the Clay Mathematical Institute MSRI summer school on Minimal Surfaces, Global theory of minimal surfaces, 175209, Clay Math. Proc., 2, Amer. Math. Soc., Providence, RI, 2005.
- [S] R. Schoen. *Estimates for stable minimal surfaces in three-dimensional manifolds*. In Seminar on minimal submanifolds volume 103 of Ann of Math Studies ; p. 111-126, Princeton University Press, Princeton, NJ 1983
- [SchS] R. Schoen and L. Simon. *Regularity of simply connected surfaces with quasiconformal Gauss map*. In Seminar on Minimal submanifolds, Annals of Math Studies, vol. 103, 127–145, Princeton University Press, Princeton, NJ 1983.
- [SS] F. Smith. *On the existence of embedded minimal 2-spheres in the 3-sphere, endowed with an arbitrary riemannian metric*. Phd thesis, Supervisor: Leon Simon, University of Melbourne, 1982.
- [SZ] G. Smith and D. Zhou. *The Morse index of the critical catenoid*, preprint available at 1609.01485v1.pdf.
- [T] H. Tran. *Index characterization for free boundary minimal surfaces*, preprint available at 1609.01651v2.pdf.
- [W2] B. White. *A local regularity theorem for mean curvature flow*, Ann. of Math. (2) 161 (2005), no. 3, 1487–1519.
- [W] B. White. *The space of minimal surfaces for varying Riemannian metrics*, Indiana Math. Journal 40 (1991), no.1, 161–200.
- [V] A. Volkman. *A monotonicity formula for free boundary surfaces with respect to the unit ball*, preprint available at abs/1402.4784.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08544

E-mail address: `dketover@math.princeton.edu`