

Exploring the $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ and $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ hidden-bottom hadronic transitions

Qi Huang,^{*} Bo Wang,[†] and Xiang Liu[‡]

*School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, China
Research Center for Hadron and CSR Physics, Lanzhou University and Institute of Modern Physics of CAS, Lanzhou 730000, China*

Dian-Yong Chen[§]

Department of Physics, Southeast University, Nanjing 210094, China

Takayuki Matsuki[¶]

*Tokyo Kasei University, 1-18-1 Kaga, Itabashi, Tokyo 173-8602, Japan
Theoretical Research Division, Nishina Center, RIKEN, Wako, Saitama 351-0198, Japan*

In this work, we investigate the hadronic loop contributions to the $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ ($J = 0, 1, 2$) along with $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ ($J = 0, 1, 2$) transitions. We predict that the branching ratios of $\Upsilon(6S) \rightarrow \chi_{b0}\phi$, $\Upsilon(6S) \rightarrow \chi_{b1}\phi$ and $\Upsilon(6S) \rightarrow \chi_{b2}\phi$ are $(0.68 \sim 4.62) \times 10^{-6}$, $(0.50 \sim 3.43) \times 10^{-6}$ and $(2.22 \sim 15.18) \times 10^{-6}$, respectively and those of $\Upsilon(6S) \rightarrow \chi_{b0}\omega$, $\Upsilon(6S) \rightarrow \chi_{b1}\omega$ and $\Upsilon(6S) \rightarrow \chi_{b2}\omega$ are $(0.15 \sim 2.81) \times 10^{-3}$, $(0.63 \sim 11.68) \times 10^{-3}$ and $(1.08 \sim 20.02) \times 10^{-3}$, respectively. Especially, some typical ratios, which reflect the relative magnitudes of the predicted branching ratios, are given, i.e., for $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ transitions, $\mathcal{R}_{10}^\phi = \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\phi]/\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\phi] \approx 0.74$, $\mathcal{R}_{20}^\phi = \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\phi]/\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\phi] \approx 3.28$, and $\mathcal{R}_{21}^\phi = \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\phi]/\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\phi] \approx 4.43$, and for $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ transitions, $\mathcal{R}_{10}^\omega = \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\omega]/\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\omega] \approx 4.11$, $\mathcal{R}_{20}^\omega = \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\omega]/\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\omega] \approx 7.06$, and $\mathcal{R}_{21}^\omega = \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\omega]/\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\omega] \approx 1.72$. With the running of BelleII in the near future, experimental measurement of these two kinds of transitions will be a potential research issue.

PACS numbers: 14.40.Pq, 13.25.Gv

I. INTRODUCTION

As an interesting research issue, experimental studies of the hadronic transitions of $\Upsilon(5S)$ have been focused on by the Belle Collaboration in the past decade. When surveying the reported hadronic transitions of $\Upsilon(5S)$, we found their general property, i.e., their observed hadronic transitions have large branching ratios. For example, Belle observed anomalous decay widths of the $\Upsilon(5S) \rightarrow \Upsilon(nS)\pi^+\pi^-$ [1], and $\Upsilon(5S) \rightarrow \chi_{bJ}\omega$ ($J = 0, 1, 2$) transitions [2]. In addition, two bottomonium-like states $Z_b(10610)$ and $Z_b(10650)$ were observed in $\Upsilon(5S) \rightarrow \Upsilon(nS)\pi^+\pi^-$ [3]. As indicated in a serial of theoretical studies [4–10], the puzzling phenomena happening on $\Upsilon(5S)$ transitions reflect an underlying mechanism mediated by a coupled channel effect since $\Upsilon(5S)$ is above the thresholds of $B_{(s)}^{(*)}\bar{B}_{(s)}^{(*)}$ [11].

In the bottomonium family, the $\Upsilon(6S)$ has the similar situation to that of $\Upsilon(5S)$. We have a reason to believe that the coupled channel effect is still important to the hadronic transitions of $\Upsilon(6S)$, whose exploration is, thus, an intriguing topic. This theme can provide us a valuable information of the coupled-channel effect on these decays.

In this work, we calculate the $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ ($J = 0, 1, 2$) along with $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ ($J = 0, 1, 2$) processes via the

hadronic loop mechanism, which is an equivalent description of the coupled channel effect [4–6, 8–10, 12–16]. By analyzing these transitions, the relative decay rates of $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ ($J = 0, 1, 2$) and $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ ($J = 0, 1, 2$), which are a typical physical quantity given by our calculation, are determined. Especially, our results show that these relative decay rates are weakly dependent on the model parameters. Thus, experimental measurement of these rates can be a crucial test of the hadronic loop mechanism in the $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ and $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ decays. In addition, we also estimate the typical values of the branching ratios of $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ and $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$, which can be measured experimentally in near future. Anyway, we would like to inspire experimenters' interest in searching for the $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ and $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ decays by our results presented in this work.

This paper is organized as follows. After introduction, we present the detailed calculation of $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ and $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ via the hadronic loop mechanism in Sec. II. The numerical results are presented in Sec. III. The paper ends with a short summary.

II. $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ AND $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ TRANSITIONS VIA HADRONIC LOOP MECHANISM

Under the hadronic loop mechanism, the $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ transitions occur via the triangle loops composed of $B_s^{(*)0}$ and $\bar{B}_s^{(*)0}$, which play a role of the bridge to connect the initial state $\Upsilon(6S)$ and final states ϕ and χ_{bJ} . In Figs. 1–3, we list the typical diagrams depicting the $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ ($J = 0, 1, 2$) transitions. For the $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ transitions, due to very different quark contents between ϕ and ω , the bridges change

^{*}Electronic address: huangqi16@lzu.edu.cn

[†]Electronic address: wangb13@lzu.edu.cn

[‡]Electronic address: xiangliu@lzu.edu.cn

[§]Electronic address: chendy@seu.edu.cn

[¶]Electronic address: matsuki@tokyo-kasei.ac.jp

to $B^{(*)}$ and $\bar{B}^{(*)}$ and the diagrams change simultaneously as in Figs. 4-6.

To calculate these diagrams at the hadron level, we adopt the effective Lagrangian approach, in which we first introduce the Lagrangians relevant to our calculation.

For the interactions between a heavy quarkonium and two heavy-light mesons, the Lagrangians are constructed based on the heavy quark effective theory. In the heavy quark limit, the light degrees of freedom s_ℓ is a good quantum number. Thus, each value of s_ℓ is assigned to a doublet formed by the states with a total angular momentum $J = s_\ell \pm 1/2$, while for the heavy quarkonium, since the degeneracy is expected under the rotations of two heavy quark spins, there is a multiplet formed by heavy quarkonia with the same angular momentum ℓ .




FIG. 1: Schematic diagrams depicting the $\Upsilon(6S) \rightarrow \chi_{b0}\phi$ process via the hadronic loop mechanism.




FIG. 2: Schematic diagrams depicting the $\Upsilon(6S) \rightarrow \chi_{b1}\phi$ process via the hadronic loop mechanism.




FIG. 3: Schematic diagrams depicting the $\Upsilon(6S) \rightarrow \chi_{b2}\phi$ process via the hadronic loop mechanism.




FIG. 4: Schematic diagrams depicting the $\Upsilon(6S) \rightarrow \chi_{b0}\omega$ process via the hadronic loop mechanism.




FIG. 5: Schematic diagrams depicting the $\Upsilon(6S) \rightarrow \chi_{b1}\omega$ process via the hadronic loop mechanism.




FIG. 6: Schematic diagrams depicting the $\Upsilon(6S) \rightarrow \chi_{b2}\omega$ process via the hadronic loop mechanism.

Therefore, under the framework of heavy quark symmetry, general forms of couplings between an S-wave or P-wave heavy quarkonium and two heavy-light mesons can be constructed as [18]

$$\begin{aligned}\mathcal{L}_s &= ig\text{Tr}\left[R^{(Q\bar{Q})}\bar{H}^{(Q\bar{Q})}\gamma^\mu\overset{\leftrightarrow}{\partial}_\mu\bar{H}^{(Q\bar{Q})}\right] + H.c., \\ \mathcal{L}_p &= ig_1\text{Tr}\left[P^{(Q\bar{Q})\mu}\bar{H}^{(Q\bar{Q})}\gamma_\mu\bar{H}^{(Q\bar{Q})}\right] + H.c.,\end{aligned}\quad (1)$$

in which $R^{(Q\bar{Q})}$ and $P^{(Q\bar{Q})}$ denote multiplets formed by bottomonia with $\ell = 0$ and $\ell = 1$, and their detailed expressions, as in Ref. [21], can be written as

$$R^{(Q\bar{Q})} = \frac{1+\not{v}}{2}\left[\gamma^\mu\gamma_\mu - \eta_b\gamma_5\right]\frac{1-\not{v}}{2},\quad (2)$$

$$\begin{aligned}P^{(Q\bar{Q})\mu} &= \frac{1+\not{v}}{2}\left[\chi_{b2}^{\mu\alpha}\gamma_\alpha + \frac{1}{\sqrt{2}}\epsilon^{\mu\alpha\beta\gamma}v_\alpha\gamma_\beta\chi_{b1\gamma}\right. \\ &\quad \left.+ \frac{1}{\sqrt{3}}(\gamma^\mu - v^\mu)\chi_{b0} + h_b^\mu\gamma_5\right]\frac{1-\not{v}}{2},\end{aligned}\quad (3)$$

respectively. $H^{(Q\bar{Q})}$ represents a doublet formed by heavy-light pseudoscalar and vector mesons [18–21]

$$H^{(Q\bar{Q})} = \frac{1+\not{v}}{2}\left[\mathcal{B}_\mu^*\gamma^\mu - \mathcal{B}\gamma^5\right],\quad (4)$$

with definitions $\mathcal{B}^{(*)\dagger} = (B^{(*)+}, B^{(*)0}, B_s^{(*)0})$ and $\mathcal{B}^{(*)} = (B^{(*)-}, \bar{B}^{(*)0}, \bar{B}_s^{(*)0})^T$ as in Ref. [13]. $H^{(Q\bar{Q})}$ corresponds to a doublet formed by heavy-light anti-mesons, which can be obtained by applying the charge conjugation operation to $H^{(Q\bar{Q})}$.

For the interaction between a light vector meson and two heavy-light mesons, the general form of the Lagrangian reads as [18, 22–26]

$$\mathcal{L}_V = i\beta\text{Tr}[H^j v^\mu(-\rho_\mu)_j \bar{H}_i] + i\lambda\text{Tr}[H^j \sigma^{\mu\nu} F_{\mu\nu}(\rho) \bar{H}_i],\quad (5)$$

where

$$\rho_\mu = i\frac{g_V}{\sqrt{2}}\mathcal{V}_\mu,\quad (6)$$

$$F_{\mu\nu}(\rho) = \partial_\mu\rho_\nu - \partial_\nu\rho_\mu + [\rho_\mu, \rho_\nu],\quad (7)$$

and a vector octet \mathcal{V} has the form

$$\mathcal{V} = \begin{pmatrix} \frac{1}{\sqrt{2}}(\rho^0 + \omega) & \rho^+ & K^{*+} \\ \rho^- & \frac{1}{\sqrt{2}}(-\rho^0 + \omega) & K^{*0} \\ K^{*-} & \bar{K}^{*0} & \phi \end{pmatrix}.\quad (8)$$

By expanding the Lagrangians in Eqs. (1) and (5), the following concrete expressions are obtained

$$\begin{aligned}\mathcal{L}_{\Upsilon\mathcal{B}^{(*)}\mathcal{B}^{(*)}} &= -ig_{\Upsilon\mathcal{B}\mathcal{B}}\Upsilon_\mu(\partial^\mu\mathcal{B}\mathcal{B}^\dagger - \mathcal{B}\partial^\mu\mathcal{B}^\dagger) \\ &\quad + g_{\Upsilon\mathcal{B}^*\mathcal{B}}\epsilon^{\mu\nu\alpha\beta}\partial_\mu\Upsilon_\nu(\mathcal{B}_\alpha^*\overset{\leftrightarrow}{\partial}_\beta\mathcal{B}^\dagger - \mathcal{B}\overset{\leftrightarrow}{\partial}_\beta\mathcal{B}_\alpha^{*\dagger}) \\ &\quad + ig_{\Upsilon\mathcal{B}^*\mathcal{B}^*}\Upsilon^\mu(\mathcal{B}_\nu^*\partial^\nu\mathcal{B}_\mu^{*\dagger} - \partial^\nu\mathcal{B}_\mu^*\mathcal{B}_\nu^{*\dagger} - \mathcal{B}_\nu^*\overset{\leftrightarrow}{\partial}_\mu\mathcal{B}^{*\nu\dagger}),\end{aligned}\quad (9)$$

$$\begin{aligned}\mathcal{L}_{\chi_{bJ}\mathcal{B}^{(*)}\mathcal{B}^{(*)}} &= -g_{\chi_{b0}\mathcal{B}\mathcal{B}}\chi_{b0}\mathcal{B}\mathcal{B}^\dagger - g_{\chi_{b0}\mathcal{B}^*\mathcal{B}^*}\chi_{b0}\mathcal{B}_\mu^*\mathcal{B}^{*\mu\dagger} \\ &\quad + ig_{\chi_{c1}\mathcal{B}\mathcal{B}}\chi_{b1}^\mu(\mathcal{B}_\mu^*\mathcal{B}^\dagger - \mathcal{B}\mathcal{B}_\mu^{*\dagger}) \\ &\quad - g_{\chi_{b2}\mathcal{B}\mathcal{B}}\chi_{b2}^{\mu\nu}\partial_\mu\mathcal{B}\partial_\nu\mathcal{B}^\dagger \\ &\quad + g_{\chi_{b2}\mathcal{B}^*\mathcal{B}^*}\chi_{b2}^{\mu\nu}(\mathcal{B}_\mu^*\mathcal{B}_\nu^{*\dagger} + \mathcal{B}_\nu^*\mathcal{B}_\mu^{*\dagger}) \\ &\quad - ig_{\chi_{b2}\mathcal{B}^*\mathcal{B}}\epsilon_{\mu\nu\alpha\beta}\partial^\alpha\chi_{b2}^{\mu\rho}(\partial_\rho\mathcal{B}^{*\nu}\partial^\beta\mathcal{B}^\dagger - \partial^\beta\mathcal{B}\partial_\rho\mathcal{B}^{*\nu\dagger}),\end{aligned}\quad (10)$$

$$\begin{aligned}\mathcal{L}_{\mathcal{B}^{(*)}\mathcal{B}^{(*)}\mathcal{V}} &= -ig_{\mathcal{B}\mathcal{B}\mathcal{V}}\mathcal{B}_i^\dagger\overset{\leftrightarrow}{\partial}^\mu\mathcal{B}^j(\mathcal{V}_\mu)_j^i \\ &\quad - 2f_{\mathcal{B}^*\mathcal{B}\mathcal{V}}\epsilon_{\mu\nu\alpha\beta}(\partial^\mu\mathcal{V}^\nu)_j^i(\mathcal{B}_i^\dagger\overset{\leftrightarrow}{\partial}^\alpha\mathcal{B}^{*\beta j} - \mathcal{B}_i^{*\beta\dagger}\overset{\leftrightarrow}{\partial}^\alpha\mathcal{B}^j) \\ &\quad + ig_{\mathcal{B}^*\mathcal{B}^*\mathcal{V}}\mathcal{B}_i^{*\nu\dagger}\overset{\leftrightarrow}{\partial}^\mu\mathcal{B}_\nu^{*j}(\mathcal{V}_\mu)_j^i \\ &\quad + 4if_{\mathcal{B}^*\mathcal{B}^*\mathcal{V}}\mathcal{B}_{i\mu}^{*\dagger}(\partial^\mu\mathcal{V}^\nu - \partial^\nu\mathcal{V}^\mu)_j^i\mathcal{B}_\nu^{*j}.\end{aligned}\quad (11)$$

With the above effective Lagrangians, we can write out the amplitudes of hadronic loop contributions to $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ ($J = 0, 1, 2$). For the $\Upsilon(6S) \rightarrow \chi_{b0}\phi$ transition, the amplitudes corresponding to Fig. 1 are

$$\begin{aligned}\mathcal{M}_{(0-1)} &= \int \frac{d^4q}{(2\pi)^4}[-ig_{\Upsilon B_s B_s}\epsilon_\mu^\mu((ik_1)_\mu - (ik_2)_\mu)] \\ &\quad \times [-ig_{B_s B_s \phi}\epsilon_{\phi\lambda}^\lambda((-ik_1)^\lambda - (iq)^\lambda)][-g_{B_s B_s \chi_{b0}}] \\ &\quad \times \frac{1}{k_1^2 - m_{B_s}^2} \frac{1}{k_2^2 - m_{B_s}^2} \frac{1}{q^2 - m_{B_s}^2} \mathcal{F}^2(q^2),\end{aligned}\quad (12)$$

$$\begin{aligned} \mathcal{M}_{(0-2)} = & \int \frac{d^4 q}{(2\pi)^4} [-g_{\Upsilon B_s^* B_s^*} \epsilon^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu} (-ip_1)_\nu ((ik_2)_\beta - (ik_1)_\beta)] \\ & \times [2f_{B_s^* B_s^* \phi} \epsilon_{\lambda\rho\delta\sigma} \epsilon_\phi^{*\lambda} (ip_2)^\rho ((-ik_1)^\delta - (iq)^\delta)] [-g_{B_s B_s \chi_{b0}}] \\ & \times \frac{-g_\alpha^\sigma + k_{1\alpha} k_1^\sigma / m_{B_s^*}^2}{k_1^2 - m_{B_s^*}^2} \frac{1}{k_2^2 - m_{B_s^*}^2} \frac{1}{q^2 - m_{B_s^*}^2} \mathcal{F}^2(q^2), \end{aligned} \quad (13)$$

$$\begin{aligned} \mathcal{M}_{(0-3)} = & \int \frac{d^4 q}{(2\pi)^4} [g_{\Upsilon B_s^* B_s^*} \epsilon^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu} (-ip_1)_\nu ((ik_2)_\beta - (ik_1)_\beta)] \\ & \times [-2f_{B_s^* B_s^* \phi} \epsilon_{\lambda\rho\delta\sigma} \epsilon_\phi^{*\lambda} (ip_2)^\rho ((-ik_1)^\delta - (iq)^\delta)] \\ & \times [-g_{B_s^* B_s^* \chi_{b0}}] \frac{1}{k_1^2 - m_{B_s^*}^2} \frac{-g_\alpha^\zeta + k_{2\alpha} k_2^\zeta / m_{B_s^*}^2}{k_2^2 - m_{B_s^*}^2} \\ & \times \frac{-g_\zeta^\sigma + q^\sigma q_\zeta / m_{B_s^*}^2}{q^2 - m_{B_s^*}^2} \mathcal{F}^2(q^2), \end{aligned} \quad (14)$$

$$\begin{aligned} \mathcal{M}_{(0-4)} = & \int \frac{d^4 q}{(2\pi)^4} [ig_{\Upsilon B_s^* B_s^*} \epsilon_\Upsilon^\mu (g_{\nu\alpha} g_{\mu\beta} (ik_2)^\nu - g_{\mu\alpha} g_{\nu\beta} (ik_1)^\nu \\ & - g_{\alpha\beta} ((ik_2)_\mu - (ik_1)_\mu))] \\ & \times [\epsilon_{\phi\lambda}^* (ig_{B_s^* B_s^* \phi} g^{\delta\sigma} ((-ik_1)^\lambda - (iq)^\lambda) \\ & + 4if_{B_s^* B_s^* \phi} (ip_2)_\rho (g^{\lambda\delta} g^{\rho\sigma} - g^{\lambda\sigma} g^{\rho\delta}))] [-g_{B_s^* B_s^* \chi_{b0}}] \\ & \times \frac{-g_\delta^\alpha + k_{1\delta} k_1^\alpha / m_{B_s^*}^2}{k_1^2 - m_{B_s^*}^2} \frac{-g^{\beta\zeta} + k_{2\beta} k_2^\zeta / m_{B_s^*}^2}{k_2^2 - m_{B_s^*}^2} \\ & \times \frac{-g_{\sigma\zeta} + q_\sigma q_\zeta / m_{B_s^*}^2}{q^2 - m_{B_s^*}^2} \mathcal{F}^2(q^2), \end{aligned} \quad (15)$$

where p_1 , p_2 and p_3 are momenta of $\Upsilon(6S)$, ϕ/ω and χ_{bJ} , and k_1 , k_2 , and q are momenta of internal $B_{(s)}^{(*)}$ and exchanged $B_{(s)}^{(*)}$, respectively. In these expressions of the decay amplitudes, the monopole form factor is introduced, by which the inner structure of interaction vertices is reflected and the off-shell effect of the exchanged bottom-strange mesons is compensated. Here, the adopted form factor is taken as $\mathcal{F}(q^2) = (m_E^2 - \Lambda^2)/(q^2 - \Lambda^2)$, with m_E being the mass of the exchanged boson and the cutoff Λ being parameterized as $\Lambda = m_E + \alpha_\Lambda \Lambda_{QCD}$ with $\Lambda_{QCD} = 0.22$ GeV as in Refs. [14–16]. We need to specify that the monopole behavior of the adopted form factor was suggested by the QCD sum rule studied in Ref. [17]. In a serial of published papers (see Refs. [4–6, 8–10, 12–16, 21, 29]), the monopole form factor was adopted to study the transitions of charmonia and bottomonia, and B decays. Thus, this approach has been tested by these successful studies.

In a similar way, we can further write out the decay amplitudes of $\Upsilon(6S) \rightarrow \chi_{b1}\phi$ and $\Upsilon(6S) \rightarrow \chi_{b2}\phi$, which are collected in Appendix. By considering the isospin symmetry, a general expression of the total amplitude of $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ with $J = 0, 1, 2$ is written as

$$\mathcal{M}_{\Upsilon(6S) \rightarrow \chi_{bJ}\phi}^{\text{Total}} = 2 \sum_j \mathcal{M}_{(J-j)}. \quad (16)$$

Then, the partial decay width reads

$$\Gamma_{\Upsilon(6S) \rightarrow \chi_{bJ}\phi} = \frac{1}{3} \frac{1}{8\pi} \frac{|\vec{p}_\phi|}{m_{\Upsilon(6S)}^2} |\overline{\mathcal{M}_{\Upsilon(6S) \rightarrow \chi_{bJ}\phi}^{\text{Total}}}|^2, \quad (17)$$

where the overline indicates the sum over polarizations of $\Upsilon(6S)$, ϕ , and χ_{b1} (or χ_{b2}) and the factor $\frac{1}{3}$ denotes the average over the polarization of initial $\Upsilon(6S)$.

In the case of $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$, the expression of the partial decay width is given by

$$\Gamma_{\Upsilon(6S) \rightarrow \chi_{bJ}\omega} = \frac{1}{3} \frac{1}{8\pi} \frac{|\vec{p}_\omega|}{m_{\Upsilon(6S)}^2} |\overline{\mathcal{A}_{\Upsilon(6S) \rightarrow \chi_{bJ}\omega}^{\text{Total}}}|^2, \quad (18)$$

with a general expression of the total amplitude of $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ as

$$\mathcal{A}_{\Upsilon(6S) \rightarrow \chi_{bJ}\omega}^{\text{Total}} = 4 \sum_j \mathcal{A}_{(J-j)} \quad (19)$$

by considering the isospin and charge symmetry. The detailed expressions of $\mathcal{A}_{(J-j)}$ are collected in Appendix.

III. NUMERICAL RESULTS

With the formulas listed in Sec. II and Appendix, we estimate the hadronic loop contributions to the $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ together with $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ ($J = 0, 1, 2$) transitions. Besides the masses taken from the Particle Data Book [11], all the other input parameters we need are the coupling constants. Since the $\Upsilon(6S)$ is above the threshold of $B_{(s)}^{(*)}\bar{B}_{(s)}^{(*)}$, the coupling constants between $\Upsilon(6S)$ and $B_{(s)}^{(*)}\bar{B}_{(s)}^{(*)}$ can be evaluated by the partial decay widths of $\Upsilon(6S) \rightarrow B_{(s)}^{(*)}\bar{B}_{(s)}^{(*)}$. In Table I we list the relevant partial decay widths given in Ref. [27] as well as the corresponding extracted coupling constants.

TABLE I: The coupling constants of $\Upsilon(6S)$ interacting with $B_{(s)}^{(*)}\bar{B}_{(s)}^{(*)}$. Here, we also list the corresponding partial decay widths provided in Ref. [27].

Final state	Decay width (MeV)	Coupling constant
$B\bar{B}$	1.32	0.654
$B\bar{B}^*$	7.59	0.077 GeV^{-1}
$B^*\bar{B}^*$	5.89	0.611
$B_s\bar{B}_s$	1.31×10^{-3}	0.043
$B_s\bar{B}_s^*$	0.136	0.023 GeV^{-1}
$B_s^*\bar{B}_s^*$	0.310	0.354

The coupling constants relevant to the interactions between χ_{bJ} and $B_{(s)}^{(*)}\bar{B}_{(s)}^{(*)}$ in the heavy quark limit are related to one gauge coupling g_1 given in Eq. (1), i.e.,

$$g_{\chi_{b0}B\bar{B}} = 2\sqrt{3}g_1\sqrt{m_{\chi_{b0}}m_B}, \quad g_{\chi_{b0}B^*\bar{B}^*} = \frac{2}{\sqrt{3}}g_1\sqrt{m_{\chi_{b0}}m_{B^*}},$$

$$g_{\chi_{b1}B\bar{B}^*} = 2\sqrt{2}g_1\sqrt{m_{\chi_{b1}}m_Bm_{B^*}}, \quad g_{\chi_{b2}B\bar{B}} = 2g_1\frac{\sqrt{m_{\chi_{b0}}}}{m_B},$$

$$g_{\chi_{b2}B\bar{B}^*} = g_1\sqrt{\frac{m_{\chi_{b2}}}{m_{B^*}^3m_B}}, \quad g_{\chi_{b2}B^*\bar{B}^*} = 4g_1\sqrt{m_{\chi_{b2}}m_{B^*}},$$

where $g_1 = -\sqrt{\frac{m_{\chi_{b0}}}{3}} \frac{1}{f_{\chi_{b0}}}$ [21] and $f_{\chi_{b0}} = 175 \pm 55$ MeV is the decay constant of χ_{b0} [28].

Similarly, the coupling constants between ϕ or ω and $B_{(s)}^{(*)} \bar{B}_{(s)}^{(*)}$ can be extracted from Eq. (5),

$$\begin{aligned} g_{B_s B_s \phi} &= g_{B_s^* B_s^* \phi} = \frac{\beta g_V}{\sqrt{2}}, \\ f_{B_s B_s \phi} &= \frac{f_{B_s^* B_s^* \phi}}{m_{B_s^*}} = \frac{\lambda g_V}{\sqrt{2}}, \\ g_{BB\omega} &= g_{B^* B^* \omega} = \frac{\beta g_V}{2}, \\ f_{BB^* \omega} &= \frac{f_{B^* B^* \omega}}{m_{B^*}} = \frac{\lambda g_V}{2}, \end{aligned}$$

with $\beta = 0.9$ and $\lambda = 0.56$ GeV⁻¹. Additionally, we have $g_V = m_\rho/f_\pi$ along with the pion decay constant $f_\pi = 132$ MeV [22–25].

With the above preparation, we can evaluate the branching ratios of the $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ and $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ transitions. However, in our model, there still exists a free parameter α_Λ , which is introduced to parameterize the cutoff Λ . Since the cutoff Λ should not be too far from the physical mass of the exchanged mesons [29], in this work we set the range $0.65 \leq \alpha_\Lambda \leq 1.15$ for $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ transitions and set $0.45 \leq \alpha_\Lambda \leq 1.15$ for $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ transitions to present the numerical results.

In Figs. 7 and 8, we illustrate the α_Λ dependence of the branching ratios of $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ and $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$, respectively, and in Figs. 9 and 10 we present the α_Λ dependence of the relative magnitudes among the branching widths of $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ and $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$, respectively.




FIG. 7: (Color online). The α_Λ dependence of the branching ratios of $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$.




FIG. 8: (Color online). The α_Λ dependence of the branching ratios of $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$.




FIG. 9: (Color online). The α_Λ dependence of the ratios $\mathcal{R}_{10}^\phi = \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\phi]/\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\phi]$, $\mathcal{R}_{20}^\phi = \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\phi]/\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\phi]$ and $\mathcal{R}_{21}^\phi = \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\phi]/\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\phi]$.




FIG. 10: (Color online). The α_Λ dependence of the ratios $\mathcal{R}_{10}^\omega = \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\omega]/\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\omega]$, $\mathcal{R}_{20}^\omega = \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\omega]/\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\omega]$ and $\mathcal{R}_{21}^\omega = \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\omega]/\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\omega]$.

Varying α_Λ between 0.65 and 1.15 in $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$, we have from Fig. 7,

$$\begin{aligned}\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\phi] &= (0.68 \sim 4.62) \times 10^{-6}, \\ \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\phi] &= (0.50 \sim 3.43) \times 10^{-6}, \\ \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\phi] &= (2.22 \sim 15.18) \times 10^{-6},\end{aligned}$$

and for α_Λ varying from 0.45 to 1.15 in $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$, we have from Fig. 8

$$\begin{aligned}\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\omega] &= (0.15 \sim 2.81) \times 10^{-3}, \\ \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\omega] &= (0.63 \sim 11.68) \times 10^{-3}, \\ \mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\omega] &= (1.08 \sim 20.02) \times 10^{-3}.\end{aligned}$$

In addition, some typical values for the relative magnitudes of the predicted branching ratios are obtained from Figs. 9 and 10, which are weakly dependent on the free parameter α_Λ , i.e.,

$$\begin{aligned}\mathcal{R}_{10}^\phi &= \frac{\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\phi]}{\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\phi]} \approx 0.74, \\ \mathcal{R}_{20}^\phi &= \frac{\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\phi]}{\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\phi]} \approx 3.28, \\ \mathcal{R}_{21}^\phi &= \frac{\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\phi]}{\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\phi]} \approx 4.43, \\ \mathcal{R}_{10}^\omega &= \frac{\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\omega]}{\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\omega]} \approx 4.11, \\ \mathcal{R}_{20}^\omega &= \frac{\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\omega]}{\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b0}\omega]} \approx 7.06, \\ \mathcal{R}_{21}^\omega &= \frac{\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b2}\omega]}{\mathcal{B}[\Upsilon(6S) \rightarrow \chi_{b1}\omega]} \approx 1.72.\end{aligned}$$

As shown in numerical results on the $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ decays, the partial decay widths of $\Upsilon(6S) \rightarrow \chi_{b0}\phi$ and $\Upsilon(6S) \rightarrow \chi_{b1}\phi$ are the same order of magnitude, while the partial decay width of $\Upsilon(6S) \rightarrow \chi_{b2}\phi$ is one order of magnitude larger than those of $\Upsilon(6S) \rightarrow \chi_{b0}\phi$ and $\Upsilon(6S) \rightarrow \chi_{b1}\phi$. On the other hand for the $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ decays, the partial decay widths of $\Upsilon(6S) \rightarrow \chi_{b1}\omega$ and $\Upsilon(6S) \rightarrow \chi_{b2}\omega$ are nearly the same order of magnitude, while the partial decay width of $\Upsilon(6S) \rightarrow \chi_{b0}\phi$ is one order of magnitude smaller than those of $\Upsilon(6S) \rightarrow \chi_{b1}\omega$ and $\Upsilon(6S) \rightarrow \chi_{b2}\omega$.

IV. SUMMARY

In the past years, the anomalous hadronic transitions like $\Upsilon(5S) \rightarrow \Upsilon(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) [1] and $\Upsilon(5S) \rightarrow \chi_{bJ}\omega$ ($J = 0, 1, 2$) [2] were reported by Belle, which has stimulated theorists' interest in revealing the underlying mechanism behind these phenomena [4–10]. As a popular and accepted opinion, the hadronic loop mechanism has been applied to explain why there exist anomalous transitions for $\Upsilon(5S)$ [4–10]. In addition, more predictions relevant to the $\Upsilon(5S)$ transition were given in Refs. [12, 13].

The main reason to introduce the hadronic loop mechanism is that $\Upsilon(5S)$ is the second observed bottomonium above the

$B\bar{B}$ threshold, where the coupled-channel effect may become important, which was tested by the studies in Refs. [4–10, 12, 13]. It is obvious that this is not the end of the whole story.

If the hadronic loop mechanism is a universal mechanism existing in higher bottomonium transitions, we have a reason to believe that this mechanism also plays an important role in higher bottomonium transitions. Considering the similarity between $\Upsilon(6S)$ and $\Upsilon(5S)$, where $\Upsilon(6S)$ is the third bottomonium with open-bottom channels, we have focused on $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ and $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ hadronic decays. Using the hadronic loop mechanism, we have estimated the branching ratios of $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ and $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$, which can reach up to 10^{-6} and 10^{-3} , respectively. In the near future, BelleII will be running near the energy range of $\Upsilon(6S)$, which makes BelleII have a great opportunity to find the $\chi_{bJ}\phi$ and $\chi_{bJ}\omega$ decay modes of $\Upsilon(6S)$. If these rare decays are observed, the hadronic loop effects can be further tested.

In this work, we have especially obtained six almost stable ratios \mathcal{R}_{10}^ϕ , \mathcal{R}_{20}^ϕ and \mathcal{R}_{21}^ϕ in addition to \mathcal{R}_{10}^ω , \mathcal{R}_{20}^ω and \mathcal{R}_{21}^ω reflecting the relative magnitudes of the $\Upsilon(6S) \rightarrow \chi_{bJ}\phi$ and $\Upsilon(6S) \rightarrow \chi_{bJ}\omega$ decays, which are weakly dependent on our model parameter α_Λ . Thus, these obtained ratios are important observable quantities. We have also suggested their experimental measurement, which is also a crucial test of our model.

We notice the recent talk of the status of SuperKEKB and the future plan of taking data at the BelleII experiment [30]. Since the collision data on $\Upsilon(6S)$ will be taken, we need to explore the possible interesting research issues about $\Upsilon(6S)$. Our present work is only one step toward the long march.

Acknowledgments

This project is supported by the National Natural Science Foundation of China under Grants No. 11222547, No. 11175073, No. 11375240, and No. 11035006, and by Chinese Academy of Sciences under the funding Y104160YQ0 and the agreement No. 2015-BH-02. XL is also supported by the National Program for Support of Young Top-notch Professionals.

Appendix

As for the $\Upsilon(6S) \rightarrow \chi_{b1}\phi$ transition, the amplitudes corresponding to Fig. 2 are

$$\begin{aligned}\mathcal{M}_{(1-1)} &= \int \frac{d^4q}{(2\pi)^4} [-ig_{\Upsilon B_s B_s} \epsilon_\mu^\mu ((ik_1)_\mu - (ik_2)_\mu)] \\ &\quad \times [-2f_{B_s B_s^* \phi} \epsilon_{\lambda\rho\delta\sigma}^* \epsilon_\phi^{*\lambda} (ip_2)^\rho ((-ik_1)^\delta - (iq)^\delta)] \\ &\quad \times [ig_{B_s B_s^* \chi_{b1}} \epsilon_{\chi_{b1}}^{*\zeta}] \frac{1}{k_1^2 - m_{B_s}^2} \frac{1}{k_2^2 - m_{B_s}^2} \\ &\quad \times \frac{-g_\zeta^\sigma + q_\zeta q^\sigma / m_{B_s^*}^2}{q^2 - m_{B_s^*}^2} \mathcal{F}^2(q^2),\end{aligned}\tag{20}$$

$$\begin{aligned}
\mathcal{M}_{(1-2)} = & \int \frac{d^4 q}{(2\pi)^4} [-g_{\Upsilon B_s^* B_s} \mathcal{E}^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu}(-ip_1)_\nu ((ik_2)_\beta - (ik_1)_\beta)] \\
& \times [\epsilon_{\phi\lambda}^* (ig_{B_s^* B_s^* \phi} g^{\delta\sigma} ((-ik_1)^\lambda - (iq)^\lambda) \\
& + 4if_{B_s^* B_s^* \phi}(ip_2)_\rho (g^{\lambda\delta} g^{\rho\sigma} - g^{\lambda\sigma} g^{\rho\delta}))] \\
& \times [ig_{B_s B_s^* \chi_{b1}} \epsilon_{\chi_{b1}}^{*\zeta}] \frac{-g_{\alpha\delta} + k_{1\alpha} k_{1\delta} / m_{B_s^*}^2}{k_1^2 - m_{B_s^*}^2} \\
& \times \frac{1}{k_2^2 - m_{B_s}^2} \frac{-g_{\sigma\zeta} + q_\sigma q_\zeta / m_{B_s^*}^2}{q^2 - m_{B_s^*}^2} \mathcal{F}^2(q^2), \quad (21)
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}_{(1-3)} = & \int \frac{d^4 q}{(2\pi)^4} [g_{\Upsilon B_s B_s^*} \mathcal{E}^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu}(-ip_1)_\nu ((ik_2)_\beta - (ik_1)_\beta)] \\
& \times [-ig_{B_s B_s \phi} \epsilon_{\phi\lambda}^* ((-ik_1)^\lambda - (iq)^\lambda)] \\
& \times [-ig_{B_s^* B_s \chi_{b1}} \epsilon_{\chi_{b1}}^{*\zeta}] \frac{1}{k_1^2 - m_{B_s}^2} \\
& \times \frac{-g_{\alpha\zeta} + k_{2\alpha} k_{2\zeta} / m_{B_s^*}^2}{k_2^2 - m_{B_s^*}^2} \frac{1}{q^2 - m_{B_s}^2} \mathcal{F}^2(q^2), \quad (22)
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}_{(1-4)} = & \int \frac{d^4 q}{(2\pi)^4} [ig_{\Upsilon B_s^* B_s^*} \epsilon_{\Upsilon}^\mu (g_{\nu\alpha} g_{\mu\beta} (ik_2)^\nu - g_{\mu\alpha} g_{\nu\beta} (ik_1)^\nu \\
& - g_{\alpha\beta} ((ik_2)_\mu - (ik_1)_\mu))] \\
& \times [2f_{B_s^* B_s \phi} \mathcal{E}_{\lambda\rho\delta\sigma} \epsilon_{\phi}^{*\lambda} (ip_2)^\rho ((-ik_1)^\delta - (iq)^\delta)] \\
& \times [-ig_{B_s^* B_s \chi_{b1}} \epsilon_{\chi_{b1}}^{*\zeta}] \frac{-g^{\alpha\sigma} + k_1^\alpha k_1^\sigma / m_{B_s^*}^2}{k_1^2 - m_{B_s^*}^2} \\
& \times \frac{-g_{\zeta}^\beta + k_{2\zeta} k_2^\beta / m_{B_s^*}^2}{k_2^2 - m_{B_s^*}^2} \frac{1}{q^2 - m_{B_s}^2} \mathcal{F}^2(q^2). \quad (23)
\end{aligned}$$

As for the $\Upsilon(6S) \rightarrow \chi_{b2}\phi$ transition, the amplitudes corresponding to Fig. 3 are

$$\begin{aligned}
\mathcal{M}_{(2-1)} = & \int \frac{d^4 q}{(2\pi)^4} [-ig_{\Upsilon B_s B_s} \epsilon_{\Upsilon}^\mu ((ik_1)_\mu - (ik_2)_\mu)] \\
& \times [-ig_{B_s B_s \phi} \epsilon_{\phi\lambda}^* ((-ik_1)^\lambda - (iq)^\lambda)] \\
& \times [-g_{B_s B_s \chi_{b2}} \epsilon_{\chi_{b2}}^{*\zeta\eta} (-iq)_\zeta (-ik_2)_\eta] \\
& \times \frac{1}{k_1^2 - m_{B_s}^2} \frac{1}{k_2^2 - m_{B_s}^2} \frac{1}{q^2 - m_{B_s}^2} \mathcal{F}^2(q^2), \quad (24)
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}_{(2-2)} = & \int \frac{d^4 q}{(2\pi)^4} [-ig_{\Upsilon B_s B_s} \epsilon_{\Upsilon}^\mu ((ik_1)_\mu - (ik_2)_\mu)] \\
& \times [-2f_{B_s B_s^* \phi} \mathcal{E}_{\lambda\rho\delta\sigma} \epsilon_{\phi}^{*\lambda} (ip_2)^\rho ((-ik_1)^\delta - (iq)^\delta)] \\
& \times [-ig_{B_s B_s^* \chi_{b2}} \mathcal{E}_{\zeta\omega\kappa\xi} \epsilon_{\chi_{b2}}^{*\zeta\eta} (ip_3)^\kappa (-iq)_\eta (-ik_2)^\xi] \\
& \times \frac{1}{k_1^2 - m_{B_s}^2} \frac{1}{k_2^2 - m_{B_s}^2} \\
& \times \frac{-g^{\sigma\omega} + q^\sigma q^\omega / m_{B_s^*}^2}{q^2 - m_{B_s^*}^2} \mathcal{F}^2(q^2), \quad (25)
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}_{(2-3)} = & \int \frac{d^4 q}{(2\pi)^4} [-g_{\Upsilon B_s^* B_s} \mathcal{E}^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu}(-ip_1)_\nu ((ik_2)_\beta - (ik_1)_\beta)] \\
& \times [2f_{B_s^* B_s \phi} \mathcal{E}_{\lambda\rho\delta\sigma} \epsilon_{\phi}^{*\lambda} (ip_2)^\rho ((-ik_1)^\delta - (iq)^\delta)] \\
& \times [-g_{B_s B_s \chi_{b2}} \epsilon_{\chi_{b2}}^{*\zeta\eta} (-iq)_\zeta (-ik_2)_\eta] \\
& \times \frac{-g_\alpha^\sigma + k_{1\alpha} k_1^\sigma / m_{B_s^*}^2}{k_1^2 - m_{B_s^*}^2} \frac{1}{k_2^2 - m_{B_s}^2} \frac{1}{q^2 - m_{B_s}^2} \mathcal{F}^2(q^2), \quad (26)
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}_{(2-4)} = & \int \frac{d^4 q}{(2\pi)^4} [-g_{\Upsilon B_s^* B_s} \mathcal{E}^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu}(-ip_1)_\nu ((ik_2)_\beta - (ik_1)_\beta)] \\
& \times [\epsilon_{\phi\lambda}^* (ig_{B_s^* B_s^* \phi} g^{\delta\sigma} ((-ik_1)^\lambda - (iq)^\lambda) \\
& + 4if_{B_s^* B_s^* \phi}(ip_2)_\rho (g^{\lambda\delta} g^{\rho\sigma} - g^{\lambda\sigma} g^{\rho\delta}))] \\
& \times [-ig_{B_s B_s^* \chi_{b2}} \mathcal{E}_{\zeta\omega\kappa\xi} \epsilon_{\chi_{b2}}^{*\zeta\eta} (ip_3)^\kappa (-iq)_\eta (-ik_2)^\xi] \\
& \times \frac{-g_{\alpha\delta} - k_{1\alpha} k_{1\delta} / m_{B_s^*}^2}{k_1^2 - m_{B_s^*}^2} \frac{1}{k_2^2 - m_{B_s}^2} \\
& \times \frac{-g_\sigma^\omega + q_\sigma q^\omega / m_{B_s^*}^2}{q^2 - m_{B_s^*}^2} \mathcal{F}^2(q^2), \quad (27)
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}_{(2-5)} = & \int \frac{d^4 q}{(2\pi)^4} [g_{\Upsilon B_s B_s^*} \mathcal{E}^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu}(-ip_1)_\nu ((ik_2)_\beta - (ik_1)_\beta)] \\
& \times [-ig_{B_s B_s \phi} \epsilon_{\phi\lambda}^* ((-ik_1)^\lambda - (iq)^\lambda)] \\
& \times [ig_{B_s^* B_s \chi_{b2}} \mathcal{E}_{\zeta\omega\kappa\xi} \epsilon_{\chi_{b2}}^{*\zeta\eta} (ip_3)^\kappa (-iq)^\xi (-ik_2)_\eta] \\
& \times \frac{1}{k_1^2 - m_{B_s}^2} \frac{-g_\alpha^\sigma + k_{2\alpha} k_2^\sigma / m_{B_s^*}^2}{k_2^2 - m_{B_s^*}^2} \\
& \times \frac{1}{q^2 - m_{B_s}^2} \mathcal{F}^2(q^2), \quad (28)
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}_{(2-6)} = & \int \frac{d^4 q}{(2\pi)^4} [g_{\Upsilon B_s B_s^*} \mathcal{E}^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu}(-ip_1)_\nu ((ik_2)_\beta - (ik_1)_\beta)] \\
& \times [-2f_{B_s B_s^* \phi} \mathcal{E}_{\lambda\rho\delta\sigma} \epsilon_{\phi}^{*\lambda} (ip_2)^\rho ((-ik_1)^\delta - (iq)^\delta)] \\
& \times [g_{B_s^* B_s \chi_{b2}} \mathcal{E}_{\zeta\omega\kappa\xi} \epsilon_{\chi_{b2}}^{*\zeta\eta} (g_{\zeta\kappa} g_{\eta\xi} + g_{\eta\kappa} g_{\zeta\xi})] \\
& \times \frac{1}{k_1^2 - m_{B_s}^2} \frac{-g_\alpha^\kappa + k_{2\alpha} k_2^\kappa / m_{B_s^*}^2}{k_2^2 - m_{B_s^*}^2} \\
& \times \frac{-g^{\sigma\xi} + q^\sigma q^\xi / m_{B_s^*}^2}{q^2 - m_{B_s^*}^2} \mathcal{F}^2(q^2), \quad (29)
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}_{(2-7)} = & \int \frac{d^4 q}{(2\pi)^4} [ig_{\Upsilon B_s^* B_s^*} \epsilon_{\Upsilon}^{\mu} (g_{\nu\alpha} g_{\mu\beta} (ik_2)^{\nu} - g_{\mu\alpha} g_{\nu\beta} (ik_1)^{\nu} \\
& - g_{\alpha\beta} ((ik_2)_{\mu} - (ik_1)_{\mu}))] \\
& \times [2f_{B_s^* B_s^* \phi} \epsilon_{\phi}^{*\lambda} (ip_2)^{\rho} ((-ik_1)^{\delta} - (iq)^{\delta})] \\
& \times [ig_{B_s^* B_s^* \chi_{b2}} \epsilon_{\chi_{b2}}^{*\zeta\eta} (ip_3)^{\kappa} (-iq)^{\xi} (-ik_2)_{\eta}] \\
& \times \frac{-g^{\alpha\sigma} + k_1^{\alpha} k_1^{\sigma}/m_{B_s^*}^2}{k_1^2 - m_{B_s^*}^2} \frac{-g^{\beta\omega} + k_2^{\beta} k_2^{\omega}/m_{B_s^*}^2}{k_2^2 - m_{B_s^*}^2} \\
& \times \frac{1}{q^2 - m_{B_s}^2} \mathcal{F}^2(q^2), \quad (30)
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}_{(2-8)} = & \int \frac{d^4 q}{(2\pi)^4} [ig_{\Upsilon B_s^* B_s^*} \epsilon_{\Upsilon}^{\mu} (g_{\nu\alpha} g_{\mu\beta} (ik_2)^{\nu} - g_{\mu\alpha} g_{\nu\beta} (ik_1)^{\nu} \\
& - g_{\alpha\beta} ((ik_2)_{\mu} - (ik_1)_{\mu}))] \\
& \times [\epsilon_{\phi\lambda}^* (ig_{B_s^* B_s^* \phi} g^{\delta\sigma} ((-ik_1)^{\lambda} - (iq)^{\lambda}) \\
& + 4if_{B_s^* B_s^* \phi} (ip_2)_{\rho} (g^{\lambda\delta} g^{\rho\sigma} - g^{\lambda\sigma} g^{\rho\delta}))] \\
& \times [g_{B_s^* B_s^* \chi_{b2}} \epsilon_{\chi_{b2}}^{*\zeta\eta} (g_{\zeta\kappa} g_{\eta\xi} + g_{\eta\kappa} g_{\zeta\xi})] \\
& \times \frac{-g_{\delta}^{\alpha} + k_{1\delta} k_1^{\alpha}/m_{B_s^*}^2}{k_1^2 - m_{B_s^*}^2} \frac{-g_{\kappa}^{\beta} + k_2^{\beta} k_2^{\kappa}/m_{B_s^*}^2}{k_2^2 - m_{B_s^*}^2} \\
& \times \frac{-g_{\sigma}^{\xi} + q_{\sigma} q^{\xi}/m_{B_s^*}^2}{q^2 - m_{B_s^*}^2} \mathcal{F}^2(q^2). \quad (31)
\end{aligned}$$

As for the $\Upsilon(6S) \rightarrow \chi_{b0}\omega$ transition, the amplitudes corresponding to Fig. 5 are

$$\begin{aligned}
\mathcal{A}_{(0-1)} = & \int \frac{d^4 q}{(2\pi)^4} [-ig_{\Upsilon BB} \epsilon_{\Upsilon}^{\mu} ((ik_1)_{\mu} - (ik_2)_{\mu})] \\
& \times [-ig_{BB\omega} \epsilon_{\omega\lambda}^* ((-ik_1)^{\lambda} - (iq)^{\lambda})] [-g_{BB\chi_{b0}}] \\
& \times \frac{1}{k_1^2 - m_B^2} \frac{1}{k_2^2 - m_B^2} \frac{1}{q^2 - m_B^2} \mathcal{F}^2(q^2), \quad (32)
\end{aligned}$$

$$\begin{aligned}
\mathcal{A}_{(0-2)} = & \int \frac{d^4 q}{(2\pi)^4} [-g_{\Upsilon B^* B} \epsilon^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu} (-ip_1)_{\nu} ((ik_2)_{\beta} - (ik_1)_{\beta})] \\
& \times [2f_{B^* B\omega} \epsilon_{\omega}^{*\lambda} (ip_2)^{\rho} ((-ik_1)^{\delta} - (iq)^{\delta})] [-g_{BB\chi_{b0}}] \\
& \times \frac{-g_{\alpha}^{\sigma} + k_{1\alpha} k_1^{\sigma}/m_{B^*}^2}{k_1^2 - m_{B^*}^2} \frac{1}{k_2^2 - m_B^2} \frac{1}{q^2 - m_B^2} \mathcal{F}^2(q^2), \quad (33)
\end{aligned}$$

$$\begin{aligned}
\mathcal{A}_{(0-3)} = & \int \frac{d^4 q}{(2\pi)^4} [g_{\Upsilon BB^*} \epsilon^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu} (-ip_1)_{\nu} ((ik_2)_{\beta} - (ik_1)_{\beta})] \\
& \times [-2f_{BB^* \omega} \epsilon_{\omega}^{*\lambda} (ip_2)^{\rho} ((-ik_1)^{\delta} - (iq)^{\delta})] \\
& \times [-g_{B^* B^* \chi_{b0}}] \frac{1}{k_1^2 - m_B^2} \frac{-g_{\alpha}^{\zeta} + k_{2\alpha} k_2^{\zeta}/m_{B^*}^2}{k_2^2 - m_{B^*}^2} \\
& \times \frac{-g_{\zeta}^{\sigma} + q^{\sigma} q_{\zeta}/m_{B^*}^2}{q^2 - m_{B^*}^2} \mathcal{F}^2(q^2), \quad (34)
\end{aligned}$$

$$\begin{aligned}
\mathcal{A}_{(0-4)} = & \int \frac{d^4 q}{(2\pi)^4} [ig_{\Upsilon B^* B^*} \epsilon_{\Upsilon}^{\mu} (g_{\nu\alpha} g_{\mu\beta} (ik_2)^{\nu} - g_{\mu\alpha} g_{\nu\beta} (ik_1)^{\nu} \\
& - g_{\alpha\beta} ((ik_2)_{\mu} - (ik_1)_{\mu}))] \\
& \times [\epsilon_{\omega\lambda}^* (ig_{B^* B^* \omega} g^{\delta\sigma} ((-ik_1)^{\lambda} - (iq)^{\lambda}) \\
& + 4if_{B^* B^* \omega} (ip_2)_{\rho} (g^{\lambda\delta} g^{\rho\sigma} - g^{\lambda\sigma} g^{\rho\delta}))] [-g_{B^* B^* \chi_{b0}}] \\
& \times \frac{-g_{\delta}^{\alpha} + k_{1\delta} k_1^{\alpha}/m_{B^*}^2}{k_1^2 - m_{B^*}^2} \frac{-g_{\kappa}^{\beta} + k_2^{\beta} k_2^{\kappa}/m_{B^*}^2}{k_2^2 - m_{B^*}^2} \\
& \times \frac{-g_{\sigma\zeta} + q_{\sigma} q_{\zeta}/m_{B^*}^2}{q^2 - m_{B^*}^2} \mathcal{F}^2(q^2). \quad (35)
\end{aligned}$$

As for the $\Upsilon(6S) \rightarrow \chi_{b1}\omega$ transition, the amplitudes corresponding to Fig. 5 are

$$\begin{aligned}
\mathcal{A}_{(1-1)} = & \int \frac{d^4 q}{(2\pi)^4} [-ig_{\Upsilon BB} \epsilon_{\Upsilon}^{\mu} ((ik_1)_{\mu} - (ik_2)_{\mu})] \\
& \times [-2f_{BB^* \omega} \epsilon_{\omega}^{*\lambda} (ip_2)^{\rho} ((-ik_1)^{\delta} - (iq)^{\delta})] \\
& \times [ig_{BB^* \chi_{b1}} \epsilon_{\chi_{b1}}^{*\zeta}] \frac{1}{k_1^2 - m_B^2} \frac{1}{k_2^2 - m_B^2} \\
& \times \frac{-g_{\zeta}^{\sigma} + q_{\zeta} q^{\sigma}/m_{B^*}^2}{q^2 - m_{B^*}^2} \mathcal{F}^2(q^2), \quad (36)
\end{aligned}$$

$$\begin{aligned}
\mathcal{A}_{(1-2)} = & \int \frac{d^4 q}{(2\pi)^4} [-g_{\Upsilon B^* B} \epsilon^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu} (-ip_1)_{\nu} ((ik_2)_{\beta} - (ik_1)_{\beta})] \\
& \times [\epsilon_{\omega\lambda}^* (ig_{B^* B^* \omega} g^{\delta\sigma} ((-ik_1)^{\lambda} - (iq)^{\lambda}) \\
& + 4if_{B^* B^* \omega} (ip_2)_{\rho} (g^{\lambda\delta} g^{\rho\sigma} - g^{\lambda\sigma} g^{\rho\delta}))] \\
& \times [ig_{BB^* \chi_{b1}} \epsilon_{\chi_{b1}}^{*\zeta}] \frac{-g_{\alpha\delta} + k_{1\alpha} k_{1\delta}/m_{B^*}^2}{k_1^2 - m_{B^*}^2} \\
& \times \frac{1}{k_2^2 - m_B^2} \frac{-g_{\sigma\zeta} + q_{\sigma} q_{\zeta}/m_{B^*}^2}{q^2 - m_{B^*}^2} \mathcal{F}^2(q^2), \quad (37)
\end{aligned}$$

$$\begin{aligned}
\mathcal{A}_{(1-3)} = & \int \frac{d^4 q}{(2\pi)^4} [g_{\Upsilon BB^*} \epsilon^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu} (-ip_1)_{\nu} ((ik_2)_{\beta} - (ik_1)_{\beta})] \\
& \times [-ig_{BB\omega} \epsilon_{\omega\lambda}^* ((-ik_1)^{\lambda} - (iq)^{\lambda})] \\
& \times [-ig_{B^* B\chi_{b1}} \epsilon_{\chi_{b1}}^{*\zeta}] \frac{1}{k_1^2 - m_B^2} \\
& \times \frac{-g_{\alpha\zeta} + k_{2\alpha} k_{2\zeta}/m_{B^*}^2}{k_2^2 - m_{B^*}^2} \frac{1}{q^2 - m_B^2} \mathcal{F}^2(q^2), \quad (38)
\end{aligned}$$

$$\begin{aligned}
\mathcal{A}_{(1-4)} = & \int \frac{d^4 q}{(2\pi)^4} [ig_{\Upsilon B^* B^*} \epsilon_{\Upsilon}^{\mu} (g_{\nu\alpha} g_{\mu\beta} (ik_2)^{\nu} - g_{\mu\alpha} g_{\nu\beta} (ik_1)^{\nu} \\
& - g_{\alpha\beta} ((ik_2)_{\mu} - (ik_1)_{\mu}))] \\
& \times [2f_{B^* B\omega} \epsilon_{\omega}^{*\lambda} (ip_2)^{\rho} ((-ik_1)^{\delta} - (iq)^{\delta})] \\
& \times [-ig_{B^* B\chi_{b1}} \epsilon_{\chi_{b1}}^{*\zeta}] \frac{-g_{\alpha}^{\sigma} + k_1^{\alpha} k_1^{\sigma}/m_{B^*}^2}{k_1^2 - m_{B^*}^2} \\
& \times \frac{-g_{\zeta}^{\beta} + k_{2\zeta} k_2^{\beta}/m_{B^*}^2}{k_2^2 - m_{B^*}^2} \frac{1}{q^2 - m_B^2} \mathcal{F}^2(q^2). \quad (39)
\end{aligned}$$

As for the $\Upsilon(6S) \rightarrow \chi_{b2}\omega$ transition, the amplitudes corresponding to Fig. 6 are

$$\begin{aligned} \mathcal{A}_{(2-1)} = & \int \frac{d^4 q}{(2\pi)^4} [-ig_{\Upsilon BB} \epsilon_{\Upsilon}^{\mu} ((ik_1)_{\mu} - (ik_2)_{\mu})] \\ & \times [-ig_{BB\omega} \epsilon_{\omega\lambda}^* ((-ik_1)^{\lambda} - (iq)^{\lambda})] \\ & \times [-g_{BB\chi_{b2}} \epsilon_{\chi_{b2}}^{*\zeta\eta} (-iq)_{\zeta} (-ik_2)_{\eta}] \\ & \times \frac{1}{k_1^2 - m_B^2} \frac{1}{k_2^2 - m_B^2} \frac{1}{q^2 - m_B^2} \mathcal{F}^2(q^2), \quad (40) \end{aligned}$$

$$\begin{aligned} \mathcal{A}_{(2-2)} = & \int \frac{d^4 q}{(2\pi)^4} [-ig_{\Upsilon BB} \epsilon_{\Upsilon}^{\mu} ((ik_1)_{\mu} - (ik_2)_{\mu})] \\ & \times [-2f_{BB^* \omega} \epsilon_{\lambda\rho\delta\sigma} \epsilon_{\omega}^{*\lambda} (ip_2)^{\rho} ((-ik_1)^{\delta} - (iq)^{\delta})] \\ & \times [-ig_{BB^* \chi_{b2}} \epsilon_{\chi_{b2}}^{*\zeta\eta} \epsilon_{\zeta\omega\kappa\xi} (ip_3)^{\kappa} (-iq)_{\eta} (-ik_2)_{\xi}] \\ & \times \frac{1}{k_1^2 - m_B^2} \frac{1}{k_2^2 - m_B^2} \\ & \times \frac{-g^{\sigma\omega} + q^{\sigma} q^{\omega} / m_{B^*}^2}{q^2 - m_{B^*}^2} \mathcal{F}^2(q^2), \quad (41) \end{aligned}$$

$$\begin{aligned} \mathcal{A}_{(2-3)} = & \int \frac{d^4 q}{(2\pi)^4} [-g_{\Upsilon B^* B} \epsilon^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu} (-ip_1)_{\nu} ((ik_2)_{\beta} - (ik_1)_{\beta})] \\ & \times [2f_{B^* B\omega} \epsilon_{\lambda\rho\delta\sigma} \epsilon_{\omega}^{*\lambda} (ip_2)^{\rho} ((-ik_1)^{\delta} - (iq)^{\delta})] \\ & \times [-g_{BB\chi_{b2}} \epsilon_{\chi_{b2}}^{*\zeta\eta} (-iq)_{\zeta} (-ik_2)_{\eta}] \\ & \times \frac{-g_{\alpha}^{\sigma} + k_{1\alpha} k_1^{\sigma} / m_{B^*}^2}{k_1^2 - m_{B^*}^2} \frac{1}{k_2^2 - m_B^2} \frac{1}{q^2 - m_B^2} \mathcal{F}^2(q^2), \quad (42) \end{aligned}$$

$$\begin{aligned} \mathcal{A}_{(2-4)} = & \int \frac{d^4 q}{(2\pi)^4} [-g_{\Upsilon B^* B} \epsilon^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu} (-ip_1)_{\nu} ((ik_2)_{\beta} - (ik_1)_{\beta})] \\ & \times [\epsilon_{\omega\lambda}^* (ig_{B^* B^* \omega} g^{\delta\sigma} ((-ik_1)^{\lambda} - (iq)^{\lambda}) \\ & + 4if_{B^* B^* \omega} (ip_2)_{\rho} (g^{\lambda\delta} g^{\rho\sigma} - g^{\lambda\sigma} g^{\rho\delta}))] \\ & \times [-ig_{BB^* \chi_{b2}} \epsilon_{\chi_{b2}}^{*\zeta\eta} \epsilon_{\zeta\omega\kappa\xi} (ip_3)^{\kappa} (-iq)_{\eta} (-ik_2)_{\xi}] \\ & \times \frac{-g_{\alpha\delta} - k_{1\alpha} k_{1\delta} / m_{B^*}^2}{k_1^2 - m_{B^*}^2} \frac{1}{k_2^2 - m_B^2} \\ & \times \frac{-g_{\sigma}^{\omega} + q_{\sigma} q^{\omega} / m_{B^*}^2}{q^2 - m_{B^*}^2} \mathcal{F}^2(q^2), \quad (43) \end{aligned}$$

$$\begin{aligned} \mathcal{A}_{(2-5)} = & \int \frac{d^4 q}{(2\pi)^4} [g_{\Upsilon BB^*} \epsilon^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu} (-ip_1)_{\nu} ((ik_2)_{\beta} - (ik_1)_{\beta})] \\ & \times [-ig_{BB\omega} \epsilon_{\omega\lambda}^* ((-ik_1)^{\lambda} - (iq)^{\lambda})] \\ & \times [ig_{B^* B\chi_{b2}} \epsilon_{\chi_{b2}}^{*\zeta\eta} \epsilon_{\zeta\omega\kappa\xi} (ip_3)^{\kappa} (-iq)_{\eta} (-ik_2)_{\xi}] \\ & \times \frac{1}{k_1^2 - m_B^2} \frac{-g_{\alpha}^{\omega} + k_{2\alpha} k_2^{\omega} / m_{B^*}^2}{k_2^2 - m_{B^*}^2} \\ & \times \frac{1}{q^2 - m_B^2} \mathcal{F}^2(q^2), \quad (44) \end{aligned}$$

$$\begin{aligned} \mathcal{A}_{(2-6)} = & \int \frac{d^4 q}{(2\pi)^4} [g_{\Upsilon BB^*} \epsilon^{\mu\nu\alpha\beta} \epsilon_{\Upsilon\mu} (-ip_1)_{\nu} ((ik_2)_{\beta} - (ik_1)_{\beta})] \\ & \times [-2f_{BB^* \omega} \epsilon_{\lambda\rho\delta\sigma} \epsilon_{\omega}^{*\lambda} (ip_2)^{\rho} ((-ik_1)^{\delta} - (iq)^{\delta})] \\ & \times [g_{B^* B^* \chi_{b2}} \epsilon_{\chi_{b2}}^{*\zeta\eta} (g_{\zeta\kappa} g_{\eta\xi} + g_{\eta\kappa} g_{\zeta\xi})] \\ & \times \frac{1}{k_1^2 - m_B^2} \frac{-g_{\alpha}^{\kappa} + k_{2\alpha} k_2^{\kappa} / m_{B^*}^2}{k_2^2 - m_{B^*}^2} \\ & \times \frac{-g^{\sigma\xi} + q^{\sigma} q^{\xi} / m_{B^*}^2}{q^2 - m_{B^*}^2} \mathcal{F}^2(q^2), \quad (45) \end{aligned}$$

$$\begin{aligned} \mathcal{A}_{(2-7)} = & \int \frac{d^4 q}{(2\pi)^4} [ig_{\Upsilon B^* B^*} \epsilon_{\Upsilon}^{\mu} (g_{\nu\alpha} g_{\mu\beta} (ik_2)^{\nu} - g_{\mu\alpha} g_{\nu\beta} (ik_1)^{\nu} \\ & - g_{\alpha\beta} ((ik_2)_{\mu} - (ik_1)_{\mu}))] \\ & \times [2f_{B^* B\omega} \epsilon_{\lambda\rho\delta\sigma} \epsilon_{\omega}^{*\lambda} (ip_2)^{\rho} ((-ik_1)^{\delta} - (iq)^{\delta})] \\ & \times [ig_{B^* B\chi_{b2}} \epsilon_{\chi_{b2}}^{*\zeta\eta} \epsilon_{\zeta\omega\kappa\xi} (ip_3)^{\kappa} (-iq)_{\eta} (-ik_2)_{\xi}] \\ & \times \frac{-g^{\alpha\sigma} + k_1^{\alpha} k_1^{\sigma} / m_{B^*}^2}{k_1^2 - m_{B^*}^2} \frac{-g^{\beta\omega} + k_2^{\beta} k_2^{\omega} / m_{B^*}^2}{k_2^2 - m_{B^*}^2} \\ & \times \frac{1}{q^2 - m_B^2} \mathcal{F}^2(q^2), \quad (46) \end{aligned}$$

$$\begin{aligned} \mathcal{A}_{(2-8)} = & \int \frac{d^4 q}{(2\pi)^4} [ig_{\Upsilon B^* B^*} \epsilon_{\Upsilon}^{\mu} (g_{\nu\alpha} g_{\mu\beta} (ik_2)^{\nu} - g_{\mu\alpha} g_{\nu\beta} (ik_1)^{\nu} \\ & - g_{\alpha\beta} ((ik_2)_{\mu} - (ik_1)_{\mu}))] \\ & \times [\epsilon_{\omega\lambda}^* (ig_{B^* B^* \omega} g^{\delta\sigma} ((-ik_1)^{\lambda} - (iq)^{\lambda}) \\ & + 4if_{B^* B^* \omega} (ip_2)_{\rho} (g^{\lambda\delta} g^{\rho\sigma} - g^{\lambda\sigma} g^{\rho\delta}))] \\ & \times [g_{B^* B^* \chi_{b2}} \epsilon_{\chi_{b2}}^{*\zeta\eta} (g_{\zeta\kappa} g_{\eta\xi} + g_{\eta\kappa} g_{\zeta\xi})] \\ & \times \frac{-g_{\delta}^{\alpha} + k_{1\delta} k_1^{\alpha} / m_{B^*}^2}{k_1^2 - m_{B^*}^2} \frac{-g^{\beta\kappa} + k_2^{\beta} k_2^{\kappa} / m_{B^*}^2}{k_2^2 - m_{B^*}^2} \\ & \times \frac{-g_{\sigma}^{\xi} + q_{\sigma} q^{\xi} / m_{B^*}^2}{q^2 - m_{B^*}^2} \mathcal{F}^2(q^2). \quad (47) \end{aligned}$$

-
- [1] K. F. Chen *et al.* [Belle Collaboration], Observation of anomalous $\Upsilon(1S)\pi^+\pi^-$ and $\Upsilon(2S)\pi^+\pi^-$ production near the $\Upsilon(5S)$ resonance, Phys. Rev. Lett. **100**, 112001 (2008).
- [2] X. H. He *et al.* [Belle Collaboration], Observation of $e^+e^- \rightarrow \pi^+\pi^-\pi^0\chi_{bJ}$ and Search for $X_b \rightarrow \omega\Upsilon(1S)$ at $\sqrt{s} = 10.867$ GeV, Phys. Rev. Lett. **113**, 142001 (2014).
- [3] A. Bondar *et al.* [Belle Collaboration], Observation of two charged bottomonium-like resonances in $\Upsilon(5S)$ decays, Phys. Rev. Lett. **108**, 122001 (2012).
- [4] D. Y. Chen, X. Liu and S. L. Zhu, Charged bottomonium-like states $Z_b(10610)$ and $Z_b(10650)$ and the $\Upsilon(5S) \rightarrow \Upsilon(2S)\pi^+\pi^-$ decay, Phys. Rev. D **84**, 074016 (2011).
- [5] C. Meng and K. T. Chao, Scalar resonance contributions to the dipion transition rates of $\Upsilon(4S, 5S)$ in the re-scattering model, Phys. Rev. D **77**, 074003 (2008).
- [6] C. Meng and K. T. Chao, Peak shifts due to $B^*\bar{B}^*$ rescattering in $\Upsilon(5S)$ dipion transitions, Phys. Rev. D **78**, 034022 (2008).
- [7] Y. A. Simonov and A. I. Veselov, Bottomonium dipion transitions, Phys. Rev. D **79**, 034024 (2009).
- [8] D. Y. Chen, J. He, X. Q. Li and X. Liu, Dipion invariant mass distribution of the anomalous $\Upsilon(1S)\pi^+\pi^-$ and $\Upsilon(2S)\pi^+\pi^-$ production near the peak of $\Upsilon(10860)$, Phys. Rev. D **84**, 074006 (2011).
- [9] D. Y. Chen, X. Liu and T. Matsuki, Explaining the anomalous $\Upsilon(5S) \rightarrow \chi_{bJ}\omega$ decays through the hadronic loop effect, Phys. Rev. D **90**, 034019 (2014).
- [10] D. Y. Chen and X. Liu, $Z_b(10610)$ and $Z_b(10650)$ structures produced by the initial single pion emission in the $\Upsilon(5S)$ decays, Phys. Rev. D **84**, 094003 (2011).
- [11] C. Patrignani *et al.* [Particle Data Group Collaboration], Review of Particle Physics, Chin. Phys. C **40**, 100001 (2016).
- [12] C. Meng and K. T. Chao, $\Upsilon(4S, 5S)$ to $\Upsilon(1S)$ eta transitions in the rescattering model and the new BaBar measurement, Phys. Rev. D **78**, 074001 (2008).
- [13] B. Wang, D. Y. Chen and X. Liu, Prediction of anomalous $\Upsilon(5S) \rightarrow \Upsilon(1^3D_J)\eta$ transitions, Phys. Rev. D **94**, 094039 (2016).
- [14] X. Liu, X. Q. Zeng and X. Q. Li, Study on contributions of hadronic loops to decays of J/ψ to vector + pseudoscalar mesons, Phys. Rev. D **74**, 074003 (2006).
- [15] X. Liu, B. Zhang and X. Q. Li, The Puzzle of excessive non- $D\bar{D}$ component of the inclusive $\psi(3770)$ decay and the long-distant contribution, Phys. Lett. B **675**, 441 (2009).
- [16] G. Li, X. h. Liu, Q. Wang and Q. Zhao, Further understanding of the non- $D\bar{D}$ decays of $\psi(3770)$, Phys. Rev. D **88**, 014010 (2013).
- [17] O. Gortchakov, M. P. Locher, V. E. Markushin and S. von Rotz, Two meson doorway calculation for $\bar{p}p \rightarrow \phi\pi$ including off-shell effects and the OZI rule, Z. Phys. A **353**, 447 (1996).
- [18] R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio and G. Nardulli, Phenomenology of heavy meson chiral Lagrangians, Phys. Rept. **281**, 145 (1997).
- [19] O. Kaymakcalan, S. Rajeev and J. Schechter, Nonabelian Anomaly and Vector Meson Decays, Phys. Rev. D **30**, 594 (1984).
- [20] Y. s. Oh, T. Song and S. H. Lee, J/ψ absorption by π and ρ mesons in meson exchange model with anomalous parity interactions, Phys. Rev. C **63**, 034901 (2001).
- [21] P. Colangelo, F. De Fazio and T. N. Pham, $B \rightarrow K^-\chi_{c0}$ decay from charmed meson rescattering, Phys. Lett. B **542**, 71 (2002).
- [22] H. Y. Cheng, C. Y. Cheung, G. L. Lin, Y. C. Lin, T. M. Yan and H. L. Yu, Chiral Lagrangians for radiative decays of heavy hadrons, Phys. Rev. D **47**, 1030 (1993).
- [23] T. M. Yan, H. Y. Cheng, C. Y. Cheung, G. L. Lin, Y. C. Lin and H. L. Yu, Heavy Quark Symmetry And Chiral Dynamics, Phys. Rev. D **46**, 1148 (1992).
- [24] M. B. Wise, Chiral Perturbation Theory For Hadrons Containing A Heavy Quark, Phys. Rev. D **45**, R2188 (1992).
- [25] G. Burdman and J. F. Donoghue, Union of chiral and heavy quark symmetries, Phys. Lett. B **280**, 287 (1992).
- [26] A. F. Falk and M. E. Luke, Strong decays of excited heavy mesons in chiral perturbation theory, Phys. Lett. B **292**, 119 (1992).
- [27] S. Godfrey and K. Moats, Bottomonium Mesons and Strategies for their Observation, Phys. Rev. D **92**, 054034 (2015).
- [28] E. V. Veliev, H. Sundu, K. Azizi and M. Bayar, Scalar Quarkonia at Finite Temperature, Phys. Rev. D **82**, 056012 (2010).
- [29] H. Y. Cheng, C. K. Chua and A. Soni, Final state interactions in hadronic B decays, Phys. Rev. D **71**, 014030 (2005).
- [30] Bo-Qun Wang, for Belle Collaboration, [Status of SuperKEKB and BelleII Experiment](#), 4th Workshop on the XYZ particles, Behang, Beijing, China, Nov. 25, 2016.