

# Strong isospin breaking at production of light scalars

N.N. Achasov and G.N. Shestakov

Laboratory of Theoretical Physics, S.L. Sobolev Institute for Mathematics, 630090 Novosibirsk, Russia

It is discussed breaking the isotopic symmetry as the tool of studying the production and nature of light scalar mesons.

## I. INTRODUCTION

The thirty seven years ago we discovered theoretically a threshold phenomenon known as the mixing of  $a_0^0(980)$  and  $f_0(980)$  resonances that breaks the isotopic invariance considerably, since the effect  $\sim \sqrt{2(M_{K^0} - M_{K^+})/M_{K^0}} \approx 0,13$  in the module of the amplitude [1]; see also Ref. [2]. This effect appears as the narrow,  $2(M_{K^0} - M_{K^+}) \approx 8$  MeV, resonant structure between the  $K^+K^-$  and  $K^0\bar{K}^0$  thresholds,  $a_0^0(980) \rightarrow K\bar{K} \rightarrow f_0(980)$  and vice versa. Since that time many new proposals were appeared, concerning both the searching it and estimating the effects related with this phenomenon [3-29].

Nowadays, this phenomenon has been discovered experimentally and studied with the help of detectors VES in Protvino [30, 31] and BESIII in Beijing [32–34] in the processes

- (a)  $\pi^- N \rightarrow \pi^- f_1(1285) N \rightarrow \pi^- f_0(980) \pi^0 N \rightarrow \pi^- \pi^+ \pi^- \pi^0 N$  [30, 31],
- (b)  $J/\psi \rightarrow \phi f_0(980) \rightarrow \phi a_0(980) \rightarrow \phi \eta \pi^0$  [32],
- (c)  $\chi_{c1} \rightarrow a_0(980) \pi^0 \rightarrow f_0(980) \pi^0 \rightarrow \pi^+ \pi^- \pi^0$  [32],
- (d)  $J/\psi \rightarrow \gamma \eta(1405) \rightarrow \gamma f_0(980) \pi^0 \rightarrow \gamma 3\pi$  [33],
- (e)  $J/\psi \rightarrow \phi f_0(980) \pi^0 \rightarrow \phi 3\pi$  [34],
- (f)  $J/\psi \rightarrow \phi f_1(1285) \rightarrow \phi f_0(980) \pi^0 \rightarrow \phi 3\pi$  [34]

It has become clear [35, 36] that the similar isospin breaking effect can appear not only due to the  $a_0^0(980) - f_0(980)$  mixing, but also for any mechanism of the production of the  $K\bar{K}$  pairs in the  $S$  wave,  $X \rightarrow K\bar{K} \rightarrow f_0(980)/a_0^0(980)$ .<sup>1</sup> Thus a new tool to study the production mechanism and nature of light scalars is emerged.

## II. THE $a_0^0(980) - f_0(980)$ MIXING

The main contribution to the  $a_0^0(980) - f_0(980)$  mixing amplitude, caused by the diagrams shown in Fig. 1, has the form

$$\Pi_{a_0^0 f_0}(m) = \frac{g_{a_0^0 K^+ K^-} g_{f_0 K^+ K^-}}{16\pi} \left[ i \left( \rho_{K^+ K^-}(m) \right. \right.$$




FIG. 1. The  $K\bar{K}$  loop mechanism of the  $a_0^0(980) - f_0(980)$  mixing.

$$\begin{aligned} & -\rho_{K^0 \bar{K}^0}(m) \Big) - \frac{\rho_{K^+ K^-}(m)}{\pi} \ln \frac{1 + \rho_{K^+ K^-}(m)}{1 - \rho_{K^+ K^-}(m)} \\ & + \frac{\rho_{K^0 \bar{K}^0}(m)}{\pi} \ln \frac{1 + \rho_{K^0 \bar{K}^0}(m)}{1 - \rho_{K^0 \bar{K}^0}(m)} \Big] \\ & \approx \frac{g_{a_0^0 K^+ K^-} g_{f_0 K^+ K^-}}{16\pi} i \left( \rho_{K^+ K^-}(m) - \rho_{K^0 \bar{K}^0}(m) \right), \end{aligned}$$

where  $m$  (invariant virtual mass of scalar resonances)  $\geq 2m_{K^0}$  and  $\rho_{K\bar{K}}(m) = \sqrt{1 - 4m_K^2/m^2}$ ; in the region  $0 \leq m \leq 2m_K$ ,  $\rho_{K\bar{K}}(m)$  should be replaced by  $i|\rho_{K\bar{K}}(m)|$ . The modulus and the phase of  $\Pi_{a_0^0 f_0}(m)$  are shown in Fig. 2. In the region between the  $K^+K^-$  and  $K^0\bar{K}^0$




FIG. 2. (a) An example of the modulus of the  $a_0^0(980) - f_0(980)$  mixing amplitude. (b) The phase of the  $a_0^0(980) - f_0(980)$  mixing amplitude.

thresholds, which is the 8 MeV wide,

$$\begin{aligned} |\Pi_{a_0^0 f_0}(m)| & \approx \frac{|g_{a_0^0 K^+ K^-} g_{f_0 K^+ K^-}|}{16\pi} \sqrt{\frac{2(m_{K^0} - m_{K^+})}{m_{K^0}}} \\ & \approx 0.127 \frac{|g_{a_0^0 K^+ K^-} g_{f_0 K^+ K^-}|}{16\pi} \simeq 0.03 \text{ GeV}^2 \\ & \approx m_K \sqrt{m_{K^0}^2 - m_{K^+}^2} \approx m_K^{3/2} \sqrt{m_d - m_u}. \end{aligned}$$

Note that  $|\Pi_{\rho^0 \omega}| \approx |\Pi_{\pi^0 \eta}| \approx 0.003 \text{ GeV}^2 \approx (m_d - m_u) \times 1 \text{ GeV}$ .

<sup>1</sup> Each such mechanism reproduces both the narrow resonant peak and the sharp jump of the phase of the amplitude between the  $K^+K^-$  and  $K^0\bar{K}^0$  thresholds.

The branching ratios of the isospin-breaking decays  $f_0(980) \rightarrow \eta\pi^0$  and  $a_0^0(980) \rightarrow \pi^+\pi^-$ , caused by the  $a_0^0(980) - f_0(980)$  mixing, are [36]

$$\begin{aligned} BR(f_0(980) \rightarrow K\bar{K} \rightarrow a_0^0(980) \rightarrow \eta\pi^0) \\ = \int \left| \frac{\Pi_{a_0^0 f_0}(m)}{D_{a_0^0}(m)D_{f_0}(m) - \Pi_{a_0^0 f_0}^2(m)} \right|^2 \\ \times \frac{2m^2 \Gamma_{a_0^0 \rightarrow \eta\pi^0}(m)}{\pi} dm \approx 0.3\%, \\ BR(a_0^0(980) \rightarrow K\bar{K} \rightarrow f_0(980) \rightarrow \pi\pi) \\ = \int \left| \frac{\Pi_{a_0^0 f_0}(m)}{D_{a_0^0}(m)D_{f_0}(m) - \Pi_{a_0^0 f_0}^2(m)} \right|^2 \\ \times \frac{2m^2 \Gamma_{f_0 \rightarrow \pi\pi}(m)}{\pi} dm \approx 0.2\%, \end{aligned}$$

where  $D_{a_0^0}(m)$  and  $D_{f_0}(m)$  are the propagators of the  $a_0^0(980)$  and  $f_0(980)$  resonances, respectively. Figure 3




FIG. 3. Mass spectra in the isospin-violating decays  $f_0(980) \rightarrow \eta\pi^0$  and  $a_0^0(980) \rightarrow \pi^+\pi^-$ , caused by the  $a_0^0(980) - f_0(980)$  mixing. The solid and dashed lines are generally similar each other. The dotted vertical lines show the locations of the  $K^+K^-$  and  $K^0\bar{K}^0$  thresholds.

shows the mass spectra correspond to the integrands in the above equations.<sup>2</sup>

### III. POLARIZATION PHENOMENA

The phase jump (see Fig. 2(b)) suggests the idea to study the  $a_0^0(980) - f_0(980)$  mixing in polarization phenomena [17, 18]. If the process amplitude with the spin




FIG. 4. Manifestation of the  $a_0(980) - f_0(980)$  mixing effect in the reaction  $\pi^-p_\uparrow \rightarrow a_0^0(980)n \rightarrow \eta\pi^0n$  on a polarized target at  $P_{lab}^\pi = 18.3$  GeV in the  $\rho_2$  and  $\pi$  exchange model. The solid (dotted) curves show the spin asymmetry  $A(0 \leq t \leq 0.025 \text{ GeV}^2, m)$  as a function of  $\eta\pi^0$  invariant mass,  $m$  (smoothed with 10 MeV mass resolution).

configuration is dominated by the  $a_0^0(980) - f_0(980)$  mixing then the spin asymmetry of the cross section jumps near the  $K\bar{K}$  thresholds. An example is the reaction  $\pi^-p_\uparrow \rightarrow (a_0^0(980) + f_0(980))n \rightarrow a_0^0(980)n \rightarrow \eta\pi^0n$  on a polarized proton target. The corresponding differential cross section has the form

$$\begin{aligned} \frac{d^3\sigma}{dt dm d\psi} = \frac{1}{2\pi} [ |M_{++}|^2 + |M_{+-}|^2 \\ + 2 \Im(M_{++}M_{+-}^*) P \cos \psi ], \end{aligned}$$

and the dimensionless normalized spin asymmetry is  $A(t, m) = 2 \Im(M_{++}M_{+-}^*) / (|M_{++}|^2 + |M_{+-}|^2)$ ,  $-1 \leq A(t, m) \leq 1$ .<sup>3</sup> Figure 4 illustrates the strong asymmetry jump which is straightforward manifestation of the  $a_0(980) - f_0(980)$  mixing amplitude interfering with the isospin preserving one in the  $\rho_2$  and  $\pi$  Regge exchange model. Details and various variants may be found in Refs. [17, 18].

These effects are still in waiting for their studies.

### IV. THE DECAY $f_1(1285) \rightarrow f_0(980)\pi^0 \rightarrow 3\pi$

Estimated are the contributions of the following mechanisms responsible for the decay  $f_1(1285) \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  [36]:

<sup>2</sup> Here we use the values of the coupling constants of the  $f_0(980)$  and  $a_0^0(980)$  resonances with the  $\pi\pi$ ,  $K\bar{K}$ , and  $\eta\pi$  channels obtained in Ref. [36] from the BESIII data on the intensities of the  $f_0(980) \rightarrow a_0^0(980)$  and  $a_0^0(980) \rightarrow f_0(980)$  transitions measured in the reactions (b) and (c) [32].

<sup>3</sup> Here  $M_{+-}$  and  $M_{++}$  are the  $s$ -channel helicity amplitude with and without nucleon helicity flip,  $\psi$  is the angle between the normal to the reaction plane formed by the momenta of the  $\pi^-$  and  $\eta\pi^0$  system, and the transverse (to the  $\pi^-$  beam axis) polarization of the proton target, and  $P$  is a degree of this polarization.

- : (1) the contribution of the  $a_0^0(980) - f_0(980)$  mixing,  $f_1(1285) \rightarrow a_0(980)\pi^0 \rightarrow (K^+K^- + K^0\bar{K}^0)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$ ,
- : (2) the contribution of the transition  $f_1(1285) \rightarrow (K^+K^- + K^0\bar{K}^0)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$ , arising due to the pointlike decay  $f_1(1285) \rightarrow K\bar{K}\pi^0$ ,
- : (3) the contribution of the transition  $f_1(1285) \rightarrow (K^*\bar{K} + \bar{K}^*K) \rightarrow (K^+K^- + K^0\bar{K}^0)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$ , where  $K^* = K^*(892)$ , and
- : (4) the contribution of the transition  $f_1(1285) \rightarrow (K_0^*\bar{K} + \bar{K}_0^*K) \rightarrow (K^+K^- + K^0\bar{K}^0)\pi^0 \rightarrow f_0(980) \rightarrow \pi^+\pi^-\pi^0$ , where  $K_0^* = K_0^*(800)$  (or  $\kappa$ ) and  $K_0^*(1430)$ .

These mechanisms break the conservation of the isospin due to the nonzero mass difference of the  $K^+$  and  $K^0$  mesons. They result in the appearance of the narrow resonance structure in the  $\pi^+\pi^-\pi^0$  mass spectrum in the region of the  $K\bar{K}$  thresholds, with the width  $\approx 2m_{K^0} - 2m_{K^+} \approx 8$  MeV. The observation of such a structure in experiment is the direct indication on the  $K\bar{K}$  loop mechanism of the breaking of the isotopic invariance.

We point out that existing data should be more precise, and it is difficult to explain them using the single specific mechanism from those listed above. Taking the decay  $f_1(1285) \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  as the example, we discuss the general approach to the description of the  $K\bar{K}$  loop mechanism of the breaking of isotopic invariance.

(1) The matter is that the  $J/\psi \rightarrow \phi f_0(980) \rightarrow \phi a_0(980) \rightarrow \phi \eta \pi^0$  [32] and  $\chi_{c1} \rightarrow a_0(980)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  [32] decays are described by the  $a_0^0(980) - f_0(980)$  mixing well enough:

$$\begin{aligned} & \frac{BR(J/\psi \rightarrow \phi f_0(980) \rightarrow \phi a_0^0(980) \rightarrow \phi \eta \pi^0)}{BR(J/\psi \rightarrow \phi f_0(980) \rightarrow \phi \pi \pi)} \\ &= (0.60 \pm 0.20(stat.) \pm 0.12(sys.) \pm 0.26(para.))\% \\ &\approx \frac{BR(f_0(980) \rightarrow K\bar{K} \rightarrow a_0^0(980) \rightarrow \eta \pi^0)}{BR(f_0(980) \rightarrow \pi \pi)}, \end{aligned}$$

$$\begin{aligned} & \frac{BR(\chi_{c1} \rightarrow a_0^0(980)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0)}{BR(\chi_{c1} \rightarrow a_0^0(980)\pi^0 \rightarrow \eta \pi^0 \pi^0)} \\ &= (0.31 \pm 0.16(stat.) \pm 0.14(sys.) \pm 0.03(para.))\% \\ &\approx \frac{BR(a_0^0(980) \rightarrow K\bar{K} \rightarrow f_0(980) \rightarrow \pi^+\pi^-)}{BR(a_0^0(980) \rightarrow \eta \pi^0)}. \end{aligned}$$

As for the  $f_1(1285) \rightarrow f_0(980)\pi^0 \rightarrow 3\pi$  decay [31], its description requires the “terrible”  $a_0^0(980) - f_0(980)$  mixing:

$$\begin{aligned} & \frac{BR(f_1(1285) \rightarrow a_0^0(980)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0)}{BR(f_1(1285) \rightarrow a_0^0(980)\pi^0 \rightarrow \eta \pi^0 \pi^0)} \\ &= (2.5 \pm 0.9)\% \\ &\approx \frac{BR(a_0^0(980) \rightarrow K\bar{K} \rightarrow f_0(980) \rightarrow \pi^+\pi^-)}{BR(a_0^0(980) \rightarrow \eta \pi^0)}, \end{aligned}$$

and, as a result, the inconvenient coupling constants of the scalar mesons with the pseudo-scalar mesons in the many cases

$$\begin{aligned} \frac{g_{f_0\pi^+\pi^-}^2}{4\pi} &= 1.2 \text{ GeV}^2, \quad \frac{g_{f_0K^+K^-}^2}{4\pi} = 5.7 \text{ GeV}^2, \\ \frac{g_{a_0^0\eta\pi^0}^2}{4\pi} &= 1.9 \text{ GeV}^2, \quad \frac{g_{a_0^0K^+K^-}^2}{4\pi} = 9.9 \text{ GeV}^2. \end{aligned}$$

For example, due to the very strong coupling of  $a_0^0(980)$  with the  $K\bar{K}$  channel, the width of the  $a_0^0(980)$  resonance in the  $\eta\pi^0$  mass spectrum turns out to be near 15 MeV.

(2) The pointlike decay  $f_1(1285) \rightarrow K\bar{K}\pi^0$  gives

$$\frac{BR(f_1(1285) \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0)}{BR(f_1(1285) \rightarrow K\bar{K}\pi)} = 0.0022$$

instead of the experimental value

$$\begin{aligned} & \frac{BR(f_1(1285) \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0)}{BR(f_1(1285) \rightarrow K\bar{K}\pi)} \\ &= 0.033 \pm 0.010. \end{aligned}$$

The  $\pi^+\pi^-\pi^0$  mass spectrum in the decay  $f_1(1285) \rightarrow (K^+K^- + K^0\bar{K}^0)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  looks similar to the curves in Fig. 3 for the  $a_0(980) - f_0(980)$  mixing case. However, it is clear that the pointlike mechanism of the decay  $f_1(1285) \rightarrow K\bar{K}\pi$  cannot by itself provide the considerable probability of the  $f_1(1285) \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  transition.

(3) The isospin-breaking decay  $f_1(1285) \rightarrow (K^*\bar{K} + \bar{K}^*K) \rightarrow (K^+K^- + K^0\bar{K}^0)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  is induced by the diagram shown in Fig. 5, because




FIG. 5. The diagram of the decay  $f_1(1285) \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  via the  $K^*\bar{K} + \bar{K}^*K$  intermediate states.

the contributions from the  $K^+K^-$  and  $K^0\bar{K}^0$  pair production are not compensated entirely. The transition  $f_1(1285) \rightarrow (K^*\bar{K} + \bar{K}^*K) \rightarrow (K^+K^- + K^0\bar{K}^0)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  gives the shape of the  $\pi^+\pi^-\pi^0$  spectrum practically coincides with the corresponding spectrum caused by the  $a_0^0(980) - f_0(980)$  mixing, but its

$$BR(f_1(1285) \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0) \approx 0.0255\%$$

is much less than the experimental value

$$\begin{aligned} & BR(f_1(1285) \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0) \\ &= (0.30 \pm 0.09)\%. \end{aligned}$$

So, the  $f_1(1285) \rightarrow (K^*\bar{K} + \bar{K}^*K) \rightarrow (K^+K^- + K^0\bar{K}^0)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  transition mechanism alone is also insufficient to understand the experimental data.

(4) The variant  $f_1(1285) \rightarrow (K_0^*(800)\bar{K} + \bar{K}_0^*(800)K) \rightarrow (K^+K^- + K^0\bar{K}^0)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  is rejected by the shapes of the  $K\pi$  and  $K\bar{K}$  mass spectra in the decay  $f_1(1285) \rightarrow K\bar{K}\pi$ . As for  $f_1(1285) \rightarrow (K_0^*(1430)\bar{K} + \bar{K}_0^*(1430)K) \rightarrow (K^+K^- + K^0\bar{K}^0)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$ , it provides the results similar to  $f_1(1285) \rightarrow (K^*\bar{K} + \bar{K}^*K) \rightarrow (K^+K^- + K^0\bar{K}^0)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  and consequently cannot describe the data alone.

## V. THE CONSISTENCY CONDITION

The isospin breaking amplitude  $\mathcal{M}_{f_1(1285) \rightarrow f_0(980)\pi^0}(m)$  can be expanded near the  $K\bar{K}$  threshold into the series in  $\rho_{K\bar{K}}(m) = \sqrt{1 - 4m_K^2/m^2}$ :

$$\begin{aligned} \mathcal{M}_{f_1(1285) \rightarrow f_0(980)\pi^0}(m) &= g_{f_0 K^+ K^-} \{ A(m) \\ &\times i[\rho_{K^+ K^-}(m) - \rho_{K^0 \bar{K}^0}(m)] + B(m)[\rho_{K^+ K^-}^2(m) \\ &- \rho_{K^0 \bar{K}^0}^2(m)] + O[\rho_{K^+ K^-}^3(m) - \rho_{K^0 \bar{K}^0}^3(m)] + \dots \}. \end{aligned}$$

With a good accuracy

$$\begin{aligned} \mathcal{M}_{f_1(1285) \rightarrow f_0(980)\pi^0}(m) &= g_{f_0 K^+ K^-} A(m) \\ &\times i[\rho_{K^+ K^-}(m) - \rho_{K^0 \bar{K}^0}(m)]. \end{aligned}$$

The amplitude  $A(m)$  contains the information about all possible mechanisms of production of the  $K\bar{K}$  system with isospin  $I = 1$  in  $S$  wave in the process  $f_1(1285) \rightarrow K\bar{K}\pi$ .

From the data on the decay  $f_1(1285) \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  one can extract the information about  $|A(m)|^2$  in the region of the  $K^+K^-$  and  $K^0\bar{K}^0$  thresholds,

$$\begin{aligned} &\frac{d\Gamma_{f_1(1285) \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0}(m)}{dm} \\ &= \frac{1}{16\pi} |\mathcal{M}_{f_1(1285) \rightarrow f_0(980)\pi^0}(m)|^2 \\ &\times p^3(m) \frac{2m^2 \Gamma_{f_0 \rightarrow \pi^+\pi^-}(m)}{\pi |D_{f_0}(m)|^2}, \end{aligned}$$

where  $p(m) = [m_{f_1}^4 - 2m_{f_1}^2(m^2 + m_\pi^2) + (m^2 - m_\pi^2)^2]^{1/2} / (2m_{f_1})$ . Moreover, the information about  $|A(m)|^2$  at  $m > 2m_K$  can be obtained from the data on the  $K\bar{K}$  mass spectra measured in the decays  $f_1(1285) \rightarrow K\bar{K}\pi$ . For instance, the  $K^+K^-$  spectrum in the decay  $f_1(1285) \rightarrow K^+K^-\pi^0$  can be represented in the form

$$\begin{aligned} &\frac{d\Gamma_{f_1(1285) \rightarrow K^+K^-\pi^0}}{dm} \\ &= \frac{2m}{\pi} \rho_{K^+K^-}(m) p^3(m) |A(m)|^2. \end{aligned}$$

Fitting the data on  $d\Gamma_{f_1 \rightarrow K^+K^-\pi^0}/dm$ , one can find the value  $|A(2m_{K^+})|^2$  and obtain the following approximate

estimate [36]

$$\begin{aligned} &\Gamma_{f_1(1285) \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0} \\ &= |A(2m_{K^+})|^2 2.59 \times 10^{-6} \text{ GeV}^5. \end{aligned}$$

Thus its comparison with the data on the decay  $f_1(1285) \rightarrow \pi^+\pi^-\pi^0$  permits one to verify their consistency with the data on the decay  $f_1(1285) \rightarrow K\bar{K}\pi$  and with the idea of the breaking of isotopic invariance caused by the mass difference of  $K^+$  and  $K^0$  mesons.

## VI. THE DECAY

$$J/\psi \rightarrow \gamma \eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma \pi^+\pi^-\pi^0$$

According to BESIII [33], the mass and the width of the  $\eta(1405)$  peak in the  $\pi^+\pi^-\pi^0$  channel are  $1409.0 \pm 1.7$  MeV and  $48.3 \pm 5.2$  MeV, respectively, while the branching ratio is

$$\begin{aligned} &BR(J/\psi \rightarrow \gamma \eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma \pi^+\pi^-\pi^0) \\ &= (1.50 \pm 0.11 \pm 0.11) \cdot 10^{-5}. \end{aligned}$$

In addition, the BESIII gives the ratio

$$\begin{aligned} &\frac{BR(\eta(1405) \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0)}{BR(\eta(1405) \rightarrow a_0^0(980)\pi^0 \rightarrow \eta\pi^0\pi^0)} \\ &= (17.9 \pm 4.2)\%, \end{aligned}$$

that rules out practically the explanation of the discovered effect by means of the  $a_0(980) - f_0(980)$  mixing.




FIG. 6. The diagram of the decay  $\eta(1405) \rightarrow (K^*\bar{K} + \bar{K}^*K) \rightarrow K\bar{K}\pi \rightarrow \pi^+\pi^-\pi^0$ . In the region of the  $\eta(1405)$  resonance all intermediate particles in the loop of this triangle diagram can lie on their mass shells. That is, in the hypothetical case of the stable  $K^*$  meson the logarithmic singularity appears in the imaginary part of the triangle diagram.

We discuss the possibility of the theoretical explanation of the large breaking of isotopic invariance in the decay  $\eta(1405) \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  by means of the anomalous Landau thresholds (the logarithmic triangle singularities), which are in the transition  $\eta(1405) \rightarrow (K^*\bar{K} + \bar{K}^*K) \rightarrow (K^+K^- + K^0\bar{K}^0)\pi^0 \rightarrow f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$  (see Fig. 6), and show that the account of the finite width of the  $K^*(892)$  ( $\Gamma_{K^* \rightarrow K\pi} \approx 50$  MeV) smoothes the logarithmic singularities in the amplitude and results in the suppression of the calculated width of the decay  $\eta(1405) \rightarrow f_0(980)\pi^0 \rightarrow 3\pi$  by the factor of 6 – 8 in comparison with the case of  $\Gamma_{K^* \rightarrow K\pi} = 0$  [35].

The accounting of the finite width of the  $K^*$  resonance, i.e., the averaging of the amplitude over the resonance Breit–Wigner distribution in accord with the spectral Källén–Lehmann representation for the propagator of the unstable  $K^*$  meson, smoothes the logarithmic singularities of the amplitude and hence makes the compensation of the contributions of the  $K^{*+}K^- + K^{*-}K^+$  and  $K^{*0}\bar{K}^0 + \bar{K}^{*0}K^0$  intermediate states more strong. This results in both the diminishing of the calculated width of the decay  $\eta(1405) \rightarrow \pi^+\pi^-\pi^0$  by a number of times in comparison with the case of  $\Gamma_{K^* \rightarrow K\pi} = 0$ , and in the concentration of the main effect of the isospin breaking in the domain of the  $\pi^+\pi^-$  invariant mass between the  $K\bar{K}$  thresholds.

Assuming the dominance of the  $\eta(1405) \rightarrow (K^*\bar{K} + \bar{K}^*K) \rightarrow K\bar{K}\pi^0$  decay, one obtains

$$BR(J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma 3\pi) \approx 1.12 \cdot 10^{-5},$$

that reasonably agrees with experiment.

## VIa. Conclusion

We also analyze the difficulties related with the assumption of the dominance of the  $\eta(1405) \rightarrow (K^*\bar{K} + \bar{K}^*K) \rightarrow K\bar{K}\pi$  decay mechanism and discuss the possible dynamics of the decay  $\eta(1405) \rightarrow \eta\pi\pi$  [35]. The decisive improvement of the experimental data on the  $K\bar{K}$ ,  $K\pi$ ,  $\eta\pi$ , and  $\pi\pi$  mass spectra in the decay of the resonance structure  $\eta(1405/1475)$  to  $K\bar{K}\pi$  and  $\eta\pi\pi$ , and on the shape of the resonance peaks themselves in the  $K\bar{K}\pi$  and  $\eta\pi\pi$  decay channels is necessary for the further establishing the  $\eta(1405) \rightarrow 3\pi$  decay mechanism.

## ACKNOWLEDGMENTS

The present work is partially supported by the Russian Foundation for Basic Research Grant No. 16-02-00065 and the Presidium of the Russian Academy of Sciences project No. 0314-2015-0011.

- 
- [1] N. N. Achasov, S. A. Devyanin, and G. N. Shestakov, Phys. Lett. B **88**, 367 (1979).
  - [2] N. N. Achasov, S. A. Devyanin, and G. N. Shestakov, Yad. Fiz. **33**, 1337 (1981) [Sov. J. Nucl. Phys. **33**, 715 (1981)]; Usp. Fiz. Nauk **142**, 361 (1984) [Sov. Phys. Usp. **27**, 161 (1984)].
  - [3] A. R. Dzierba, in *Proceedings of the Second Workshop on Physics and Detectors for DAΦNE'95, Frascati, 1995*, edited by R. Baldini, F. Bossi, G. Capon, and G. Pancheri, Frascati Physics Series Vol. 4 (INFN, Frascati, 1996), p. 99.
  - [4] N. N. Achasov and G. N. Shestakov, Phys. Rev. D **56**, 212 (1997); Yad. Fiz. **60**, 1669 (1997) [Phys. At. Nucl. **60**, 1522 (1997)].
  - [5] B. Kerbikov and F. Tabakin, Phys. Rev. C **62**, 064601 (2000).
  - [6] F. E. Close and A. Kirk, Phys. Lett. B **489**, 24 (2000).
  - [7] V. Yu. Grishina, L. A. Kondratyuk, M. Büscher, W. Cassing, and H. Ströher, Phys. Lett. B **521**, 217 (2001).
  - [8] N. N. Achasov and A. V. Kiselev, Phys. Lett. B **534**, 83 (2002).
  - [9] D. Black, M. Harada, and J. Schechter, Phys. Rev. Lett. **88**, 181603 (2002).
  - [10] A. E. Kudryavtsev, V. E. Tarasov, J. Haidenbauer, C. Hanhart, and J. Speth, Phys. Rev. C **66**, 015207 (2002); Yad. Fiz. **66**, 1994 (2003) [Phys. At. Nucl. **66**, 1946 (2003)].
  - [11] M. Buescher, F. P. Sassen, N. N. Achasov, and L. Kondratyuk, arXiv:hep-ph/0301126.
  - [12] L. A. Kondratyuk, E. L. Bratkovskaya, V. Yu. Grishina, M. Büscher, W. Cassing, and H. Ströher, Yad. Fiz. **66**, 155 (2003) [Phys. At. Nucl. **66**, 152 (2003)].
  - [13] C. Hanhart, in *Scalar Mesons: An Interesting Puzzle for QCD*, edited by A. H. Fariborz, AIP Conf. Proc. No. 688 (AIP, New York, 2003), p. 61; Phys. Rep. **397**, 155 (2004).
  - [14] C. Amsler and N. A. Törnqvist, Phys. Rep. **389**, 61 (2004).
  - [15] M. Buescher, Acta Phys. Pol. B **35**, 1055 (2004).
  - [16] Z. G. Wang, W. M. Yang, and S. L. Wan, Eur. Phys. J. C **37**, 223 (2004).
  - [17] N. N. Achasov and G. N. Shestakov, Phys. Rev. Lett. **92**, 182001 (2004).
  - [18] N. N. Achasov and G. N. Shestakov, Phys. Rev. D **70**, 074015 (2004).
  - [19] J. J. Wu, Q. Zhao, and B. S. Zou, Phys. Rev. D **75**, 114012 (2007).
  - [20] J. J. Wu and B. S. Zou, Phys. Rev. D **78**, 074017 (2008).
  - [21] J. J. Wu, X. H. Liu, Q. Zhao, and B. S. Zou, Phys. Rev. Lett. **108**, 081803 (2012).
  - [22] F. Aceti, W. H. Liang, E. Oset, J. J. Wu, and B. S. Zou, Phys. Rev. D **86**, 114007 (2012).
  - [23] X. G. Wu, J. J. Wu, Q. Zhao, and B. S. Zou, Phys. Rev. D **87**, 014023 (2013).
  - [24] L. Roca, Phys. Rev. D **88**, 014045 (2013).
  - [25] F. E. Close and A. Kirk, Phys. Rev. D **91**, 114015 (2015).
  - [26] T. Sekihara and S. Kumano, Phys. Rev. D **92**, 034010 (2015).
  - [27] T. Sekihara and S. Kumano, J. Phys. Soc. Jpn. Conf. Proc. **8**, 022006 (2015).
  - [28] F. Aceti, J. M. Dias, and E. Oset, Eur. Phys. J. A **51**, 48 (2015).
  - [29] W. Wang, Phys. Lett. B **759**, 501 (2016).
  - [30] V. Dorofeev *et al.*, Eur. Phys. J. A **38**, 149 (2008).
  - [31] V. Dorofeev *et al.*, Eur. Phys. J. A **47**, 68 (2011).
  - [32] M. Ablikim *et al.*, Phys. Rev. D **83**, 032003 (2011).
  - [33] M. Ablikim *et al.*, Phys. Rev. Lett. **108**, 182001 (2012).
  - [34] M. Ablikim *et al.*, Phys. Rev. D **92**, 012007 (2015).
  - [35] N. N. Achasov, A. A. Kozhevnikov, and G. N. Shestakov, Phys. Rev. D **92**, 036003 (2015).
  - [36] N. N. Achasov, A. A. Kozhevnikov, and G. N. Shestakov,

Phys. Rev. D **93**, 114027 (2016).