arXiv:1701.05678v1 [hep-ph] 20 Jan 2017

Study of radially excited $D_s(2^1S_0)$ and $D_{sJ}^+(2632)$

Yu Tian, Ze Zhao and Ailin Zhang*
Department of Physics, Shanghai University, Shanghai 200444, China

The $J^P = 0^-$ radial excitation $D_s(2^1S_0)$ is anticipated to have mass 2650 MeV (denoted with $D_s(2650)$) though it has not been observed. $D_s(2650)$ is anticipated to be observed in inclusive e^+e^- and pp collisions in D^*K channel. $D_s(2650)$ can be produced from hadronic decays of higher excited resonances. In a $^3P_0$ model, hadronic production of $D_s(2650)$ has been studied, relevant decay widths have been estimated. The hadronic decays of $D_s(3P)$ have been explored in detail, and the dominant decay channels have been pointed out. Two hadronic decay channels of $D_{sJ}^+(2632)$ were observed by SELEX, but have not been observed by any other experiment. Hadronic decays of $D_{sJ}^+(2632)$ in different assignments have been explored in the $^3P_0$ model. In view of the hadronic decay manner of $D_{sJ}^+(2632)$, we conclude that $D_{sJ}^+(2632)$ cannot be a conventional charmed strange $c\bar{s}$ meson.

PACS numbers: 13.25.Ft; 12.39.-x
Keywords: $^3P_0$ model, Hadronic decay, $D_s(2650)$, $D_{sJ}^+(2632)$

I. INTRODUCTION

The lowest lying $1S$ and $1P$ excited D and D_s states are believed established [1, 2]. In recent years, more and more higher excited D and D_s resonances have been observed though some of them have not been identified. Study of their spectroscopy, production and decay will be helpful for their identification and quark dynamics.

For D resonances, $D(2550)$, $D^*(2600)$, $D(2750)$ and $D^*(2760)$ were observed in inclusive e^+e^- and pp collisions by BaBar [3] and LHCb [4] collaborations, respectively. $D^*(2760)$ was subsequently found consisting of $D_1^*(2760)$ and $D_3^*(2760)$ [5, 6].

For D_s resonances, $D_{s1}^*(2700)$ and $D_{sJ}^*(2860)$ were observed by Belle [7] and BaBar [8] collaborations. $D_{sJ}^*(2860)$ was also found consisting of $D_{s1}^*(2860)$ and $D_{s3}^*(2860)$ [9, 10]. According to Refs. [11–20], we have a tentative assignment of these D and D_s resonances given in Table I

When these highly excited D and D_s resonances are examined, it is interesting to note that the masses of $D_{s1}^*(2700)$, $D_{s1}^*(2860)$ and $D_{s3}^*(2860)$ are about 100 MeV higher than those of $D^*(2600)$, $D_1^*(2760)$ and $D_3^*(2760)$, respectively. Therefore, two missing D_s with masses 2650 MeV and 2850 MeV are expected to correspond to the observed $D(2550)$ and $D(2750)$. In Ref. [16], $D_{sJ}(2850)^{\pm}$ has been predicted with $\Gamma_{total} = 125.1$ MeV and $\Gamma_{total} = 51.6$ MeV corresponding to $(J^P = 2^-, j_q = \frac{3}{2})$ and $(J^P = 2^-, j_q = \frac{5}{2})$, respectively.

$D(2550)$ was suggested as the $J^P = 0^-$ radial excitation $D(2^1S_0)$ [11, 12, 15, 16], and $D(2750)$ was suggested as the $J^P = 2^-$ admixture of $D(1^1D_2)$ and $D(1^3D_2)$ [12, 14, 16]. Therefore, the two missing D_s are expected to have $J^P = 0^-$ and $J^P = 2^-$. In this paper, the $D_s(2^1S_0)$ with mass around 2650 MeV will be denoted as $D_s(2650)$ and be studied. The $J^P = 2^-$ D_s resonance will not be addressed for an unclear mixing.

TABLE I: Tentative assignment of $2S$ and $1D$ excited D and D_s resonances.

$\backslash n^{2S+1}L_J$	2^1S_0	2^3S_1	$1^{1,3}D_2$	1^3D_1	1^3D_3
D	$D(2550)$	$D^*(2600)$	$D(2750)$	$D_1^*(2760)$	$D_3^*(2760)$
D_s	$--$	$D_{s1}^*(2700)$	$--$	$D_{s1}^*(2860)$	$D_{s3}^*(2860)$

In 2600 $\sim$ 2700 MeV energy region, a surprisingly narrow charmed strange state, $D_{sJ}^+(2632)$, was first reported by SELEX [21] in $D_s^+\eta$ and D^0K^+ decay channels: $D_{sJ}^+(2632) \to D^0K^+, D_s^+\eta$. $D_{sJ}(2632)$ was observed with mass $M = 2632.5 \pm 1.7$ MeV and $\Gamma < 17$ MeV. In particular, it has an exotic relative branching ratio $\Gamma(D^0K^+)/\Gamma(D_s^+\eta) = 0.14 \pm 0.06$. However, subsequent search for $D_{sJ}^+(2632)$ by BaBar [22], FOCUS or BELLE has not found any evidence.

Since the report of $D_{sJ}^+(2632)$, there appear many interpretations of this state. In Refs. [23–25], $D_{sJ}^+(2632)$ was interpreted as the $J^P = 1^-$ first radial excitation of $D_s^*(2112)$. The special decay manner was attributed to the node structure of the wave functions in Ref. [23] or to the couple channels effect in Ref. [25]. However, the assignment of $D_{sJ}^+(2632)$ with 2^+ $D_s(^3P_2)$ or 1^- $D_s(2^3S_1)$ was pointed out impossible in Refs. [26, 27], and the assignment of $D_{sJ}^+(2632)$ with $D_s(1^3D_1)$ was advocated in Refs. [27, 28]. In the frame of heavy quark effective theory (HQET) sum rule [29], it was pointed out that $D_{sJ}^+(2632)$ was unlikely to be a conventional orbitally higher excited $c\bar{s}$ resonance.

$D_{sJ}^+(2632)$ was alternatively interpreted as a four-quark state in Refs. [30–32]. The narrow width and exotic decay manner were interpreted in this assignment. However, the predicted partners and more decay channels of $D_{sJ}^+(2632)$ have not been observed. In Refs. [24, 33], the authors concluded that $D_{sJ}^+(2632)$ was an experimental artefact.

The $^3P_0$ model is a phenomenological method first

*Corresponding author: zhangal@staff.shu.edu.cn

put forth by Micu [34] and subsequently developed by Yaouanc *et al.* [35–37]. It has been employed successfully to explore Okubo-Zweig-Iizuka-allowed (OZI-allowed) hadronic decays of hadrons. Hadronic decays of the $J^P = 0^-$ radial excitation D_s with $M = 2673$ MeV and $M = 2650$ MeV have been explored in the $^3P_0$ model [12, 38], but the hadronic production of this missing $D_s(2650)$ from higher excited resonances has not been explored. As that in Refs. [18, 19], it is interesting to explore the hadronic production of $D_s(2650)$ from higher excited resonances.

As for $D_{sJ}^+(2632)$, its hadronic decay has not yet been systematically studied in the $^3P_0$ model. In order to identify $D_{sJ}^+(2632)$, it is required to explore the hadronic decay features of $D_{sJ}^+(2632)$ no matter whether $D_{sJ}^+(2632)$ is an experimental artefact or not.

The paper is organized as follows. In the second section, the $^3P_0$ model and the parameters are briefly introduced. In Sec. III, the numerical results relevant to hadronic decays of $D_s(2650)$, $D_s(3P)$ and $D_{sJ}^+(2632)$ are presented. The summaries and discussions are presented in Sec. IV.

II. $^3P_0$ MODEL

Among models for hadronic decays, the $^3P_0$ model is popularly known as a quark-pair creation (QPC) model. Since the propose of $^3P_0$ model [34–37], it has been extensively applied to mesons and baryons with success. To proceed with a practical computation, the formula in Refs. [19, 39] are presented. In the $^3P_0$ model, the hadronic decay width of $A \to BC$ is

$$\Gamma = \pi^2 \frac{|\vec{p}|}{M_A^2} \sum_{JL} |\mathcal{M}^{JL}|^2 \tag{1}$$

where $\vec{p}$ is the momentum of the final mesons B and C in the initial meson A's center-of-mass frame

$$|\vec{p}| = \frac{\sqrt{[m_A^2 - (m_B - m_C)^2][m_A^2 - (m_B + m_C)^2]}}{2m_A} \tag{2}$$

and M^{JL} is the partial wave amplitude of $A \to BC$. The partial wave amplitude M^{JL} is obtained from the helicity amplitude $\mathcal{M}^{M_{J_A} M_{J_B} M_{J_C}}$ in terms of the Jacob-Wick formula as follows

$$\begin{aligned}\mathcal{M}^{JL}(A \to BC) = & \frac{\sqrt{2L+1}}{2J_A+1} \\ & \times \sum_{M_{J_B}, M_{J_C}} \langle L0JM_{J_A} | J_A M_{J_A} \rangle \\ & \times \langle J_B M_{J_B} J_C M_{J_C} | J, JM_{J_A} \rangle \\ & \times \mathcal{M}^{M_{J_A} M_{J_B} M_{J_C}}(\vec{p})\end{aligned} \tag{3}$$

where $\vec{J} = \vec{J_B} + \vec{J_C}$, $\vec{J_A} = \vec{J_B} + \vec{J_C} + \vec{L}$ and $M_{J_A} = M_{J_B} + M_{J_C}$. The helicity amplitude reads

$$\begin{aligned}&\delta^3(\vec{p}_B + \vec{p}_C)\mathcal{M}^{M_{J_A} M_{J_B} M_{J_C}} \\ &= \sqrt{8E_A E_B E_C}\gamma \sum_{\substack{M_{L_A}, M_{S_A}, \\ M_{L_B}, M_{S_B}, \\ M_{L_C}, M_{S_C}, m}} \langle L_A M_{L_A} S_A M_{S_A} | J_A M_{J_A} \rangle \\ &\times \langle L_B M_{L_B} S_B M_{S_B} | J_B M_{J_B} \rangle \langle L_C M_{L_C} S_C M_{S_C} | J_C M_{J_C} \rangle \\ &\times \langle 1m; 1-m | 00 \rangle \langle \chi^{13}_{S_B M_{S_B}} \chi^{24}_{S_C M_{S_C}} | \chi^{12}_{S_A M_{S_A}} \chi^{34}_{1-m} \rangle \\ &\times \langle \varphi_B^{13} \varphi_C^{24} | \varphi_A^{12} \varphi_0^{34} \rangle I^{M_{L_A}, m}_{M_{L_B}, M_{L_C}}(\vec{p})\end{aligned} \tag{4}$$

where $\vec{p}_B$ and $\vec{p}_C$ ($\vec{p}_B$=-$\vec{p}_C$=$\vec{p}$) are the momenta of final mesons B and C in A's center-of-mass frame, respectively. γ is a phenomenological parameter, which indicates the strength of the quark-pair creation from the vacuum. $I^{M_{L_A}, m}_{M_{L_B}, M_{L_C}}(\vec{p})$ is a momentum integral

$$\begin{aligned}I^{M_{L_A}, m}_{M_{L_B}, M_{L_C}}(\vec{p}) = & \int d\vec{k}_1 d\vec{k}_2 d\vec{k}_3 d\vec{k}_4 \\ & \times \delta^3(\vec{k}_1 + \vec{k}_2 - \vec{p}_A)\delta^3(\vec{k}_3 + \vec{k}_4) \\ & \times \delta^3(\vec{p}_B - \vec{k}_1 - \vec{k}_3)\delta^3(\vec{p}_C - \vec{k}_2 - \vec{k}_4) \\ & \times \Psi^*_{n_B L_B M_{L_B}}(\vec{k}_{13}) \Psi^*_{n_c L_C M_{L_c}}(\vec{k}_{24}) \\ & \times \Psi_{n_A L_A M_{LA}}(\vec{k}_{12}) Y_{1m}\left(\vec{k}_{34}\right)\end{aligned} \tag{5}$$

with a relative momentum of quark i and quark j: $\vec{k}_{ij} = \frac{m_j \vec{k}_i - m_i \vec{k}_j}{m_i + m_j}$ $(i, j = 1, 2, 3, 4)$. $\Psi_{nLM_L}(\vec{k})$ are simple harmonic oscillator (SHO) wave functions for the mesons, and $y_{1m}(\vec{k}) = |\vec{k}| Y_{1m}(\Omega)$ is the solid harmonic polynomial of the created P-wave quark pair.

In terms of $9j$ symbols [37], the flavor matrix element reads

$$\begin{aligned}\langle \varphi_B^{13} \varphi_C^{24} | \varphi_A^{12} \varphi_0^{34} \rangle = & \sum_{I, I^3} \langle I_C I_C^3; I_B I_B^3 | I_A, I_A^3 \rangle \\ & \times [(2I_B + 1)(2I_C + 1)(2I_A + 1)]^{1/2} \\ & \times \begin{Bmatrix} I_1 & I_3 & I_B \\ I_2 & I_4 & I_C \\ I_A & 0 & I_A \end{Bmatrix}\end{aligned} \tag{6}$$

where I_i $(i = 1, 2, 3, 4)$ is the isospin of the four u, d, s or c quark. I_A, I_B and I_C are the isospins of the mesons A, B and C, respectively. I_A^3, I_B^3 and I_C^3 are the third isospin components of the mesons, and $\langle I_C I_C^3; I_B I_B^3 | I_A, I_A^3 \rangle$ is the isospin matrix element.

The spin matrix element reads

$$\begin{aligned}&\left\langle \chi^{13}_{S_B M_{S_B}} \chi^{24}_{S_C M_{S_C}} | \chi^{12}_{S_A M_{S_A}} \chi^{34}_{1-m} \right\rangle \\ &= (-1)^{S_C+1} [3(2S_B + 1)(2S_C + 1)(2S_A + 1)]^{1/2} \\ &\times \sum_{S, M_S} \langle S_B M_{S_B} S_C M_{S_C} | S M_S \rangle \langle S M_S | S_A M_{S_A}; 1, -m \rangle \\ &\times \begin{Bmatrix} 1/2 & 1/2 & S_B \\ 1/2 & 1/2 & S_C \\ S_A & 1 & S \end{Bmatrix}.\end{aligned} \tag{7}$$

With these formula in hand, we proceed with our calculation. In the practical calculation, the parameters have been employed as follows. The masses of the constituent quarks are taken to be, $m_c = 1628$ MeV, $m_u = m_d = 220$ MeV, and $m_s = 419$ MeV [12]. The masses and the effective scale parameters β of relevant mesons are presented in Table II [1, 12], where a common value $\beta = 0.4$ GeV is employed for all light flavor mesons. For resonances included in PDG, their masses in PDG are employed (unlike those in Ref. [12]), while for resonances not included in PDG or not observed by experiment, their masses and β are taken from Ref. [12]. $\gamma = 6.95$ is employed in this paper, where γ is $\sqrt{96\pi}$ times as the $\gamma = 0.4$ adopted in Ref. [12] for a different definition. The meson flavor follows the convention [12]: $D^0 = c\bar{u}$, $D^+ = -c\bar{d}$, $D_s^+ = -c\bar{s}$, $K^+ = -u\bar{s}$, $K^- = s\bar{u}$, $\phi = -s\bar{s}$, and $\eta = (u\bar{u} - d\bar{d})/2 - s\bar{s}/\sqrt{2}$.

III. HADRONIC DECAYS

A. $D_s(2650)$

Two possible hadronic decay channels and partial decay widths of the $J^P = 0^-$ radial excitation D_s with $M = 2650$ MeV ($D_s(2650)$) were calculated in the $^3P_0$ model [38]:

$$\Gamma(D_s(2^1S_0) \to D^*K) = 78 \text{ MeV},$$
$$\Gamma(D_s(2^1S_0) \to D_s^*\eta) = 0.$$

Once the $J^P = 0^-$ radial excitation D_s has a mass $M = 2673$ MeV, the partial decay widths become [38]:

$$\Gamma(D_s(2^1S_0) \to D^*K) = 94 \text{ MeV},$$
$$\Gamma(D_s(2^1S_0) \to D_s^*\eta) = 0.6 \text{ MeV}.$$

In Ref. [12], the results were improved as follows

$$\Gamma(D_s(2^1S_0) \to D^*K) = 73.6 \text{ MeV},$$
$$\Gamma(D_s(2^1S_0) \to D_s^*\eta) = 0.$$

D^*K decay channel is the dominant hadronic decay of $D_s(2650)$, which may be used as a typical evidence to identify the $J^P = 0^-$ radial excitation D_s. $D(2550)$ was observed in inclusive e^+e^- and pp collisions, $D_s(2650)$ is naturally anticipated to be observed in these inclusive e^+e^- and pp collisions in D^*K channel.

In other ways, $D_s(2650)$ can be produced from the hadronic decays of higher excited D and D_s mesons through $D_s(2650)K$ or $D_s(2650)\eta$ channels, respectively. Relevant hadronic partial decay widths have been estimated and presented in Table III.

B. Hadronic decay of $D_s(3P)$ resonances

$D_s(2650)$ can be produced from hadronic decays of higher excited D and D_s mesons. Hadronic decays of $D(2D)$ and $D_s(2D)$ have been explored in the $^3P_0$ model in Refs. [18, 19]. Hadronic decays of $D_s(3P)$ have been studied in Ref. [12], where some decay channels were ignored. In this subsection, hadronic decays of $D_s(3P)$ resonances as an example are explored in detail.

In the $D_s(3P)$ multiplets, $D_s(3^3P_1)$ and $D_s(3^1P_1)$ may mix with each other. For lack of information, the detail of the mixing of $D_s(3^3P_1)$ with $D_s(3^1P_1)$ is not clear. Therefore, the OZI-allowed hadronic decay channels of $D_s(3^3P_0)$, $D_s(3^1P_1)$, $D_s(3^3P_1)$ and $D_s(3^3P_2)$ are studied separately.

The masses of $D_s(3P)$ are employed as follows [12]: $M(D_s(3^3P_0)) = 3412$ MeV, $M(D_s(3^1P_1)) = M(D_s(3^3P_1)) = 3425$ MeV, $M(D_s(3^3P_2)) = 3439$ MeV. The masses of $D_s(3^1P_1)$ and $D_s(3^3P_1)$ are assumed equal to the mean mass value of $D_s(3P_1)$ and $D_s(3P_1')$ in Ref. [12]. The final decay widths are given in Tables IV-VII. All the four $D_s(3P)$ resonances have large total decay widths, which are larger than those in Ref. [12].

From Table IV, $\Gamma_{total}(D_s(3^3P_0)) = 162.3$ MeV and the dominant decay channels of $D_s(3^3P_0)$ are: $(1^1D_2)D(2750)K$, DK, $D(2550)K$ and $D_1(2420)K^*$.

From Table V, $\Gamma_{total}(D_s(3^1P_1)) = 260.4$ MeV and the dominant decay channels of $D_s(3^1P_1)$ are: $D_2^*(2460)K$, $D_3^*(2760)K$, $DK^*(1410)$ and $D_2^*(2460)K^*$.

From Table VI, $\Gamma_{total}(D_s(3^3P_1)) = 204.8$ MeV and the dominant decay channels of $D_s(3^3P_1)$ are: $D_1(2430)K^*$, $D_3^*(2760)K$, $D(2600)K$ and $D_2^*(2460)K$.

From Table VII, $\Gamma_{total}(D_s(3^3P_2)) = 307.9$ MeV and the dominant decay channels of $D_s(3^3P_2)$ are: $D^*K^*(1410)$, $D_2^*(2460)K^*$, $DK^*(1410)$ and $D(2600)K$.

C. $D_{sJ}^+(2632)$

In Table I, $D_{s1}^*(2700)$ and $D_{s3}^*(2860)$ are tentatively identified with $D_s(2^3S_1)$ and $D_s(1^3D_1)$, respectively. However, as introduced in the first section, $D_{sJ}^+(2632)$ was once regarded as a $D_s(2^3S_1)$, $D_s(1^3D_1)$ or their admixture if its lower mass is ignored. In order to have a look at these possibilities, let's explore the hadronic decay of $D_{sJ}^+(2632)$. The mixing of $D_s(2^3S_1)$ and $D_s(1^3D_1)$ will not be taken into account for its unclear mixing feature.

In $D_s(2^3S_1)$ or $D_s(1^3D_1)$ assignment, $D_{sJ}^+(2632)$ may decay into D^0K^+, D^+K^0, $D_s\eta$, $D^{*0}K^+$ and $D^{*+}K^0$ final states. The partial and total decay widths are given in Table VIII.

In the case of a $D_s(2^3S_1)$, we have the following ratios:

$$\frac{\Gamma(DK)}{\Gamma(D^*K)} = 0.79; \quad \frac{\Gamma(D_s^+\eta)}{\Gamma(D^*K)} = 0.08. \tag{8}$$

In the case of a $D_s(1^3D_1)$, we have the following ratios:

$$\frac{\Gamma(DK)}{\Gamma(D^*K)} = 3.95; \quad \frac{\Gamma(D_s^+\eta)}{\Gamma(D^*K)} = 0.25. \tag{9}$$

TABLE II: Masses and β values of mesons (MeV).

States	Mass	β	States	Mass	β
$(1^1S_0)K^{\pm}$	493.677	400	$(1P_1)K_1(1270)^{0(\pm)}$	1272	400
$(1^1S_0)K^0$	497.614	400	$(1P_1')K_1(1400)^{0(\pm)}$	1403	400
$(1^3S_1)K^{*\pm}$	891.66	400	$(1^3P_2)K_2^*(1430)^0$	1432.4	400
$(1^3S_1)K^{*0}$	895.81	400	$(1^3P_2)K_2^*(1430)^{\pm}$	1425.6	400
$(1^1S_0)\eta$	547.862	400	$(1^3P_0)K_0^*(1430)^{0(\pm)}$	1425	400
$(1^3S_1)\phi$	1019.461	400	$(1P_1)D_1(2430)^{0(\pm)}$	2427	475
$(1^1S_0)D^{\pm}$	1869.61	601	$(1P_1')D_1(2420)^0$	2421.4	475
$(1^1S_0)D^0$	1864.84	601	$(1P_1')D_1(2420)^{\pm}$	2423.2	475
$(1^3S_1)D^{*\pm}$	2010.26	516	$(1^3P_0)D_0(2400)^0$	2318	516
$(1^3S_1)D^{*0}$	2006.96	516	$(1^3P_0)D_0(2400)^{\pm}$	2403	516
$(1^1S_0)D_s$	1968.30	651	$(1^3P_2)D_2(2460)^{\pm}$	2464.3	437
$(1^3S_1)D_S^*$	2112.1	562	$(1^3P_2)D_2(2460)^0$	2462.6	437
$(2^1S_0)K(1460)^{0(\pm)}$	1460	400	$(2^3S_1)D^*(2600)^0$	2608.7	434
$(2^3S_1)K^*(1410)^{0(\pm)}$	1414	400	$(2^3S_1)D^*(2600)^+$	2621.3	434
$(2^1S_0)D(2550)^{0(\pm)}$	2539.4	450	$(1^3D_1)D_1^*(2760)^0$	2763.3	456
$(1^3P_0)D_{s0}(2317)$	2317.7	542	$(1^3D_1)D_1^*(2760)^+$	2769.7	456
$(1P_1)D_{s1}(2460)$	2459.5	498	$(1^3D_3)D_3^*(2760)^0$	2763.3	407
$(1P_1')D_{s1}(2536)$	2535.10	498	$(1^3D_3)D_3^*(2760)^+$	2769.7	407
$(1^3P_2)D_{s2}(2573)$	2571.9	464	$(1^1D_2)D(2750)^0$	2752.4	428
$(2^3S_1)D_{s1}^*(2700)$	2708.3	458	$(1^1D_2)D(2750)^+$	2752.4	428
$(1^3D_1)D_{s1}^*(2860)$	2859	469	$(1^3D_2)D(2750)^0$	2752.4	428
$(1^3D_3)D_{s3}^*(2860)$	2860.5	426	$(1^3D_2)D(2750)^+$	2752.4	428

TABLE III: Hadronic partial decay widths of $D_s(2650)$ from highly excited resonances (MeV)

Mode	Width	Mode	Width
$D(4^3S_1) \to D_s(2650)K$	0.1	$D_s(4^3S_1) \to D_s(2650)\eta$	0.0
$D(5^3S_1) \to D_s(2650)K$	0.0	$D_s(5^3S_1) \to D_s(2650)\eta$	0.0
$D(3^3P_0) \to D_s(2650)K$	1.8	$D_s(3^3P_0) \to D_s(2650)\eta$	0.1
$D(3^3P_2) \to D_s(2650)K$	2.8	$D_s(3^3P_2) \to D_s(2650)\eta$	2.4
$D(4^3P_0) \to D_s(2650)K$	0.2	$D_s(4^3P_0) \to D_s(2650)\eta$	0.0
$D(4^3P_2) \to D_s(2650)K$	0.0	$D_s(4^3P_2) \to D_s(2650)\eta$	0.2
$D(2^3D_1) \to D_s(2650)K$	6.1	$D_s(2^3D_1) \to D_s(2650)\eta$	4.7
$D(2^3D_3) \to D_s(2650)K$	0.0	$D_s(2^3D_3) \to D_s(2650)\eta$	0.1
$D(3^3D_1) \to D_s(2650)K$	0.0	$D_s(3^3D_1) \to D_s(2650)\eta$	0.5
$D(3^3D_3) \to D_s(2650)K$	1.4	$D_s(3^3D_3) \to D_s(2650)\eta$	1.0
$D(1^3F_2) \to D_s(2650)K$	$--$	$D_s(1^3F_2) \to D_s(2650)\eta$	0.0
$D(2^3F_2) \to D_s(2650)K$	4.5	$D_s(2^3F_2) \to D_s(2650)\eta$	3.0
$D(2^3F_4) \to D_s(2650)K$	0.6	$D_s(2^3F_4) \to D_s(2650)\eta$	0.8
$D(1^3G_3) \to D_s(2650)K$	0.6	$D_s(1^3G_3) \to D_s(2650)\eta$	0.6
$D(1^3G_5) \to D_s(2650)K$	0.0	$D_s(1^3G_5) \to D_s(2650)\eta$	0.0
$D(2^3G_3) \to D_s(2650)K$	1.6	$D_s(2^3G_3) \to D_s(2650)\eta$	1.0
$D(2^3G_5) \to D_s(2650)K$	0.7	$D_s(2^3G_5) \to D_s(2650)\eta$	1.0

Obviously, the decay manner of $D_{sJ}^+(2632)$ is different in different assignments. In fact, this decay manner can be employed to distinguish $D_s(2^3S_1)$ from $D_s(1^3D_1)$.

In particular, the estimated ratios $\frac{\Gamma(D^0K^+)}{\Gamma(D_s^+\eta)}$, 5.02 and 7.90, are much larger than the observed 0.14 ± 0.06. Obviously, $D_{sJ}^+(2632)$ cannot be a conventional charmed strange $c\bar{s}$ meson.

IV. CONCLUSIONS AND DISCUSSIONS

The missing $J^P = 0^-$ radial excitation $D_s(2650)$ is anticipated to have mass 2650 MeV. D^*K channel is the dominant hadronic decay channel of $D_s(2650)$, and $D_s(2650)$ is anticipated to be observed in inclusive e^+e^- and pp collisions in D^*K channel. $D_s(2650)$ can be produced from higher excited D through $D_s(2650)K$ channel or from D_s meson through $D_s(2650)\eta$ channel.

Hadronic decays of $D_s(3P)$ have been studied. All the hadronic decay channels are given, and the decay

TABLE IV: Hadronic decay widths of $D_s(3^3P_0)$ (MeV).

Channels	Width	Channels	Width
D^0K^+	10.5	$D_1(2420)^+K^0$	0.1
D^+K^0	10.3	$D_{s1}(2536)\eta$	0.3
$D_s\eta$	0.2	$D_1(2430)^0K^+$	7.8
$D(2550)^0K^+$	9.8	$D_1(2430)^+K^0$	7.6
$D(2550)^+K^0$	9.7	$D_{s1}(2460)\eta$	0.2
$D^0K(1460)^+$	5.5	$D_1(2420)^0K^{*+}$	8.1
$D^+K(1460)^0$	6.9	$D_1(2420)^+K^{*0}$	8.2
$D^{*0}K^{*+}$	3.2	$D_1(2430)^0K^{*+}$	5.7
$D^{*+}K^{*0}$	2.8	$D_1(2430)^+K^{*0}$	5.7
$D_s^*\phi$	4.0	$D^{*0}K_1(1270)^+$	1.1
$D_0^*(2400)^0K^{*+}$	0.6	$D^{*+}K_1(1270)^0$	1.1
$D_0^*(2400)^+K^{*0}$	2.2	$D^{*0}K_1(1400)^+$	0.3
$D_{s0}^*(2317)\phi$	2.0	$D_2^*(2460)^0K^{*+}$	3.0
$D^0K_1(1270)^+$	0.8	$D_2^*(2460)^+K^{*0}$	2.5
$D^+K_1(1270)^0$	0.6	$D(2750)^0K^+$	19.3
$D^0K_1(1400)^+$	1.5	$D(2750)^+K^0$	18.8
$D^+K_1(1400)^0$	1.7	$(2^1S_0)D_s(2650)\eta$	0.1
$D_1(2420)^0K^+$	0.1	Γ_{total}	162.3

widths have been estimated. The four $D_s(3P)$ resonances have large total decay widths. The dominant decay channels of $D_s(3^3P_0)$ are: $(1^1D_2)D(2750)K$, DK, $D(2550)K$ and $D_1(2420)K^*$. The dominant decay channels of $D_s(3^1P_1)$ are: $D_2^*(2460)K$, $D_3^*(2760)K$, $DK^*(1410)$ and $D_2^*(2460)K^*$. The dominant decay channels of $D_s(3^3P_1)$ are: $D_1(2430)K^*$, $D_3^*(2760)K$, $D(2600)K$ and $D_2^*(2460)K$. The dominant decay channels of $D_s(3^3P_2)$ are: $D^*K^*(1410)$, $D_2^*(2460)K^*$, $DK^*(1410)$ and $D(2600)K$.

$D_{sJ}^+(2632)$ has a mass comparable to those of predicted $D_s(2^1S_0)$, $D_s(2^3S_1)$ and $D_s(1^3D_1)$. However, $D_{sJ}^+(2632)$ has an exotic ratio $\frac{\Gamma(D^0K^+)}{\Gamma(D_s^+\eta)} = 0.14 \pm 0.06$, which is much smaller than the predicted ones in these possibilities. In short, $D_{sJ}^+(2632)$ cannot be a conventional charmed strange $c\bar{s}$ meson, which keeps rooms for $D_s(2650)$, $D_{s1}^*(2700)$ and $D_{s1}^*(2860)$.

Acknowledgments

This work is supported by National Natural Science Foundation of China under the grants: 11475111 and 11075102. It is also supported by the Innovation Program of Shanghai Municipal Education Commission under the grant No. 13ZZ066.

[1] C. Patrignani *et al.* (Particle Data Group), Chin. Phys. C, **40**, 100001 (2016).
[2] S. Godfrey and N. Isgur, Phys. Rev. D **32**, 189 (1985).
[3] P.del Amo Sanchez *et al.* (BABAR Collaboration), Phys. Rev. D 82, 111101 (2010).
[4] R. Aaij *et al.* (LHCb collaboration), J. High Energy Phys. 09, 145 (2013).
[5] R. Aaij *et al.* (LHCb collaboration), Phys. Rev. D **91**, 092002 (2015).
[6] R. Aaij *et al.* (LHCb collaboration), Phys. Rev. D **92**, 032002 (2015).
[7] J. Brodzicka, *et al.* (Belle Collaboration), Phys. Rev. Lett. 100, 092001 (2008).
[8] B. Aubert, *et al.* (BaBar Collaboration), Phys. Rev. D **80**, 092003 (2009).
[9] R. Aaij *et al.* (LHCb collaboration), Phys. Rev. D **90**, 072003 (2014).
[10] R. Aaij et al. (LHCb collaboration), Phys. Rev. Lett. **113**, 162001 (2014).
[11] P. Colangelo, F. De Fazio, F. Giannuzzi and S. Nicotri, Phys. Rev. D **86**, 054024 (2012).
[12] S. Godfrey and K. Moats, Phys Rev. D **93**, 034035 (2016).
[13] Zhi-Feng Sun, Jie-Sheng Yu, Xiang Liu and Takayuki Matsuki, Phys. Rev. D **82**, 111501 (R) (2010).
[14] Xian-Hui Zhong, Phys. Rev. D **82**, 114014 (2010).
[15] Zhi-Gang Wang, Phys. Rev. D **83**, 014009 (2011).
[16] Bing Chen, Ling Yuan and Ailin Zhang, Phys. Rev. D **83**, 114025 (2011).
[17] Bing Chen, Xiang Liu and Ailin Zhang, Phys. Rev. D **92**, 034005 (2015).
[18] Jing Ge, Dan-Dan Ye and Ailin Zhang, Eur. Phys. J. C **75**, 178 (2015).
[19] Ze Zhao, Yu Tian and Ailin Zhang, Phys. Rev. D **94**, 114035 (2016).
[20] Hua-Xing Chen, Wei Chen, Xiang Liu, Yan-Rui Liu and Shi-Lin Zhu, arXiv:1609.08928.
[21] A.V. Evdokimov *et al.* (SELEX Collaboration), Phys. Rev. Lett, **93**, 242001 (2004).
[22] B. Aubert *et al.* (BABAR Collaboration), arXiv: hep-ex/0408087.
[23] Kuang-Ta Chao, Phys. Lett. B **599**, 43 (2004).

TABLE V: Hadronic decay widths of $D_s(3^1P_1)$ (MeV).

Channels	Width	Channels	Width
D^0K^{*+}	3.9	$D_{s1}(2460)\eta$	0.0
D^+K^{*0}	3.7	$D_1(2420)^0K^{*+}$	7.9
$D_s\phi$	0.1	$D_1(2420)^+K^{*0}$	8.1
$D^0K^*(1410)^+$	13.0	$D_1(2430)^0K^{*+}$	9.3
$D^+K^*(1410)^0$	12.4	$D_1(2430)^+K^{*0}$	9.3
$D^{*0}K^+$	7.7	$D^{*0}K_1(1270)^+$	8.1
$D^{*+}K^0$	7.6	$D^{*+}K_1(1270)^0$	8.4
$D_s^*\eta$	0.2	$D^{*0}K_1(1400)^+$	2.6
$D^{*0}K^{*+}$	2.7	$D^{*+}K_1(1400)^0$	1.8
$D^{*+}K^{*0}$	2.4	$D^0K_2^*(1430)^+$	3.1
$D_s^*\phi$	2.0	$D^+K_2^*(1430)^0$	3.2
$D^{*0}K^*(1410)^+$	15.9	$D_2^*(2460)^0K^+$	14.6
$D^{*+}K^*(1410)^0$	4.5	$D_2^*(2460)^+K^0$	14.4
$D_0^*(2400)^0K^+$	2.5	$D_{s2}^*(2573)\eta$	3.9
$D_0^*(2400)^+K^0$	1.3	$D_2^*(2460)^0K^{*+}$	13.0
$D_{s0}^*(2317)\eta$	0.2	$D_2^*(2460)^+K^{*0}$	12.4
$D^0K_0^*(1430)^+$	1.4	$D^*(2600)^0K^+$	11.9
$D^+K_0^*(1430)^0$	1.6	$D^*(2600)^+K^0$	12.0
$D_0^*(2400)^0K^{*+}$	0.0	$D(2750)^0K^+$	0.0
$D_0^*(2400)^+K^{*0}$	0.0	$D(2750)^+K^0$	0.0
$D_{s0}^*(2317)\phi$	0.0	$D_1^*(2760)^0K^+$	0.7
$D^0K_1(1270)^+$	0.0	$D_1^*(2760)^+K^0$	0.7
$D^+K_1(1270)^0$	0.0	$D_3^*(2760)^0K^+$	14.4
$D^0K_1(1400)^+$	0.0	$D_3^*(2760)^+K^0$	13.1
$D^+K_1(1400)^0$	0.0	$D_{s1}^*(2860)\eta$	0.5
$D_1(2420)^0K^+$	0.0	$D_{s3}^*(2860)\eta$	0.0
$D_1(2420)^+K^0$	0.0	$D_{s1}^*(2700)\eta$	3.5
$D_{s1}(2536)\eta$	0.0	Γ_{total}	260.4
$D_1(2430)^0K^+$	0.0		
$D_1(2430)^+K^0$	0.4		

[24] T. Barnes, F.E. Close, J.J. Dudek, S. Godfrey, and E.S. Swanson, Phys. Lett. B **600**, 223 (2004).

[25] E.van Beveren and George Rupp, Phys. Rev. Lett. **93**, 202001 (2004).

[26] Ailin Zhang, Phys. Rev. D **72**, 017902 (2005).

[27] Chao-Hsi Chang, C. S. Kim and Guo-Li Wang, Phys. Lett. B **623**, 218 (2005).

[28] Bing Chen, Deng-Xia Wang and Ailin Zhang, Phys. Rev. D **80**, 071502 (2009).

[29] Ming-Qiu Huang and Dao-Wei Wang, Phys. Rev. D **71**, 114015 (2005).

[30] Y.-R. Liu, Shi-Lin Zhu, Y.-B. Dai and C. Liu, Phys. Rev. D **70**, 094009 (2004).

[31] L. Maiani, F. Piccinini, A. D. Polosa and V. Riquer, Phys. Rev. D **70**, 054009 (2004).

[32] Yu-Qi Chen and Xue-Qian Li, Phys. Rev. Lett **93**, 232001 (2004).

[33] E.S. Swanson, Phys. Rept. **429**, 243 (2006).

[34] L. Micu, Nucl. Phys. B **10**, 521 (1969).

[35] A. Le Yaouanc, L. Oliver, O. Pène and J.C. Raynal, Phys. Rev. D **8**, 2223 (1973); **9**, 1415 (1974); **11**, 1272 (1975).

[36] A. Le Yaouanc, L. Oliver, O. Pène and J.C. Raynal, Phys. Lett. B **71**, 397 (1977); **72**, 57 (1977).

[37] A. Le Yaouanc, L. Oliver, O. Pène and J.C. Raynal, *Hadron Transitions in the Quark Model* (Gordon and Breach Science Publishers, New York, 1987).

[38] S. Godfrey and K. Moats, Phys Rev. D **90**, 117501 (2014).

[39] Ze Zhao, Dan-Dan Ye and Ailin Zhang, Phys. Rev. D **94**, 114020 (2016).

TABLE VI: Hadronic decay widths of $D_s(3^3P_1)$ (MeV).

Channels	Width	Channels	Width
D^0K^{*+}	4.1	$D_{s1}(2460)\eta$	0.0
D^+K^{*0}	3.8	$D_1(2420)^0K^{*+}$	4.5
$D_s\phi$	0.2	$D_1(2420)^+K^{*0}$	4.6
$D^0K^*(1410)^+$	7.2	$D_1(2430)^0K^{*+}$	11.9
$D^+K^*(1410)^0$	6.6	$D_1(2430)^+K^{*0}$	12.0
$D^{*0}K^+$	8.5	$D^{*0}K_1(1270)^+$	7.6
$D^{*+}K^0$	8.3	$D^{*+}K_1(1270)^0$	8.0
$D_s^*\eta$	0.1	$D^{*0}K_1(1400)^+$	2.3
$D^{*0}K^{*+}$	3.7	$D^{*+}K_1(1400)^0$	1.7
$D^{*+}K^{*0}$	3.3	$D^0K_2^*(1430)^+$	2.5
$D_s^*\phi$	2.0	$D^+K_2^*(1430)^0$	2.8
$D^{*0}K^*(1410)^+$	0.0	$D_2^*(2460)^0K^+$	9.1
$D^{*+}K^*(1410)^0$	0.0	$D_2^*(2460)^+K^0$	9.0
$D_0^*(2400)^0K^+$	0.0	$D_{s2}^*(2573)\eta$	2.6
$D_0^*(2400)^+K^0$	0.0	$D_2^*(2460)^0K^{*+}$	3.6
$D_{s0}^*(2317)\eta$	0.0	$D_2^*(2460)^+K^{*0}$	3.2
$D^0K_0^*(1430)^+$	0.0	$D^*(2600)^0K^+$	9.8
$D^+K_0^*(1430)^0$	0.0	$D^*(2600)^+K^0$	9.3
$D_0^*(2400)^0K^{*+}$	0.3	$D(2750)^0K^+$	6.7
$D_0^*(2400)^+K^{*0}$	1.7	$D(2750)^+K^0$	6.5
$D_{s0}^*(2317)\phi$	1.8	$D_1^*(2760)^0K^+$	0.1
$D^0K_1(1270)^+$	0.0	$D_1^*(2760)^+K^0$	0.1
$D^+K_1(1270)^0$	0.0	$D_3^*(2760)^0K^+$	12.0
$D^0K_1(1400)^+$	1.2	$D_3^*(2760)^+K^0$	11.0
$D^+K_1(1400)^0$	1.4	$D_{s1}^*(2860)\eta$	0.3
$D_1(2420)^0K^+$	1.7	$D_{s3}^*(2860)\eta$	0.0
$D_1(2420)^+K^0$	1.6	$D_{s1}^*(2700)\eta$	2.3
$D_{s1}(2536)\eta$	0.2	Γ_{total}	204.8
$D_1(2430)^0K^+$	1.8		
$D_1(2430)^+K^0$	1.8		

TABLE VII: Hadronic decay widths of $D_s(3^3P_2)$ (MeV).

Channels	Width	Channels	Width
D^0K^+	1.0	$D^{*0}K_0^*(1430)^+$	0.3
D^+K^0	1.1	$D^{*+}K_0^*(1430)^0$	0.2
$D_s\eta$	0.3	$D_1(2420)^0K^{*+}$	4.1
$D(2550)^0K^+$	2.6	$D_1(2420)^+K^{*0}$	4.3
$D(2550)^+K^0$	2.7	$D_1(2430)^0K^{*+}$	3.1
$D^0K(1460)^+$	5.3	$D_1(2430)^+K^{*0}$	3.0
$D^+K(1460)^0$	5.0	$D^{*0}K_1(1270)^+$	3.1
D^0K^{*+}	2.3	$D^{*+}K_1(1270)^0$	3.2
D^+K^{*0}	2.2	$D^{*0}K_1(1400)^+$	1.8
$D_s\phi$	0.0	$D^{*+}K_1(1400)^0$	1.6
$D^0K^*(1410)^+$	11.1	$D^0K_2^*(1430)^+$	1.1
$D^+K^*(1410)^0$	10.9	$D^+K_2^*(1430)^0$	1.0
$D(2550)^0K^{*+}$	0.0	$D_2^*(2460)^0K^+$	6.5
$D(2550)^+K^{*0}$	0.0	$D_2^*(2460)^+K^0$	6.5
$D^{*0}K^+$	3.8	$D_{s2}^*(2573)\eta$	1.8
$D^{*+}K^0$	3.8	$D^{*0}K_2^*(1430)^+$	1.6
$D_s^*\eta$	0.3	$D_2^*(2460)^0K^{*+}$	25.0
$D^{*0}K^{*+}$	4.8	$D_2^*(2460)^+K^{*0}$	24.9
$D^{*+}K^{*0}$	4.4	$D^*(2600)^0K^+$	7.1
$D_s^*\phi$	1.7	$D^*(2600)^+K^0$	7.6
$D^{*0}K^*(1410)^+$	45.2	$D(2750)^0K^+$	2.0
$D^{*+}K^*(1410)^0$	44.5	$D(2750)^+K^0$	2.0
$D_0^*(2400)^0K^{*+}$	0.1	$D(2750)^0K^+$	0.4
$D_0^*(2400)^+K^{*0}$	1.1	$D(2750)^+K^0$	0.4
$D_{s0}^*(2317)\phi$	1.3	$D_1^*(2760)^0K^+$	0.2
$D^0K_1(1270)^+$	2.9	$D_1^*(2760)^+K^0$	0.2
$D^+K_1(1270)^0$	2.7	$D_3^*(2760)^0K^+$	4.7
$D^0K_1(1400)^+$	1.6	$D_3^*(2760)^+K^0$	4.3
$D^+K_1(1400)^0$	1.5	$D_s(2650)\eta$	2.4
$D_1(2420)^0K^+$	1.9	$D_{s1}^*(2860)\eta$	0.0
$D_1(2420)^+K^0$	1.8	$D_{s3}^*(2860)\eta$	0.0
$D_{s1}(2536)\eta$	0.1	$D_{s1}^*(2700)\eta$	3.0
$D_1(2430)^0K^+$	4.9	Γ_{total}	307.9
$D_1(2430)^+K^0$	4.9		
$D_{s1}(2460)\eta$	2.7		

TABLE VIII: Partial and total widths of $D_{sJ}^+(2632)$ (MeV) and $\Gamma(D^0K^+)/\Gamma(D_s^+\eta)$ in different assignments.

Assignment	D^0K^+	D^+K^0	$D_s\eta$	$D^{*0}K^+$	$D^{*+}K^0$	Γ_{total}	$\frac{\Gamma(D^0K^+)}{\Gamma(D_s^+\eta)}$
$D_{sJ}^+(2632)(2^3S_1)$	17.8	17.5	3.5	22.9	21.6	83.2	5.02
$D_{sJ}^+(2632)(1^3D_1)$	34.2	33.4	4.3	8.8	8.3	88.9	7.90