

Lepton Flavour Violation in Left-Right Theory

Pavel Fileviez Pérez¹, Clara Murgui²

¹*CERCA, Physics Department, Case Western Reserve University,*

Rockefeller Bldg. 2076 Adelbert Rd. Cleveland, OH 44106, USA

²*Departamento de Física Teórica, IFIC, Universitat de Valencia-CSIC, E-46071, Valencia, Spain*

We investigate the predictions for lepton flavour number violating processes in the context of a simple left-right symmetric theory. In this context neutrinos are Majorana fermions and their masses are generated at the quantum level through the Zee mechanism using the simplest Higgs sector. We show that the right handed neutrinos are generically light and can give rise to large lepton flavour violating contributions to rare processes. We discuss the correlation between the collider constraints and the predictions for lepton flavour violating processes. We find that using the predictions for $\mu \rightarrow e\gamma$ and $\mu \rightarrow e$ conversion together with the collider signatures one could test this theory in the near future.

I. INTRODUCTION

The Large Hadron Collider (LHC) has discovered the last missing piece of the Standard Model (SM) of particle physics. The discovery of the Brout-Englert-Higgs boson was crucial to establish the SM as one of the most important theories of nature. Today, we believe that the SM should be an effective theory to explain most of the current experimental results. However, it is well-known that one cannot explain in this context, for example, the hierarchy of the fermion masses, the origin of neutrino masses, the origin of Parity and CP violation, the nature of dark matter and the baryon-asymmetry in the Universe.

There are many ideas for physics beyond the Standard Model which can help us to define a new theory to describe the new energy scale, TeV scale, which is currently explored by the Large Hadron Collider. In the context of left-right symmetric theories [1–5], proposed by J. Pati, A. Salam, R. Mohapatra and G. Senjanović, one can explain some of the open issues of the SM. In this context, the spontaneous breaking of Parity is naturally explained and one can understand why at the low scale the weak interactions are $V - A$ interactions. These theories predict the existence of right-handed neutrinos in nature which play a crucial role to generate neutrino masses. In the context of left-right symmetric theories, neutrinos can be Dirac fermions [3] or Majorana fermions [5]. In the Majorana case one can make use of the see-saw mechanism [5, 6] to understand the smallness of the neutrino masses. These theories can give rise to many interesting signatures at colliders and low energy experiments, see for example Refs. [7–28] for different phenomenological studies.

Recently, we have proposed a simple left-right symmetric theory [29] where the Majorana neutrino masses are generated through the Zee-mechanism [30]. In this context the charged fermion masses are generated at tree level as in the SM, while the neutrino masses are generated at one-loop level. This theory has the simplest Higgs sector needed to generate Majorana masses for neutrinos and to realize the spontaneous breaking of the local left-right symmetry obtaining the SM at the weak scale. In Ref. [29] we have proposed this new theory and investigated the main collider signatures which can help us to test the theory at the Large Hadron Collider.

In this article we investigate in details the properties of the right-handed neutrinos in the theory proposed in Ref. [29] and the predictions for lepton flavor violating (LFV) processes such as the rare decays $e_i \rightarrow e_j \gamma$ and $\mu \rightarrow e$ conversion. The current experimental bounds from the LFV experiments provide non-trivial bounds on lepton flavour violating interactions present in different theories (see Ref. [31] for a review of LFV experiments). The next generation of LFV experiments will set very strong bounds on the branching fractions for these rare decays and we investigate the impact of these results in our model, in which one has several contributions to LFV processes: the interactions between the W_L^\pm or W_R^\pm with the charged leptons and the neutrinos, and the L-violating Higgs interactions. We show that, in this context, one can have very large contributions to LFV processes in agreement with all experimental constraints. Together with the collider signatures studied in Ref. [29] these results can be used to test this theory in current and future experiments.

II. SIMPLE LEFT-RIGHT SYMMETRIC THEORY

Recently, we have proposed in Ref. [29] a simple left-right symmetric theory based on the gauge symmetry

$$G_{LR} = SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$$

where the Majorana neutrino masses are generated at the quantum level. As in any left-right symmetric theory the matter fields live in the following representations

$$Q_L = \begin{pmatrix} u_L \\ d_L \end{pmatrix} \sim (3, 2, 1, 1/3), \quad Q_R = \begin{pmatrix} u_R \\ d_R \end{pmatrix} \sim (3, 1, 2, 1/3),$$

$$\ell_L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix} \sim (1, 2, 1, -1), \quad \text{and} \quad \ell_R = \begin{pmatrix} \nu_R \\ e_R \end{pmatrix} \sim (1, 1, 2, -1).$$

and the Higgs sector is composed of four Higgses: a bi-doublet needed to generate charged fermion masses, a charged singlet and two doublets required to break the left-right symmetry and generate Majorana neutrino masses in a minimal way,

$$\Phi = \begin{pmatrix} \phi_1^0 & \phi_2^+ \\ \phi_1^- & \phi_2^0 \end{pmatrix} \sim (1, 2, 2, 0), \quad \delta^+ \sim (1, 1, 1, 2),$$

$$H_L = \begin{pmatrix} h_L^+ \\ h_L^0 \end{pmatrix} \sim (1, 2, 1, 1) \quad \text{and} \quad H_R = \begin{pmatrix} h_R^+ \\ h_R^0 \end{pmatrix} \sim (1, 1, 2, 1).$$

A. Charged Fermion Masses

As in any left-right symmetric theory the charged fermions acquire mass at tree level once the Higgs bi-doublet gets a vacuum expectation value. Using the interactions

$$- \mathcal{L}_{LR}^D = \overline{Q}_L (Y_1 \Phi + Y_2 \tilde{\Phi}) Q_R + \bar{\ell}_L (Y_3 \Phi + Y_4 \tilde{\Phi}) \ell_R + \text{h.c.}, \quad (1)$$

where $\tilde{\Phi} = \sigma_2 \Phi^* \sigma_2$, one finds the following charged fermion masses after electroweak symmetry breaking

$$M_U = Y_1 v_1 + Y_2 v_2^*, \quad (2)$$

$$M_D = Y_1 v_2 + Y_2 v_1^*, \quad (3)$$

$$M_E = Y_3 v_2 + Y_4 v_1^*. \quad (4)$$

Here, v_1 and v_2 are the vacuum expectation values for the fields ϕ_1^0 and ϕ_2^0 , respectively. Notice that the Y_1 and Y_2 can be written as linear combinations of the mass matrices for the up and down quarks, while one has more freedom in the expression for charged lepton masses due to the presence of two Yukawa couplings.

B. Neutrino Masses

In the left-right theory with only three Higgses, Φ , H_L and H_R , the total lepton number is conserved after symmetry breaking and neutrinos are Dirac massive fermions with mass given by

$$M_\nu^D = Y_3 v_1 + Y_4 v_2^*. \quad (5)$$




FIG. 1: Neutrino mass generation at the quantum level.

Notice that using the freedom in Eqs.(4) and (5) one can have a consistent scenario for Dirac neutrinos in this context [3]. However, the neutrino masses are very small and they could be Majorana fermions. In the theory proposed in Ref. [29] Majorana neutrino masses are generated at one-loop level using the following interactions

$$-\mathcal{L}_{LR}^M = \lambda_L \ell_L \ell_L \delta^+ + \lambda_R \ell_R \ell_R \delta^+ + \lambda_1 H_L^T i \sigma_2 \Phi H_R \delta^- + \lambda_2 H_L^T i \sigma_2 \tilde{\Phi} H_R \delta^- + \text{h.c.} \quad (6)$$

See Fig.1 for the one-loop contribution to neutrino masses in the unbroken phase. Notice that both the left-handed and right-handed neutrinos acquire masses at one-loop level and their masses are proportional to the vacuum expectation values of h_L^0 and h_R^0 , i.e. $v_L/\sqrt{2}$ and $v_R/\sqrt{2}$. The neutrino mass matrix in the basis $(\nu_L, (\nu^C)_L)$ is given by

$$\mathcal{M}_\nu = \begin{pmatrix} M_\nu^L & M_\nu^D \\ (M_\nu^D)^T & M_\nu^R \end{pmatrix}, \quad (7)$$

where M_ν^L and M_ν^R are generated at one-loop level while M_ν^D is generated at tree level. The explicit forms of M_ν^L and M_ν^R are given by

$$(M_\nu^L)^{\alpha\gamma} = \frac{1}{4\pi^2} \lambda_L^{\alpha\beta} m_{e\beta} \sum_i \text{Log} \left(\frac{M_{h_i}^2}{m_{e\beta}^2} \right) V_{5i} \left[(Y_3^\dagger)^{\beta\gamma} V_{2i}^* - (Y_4^\dagger)^{\beta\gamma} V_{1i}^* \right] + \alpha \leftrightarrow \gamma, \quad (8)$$

$$(M_\nu^R)^{\alpha\gamma} = \frac{1}{4\pi^2} \lambda_R^{\alpha\beta} m_{e\beta} \sum_i \text{Log} \left(\frac{M_{h_i}^2}{m_{e\beta}^2} \right) V_{5i} \left[(Y_3)^{\beta\gamma} V_{1i}^* - (Y_4)^{\beta\gamma} V_{2i}^* \right] + \alpha \leftrightarrow \gamma. \quad (9)$$

Here V_{ij} defines the mixing between the charged Higgses in the theory and M_{h_i} their physical masses (see Ref. [29] for more details). In our notation the neutrino mass matrix is diagonalized by the following matrix

$$\begin{pmatrix} \nu_L \\ (\nu^C)_L \end{pmatrix} \rightarrow \begin{pmatrix} V_\nu & A \\ B & V_N \end{pmatrix} \begin{pmatrix} \nu_L \\ N_L \end{pmatrix} = \begin{pmatrix} V_\nu \nu_L + A N_L \\ B \nu_L + V_N N_L \end{pmatrix}, \quad (10)$$

which is useful to obtain all physical interactions. Henceforth we are neglecting the mixing between the left-handed and right-handed neutrinos because it is very small. In general the vevs as well as the Yukawas are free parameters and one cannot predict anything about the magnitude of the masses. However, for the theory to be consistent one needs to assume (see Ref. [29]) that $v_2 \ll v_1$ and $Y_3 \ll Y_4$. In this limit, the mass matrix for charged leptons can be approximated by $M_E \approx Y_4 v_1^*$ and M_ν^R as

$$(M_\nu^R)^{\alpha\beta} = \frac{1}{4\pi^2} \lambda_R^{\alpha\beta} \frac{m_{e_\beta}^2}{v_1^*} \sum_i \text{Log} \left(\frac{M_{h_i}^2}{m_{e_\beta}^2} \right) V_{5i} V_{2i}^* + \alpha \leftrightarrow \beta. \quad (11)$$

From the above relation one can extract predictions about the hierarchy of the sterile neutrino masses. Notice that Eq.(11) is traceless due to the product of the antisymmetric Yukawa $\lambda_R^{\alpha\beta}$ with the symmetric mass matrix of the charged leptons, which is assumed without loss of generality to be diagonal. On the other hand, the mass matrix in Eq. (11) is sensible to the difference between pairs of charged lepton masses squared. Since the difference between the muon mass and the other charged lepton masses is one order of magnitude smaller than the rest of differences, the muon neutrino N_μ is predicted to be at least two orders of magnitude lighter than the electron neutrino N_e and the tauon neutrino N_τ . Bringing together both statements and taking into account the invariance of the trace, the following conclusion about the hierarchy of the masses can be drawn: the model predicts that the muon sterile neutrino is much lighter than the others, which therefore have almost degenerated masses.

In order to study qualitatively the order of magnitude of the sterile neutrino masses, one could assume as a good approximation that the product of the charged Higgses mixing matrices is of the order of one, due to the unitarity nature of the mixing matrix. Notice that unitarity further constraints the sum of the logarithm over the different five physical charged Higgses in the theory, making this term only sensible to twice the difference of the order of magnitude between the lightest and the heaviest charged Higgses. Let us call this factor Δ , which will represent the contribution of the logarithms in Eq.(11). Hence, in the limit $v_2 \ll v_1$ and $Y_3 \ll Y_4$, and assuming that $v_1 \approx v = 246$ GeV, Eq. (11) can be rewritten as

$$(M_\nu^R)^{\alpha\beta} \approx \frac{\Delta}{4\pi^2 v_1^*} \lambda_R^{\alpha\beta} (m_{e_\alpha}^2 - m_{e_\beta}^2). \quad (12)$$




FIG. 2: Predictions for the sterile neutrino masses as a function of Δ . The blue, orange and red points correspond to the electron N_e , muon N_μ , tauon N_τ sterile neutrinos, respectively. For the scan, the entries of the Yukawa λ_R have been taken randomly ranging between 0 to $2\sqrt{\pi}$. The vertical green line shows the estimated upper bound Δ_{upper} .

We can estimate a theoretical upper limit for the factor Δ by assuming the extreme case in which the lightest charged scalar lives at the electroweak scale and the heaviest one at the Plank scale. In this scenario Δ is given by, $\Delta_{\text{upper}} \sim \text{Log} \left(\frac{10^{19} \text{ GeV}}{10^2 \text{ GeV}} \right)^2 = 2 \times 17 \times \text{Log}(10)$. The relation in Eq. (12) is one of the main predictions of the model, since this correlation between the sterile neutrino masses and the Yukawa coupling λ_R constrains strongly, on one hand, the hierarchy of the sterile neutrino masses and, on the other hand, allows us to estimate an upper bound for the right-handed neutrino masses.

In Fig. 2 we show the correlation between the sterile neutrino masses M_{N_i} , given by the eigenvalues of Eq. (12) and the factor Δ . The scattered points correspond to different values for the entries of λ_R , which range randomly from $[0, 2\sqrt{\pi}]$, according to perturbativity. As we can see in Fig. 2, the model predicts very light sterile neutrinos with the theoretically predicted hierarchy. We emphasize again the relevance of the antisymmetric nature of the Yukawa λ_R matrix since one has only three free parameters, $\lambda_R^{e\mu}$, $\lambda_R^{\mu\tau}$ and $\lambda_R^{e\tau}$. Our main result here is that the right-handed neutrinos are generically light. This prediction will be very important to study the predictions for lepton flavour violating processes.

C. Charged Gauge Boson Masses

In the basis (W_L^+, W_R^+) the charged gauge boson mass matrix reads as

$$\mathcal{M}_\pm^2 = \begin{pmatrix} \frac{g_L^2}{2}(\frac{1}{2}v_L^2 + v^2) & -g_L g_R v_1 v_2 \\ -g_L g_R v_1 v_2 & \frac{g_R^2}{2}(\frac{1}{2}v_R^2 + v^2) \end{pmatrix}, \quad (13)$$

where $v^2 = v_1^2 + v_2^2$. The mass of the W_R -like charged gauge boson is $M_{W_R} \approx g_R v_R / 2$. Using the charged current interactions

$$-\mathcal{L}_{LR}^{CC} \supset \frac{g_L}{\sqrt{2}} \bar{\nu}_L W_L^+ e_L + \frac{g_R}{\sqrt{2}} \bar{\nu}_R W_R^+ e_R + \text{h.c.} , \quad (14)$$

and the definition in Eq.(10) one can study lepton flavour violation in the leptonic sector mediated by the gauge bosons. Recently, the LHC experiments have set bounds on the mass of these gauge bosons. See Ref. [32] for the lower experimental bound, $M_{W_R} > 4.1$ TeV, on the mass of the W_R -like gauge boson.

III. LEPTON FLAVOUR VIOLATING PROCESSES

In the theory proposed in Ref. [29] there are several sources of lepton flavour violation:

- The physical interactions between the W_L^\pm , the charged leptons and the neutrinos.
- The physical interactions between the W_R^\pm , the charged leptons and the neutrinos.
- In Eq.(1) we cannot simultaneously diagonalize the Yukawa couplings Y_3 and Y_4 , and the Higgs interactions violate the family lepton numbers.
- The Yukawa interactions in Eq.(6) violate the global $U(1)_{L_i}$ as well.

In this section we investigate the predictions for lepton flavour violating processes such as $e_i \rightarrow e_j \gamma$ taking into account the different sources for L_i violation. Notice that, among the different sources violating lepton flavour, there are the charged Higgses in the bi-doublet. In this work we will not focus on them since they also contribute to $\Delta F = 2$ hadronic changing neutral current effects which are very constrained and, therefore, these Higgses have to be very heavy [10]. However, in the context of the recently proposed left-right symmetric model [29], the singly charged Higgs could be relatively light and can induce large contributions to lepton flavour violating processes.

A. LFV $e_i \rightarrow e_j \gamma$ Processes

In this section we investigate the predictions for the lepton flavour violating processes in order to understand the testability of the theory proposed in Ref. [29]. The current experimental bounds on the branching ratios for the $e_i \rightarrow e_j \gamma$ processes are

$$\text{Br}(\mu \rightarrow e \gamma) < 4.2 \times 10^{-13} [33], \text{Br}(\tau \rightarrow e \gamma) < 3.3 \times 10^{-8} [34], \text{Br}(\tau \rightarrow \mu \gamma) < 4.4 \times 10^{-8} [34].$$




FIG. 3: Contributions to $\mu \rightarrow e\gamma$. Topology (a) mediated by a charged gauge boson, W_L^\pm or W_R^\pm , and topology (b) mediated by a singly charged Higgs.

As it is well known, the amplitude for the process $\mu \rightarrow e\gamma$ can be written as

$$\mathcal{A}(\mu \rightarrow e\gamma) = i\bar{u}_e(p-q)\epsilon_\nu^*\sigma^{\nu\mu}q_\mu[A_R P_R + A_L P_L]u_\mu(p), \quad (15)$$

where p^μ and q^μ are the muon and photon quadrimomenta, respectively, and the decay width reads as

$$\Gamma(\mu \rightarrow e\gamma) = \frac{m_\mu^3}{16\pi}(|A_L|^2 + |A_R|^2). \quad (16)$$

The branching ratio can be computed using the relation

$$\text{Br}(\mu \rightarrow e\gamma) \equiv \frac{\Gamma(\mu \rightarrow e\gamma)}{\Gamma(\mu \rightarrow e\nu_\mu\bar{\nu}_e) + \Gamma(\mu \rightarrow e\gamma)}, \quad (17)$$

where

$$\Gamma(\mu \rightarrow e\nu_\mu\bar{\nu}_e) = \frac{m_\mu^5 G_F^2}{192\pi^3}. \quad (18)$$

In our model there are several contributions to the coefficients A_L and A_R relevant for the decay width. In Fig. 3 we show the Feynman graphs for the different contributions mediated by the charged gauge bosons and the charged Higgses. Here we will investigate the predictions for each contribution in order to understand the testability of the theory in current and future experiments.

1. LFV induced by W_i^\pm gauge bosons

The W_L^\pm and W_R^\pm contributions to the process $\mu \rightarrow e\gamma$, neglecting the electron mass, read as

$$A_L^{W_R} \approx g_R^2 \frac{e m_\mu}{64\pi^2 M_{W_R}^2} \sum_i (V_N)_{ei} (V_N^*)_{\mu i} F\left(\frac{m_{N_i}^2}{m_{W_R}^2}\right), \quad (19)$$

$$A_R^{W_L} \approx g_L^2 \frac{e m_\mu}{64\pi^2 M_{W_L}^2} \sum_i (V_\nu)_{ei} (V_\nu^*)_{\mu i} F\left(\frac{m_{\nu_i}^2}{m_{W_L}^2}\right), \quad (20)$$

where we have neglected the mixing between the neutrinos. The loop scalar function $F(x)$ is defined as

$$F(x) = \frac{1}{6(1-x)^4} (10 - 43x + 78x^2 - 49x^3 + 18x^3 \text{Log}(x) + 4x^4), \quad (21)$$

which has the following limits,

$$F(x)_{x \rightarrow \infty} \sim \frac{2}{3} + 3 \frac{\text{Log}(x)}{x}, \quad F(x)_{x \rightarrow 0} \sim \frac{5}{3} - \frac{1}{2}x, \quad F(x)_{x \rightarrow 1} \sim \frac{17}{12} + \frac{3}{20}(1-x). \quad (22)$$

Notice that when W_L^\pm is the mediator of the process, the Standard Model neutrinos are the ones contributing into the amplitude. In the second contribution, the W_R^\pm and the sterile neutrinos are inside the loop. In the case of W_L^\pm , $x \equiv m_{\nu_i}^2/M_{W_L}^2 \rightarrow 0$ due to the smallness of the Standard Model neutrino masses and the function $F(x \rightarrow 0)$ characterizing the loop behaves almost as a constant. Hence, $A_R^{W_L} \sim 0$ due to the unitarity of V_ν , i.e. the so-called GIM suppression. For the W_R gauge boson, however, the GIM suppression can be avoided if the sterile neutrinos are heavy enough to spoil the suppression coming from the unitarity relations. Each of the limits of $F(x)$ leads to a different amplitude, shown in Table I. The largest possible

TABLE I: Different limits for $A_L^{W_R}$ amplitud.

Limit	$A_L^{W_R}$
$m_{N_i} \ll M_{W_R}$	$-g_R^2 \frac{e}{128\pi^2} \frac{m_\mu}{M_{W_R}^4} \sum_i (V_N)_{ei} (V_N^*)_{\mu i} m_{N_i}^2$
$m_{N_i} \gg M_{W_R}$	$g_R^2 m_\mu \frac{3e}{64\pi^2} \sum_i \text{Log}\left(\frac{m_{N_i}^2}{M_{W_R}^2}\right) (V_N)_{ie} (V_N^*)_{i\mu} \frac{1}{m_{N_i}^2}$
$m_{N_i} \sim M_{W_R}$	$g_R^2 \frac{3}{1280\pi^2} \frac{m_\mu}{M_{W_R}^2} \sum_i (V_N)_{ei} (V_N^*)_{\mu i} \frac{M_{W_R}^2 - m_{N_i}^2}{M_{W_R}^2}$

contribution corresponds to the case where both, the masses of the sterile neutrinos and the mass of the charged gauge boson, are of the same order of magnitude, i.e. $m_{N_i} \sim M_{W_R}$. We illustrate this scenario in Fig. 4 by showing the prediction on the branching ratio $e_i \rightarrow e_j \gamma$ (purple points) as a function of M_{W_R} for different values of the sterile neutrino masses of the same order of magnitude as M_{W_R} . In the context of our LR-model in Ref. [29], neutrinos are predicted to be very light; $m_{N_i}/M_{W_R} \leq 10^{-4}$. Therefore, as we also show in Fig. 4, blue points, the GIM suppression occurs and the predictions for the branching ratio are far away from the current and even future experimental reach. In order to complete our discussion we show the predictions for $g = 2$ which are always very small.




FIG. 4: Predictions on the branching ratio of $e_i \rightarrow e_j \gamma$ and $g - 2$ anomalous momentum of the muon mediated by a right handed charged gauge boson as a function of its mass M_{W_R} . The blue points represent the predictions on the branching ratio in the context of our model, whereas the purple points correspond to the most optimistic case to observe LNV in which the right handed neutrinos and the W_R boson masses are of the same order of magnitude. For all points, $\lambda_R \in [0, 2\sqrt{\pi}]$ (perturbative scenario) and the factor $\Delta \in [0, \Delta_{\text{upper}}]$. The red lines represent the experimental lower bounds on the different processes, $\text{Br}(\mu \rightarrow e\gamma) < 4.2 \times 10^{-13}$ [33], $\text{Br}(\tau \rightarrow e\gamma) < 3.3 \times 10^{-8}$ [34], $\text{Br}(\tau \rightarrow \mu\gamma) < 4.4 \times 10^{-8}$ [34] and $\Delta a_\mu < 287 \times 10^{-11}$ [35]. The orange dashed lines represent the projected limits 6×10^{-14} for $\mu \rightarrow e\gamma$ [36], 3×10^{-9} for $\mu \rightarrow e\gamma$ [37], and $\sim 10^{-9}$ for $\tau \rightarrow e\gamma$ [37]. Finally, the green vertical line shows the lower bound on the W_R gauge boson [32], which only applies to the blue points. Here, left-right symmetry has been assumed in the gauge sector, i.e. $g_L = g_R$.

2. LFV mediated by δ^\pm Charged Higgses

The charged Higgs, δ^\pm , can be light and induce large contributions to the lepton flavour violating processes. The amplitude for the $\mu \rightarrow e\gamma$ process can be written as

$$A_L^{\delta^+} = \frac{e}{4\pi^2} \frac{m_\mu}{m_{\delta^+}^2} \sum_i \sum_{c,d} (\lambda_R^*)^{ce} \lambda_R^{d\mu} V_N^{ci} (V_N^*)^{di} G\left(\frac{m_{N_i}^2}{m_{\delta^+}^2}\right), \quad (23)$$

$$A_R^{\delta^+} = \frac{e}{4\pi^2} \frac{m_\mu}{m_{\delta^+}^2} \sum_i \sum_{c,d} (\lambda_L^*)^{ce} \lambda_L^{d\mu} (V_\nu^*)^{ci} V_\nu^{di} G\left(\frac{m_{\nu_i}^2}{m_{\delta^+}^2}\right), \quad (24)$$

where, as in previous discussion, fermion masses in the loop have been neglected, and the scalar function $G(x)$ is defined as

$$G(x) = \frac{1 - 6x + 3x^2 + 2x^3 - 6x^2 \text{Log}(x)}{12(1 - x)^4}. \quad (25)$$

Notice that, as one can see in Fig. 3 (b) as well as in the above equation, the N_i and δ^+ decouple in the limit where their masses are heavy so that the amplitude contributing to LFV is suppressed. Therefore, only when both masses are light, LFV processes can have a large effect. As we have already commented, the crucial point here is that λ_R has flavour indices and therefore breaks the unitarity relations. However, LNV processes not protected by the GIM suppression or any given internal symmetry are dangerous in the sense that they could easily be in conflict with the strong current experimental bounds. It is remarkable that, despite of losing the GIM protection, the consistency of the left-right symmetric model with a charged scalar introduced in Ref. [29] is ensured due to the light sterile neutrinos that this theory predicts. In this article we neglect the mixing between δ^+ and the other charged Higgses because it is always small. The term in the scalar potential which define this mixing is $H_L^T i\sigma_2 \Phi H_R \delta^-$, and since the charged Higgses in the bidoublet are very heavy, the mixing angle is always very small.

In Fig. 5 we show the predictions on the branching ratio for $e_i \rightarrow e_j \gamma$ mediated by δ^+ as a function of the mass of the charged scalar. The points plotted range from $\lambda_R \in [0, 0.1]$, i.e. the Yukawa coupling has been assumed to be perturbative, and the Δ factor from $\Delta \in [0, \Delta_{\text{upper}}]$. Notice that, due to the relation shown in Eq. (12), the sterile neutrino masses as well as the rotation matrix V_N can be extracted from the coupling λ_R and the charged fermion masses. We neglected the contributions mediated by the charged gauge bosons since they are very small. In Fig. 5 one can see that the limits on the branching ratio for the process $\mu \rightarrow e \gamma$ impose non-trivial bounds on the mass of the charged Higgs generating neutrino masses at the one-loop level. However, generically the charged Higgs δ^\pm can be light enough to avoid the LFV bounds and be produced at the Large Hadron Collider with a large cross section. In Fig. 5 we also show the numerical results for $g = 2$, which are very small, in order to complete our discussion.




FIG. 5: Predictions on the branching ratio of the process $e_i \rightarrow e_j \gamma$ mediated by a singly charged scalar as a function of the M_{δ^+} . Here, the Yukawa $\lambda_R \in [0, 0.1]$ and the factor $\Delta \in [0, \Delta_{\text{upper}}]$. The red lines represent the experimental lower bounds on the different processes, $\text{Br}(\mu \rightarrow e \gamma) < 4.2 \times 10^{-13}$ [33], $\text{Br}(\tau \rightarrow e \gamma) < 3.3 \times 10^{-8}$ [34], $\text{Br}(\tau \rightarrow \mu \gamma) < 4.4 \times 10^{-8}$ [34] and $\Delta a_\mu < 287 \times 10^{-11}$ [35]. The orange dashed lines represent the projected limits 6×10^{-14} for $\mu \rightarrow e \gamma$ [36], 3×10^{-9} for $\mu \rightarrow e \gamma$ [37], and $\sim 10^{-9}$ for $\tau \rightarrow e \gamma$ [37].

B. $\mu \rightarrow e$ conversion

The process of $\mu \rightarrow e$ conversion was first studied by Weinberg and Feinberg [38]. See also Refs. [39–43]. The experimental lower bounds for this process are stronger than for the $\mu \rightarrow e \gamma$ process

$$R_{\mu \rightarrow e}^{\text{Ti}} < 4.3 \times 10^{-12} [44], R_{\mu \rightarrow e}^{\text{Au}} < 7 \times 10^{-13} [45], R_{\mu \rightarrow e}^{\text{Pb}} < 4.6 \times 10^{-11} [46],$$

and, moreover, these constraints are expected to be improved by several orders of magnitude in a near future according to some future experiments such as DeeMe at J-PARC [47], with an expected sensitivity of 10^{-14} , Mu2e at Fermilab [49], with 6×10^{-17} , and COMET at J-PARC [48], with 10^{-16} . These competitive lower bounds make the process $\mu - e$ conversion very attractive to constrain models predicting lepton flavour violating interactions.

From an effective theory point of view, the effective interactions contributing to this process in the




FIG. 6: Diagrammatic contributions to μ to e conversion. The black spot symbolizes the effective vertex.

context of our left-right symmetric model with a singly charged Higgs can be written as

$$-\mathcal{L}_{\text{eff}}^{(q)} = C_{DR} m_\mu \bar{e} \sigma^{\rho\nu} P_L \mu F_{\rho\nu} + \sum_q C_{VR}^{(q)} \bar{e} \gamma^\rho P_R \mu (\bar{q} \gamma_\rho q) + \text{h.c.} \quad (26)$$

Among the gauge boson contributions to the above lagrangian, shown in Fig. 6, we only consider the photonic contribution since the contribution of the massive gauge bosons is suppressed by their masses squared. As a good approximation, we assume that the transferred momentum t of the process shown in Fig. 6 is of the order of $t \sim m_\mu^2$, which would correspond to an elastic collision. The Wilson coefficients in Eq.(26) in the context of our left-right symmetric model are written explicitly in the appendix. Notice that here, unlike in the $\mu \rightarrow e\gamma$ case, the photon is off-shell. However, since $t \sim m_\mu^2$, the transferred momentum can be neglected inside the loop compared to the charged scalar/gauge bosons masses, giving the same result as the one showed in the section of $\mu \rightarrow e\gamma$. Here the contributions mediated by the charged gauge bosons have been neglected in front of the contribution of the singly charged Higgs, as we have justified before. The above lagrangian is defined at the quark level. However, we are interested in low energy processes involving nuclei. In order to perform the matching between both energy regimes we follow Refs. [41] and [42], where the matching between quark and parton level is made by introducing the following form factors

$$\bar{q} \gamma_\rho q \rightarrow f_{VN}^{(q)} \bar{\Psi}_N \gamma_\rho \Psi_N, \quad (27)$$

defined as

$$f_{Vp}^{(u)} = 2, \quad f_{Vp}^{(d)} = 1, \quad f_{Vp}^{(s)} = 0, \quad (28)$$

$$f_{Vn}^{(u)} = 1, \quad f_{Vn}^{(d)} = 2, \quad f_{Vn}^{(s)} = 0. \quad (29)$$

and thus the new effective couplings $\tilde{C}_{VR}^{(N)}$ read as

$$\tilde{C}_{VR}^{(p)} = \sum_{q=u,d,s} C_{VR}^{(q)} f_{Vp}^{(q)}, \quad (30)$$

$$\tilde{C}_{VR}^{(n)} = \sum_{q=u,d,s} C_{VR}^{(q)} f_{Vn}^{(q)}. \quad (31)$$

The muon conversion rate reads as [42]

$$\Gamma_{\text{conv}} = \frac{m_\mu^5}{4} \left| C_{DR} D + \tilde{C}_{VR}^{(p)} 4V^{(p)} + \tilde{C}_{VR}^{(n)} 4V^{(n)} \right|^2, \quad (32)$$

where the dimensionless integrals D and $V^{(N)}$ represent the overlap of electron and muon wavefunctions and depend on the nucleus involved. Their explicit expressions are given by [42]

$$D = \frac{4}{\sqrt{2}} m_\mu \int_0^\infty dr r^2 [-E(r)] (g_e^- f_\mu^- + f_e^- g_\mu^-), \quad (33)$$

$$V^{(p)} = \frac{1}{2\sqrt{2}} \int_0^\infty dr r^2 Z \rho^{(p)} (g_e^- g_\mu^- + f_e^- f_\mu^-), \quad (34)$$

$$V^{(n)} = \frac{1}{2\sqrt{2}} \int_0^\infty dr r^2 (A - Z) \rho^{(n)} (g_e^- g_\mu^- + f_e^- f_\mu^-), \quad (35)$$

where $\rho^{(N)}$ is the density of the nucleon, $E(r)$ refers to the electric field, which is obtained by integrating the Maxwell equation

$$E(r) = \frac{Ze}{r^2} \int_0^r r'^2 \rho^{(p)}(r') dr', \quad (36)$$

and the functions g_μ^- , f_μ^- , g_e^- and f_e^- correspond to the 1s muon wave functions and electron wave functions, respectively, according to the nomenclature used in Ref. [40]. The values of the dimensionless integrals D and $V^{(N)}$ along with the capture rates of the named isotopes are listed in Table II.

Isotop	D [42]	$V^{(p)}$ [42]	$V^{(n)}$ [42]	$\Gamma_{\text{capt}} (10^6 s^{-1})$ [51]
$^{27}_{13}\text{Al}$	0.360	0.0160	0.0171	0.69
$^{48}_{22}\text{Ti}$	0.0867	0.0398	0.0482	2.59
$^{197}_{79}\text{Au}$	0.178	0.0917	0.127	13.07
$^{207}_{82}\text{Pb}$	0.160	0.0828	0.119	13.45

TABLE II: For the dimensionless integrals, the average of the numerical values computed through different methodes has been taken (see Ref. [42]).




FIG. 7: Prediction on the branching ratio of the process $\mu - e$ conversion in different nuclei as a function of the mass of the singly charged Higgs. Here, the Yukawa couplings λ_R range from 0 to 0.1 and Δ from 0 to Δ_{upper} randomly. The red lines show the upper experimental bound for $\mu - e$ conversion, according to the nucleus. The dashed lines represent the projected sensitivities of the experiments DeeMe at J-PARC with 10^{-14} [47] (green line), and COMET at J-PARC with 10^{-16} [48] and Mu2e at Fermilab with 6×10^{-17} [49] (magenta line).

Finally, the branching ratio is usually expressed as the conversion rate normalized by the muon capture rate

$$B_{\mu \rightarrow e}(Z) = \frac{\Gamma_{\text{conv}}(Z, A)}{\Gamma_{\text{capt}}(Z, A)}. \quad (37)$$

In Fig. 7 we show the prediction on the $\mu \rightarrow e$ process in the context of our left-right symmetric model as a function of the charged scalar mass for the isotopes $^{27}_{13}\text{Al}$, $^{48}_{22}\text{Ti}$, $^{197}_{79}\text{Au}$ and $^{207}_{82}\text{Pb}$. The current experimental bounds are represented by the red line and with dashed lines we show the projected limits. As one can see, the current bounds constrain only a small region of the parameter space, while the projected bounds will constrain the model in a significant way. It is important to emphasize that in this theory one can have large lepton flavour violating effects in a consistent way in agreement with all experimental constraints. Then, combining the results for the $e_i \rightarrow e_j \gamma$ and the $\mu \rightarrow e$ conversion one can hope to test these predictions in the near future.

In this model one can have new contributions to neutrinoless double beta decay once the singly charged Higgs, δ^\pm , mixes with the charged Higgses in the bi-doublet Higgs. Since the charged Higgses in the

bi-doublet have to be heavy to avoid large flavour violation in the quark sector and the mixing angle is small the contribution to neutrinoless double beta decay is very small. Therefore, here one has the usual contributions in the context of the left-right theory but taking into account that the right-handed neutrinos are light. We have investigated the $e_i^\pm \rightarrow e_j^\pm e_k^\mp e_l^\pm$, and the predictions are quite below the current and projected experimental bounds.

IV. SUMMARY

We have discussed the main features of a simple left-right symmetric theory [29] where the neutrinos are Majorana fermions and their masses are generated at one-loop level. This theory predicts that the right-handed neutrinos are generically light and the existence of new interactions violating lepton number which can give rise to large contributions to flavor violating processes such as $\mu \rightarrow e\gamma$ and $\mu \rightarrow e$ conversion. We have investigated the lepton flavour violating contributions mediated by the new charged gauge bosons W_R^\pm . These contributions are very small in our model due to the GIM suppression. However, in other models where the right-handed neutrinos are heavy and close in mass to the W_R^\pm gauge bosons, these contributions can be very large motivating the search for the LFV processes.

Our left-right symmetric theory predicts the existence of a new singly charged Higgs which can give rise to very large flavour number violation effects. Since the charged scalar could be in principle very light due to the lack of strong collider constraints, one can have large contributions to LFV processes such as $\mu \rightarrow e\gamma$ and $\mu \rightarrow e$ conversion. We have shown that both processes provide non-trivial bounds on this model and thanks to the possibility of improving the experimental bounds in the near future one can hope to test these predictions. Together with the usual collider signatures from the W_R^\pm and right-handed neutrinos decays, one can use the predictions for lepton flavour number violating processes to realize the testability of this theory.

Acknowledgments: The work of C.M. has been supported in part by the Spanish Government and ERDF funds from the EU Commission [Grants No. FPA2014-53631-C2-1-P and SEV-2014-0398] and “La Caixa-Severo Ochoa” scholarship.

V. APPENDICES

A. Form factors

The relevant coefficients for the study of $\mu \rightarrow e$ conversion are given by

$$\begin{aligned}
C_{VR}^{(q)} &= \frac{Q_q e^2}{4\pi^2} \frac{1}{t} \sum_{N_i} \sum_{c,d} (\lambda_R^*)^{ce} \lambda_R^{d\mu} V_N^{ci} (V_N^*)^{di} \left(\frac{3}{4} + m_{\delta+}^2 \left(\frac{1}{2(m_{\delta+}^2 - m_{N_i}^2)} - \frac{1}{t} \right) \right. \\
&\quad \left. + \frac{m_{N_i}^2}{t} + \frac{((m_{N_i}^2 - m_{\delta+}^2)^2 + m_{N_i}^2 t)}{t} C_0[0, 0, t, m_{\delta+}, m_{\delta+}, m_{N_i}] \right. \\
&\quad \left. + \text{Log} \left(\frac{m_{N_i}^2}{m_{\delta+}^2} \right) \frac{2m_{N_i}^6 + 2m_{N_i}^2 m_{\delta+}^4 + m_{N_i}^4 (t - 4m_{\delta+}^2)}{2(m_{N_i}^2 - m_{\delta+}^2)^2 t} + \frac{(2m_{N_i}^2 - 2m_{\delta+}^2 + t) \Lambda[t, m_{\delta+}, m_{\delta+}]}{2t} \right), \tag{38}
\end{aligned}$$

and

$$\begin{aligned}
C_{DR} &= \frac{e}{8\pi^2} \sum_{N_i} \sum_{c,d} (\lambda_R^*)^{ce} \lambda_R^{d\mu} V_N^{ci} (V_N^*)^{di} \\
&\quad \left(\frac{12m_{N_i}^4 + 12m_{\delta+}^4 - 3m_{\delta+}^2 t + m_{N_i}^2 (5t - 24m_{\delta+}^2)}{4(m_{N_i}^2 - m_{\delta+}^2)t^2} + \frac{(6m_{N_i}^2 + 6m_{\delta+}^2 + t) \Lambda[t, m_{\delta+}, m_{\delta+}]}{2t^2} \right. \\
&\quad \left. + m_{N_i}^2 \text{Log} \left(\frac{m_{N_i}^2}{m_{\delta+}^2} \right) \frac{6m_{N_i}^4 - 6m_{\delta+}^4 - 2m_{\delta+}^2 t + m_{N_i}^2 (t - 12m_{\delta+}^2)}{2(m_{N_i}^2 - m_{\delta+}^2)^2 t^2} \right. \\
&\quad \left. + \frac{3m_{N_i}^4 + 3m_{\delta+}^4 - m_{\delta+}^2 t + m_{N_i}^2 (2t - 6m_{\delta+}^2)}{t^2} C_0[0, 0, t, m_{\delta+}, m_{\delta+}, m_{N_i}] \right), \tag{39}
\end{aligned}$$

where

$$\Lambda[t, m_{\delta+}, m_{\delta+}] \equiv \sqrt{1 - \frac{4m_{\delta+}^2}{t}} \text{Log} \left(\frac{2m_{\delta+}^2}{2m_{\delta+}^2 - \left(1 + \sqrt{1 - \frac{2m_{\delta+}^2}{t}} \right) t} \right), \tag{40}$$

and the Passarino–Veltman C_0 function is given by

$$\begin{aligned}
C_0[0, 0, t, m_{\delta+}, m_{\delta+}, m_{N_i}] = & \frac{1}{t} \left[-\text{DiLog} \left(\frac{(m_{N_i}^2 - m_{\delta+}^2)^2}{(m_{N_i}^2 - m_{\delta+}^2)^2 + m_{N_i}^2 t}, t \right) \right. \\
& + \text{DiLog} \left(\frac{(m_{N_i}^2 - m_{\delta+}^2)(m_{N_i}^2 - m_{\delta+}^2 + t)}{(m_{N_i}^2 - m_{\delta+}^2)^2 + m_{N_i}^2 t}, (m_{N_i} - m_{\delta+})t(m_{N_i}^2 - m_{\delta+}^2 + t) \right) \\
& + \text{DiLog} \left(\frac{2(m_{N_i} - m_{\delta+})(m_{N_i} + m_{\delta+})}{2m_{N_i}^2 - 2m_{\delta+}^2 + t - \sqrt{t(t - 4m_{\delta+}^2)}}, (m_{N_i} - m_{\delta+})t \right) \\
& - \text{DiLog} \left(-\frac{2(m_{N_i}^2 - m_{\delta+}^2 + t)}{2m_{\delta+}^2 - 2m_{N_i}^2 - t + \sqrt{t(t - 4m_{\delta+}^2)}}, (m_{N_i}^2 - m_{\delta+}^2 + t)t \right) \\
& + \text{DiLog} \left(\frac{2(m_{N_i} - m_{\delta+})(m_{N_i} + m_{\delta+})}{2m_{N_i}^2 - 2m_{\delta+}^2 + t + \sqrt{t(t - 4m_{\delta+}^2)}}, (m_{\delta+} - m_{N_i})t \right) \\
& \left. - \text{DiLog} \left(\frac{2(m_{N_i}^2 - m_{\delta+}^2 + t)}{2m_{N_i}^2 - 2m_{\delta+}^2 + t + \sqrt{t(t - 4m_{\delta+}^2)}}, t(m_{\delta+}^2 - m_{N_i}^2 - t) \right) \right]. \quad (41)
\end{aligned}$$

However, in the elastic limit $t \sim m_\mu^2$, one can neglect the variable t since $m_\mu^2 \sim 0$ compared to $m_{\delta+}$ running inside the loop. Hence, C_{DR} corresponds to the amplitude of $\mu \rightarrow e\gamma$ given by Eq.(23) with some arrangements of the factors to be consistent with the notation used in Eqs.(26) and (32),

$$C_{DR} = \frac{e}{8\pi^2} \frac{1}{m_{\delta+}^2} \sum_i \sum_{c,d} (\lambda_R^*)^{ce} \lambda_R^{d\mu} V_N^{ci} (V_N^*)^{di} G \left(\frac{m_{N_i}^2}{m_{\delta+}^2} \right), \quad (42)$$

where the function $G(x)$ is defined in Eq.(25), and the vectorial form factor becomes zero in this limit, i.e. $C_{VR}^{(q)} = 0$. We have used the Package-X [52] to perform the loop calculations.

B. Relevant Feynman rules

Here we list some of the most relevant Feynman rules for our study

- $\bar{\nu}_i e_j W_R^+ : i \frac{g_R}{\sqrt{2}} (B^\dagger)^{ij} \gamma_\mu P_R,$
- $\bar{N}_i e_j W_R^+ : i \frac{g_R}{\sqrt{2}} (V_N^\dagger)^{ij} \gamma_\mu P_R,$
- $\bar{\nu}_i e_j W_L^+ : i \frac{g_L}{\sqrt{2}} (V_\nu^\dagger)^{ij} \gamma_\mu P_L,$
- $\bar{N}_i e_j W_L^+ : i \frac{g_L}{\sqrt{2}} (A^\dagger)^{ij} \gamma_\mu P_L,$

- $\bar{\nu}_i e_j \delta^+ : 2i \sum_c (B^*)^{ci} \lambda_R^{cj} P_R + V_\nu^{ci} \lambda_L^{cj} P_L,$
- $\bar{N}_i e_j \delta^+ : 2i \sum_c (V_N^*)^{ci} \lambda_R^{cj} P_R + A^{ci} \lambda_L^{cj} P_L.$

Here we have neglected the mixing between the W_L and W_R gauge bosons for simplicity.

-
- [1] J. C. Pati and A. Salam, “Lepton Number as the Fourth Color,” Phys. Rev. D **10** (1974) 275 Erratum: [Phys. Rev. D **11** (1975) 703].
 - [2] R. N. Mohapatra and J. C. Pati, “A Natural Left-Right Symmetry,” Phys. Rev. D **11** (1975) 2558.
 - [3] G. Senjanovic, “Spontaneous Breakdown of Parity in a Class of Gauge Theories,” Nucl. Phys. B **153** (1979) 334.
 - [4] G. Senjanovic and R. N. Mohapatra, “Exact Left-Right Symmetry and Spontaneous Violation of Parity,” Phys. Rev. D **12** (1975) 1502.
 - [5] R. N. Mohapatra and G. Senjanovic, “Neutrino Mass and Spontaneous Parity Violation,” Phys. Rev. Lett. **44** (1980) 912.
 - [6] P. Minkowski, “ $\mu \rightarrow e$ Gamma At A Rate Of One Out Of 1-Billion Muon Decays?,” Phys. Lett. B **67** (1977) 421; T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe*, eds. O. Sawada et al., (KEK Report 79-18, Tsukuba, 1979), p. 95; M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, eds. P. van Nieuwenhuizen et al., (North-Holland, 1979), p. 315; S.L. Glashow, in *Quarks and Leptons*, Cargèse, eds. M. Lévy et al., (Plenum, 1980), p. 707.
 - [7] W. Y. Keung and G. Senjanovic, “Majorana Neutrinos and the Production of the Right-handed Charged Gauge Boson,” Phys. Rev. Lett. **50** (1983) 1427.
 - [8] P. S. B. Dev, R. N. Mohapatra and Y. Zhang, “Probing the Higgs Sector of the Minimal Left-Right Symmetric Model at Future Hadron Colliders,” JHEP **1605** (2016) 174 doi:10.1007/JHEP05(2016)174 [[arXiv:1602.05947](#) [hep-ph]].
 - [9] Y. Zhang, H. An, X. Ji and R. N. Mohapatra, “General CP Violation in Minimal Left-Right Symmetric Model and Constraints on the Right-Handed Scale,” Nucl. Phys. B **802** (2008) 247 doi:10.1016/j.nuclphysb.2008.05.019 [[arXiv:0712.4218](#) [hep-ph]].
 - [10] Y. Zhang, H. An, X. Ji and R. N. Mohapatra, “General CP Violation in Minimal Left-Right Symmetric Model and Constraints on the Right-Handed Scale,” Nucl. Phys. B **802** (2008) 247 [[arXiv:0712.4218](#) [hep-ph]].
 - [11] A. Maiezza, M. Nemevsek, F. Nesti and G. Senjanovic, “Left-Right Symmetry at LHC,” Phys. Rev. D **82** (2010) 055022 [[arXiv:1005.5160](#) [hep-ph]].
 - [12] M. Blanke, A. J. Buras, K. Gemmler and T. Heidsieck, “Delta $F = 2$ observables and $B \rightarrow X_q \gamma$ decays in the Left-Right Model: Higgs particles striking back,” JHEP **1203** (2012) 024 [[arXiv:1111.5014](#) [hep-ph]].
 - [13] S. Bertolini, A. Maiezza and F. Nesti, “Present and Future K and B Meson Mixing Constraints on TeV Scale Left-Right Symmetry,” Phys. Rev. D **89** (2014) no.9, 095028 [[arXiv:1403.7112](#) [hep-ph]].

- [14] A. Maiezza, G. Senjanovic and J. C. Vasquez, “Higgs Sector of the Left-Right Symmetric Theory,” [arXiv:1612.09146](#) [hep-ph].
- [15] M. Nemevsek, G. Senjanovic and V. Tello, “Connecting Dirac and Majorana Neutrino Mass Matrices in the Minimal Left-Right Symmetric Model,” *Phys. Rev. Lett.* **110** (2013) no.15, 151802 doi:10.1103/PhysRevLett.110.151802 [[arXiv:1211.2837](#) [hep-ph]].
- [16] V. Tello, M. Nemevsek, F. Nesti, G. Senjanovic and F. Vissani, “Left-Right Symmetry: from LHC to Neutrinoless Double Beta Decay,” *Phys. Rev. Lett.* **106** (2011) 151801 doi:10.1103/PhysRevLett.106.151801 [[arXiv:1011.3522](#) [hep-ph]].
- [17] W. C. Huang and J. Lopez-Pavon, “On neutrinoless double beta decay in the minimal left-right symmetric model,” *Eur. Phys. J. C* **74** (2014) 2853 doi:10.1140/epjc/s10052-014-2853-z [[arXiv:1310.0265](#) [hep-ph]].
- [18] G. Senjanovic and V. Tello, “Right Handed Quark Mixing in Left-Right Symmetric Theory,” *Phys. Rev. Lett.* **114** (2015) no.7, 071801 doi:10.1103/PhysRevLett.114.071801 [[arXiv:1408.3835](#) [hep-ph]].
- [19] A. Maiezza, M. Nemevsek, F. Nesti and G. Senjanovic, “Left-Right Symmetry at LHC,” *Phys. Rev. D* **82** (2010) 055022 doi:10.1103/PhysRevD.82.055022 [[arXiv:1005.5160](#) [hep-ph]].
- [20] M. Nemevsek, F. Nesti, G. Senjanovic and Y. Zhang, “First Limits on Left-Right Symmetry Scale from LHC Data,” *Phys. Rev. D* **83** (2011) 115014 doi:10.1103/PhysRevD.83.115014 [[arXiv:1103.1627](#) [hep-ph]].
- [21] C. Guo, S. Y. Guo, Z. L. Han, B. Li and Y. Liao, “Hunting for Heavy Majorana Neutrinos with Lepton Number Violating Signatures at LHC,” [arXiv:1701.02463](#) [hep-ph].
- [22] F. F. Deppisch, C. Hati, S. Patra, P. Pritimita and U. Sarkar, “Neutrinoless Double Beta Decay in Left-Right Symmetry with Universal Seesaw,” [arXiv:1701.02107](#) [hep-ph].
- [23] M. Mitra, R. Ruiz, D. J. Scott and M. Spannowsky, *Phys. Rev. D* **94** (2016) no.9, 095016 doi:10.1103/PhysRevD.94.095016 [[arXiv:1607.03504](#) [hep-ph]].
- [24] D. Borah and A. Dasgupta, “Charged lepton flavour violation and neutrinoless double beta decay in left-right symmetric models with type I+II seesaw,” *JHEP* **1607** (2016) 022 doi:10.1007/JHEP07(2016)022 [[arXiv:1606.00378](#) [hep-ph]].
- [25] A. Maiezza, M. Nemevsek and F. Nesti, “Perturbativity and mass scales in the minimal left-right symmetric model,” *Phys. Rev. D* **94** (2016) no.3, 035008 doi:10.1103/PhysRevD.94.035008 [[arXiv:1603.00360](#) [hep-ph]].
- [26] D. Alva, T. Han and R. Ruiz, “Heavy Majorana neutrinos from $W\gamma$ fusion at hadron colliders,” *JHEP* **1502** (2015) 072 doi:10.1007/JHEP02(2015)072 [[arXiv:1411.7305](#) [hep-ph]].
- [27] T. Han, I. Lewis, R. Ruiz and Z. g. Si, “Lepton Number Violation and W' Chiral Couplings at the LHC,” *Phys. Rev. D* **87** (2013) no.3, 035011 Erratum: [*Phys. Rev. D* **87** (2013) no.3, 039906] doi:10.1103/PhysRevD.87.035011, 10.1103/PhysRevD.87.039906 [[arXiv:1211.6447](#) [hep-ph]].
- [28] C. H. Lee, P. S. Bhupal Dev and R. N. Mohapatra, *Phys. Rev. D* **88** (2013) no.9, 093010 doi:10.1103/PhysRevD.88.093010 [[arXiv:1309.0774](#) [hep-ph]].
- [29] P. Fileviez Perez, C. Murgui and S. Ohmer, “Simple Left-Right Theory: Lepton Number Violation at the LHC,” *Phys. Rev. D* **94** (2016) no.5, 051701 doi:10.1103/PhysRevD.94.051701 [[arXiv:1607.00246](#) [hep-ph]].
- [30] A. Zee, “A Theory of Lepton Number Violation, Neutrino Majorana Mass, and Oscillation,” *Phys. Lett. B* **93**

- (1980) 389 Erratum: [Phys. Lett. B **95** (1980) 461].
- [31] R. H. Bernstein and P. S. Cooper, Phys. Rept. **532** (2013) 27 doi:10.1016/j.physrep.2013.07.002 [[arXiv:1307.5787](#) [hep-ex]].
 - [32] V. Khachatryan *et al.* [CMS Collaboration], “Search for heavy gauge W’ boson in events with an energetic lepton and large missing transverse momentum at sqrt(s) = 13 TeV,” [arXiv:1612.09274](#) [hep-ex].
 - [33] J. Adam *et al.* [MEG Collaboration], “New constraint on the existence of the $\mu^+ \rightarrow e^+ \gamma$ decay,” Phys. Rev. Lett. **110** (2013) 201801 doi:10.1103/PhysRevLett.110.201801 [[arXiv:1303.0754](#) [hep-ex]].
 - [34] B. Aubert *et al.* [BaBar Collaboration], “Searches for Lepton Flavor Violation in the Decays $\tau^+ \rightarrow e + \gamma$ and $\tau^+ \rightarrow \mu^+ \gamma$,” Phys. Rev. Lett. **104** (2010) 021802 doi:10.1103/PhysRevLett.104.021802 [[arXiv:0908.2381](#) [hep-ex]].
 - [35] T. Blum, A. Denig, I. Logashenko, E. de Rafael, B. Lee Roberts, T. Teubner and G. Venanzoni, [arXiv:1311.2198](#) [hep-ph].
 - [36] A. M. Baldini *et al.*, “MEG Upgrade Proposal,” [arXiv:1301.7225](#) [physics.ins-det].
 - [37] T. Aushev *et al.*, “Physics at Super B Factory,” [arXiv:1002.5012](#) [hep-ex].
 - [38] S. Weinberg and G. Feinberg, “Electromagnetic Transitions Between mu Meson and Electron,” Phys. Rev. Lett. **3** (1959) 111. doi:10.1103/PhysRevLett.3.111
 - [39] W. J. Marciano and A. I. Sanda, “The Reaction mu- Nucleus to e- Nucleus in Gauge Theories,” Phys. Rev. Lett. **38** (1977) 1512. doi:10.1103/PhysRevLett.38.1512
 - [40] A. Czarnecki, W. J. Marciano and K. Melnikov, “Coherent muon electron conversion in muonic atoms,” AIP Conf. Proc. **435** (1998) 409 doi:10.1063/1.56214 [[hep-ph/9801218](#)].
 - [41] V. Cirigliano, R. Kitano, Y. Okada and P. Tuzon, “On the model discriminating power of $\mu \rightarrow e$ conversion in nuclei,” Phys. Rev. D **80** (2009) 013002 doi:10.1103/PhysRevD.80.013002 [[arXiv:0904.0957](#) [hep-ph]].
 - [42] R. Kitano, M. Koike and Y. Okada, “Detailed calculation of lepton flavor violating muon electron conversion rate for various nuclei,” Phys. Rev. D **66** (2002) 096002 Erratum: [Phys. Rev. D **76** (2007) 059902] doi:10.1103/PhysRevD.76.059902, 10.1103/PhysRevD.66.096002 [[hep-ph/0203110](#)].
 - [43] V. Cirigliano, A. Kurylov, M. J. Ramsey-Musolf and P. Vogel, Phys. Rev. D **70** (2004) 075007 doi:10.1103/PhysRevD.70.075007 [[hep-ph/0404233](#)].
 - [44] C. Dohmen *et al.* [SINDRUM II Collaboration], “Test of lepton flavor conservation in $\mu \rightarrow e$ conversion on titanium,” Phys. Lett. B **317** (1993) 631. doi:10.1016/0370-2693(93)91383-X
 - [45] W. H. Bertl *et al.* [SINDRUM II Collaboration], “A Search for muon to electron conversion in muonic gold,” Eur. Phys. J. C **47** (2006) 337. doi:10.1140/epjc/s2006-02582-x
 - [46] W. Honecker *et al.* [SINDRUM II Collaboration], “Improved limit on the branching ratio of $\mu \rightarrow e$ conversion on lead,” Phys. Rev. Lett. **76** (1996) 200. doi:10.1103/PhysRevLett.76.200
 - [47] M. Aoki [DeeMe Collaboration], “A new idea for an experimental search for $\mu - e$ conversion,” PoS ICHEP **2010** (2010) 279.
 - [48] Y. G. Cui *et al.* [COMET Collaboration], “Conceptual design report for experimental search for lepton flavor violating $\mu - e$ conversion at sensitivity of 10^{-16} with a slow-extracted bunched proton beam (COMET),”

KEK-2009-10.

- [49] L. Morescalchi, “The Mu2e Experiment at Fermilab,” PoS DIS **2016** (2016) 259 [[arXiv:1609.02021](#) [hep-ex]].
- [50] Y. Kuno *et al.* [PRISM/PRIME Group], “Letter of Intent, An Experimental Search for a $\mu \rightarrow e$ Conversion at Sensitivity of the Order of 10^{18} with a Highly Intense Muon Source: PRISM J-PARC J-PARC. Letter of Intent (2006).”
- [51] T. Suzuki, D. F. Measday and J. P. Roalsvig, “Total Nuclear Capture Rates for Negative Muons,” Phys. Rev. C **35** (1987) 2212. doi:10.1103/PhysRevC.35.2212
- [52] H. H. Patel, “Package-X 2.0: A Mathematica package for the analytic calculation of one-loop integrals,” [arXiv:1612.00009](#) [hep-ph]; H. H. Patel, “Package-X: A Mathematica package for the analytic calculation of one-loop integrals,” Comput. Phys. Commun. **197** (2015) 276 doi:10.1016/j.cpc.2015.08.017 [[arXiv:1503.01469](#) [hep-ph]].