

Means Moments and Newton's Inequalities

R. Sharma, A. Sharma, R. Saini and G. Kapoor

Department of Mathematics & Statistics

Himachal Pradesh University

Shimla - 171005,

India

email: rajesh.sharma.hpn@nic.in

Abstract. It is shown that Newton's inequalities and the related Maclaurin's inequalities provide several refinements of the fundamental Arithmetic mean - Geometric mean - Harmonic mean inequality in terms of the means and variance of positive real numbers. We also obtain some inequalities involving third and fourth central moments of real numbers.

AMS classification. 60E15

Key words and phrases. Arithmetic mean, Geometric mean, Moments, Newton's identities.

1 Introduction

Let x_1, x_2, \dots, x_n denote n real numbers. Their r th moment and r th central moment are respectively the numbers

$$m'_r = \frac{1}{n} \sum_{i=1}^n x_i^r \quad (1.1)$$

and

$$m_r = \frac{1}{n} \sum_{i=1}^n (x_i - m'_1)^r, \quad (1.2)$$

where $r = 1, 2, \dots$. Note that $m_1 = 0$, m'_1 is the arithmetic mean and m_2 is the variance of n real numbers x_1, x_2, \dots, x_n . It is customary to denote m'_1 and m_2 respectively by A and s^2 . The Geometric mean (G) and Harmonic mean (H) of n positive real numbers x_1, x_2, \dots, x_n are respectively the numbers

$$G = \left(\prod_{i=1}^n x_i \right)^{\frac{1}{n}} \quad \text{and} \quad H = \left(\frac{1}{n} \sum_{i=1}^n \frac{1}{x_i} \right)^{-1}.$$

The k^{th} elementary symmetric function C_k of x_1, x_2, \dots, x_n is the sum of the products taken k at a time of different x_i 's. The k^{th} elementary symmetric mean is the number

$$S_k = \frac{1}{\binom{n}{k}} C_k. \quad (1.3)$$

We write $S_0 = C_0 = 1$.

A well known theorem of Newton (1707) says that elementary symmetric means satisfy the inequality

$$S_k^2 \geq S_{k+1} S_{k-1}, \quad (1.4)$$

for all $k = 1, 2, \dots, n$, and equality holds if and only if the numbers x_i 's are all equal.

A related theorem due to Maclaurin (1729) asserts that the elementary symmetric means of n positive real numbers satisfy the following inequalities string:

$$S_1 \geq S_2^{\frac{1}{2}} \geq \dots \geq S_n^{\frac{1}{n}}. \quad (1.5)$$

Equality occurs if and only if x_i 's are all equal. One can easily see that the inequalities (1.5) also follow from the inequalities (1.4). For more detail, see Hardy et al (1952).

Newton's identities give the relations between the elementary symmetric functions and the sum of the powers of x_1, x_2, \dots, x_n . Let

$$\alpha_k = \sum_{i=1}^n x_i^k,$$

and write $\alpha_0 = 1$. Then Newton's identities say that

$$kC_k = \sum_{i=1}^k (-1)^{i-1} C_{k-i} \alpha_i. \quad (1.6)$$

The well known Arithmetic mean - Geometric mean - Harmonic mean inequality (A-G-H inequality) says that $H \leq G \leq A$. Alternative proofs, refinements and generalizations of A-G-H inequality have been studied extensively in literature over roughly the last two centuries. See Bullen (2003). Note that $S_1 = A$ and $S_n^{\frac{1}{n}} = G$. The Maclaurin inequalities (1.5) therefore provide the refinements of the A-G inequality. These refinements are in terms of the symmetric means S_k . On using Newton's identity we can find the relation between symmetric means and moments of the n real numbers. Then, the Maclaurin inequalities can be used to find the refinements of the A-G inequality in terms of the expressions involving means and moments. The purpose of the present note is to point out some such inequalities which remains unnoticed in the literature in the present explicit forms.

Our first theorem gives a refinement of the A-G-H inequality in terms of expressions involving arithmetic mean and harmonic mean (see Theorem 2.1, below). A refinement of the A-G inequality involving second moment is given. This also provides an upper bound for the variance, and Brunk inequalities (Theorem 2.2). A further refinement of this inequality involves harmonic mean (Theorem 2.3). We derive bounds for the third and fourth moments and central moments of statistical interest (Theorem 2.4 and 2.7). The refinements of A-G inequality in terms of first four moments are also given (Theorem 2.5, 2.6 and 2.8).

2 Main results

Theorem 2.1. For n positive real numbers x_1, x_2, \dots, x_n and with notations as above, we have

$$H \leq H \left(\frac{A}{H} \right)^{\frac{1}{n}} \leq G \leq \left(\frac{H}{A} \right)^{\frac{1}{n}} A \leq A. \quad (2.1)$$

Proof. The third inequality (2.1) follows from the inequality $S_1 \geq S_{n-1}^{\frac{1}{n-1}}$ in (1.5) and the fact that from (1.3) we have $S_1 = A$ and $S_{n-1}H = G^n$. The second inequality (2.1) follows on applying third inequality (2.1) to the numbers $\frac{1}{x_1}, \frac{1}{x_2}, \dots, \frac{1}{x_n}$. The extreme inequalities are immediate. ■

It may be noted here that from (2.1) we also have

$$\left(\frac{G}{H} \right)^{\frac{1}{n-1}} G \leq A \leq \left(\frac{G}{H} \right)^{n-1} G.$$

and

$$\left(\frac{G}{A} \right)^{n-1} G \leq H \leq \left(\frac{G}{A} \right)^{\frac{1}{n-1}} G.$$

Theorem 2.2. For n positive real numbers x_1, x_2, \dots, x_n and with notations as above, we have

$$G \leq \sqrt{\frac{nA^2 - m'_2}{n-1}} \leq A. \quad (2.2)$$

Proof. It follows from the inequality $S_2 \geq (S_n)^{\frac{2}{n}}$ in (1.5) that for n positive real numbers x_1, x_2, \dots, x_n the inequality

$$\frac{2}{n(n-1)} \sum_{i < j}^n x_i x_j \geq G^2, \quad (2.3)$$

holds true. From (1.6), $2C_2 = C_1\alpha_1 - C_0\alpha_2$ where $C_0 = 1$, $C_1 = \alpha_1 = \sum x_i$, $\alpha_2 = \sum x_i^2$ and $C_2 = \sum_{i < j} x_i x_j$. Therefore, on using (1.1), we get that

$$2 \sum_{i < j}^n x_i x_j = \left(\sum_{i=1}^n x_i \right)^2 - \sum_{i=1}^n x_i^2 = n(nA^2 - m'_2). \quad (2.4)$$

Insert (2.4) in (2.3); a little computation leads to first inequality (2.2). The second inequality (2.2) follows from the fact that $m'_2 \geq A^2$. ■

The inequality (2.1) can be written in several different ways, for example the inequality

$$G \leq A \sqrt{1 - \frac{1}{n-1} \left(\frac{s}{A}\right)^2}$$

follows from (2.1) on using $m'_2 = s^2 + A^2$. We also have

$$m'_2 \leq nA^2 - (n-1)G^2 \quad (2.5)$$

and

$$s^2 \leq (n-1)(A^2 - G^2). \quad (2.6)$$

We note that Brunk's inequalities (1959) follow from the inequality (2.6). Let $a \leq x_i \leq b$ for all $i = 1, 2, \dots, n$. From (2.6), $s^2 \leq (n-1)A^2$. Apply this to n positive numbers $x_i - a$, we get Brunk's inequality, $s \leq \sqrt{n-1}(A - a)$. Likewise, the inequality $s \leq \sqrt{n-1}(b - A)$ follows from (2.6). For the related variance upper bounds, see Bhatia and Davis (2000), Sharma (2008) and Sharma et al (2010).

Theorem 2.3. With conditions as in above theorem and for $n \geq 3$, we have

$$G \leq \left(\frac{G^n}{H}\right)^{\frac{1}{n-1}} \leq \sqrt{\frac{nA^2 - m'_2}{n-1}} \leq A. \quad (2.7)$$

Proof. From (1.5), $S_2 \geq (S_{n-1})^{\frac{2}{n-1}}$. Therefore, for n positive real numbers x_1, x_2, \dots, x_n the inequality

$$\frac{2}{n(n-1)} \sum_{i < j}^n x_i x_j \geq \left(\frac{C_{n-1}}{n}\right)^{\frac{2}{n-1}}, \quad (2.8)$$

holds true. Also

$$C_{n-1} = \prod_{i=1}^n x_i \sum_{i=1}^n \frac{1}{x_i} = n \frac{G^n}{H}. \quad (2.9)$$

Insert (2.4) and (2.9) in (2.8); a little computation leads to the second inequality (2.7)

■

From (2.7), we also have

$$G \leq \left(\frac{G^n}{H}\right)^{\frac{1}{n-1}} \leq A \sqrt{1 - \frac{1}{n-1} \left(\frac{s}{A}\right)^2}$$

$$m'_2 \leq nA^2 - (n-1) \left(\frac{G^n}{H} \right)^{\frac{2}{n-1}}. \quad (2.10)$$

and

$$s^2 \leq (n-1) \left(A^2 - \left(\frac{G^n}{H} \right)^{\frac{2}{n-1}} \right). \quad (2.11)$$

The inequalities (2.10) and (2.11) provide refinements of the inequalities (2.5) and (2.6), respectively.

Theorem 2.4. For n real numbers x_1, x_2, \dots, x_n and $n \geq 3$, we have

$$Am'_3 \leq \frac{1}{2(n-1)} [(n-2)m_2'^2 + n(n+1)m_2'A^2]. \quad (2.12)$$

Proof. For $k = 2$, the inequality (1.4) gives $S_2^2 \geq S_1S_3$. Therefore, for n real numbers x_1, x_2, \dots, x_n , we have

$$\left(\frac{2}{n(n-1)} \sum_{i < j}^n x_i x_j \right)^2 \geq \frac{6A}{n(n-1)(n-2)} \sum_{i < j < k}^n x_i x_j x_k. \quad (2.13)$$

Further, from (1.6), $3C_3 = C_2\alpha_1 - C_1\alpha_2 + \alpha_3$ where $\alpha_3 = \sum x_i^3$ and $C_3 = \sum_{i < j < l} x_i x_j x_l$.

Therefore, on using (2.3) and (1.1), we find that

$$\begin{aligned} 3 \sum_{i < j < k}^n x_i x_j x_k &= \sum_{i=1}^n x_i^3 + \frac{1}{2} \left(\sum_{i=1}^n x_i \right)^3 - \frac{3}{2} \sum_{i=1}^n x_i \sum_{i=1}^n x_i^2 \\ &= n \left(m'_3 + \frac{n^2}{2} A^3 - \frac{3}{2} n A m'_2 \right). \end{aligned} \quad (2.14)$$

On inserting (2.4) and (2.14) in (2.13) and simplifying the resulting expressions; we immediately get (2.12). ■

The twin inequality

$$Am_3 \leq \frac{n-2}{2(n-1)} s^4 + \frac{(n-2)}{2} A^2 s^2 \quad (2.15)$$

follows from (2.12) on using the relations $m'_2 = s^2 + A^2$ and $m'_3 = m_3 + 3As^2 + A^3$. It may be noted here that

$$3 \sum_{i < j < k}^n x_i x_j x_k = n \left(m_3 - \frac{3(n-2)}{2} A s^2 + \frac{(n-1)(n-2)}{2} A^3 \right)$$

and the inequality

$$m_3 \leq \frac{n-2}{2(n-1)} \frac{s^4}{A} + \frac{(n-2)}{2} A s^2$$

holds true for positive real numbers x_1, x_2, \dots, x_n .

Corollary 2.1. Under the conditions of Theorem 2.2, we have

$$m_3 \leq (n-2) A s^2 \leq (n-1)(n-2) A^3. \quad (2.16)$$

Proof. The inequality (2.16) follows from (2.15) and the fact that for positive real numbers x_1, x_2, \dots, x_n , we have $A > 0$ and $s \leq \sqrt{n-1} A$. ■

Note that the second inequality (2.16) implies that for n real numbers $a \leq x_i \leq b$, $i = 1, 2, \dots, n$, we have $m_3 \leq (n-1)(n-2)(b-A)^3$ and $m_3 \leq (n-1)(n-2)(A-a)^3$. Thus, we get inequalities analogous to Brunk's inequalities (1959) for third central moment. Likewise, from the first inequality (2.16), we have $m_3 \leq (n-2)(b-A)s^2$ and $m_3 \leq (n-2)(A-a)s^2$. Also, see Sharma et al (2012).

Theorem 2.5. With conditions as in Theorem 2.3, we have

$$G \leq A \left(\frac{n^2}{(n-1)(n-2)} \left(1 - \frac{3}{n} \frac{m'_2}{A^2} + \frac{2}{n^2} \frac{m'_3}{A^3} \right) \right)^{\frac{1}{3}}. \quad (2.17)$$

Proof. It follows from the inequality $S_3 \geq S_n^{\frac{3}{n}}$ in (1.5) that for n positive real numbers x_1, x_2, \dots, x_n , we have

$$\frac{6}{n(n-1)(n-2)} \sum_{i < j < k}^n x_i x_j x_k \geq G^3. \quad (2.18)$$

Insert (2.14) in (2.18); a little computation leads to (2.17). ■

From (2.17), we also have

$$G \leq A \left(1 - \frac{3}{n-1} \left(\frac{s}{A} \right)^2 + \frac{2}{(n-1)(n-2)} \frac{m_3}{A^3} \right)^{\frac{1}{3}}$$

$$m'_3 \geq \frac{1}{2} [3nAm'_2 - n^2A^3 + (n-1)(n-2)G^3]$$

and

$$m_3 \geq \frac{(n-1)(n-2)}{2} \left[\frac{3}{(n-1)} A s^2 - (A^3 - G^3) \right].$$

Theorem 2.6. With conditions as in above theorem and for $n \geq 4$, we have

$$G \leq \left(\frac{G^n}{H} \right)^{\frac{1}{n-1}} \leq A \left(\frac{n^2}{(n-1)(n-2)} \left(1 - \frac{3}{n} \frac{m'_2}{A^2} + \frac{2}{n^2} \frac{m'_3}{A^3} \right) \right)^{\frac{1}{3}}. \quad (2.19)$$

Proof. It follows from inequality $S_3 \geq S_{n-1}^{\frac{3}{n-1}}$ in (1.5) that for n positive real numbers x_1, x_2, \dots, x_n , we have

$$\frac{6}{n(n-1)(n-2)} \sum_{i < j < k}^n x_i x_j x_k \geq \left(\frac{G^n}{H} \right)^{\frac{3}{n-1}}. \quad (2.20)$$

Insert (2.14) in (2.20); a little computation leads to (2.19). ■

From (2.19), we also have

$$G \leq \left(\frac{G^n}{H} \right)^{\frac{1}{n-1}} \leq A \left(1 - \frac{3}{n-1} \left(\frac{s}{A} \right)^2 + \frac{2}{(n-1)(n-2)} \frac{m_3}{A^3} \right)^{\frac{1}{3}},$$

$$m'_3 \geq \frac{1}{2} \left[3nAm'_2 - n^2A^3 + (n-1)(n-2) \left(\frac{G^n}{H} \right)^{\frac{3}{n-1}} \right]$$

and

$$m_3 \geq \frac{(n-1)(n-2)}{2} \left[\frac{3}{(n-1)} As^2 - \left(A^3 - \left(\frac{G^n}{H} \right)^{\frac{3}{n-1}} \right) \right].$$

Recently, Sharma and Bhandari (2015) have shown that the Newton inequality provides an inequality involving skewness and kurtosis,

$$1 + \frac{m_3^2}{m_2^3} \leq \frac{m_4}{m_2^2} \leq \frac{1}{2} \frac{n-3}{n-2} \frac{m_3^2}{m_2^3} + \frac{n}{2}, \quad n \geq 3.$$

Sharma et al (2015) have obtained bounds for the fourth central moment. Here we show that Newton inequality provide several other inequalities involving first four moments.

Theorem 2.7. For n real numbers x_1, x_2, \dots, x_n and $n \geq 4$, we have

$$(nA^2 - m'_2) m'_4 \geq \frac{1}{6(n-3)} \left(\begin{aligned} & n^4 A^6 - n^3 (n+4) m'_2 A^4 + 4n^2 (n-1) m'_3 A^3 + 9n^2 m_2'^2 A^2 \\ & + 4n(n-5) m'_2 m'_3 A - 3n(n-2) m_2'^3 - 4(n-3) m_3'^2 \end{aligned} \right). \quad (2.21)$$

Proof. The inequality (2.21) follows from the inequality $S_3^2 \geq S_2 S_4$ in (1.4). Note that from (1.6), $4C_4 = C_3 \alpha_1 - C_2 \alpha_2 + C_1 \alpha_3 - C_0 \alpha_4$. Therefore, on using (2.3), (2.14) and (1.1), we get that

$$24 \sum_{i < j < k < l}^n x_i x_j x_k x_l = n^4 A^4 - 6n^3 m'_2 + 8n^2 Am'_3 + 3n^2 m_2'^2 - 6nm'_4.$$

■

Theorem 2.8. For n positive real numbers x_1, x_2, \dots, x_n and $n \geq 4$, we have

$$G^4 \leq \frac{1}{(n-1)(n-2)(n-3)} (n^3 m_1'^4 - 6n^2 m_2' m_1'^2 + 8n m_3' m_1' + 3n m_2'^2 - 6m_4') \quad (2.22)$$

Proof. From (1.5), $S_4 \geq (S_n)^{\frac{4}{n}}$. Therefore, for n positive real numbers x_1, x_2, \dots, x_n the inequality

$$\frac{24}{n(n-1)(n-2)(n-3)} \sum_{i < j < k < l}^n x_i x_j x_k x_l \geq G^4,$$

holds true. ■

The inequality (2.22) also gives

$$G \leq A \left(1 - \frac{6}{n-1} \left(\frac{s}{A} \right)^2 + \frac{8}{(n-1)(n-2)} \frac{m_3}{A^3} + \frac{3(ns^4 - 2m_4)}{(n-1)(n-2)(n-3)A^4} \right)^{\frac{1}{4}}.$$

Acknowledgements. The authors are grateful to Prof. Rajendra Bhatia for the useful discussions and suggestions, and I.S.I. Delhi for a visit in February 2017 when this work was done. The support of the UGC-SAP is also acknowledged.

References

- [1] Bhatia R, Davis C, A better bound on the variance, Amer. Math. Monthly, 107, 353-357, (2000) .
- [2] Brunk H. D, Note on two papers of K. R. Nair, Journal of the Indian Society of Agricultural Statistics, 11, 186-189, (1959).
- [3] Hardy G, Littlewood J E and Polya G, Inequalities, Cambridge Mathematical Library, (1952).
- [4] Bullen P. S, Handbook of Means and Their Inequalities, Springer, (2003).

- [5] Maclaurin C, A second letter to Martin Folkes, Esq. ;concerning the roots of equations, with the demonstration of other rules in algebra, phil. Transactions, 36 , 59-96, (1729).
- [6] Newton I, Arithmetica universalis: sive de compositione et resolutione arithmetica liber, (1707).
- [7] Sharma R, Some more inequalities for Arithmetic Mean, Harmonic Mean and Variance, JMI, 2, 109-114, (2008) .
- [8] Sharma R, Gupta M, Kapoor G, Some better bounds on variance with applications, JMI,4, 355-363, (2010).
- [9] Sharma R, Bhandari R, Gupta M, Inequalities related to the Cauchy-Schwarz inequality, Sankhya 74(A), 101-111, (2012)
- [10] Sharma R, Bhandari R, Skewness Kurtosis and Newton's Inequalities , Rocky Mountain Journal of Mathematics, 45, 1639-1643, (2015) .
- [11] Sharma R, Kumar R, Saini R, Kapoor G, Bounds on spread of matrices related to fourth central moment, Bull. Malays. Math. Sci. Soc. DOI 10.1007/540840-015-0267-1, (2015).