

COHOMOLOGY OF UNIPOTENT GROUP SCHEMES

ERIC M. FRIEDLANDER*

ABSTRACT. We verify that universal classes in the cohomology of GL_N determine explicit cohomology classes of Frobenius kernels $G_{(r)}$ of various linear algebraic groups G . We consider the relationship of $\varprojlim_r H^*(U_{(r)}, k)$ to the rational cohomology $H^*(U, k)$ of many unipotent algebraic groups U . The second half of this paper investigates in detail the cohomology of Frobenius kernels $(U_3)_{(r)}$ of the Heisenberg group $U_3 \subset GL_3$.

CONTENTS

0. Introduction	1
1. 1-parameter subgroups, exponential type, and cohomology	3
2. Stabilization of $H^\bullet(U_{(r)}, k)$ with respect to r	10
3. The map $\bar{\eta}_{U_J/\Gamma_3, r} : \bar{S}^*((U_J/\Gamma_3)_{(r)}) \rightarrow H^\bullet((U_J/\Gamma_3)_{(r)}, k)$	14
4. The map $\bar{\eta}_{U_3, r} : \bar{S}^*((U_3)_{(r)}) \rightarrow H^\bullet((U_3)_{(r)}, k)$	23
5. Questions	28
References	28

0. INTRODUCTION

We consider linear algebraic groups G defined over a field of characteristic $p > 0$ and their Frobenius kernels $G_{(r)}$. We investigate the rational cohomology algebra $H^*(G, k)$ of G and the cohomology algebra $H^*(G_{(r)}, k)$. Our results are of two types. The first two sections are of a general nature, applying to a wide class of unipotent groups. The next two sections provide more detailed information for $H^*((U_3)_{(r)}, k)$, where $U_3 \subset GL_3$ is the Heisenberg group.

The cohomology of groups has played an important role in various aspects of topology, number theory, and algebraic geometry. Our initial interest was generated by the foundational work of D. Quillen [19], [20] and the connections with algebraic K-theory as also developed by Quillen [21]. Subsequently, thanks to the work of many mathematicians beginning with J. Alperin - L. Evens [2], J. Carlson [4], and E. Cline - B. Parshall - L. Scott [6], cohomology of groups has evolved into a useful tool (“support varieties”) for the study of representations of finite group schemes. We would be amiss not to mention work of the author with B. Parshall (e.g., [10]), A. Suslin (e.g., [13]), and J. Pevtsova (e.g., [12]).

Date: March 6, 2024.

2010 Mathematics Subject Classification. 20G05, 20C20, 20G10.

Key words and phrases. rational cohomology, Frobenius kernels, unipotent algebraic groups.

* partially supported by the Simons Foundation.

Considerable progress has been made in computing the cohomology of infinitesimal group schemes of height 1, beginning with work of Friedlander-Parshall [10]; this was extended by C. Drupieski - D. Nakano - N. Ngo [8] to a determination of the cohomology algebra $H^*((U_J)_1, k)$ for unipotent radicals U_J of parabolic subgroups of reductive groups (as considered in this paper) for sufficiently large primes p , and further investigated by J. Carlson - D. Nakano [5] for small primes. The focus of this paper is the challenge of achieving explicit computations of the cohomology of infinitesimal group schemes of height > 1 .

The work of A. Suslin, C. Bendel, and the author [23], [24] gives a qualitative description of $H^*(G_{(r)}, k)$ for the cohomology of Frobenius kernels of any linear algebraic group G . A key step in this description for G utilizes universal classes for GL_N constructed in [13] leading to a map $\bar{\phi}_{GL_N} : k[V_r(GL_N)] \rightarrow H^*((GL_N)_{(r)}, k)$, complementing in the special case of $G = GL_N$, the canonical map $\psi : H^*(G_{(r)}, k) \rightarrow k[V_r(G)]$; here, $k[V_r(G)]$ is the coordinate algebra of the variety of 1-parameter subgroups $\mathbb{G}_{a(r)} \rightarrow G$.

In Section 1, we extend this construction to various linear algebraic groups G with an embedding $G \rightarrow GL_N$ of exponential type. In particular, Theorem 1.7 establishes a map $\phi_{G,r}$ from an explicit symmetric algebra determined by $\mathfrak{g} = \text{Lie}(G)$ to $H^*(G_{(r)}, k)$ whose image is highly non-trivial. For U_J the unipotent radical of a parabolic subgroup of GL_N , Proposition 1.9 shows that $\phi_{U_J,r}$ determines the map $\bar{\phi}_{U_J,r}$ satisfying the properties established for $\bar{\phi}_{GL_N,r}$ in [23], [24].

The author's original motivation for the study of $H^*(U_{(r)}, k)$ was to investigate the feasibility of a cohomology-based theory of support varieties for a unipotent algebraic group U complementing his theory using 1-parameter subgroups [9] for much more general linear algebraic groups. One reason for restricting our attention to unipotent algebraic groups is that $H^*(G, k)$ is trivial for many linear algebraic groups (for example, if G is simple) but is never trivial if G is unipotent. Our computations show such a cohomology-based theory even for unipotent linear algebraic groups is unlikely; see, for example, Corollary 2.7 in conjunction with Theorem 4.5.

We present in Section 2 various results concerning the limiting behavior of the cohomology of Frobenius kernels $U_{(r)}$ of a unipotent algebraic group U as r increases. In particular, Theorem 2.2 established that the inverse limit with respect to r of such Frobenius kernels is additively isomorphic to the rational cohomology of U ; this verification is the first of several occasions in this paper where we utilize the Andersen-Jantzen spectral sequence [1] for the cohomology of a connected group scheme. The results of this Section 2 formulate the general principal that the explicit cohomology classes considered in other sections of this paper “vanish in the limit.”

A second motivation for our calculations is internal, within the general framework of cohomology of groups. Let G be a simple algebraic group, $U_J \subset P_J \subset G$ the unipotent radical of a parabolic subgroup, and $\{\Gamma_v, v \geq 1\}$ the descending central series of U_J . This descending central series is well described by H. Azad, M. Barry, and G. Seitz in [3]. Using the Lyndon-Hochschild-Serre spectral sequence [15] for the central extension $1 \rightarrow \Gamma_2/\Gamma_3 \rightarrow U_J \rightarrow U_J/\Gamma_2 \rightarrow 1$, we construct in Definition 3.8 the map of graded k -algebras

$$\eta_{U_J/\Gamma_3} : S^*((U_J/\Gamma_3)_{(r)}) \rightarrow H^*((U_J/\Gamma_3)_{(r)}, k)$$

where $S^*((U_J/\Gamma_3)_{(r)})$ is a polynomial algebra with generating subspaces $\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_2)^{\#(\ell+1)}[2]$ (in cohomology degree 2, where the Frobenius twist $(-)^{(\ell+1)}$ indicates the torus action) and $\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}]$. The subtlety here is the existence of the choice of an intrinsic map η_{U_J/Γ_3} . This is established with the help of the Andersen-Jantzen spectral sequence.

Proposition 3.11 verifies that η_{U_J/Γ_3} factors through $\bar{\eta}_{U_J/\Gamma_3,r} : \bar{S}^*((U_J/\Gamma_3)_{(r)}) \rightarrow H^*((U_J/\Gamma_3)_{(r)}, k)$, reflecting the kernel of the inflation map $H^*((U_J/\Gamma_2)_{(r)}, k) \rightarrow H^*((U_J/\Gamma_3)_{(r)}, k)$; this kernel has explicit generators given by (3.3.2) arising from a non-zero differential in the spectral sequence. A key ingredient in this construction is the action of the Steenrod algebra on the LHS spectral sequence which enables us to identify permanent cycles.

We view the map $\bar{\eta}_{U_J/\Gamma_3,r}$ as a good “explicit” model for $H^*((U_J/\Gamma_3)_{(r)}, k)$. As summarized in Theorem 4.5, $\bar{\eta}_{U_3,r} = \bar{\phi}_{U_3,r} : \bar{S}^*((U_3)_{(r)}) \rightarrow H^*((U_3)_{(r)}, k)$ is a map from an integrally closed domain with known generators and relations to $H^*((U_3)_{(r)}, k)$ which is a.) injective, b.) surjective onto p -th powers, and c.) has associated graded map $gr(\bar{\eta}_{U_3,r})$ which is both injective and surjective onto p -th powers.

We anticipate that similar arguments should apply to $H^*((U_J/\Gamma_{v+1})_{(r)}, k)$ for any $v \geq 2$ once a suitable action of the Steenrod algebra on the Andersen-Jantzen spectral sequence [1] is established. This is only one of the many challenges left unanswered in the present paper, and appears as Question 3 in the list of seven questions given in Section 5.

In what follows, k denotes an algebraically closed field of characteristic $p > 2$. We denote by $H^*(G, k)$ the (rational, or Hochschild) cohomology of an affine group scheme G over k and by $H^\bullet(G, k) \subset H^*(G, k)$ the commutative subalgebra of cohomology classes of even degree. We use $V^\#$ to denote the k -linear dual of a k -vector space V . Other than in Section 2 where we consider the effect of increasing r , we fix an arbitrary positive integer r .

We thank Robert Guralnick for helpful discussions. We especially express our gratitude and admiration to the patient referee for detailed and constructive corrections.

1. 1-PARAMETER SUBGROUPS, EXPONENTIAL TYPE, AND COHOMOLOGY

In this section, we extend the formulation of the map of k -algebras

$$(1.0.1) \quad \phi_{GL_N,r} : S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{gl}_N^{\#(r)}[2p^{r-\ell-1}])) \rightarrow H^\bullet(GL_{N(r)}, k)$$

given by A. Suslin and the author in [13] to the Frobenius kernels $U_{(r)}$ of various unipotent subgroups $U \subset GL_N$. In contrast to our subsequent constructions, this extension involves little computation. Throughout this discussion, r will denote an arbitrary positive integer. The main result of this section, Theorem 1.7 gives sufficient conditions for a simple group G given together with an embedding $i : G \subset GL_N$ to admit an induced map $\phi_{G,r}$ which in turn determine an induced map for certain unipotent subgroups $U_J \subset G$.

Recall that for any linear algebraic group G over k , the r -th iterate $F^r : G \rightarrow G^{(r)}$ of the Frobenius map $F : G \rightarrow G^{(1)}$ admits a scheme theoretic kernel $G_{(r)} \equiv \ker\{F^r\}$ which is an infinitesimal group scheme of height r . The coordinate algebra $k[G_{(r)}]$ of $G_{(r)}$ equals the finite dimensional commutative Hopf algebra $k[G]/I^{p^r}$, where I is the maximal ideal at the identity of G and where I^{p^r} is the ideal generated

by $\{f^{p^r}, f \in I\}$. A (rational) $G_{(r)}$ -module is a comodule for $k[G_{(r)}]$ or, equivalently, a module for $kG_{(r)} \equiv (k[G_{(r)}])^\#$ (the k -linear dual of $k[G_{(r)}]$ with its inherited Hopf algebra structure). For G defined over \mathbb{F}_{p^r} , we may view the Frobenius map F^r as an endomorphism of G and $G_{(r)} \subset G$ as the kernel of $F^r : G \rightarrow G$.

The universal, GL_N -invariant classes $e_{r-\ell} = e_{r-\ell}^{(0)} \in H^{2p^{r-\ell-1}}(GL_N, \mathfrak{gl}_N^{(r-\ell)})$ of [13] and their ℓ -th Frobenius twist $e_{r-\ell}^{(\ell)} \in H^{2p^{r-\ell-1}}(GL_N, \mathfrak{gl}_N^{(r)})$ (i.e., pull-back along the ℓ -th iterate F^ℓ of the Frobenius morphism $F : GL_N \rightarrow GL_N$) are elements in the rational cohomology of the (reductive) algebraic group GL_N . The restriction of $e_{r-\ell}^{(\ell)}$ to $GL_{N(r)}$,

$$(1.0.2) \quad (e_{r-\ell}^{(\ell)})_{(r)} \in H^{2p^{r-\ell-1}}(GL_{N(r)}, \mathfrak{gl}_N^{(r)}) \simeq H^{2p^{r-\ell-1}}(GL_{N(r)}, k) \otimes \mathfrak{gl}_N^{(r)},$$

can be identified with a GL_N -equivariant map

$$\mathfrak{gl}_N^{(r)}[2p^{r-\ell-1}] \rightarrow H^{2p^{r-\ell-1}}(GL_{N(r)}, k)$$

(vanishing on the dual trace class $Tr^{(r)} \in \mathfrak{gl}_N^{(r)}$), thereby determining the GL_N -equivariant map of commutative k -algebras (1.0.1). For $\ell < r$, the Frobenius map F^ℓ restricts to $F^\ell : GL_{N(r)} \rightarrow GL_{N(r)}$ and factors as

$$GL_{N(r)} \twoheadrightarrow GL_{N(r)}/GL_{N(\ell)} \simeq GL_{N(r-\ell)} \subset GL_{N(r)}.$$

The Frobenius twist $(e_{r-\ell}^{(\ell)})_{(r)}$ can thus be realized as the pull-back along $GL_{N(r)} \twoheadrightarrow GL_{N(r-\ell)}$ of $(e_{r-\ell}^{(0)})_{(r-\ell)} \in H^{2p^{r-\ell-1}}(GL_{N(r-\ell)}, \mathfrak{gl}_N^{(r-\ell)})$.

The following proposition summarizes some of the basic properties of these universal classes (whose proofs can be found in [13] and [23]).

Proposition 1.1. *With notation as above,*

- (1) $(e_{r-\ell}^{(\ell)})_{(\ell+1)} \neq 0$ in $H^{2p^{r-\ell-1}}(GL_{N(\ell+1)}, \mathfrak{gl}_N^{(r)})$.
- (2) $(e_{r-\ell}^{(\ell)})_{(\ell)} = 0$ in $H^{2p^{r-\ell-1}}(GL_{N(\ell)}, \mathfrak{gl}_N^{(r)})$.
- (3) $e_{r-\ell} \in H^{2p^{r-\ell-1}}(GL_N, \mathfrak{gl}_N^{(r-\ell)})$ restricts via the “standard inclusion” $GL_{N-1} \subset GL_N$ to $e_{r-\ell} \in H^{2p^{r-\ell-1}}(GL_{N-1}, \mathfrak{gl}_N^{(r-\ell)})$.
- (4) For any root subgroup $E_{i,j} : \mathbb{G}_a \rightarrow GL_N$ (with $i < j$), the restriction of $e_{r-\ell}^{(\ell)}$ to $H^{2p^{r-\ell-1}}(\mathbb{G}_{a(1)}, \mathfrak{gl}_N^{(r)})$ equals $x_1^{p^{r-\ell-1}} \otimes X_{i,j}^{(r)}$, where $x_1 \in H^2(\mathbb{G}_{a(1)}, k)$ is the “canonical generator”.

Following [23] (specifically, the notation of the proof of Proposition 5.1 of [23]), we use the following notation: we identify $S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{gl}_N^{(r)}[2p^{r-\ell-1}]))$ with the algebra of functions on the affine space $\mathbb{A}^{rN^2} = \prod_{\ell=0}^{r-1} M_{n,n}$, identifying $X^{i,j}(\ell) \in \mathfrak{gl}_N^{(r)}[2p^{r-\ell-1}]$ with the (i, j) coordinate function of the ℓ -th factor.

For any affine group scheme G over k , we use the notation $V_r(G)$ for the affine scheme of 1-parameter subgroups of G of height r (i.e., homomorphisms $\mathbb{G}_{a(r)} \rightarrow G$ of group schemes over k) with coordinate algebra $k[V_r(G)]$ as in [23].

We state two theorems of Suslin-Friedlander-Bendel. The first originates from the observation in [23] that every infinitesimal 1-parameter subgroup $\psi : \mathbb{G}_{a(r)} \rightarrow GL_N$ is uniquely of the form

$$\exp \underline{B} \equiv \prod_{s=0}^{r-1} \exp_{B_s} \circ F^s : \mathbb{G}_{a(r)} \rightarrow GL_N, \quad t \mapsto \prod_{s=0}^{r-1} \exp(t^{p^s} \cdot B_s)$$

for some r -tuple $\underline{B} = (B_0, \dots, B_{r-1})$ of p -nilpotent, pair-wise commuting elements of \mathfrak{gl}_N .

Theorem 1.2. ([23, 5.1]) *The map $\phi_{GL_N, r}$ of (1.0.1) factors as*

(1.2.1)

$$\phi_{GL_N, r} = \bar{\phi}_{GL_N, r} \circ q : S^*(\oplus_{\ell=0}^{r-1} (\mathfrak{gl}_N^{\#(r)}[2p^{r-\ell-1}])) \twoheadrightarrow k[V_r(GL_N)] \rightarrow H^\bullet(GL_{N(r)}, k).$$

Here, q is the quotient by the ideal generated by the relations

$$\begin{aligned} R_{i,j,\ell,\ell'} &= \sum_t X^{i,t}(\ell) \cdot X^{t,j}(\ell') - X^{i,t}(\ell') \cdot X^{t,j}(\ell) \\ S_{i,j,\ell} &= \sum_{t_1, \dots, t_{p-1}} X^{i,t_1}(\ell) \cdot X^{t_1,t_2}(\ell) \cdots X^{t_{p-1},j}(\ell) \end{aligned}$$

for all i, j, ℓ, ℓ' as in [23, 5.1]. Thus, $V_r(GL_N)$ is identified with the k -scheme of r -tuples of p -nilpotent matrices (in view of relations $\{S_{i,j,\ell}\}$) which are pair-wise commuting (in view of relations $\{R_{i,j,\ell,\ell'}\}$).

Theorem 1.2 can be viewed as a complement (in the special case $G = GL_{N(r)}$) to the following theorem.

Theorem 1.3. [24, 5.2], [23, 5.2] *Fix some integer $r \geq 1$. Then for any infinitesimal group scheme H of height $\leq r$, there is a natural homomorphism of commutative k -algebras*

$$(1.3.1) \quad \psi : H^\bullet(H, k) \rightarrow k[V_r(H)]$$

whose kernel is nilpotent and whose image contains all p^r -th powers of elements of $k[V_r(H)]$. If $H = G_{(r)}$, the r -th Frobenius kernel of some linear algebraic group G , then we denote ψ by $\psi_{G,r} : H^\bullet(G_{(r)}, k) \rightarrow k[V_r(G)]$.

The map $\psi_{G,r}$ is G -equivariant.

In the special case of $G = GL_{N(r)}$, the composition

$$\psi_{GL_N, r} \circ \bar{\phi}_{GL_N, r} : k[V_r(GL_N)] \rightarrow k[V_r(GL_N)]$$

is the r -th iterate of the Frobenius map. In particular, $\psi_{GL_N, r}(\phi_{GL_N, r}(X^{i,j}(\ell))) = (X^{i,j}(\ell))^{p^r}$.

Remark 1.4. The assertion of Theorem 1.3 of G -equivariance of $\psi_{G,r}$ arises from the naturality of ψ , in particular the commutativity of the first displayed square of the proof of Theorem 1.14 of [23].

As shown in [23], $V_r(G)$ has a natural grading given by the monoid action of (right) composition by $V_r(\mathbb{G}_{a(r)})$ on $V_r(G)$; namely, one restricts this action of $\mathbb{A}^r \simeq V_r(\mathbb{G}_{a(r)})$ to the linear polynomials $\mathbb{A}^1 \subset \mathbb{A}^r$. With this grading, $X^{i,j}(\ell) \in k[V_r(GL_N)]$ has grading $p^{r-\ell-1}$ mapping via $\bar{\phi}_{GL_N, r}$ to a cohomology class of degree $2p^{r-\ell-1}$, then further mapping via $\psi_{GL_N, r}$ to $F^r(X^{i,j}(\ell))$ with degree $p^r \cdot p^{r-\ell-1}$.

We recall that a closed embedding $G \rightarrow GL_N$ of a linear algebraic group G is said to be of **exponential type** if the map of schemes $\exp : \mathbb{G}_a \times \mathcal{N}_p(\mathfrak{gl}_N) \rightarrow V_r(GL_N)$ given by the usual truncated exponential map restricts to $\mathcal{E} : \mathbb{G}_a \times \mathcal{N}_p(\mathfrak{g}) \rightarrow V_r(G)$. (As usual, for any p -restricted Lie algebra \mathfrak{g} , we denote the p -operator as $(-)^{[p]} : \mathfrak{g} \rightarrow \mathfrak{g}$ and we denote by $\mathcal{N}_p(\mathfrak{g}) \subset \mathfrak{g}$ the subvariety whose k points are elements $X \in \mathfrak{g}$ such that $X^{[p]} = 0$.) For any G equipped with such an embedding, every infinitesimal 1-parameter subgroup $\mathbb{G}_{a(r)} \rightarrow G$ is uniquely of the form $\prod_{s=0}^{r-1} \mathcal{E}_{B_s} \circ F^s$

for some $\underline{B} \in \mathcal{C}_r(\mathcal{N}_p(\mathfrak{g}))$ by [22, 2.5]; here, $\mathcal{C}_r(\mathcal{N}_p(\mathfrak{g}))$ is the variety of r -tuples (B_0, \dots, B_{r-1}) of p -nilpotent, pairwise commuting elements of \mathfrak{g} .

Proposition 1.5. *Let $i : G \rightarrow GL_N$ be a closed embedding of exponential type for some linear algebraic group G . Then the following square is a cartesian square of closed immersions*

$$(1.5.1) \quad \begin{array}{ccc} V_r(G) & \xrightarrow{i_V} & V_r(GL_N) \\ \downarrow & & \downarrow \\ \mathfrak{g}^{\times r} & \xrightarrow{i} & \mathfrak{gl}_N^{\times r}. \end{array}$$

In other words, we have a cocartesian square of quotient maps of k -algebras

$$(1.5.2) \quad \begin{array}{ccc} S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{gl}_N^{\#(r)}[2p^{r-\ell-1}])) & \xrightarrow{q_{GL_N, r}} & k[V_r(GL_N)] \\ \downarrow i^* & & \downarrow i_V^* \\ S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{g}^{\#(r)}[2p^{r-\ell-1}])) & \xrightarrow{q_{G, r}} & k[V_r(G)]. \end{array}$$

Proof. The condition that $i : G \rightarrow GL_N$ is of exponential type enables us to identify the embedding $V_r(G) \rightarrow V_r(GL_N)$ of schemes representing height r infinitesimal 1-parameter subgroups with the embedding of schemes of r -tuples of p -nilpotent, pair-wise commuting elements of respective Lie algebras. Using this, we verify that (1.5.1) arises as a cartesian square of representable functors. Namely, we verify that the defining relations $\{R_{i,j,\ell,\ell'}, S_{i,j,\ell}\}$ in $S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{gl}_N^{\#(r)}[2p^{r-\ell-1}]))$ (for $V_r(GL_N) \subset (\mathfrak{gl}_N)^{\times r}$) have image in $S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{g}^{\#(r)}[2p^{r-\ell-1}]))$ (i.e., restrictions to $(\mathfrak{g})^{\times r} \subset (\mathfrak{gl}_N)^{\times r}$) which generate defining relations for $V_r(G) \subset (\mathfrak{gl}_N)^{\times r}$. This follows from the observation for $X, Y \in \mathfrak{g}$ that the condition that X, Y commute in \mathfrak{g} is the same as the condition that their images commute in \mathfrak{gl}_N , and the condition that $X^{[p]} = 0$ is the condition that the image of X in \mathfrak{gl}_N has p -th power 0.

The cartesian square (1.5.1) is equivalent to the cocartesian square (1.5.2) of coordinate algebras thanks to the anti-equivalence of categories relating affine k -schemes and finitely generated commutative k -algebras. \square

We thank R Guralnick for explaining the following result of S. Garibaldi given in [14].

Proposition 1.6. [14, Prop 8.1] *If G is a simple algebraic group for which $p > 2$ is a very good prime (i.e., for type A_{n-1} , p does not divide n ; for type G_2, F_4, E_6, E_7 , $p > 3$; for type E_8 , $p > 5$), then there exist a closed embedding $i : G \rightarrow GL_N$ such that the induced map $i : \mathfrak{g} \rightarrow \mathfrak{gl}_N$ admits a unique G -equivariant splitting $\tau : \mathfrak{gl}_N \rightarrow \mathfrak{g}$.*

Garibaldi's result is proved for a split, almost simple algebraic group over an arbitrary field F and uses an embedding $G \subset GL_N$ associated to a representation defined over F . The splitting is also defined over F .

The following theorem can be interpreted as giving a lower bound on the “size” of $H^\bullet(G_{(r)}, k)$.

Theorem 1.7. *Let G be a simple algebraic group and assume that $p > 2$ is very good for G . Assume give some embedding $i : G \subset GL_N$ as in Proposition 1.6 which*

is also an embedding of exponential type. Let $\tau : \mathfrak{gl}_N \rightarrow \mathfrak{g} = \text{Lie}(G)$ be the unique G -equivariant splitting of $i : \mathfrak{g} \rightarrow \mathfrak{gl}_N$. Set $\phi_{G,r} = i^* \circ \phi_{GL_N,r} \circ \tau^*$. Then

$$\psi_{G,r} \circ \phi_{G,r} : S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{g}^{\#(r)}[2p^{r-\ell-1}])) \rightarrow H^\bullet(G_{(r)}, k) \rightarrow k[V_r(G)]$$

has image containing all p^r -th powers.

Consider the unipotent radical U_J of $P_J \subset G$ for some subset J of the set of fundamental positive roots Π of a chosen root system for G (given by a choice $B \subset G$ of Borel subgroup with maximal torus T), denote by $\tau_{U_J} : \mathfrak{g} \rightarrow \mathfrak{u}_J$ the unique T -equivariant splitting of $i_{U_J} : \mathfrak{u} \rightarrow \mathfrak{g}$, and set $\phi_{U_J,r} = i_{U_J}^* \circ \phi_{G,r} \circ \tau_{U_J}^*$. Then the composition

$$\psi_{U_J,r} \circ \phi_{U_J,r} : S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J^{\#(r)}[2p^{r-\ell-1}])) \rightarrow H^\bullet((U_J)_{(r)}, k) \rightarrow k[V_r(U_J)]$$

has image containing all p^r -th powers and is injective when restricted to each $\mathfrak{u}_J^{\#(r)}[2p^{r-\ell-1}]$.

Proof. Consider the following diagram, where F^r denotes the r -th power of the Frobenius map:
(1.7.1)

$$\begin{array}{ccccccc}
 & & \phi_{GL_N,r} & & F^r & & \\
 & \nearrow & & \searrow & & \nearrow & \\
 S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{gl}_N^{\#(r)}[2p^{r-\ell-1}])) & \xrightarrow{q_{GL_N,r}} & k[V_r(GL_N)] & \xrightarrow{\phi_{GL_N,r}} & H^\bullet(GL_{N(r)}, k) & \xrightarrow{\psi_{GL_N,r}} & k[V_r(GL_N)] \\
 \uparrow \tau^* & & \downarrow i_V^* & & \downarrow i^* & & \downarrow i_V^* \\
 S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{g}^{\#(r)}[2p^{r-\ell-1}])) & \xrightarrow{q_{G,r}} & k[V_r(G)] & & H^\bullet(G_{(r)}, k) & \xrightarrow{\psi_{G,r}} & k[V_r(G)] \\
 & \searrow & & \nearrow & & \searrow & \\
 & & \phi_{G,r} & & F^r & &
 \end{array}$$

Commutativity of the left square is a consequence of Proposition 1.5 and commutativity of the right square follows from the functoriality of ψ ; commutativity of the left rectangle follows from the definition of $\phi_{G,r}$, whereas commutativity of the right rectangle is a consequence of the functoriality of F^r for maps of varieties defined over \mathbb{F}_{p^r} (granted that Garibaldi's splitting is defined over \mathbb{F}_p , as observed after the statement of Proposition 1.6).

To show that $\psi_{G,r} \circ \phi_{G,r}$ has image containing all p^r -th powers, we observe that $F^r : k[V_r(G)] \rightarrow k[V_r(G)]$ has image containing all p^r -th powers. Thus, a simple diagram chase around the commutative diagram (1.7.1) using the surjectivity of $q_{G,r}$ verifies this assertion.

Consider now the following diagram obtained by replacing $i : G \rightarrow GL_N$ in (1.7.1) by $i_{U_J} : U_J \rightarrow G$:
(1.7.2)

$$\begin{array}{ccccc}
& \xrightarrow{\phi_{G,r}} & & \xrightarrow{F^r} & \\
S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{g}^{\#(r)}[2p^{r-\ell-1}])) & \xrightarrow{q_{G,r}} & k[V_r(G)] & \xrightarrow{\psi_{G,r}} & H^\bullet(G_{(r)}, k) \xrightarrow{\psi_{G,r}} k[V_r(G)] \\
\uparrow \tau_{U_J}^* & & \downarrow i_V^* & & \downarrow i_V^* \\
S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J^{\#(r)}[2p^{r-\ell-1}])) & \xrightarrow{q_{U_J,r}} & k[V_r(U_J)] & \xrightarrow{\psi_{U_J,r}} & H^\bullet((U_J)_{(r)}, k) \xrightarrow{\psi_{U_J,r}} k[V_r(U_J)] \\
& \xrightarrow{\phi_{U_J,r}} & & \xrightarrow{F^r} &
\end{array}$$

The previous argument applies with only notational changes to verify the corresponding assertions for $\psi_{U_J,r} \circ \phi_{U_J,r}$. \square

Example 1.8. As stated in [23, 1.8], the classical simple algebraic groups Sp_{2n} , SO_n and SL_n admit embeddings of exponential type. Namely, one considers a vector space (of dimension $2n$ for Sp_{2n} , of dimension n for O_n) equipped with a non-degenerate bilinear form and one takes the embedding given by considering those linear isomorphisms preserving the form. These embeddings $i : G \rightarrow GL_N$ are defined over \mathbb{F}_p and also satisfy the condition that $i : \mathfrak{gl}_N \rightarrow \mathfrak{g}$ admits a (unique) G -equivariant splitting.

For $U_J \subset GL_N$, we have the following natural strengthening of Theorem 1.7. We denote by $T_N \subset GL_N$ the maximal torus of diagonal matrices.

Proposition 1.9. *Let $i_{U_J} : U_J \rightarrow GL_N$ be the inclusion of the unipotent radical of a parabolic subgroup of GL_N . Then we have the following T_N -equivariant, commutative diagram, a stronger version of the diagram obtained from (1.7.1):*
(1.9.1)

$$\begin{array}{ccccc}
& \xrightarrow{\phi_{GL_N,r}} & & \xrightarrow{F^r} & \\
S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{gl}_N^{\#(r)}[2p^{r-\ell-1}])) & \xrightarrow{q_{GL_N,r}} & k[V_r(GL_N)] & \xrightarrow{\psi_{GL_N,r}} & H^\bullet(GL_N(r), k) \xrightarrow{\psi_{GL_N,r}} k[V_r(GL_N)] \\
\downarrow i_{U_N}^* & & \downarrow i_V^* & & \downarrow i_V^* \\
S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J^{\#(r)}[2p^{r-\ell-1}])) & \xrightarrow{q_{U_J,r}} & k[V_r(U_J)] & \xrightarrow{\psi_{U_J,r}} & H^\bullet((U_J)_{(r)}, k) \xrightarrow{\psi_{U_J,r}} k[V_r(U_J)] \\
& \xrightarrow{\phi_{U_J,r}} & & \xrightarrow{F^r} &
\end{array}$$

In particular, any element in the kernel of $\bar{\phi}_{U_J,r}$ has p^r -th power 0.

Proof. Observe that the T -weights of $H^\bullet((U_J)_{(r)}, k)$ are all negatives of roots in the lattice generated by the roots of U_N , so that $i_{U_J}^* \circ \phi_{GL_N,r}$ restricted to each $\mathfrak{u}_J^{\#(r)}[2p^{r-\ell-1}]$ must factor as a T -equivariant map through the projection $\mathfrak{gl}_N^{\#(r)}[2p^{r-\ell-1}] \rightarrow \mathfrak{u}_J^{\#(r)}[2p^{r-\ell-1}]$. This determines the map $\phi_{U_J,r}$ making commutative the left rectangle (i.e., double square) of (1.9.1). In view of the surjectivity of $q_{GL_N,r}$, the map $\bar{\phi}_{U_J,r}$ is uniquely determined making the middle square commute.

Recall that the composition of F with the geometric Frobenius is the p -th power map. This implies that any element in the kernel of F^r , and thus any element in the kernel of $\phi_{U,r}$, has p^r -th power 0. \square

The next proposition helps to pin down $\phi_{G,r}$.

Proposition 1.10. *Let $i_{U_J} : U_J \rightarrow GL_N$ be the inclusion of the unipotent radical of a parabolic subgroup of GL_N . Denote by $\Gamma_2 \subset U_J$ the commutator subgroup of U_J with Lie algebra γ_2 , and denote by $q : U_J \rightarrow U_J/\Gamma_2$ the projection. Then the following square commutes for each ℓ , $0 \leq \ell < r$:*

$$(1.10.1) \quad \begin{array}{ccccc} \mathfrak{u}_J^{\#(r)}[2p^{r-\ell-1}] & \xrightarrow{=} & \mathfrak{u}_J^{\#(r)}[2p^{r-\ell-1}] & \xrightarrow{\phi_{U_J,r}} & H^{2p^{r-\ell-1}}((U_J)_{(r)}, k) \\ \uparrow q^* & & & & \uparrow q^* \\ (\mathfrak{u}_J/\gamma_2)^{\#(r)}[2p^{r-\ell-1}] & \longrightarrow & S^{p^{r-\ell-1}}((\mathfrak{u}_J/\gamma_2)^{\#(\ell+1)}[2]) & \longrightarrow & H^{2p^{r-\ell-1}}((U_J/\Gamma_2)_{(r)}, k). \\ & \searrow \phi_{U_J/\Gamma_2,r} & & & \end{array}$$

The lower left map of (1.10.1) is a special case (for $V = (\mathfrak{u}_J/\gamma_2)^{\#(\ell+1)}$) of the natural embedding $V^{(r-\ell-1)} \subset S^{p^{r-\ell-1}}(V)$ sending $v \in V$ to its $p^{r-\ell-1}$ -st power. We have abused notation by using $\phi_{U_J,r}$ to denote the restriction to the indicated summand of the map of (1.9.1) with this name.

For a minimal weight α of \mathfrak{u}_J , the map $\phi_{U/\Gamma_2,r}$ restricted to the weight space $k \cdot X^\alpha(\ell)$ of weight $p^r \alpha$ is explicitly described as the projection $(\mathfrak{u}_J/\gamma_2)^{\#(r)}[2p^{r-\ell-1}] \rightarrow k \cdot X^\alpha(\ell)$, followed by the map sending $X^\alpha(\ell)$ to the $p^{r-\ell-1}$ -st power of the natural class in $H^2(\mathbb{G}_{a(r)}, k)$, followed by the inflation map $H^*(\mathbb{G}_{a(r)}, k) \rightarrow H^*((U_J/\Gamma_2)_{(r)}, k)$ induced by the quotient map $U_J/\Gamma_2 \rightarrow \mathbb{G}_a$ onto the root subgroup indexed by the root α .

Proof. We first observe the commutativity of

$$(1.10.2) \quad \begin{array}{ccc} \mathfrak{g}_N^{\#(r)}[2p^{r-1}] & \xrightarrow{\phi_{GL_N,r}} & H^{2p^{r-1}}(GL_{N(r)}, k) \\ \downarrow i_{U_J}^* & & \downarrow i_{U_J}^* \\ \mathfrak{u}_J^{\#(r)}[2p^{r-1}] & \xrightarrow{\phi_{U_J,r}} & H^{2p^{r-1}}((U_J)_{(r)}, k) \\ \uparrow q_\alpha^* & & \uparrow q_\alpha^* \\ (\mathfrak{e}_\alpha)^{\#(r)}[2p^{r-1}] & \xrightarrow{\phi_{E_\alpha,r}} & H^{2p^{r-1}}((E_\alpha)_{(r)}, k) \end{array}$$

where $\mathbb{G}_a \simeq E_\alpha \subset U_J$ is the root subgroup associated to a minimal root α and $q_\alpha : U_J/\Gamma_2 \rightarrow E_\alpha$ is the projection, a group homomorphism since α is minimal. This commutativity of (1.10.2) follows from Proposition 1.1(4) and the fact $i_\alpha : \mathbb{G}_a \rightarrow U_J/\Gamma_2$ is left inverse to q_α . This implies the commutativity of (1.10.1) for $\ell = 0$

For $\ell > 0$, we apply (1.10.2) with r replaced by $r - \ell$ and use the commutativity of

$$(1.10.3) \quad \begin{array}{ccc} \mathbf{u}_J^{\#(r-\ell)}[2p^{r-\ell-1}] & \longrightarrow & H^{2p^{r-\ell-1}}((U_J)_{(r-\ell)}, k) \\ \downarrow (-)^\ell & & \downarrow \ell_* \\ \mathbf{u}_J^{\#(r)}[2p^{r-1}] & \longrightarrow & H^{2p^{r-1}}((U_J)_{(r)}, k) \end{array}$$

which is a direct consequence of the definition of $\phi_{GL_N, r}$. \square

2. STABILIZATION OF $H^\bullet(U_{(r)}, k)$ WITH RESPECT TO r

Part of the author's motivation for considering $H^*(U_{(r)}, k)$ was the hope that some form of “continuous cohomology” for the unipotent algebraic group U would prove useful in the study of the (rational) representations of U . This requires understanding the limiting behavior of $H^*(U_{(r)}, k)$ as r increases. Earlier computational information for $H^*(U_{(r)}, k)$ (especially in [23], [24]) shed little if any light on this limiting behavior.

We begin by recalling the following spectral sequence formulated by H. Andersen and J. Jantzen.

Proposition 2.1. [1], [16, §9] *Let H be an irreducible affine group scheme (over k) and let $I_1 \subset k[H]$ denote the maximal ideal at the identity of H . The filtration of $k[H]$ by powers of I_1 leads to an associated graded Hopf algebra which is the coordinate algebra of the vector group scheme $\text{gr}(H)$. For any rational H -module M , there is a naturally associated convergent spectral sequence*

$$(2.1.1) \quad {}^{AJ}E_1^{i,j}(H) = H^{i+j}(\text{gr}(H), k)_i \otimes M \Rightarrow H^{i+j}(H, M),$$

where $H^*(\text{gr}(H), k)_i$ is the cohomology algebra of the i^{th} graded summand of the Hochschild complex of $\text{gr}(H)$.

If G is a irreducible linear algebraic group (hence, a reduced affine group scheme of finite type over k) and $p \neq 2$, then ${}^{AJ}E_1^{i,j}(G)$ can be identified with the direct sum of tensor products of the form

$$(2.1.2) \quad S^{a_1}(\mathfrak{g}^{\#(1)}[2]) \otimes S^{a_2}(\mathfrak{g}^{\#(2)}[2]) \otimes \cdots \otimes \Lambda^{b_1}(\mathfrak{g}^{\#(1)}[1]) \otimes \Lambda^{b_2}(\mathfrak{g}^{\#(1)}[1]) \otimes \cdots$$

where the sum is over all sequences $\{a_n\}, \{b_n\}$ with each $a_n \geq 0$, each $b_n \geq 0$ and

$$i = \sum_{n \geq 1} (a_n p^n + b_n p^{n-1}), \quad i + j = \sum_{n \geq 1} (2a_n + b_n).$$

Moreover, for any $r \geq 1$, ${}^{AJ}E_1^{i,j}(G_{(r)})$ can be identified with the direct sum of those tensor products of the form (2.1.2) with $a_n = b_n = 0$, $n > r$.

The following theorem shows for a large class of unipotent algebraic groups that the rational cohomology equals the “continuous cohomology”, thereby motivating the study of this continuous cohomology.

Theorem 2.2. *Let G be a simple algebraic group provided with a choice of Borel subgroup $B \subset G$ with maximal torus T and unipotent radical U . Let $U_1 \subset U$ be*

T -stable closed subgroup, $U_2 \subset U_1$ a T -stable, normal closed subgroup of U_1 , and consider $V \equiv U_1/U_2$. The natural map

$$H^*(V, k) \rightarrow \varprojlim_r H^*(V_{(r)}, k)$$

is an isomorphism.

Proof. Let $0 \neq \zeta \in H^d(V, k)$ be a T -eigenvector of weight $\omega = \sum_{m=1}^{\ell} w_m \alpha_m$, $w_m \geq 0$, an element in the positive cone of the root lattice for $\mathfrak{v} = \text{Lie}(V)$; here, $\alpha_1, \dots, \alpha_{\ell}$ are the simple roots determined by $B \subset G$. One verifies by inspection that the restriction map ${}^{AJ}E_1^{*,*}(V) \rightarrow {}^{AJ}E_1^{*,*}(V_{(r)})$ is an isomorphism of ω -weight spaces $({}^{AJ}E_1^{*,*}(V))_{\omega} \rightarrow ({}^{AJ}E_1^{*,*}(V_{(r)}))_{\omega}$ provided that p^{r-1} is greater than any of w_1, \dots, w_{ℓ} . Namely, using (2.1.2), we can see that this condition on each w_i implies that there can be no contribution from a tensor factor in $S^{a_i}(\mathfrak{v}^{\#(i)}[2])$ or in $\Lambda^{b_j}(\mathfrak{v}^{\#(j-1)}[1])$ for $i > r$ or for $j > r + 1$. Observe that the spectral sequences $\{{}^{AJ}E_s^{i,j}(V), s \geq 1\}$ and $\{{}^{AJ}E_s^{i,j}(V_{(r)}), s \geq 1\}$ split (additively) as a direct sum of spectral sequences indexed by the weights in the positive cone of the root lattice for \mathfrak{v} .

This implies that the restriction map induces an isomorphism $(H^*(V, k))_{\omega} \rightarrow (H^*(V_{(r)}, k))_{\omega}$ whenever p^{r-1} is greater than $\max\{w_i\}$, the maximum of w_1, \dots, w_{ℓ} in the expression for ω . Thus, the restriction map $(H^*(V, k))_{\omega} \rightarrow \varprojlim_r (H^*(V_{(r)}, k))_{\omega}$ is an isomorphism for all weights ω in the positive cone of the root lattice for \mathfrak{v} , so that $H^*(V, k) \rightarrow \varprojlim_r H^*(V_{(r)}, k)$ is also an isomorphism. \square

We easily verify that the injectivity statement of Theorem 2.2 extends to cohomology of V with coefficients in a finite dimensional V -module having a compatible torus action.

Corollary 2.3. *Retain the notation of Theorem 2.2 and let M be a finite dimensional rational $V \rtimes T$ -module. Then the natural map*

$$H^*(V, M) \rightarrow \varprojlim_r H^*(V_{(r)}, M)$$

is injective.

Moreover, if $\zeta_r \in H^d(V_{(r)}, M)$ is the restriction of some $\zeta_s \in H^d(V_{(s)}, M)$ for all $s \geq r$, then there exists some $\zeta \in H^d(V, M)$ which restricts to ζ_r .

Proof. We repeat the argument of the proof of Theorem 2.2 for A-J spectral sequences $\{{}^{AJ}E_s^{i,j}(V, M), s \geq 1\}$ and $\{{}^{AJ}E_s^{i,j}(V_{(r)}, M), s \geq 1\}$. The action of V on the associated graded group of M is trivial, so that the E_1 terms are obtained from those for coefficients equal to k by tensoring with M . These spectral sequences now split as a direct sum of spectral sequences indexed by weights given as the sum of a weight of M and an element in the positive cone of the root lattice for \mathfrak{v} . As in the proof of Theorem 2.2, we conclude that the restriction map ${}^{AJ}E_1^{*,*}(V, M) \rightarrow {}^{AJ}E_1^{*,*}(V_{(r)}, M)$ is an isomorphism of ω -weight spaces $({}^{AJ}E_1^{*,*}(V, M))_{\omega} \rightarrow ({}^{AJ}E_1^{*,*}(V_{(r)}, M))_{\omega}$ provided that p^{r-1} does not divide the coefficient of the linear expansion of any weight of the form $w + w'$, where w' is a weight of M . The remainder of the proof is a repetition of that of Theorem 2.2. \square

In order to investigate how the map $\phi_{GL_N, r}$ in (1.0.1) behaves as r increases, we introduce in the next proposition the map ρ .

Proposition 2.4. *Define the degree preserving map of graded k -algebras*

$$(2.4.1) \quad \rho : S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{gl}_N^{(r)\#}[2p^{r-\ell-1}])) \rightarrow S^*(\oplus_{\ell=0}^{r-2}(\mathfrak{gl}_N^{(r-1)\#}[2p^{r-\ell-2}]))$$

by sending $X^{s,t}(\ell) \in \mathfrak{gl}_N^{(r)\#}[2p^{r-\ell-1}]$ to the p -th power $(X^{s,t}(\ell))^p \in S^p(\mathfrak{gl}_N^{(r-1)\#}[2p^{r-\ell-2}])$ if $\ell < r-1$ and to 0 if $\ell = r-1$. Then ρ fits in the GL_N -equivariant commutative square

$$(2.4.2) \quad \begin{array}{ccc} S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{gl}_N^{(r)\#}[2p^{r-\ell-1}])) & \xrightarrow{\phi_{GL_N,r}} & H^\bullet(GL_{N(r)}, k) \\ \rho \downarrow & & \downarrow res \\ S^*(\oplus_{\ell=0}^{r-2}(\mathfrak{gl}_N^{(r-1)\#}[2p^{r-\ell-2}])) & \xrightarrow{\phi_{GL_N,r-1}} & H^\bullet(GL_{N(r-1)}, k). \end{array}$$

Proof. We first show that $\phi_{GL_N,r}(X^{s,t}(0)) \in H^{2p^{r-1}}(GL_{N(r)}, k)$ restricts to the p -th power of $\phi_{GL_N,r-1}(X^{s,t}(0)) \in H^{2p^{r-2}}(GL_{N(r-1)}, k)$. By [23, 3.4], both of these classes restrict to the p^{r-1} -st power of the image of $\phi_{GL_N,1}(X^{s,t}(0))$ in $H^2(GL_{N(1)}, k)$. Thus, the outer rectangle and the lower square of the following diagram commutes:

$$(2.4.3) \quad \begin{array}{ccc} \mathfrak{gl}_N^{(r)\#}[2p^{r-1}] & \xrightarrow{e_r} & H^{2p^{r-1}}(GL_{N(r)}, k) \\ \rho \downarrow & & \downarrow res \\ S^p(\mathfrak{gl}_N^{(r-1)\#}[2p^{r-2}]) & \xrightarrow{e_{r-1}} & H^{2p^{r-1}}(GL_{N(r-1)}, k) \\ (-)^{\rho^{r-2}} \downarrow & & \downarrow res \\ S^{p^{r-1}}(\mathfrak{gl}_N^{(1)\#}[2]) & \xrightarrow{e_1} & H^{2p^{r-1}}(GL_{N(1)}, k) \end{array}$$

The images in $H^{2p^{r-1}}(GL_{N(r-1)}, k)$ of the two compositions in the upper square of (2.4.3) are each irreducible GL_N -modules (copies of $(\mathfrak{gl}_N^{(r)\#}[2p^{r-1}])/(k \cdot Tr^{(r)})$) which restrict non-trivially to $H^{2p^{r-1}}(GL_{N(1)}, k)$. Using the form of the E_1 -term of the A-J spectral sequence (2.1.1), we conclude that there is a unique copy of $(\mathfrak{gl}_N^{(r)\#}[2p^{r-1}])/(k \cdot Tr^{(r)})$ in ${}^{AJ}E_1^{*,*}(GL_{N(r-1)})$ of cohomological degree $2p^{r-1}$, so that these images are equal. The functoriality of (2.1.1) with respect to $GL_{N(1)} \rightarrow GL_{N(r)}$ now implies that the upper square of (2.4.3) must also commute.

By definition of $e_{r-\ell}^{(\ell)}$ as the pull-back via $F^\ell : GL_N \rightarrow GL_N$ of $e_{r-\ell}$, we have the commutativity of the following square

$$(2.4.4) \quad \begin{array}{ccc} \mathfrak{gl}_N^{(r-\ell)\#}[2p^{r-\ell-1}] & \xrightarrow{e_{r-\ell}} & H^{2p^{r-\ell-1}}(GL_{N(r-\ell)}, k) \\ (-)^{(\ell)} \downarrow & & \downarrow (F^\ell)^* \\ \mathfrak{gl}_N^{(r)\#}[2p^{r-\ell-1}] & \xrightarrow{e_{r-\ell}^{(\ell)}} & H^{2p^{r-\ell-1}}(GL_{N(r)}, k). \end{array}$$

Consequently, pulling back via F^ℓ the commutative upper square of (2.4.3) with r replaced by $r-\ell$ determines the following commutative square for each $\ell, 0 \leq \ell < r$:

$$(2.4.5) \quad \begin{array}{ccc} \mathfrak{gl}_N^{\#(r)}[2p^{r-\ell-1}] & \xrightarrow{e_{r-\ell}^{(\ell)}} & H^{2p^{r-\ell-1}}(GL_{N(r)}, k) \\ \rho \downarrow & & \downarrow res \\ S^p(\mathfrak{gl}_N^{\#(r-1)}[2p^{r-\ell-2}]) & \xrightarrow{e_{r-\ell-1}^{(\ell)}} & H^{2p^{r-\ell-1}}(GL_{N(r-1)}, k). \end{array}$$

The proposition now follows since the maps of (2.4.2) are maps of k -algebras and the commutativity of (2.4.5) implies the commutativity of (2.4.2) on generators. \square

The proof of the following lemma (including the definitions of ρ_G and ρ_{U_J}) is immediate from the definitions.

Lemma 2.5. *Let $U_J \subset G$ satisfy the hypotheses of Theorem 1.7. Define*

$$(2.5.1) \quad \rho_G : S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{g}^{\#(r)}[2p^{r-\ell-1}])) \rightarrow S^*(\oplus_{\ell=0}^{r-2}(\mathfrak{g}^{\#(r-1)}[2p^{r-\ell-2}]))$$

by sending $X \in \mathfrak{g}^{\#(r)}[2p^{r-\ell-1}]$ to $X^p \in S^p(\mathfrak{g}^{\#(r-1)}[2p^{r-\ell-2}])$ if $\ell < r-1$ and to 0 if $\ell = r-1$. Define $\rho_{U_J} : S^(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J^{\#(r)}[2p^{r-\ell-1}])) \rightarrow S^*(\oplus_{\ell=0}^{r-2}(\mathfrak{u}_J^{\#(r-1)}[2p^{r-\ell-2}]))$ similarly. Then ρ, ρ_G, ρ_{U_J} fit in the following commutative diagram*

$$(2.5.2) \quad \begin{array}{ccccc} S^*(\oplus_{\ell=0}^r(\mathfrak{u}_J^{\#(r)}[2p^{r-\ell-1}])) & \xrightarrow{\tau_{U_J}^*} & S^*(\oplus_{\ell=0}^r(\mathfrak{g}^{\#(r)}[2p^{r-\ell-1}])) & \xrightarrow{\tau_G^*} & S^*(\oplus_{\ell=0}^r(\mathfrak{gl}_N^{\#(r)}[2p^{r-\ell-1}])) \\ \downarrow \rho_{U_J} & & \downarrow \rho_G & & \downarrow \rho \\ S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J^{\#(r-1)}[2p^{r-\ell-2}])) & \xrightarrow{\tau_{U_J}^*} & S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{g}^{\#(r-1)}[2p^{r-\ell-2}])) & \xrightarrow{\tau_G^*} & S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{gl}_N^{\#(r-1)}[2p^{r-\ell-2}])). \end{array}$$

The following extension of Proposition 2.4 to G and U_J now follows easily.

Proposition 2.6. *Retain the hypotheses and notation of Theorem 1.7. Then the following squares commute*

$$(2.6.1) \quad \begin{array}{ccc} S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{g}^{\#(r)}[2p^{r-\ell-1}])) & \xrightarrow{\phi_{G,r}} & H^\bullet(G_{(r)}, k) \\ \rho_G \downarrow & & \downarrow res \\ S^*(\oplus_{\ell=0}^{r-2}(\mathfrak{g}^{\#(r-1)}[2p^{r-\ell-2}])) & \xrightarrow{\phi_{G,r-1}} & H^\bullet(G_{(r-1)}, k). \end{array}$$

$$(2.6.2) \quad \begin{array}{ccc} S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J^{\#(r)}[2p^{r-\ell-1}])) & \xrightarrow{\phi_{U_J,r}} & H^\bullet((U_J)_{(r)}, k) \\ \rho_{U_J} \downarrow & & \downarrow res \\ S^*(\oplus_{\ell=0}^{r-2}(\mathfrak{u}_J^{\#(r-1)}[2p^{r-\ell-2}])) & \xrightarrow{\phi_{U_J,r-1}} & H^\bullet((U_J)_{(r-1)}, k). \end{array}$$

Proof. We readily verify that $\phi_{G,r}, \phi_{G,r-1}$ provide two faces of a commutative “cube” with three other faces provided by (2.4.2), the right square of (2.5.2), and the square resulting from the functoriality of the restriction map with respect to $G \rightarrow GL_N$. The remaining face is (2.6.1).

The commutativity of (2.6.2) is verified similarly, replacing $G \rightarrow GL_N$ by $U_J \rightarrow G$. \square

As summarized in [16, I.9.9], the natural map $H^*(G, M) \rightarrow \varprojlim_r H^*((G_{(r)}, M))$ is an isomorphism for all finite dimensional G -modules M ; a key ingredient in the proof of that isomorphism is the fact that $H^*(G, k)$ is isomorphic to k (in degree 0). Although $H^*(U_J, k)$ is non-trivial, we do have the following vanishing result.

Corollary 2.7. *Retain the hypotheses and notation of Theorem 1.7. Then the sub-algebra*

$$\varprojlim_r \text{im}\{\phi_{U_J, r} : S^*(\oplus_{\ell=0}^{r-1}(\mathbf{u}_J^{\#(r)}[2p^{r-\ell-1}])) \rightarrow H^*((U_J)_{(r)}, k)\}$$

of $\varprojlim_r H^*((U_J)_{(r)}, k) \simeq H^*(U_J, k)$ is isomorphic to k (in degree 0).

Proof. The left exactness of \varprojlim_r implies that

$$\varprojlim_r \text{im}\{\phi_{U_J, r} : S^*(\oplus_{\ell=0}^{r-1}(\mathbf{u}_J^{\#(r)}[2p^{r-\ell-1}])) \rightarrow H^*((U_J)_{(r)}, k)\}$$

is a subalgebra of $\varprojlim_r H^*((U_J)_{(r)}, k)$ which is isomorphic to $H^*(U_J, k)$ by Theorem 2.2.

By the commutativity of (2.4.2), (2.6.1), and (2.6.2), we observe that no homogeneous element of degree d , $0 < d < p^s$, in $S^*(\oplus_{\ell=0}^{r-1}(\mathbf{u}_J^{\#(r)}[2p^{r-\ell-1}]))$ lies in the image of ρ^s for any $r > s$. Consequently, there can not exist some non-zero element $\{x_r\} \in \varprojlim_r \text{im}\{\phi_{U_J, r} : S^*(\oplus_{\ell=0}^{r-1}(\mathbf{u}_J^{\#(r)}[2p^{r-\ell-1}])) \rightarrow H^*((U_J)_{(r)}, k)\}$ of degree d : such an element $\{x_r\}$ in the inverse limit must satisfy $\rho_{r'-r}(x_{r'}) = x_r$ for all $r' \geq r \geq 0$ and would have to have some $x_r \neq 0$, $r > s$; then $\{x_r\}$ would have to satisfy $x_{r'} \neq 0$ for all $r' \geq r$ (including $r' = r + s$), contradicting the above observation. \square

As we shall see in the next section, the image of the composition

$$\varprojlim_r S^*(\oplus_{\ell=0}^{r-1}((\mathbf{u}_3/\gamma_2)^{\#(\ell+1)}[2])) \rightarrow \varprojlim_r H^*((U_3/\Gamma_2)_{(r)}, k) \rightarrow \varprojlim_r H^*((U_3)_{(r)}, k)$$

(which factors through $\varprojlim_r S^*(\oplus_{\ell=0}^{r-1}(\mathbf{u}_3^{\#(\ell+1)}[2])) \rightarrow \varprojlim_r H^*((U_3)_{(r)}, k)$) is highly non-trivial, where $\mathbb{G}_a \simeq \Gamma_2 \subset U_3$ is the center of U_3 .

3. THE MAP $\bar{\eta}_{U_J/\Gamma_3, r} : \bar{S}^*((U_J/\Gamma_3)_{(r)}) \rightarrow H^\bullet((U_J/\Gamma_3)_{(r)}, k)$

In this section, we consider groups of the form U_J/Γ_3 , where $\Gamma_3 \equiv \Gamma_3(U_J)$ is the third stage of the descending central series of the unipotent radical of a parabolic subgroup $P_J \subset G$ of a simple algebraic group G . Our primary tool is the Lyndon-Hochschild-Serre spectral sequence for a central extension (see Proposition 3.2) together with the action of the mod- p Steenrod algebra on this spectral sequence.

We recall from [3] the description due to H. Azad, M. Barry, and G. Seitz of the terms of the descending central series of the unipotent radical U_J of $P_J \subset G$ for some subset J of the set of fundamental positive roots Π of a chosen root system for G . For a positive root $\beta \in \Sigma^+ - \Sigma_J^+$ (where Σ^+ is the set of positive roots for the root system of $G \supset B \supset T$ and Σ_J^+ is the set of positive roots for the root system of $L_J = P_J/U_J \supset T_J$), we adopt the terminology of [3]: write $\beta = \beta_J + \beta_{J'}$ where β_J is a sum $\sum_i c_i \alpha_i$ with each $\alpha_i \in J$ and $\beta_{J'}$ is a sum $\sum_j d_j \alpha_j$ with each $\alpha_j \in \Pi - J$; then the height of β is defined to be $\sum_i c_i + \sum_j d_j$, the level of β is defined to be $\sum_j d_j$ and the shape of β is defined to be $\beta_{J'}$.

Proposition 3.1. (summary of §2 of [3]) *Let G be a simple algebraic group of adjoint type, and $P = P_J \subset G$ a parabolic subgroup, L_J its Levi factor, and U_J its unipotent radical for some subset $J \subset \Pi$. As usual, assume $p > 2$; for G of type G_2 , assume $p > 3$. Consider the descending central series for U_J :*

$$\cdots \subset \Gamma_{v+1} = [U_J, \Gamma_v] \subset \cdots \subset \Gamma_2 = [U_J, U_J] \subset \Gamma_1 = U_J.$$

For any $v > 1$, we have the central extension with a natural action of L_J :

$$(3.1.1) \quad 1 \rightarrow \Gamma_v/\Gamma_{v+1} \rightarrow U_J/\Gamma_{v+1} \rightarrow U_J/\Gamma_v \rightarrow 1.$$

The commutative group Γ_v/Γ_{v+1} is a direct product of irreducible L_J -modules $V_{\mathcal{S}}$ indexed by shapes \mathcal{S} of level v ; each $V_{\mathcal{S}}$ is T isomorphic to a product of $U_{-\beta}$ indexed by $\beta \in \Sigma^+ - \Sigma_J^+$ of shape \mathcal{S} and level v , where $U_{-\beta}$ is the root subgroup with T -weight $-\beta$; $V_{\mathcal{S}}$ is a high weight L_J -module with highest weight $-\beta_{\mathcal{S}}^o$, where $\beta_{\mathcal{S}}^o$ is the unique root of minimal height and shape \mathcal{S} .

In particular, if P_J is the minimal parabolic (i.e., equal to the given Borel subgroup $B \subset G$, corresponding to $J = \emptyset$), then Γ_v/Γ_{v+1} is T -isomorphic to $\prod U_{-\beta}$ where the product is indexed by $\beta \in \Sigma^+$ of level v .

In what follows, we shall denote $\text{Lie}(\Gamma_v(U_J))$ by γ_v . As in Section 1, r will denote a fixed (but arbitrary) positive integer.

In order to fix notation and T -weights for Frobenius twists, we recall the known computation of $H^*(\mathbb{G}_a, k)$ and $H^*(\mathbb{G}_{a(r)}, k)$ (see, for example, [16, Ch I.4] or [7]): there is a natural isomorphism of graded commutative k -algebras

$$(3.1.2) \quad S^*(V^{(1)}[2]) \otimes \Lambda^*(V[1]) \xrightarrow{\sim} H^*(\mathbb{G}_a, k),$$

where $V = H^1(\mathbb{G}_a, k)$ is a countable k -vector space spanned by $\lambda_1, \lambda_2, \dots, \lambda_s, \dots$ with Frobenius action $F^*(\lambda_s) = \lambda_{s+1}$, $\Lambda^*(V[1])$ is the exterior algebra on V placed in degree 1, and $S^*(V^{(1)}[2])$ is the polynomial algebra on the Frobenius twist $V^{(1)}$ of V placed in degree 2 spanned by $x_1, x_2, \dots, x_s, \dots$, the Bocksteins of the λ_i . The action of multiplication by $c \in k$ on \mathbb{G}_a induces an action on $H^*(\mathbb{G}_a, k)$ given by $c^*(\lambda_i) = c^{p^i-1} \lambda_i$, $c^*(x_i) = c^{p^i} x_i$. This indexing is that of [7] and [24, Thm 1.3]. We recall that the cohomology algebra $H^*(\mathbb{G}_{a(r)}, k)$ of the r^{th} Frobenius kernel $\mathbb{G}_{a(r)}$ of \mathbb{G}_a can be identified with the quotient of $H^*(\mathbb{G}_a, k)$ obtained by setting $\lambda_s = 0 = x_s$ for $s > r$. This finitely generated cohomology algebra admits the natural action of the mod- p Steenrod algebra \mathcal{A}_p (see [24, 1.7] for an explicit description of the action of the generators $\mathcal{P}^i, \beta \mathcal{P}^i$ of \mathcal{A}_p on $H^*(\mathbb{G}_a, k)$).

Observe that U_J/Γ_2 is a product $\mathbb{G}_a^{\times s}$ of copies of \mathbb{G}_a , so that its cohomology and that of $(\mathbb{G}_a^{\times s})_{(r)}$ are determined by the above computation and the Künneth theorem. In particular, we conclude that there is a natural injective map

$$(3.1.3) \quad \eta_{U_J/\Gamma_2, r} : S^*(\oplus_{\ell=0}^{r-1} (\mathbf{u}_J/\gamma_2)^{\#(\ell+1)}[2]) \rightarrow H^*((U_J/\Gamma_2)_{(r)}, k)$$

with left inverse $H^*((U_J/\Gamma_2)_{(r)}, k) \rightarrow S^*(\oplus_{\ell=0}^{r-1} (\mathbf{u}_J/\gamma_2)^{\#(\ell+1)}[2])$ whose kernel consists of elements with p -th power 0.

We designate T -eigenvector generators for

$$H^*(U_J/\Gamma_2, k) \simeq S^*(\oplus_{\ell=0}^{\infty} (\mathbf{u}_J/\gamma_2)^{\#(\ell+1)}[2]) \otimes \Lambda^*(\oplus_{\ell=0}^{\infty} (\mathbf{u}_J/\gamma_2)^{\#(\ell)}[1])$$

by $x_{\alpha}^{(\ell)}$ of cohomological degree 2 and T -weight $p^{\ell+1}\alpha$ and $y_{\alpha}^{(\ell)}$ of cohomological degree 1 and T -weight $p^{\ell}\alpha$; here, α ranges over roots of U_J of level 1 and ℓ is a non-negative integer satisfying $0 \leq \ell$. Similarly, we designate generators for

$$H^*(\Gamma_2/\Gamma_3, k) \simeq S^*(\oplus_{\ell=0}^{\infty} ((\gamma_2/\gamma_3)^{\#(\ell+1)})[2]) \otimes \Lambda^*(\oplus_{\ell=0}^{\infty} (\gamma_2/\gamma_3)^{\#(\ell)}[1])$$

by $x_\beta^{(\ell)}$ of cohomological 2 and $y_\beta^{(\ell)}$ of cohomological degree 1, with $0 \leq \ell$ and with β ranging over T -weights of U_J of level 2. Consequently,

$$(3.1.4) \quad E_2^{*,*}(U_J/\Gamma_3) = H^*(U_J/\Gamma_2, k) \otimes H^*(\Gamma_2/\Gamma_3, k)$$

equals

$$S^*(\oplus_{\ell=0}^\infty (\mathfrak{u}_J/\gamma_3)^{\#(\ell+1)}[2]) \otimes \Lambda^*(\oplus_{\ell=0}^\infty (\mathfrak{u}_J/\gamma_3)^{\#(\ell)}[1]),$$

where $E_2^{*,*}(U_J/\Gamma_3)$ is the spectral sequence considered in the following proposition.

The indexing we adopt (for example, in Proposition 3.2) relates to the above indexing as follows for cohomology classes of \mathbb{G}_a : λ_i corresponds to $y_\alpha^{(\ell)}$ and x_i corresponds to $x_\alpha^{(\ell)}$ with $\ell = i - 1$. The summation $\sum_{\alpha+\alpha'=\beta}$ indicates a sum of pairs of roots α, α' with the property that $[X_\alpha, X_{\alpha'}] = X_\beta \in \mathfrak{u}_J/\gamma_{v+1}$.

Proposition 3.2. *Retain the notation and hypotheses of Proposition 3.1. Consider the T -equivariant Lyndon-Hochschild-Serre spectral sequence [15] for the extension $1 \rightarrow \Gamma_2/\Gamma_3 \rightarrow U_J/\Gamma_3 \rightarrow U_J/\Gamma_2 \rightarrow 1$:*

$$(3.2.1) \quad E_2^{a,b}(U_J/\Gamma_3) = H^a(U_J/\Gamma_2, k) \otimes H^b(\Gamma_2/\Gamma_3, k) \Rightarrow H^{a+b}(U_J/\Gamma_3, k).$$

For any $\ell \geq 0$, $j \geq 0$, and any β a weight of level 2:

(1)

$$d_2^{0,1}(y_\beta^{(\ell)}) = \sum_{\alpha+\alpha'=\beta} y_\alpha^{(\ell)} \wedge y_{\alpha'}^{(\ell)} \in H^2(U_J/\Gamma_2, k).$$

(2) $d_{2p^j+1}^{0,2p^j}((x_\beta^{(\ell)})^{p^j}) = \sum_{\alpha+\alpha'=\beta} \{(x_\alpha^{(\ell)})^{p^j} \otimes y_{\alpha'}^{(\ell+1+j)} - (x_{\alpha'}^{(\ell)})^{p^j} \otimes y_\alpha^{(\ell+1+j)}\}$ is non-zero in $H^{2p^j+1+1}(U_J/\Gamma_2, k)$. Thus, $(x_\beta^{(\ell)})^{p^j} \in H^{2p^j}(\Gamma_2/\Gamma_3, k)$ does not lie in the image of $H^\bullet(U_J/\Gamma_3, k)$.

(3) $\beta \mathcal{P}^{p^j}(\sum_{\alpha+\alpha'=\beta} \{(x_\alpha^{(\ell)})^{p^j} \otimes y_{\alpha'}^{(\ell+1+j)} - (x_{\alpha'}^{(\ell)})^{p^j} \otimes y_\alpha^{(\ell+1+j)}\})$ equals

$$(3.2.2) \quad \sum_{\alpha+\alpha'=\beta} \{(x_\alpha^{(\ell)})^{p^{j+1}} \otimes x_{\alpha'}^{(\ell+1+j)} - (x_{\alpha'}^{(\ell)})^{p^{j+1}} \otimes x_\alpha^{(\ell+1+j)}\},$$

non-zero in $H^{2p^{j+1}+2}(U_J/\Gamma_2, k)$. The expression (3.2.2) maps to 0 in $H^{2p^{j+1}+2}(U_J/\Gamma_3, k)$.

Proof. Assume first that U_J equals U_3 and consider the extensions

$$(\mathbb{G}_a)_{(i)}/(\mathbb{G}_a)_{(i-1)} \rightarrow (U_3)_{(i)}/(U_3)_{(i-1)} \rightarrow (\mathbb{G}_a^{\times 2})_{(i)}/(\mathbb{G}_a^{\times 2})_{(i-1)}.$$

Since $(\mathbb{G}_a)_{(i)}/(\mathbb{G}_a)_{(i-1)}$ lies in the commutator of $(U_3)_{(i)}/(U_3)_{(i-1)}$, we conclude that the map $H^1((U_3)_{(i)}/(U_3)_{(i-1)}, k) \rightarrow H^1((\mathbb{G}_a)_{(i)}/(\mathbb{G}_a)_{(i-1)}, k)$ is 0, so that the differential $d_2^{0,1}$ must be non-zero for each of these extensions. We thus conclude using induction on r and the fact that differentials commute with the action of T_3 that $d_2^{0,1}(y_\beta^{(\ell)})$ is a sum of the form given in (1) with non-zero coefficients of each summand $y_\alpha^{(\ell)} \wedge y_{\alpha'}^{(\ell)}$. These coefficients must be equal for varying ℓ using functoriality with respect to Frobenius maps $U_{3(i)} \rightarrow U_{3(i+1)}$. Defining our group schemes over \mathbb{Z} , we conclude these coefficients must be ± 1 . We have chosen the ordering of the pairs α, α' so that these coefficients are all +1.

For a more general U_J/Γ_3 , we consider a pair α, α' with $\alpha + \alpha' = \beta$ and define the subgroup $R_{\alpha, \alpha'} \subset U_J/\Gamma_3$ to be the subgroup generated by the root subgroups

$U_\alpha, U_{\alpha'}$. Thus, $R_{\alpha, \alpha'} \simeq U_3$ with center U_β . The functoriality of the LHS spectral sequence implies that

$$(3.2.3) \quad \begin{array}{ccc} H^1(\Gamma_2/\Gamma_3, k) & \longrightarrow & H^1(U_\beta, k) \\ \downarrow d_2^{0,1} & & \downarrow d_2^{0,1} \\ H^2(U_J/\Gamma_2, k) & \longrightarrow & H^2(U_\alpha \times U_{\alpha'}, k) \end{array}$$

commutes, so that $d_2^{0,1}(y_\beta^{(\ell)})$ must be given by (1) plus additional terms. Yet there are no other eigenvectors of T -weight $p^\ell \beta$ in $H^2(U_J/\Gamma_2, k)$, so there can be no additional terms in the formula for $d_2^{0,1}(y_\beta^{(\ell)})$.

For $j = 0$, (2) follows from the equality $x_\beta^{(\ell)} = (\beta \mathcal{P}^0)(y_\beta^{(\ell)})$, the fact that $\beta \mathcal{P}^0$ commutes with transgression ([17]), and the Cartan formula

$$\beta \mathcal{P}^j(u \cdot v) = \sum_{i=0}^j ((\beta \mathcal{P}^i(u) \cdot \otimes \mathcal{P}^{j-i}(v) - (\mathcal{P}^i(u) \cdot (\beta \mathcal{P}^{j-i}(v)))(v)).$$

In particular, this tells us that

$$\beta \mathcal{P}^0(y_\alpha^{(\ell)} \wedge y_{\alpha'}^{(\ell)}) = \beta \mathcal{P}^0(y_\alpha^{(\ell)}) \otimes \mathcal{P}^0(y_{\alpha'}^{(\ell)}) - \mathcal{P}^0(y_\alpha^{(\ell)}) \otimes \beta \mathcal{P}^0(y_{\alpha'}^{(\ell)}) = x_\alpha^{(\ell)} \otimes y_{\alpha'}^{(\ell+1)} - x_{\alpha'}^{(\ell)} \otimes y_\alpha^{(\ell+1)}.$$

To prove (2) for $j > 0$, we recall that \mathcal{P}^{p^i} applied to $(x_\beta^{(\ell)})^{p^i}$ equals $(x_\beta^{(\ell)})^{p^{i+1}}$. Using the fact that the Steenrod action commutes with the differentials in the spectral sequence and repeated applications of the Cartan formula, we verify (2) by computing $d_{2p^j+1}^{0,2p^j}((x_\beta^{(\ell)})^{p^j})$, the result of applying $d_{2p^j+1}^{0,2p^j}$ to $(\mathcal{P}^{p^{j-1}} \circ \dots \circ \mathcal{P}^1 \circ \beta \mathcal{P}^0)(y_\beta^{(\ell)})$. The fact that $d_{2p^j+1}^{0,2p^j}((x_\beta^{(\ell)})^{p^j}) \neq 0$ follows from the explicit computation of $H^\bullet(U_J/\Gamma_2, k)$. Because some differential in the spectral sequence is non-vanishing on $(x_\beta^{(\ell)})^{p^j}$, it does not lie in the image of $H^\bullet(U_J/\Gamma_3, k)$.

The computation of assertion (3) follows from the Cartan formula for $\beta \mathcal{P}^{p^j}$ and the detailed description of \mathcal{P}^i and $\beta \mathcal{P}^i$ given in [24, 1.7]. The non-vanishing of (3.2.2) follows once again from the explicit computation of $H^\bullet(U_J/\Gamma_2, k)$. \square

The restriction map for the embedding $(U_J/\Gamma_3)_{(r)} \rightarrow U_J/\Gamma_3$ determines a map from the spectral sequence (3.2.1) to the spectral sequence

$$(3.2.4) \quad E_2^{a,b}((U_J/\Gamma_3)_{(r)}) = H^a((U_J/\Gamma_2)_{(r)}, k) \otimes H^b((\Gamma_2/\Gamma_3)_{(r)}, k) \Rightarrow H^{a+b}((U_J/\Gamma_3)_{(r)}, k).$$

considered in the next proposition. On E_2 -terms, this map sends $y_\beta^{(\ell)}, x_\beta^{(\ell)}, y_\alpha^{(\ell)}, x_\alpha^{(\ell)}$ to 0 for $\ell \geq r$.

To exhibit the action of T , we retain the indexing of Proposition 3.2, viewing $E_2^{*,*}((U_J/\Gamma_3)_{(r)})$ as the tensor product

$$(3.2.5) \quad S^*(\oplus_{\ell=0}^{r-1} (\mathfrak{u}_J/\gamma_2)^{\#(\ell+1)}[2]) \otimes \Lambda^*(\oplus_{\ell=0}^{r-1} (\mathfrak{u}_J/\gamma_2)^{\#(\ell)}[1]) \otimes S^*(\oplus_{\ell=0}^{r-1} ((\gamma_2/\gamma_3)^{\#(\ell+1)})[2]) \otimes \Lambda^*(\oplus_{\ell=0}^{r-1} (\gamma_2/\gamma_3)^{\#(\ell)}[1]).$$

Proposition 3.3. *Retain the notation and hypotheses of Proposition 3.1, and consider the spectral sequence (3.2.4) for the central extension*

$$(3.3.1) \quad 1 \rightarrow (\Gamma_2/\Gamma_3)_{(r)} \rightarrow (U_J/\Gamma_3)_{(r)} \rightarrow (U_J/\Gamma_2)_{(r)} \rightarrow 0.$$

For any β of level 2,

- (1) $(x_\beta^{(\ell)})^{p^j} \in S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(\ell+1)}[2]) \subset E_2^{0,*}((U_J/\Gamma_3)_{(r)})$ is a permanent cycle if and only if $\ell + 1 + j \geq r$.
- (2) For any $\ell, j \geq 0$ with $\ell + 1 + j < r$,
- (3.3.2)
$$\sum_{\alpha+\alpha'=\beta, \alpha<\alpha'} \{(x_\alpha^{(\ell)})^{p^{j+1}} \otimes x_{\alpha'}^{(\ell+1+j)} - (x_{\alpha'}^{(\ell)})^{p^{j+1}} \otimes x_\alpha^{(\ell+1+j)}\} = 0 \in H^{2p^{j+1}+2}((U_J/\Gamma_3)_{(r)}, k).$$
- (3) For $U_J = U_3$ (with $\Gamma_3 = 1$), the $p^{r-\ell-j-1}$ -st power of relation (3.3.2) in $H^\bullet((U_3)_{(r)}, k)$ is the restriction to $(U_3)_{(r)}$ of the relation $X^{1,2}(\ell) \cdot X^{2,3}(\ell') - X^{2,3}(\ell) \cdot X^{1,2}(\ell')$ of Theorem 1.2.

Proof. The vanishing of $y_\alpha^{(\ell)}, \ell \geq r$ together with Proposition 3.2(2) immediately implies that $(x_\beta^{(\ell)})^{p^j}$ is a permanent cycle if $\ell + 1 + j \geq r$. Conversely, if $\ell + 1 + j < r$, then Proposition 3.2 tells us that $d_{2p^j}^{0,2p^j}$ does not vanish on $(x_\beta^{(\ell)})^{p^j}$.

To conclude (2), we first recall that Proposition 3.2(2) implies that

$$(3.3.3) \quad \sum_{\alpha+\alpha'=\beta} \{(x_\alpha^{(\ell)})^{p^j} \otimes y_{\alpha'}^{(\ell+1+j)} - (x_{\alpha'}^{(\ell)})^{p^j} \otimes y_\alpha^{(\ell+1+j)}\} \in H^{2p^{j+1}}(U_J/\Gamma_2)_{(r)}, k)$$

is non-zero and in the image of a differential in the spectral sequence (3.2.4). This implies that (3.3.3) lies in the kernel of the inflation map. One obtains the relation of (3.3.2) by applying $\beta\mathcal{P}^j$ to (3.3.3) to obtain a class in $H^{2p^{j+1}+2}((U_J/\Gamma_2)_{(r)}, k)$ whose inflation gives (3.3.2); since $\beta\mathcal{P}^j$ commutes with the inflation map, we conclude the asserted vanishing.

Assertion (2) follows from Proposition 3.2(2), since $\sum_{\alpha+\alpha'=\beta} \{(x_\alpha^{(\ell)})^{p^{j+1}} \otimes y_{\alpha'}^{(\ell+1+j)} - (x_{\alpha'}^{(\ell)})^{p^{j+1}} \otimes y_\alpha^{(\ell+1+j)}\}$ is a boundary and the restriction map commutes with the Bockstein.

The fact that the $p^{r-\ell-j-1}$ -st power of (3.3.2) equals the relation $X^{1,2}(\ell) \cdot X^{2,3}(\ell') - X^{2,3}(\ell) \cdot X^{1,2}(\ell')$ of Theorem 1.2 is immediate from the identification of $X^{1,2}(\ell)$ with $(x_\alpha^{(\ell)})^{p^{r-\ell-1}}$ and $X^{2,3}(\ell)$ with $(x_{\alpha'}^{(\ell)})^{p^{r-\ell-1}}$. \square

We view $S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(\ell+1)}[2])$ as a subalgebra of $E_2^{0,*}((U_J/\Gamma_3)_{(r)})$ using the identification (3.2.5). Proposition 3.3(1) tells us that the subalgebra

$$S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}]) \subset S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(\ell+1)}[2])$$

(defined as the image of the endomorphism $S^*(\oplus_{\ell=0}^{r-1}F^{r-\ell-1})$ on $S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(\ell+1)}[2])$) consists of permanent cycles in the spectral sequence (3.2.4). The following corollary tells us that this subalgebra is the intersection of the permanent cycles in $E_2^{*,*}((U_J/\Gamma_3)_{(r)})$ with $S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(\ell+1)}[2])$. What this corollary does *not* do is identify all permanent cycles of $E_2^{*,*}((U_J/\Gamma_3)_{(r)})$.

Corollary 3.4. *If $z \in S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(\ell+1)}[2])$ does not lie in the subalgebra $S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}])$, then there exists some differential of (3.2.4) which is non-zero on z .*

Proof. We employ the fact that the differentials in the spectral sequence are k -linear derivations. Let $\{\beta_i, i \in I\}$ be the set of positive roots of U_J of level 2. Consider a monomial $w = \prod_{i \in I} \prod_{\ell=0}^{r-1} (x_{\beta_i}^{(\ell)})^{n_{i,\ell}}$ with some $n_{i,\ell}$ not divisible by $p^{r-\ell-1}$. Let p^j be the smallest power of p such that p^j divides some $n_{i,\ell}$, and p^{j+1} does not divide $n_{i,\ell}$. Then $d_{2p^j+1}^{0, \sum_{\ell} 2n_{i,\ell}}(w)$ is a sum of non-zero terms indexed by those i, ℓ with p^j

but not p^{j+1} dividing $n_{i,\ell}$, each summand having a different T -weight. Thus, w is not a permanent cycle.

More generally, different such monomials have different T -weights, so that no non-trivial sum of such monomials is a permanent cycle. \square

Notation 3.5. Let $U_N \subset GL_N$ denote the subgroup of strictly upper triangular matrices. The generators $X^{i,j}(\ell) \in S^*((\mathfrak{u}_N/\gamma_3)^{\#(r)}[2p^{r-\ell-1}])$ (as discussed following Proposition 1.1) correspond to $(x_{i,j}^{(\ell)})^{p^{r-\ell-1}} \in E_2^{*,*}((U_N/\Gamma_3)_{(r)})$ of Proposition 3.3.

We extend this notation, denoting generators of $S^*((\mathfrak{u}_J/\gamma_3)^{\#(r)}[2p^{r-\ell-1}])$ by $X^\beta(\ell) \in S^*((\mathfrak{u}_J/\gamma_3)^{\#(r)}[2p^{r-\ell-1}])$. These classes have the same weight and cohomological degree as the classes $(x_\beta^{(\ell)})^{p^{r-\ell-1}} \in E_2^{*,*}((U_J/\Gamma_3)_{(r)})$. We shall see that the representative in $E_2^{*,*}((U_J/\Gamma_3)_{(r)})$ of $\eta_{U_J/\Gamma_3,r}(X^\beta(\ell)) \in H^*((U_J/\Gamma_3)_{(r)}, k)$ is $(x_\beta^{(\ell)})^{p^{r-\ell-1}}$.

The uniqueness given in the following proposition enables us to specify the map $\eta_{U_J/\Gamma_3,r}$. We are particularly interested in the special case $U_J/\Gamma_3 = U_3$.

Proposition 3.6. *Retain the notation and hypotheses of Proposition 3.1. Assume each root β of U_J of level 2 can be written uniquely as a sum $\alpha + \alpha'$ of roots U_J of level 1. Then there exists a unique T -equivariant k -linear map*

$$\eta_r : (\mathfrak{u}_J/\gamma_3)^{\#(r)}[2p^{r-1}] \rightarrow H^{2p^{r-1}}((U_J/\Gamma_3)_{(r)}, k)$$

which fits in the following commutative diagram (3.6.1)

$$\begin{array}{ccccc} (\mathfrak{u}_J/\gamma_2)^{\#(r)}[2p^{r-1}] & \longrightarrow & (\mathfrak{u}_J/\gamma_3)^{\#(r)}[2p^{r-1}] & \longrightarrow & (\gamma_2/\gamma_3)^{\#(r)}[2p^{r-1}] \\ \downarrow & & \downarrow \eta_r & & \downarrow \\ H^{2p^{r-1}}((U_J/\Gamma_2)_{(r)}, k) & \longrightarrow & H^{2p^{r-1}}((U_J/\Gamma_3)_{(r)}, k) & \longrightarrow & H^{2p^{r-1}}((\Gamma_2/\Gamma_3)_{(r)}, k). \end{array}$$

Here, the left and right vertical maps are given by the inclusions $S^*((\mathfrak{u}_J/\gamma_2)^{\#(1)}[2]) \rightarrow H^\bullet((U_J/\Gamma_2)_{(r)}, k)$ and $S^*((\gamma_2/\gamma_3)^{\#(1)}[2]) \rightarrow H^\bullet((\Gamma_2/\Gamma_3)_{(r)}, k)$, the upper horizontal maps are the evident ones, the lower horizontal maps are those given by functoriality.

Furthermore, the map η_r fits in a commutative square

$$(3.6.2) \quad \begin{array}{ccc} (\mathfrak{u}_J/\gamma_3)^{\#(r)}[2p^{r-1}] & \xrightarrow{\eta_r} & H^{2p^{r-1}}((U_J/\Gamma_3)_{(r)}, k) \\ \downarrow & & \downarrow \\ (\mathfrak{u}_J)^{\#(r)}[2p^{r-1}] & \xrightarrow{(\phi_{U_J,r})|} & H^{2p^{r-1}}((U_J)_{(r)}, k) \end{array}$$

whose vertical maps are inflation maps and whose lower horizontal map is the restriction of the map $\phi_{U,r}$ of Theorem 1.7 with $U = U_J$ satisfying the hypotheses of this proposition.

Proof. The existence of some η fitting in the commutative diagram (3.6.1) is implied by Proposition 3.3(1). To prove the uniqueness of η , it suffices to verify for each root β of U_J of level 2 that the T -weight space of $H^{2p^{r-1}}((U_J/\Gamma_3)_{(r)}, k)$ of weight $p^r\beta$ is

1-dimensional. This would imply the uniqueness of the choice of class $\eta(X^\beta(0)) \in H^{2p^{r-1}}((U_J/\Gamma_3)_{(r)}, k)$ fitting in the commutative diagram (3.6.1).

We search in ${}^{AJ}E_1^{*,*}$ as given in (2.1.2) for T -weight vectors with T -weight $p^r\beta$ and cohomology degree $2p^{r-1}$ other than $(x_\beta^{(0)})^{p^{r-1}} \in E_2^{0,2p^{r-1}}$. Consider a simple tensor of the specified weight and degree, in other words a monomial z in x 's and y 's. Because the weight of $y_\alpha^{(0)} \wedge y_{\alpha'}^{(0)}$ is not divisible by p , this does not divide the monomial z . None of the factors of the monomial z of weight $p^r\beta$ and degree $2p^{r-1}$ can be of the form $x^{(\ell)}$ for $\ell > 1$ or for $y^{(\ell)}$ for $\ell > 2$ because such a factor would increase the weight too "fast" with respect to increase of the resulting degree by either 2 or 1.

Thus, the only allowable weight vectors of cohomology degree $2p^{r-1}$ are scalar multiples of $(x_\beta^{(0)})^{p^{r-1}}$ and $(x_\beta^{(0)})^{p^{r-1}-1} \otimes y_\alpha^{(1)} \wedge y_{\alpha'}^{(1)}$. Since $y_\beta^{(i)} \in E_2^{0,1}$ is a permanent cycle, we conclude that $d_2((x_\beta)^{(i)}) = 0$ (using the fact that differentials in the spectral sequence commute with Bocksteins). Consequently, the value of the derivation $d_2^{0,2p^{r-1}-1}$ applied to $(x_\beta^{(0)})^{p^{r-1}-1} \otimes y_\beta^{(1)}$ equals $(x_\beta^{(0)})^{p^{r-1}-1} \otimes y_{\alpha_1}^{(1)} \wedge y_{\alpha_2}^{(1)}$. We conclude that the class of $(x_\beta^{(0)})^{p^{r-1}} \in E_\infty^{0,2p^{r-1}}$ spans the $p^r\beta$ weight space of degree $2p^{r-1}$ of $E_\infty^{0,2p^{r-1}}$. This implies that the $p^r\beta$ weight space of $H^{2p^{r-1}}((U_J/\Gamma_3)_{(r)}, k)$ is 1-dimensional.

Finally, the commutativity of (3.6.2) follows from the T -equivariance of η and the uniqueness assertion of Theorem 1.7. \square

Definition 3.7. Retain the notation and hypotheses of Propositions 3.6. We define $S^*((U_J/\Gamma_v)_{(r)})$ to be

$$(3.7.1) \quad S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_2)^{\#(\ell+1)}[2]) \otimes_{S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_2)^{\#(r)}[2p^{r-\ell-1}])} S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_v)^{\#(r)}[2p^{r-\ell-1}]).$$

In other words, $S^*((U_J/\Gamma_v)_{(r)})$ is the coproduct in the category of commutative k algebras of $S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_2)^{\#(\ell+1)}[2])$ and $S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_v)^{\#(r)}[2p^{r-\ell-1}])$ over $S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_2)^{\#(r)}[2p^{r-\ell-1}])$. The T -equivariant splitting $\mathfrak{u}_J/\gamma_v \simeq (\mathfrak{u}_J/\gamma_2) \oplus (\gamma_2/\gamma_v)$ gives the T -equivariant splitting

$$(3.7.2) \quad S^*((U_J/\Gamma_v)_{(r)}) \simeq S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_2)^{\#(\ell+1)}[2]) \otimes S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_v)^{\#(r)}[2p^{r-\ell-1}]).$$

Definition 3.8. Retain the notation and hypotheses of Propositions 3.1. We define

$$(3.8.1) \quad \tilde{g} : S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}]) \rightarrow H^\bullet((U_J/\Gamma_3)_{(r)}, k)$$

to be the map of k -algebras determined by the ℓ -th Frobenius twists

$$\eta_{r-\ell}^{(\ell)} : (\mathfrak{u}_J/\gamma_3)^{\#(r)}[2p^{r-\ell-1}] \rightarrow H^{2p^{r-\ell-1}}((U_J/\Gamma_3)_{(r)}, k) \text{ of the maps}$$

$$\eta_{r-\ell} : (\mathfrak{u}_J/\gamma_3)^{\#(r-\ell)}[2p^{r-\ell-1}] \rightarrow H^{2p^{r-\ell-1}}((U_J/\Gamma_3)_{(r-\ell)}, k) \text{ constructed in Proposition 3.6 (with } r \text{ replaced by } r-\ell).$$

We define

$$(3.8.2) \quad \tilde{f} : S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_2)^{\#(\ell+1)}[2]) \rightarrow H^\bullet((U_J/\Gamma_2)_{(r)}, k) \rightarrow H^\bullet((U_J/\Gamma_3)_{(r)}, k)$$

to be the composition of $\eta_{U_J/\Gamma_2, r}$ given in (3.1.3) and the inflation map.

We define

$$(3.8.3) \quad \eta_{U_J/\Gamma_3, r} = \tilde{f} \circ \tilde{g} : S^*((U_J/\Gamma_3)_{(r)}) \rightarrow H^\bullet((U_J/\Gamma_3)_{(r)}, k)$$

to be the coproduct of \tilde{g} and the map \tilde{f} . The maps \tilde{g} and \tilde{f} agree on $S^*((\oplus_{\ell=0}^{r-1}(\mathbf{u}_J/\gamma_2)^{\#(\ell+1)}[2]p^{r-\ell-1}))$ by commutativity of the left square of (3.6.1), so that $\eta_{U_J/\Gamma_3,r}$ is well defined.

We give $S^*((U_J/\Gamma_3)_{(r)})$ the decreasing filtration whose subalgebra of level $2i$ is the coproduct of $S^{\geq i}(\oplus_{\ell=0}^{r-1}(\mathbf{u}_J/\gamma_2)^{\#(\ell+1)}[2])$ and $S^*((\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2]p^{r-\ell-1}))$ over $S^{\geq i}(\oplus_{\ell=0}^{r-1}(\mathbf{u}_J/\gamma_2)^{\#(r)}[2]p^{r-\ell-1}))$.

Proposition 3.9. *The map $\eta_{U_J/\Gamma_3,r}$ is a naturally defined T -equivariant map of filtered k -algebras, where we give $H^\bullet((U_J/\Gamma_3)_{(r)}, k)$ the LHS filtration of Proposition 3.3 and $S^*((U_J/\Gamma_3)_{(r)})$ the filtration described immediately above.*

Furthermore,

$$(3.9.1) \quad gr(\eta_{U_J/\Gamma_3,r}) = f \odot g : S^*((U_J/\Gamma_3)_{(r)}) \rightarrow gr\{H^\bullet((U_J/\Gamma_3)_{(r)}, k)\}$$

where $g : S^*((\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2]p^{r-\ell-1})) \rightarrow E_\infty^{0,*}$ is defined to be the composition of \tilde{g} with the natural map $H^\bullet((U_J/\Gamma_3)_{(r)}, k) \rightarrow E_\infty^{0,*}$ and $f : S^*((U_J/\Gamma_2)_{(r)}) \rightarrow H^\bullet((U_J/\Gamma_2)_{(r)}, k) \rightarrow E_\infty^{0,*}$ is the map whose composition with the natural inclusion $E_\infty^{0,*} \rightarrow H^\bullet((U_J/\Gamma_3)_{(r)}, k)$ equals \tilde{f} .

Proof. The map \tilde{f} is induced by $U_J/\Gamma_3 \rightarrow U_J/\Gamma_2$ and thus is filtration preserving. Observe that $1 \otimes S^*((\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2]p^{r-\ell-1})) \subset S^*((U_J/\Gamma_3)_{(r)})$ has filtration degree 0 as does $E_\infty^{0,*}$, so that \tilde{g} is also a map of filtered algebras. The multiplicative property of these filtrations thus implies that $\eta_{U_J/\Gamma_3,r}$ itself is a map of filtered algebras.

The identification of $gr\{\eta_{U_J/\Gamma_3,r}\}$ is verified by proving the commutativity of the following two diagrams

$$(3.9.2) \quad \begin{array}{ccccc} S^*((\oplus_{\ell=0}^{r-1}(\mathbf{u}_J/\gamma_2)^{\#(\ell+1)}[2])) & \xrightarrow{=} & S^*((\oplus_{\ell=0}^{r-1}(\mathbf{u}_J/\gamma_2)^{\#(\ell+1)}[2])) & \longrightarrow & S^*((U_J/\Gamma_3)_{(r)}) \\ \downarrow \eta_{U_J/\Gamma_2,r} & & \downarrow & & \downarrow \eta_{U_J/\Gamma_3,r} \\ H^\bullet((U_J/\Gamma_2)_{(r)}, k) & \longrightarrow & E_\infty^{0,*}((U_J/\Gamma_3)_{(r)}) & \longrightarrow & H^\bullet((U_J/\Gamma_3)_{(r)}, k), \end{array}$$

$$(3.9.3) \quad \begin{array}{ccccc} S^*((U_J/\Gamma_3)_{(r)}) & \longrightarrow & S^*((\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2]p^{r-\ell-1})) & \longrightarrow & S^*((\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(\ell+1)}[2])) \\ \downarrow \eta_{U_J/\Gamma_3,r} & & \downarrow g & & \downarrow \tilde{g} \\ H^\bullet((U_J/\Gamma_3)_{(r)}, k) & \longrightarrow & E_\infty^{0,*}((U_J/\Gamma_3)_{(r)}) & \longrightarrow & H^\bullet((\Gamma_2/\Gamma_3)_{(r)}, k) \end{array}$$

By Proposition 3.3(2), the left square of (3.9.2) commutes. By definition of $\eta_{U_J/\Gamma_3,r}$, the right square of (3.9.2) also commutes.

The commutativity of the outer square of (3.9.3) arises from the naturality of the restriction maps for $\Gamma_2/\Gamma_3 \rightarrow U_J/\Gamma_3$. The commutativity of the two squares of (3.9.3) thus follows from the fact that $S^*((\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2]p^{r-\ell-1}))$ is the image of the upper composition of (3.9.3) and the fact that $E_\infty^{0,*}((U_J/\Gamma_3)_{(r)})$ is the image of the restriction map $H^\bullet((U_J/\Gamma_3)_{(r)}, k) \rightarrow H^\bullet((\Gamma_2/\Gamma_3)_{(r)}, k)$ by a standard property of Grothendieck spectral sequences. \square

Definition 3.10. We define $Q((U_J/\Gamma_2)_{(r)})$ to be the quotient of $S^*((\oplus_{\ell=0}^{r-1}(\mathbf{u}_J/\gamma_2)^{\#(\ell+1)}[2]))$ by the ideal generated by the elements of (3.3.2), which we denote by J_2 :

$$Q((U/\Gamma_2)_{(r)}) \equiv S^*((\oplus_{\ell=0}^{r-1}(\mathbf{u}_J/\gamma_2)^{\#(\ell+1)}[2]))/J_2.$$

We define $\overline{S}^*((U_J/\Gamma_3)_{(r)})$ to be the tensor product of $S^*((U_J/\Gamma_3)_{(r)})$ and $Q((U_J/\Gamma_2)_{(r)})$ over $S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_2)^{\#(\ell+1)}[2])$.

The T -equivariant splitting $\mathfrak{u}_J/\gamma_3 \simeq (\mathfrak{u}_J/\gamma_2) \oplus (\gamma_2/\gamma_3)$ gives the T equivariant splitting

$$(3.10.1) \quad \overline{S}^*((U_J/\Gamma_3)_{(r)}) \simeq Q((U_J/\Gamma_2)_{(r)}) \otimes S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}]).$$

We view $\overline{S}^*((U_J/\Gamma_3)_{(r)})$ as

$$\overline{S}^*((U_J/\Gamma_3)_{(r)}) \simeq S^*((U_J/\Gamma_3)_{(r)})/I_3$$

where $I_3 \subset S^*((U_J/\Gamma_3)_{(r)})$ is the ideal generated by the relations (3.3.2).

Proposition 3.11. *The map $\eta_{U_J/\Gamma_3,r}$ of Proposition 3.9 factors through the quotient $S^*((U_J/\Gamma_3)_{(r)}) \twoheadrightarrow \overline{S}^*((U_J/\Gamma_3)_{(r)})$, thereby determining the map of k -algebras*

$$(3.11.1) \quad \overline{\eta}_{U_J/\Gamma_3,r} = \tilde{f}_Q \odot \tilde{g} : \overline{S}^*((U_J/\Gamma_3)_{(r)}) \rightarrow H^\bullet((U_J/\Gamma_3)_{(r)}, k),$$

where \tilde{f}_Q factors the map \tilde{f} of (3.8.2) via the natural surjection $S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_2)^{\#(\ell+1)}[2]) \rightarrow Q((U_J/\Gamma_2)_{(r)})$.

The map $\overline{\eta}_{U_J/\Gamma_3,r}$ is a map of filtered algebras for the LHS filtration. Moreover,

$$gr\{\overline{\eta}_{U_J/\Gamma_3,r}\} = f_Q \odot g,$$

where $f_Q : Q((U_J/\Gamma_2)_{(r)}) \rightarrow E_\infty^{*,0}$ composed with the natural map $E_\infty^{*,0} \rightarrow H^*((U_J/\Gamma_3)_{(r)}, k)$ equals \tilde{f}_Q .

Proof. Because $\eta_{U_J,3}(I_3) = 0$, $\eta_{U_J,3}$ induces $\overline{\eta}_{U_J,3}$. Since I_3 is generated by elements of J_2 , we conclude that $\overline{\eta}_{U_J,3}$ is given as indicated in (3.11.1). Observe that $Q^*((U_J/\Gamma_2)_{(r)}) = S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_2)^{\#(\ell+1)}[2])/J_2$ inherits an LHS filtration because J_2 is a filtered ideal.

Thus, to prove that $\overline{\eta}_{U_J/\Gamma_3,r}$ is filtration preserving, it suffices to observe that

- (1) $\overline{\eta}_{U_J/\Gamma_3,r}$ restricted to $Q^*((U_J/\Gamma_2)_{(r)})$ is a filtered map of algebras for the LHS filtration because it is induced by the inflation map for $(U_J/\Gamma_3)_{(r)} \rightarrow (U_J/\Gamma_2)_{(r)}$.
- (2) $\eta_{U_J/\Gamma_3,r}$ restricted to $S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(\ell+1)}[2])$ is also a filtered map of algebras for the LHS filtration, as shown in the proof of Propositions 3.9.

The identification of $gr\{\overline{\eta}_{U_J/\Gamma_3,r}\}$ follows from Proposition 3.9 and the fact that the LHS filtration on $\overline{S}^*((U_J/\Gamma_3)_{(r)})$ is induced by taking the quotient of the filtration for $S^*((U_J/\Gamma_3)_{(r)})$ by an ideal which is of the form $J_2 \otimes S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}])$. \square

The Andersen-Jantzen spectral sequence of Proposition 2.1 admits a natural action of T whose T -weights are identified using (2.1.2). We envision that this spectral sequence should enable the extension to U_J/Γ_{v+1} of our considerations of U_J/Γ_3 .

For U_J/Γ_3 , we verify below that the elements of $S^*((U_J/\Gamma_3)_{(r)})$ are permanent cycles of the AJ spectral sequence. It would be of interest to know whether or not all permanent cycles in $S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_3)^{\#(\ell+1)}[2])$ belong to $S^*((U_J/\Gamma_3)_{(r)})$.

Proposition 3.12. *Retain the hypotheses and notation of Proposition 3.1. Every element*

$$z \in S^*((U_J/\Gamma_3)_{(r)}) \subset S^*(\oplus_{\ell=0}^{r-1}(\mathfrak{u}_J/\gamma_3)^{\#(\ell+1)}[2]) \subset {}^{AJ}E_1^{*,*}((U_J/\Gamma_3)_{(r)})$$

is a permanent cycle for the AJ spectral sequence.

Moreover, the map $\eta_{U_J/\Gamma_3,r} : S^*((U_J/\Gamma_3)_{(r)}) \rightarrow H^\bullet((U_J/\Gamma_3)_{(r)}, k)$ sends $z \in S^*((U_J/\Gamma_3)_{(r)})$ to the cohomology class $\eta_{U_J/\Gamma_3,r}(z)$ represented at the E_1^{**} -page of the AJ spectral sequence by the image of z in ${}^{AJ}gr(S^*((U_J/\Gamma_3)_{(r)}) \simeq S^*((U_J/\Gamma_3)_{(r)}) \subset {}^{AJ}E_1^{*,*}((U_J/\Gamma_3)_{(r)})$.

Proof. The map $\eta_{U_J/\Gamma_2,r} : S^*((U_J/\Gamma_2)_{(r)}) \rightarrow H^\bullet((U_J/\Gamma_2)_{(r)}, k)$ of Definition 3.8 is the tensor power of natural embeddings $S^*(H^2(\mathbb{G}_{a(r)}, k)) \rightarrow H^\bullet(\mathbb{G}_{a(r)}, k)$ as in (3.1.2) and so can be identified with ${}^{AJ}gr(\eta_{U_J/\Gamma_2,r})$.

Recall that $S^*((U_J/\Gamma_3)_{(r)})$ is the tensor product of its restrictions to $S^*(\oplus_{\ell=0}^{r-1} (\mathfrak{u}_J/\gamma_2)^{\#(\ell+1)}[2])$ and $S^*(\oplus_{\ell=0}^{r-1} (\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}])$. Using functoriality for the map $U_J/\Gamma_3 \rightarrow U_J/\Gamma_2$ and multiplicativity of $\eta_{U/\Gamma_3,r}$, we invoke a simple induction argument to conclude that it suffices to prove the assertions for $z \in S^*(\oplus_{\ell=0}^{r-1} (\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}])$.

Let β be a root of U_J of level 2 and $0 \leq \ell < r$. Consider $\eta_{U/\Gamma_3,r-\ell}$ applied to $(X^\beta(0))$ (corresponding to $(x_\beta^{(0)})^{p^{r-1}} \in E_1^{*,*}$ as in Notation 3.5), giving an element of $H^{2p^{r-\ell-1}}((U_J/\Gamma_3)_{(r-\ell)}, k)$ of weight $p^{r-\ell-1} \cdot \beta$. An inspection of (2.1.2) verifies that the only possible representative in ${}^{AJ}E_1^{*,*}((U_J/\Gamma_3)_{(r-\ell)})$ of this weight and degree is $(x_\beta^{(0)})^{p^{r-1-\ell}}$.

The following square commutes

$$(3.12.1) \quad \begin{array}{ccc} S^*((U_J/\Gamma_3)_{(r-\ell)}) & \xrightarrow{F^{\ell*}} & \overline{S}^*((U_J/\Gamma_3)_{(r)}) \\ \eta_{U/\Gamma_3,r-\ell} \downarrow & & \downarrow \eta_{U/\Gamma_3,r} \\ H^*((U_J/\Gamma_3)_{(r-\ell)}, k) & \xrightarrow{F^{\ell*}} & H^*((U_J/\Gamma_3)_{(r)}, k) \end{array}$$

because the maps η are defined over \mathbb{F}_p . Moreover, $(F^\ell)^*$ determines a map of AJ spectral sequences, so that the representative in ${}^{AJ}E_1^{*,*}((U_J/\Gamma_3)_{(r-\ell)})$ of $\eta_{U_J/\Gamma_3,r-\ell}(X^\beta(0))$ is sent to the representative in ${}^{AJ}E_1^{*,*}((U_J/\Gamma_3)_{(r)})$ of $\eta_{U_J/\Gamma_3,r}(X^\beta(\ell))$. In particular, $(x_\beta^{(\ell)})^{p^{r-1-\ell}} \in S^*((U_J/\Gamma_3)_{(r)})$ is a permanent cycle in the AJ spectral sequence representing $\eta_{U/\Gamma_3,r}(X^\beta(\ell))$ for each β and each $0 \leq \ell < r$. Now, using multiplicativity, we conclude the assertions of the proposition for $z \in S^*(\oplus_{\ell=0}^{r-1} (\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}])$ and thus the proposition as stated. \square

As a corollary of Proposition 3.12, we conclude the following description $gr\{\eta_{U_J/\Gamma_3,r}\}$. This is of some interest for the AJ-filtration is intrinsic to U_J/Γ_3 .

Corollary 3.13. *As for $gr\{\eta_{U_J/\Gamma_3,r}\}$ in Proposition 3.9,*

$${}^{AJ}gr\{\eta_{U_J/\Gamma_3,r}\} = f \odot g : S^*((U_J/\Gamma_3)_{(r)}) \rightarrow {}^{AJ}gr\{H^*((U_J/\Gamma_3)_{(r)}, k)\}.$$

4. THE MAP $\overline{\eta}_{U_3,r} : \overline{S}^*((U_3)_{(r)}) \rightarrow H^\bullet((U_3)_{(r)}, k)$

We apply the results of the previous section to the Heisenberg group U_3 and its Frobenius kernels $(U_3)_{(r)}$.

Example 4.1. $\overline{S}^*((U_3)_{(r)})$ is generated by elements $X^{1,3}(\ell) = (x_{1,3}^{(\ell)})^{p^{r-\ell-1}} \in (\gamma_2)^{\#(r)}[2p^{r-\ell-1}]$ and $(X^{1,2}(\ell))^{p^{-r+\ell+1}} = x_{1,2}^{(\ell)}$, $(X^{2,3}(\ell))^{p^{-r+\ell+1}} = x_{2,3}^{(\ell)} \in (\mathfrak{u}_3/\gamma_2)^{\#(\ell+1)}[2]$

with $0 \leq \ell < r$. A set of relations for $\overline{S}^*((U_3)_{(r)})$ is given by the special case of (3.3.2):

$$(4.1.1) \quad (x_{1,2}^{(\ell)})^{p^{j+1}} \otimes x_{2,3}^{(\ell+1+j)} - (x_{2,3}^{(\ell)})^{p^{j+1}} \otimes x_{1,3}^{(\ell+1+j)} = 0 \in H^{2p^{j+1}+2}((U_3)_{(r)}, k)$$

for each $\ell, j \geq 0$ such that $\ell + 1 + j < r$.

Generators and relations for $Q((U_3/\Gamma_2)_{(r)})$ are obtained from those for $\overline{S}^*((U_3)_{(r)})$ by setting each $X^{1,3}(\ell)$ equal to 0.

Composition with the quotient map $U_J/\Gamma_3 \rightarrow U_J/\Gamma_2$ determines $V_r(U_J/\Gamma_3) \rightarrow V_r(U_J/\Gamma_2)$ and thus the map of k -algebras $k[V_r(U_J/\Gamma_2)] \rightarrow k[V_r(U_J/\Gamma_3)]$. We denote the image of this map by $\overline{k[V_r(U_J/\Gamma_3)]} \subset k[V_r(U_J/\Gamma_3)]$.

In the proof of the following proposition we use the description of $k[V_r(U_J/\Gamma_3)]$ in terms of generators and relations which follows immediately from Proposition 1.5 and the explicit description of $k[V_r(GL_N)]$ given in Theorem 1.2. Namely, $k[V_r(U_J/\Gamma_3)]$ is generated by $X^{i,j}(\ell)$ where $1 \leq i \leq j \leq 3$ and $0 \leq \ell < r$; a set of relations is given by

$$(4.1.2) \quad X^{1,2}(\ell) \cdot X^{2,3}(\ell') - X^{2,3}(\ell) \cdot X^{1,2}(\ell'), \quad 0 \leq \ell < \ell'.$$

Proposition 4.2. *The coordinate algebra $k[V_r(U_J/\Gamma_3)]$ admits a natural tensor product decomposition as k -algebras,*

$$k[V_r(U_J/\Gamma_3)] = \overline{k[V_r(U_3/\Gamma_2)]} \otimes S^*(\oplus_{\ell=0}^{r-1} (\gamma_2/\gamma_3)^{\#(r)} [2p^{r-\ell-1}]).$$

In the special case $U_J = U_3$, $\overline{k[V_r(U_3/\Gamma_2)]}$ is an integral domain smooth outside of the origin with field of fractions a purely transcendental extension of transcendence degree $r + 1$.

Consequently, $k[V_r(U_3)]$ is an integrally closed domain of dimension $2r + 1$.

Proof. We identify $k[V_r(U_J/\Gamma_2)]$ with

$$(4.2.1) \quad S^*(\oplus_{\ell=0}^{r-1} (\mathbf{u}_J/\gamma_2)^{\#(r)} [2p^{r-\ell-1}]) \simeq k[X^{1,2}(\ell), X^{2,3}(\ell'); 0 \leq \ell, \ell' < r]$$

and $\overline{k[V_r(U_3/\Gamma_2)]}$ with the quotient of $k[V_r(U_J/\Gamma_2)]$ by the relations (4.1.2). The tensor product decomposition is immediate from the observation that these relations do not involve elements of $S^*(\oplus_{\ell=0}^{r-1} (\gamma_2/\gamma_3)^{\#(r)} [2p^{r-\ell-1}])$.

If $r = 1$, then $V_r(U_3) = \mathbf{u}_3$ so that $\overline{k[V_1(U_3/\Gamma_2)]} \simeq k[X^{1,3}(0), X^{2,3}(0)]$. For the remainder of the proof, we assume $r > 1$.

For any $\ell_1, 0 \leq \ell_1 < r$, the algebra $\overline{k[V_r(U_3/\Gamma_2)]}[(X^{2,3}(\ell_1))^{-1}]$ is isomorphic to

$$k[X^{2,3}(\ell), 0 \leq \ell < r; X^{1,2}(\ell_1)][(X^{2,3}(\ell_1))^{-1}],$$

since $X^{1,2}(\ell) = X^{1,2}(\ell_1)(X^{2,3}(\ell_1))^{-1}X^{2,3}(\ell)$; similarly, for any $\ell_0, 0 \leq \ell_0 < r$, the algebra $\overline{k[V_r(U_3/\Gamma_2)]}[(X^{1,2}(\ell_0))^{-1}]$ is isomorphic to

$$k[X^{1,2}(\ell), 0 \leq \ell < r; X^{2,3}(\ell_0)][(X^{1,2}(\ell_0))^{-1}].$$

This verifies the computation of the field of fractions of $\overline{k[V_r(U_3/\Gamma_2)]}$ and shows that $\overline{k[V_r(U_3/\Gamma_2)]}$ is smooth outside the common zeros of $\{X^{1,2}(\ell_0), X^{2,3}(\ell_1); 0 \leq \ell_0, \ell_1 < r\}$; namely, the origin. A theorem of Serre (see [18, Thm 39]) tells us that $\overline{k[V_r(U_3/\Gamma_2)]}$ is an integrally closed domain since the codimension of this zero locus is at least 2. \square

Proposition 4.3. *There is a naturally constructed injective map*

$$(4.3.1) \quad \theta_{U_J/\Gamma_3, r} : k[V_r(U_J/\Gamma_3)] \rightarrow \overline{S}^*((U_J/\Gamma_3)_{(r)})$$

such that $\eta_{U_J/\Gamma_3, r} : S^*((U_J/\Gamma_3)_{(r)}) \rightarrow H^\bullet((U_J/\Gamma_3)_{(r)}, k)$ as defined in Definition 3.8 factors through $\overline{\eta}_{U_J/\Gamma_3, r} \circ \theta_{U_J/\Gamma_3, r} : k[V_r(U_J/\Gamma_3)] \rightarrow H^\bullet((U_J/\Gamma_3)_{(r)}, k)$, where $\overline{\eta}_{U_J/\Gamma_3, r}$ is given in Proposition 3.11.

In the special case $U_J = U_3$,

$$\overline{\eta}_{U_3, r} \circ \theta_{U_3, r} = \overline{\phi}_{U_3, r} : k[V_r(U_3)] \rightarrow H^\bullet((U_3)_{(r)}, k),$$

where $\overline{\phi}_{U_3, r}$ is given in Proposition 1.9.

Proof. Consider the defining quotient map $q_{U_J/\Gamma_3, r} : S^*(\oplus_{\ell=0}^{r-1}(\mathbf{u}_J/\gamma_3)^{\#(r)}[2p^{r-\ell-1}]) \rightarrow k[V_r(U_J/\Gamma_3)]$. We see by inspection that the kernel of $q_{U_J/\Gamma_3, r}$ is generated by the intersection of $J_2 \subset S^*(\oplus_{\ell=0}^{r-1}(\mathbf{u}_J/\gamma_2)^{\#(r)}[2p^{r-\ell-1}])$ with $S^*(\oplus_{\ell=0}^{r-1}(\mathbf{u}_J/\gamma_2)^{\#(r)}[2p^{r-\ell-1}])$. This determines $\theta_{U_J/\Gamma_3, r}$ fitting in the commutative square

$$(4.3.2) \quad \begin{array}{ccc} S^*(\oplus_{\ell=0}^{r-1}(\mathbf{u}_J/\gamma_3)^{\#(r)}[2p^{r-\ell-1}]) & \xrightarrow{q_{U_J/\Gamma_3, r}} & k[V_r(U_J/\Gamma_3)] \\ \downarrow & & \downarrow \theta_{U_J/\Gamma_3, r} \\ S^*((U_J/\Gamma_3)_{(r)}) & \longrightarrow & \overline{S}^*((U_J/\Gamma_3)_{(r)}). \end{array}$$

The fact that the kernel of $q_{U_J/\Gamma_3, r}$ is the intersection of I_3 with $S^*(\oplus_{\ell=0}^{r-1}(\mathbf{u}_J/\gamma_3)^{\#(r)}[2p^{r-\ell-1}])$ (also generated by the intersection of $J_2 \subset S^*(\oplus_{\ell=0}^{r-1}(\mathbf{u}_J/\gamma_2)^{\#(r)}[2p^{r-\ell-1}])$) implies the injectivity of $\theta_{U_J/\Gamma_3, r}$.

The equality $\eta_{U_3, r} = \phi_{U_3, r}$ of Proposition 3.6 together with the surjectivity of $q_{U_J/\Gamma_3, r}$ implies the equality $\overline{\eta}_{U_3, r} \circ \theta_{U_3, r} = \overline{\phi}_{U_3, r}$. \square

We next observe that the tensor product decomposition of Proposition 4.2 is respected by the map θ_{U_J/Γ_3} . This enables us to show in the following proposition that $\theta_{U_3, r} : k[V_r(U_3)] \rightarrow \overline{S}^*((U_3)_{(r)})$ is a finite map of integral domains.

Proposition 4.4. *The map $\theta_{U_3, r}$ of Proposition 4.3 can be written as the tensor product*

$$\begin{aligned} & \overline{\theta}_{U_3, r} \otimes 1 : \overline{k[V_r(U_3/\Gamma_3)]} \otimes S^*((\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}]) \rightarrow \\ & \rightarrow Q^*((U_3/\Gamma_2)_{(r)}) \otimes S^*((\gamma_2)^{\#(r)}[2p^{r-\ell-1}]) = \overline{S}^*((U_3)_{(r)}). \end{aligned}$$

The map $\overline{\theta}_{U_3, r} : \overline{k[V_r(U_3)]} \rightarrow Q^*((U_3/\Gamma_2)_{(r)})$ is a finite map of integral domains of degree $p^{\frac{(r+2)(r-1)}{2}}$ obtained by taking $p^{r-\ell-1}$ -st roots of $X^{1,2}(\ell), X^{2,3}(\ell)$ for each $\ell, 0 \leq \ell < r$. Thus, $\theta_{U_3, r} : k[V_r(U_3)] \rightarrow \overline{S}^*((U_3)_{(r)})$ is a finite map of integral domains.

Proof. The fact that $\theta_{U_J/\Gamma_3, r}$ is a tensor product of the form $\overline{\theta}_{U_3, r} \otimes 1$ arises from the fact that the tensor decomposition of $k[V_r(U_3)]$ in Proposition 4.2 and that of $\overline{S}^*((U_3)_{(r)})$ in (3.10.1) both arise because the relations do not involve weights of level 2. The fact that $\theta_{U_J/\Gamma_3, r}$ is essentially the identity on $S^*((\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}])$ can be traced back to the definition of $X^{i,j}(\ell) \in \mathfrak{g}_N^{\#(r)}[2p^{r-\ell-1}]$ prior to Theorem 1.2.

The k -algebra $k[V_r(U_3)]$ is an integral domain by Proposition 4.2. Arguing as in the proof of Proposition 4.2, we verify that $Q^*((U_3/\Gamma_2)_{(r)}[(x_\alpha^{(0)})^{-1}])$ is the localization of the polynomial algebra on generators $x_{\alpha'}^{(\ell)}, 0 \leq \ell < r; Y^{1,2}(0)$ with $x_\alpha^{(0)}$ inverted. Thus, to show that $Q^*((U_3/\Gamma_2)_{(r)})$ is a domain it suffices to show that the localization map $Q^*((U_3/\Gamma_2)_{(r)}) \rightarrow Q^*((U_3/\Gamma_2)_{(r)}[(x_\alpha^{(0)})^{-1}])$ is injective. This is verified by examining the relations (3.3.2) to show that $x_\alpha^{(0)} \in Q^*((U_3/\Gamma_2)_{(r)})$ is not a zero-divisor.

Because $k[V_r(U_3)]$ is a domain, $F^r = \psi_{U_{3,r}} \circ \bar{\eta}_{U_{3,r}} \circ \theta_{U_{3,r}} : k[V_r(U_3)] \rightarrow k[V_r(U_3)]$ is injective and thus $\theta_{U_{3,r}}$ is injective. Since $S^*(\bigoplus_{\ell=0}^{r-1} \mathfrak{u}_3)^{\#(r)}[2p^{r-\ell-1}] \rightarrow S^*((U_3)_{(r)})$ is obtained by taking $p^{r-\ell-1}$ -st roots of $(x_\alpha^{(\ell)})^{p^{r-\ell-1}}, (x_{\alpha'}^{(\ell)})^{p^{r-\ell-1}}$ for each $\ell, 0 \leq \ell < r$, we conclude that $Q^*((U_3/\Gamma_2)_{(r)})$ is similarly obtained from $\overline{k[V_r(U_3/\Gamma_2)]}$.

To compute the degree of $\bar{\theta}_{U_{3,r}}$, we consider the map $\overline{k[V_r(U_3/\Gamma_2)]}[(X^{1,2}(0))^{-1}] \rightarrow Q^*((U_3/\Gamma_2)_{(r)}[(x_\alpha^{(0)})^{-1}])$ and utilize the facts that $x_{\alpha'}^{(\ell)}$ is the $p^{r-\ell-1}$ -th root of the image of $X^{2,3}(\ell)$ and that $x_\alpha^{(0)}$ is the p^{r-1} -st root of the image of $X^{1,2}(0)$ (using the notation of Proposition 4.2).

Finally, since $\theta_{U_{3,r}} = \bar{\theta}_{U_{3,r}} \otimes 1$, the fact that $\bar{\theta}_{U_{3,r}}$ is a finite map of integral domains implies that $\theta_{U_{3,r}}$ is also a finite map of integral domains \square

The following theorem summarizes what we know about $\bar{\eta}_{U_{3,r}}$.

Theorem 4.5. *Retain the hypotheses and notation of Proposition 3.1.*

- (1) $\bar{\eta}_{U_{3,r}} = \phi_{U_{3,r}}$.
- (2) $\bar{\eta}_{U_{3,r}} : \bar{S}^*((U_3)_{(r)}) \rightarrow H^\bullet((U_3)_{(r)}, k)$ is injective.
- (3) $\bar{\eta}_{U_{3,r}}$ is surjective onto p^r -th powers of elements of $H^\bullet((U_3)_{(r)}, k)$.
- (4) $gr(\bar{\eta}_{U_{3,r}})$ factors through

$$f_Q \otimes g : \bar{S}^*((U_3)_{(r)}) \simeq Q^*((U_3/\Gamma_2)_{(r)}) \otimes S^*(\bigoplus_{\ell=0}^{r-1} (\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}]) \rightarrow E_\infty^{*,0} \otimes E_\infty^{0,*}.$$

- (5) $gr(\bar{\eta}_{U_{3,r}}) : \bar{S}^*((U_3)_{(r)}) \rightarrow gr\{H^\bullet((U_3)_{(r)}, k)\}$ is injective.
- (6) $gr(\bar{\eta}_{U_J/\Gamma_{3,r}})$ is surjective onto p -th powers of elements of $gr\{H^\bullet((U_3)_{(r)}, k)\}$.

Proof. The equality $\bar{\eta}_r = (\phi_{U_{3,r}})_|$ of (3.6.2) implies that $\bar{\eta}_{U_{3,r}} = \phi_{U_{3,r}}$, since $\eta_{U_{3,r}}, \phi_{U_{3,r}}$ are determined by the Frobenius twists of the basic maps $\bar{\eta}_r, (\phi_{U_{3,r}})_|$.

By Proposition 1.9, $F^r = \psi_{U_{3,r}} \circ \bar{\phi}_{U_{3,r}} : k[V_r(U_3)] \rightarrow k[V_r(U_3)]$. Since $k[V_r(U_3)]$ is a domain by Proposition 4.2, F^r and thus also $\bar{\phi}_{U_{3,r}}$ are injective. Since the p^{r-1} -st power of each element in $S^*((U_3)_{(r)})$ lies in the image of $S^*(\bigoplus_{\ell=0}^{r-1} \mathfrak{u}_3)^{\#(r)}[2p^{r-\ell-1}]$, the commutativity of (4.3.2) and the surjectivity of $S^*((U_J/\Gamma_3)_{(r)}) \rightarrow \bar{S}^*((U_J/\Gamma_3)_{(r)})$ imply that the p^{r-1} -st power of an element in $\bar{S}^*((U_3)_{(r)})$ lies in the image of $\theta_{U_{3,r}}$. Thus, any element in the kernel of $\bar{\eta}_{U_{3,r}}$ must have p^{r-1} -st power which is in the kernel of $\bar{\phi}_{U_{3,r}}$ which we have observed is trivial. Since $\bar{S}^*((U_3)_{(r)})$ is a domain by Proposition 4.4, we conclude that $\bar{\eta}_{U_{3,r}}$ must be injective.

The surjectivity statement of (3) follows from the fact that the composition $F^r = \psi_{U_{3,r}} \circ \bar{\phi}_{U_{3,r}}$ is surjective onto p^r -th powers.

The factorization of $f \circ g$ through $f \otimes g : S^*((U_3)_{(r)}) \otimes S^*(\bigoplus_{\ell=0}^{r-1} (\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}]) \rightarrow E_\infty^{*,0} \otimes E_\infty^{0,*}$ is given by the proof of Proposition 3.9. This is easily seen to determine the factorization of $gr(\bar{\eta}_{U_{3,r}})$ as asserted in (4).

Granted the injectivity (1), to prove the injectivity of $gr(\bar{\eta}_{U_{3,r}})$ as asserted in (5) it suffices to show that both \tilde{f}_Q and \tilde{g} are filtration level preserving. Since the filtration level on $Q((U_J/\Gamma_2)_{(r)})$ equals the cohomological degree and \tilde{f}_Q can only increase filtration level (since it is a map of filtered algebras), \tilde{f}_Q preserves filtration level. Since the composition of $\tilde{g} : S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}]) \rightarrow H^\bullet((U_3)_{(r)}, k)$ with the natural map $E_\infty^{0,*}$ is injective, we conclude that \tilde{g} sends every non-zero element to a cohomology class of filtration level 0 and therefore must also preserve filtration level.

Finally, to prove that $gr(\bar{\eta}_{U_{3,r}})$ is surjective onto p -th powers, it suffices to prove that the p -th power of any permanent cycle, $z \in Z_2^{*,*} \subset E_2^{*,*}$, lies in the image of $S^*((U_3)_{(r)})$. Write $z = x + y$ with x defined as the sum of those summands of z lying in $S^*(\oplus_{\ell=0}^{r-1}(\mathbf{u}_3)^{\#(\ell+1)}[2])$ and y as the sum of the remaining summands of z , where z is written as a sum of terms using the decomposition (3.1.4). Then $y^p = 0 \in E_2^{*,*}$, so that $z^p = x^p \in Z_2^{*,*} \cap S^*(\oplus_{\ell=0}^{r-1}(\mathbf{u}_3)^{\#(\ell+1)}[2])$. By Corollary 2.4, either z^p is a boundary or $z^p \in S^*((U_3)_{(r)})$. This completes the proof of (6). \square

Observe that the (adjoint) action of T on $S^*((U_J)_{(r)})$ induces an action on $\bar{S}^*((U_J)_{(r)}) = S^*((U_J)_{(r)})/I_3$ which restricts to an action on the tensor factor $Q((U_J/\Gamma_2)_{(r)})$ of $\bar{S}^*((U_J)_{(r)})$ because the relations (3.2.2) are generated by T -eigenvalues. The following proposition is a consequence of the description of $\theta_{U_{3,r}}$ given in Proposition 4.4.

Proposition 4.6. *With respect to the above action, we have the equality*

$$H^0(T_{3(r)}, Q((U_3/\Gamma_2)_{(r)})) = \overline{k[V_r(U_3)]}.$$

Consequently, the restriction of $\bar{\eta}_{U_{3,r}} : \bar{S}^*((U_3)_{(r)}) \rightarrow H^\bullet((U_3)_{(r)}, k)$ to $(T_3)_{(r)}$ -invariants has the form

$$(4.6.1) \quad k[V_r(U_3)] \rightarrow H^\bullet((B_3)_{(r)}, k).$$

Proof. By definition of $X^{i,j}(\ell)$ as an element of $\mathfrak{gl}_N^{\#(r)}[2p^{r-\ell-1}]$ (see Theorem 1.2), we see that the images in $\overline{k[V_r(U_3)]} \subset Q((U_3/\Gamma_2)_{(r)})$ of $X^{i,j}(\ell)$ for $1 \leq i < j \leq 3$ are $T_{3(r)}$ invariant. On the other hand, using Proposition 4.4 we write $Q((U_3/\Gamma_2)_{(r)})$ as a free module over $\overline{k[V_r(U_3)]}$ generated by powers of the $p^{r-\ell-1}$ -st roots of the images of $X^{i,j}(\ell)$ viewed as images of elements of $S^*(\oplus_{\ell=0}^{r-1}\mathbf{u}^{\#(\ell+1)}[2])$. If a power of such a root is not divisible by $p^{r-\ell-1}$, then it is a non-trivial eigenvector for the semi-simple action of $T_{3(r)}$. This implies the asserted equality.

Since $\bar{S}^*((U_3)_{(r)}) \simeq Q((U_3/\Gamma_2)_{(r)}) \otimes S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}])$, we conclude that $H^0(T_{3(r)}, \bar{S}^*((U_3)_{(r)}))$ equals $\overline{k[V_r(U_3)]} \otimes S^*(\oplus_{\ell=0}^{r-1}(\gamma_2/\gamma_3)^{\#(r)}[2p^{r-\ell-1}])$ which equals $k[V_r(U_3)]$ by Proposition 4.2. The spectral sequence for the extension

$$1 \rightarrow (U_3)_{((r)} \rightarrow (B_3)_{(r)} \rightarrow (T_3)_{(r)} \rightarrow 1$$

and the semi-simplicity of $(T_3)_{(r)}$ imply the equality

$$H^0((T_3)_{(r)}, H^*((U_3)_{(r)}, k)) = H^*((B_3)_{(r)}, k).$$

This implies that $H^0(T_{(r)}, -)$ applied to $\bar{\eta}_{U_{3,r}}$ yields $k[V_r(U_3)] \rightarrow H^\bullet((B_3)_{(r)}, k)$. \square

5. QUESTIONS

Here are a few of the many questions encountered, but not answered, in this paper.

Question 5.1. *For V as in Theorem 2.2, does there exist a non-nilpotent cohomology class $\alpha \in H^*(V, k)$ each of whose restrictions $i_r^*(\alpha) \in H^*(V_{(r)}, k)$ is nilpotent?*

Question 5.2. *Under what conditions on the unipotent group V is the image of the restriction map $H^*(V, k) \rightarrow H^*(V_{(r)}, k)$ finitely generated?*

Question 5.3. *Are there natural Steenrod operations on the Andersen-Jantzen spectral sequence which satisfy the usual relationship with respect to differentials including the Kudo transgression theorem?*

Question 5.4. *Can one compare the LHS-filtration and the AJ-filtration on the Hochschild complex $C^*((U_J/\Gamma_3)_{(r)})$?*

Question 5.5. *Is the map $\bar{\phi}_{U_3, r} : k[V_r(U_3)] \rightarrow (H^*((U_3)_{(r)}, k))_{red}$ an isomorphism?*

Question 5.6. *Under what conditions on the unipotent algebraic group U is the k -algebra $k[V_r(U)]$ reduced (i.e., has no non-trivial nilpotent elements) for all $r > 0$?*

Question 5.7. *Can we establish the unipotent analogue of the “matrix p -th power relation”*

$$S_{i,j,\ell} = \sum_{t_1, \dots, t_{p-1}} X^{i,t_1}(\ell) \cdot X^{t_1,t_2}(\ell) \cdots X^{t_{p-1},j}(\ell)$$

using the AJ spectral sequence.

REFERENCES

- [1] H. Andersen, J. Jantzen, *Cohomology of induced representations for algebraic groups*, Math. Ann. **269** (1985), 487-525.
- [2] J. Alperin, L. Evens, *Representations, resolutions, and Quillen’s dimension theorem*, J. Pure and Appl. Algebra **22** (1981), 1 - 9.
- [3] H. Azad, M. Barry, G. Seitz, *On the structure of parabolic subgroups*, Comm. Algebra **18** (1990), 551-562.
- [4] J. Carlson, *The varieties and the cohomology ring of a module*, J. Algebra **85** (1983), 101-143.
- [5] J. Carlson, D. Nakano, *On the structure of cohomology rings of p -nilpotent Lie algebras*, Transform. Groups **19** (2104), 721-734.
- [6] E. Cline, B. Parshall, L. Scott, *Cohomology, hyperalgebras, and representations*, J. Alg. **63** (1980), 98 - 123.
- [7] E. Cline, B. Parshall, L. Scott, W. van der Kallen, *Rational and generic cohomology*, Invent. Math **39** (1977), 143-163.
- [8] C. Drupieski, D. Nakano, N. Ngo, *Cohomology for infinitesimal unipotent groups and quantum groups*, Transform. Groups **17**(2012), 393-416.
- [9] E. Friedlander, *Support varieties for rational representations*, Compositio Math **151** (2015), 765-792.
- [10] E. Friedlander, B. Barshall, *On the cohomology of algebraic and related finite groups*, Invent. Math **74** (1983), 85-117.
- [11] E. Friedlander, B. Barshall, *Cohomology of infinitesimal and discrete groups*, Math. Ann **273** (1986), 353-374.
- [12] E. Friedlander, J. Pevtsova, *Π -supports for modules for finite group schemes*, Duke. Math. J. **139** (2007), 317-368.
- [13] E. Friedlander, A. Suslin, *Cohomology of finite group schemes over a field*, Invent. Math. **127**(1997), 209-270.

- [14] S. Garibaldi, *Vanishing of trace forms in low characteristic*, Algebra and Number Theory **3** (2009), 543-566.
- [15] G. Hochschild, J.-P. Serre, *Cohomology of group extensions*, Trans. A.M.S. **74** (1953), 110-134.
- [16] J.C. Jantzen, *Representations of Algebraic groups. Second edition*, Math Surveys and Monographs **107**, AMS (2003).
- [17] T. Kudo, *A transgression theorem*, Mem. Fac. Sci. Kyusyu Univ **9** (1956), 79-81.
- [18] H. Matsumura, *Commutative Algebra*, Benjamin, N.Y (1970).
- [19] D. Quillen, *The spectrum of an equivariant cohomology ring: I*, Ann. of Math **94**(1971), 549-572.
- [20] D. Quillen, *The spectrum of an equivariant cohomology ring: II*, Ann. of Math **94**(1971), 573-602.
- [21] D. Quillen, *On the cohomology and K-theory of the general linear groups over a finite field*, Ann. of Math **96**(1972), 552-586.
- [22] P. Sobaje, *Exponentiation of commuting nilpotent varieties*, J. Pure and Appl. Algebra **219**, (2015), 2206 - 2217.
- [23] A. Suslin, E. Friedlander, C. Bendel, *Infinitesimal 1-parameter subgroups and cohomology*, J. Amer. Math. Soc. **10** (1997), 693-728.
- [24] A. Suslin, E. Friedlander, C. Bendel *Support varieties for infinitesimal group schemes*, J. Amer. Math. Soc. **10** (1997), 729-759.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA
E-mail address: `ericmf@usc.edu`