

On discrimination between two close distribution tails.

Rodionov I. V.*

March 11, 2022

1 Introduction. Main result.

Statistics deals often with discrimination of close distributions based on censored or truncated data, in particular, for high-risk insurances and reliability problems. The situation when one observes data exceeding a pre-determined threshold is well-studied, see [1], [2], [3] and references therein. On the other hand statistics of extremes says that only higher order statistics should be used for discrimination of close distribution tails, wherein moderate sample values can be modeled with standard statistical tools. In particular, such approach for distributions from Gumbel maximum domain of attraction (for the definitions see [4]) is considered in [5], [6], [7]. As well, any estimators of the extreme value indices γ and ρ (see [8]) can be used also to discriminate the distribution tails. Notice that we do not assume belonging the corresponding distribution function to a maximum domain of attraction.

Definition 1 *The distribution functions F and G are said to be satisfied the condition $B(F, G)$ if for some $\varepsilon > 0$ and x_0*

$$\frac{1 - G(x)}{(1 - F(x))^{1-\varepsilon}} \text{ is nondecreasing with } x > x_0. \quad (1)$$

Denote by $\Theta(F_0)$ the class of continuous distribution functions F_1 satisfying either $B(F_1, F_0)$ or $B(F_0, F_1)$. Consider the simple hypothesis $H_0 : F = F_0$ and the alternative hypothesis $H_1 : F \in \Theta(F_0)$, where F_0 is continuous. Notice that if distribution functions F, G satisfy either $B(F, G)$ or $B(G, F)$ for some $\varepsilon > 0$ then it holds for all ε_1 , $0 < \varepsilon_1 < \varepsilon$. So denote

$$\varepsilon(F, G) = \max\{\varepsilon : F, G \text{ satisfy either } B(F, G) \text{ or } B(G, F) \text{ for } \varepsilon\}.$$

*Moscow State University, Faculty of Mathematics and Mechanics and Moscow Institute of Physics and Technology, Faculty of Innovations and High Technologies. E-mail: vecsell@gmail.com

Denote by $\Theta_\varepsilon(F_0)$ the class of continuous distribution functions F_1 satisfying either $B(F_1, F_0)$ or $B(F_0, F_1)$ with $\varepsilon(F_0, F_1) \geq \varepsilon$ and consider another alternative hypothesis $H_{1,\varepsilon} : F \in \Theta_\varepsilon(F_0)$. Let X_1, \dots, X_n be i.i.d. random variables with a common distribution function F . Denote by $X_{(1)} \leq \dots \leq X_{(n)}$ the order statistics for them. Introduce the Hill-like statistics

$$R_{k,n} = \ln(1 - F_0(X_{(n-k)})) - \frac{1}{k} \sum_{i=n-k+1}^n \ln(1 - F_0(X_{(i)})).$$

which we are going to use for the problem of discrimination between the two introduced above hypotheses when k higher order statistics are known. Remark that if F_0 is Pareto distribution function with parameter γ , then

$$R_{k,n} \stackrel{d}{=} \gamma_H / \gamma,$$

where γ_H is the Hill estimator of γ . If furthermore F_0 belongs to Fréchet max-domain of attraction, then $R_{k,n}$ behaves asymptotically as γ_H / γ , that is, their ratio tends to one as $n \rightarrow \infty$. We will show that the distributions of $R_{k,n}$ if either H_0 or H_1 fulfilled are different which can give a statistical for discrimination the hypotheses. The following two results describe the behavior of $R_{k,n}$ as $k, n \rightarrow \infty$ with $k < n$ provided H_0 or H_1 is fulfilled.

Theorem 1 *If H_0 holds then*

$$\sqrt{k}(R_{k,n} - 1) \xrightarrow{d} \xi \text{ as } k, n \rightarrow \infty,$$

where ξ is standard normal random variable, i.e. $\xi \sim N(0, 1)$.

This theorem gives obvious goodness-of-fit test for the tail of F . Besides, the following result provides some information about the consistency of this test. Assume that H_0 does not hold and F is equal to F_1 which is different from F_0 . Denote x^* , the right endpoint of F_1 , that is, $x^* = \sup\{x : F_1(x) < 1\}$. Assume that F_0 and any $F_1 \in \Theta(F_0)$ have the same right endpoint (how to discriminate distributions with different endpoints, see [10], [4]). Further consider $x^* = +\infty$, otherwise change variables $y = 1/(x^* - x)$ gives the assumption. The following theorem shows consistency of the proposed test.

Theorem 2 (i) *If H_1 holds then*

$$\sqrt{k_n}|R_{k_n,n} - 1| \xrightarrow{d} +\infty$$

provided $k_n \rightarrow \infty$, $k_n/n \rightarrow 0$ as $n \rightarrow \infty$.

(ii) *If $H_{1,\varepsilon}$ holds then under the same conditions*

$$\inf_{F_1 \in \Theta_\varepsilon(F_0)} \sqrt{k_n}|R_{k_n,n} - 1| \xrightarrow{d} +\infty.$$

The considered test makes it possible to discriminate, for example, two normal distributions with different variances, but we should weaken the condition (1) to discriminate two normal distributions with the same variance and different means. But weakening the condition (1) imposes some conditions on behavior of the sequence k_n .

Definition 2 *The distribution functions F and G are said to satisfy the condition $C(F, G)$ if for some $\varepsilon > 0$ and x_0*

$$\frac{1 - G(x)}{(1 - F(x))(-\ln(1 - F(x)))^\varepsilon} \text{ is nondecreasing, } x > x_0. \quad (2)$$

Denote by $\Theta'(F_0)$ the class of continuous distribution functions F_1 satisfying either $C(F_1, F_0)$ or $C(F_0, F_1)$ and the following condition: for some $\delta \in (0, 1)$

$$1 - F_1(x) \leq (1 - F_0(x))^\delta, \quad x > x_0. \quad (3)$$

See, if distribution functions F, G satisfy either $C(F, G)$ or $C(G, F)$ for some $\varepsilon > 0$ then it holds for all $\varepsilon_1, 0 < \varepsilon_1 < \varepsilon$. Denote

$$\varepsilon'(F, G) = \max\{\varepsilon : F, G \text{ satisfy either } C(F, G) \text{ or } C(G, F) \text{ for } \varepsilon\}.$$

Denote by $\Theta'_\varepsilon(F_0)$ the class of continuous distribution functions F_1 satisfying (3) and either $C(F_1, F_0)$ or $C(F_0, F_1)$ with $\varepsilon'(F_0, F_1) \geq \varepsilon$. As before, consider the simple hypothesis $H_0 : F = F_0$ and two alternative hypotheses $H'_1 : F \in \Theta'(F_0)$, $H'_{1,\varepsilon} : F \in \Theta'_\varepsilon(F_0)$ with continuous F_0 .

Theorem 3 (i) *If H'_1 holds then*

$$\sqrt{k_n}|R_{k_n,n} - 1| \xrightarrow{d} +\infty$$

provided $k_n/n \rightarrow 0, k_n^{1/2-\alpha}/\ln n \rightarrow +\infty$, for some $\alpha \in (0, 1/2)$, as $n \rightarrow \infty$.

(ii) *If $H'_{1,\varepsilon}$ holds then under the same conditions*

$$\inf_{F_1 \in \Theta'_\varepsilon(F_0)} \sqrt{k_n}|R_{k_n,n} - 1| \xrightarrow{d} +\infty.$$

2 Auxiliary results and proofs.

2.1 Auxiliary results.

Since R_n depends on the higher order statistics we cannot immediately use independence of the random variables (X_1, \dots, X_n) . Therefore consider the conditional distribution of R_n given $X_{(n-k)} = q$ applying the following lemma.

Lemma 1 ([4]) *Let X, X_1, \dots, X_n be i.i.d. random variables with common distribution function F , and let $X_{(1)} \leq \dots \leq X_{(n)}$ be the n th order statistics. For any $k = 1 \dots n-1$, the conditional joint distribution of $\{X_{(i)}\}_{i=n-k+1}^n$ given $X_{(n-k)} = q$ is equal to the (unconditional) joint distribution of the corresponding set $\{X_{(i)}^*\}_{i=1}^k$ of order statistics for i.i.d. random variables $\{X_i^*\}_{i=1}^k$ having the distribution function*

$$F_q(x) = P(X \leq x | X > q) = \frac{F(x) - F(q)}{1 - F(q)}, \quad x > q.$$

We call $F_q(x)$, $x > q$, the tail distribution function linked with the distribution function F . Consider two continuous distribution functions F and G and a random variable ξ_q with distribution function G_q , where $q \in \mathbb{R}$ is some parameter. Let

$$\eta_q = \ln \left(\frac{1 - F(q)}{1 - F(\xi_q)} \right).$$

Clear, $\eta_q \geq 0$ for all $q \in \mathbb{R}$.

The crucial point in the proof of Theorem 2 is studying of asymptotical behavior of η_q .

Proposition 1 *Let F_q and G_q are tail distribution functions of F and G respectively. Then*

- (i) *If for some x_0 , $q > x_0$, and any $x > q$, $F_q(x) = G_q(x)$, then η_q is standard exponential.*
- (ii) *$G_q(x) \geq F_q(x)$ for any $x > q$ if and only if η_q is stochastically smaller than a standard exponential random variable.
 $G_q(x) \leq F_q(x)$ for any $x > q$ if and only if η_q is stochastically larger than a standard exponential random variable.*
- (iii) *$G_q(x) \geq F_q(x)$ for any $x > q \geq x_0$ and some x_0 if and only if $(1 - G(x))/(1 - F(x))$ is nonincreasing function as $x > x_0$.*

2.2 Proof of Proposition 1.

(i) Let $F_q(x) = G_q(x)$ for all $x > q$, then we have for the distribution function of η_q ,

$$\begin{aligned} P(\eta_q \leq y) &= P\left(\ln \left(\frac{1 - F(q)}{1 - F(\xi_q)} \right) \leq y\right) = P\left(\frac{1 - F(q)}{1 - F(\xi_q)} \leq e^y\right) = \\ &= P\left(F(\xi_q) \leq 1 - (1 - F(q))e^{-y}\right) = P\left(\xi_q \leq F^{\leftarrow}\left(1 - \frac{1 - F(q)}{e^y}\right)\right). \end{aligned} \quad (4)$$

Furthermore, for the same x ,

$$P\left(\xi_q \leq F^{\leftarrow}\left(1 - \frac{1 - F(q)}{e^y}\right)\right) = \frac{F\left(F^{\leftarrow}\left(1 - \frac{1 - F(q)}{e^y}\right)\right) - F(q)}{1 - F(q)} = 1 - e^{-y}.$$

(ii) Now assume that for all $x > q$ and some $q \in \mathbb{R}$, $G_q(x) \geq F_q(x)$. Then from (4), since $1 - (1 - F(q))e^{-y} \geq F(q)$ for all $y \geq 0$ it follows that

$$P(\eta_q \leq y) = \frac{G\left(F^{\leftarrow}\left(1 - \frac{1 - F(q)}{e^y}\right)\right) - G(q)}{1 - G(q)} \geq$$

$$\frac{F\left(F^{\leftarrow}\left(1 - \frac{1-F(q)}{e^y}\right)\right) - F(q)}{1 - F(q)} = 1 - e^{-y}. \quad (5)$$

Conversely, assume that η_q is stochastically smaller than a standard exponential random variable, that is, $P(\eta_q \leq y) \geq 1 - e^{-y}$ for all $y \geq 0$. With (4) we get that

$$\begin{aligned} \frac{G\left(F^{\leftarrow}\left(1 - \frac{1-F(q)}{e^y}\right)\right) - G(q)}{1 - G(q)} &\geq 1 - e^{-y} \iff \frac{1 - G\left(F^{\leftarrow}\left(1 - \frac{1-F(q)}{e^y}\right)\right)}{1 - G(q)} \leq e^{-y} \\ &\iff G\left(F^{\leftarrow}\left(1 - \frac{1 - F(q)}{e^y}\right)\right) \leq 1 - \frac{1 - G(q)}{e^y} \iff \\ &F^{\leftarrow}\left(1 - \frac{1 - F(q)}{e^y}\right) \leq G^{\leftarrow}\left(1 - \frac{1 - G(q)}{e^y}\right). \end{aligned}$$

Denote $z_F = F^{\leftarrow}(1 - e^{-y}(1 - F(q)))$ and $z_G = G^{\leftarrow}(1 - e^{-y}(1 - G(q)))$. Since $F(z_F) = 1 - e^{-y}(1 - F(q))$ and $G(z_G) = 1 - e^{-y}(1 - G(q))$, we have,

$$e^{-y} = \frac{1 - G(z_G)}{1 - G(q)} = \frac{1 - F(z_F)}{1 - F(q)}.$$

Further, since $z_F \leq z_G$ then

$$\frac{1 - F(z_F)}{1 - F(q)} = \frac{1 - G(z_G)}{1 - G(q)} \leq \frac{1 - G(z_F)}{1 - G(q)}.$$

This observation completes the proof since $z_F \in [q, \infty)$. The proof of the second assertion is similar.

(iii) We have,

$$\frac{G(x) - G(q)}{1 - G(q)} \geq \frac{F(x) - F(q)}{1 - F(q)} \quad \forall x > q \geq x_0 \iff \frac{1 - G(x)}{1 - G(q)} \leq \frac{1 - F(x)}{1 - F(q)} \quad \forall x > q \geq x_0 \iff$$

$$\frac{1 - G(x)}{1 - F(x)} \leq \frac{1 - G(q)}{1 - F(q)} \quad \forall x > q \geq x_0 \iff \frac{1 - G(x)}{1 - F(x)} \text{ is nonincreasing for all } x > x_0. \blacksquare$$

2.3 Proof of Theorem 1.

Under the conditions of Theorem 1, $F_0(X_1)$ is uniformly distributed on $[0, 1]$, that is, $F_0(X_1) \sim U[0, 1]$, hence $-\ln(1 - F_0(X))$ is standard exponential random variable. It follows from Rényi's representation (see [4]), that

$$\left\{-\ln(1 - F_0(X_{(n-i)})) + \ln(1 - F_0(X_{(n-k)}))\right\}_{i=0}^{k-1} \stackrel{d}{=} \left\{\sum_{j=i+1}^k \frac{E_{n-j+1}}{j}\right\}_{i=0}^{k-1},$$

where E_1, E_2, \dots are independent standard exponential variables. Therefore the distribution of the left-hand side does not depend on n and

$$\{-\ln(1 - F_0(X_{(n-i)})) + \ln(1 - F_0(X_{(n-k)}))\}_{i=0}^{k-1} \stackrel{d}{=} \{E_{(k-i)}\}_{i=0}^{k-1},$$

where $E_{(1)} \leq \dots \leq E_{(k)}$ are the n th order statistics of the sample $\{E_i\}_{i=1}^k$. Finally we have,

$$\sqrt{k}(R_{k,n} - 1) \stackrel{d}{=} \sqrt{k} \left(\frac{1}{k} \sum_{i=0}^{k-1} E_{(k-i)} - 1 \right) = \sqrt{k} \left(\frac{1}{k} \sum_{j=1}^k E_j - 1 \right),$$

and the assertion follows from the Central Limit Theorem.

2.4 Proof of Theorem 2.

We first prove (i). The steps of the proof are similar to corresponding steps in [6] and [7]. Consider asymptotic behavior of $R_{k_n,n}$ as $n \rightarrow \infty$. Denote

$$Y_i = \ln(1 - F_0(q)) - \ln(1 - F_0(X_i^*)),$$

where $\{X_i^*\}_{i=1}^{k_n}$ are i.i.d. random variables introduced in Lemma 1 with the distribution function

$$F_q(x) = \frac{F_1(x) - F_1(q)}{1 - F_1(q)}, \quad q < x.$$

Taking $F = F_0$ and $G = F_1$ we have, $Y_i \stackrel{d}{=} \eta_q$, $i \in \{1, \dots, k_n\}$. Notice that, in view of Lemma 1, the joint distribution of order statistics $\{Y_{(i)}\}_{i=1}^{k_n}$ of the sample $\{Y_j\}_{j=1}^{k_n}$ is equal to the joint conditional distribution of order statistics $\{Z_{(j)}\}_{j=1}^{k_n}$ of $\{Z_j\}_{j=1}^{k_n}$ given $X_{(n-k_n)} = q$, where

$$Z_j = \ln(1 - F_0(X_{(n-k_n)})) - \ln(1 - F_0(X_{(n-j+1)})), \quad j = 1, \dots, k_n.$$

Clear,

$$R_{n,k_n} = \frac{1}{k_n} \sum_{i=1}^{k_n} Z_i.$$

So, the conditional distribution of $R_{k_n,n}$ given $X_{(n-k)} = q$ is equal to the distribution of $\frac{1}{k_n} \sum_{i=1}^{k_n} Y_i$. Further, distribution functions F_1 and F_0 satisfy $B(F_0, F_1)$ or $B(F_1, F_0)$. First suppose that the condition $B(F_0, F_1)$ holds for some $\varepsilon > 0$ and x_0 . Since $x^* = +\infty$, $X_{(n-k_n)} \rightarrow +\infty$ a.s., we may consider the case $q > x_0$ only. Proposition 1 (iii) implies, that

$$\frac{1 - F_1(x)}{1 - F_1(x_0)} \geq \frac{(1 - F_0(x))^{1-\varepsilon}}{(1 - F_0(x_0))^{1-\varepsilon}}, \quad x > x_0.$$

With (5), we get that,

$$P(Y_1 \leq x) = 1 - \frac{1 - F_1 \left(F_0^{\leftarrow} \left(1 - \frac{1 - F_0(q)}{e^x} \right) \right)}{1 - F_1(q)} \leq$$

$$1 - \frac{\left(1 - F_0 \left(F_0^{\leftarrow} \left(1 - \frac{1 - F_0(q)}{e^x} \right) \right) \right)^{1-\varepsilon}}{(1 - F_0(q))^{1-\varepsilon}} = 1 - e^{-(1-\varepsilon)x},$$

hence Y_1 is stochastically larger than a random variable $E \sim \text{Exp}(1 - \varepsilon)$, write $Y_1 \gg E$. Further, let E_1, \dots, E_{k_n} are i.i.d. random variables with distribution function $H(x) = 1 - e^{-(1-\varepsilon)x}$, then

$$\sqrt{k_n} \left(\frac{1}{k_n} \sum_{i=1}^{k_n} Y_i - 1 \right) \gg \sqrt{k_n} \left(\frac{1}{k_n} \sum_{i=1}^{k_n} E_i - 1 \right). \quad (6)$$

Since (6) holds for all $q > x_0$, and $X_{(n-k_n)} \rightarrow +\infty$ a.s. as $n \rightarrow \infty$, we have under the conditions of Theorem 2, that

$$\sqrt{k_n}(R_{k_n,n} - 1) \gg \sqrt{k_n} \left(\frac{1}{k_n} \sum_{i=1}^{k_n} E_i - 1 \right). \quad (7)$$

It follows from Lindeberg-Feller theorem, that

$$(1 - \varepsilon) \sqrt{k_n} \left(\frac{1}{k_n} \sum_{i=1}^{k_n} E_i - \frac{1}{1 - \varepsilon} \right) \xrightarrow{d} \xi \sim N(0, 1), \quad n \rightarrow \infty,$$

therefore

$$\sqrt{k_n} \left(\frac{1}{k_n} \sum_{i=1}^{k_n} E_i - 1 \right) \xrightarrow{P} +\infty, \quad n \rightarrow \infty. \quad (8)$$

Finally, with (7), we have,

$$\sqrt{k_n}(R_{k_n,n} - 1) \xrightarrow{P} +\infty, \quad n \rightarrow \infty.$$

If the condition $B(F_0, F_1)$ holds, then

$$\sqrt{k_n}(R_{k_n,n} - 1) \xrightarrow{P} -\infty, \quad n \rightarrow \infty,$$

and the proof is similar. The second assertion easily follows from (7) and (8).

2.5 Proof of Theorem 3.

Firstly we prove (i). Denote $\overline{F}(x) = 1 - F(x)$. In notation of the proof of Theorem 2, find the distribution of Y_1 . First assume that $C(F_0, F_1)$ holds. With (5) and Proposition 1 (iii) we have,

$$P(Y_1 \leq x) = 1 - \frac{\overline{F}_1(\overline{F}_0^{\leftarrow}(\overline{F}_0(q)e^{-x}))}{\overline{F}_1(q)} \leq$$

$$1 - \frac{\overline{F}_0(q)e^{-x}(-\ln(\overline{F}_0(q)e^{-x}))^\varepsilon}{\overline{F}(q)(-\ln \overline{F}_0(q))^\varepsilon} = 1 - e^{-x} \left(1 + \frac{x}{-\ln \overline{F}_0(q)}\right)^\varepsilon.$$

For $\varepsilon, c \in (0, 1)$,

$$(1 + cx)^\varepsilon \geq 1 + c\varepsilon - c\varepsilon e^{-x}, \quad x \geq 0,$$

and $G(x) = 1 - e^{-x}(1 + c\varepsilon - c\varepsilon e^{-x})$ is the distribution function. Hence,

$$P(Y_1 \leq x) \geq 1 - e^{-x} - \frac{\varepsilon}{-\ln \overline{F}_0(q)}(1 - e^{-x}).$$

Further, let $\zeta, \zeta_1, \dots, \zeta_{k_n}$ be i.i.d. random variables with this distribution function. Therefore, like the proof of Theorem 2,

$$\sqrt{k_n} \left(\frac{1}{k_n} \sum_{i=1}^{k_n} Y_i - 1 \right) \gg \sqrt{k_n} \left(\frac{1}{k_n} \sum_{i=1}^{k_n} \zeta_i - 1 \right). \quad (9)$$

Clear,

$$E\zeta = 1 + \frac{\varepsilon}{-2 \ln \overline{F}_0(q)}, \quad Var \zeta = 1 - \left(\frac{\varepsilon}{-2 \ln \overline{F}_0(q)} \right)^2,$$

so we have,

$$\sqrt{k_n} \left(\frac{1}{k_n} \sum_{i=1}^{k_n} \zeta_i - 1 \right) = \sqrt{k_n} \left(\frac{1}{k_n} \sum_{i=1}^{k_n} \zeta_i - E\zeta \right) + \sqrt{k_n} \frac{\varepsilon}{-2 \ln \overline{F}_0(q)}. \quad (10)$$

Consider now the statistic $\sqrt{k_n}/\ln \overline{F}_0(X_{(n-k_n)})$. Denote $R_i = \overline{F}_1(X_i)$, $i = 1, \dots, n$. Since \overline{F}_1 is continuous, R_1, \dots, R_n are i.i.d. standard uniform random variables and $R_{(k_n)} = \overline{F}_1(X_{(n-k_n)})$. Theorem 2.2.1 [4] implies, that

$$\frac{n}{\sqrt{k_n}} \left(R_{(k_n)} - \frac{k_n}{n} \right) \xrightarrow{d} N(0, 1), \quad n \rightarrow \infty. \quad (11)$$

Using the delta method (see [11]) for the function $f(x) = -x/\ln x$, we have

$$\frac{n}{\sqrt{k_n}} \left(\frac{R_{(k_n)}}{-\ln R_{(k_n)}} - \frac{k_n/n}{-\ln(k_n/n)} \right) \xrightarrow{P} 0, \quad n \rightarrow \infty,$$

since under the conditions of theorem

$$f' \left(\frac{k_n}{n} \right) = -\frac{1}{\ln(n/k_n)} + \frac{1}{(\ln(n/k_n))^2} \rightarrow 0, \quad n \rightarrow \infty.$$

Further,

$$\begin{aligned} & \frac{n}{\sqrt{k_n}} \left(\frac{R_{(k_n)}}{\ln R_{(k_n)}} - \frac{k_n/n}{\ln(k_n/n)} \right) = \\ & \frac{n}{\sqrt{k_n}} \left(\frac{R_{(k_n)}}{\ln R_{(k_n)}} - \frac{k_n/n}{\ln(R_{(k_n)})} \right) + \sqrt{k_n} \left(\frac{1}{\ln R_{(k_n)}} - \frac{1}{\ln(k_n/n)} \right), \end{aligned}$$

and (11) implies that the first summand in the right hand side tends to 0 in probability. Therefore,

$$\sqrt{k_n} \left(\frac{1}{\ln R_{(k_n)}} - \frac{1}{\ln(k_n/n)} \right) \xrightarrow{P} 0, \quad n \rightarrow \infty,$$

and under the conditions of Theorem 3,

$$\frac{\sqrt{k_n}}{-\ln R_{(k_n)}} = \sqrt{k_n} \left(\frac{1}{-\ln R_{(k_n)}} - \frac{1}{-\ln(k_n/n)} \right) + \frac{\sqrt{k_n}}{-\ln(k_n/n)} \xrightarrow{P} +\infty, \quad n \rightarrow \infty.$$

On the other hand, from (3) it follows that

$$\frac{\sqrt{k_n}}{-\ln \bar{F}_0(X_{(n-k_n)})} = \frac{\sqrt{k_n}}{-\ln \bar{F}_0(\bar{F}_1^{\leftarrow}(R_{(k_n)}))} \geq \frac{\sqrt{k_n}}{-\delta^{-1} \ln R_{(k_n)}} \xrightarrow{P} +\infty \quad (12)$$

as $n \rightarrow \infty$. Further, it follows from the Law of large numbers for triangular arrays (see [9]), that for any $\epsilon > 0$

$$\sqrt{k_n} \left(\frac{1}{k_n} \sum_{i=1}^{k_n} \zeta_i - 1 \right) = o_P(k_n^\epsilon), \quad n \rightarrow \infty.$$

It means that the term in the left hand side is asymptotically smaller in probability than k_n^ϵ . Hence for any q , given $X_{(n-k_n)} = q$

$$\sqrt{k_n} \left(\frac{1}{k_n} \sum_{i=1}^{k_n} Y_i - 1 \right) \xrightarrow{P} +\infty, \quad n \rightarrow \infty,$$

and finally,

$$\sqrt{k_n} (R_{k_n, n} - 1) \xrightarrow{P} +\infty, \quad n \rightarrow \infty.$$

If the condition $C(F_1, F_0)$ holds, then

$$\sqrt{k_n} (R_{k_n, n} - 1) \xrightarrow{P} -\infty, \quad n \rightarrow \infty,$$

and the proof is the same. The second assertion clearly follows from (9), (10) and (12).

References

- [1] *Dufour R., Maag U.R.* Distribution Results for Modified Kolmogorov-Smirnov Statistics for Truncated or Censored Samples. — *Technometrics*, 1978, v. 20, p. 29–32.
- [2] *Guilbaud O.* Exact Kolmogorov-Type Test for Left-Truncated and/or Right-Censored Data. — *Journal of American Statistical Association*, 1998, v. 83, p. 213–221.

- [3] *Chernobai A., Menn C., Rachev S. T., Truck S.* Estimation of operational value-at-risk in the presence of minimum collection thresholds. — Tech. Rep., University of California, Santa Barbara, Calif, USA, 2005.
- [4] *Ferreira A., Haan L. de.* Extreme value theory. An introduction. N. Y.: Springer, Springer Series in Operations Research and Financial Engineering, 2006.
- [5] *Gardes L., Girard S., Guillou A.* Weibull tail-distributions revisited: a new look at some tail estimators. — Journal of Statistical Planning and Inference, 2009, v. 141, p. 429–444.
- [6] *Rodionov I. V.* A discrimination test for tails of Weibull-like distributions. — to appear in Probability Theory and its Applications.
- [7] *Rodionov I. V.* Discrimination of close hypotheses on distribution tails using higher order statistics. — to appear in Extremes.
- [8] *Haan L. de, Resnick S.* Second-order regular variation and rates of convergence in extreme value theory. — The Annals of Probability, 1996, v. 24, i. 1, p. 97–124.
- [9] *Mikusheva A. E.* The law of large numbers and the logarithmic law for arrays. — Fundamental and applied mathematics, 2000, v. 6, i. 1, p. 195–206 (in Russian).
- [10] *Falk M.* Some best estimators for distributions with finite endpoint. — Statistics, 1995, v. 27, p. 115–125.
- [11] *Resnick S. I.* A probability path. Birkhauser, 1999.