

How to Escape Saddle Points Efficiently

Chi Jin^{*} Rong Ge[†] Praneeth Netrapalli[‡] Sham M. Kakade[§]
 Michael I. Jordan[¶]

April 9, 2021

Abstract

This paper shows that a perturbed form of gradient descent converges to a second-order stationary point in a number of iterations which depends only poly-logarithmically on dimension (i.e., it is almost “dimension-free”). The convergence rate of this procedure matches the well-known convergence rate of gradient descent to first-order stationary points, up to log factors. When all saddle points are non-degenerate, all second-order stationary points are local minima, and our result thus shows that perturbed gradient descent can escape saddle points almost for free.

Our results can be directly applied to many machine learning applications, including deep learning. As a particular concrete example of such an application, we show that our results can be used directly to establish sharp global convergence rates for matrix factorization. Our results rely on a novel characterization of the geometry around saddle points, which may be of independent interest to the non-convex optimization community.

1 Introduction

Given a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$, gradient descent aims to minimize the function via the following iteration:

$$\mathbf{x}_{t+1} = \mathbf{x}_t - \eta \nabla f(\mathbf{x}_t),$$

where $\eta > 0$ is a step size. Gradient descent and its variants (e.g., stochastic gradient) are widely used in machine learning applications due to their favorable computational properties. This is notably true in the deep learning setting, where gradients can be computed efficiently via back-propagation [Rumelhart et al., 1988].

Gradient descent is especially useful in high-dimensional settings because the number of iterations required to reach a point with small gradient is independent of the dimension (“dimension-free”). More precisely, for a function that is ℓ -gradient Lipschitz (see Definition 1), it is well known that gradient descent finds an ϵ -first-order stationary point (i.e., a point \mathbf{x} with $\|\nabla f(\mathbf{x})\| \leq \epsilon$)

^{*}University of California, Berkeley. Email: chijin@cs.berkeley.edu

[†]Duke University. Email: rongge@cs.duke.edu

[‡]Microsoft Research, India. Email: praneeth@microsoft.com

[§]University of Washington. Email: sham@cs.washington.edu

[¶]University of California, Berkeley. Email: jordan@cs.berkeley.edu

within $\ell(f(\mathbf{x}_0) - f^*)/\epsilon^2$ iterations [Nesterov, 1998], where \mathbf{x}_0 is the initial point and f^* is the optimal value of f . This bound does not depend on the dimension of \mathbf{x} . In convex optimization, finding an ϵ -first-order stationary point is equivalent to finding an approximate global optimum.

In non-convex settings, however, convergence to first-order stationary points is not satisfactory. For non-convex functions, first-order stationary points can be global minima, local minima, saddle points or even local maxima. Finding a global minimum can be hard, but fortunately, for many non-convex problems, it is sufficient to find a local minimum. Indeed, a line of recent results show that, in many problems of interest, either all local minima are global minima (e.g., in tensor decomposition [Ge et al., 2015], dictionary learning [Sun et al., 2016a], phase retrieval [Sun et al., 2016b], matrix sensing [Bhojanapalli et al., 2016, Park et al., 2016], matrix completion [Ge et al., 2016], and certain classes of deep neural networks [Kawaguchi, 2016]). Moreover, there are suggestions that in more general deep networks most of the local minima are as good as global minima [Choromanska et al., 2014].

On the other hand, saddle points (and local maxima) can correspond to highly suboptimal solutions in many problems [see, e.g., Jain et al., 2015, Sun et al., 2016b]. Furthermore, Dauphin et al. [2014] argue that saddle points are ubiquitous in high-dimensional, non-convex optimization problems, and are thus the main bottleneck in training neural networks. Standard analysis of gradient descent cannot distinguish between saddle points and local minima, leaving open the possibility that gradient descent may get stuck at saddle points, either asymptotically or for a sufficiently long time so as to make training times for arriving at a local minimum infeasible. Ge et al. [2015] showed that by adding noise at each step, gradient descent can escape all saddle points in a polynomial number of iterations, provided that the objective function satisfies the strict saddle property (see Assumption A2). Lee et al. [2016] proved that under similar conditions, gradient descent with random initialization avoids saddle points even without adding noise. However, this result does not bound the number of steps needed to reach a local minimum.

Though these results establish that gradient descent can find local minima in a polynomial number of iterations, they are still far from being efficient. For instance, the number of iterations required in Ge et al. [2015] is at least $\Omega(d^4)$, where d is the underlying dimension. This is significantly suboptimal compared to rates of convergence to first-order stationary points, where the iteration complexity is dimension-free. This motivates the following question: **Can gradient descent escape saddle points and converge to local minima in a number of iterations that is (almost) dimension-free?**

In order to answer this question formally, this paper investigates the complexity of finding ϵ -second-order stationary points. For ρ -Hessian Lipschitz functions (see Definition 5), these points are defined as [Nesterov and Polyak, 2006]:

$$\|\nabla f(\mathbf{x})\| \leq \epsilon, \quad \text{and} \quad \lambda_{\min}(\nabla^2 f(\mathbf{x})) \geq -\sqrt{\rho\epsilon}.$$

Under the assumption that all saddle points are strict (i.e., for any saddle point \mathbf{x}_s , $\lambda_{\min}(\nabla^2 f(\mathbf{x}_s)) < 0$), all second-order stationary points ($\epsilon = 0$) are local minima. Therefore, convergence to second-order stationary points is equivalent to convergence to local minima.

This paper studies gradient descent with phasic perturbations (see Algorithm 1). For ℓ -smooth functions that are also Hessian Lipschitz, we show that perturbed gradient descent will converge to an ϵ -second-order stationary point in $\tilde{O}(\ell(f(\mathbf{x}_0) - f^*)/\epsilon^2)$, where $\tilde{O}(\cdot)$ hides polylog factors. This guarantee is almost dimension free (up to $\text{polylog}(d)$ factors), answering the above highlighted question affirmatively. Note that this rate is exactly the same as the well-known convergence rate

Algorithm 1 Perturbed Gradient Descent (Meta-algorithm)

```
for  $t = 0, 1, \dots$  do  
  if perturbation condition holds then  
     $\mathbf{x}_t \leftarrow \mathbf{x}_t + \xi_t, \quad \xi_t \text{ uniformly } \sim \mathbb{B}_0(r)$   
   $\mathbf{x}_{t+1} \leftarrow \mathbf{x}_t - \eta \nabla f(\mathbf{x}_t)$ 
```

of gradient descent to first-order stationary points [Nesterov, 1998], up to log factors. Furthermore, our analysis admits a maximal step size of up to $\Omega(1/\ell)$, which is the same as that in analyses for first-order stationary points.

As many real learning problems present strong *local* geometric properties, similar to strong convexity in the global setting [see, e.g. Bhojanapalli et al., 2016, Sun and Luo, 2016, Zheng and Lafferty, 2016], it is important to note that our analysis naturally takes advantage of such local structure. We show that when local strong convexity is present, the ϵ -dependence goes from a polynomial rate, $1/\epsilon^2$, to linear convergence, $\log(1/\epsilon)$. As an example, we show that sharp global convergence rates can be obtained for matrix factorization as a direct consequence of our analysis.

1.1 Our Contributions

This paper presents the first sharp analysis that shows that (perturbed) gradient descent finds an approximate second-order stationary point in at most $\text{polylog}(d)$ iterations, thus escaping all saddle points efficiently. Our main technical contributions are as follows:

- For ℓ -gradient Lipschitz, ρ -Hessian Lipschitz functions (possibly non-convex), gradient descent with appropriate perturbations finds an ϵ -second-order stationary point in $\tilde{O}(\ell(f(\mathbf{x}_0) - f^*)/\epsilon^2)$ iterations. This rate matches the well-known convergence rate of gradient descent to first-order stationary points up to log factors.
- Under a strict-saddle condition (see Assumption A2), this convergence result directly applies for finding local minima. This means that gradient descent can escape all saddle points with only logarithmic overhead in runtime.
- When the function has local structure, such as local strong convexity (see Assumption A3.a), the above results can be further improved to linear convergence. We give sharp rates that are comparable to previous problem-specific local analysis of gradient descent with smart initialization (see Section 1.2).
- All the above results rely on a new characterization of the geometry around saddle points: points from where gradient descent gets stuck at a saddle point constitute a thin “band.” We develop novel techniques to bound the volume of this band. As a result, we can show that after a random perturbation the current point is very unlikely to be in the “band”; hence, efficient escape from the saddle point is possible (see Section 5).

1.2 Related Work

Over the past few years, there have been many problem-specific convergence results for non-convex optimization. One line of work requires a smart initialization algorithm to provide a coarse estimate lying inside a local neighborhood, from which popular local search algorithms

Algorithm	Iterations	Oracle
Ge et al. [2015]	$O(\text{poly}(d/\epsilon))$	Gradient
Levy [2016]	$O(d^3 \cdot \text{poly}(1/\epsilon))$	Gradient
This Work	$O(\log^4(d)/\epsilon^2)$	Gradient
Agarwal et al. [2016]	$O(\log(d)/\epsilon^{7/4})$	Hessian-vector product
Carmon et al. [2016]	$O(\log(d)/\epsilon^{7/4})$	Hessian-vector product
Carmon and Duchi [2016]	$O(\log(d)/\epsilon^2)$	Hessian-vector product
Nesterov and Polyak [2006]	$O(1/\epsilon^{1.5})$	Hessian
Curtis et al. [2014]	$O(1/\epsilon^{1.5})$	Hessian

Table 1: Oracle model and iteration complexity to second-order stationary point

enjoy fast local convergence [see, e.g., Netrapalli et al., 2013, Candes et al., 2015, Sun and Luo, 2016, Bhojanapalli et al., 2016]. While there are not many results that show global convergence for non-convex problems, Jain et al. [2015] show that gradient descent yields global convergence rates for matrix square-root problems. Although these results give strong guarantees, the analyses are heavily tailored to specific problems, and it is unclear how to generalize them to a wider class of non-convex functions.

For general non-convex optimization, there are a few previous results on finding second-order stationary points. These results can be divided into the following three categories, where, for simplicity of presentation, we only highlight dependence on dimension d and ϵ , assuming that all other problem parameters are constant from the point of view of iteration complexity:

Hessian-based: Traditionally, only second-order optimization methods were known to converge to second-order stationary points. These algorithms rely on computing the Hessian to distinguish between first- and second-order stationary points. Nesterov and Polyak [2006] designed a cubic regularization algorithm which converges to an ϵ -second-order stationary point in $O(1/\epsilon^{1.5})$ iterations. Trust region algorithms [Curtis et al., 2014] can also achieve the same performance if the parameters are chosen carefully. These algorithms typically require the computation of the inverse of the full Hessian per iteration, which can be very expensive.

Hessian-vector-product-based: A number of recent papers have explored the possibility of using only Hessian-vector products instead of full Hessian information in order to find second-order stationary points. These algorithms require a Hessian-vector product oracle: given a function f , a point \mathbf{x} and a direction \mathbf{u} , the oracle returns $\nabla^2 f(\mathbf{x}) \cdot \mathbf{u}$. Agarwal et al. [2016] and Carmon et al. [2016] presented accelerated algorithms that can find an ϵ -second-order stationary point in $O(\log d/\epsilon^{7/4})$ steps. Also, Carmon and Duchi [2016] showed by running gradient descent as a subroutine to solve the subproblem of cubic regularization (which requires Hessian-vector product oracle), it is possible to find an ϵ -second-order stationary point in $O(\log d/\epsilon^2)$ iterations. In many applications such an oracle can be implemented efficiently, in roughly the same complexity as the gradient oracle. Also, when the function has a Hessian Lipschitz property such an oracle can be approximated by differ-

entiating the gradients at two very close points (although this may suffer from numerical issues, thus is seldom used in practice).

Gradient-based: Another recent line of work shows that it is possible to converge to a second-order stationary point without any use of the Hessian. These methods feature simple computation per iteration (only involving gradient operations), and are closest to the algorithms used in practice. Ge et al. [2015] showed that stochastic gradient descent could converge to a second-order stationary point in $\text{poly}(d/\epsilon)$ iterations, with polynomial of order at least four. This was improved in Levy [2016] to $O(d^3 \cdot \text{poly}(1/\epsilon))$ using normalized gradient descent. The current paper improves on both results by showing that perturbed gradient descent can actually find an ϵ -second-order stationary point in $O(\text{polylog}(d)/\epsilon^2)$ steps, which matches the guarantee for converging to first-order stationary points up to polylog factors.

2 Preliminaries

In this section, we will first introduce our notation, and then present some definitions and existing results in optimization which will be used later.

2.1 Notation

We use bold upper-case letters \mathbf{A}, \mathbf{B} to denote matrices and bold lower-case letters \mathbf{x}, \mathbf{y} to denote vectors. \mathbf{A}_{ij} means the $(i, j)^{\text{th}}$ entry of matrix \mathbf{A} . For vectors we use $\|\cdot\|$ to denote the ℓ_2 -norm, and for matrices we use $\|\cdot\|$ and $\|\cdot\|_F$ to denote spectral norm and Frobenius norm respectively. We use $\sigma_{\max}(\cdot), \sigma_{\min}(\cdot), \sigma_i(\cdot)$ to denote the largest, the smallest and the i -th largest singular values respectively, and $\lambda_{\max}(\cdot), \lambda_{\min}(\cdot), \lambda_i(\cdot)$ for corresponding eigenvalues.

For a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$, we use $\nabla f(\cdot)$ and $\nabla^2 f(\cdot)$ to denote its gradient and Hessian, and f^* to denote the global minimum of $f(\cdot)$. We use notation $O(\cdot)$ to hide only absolute constants which do not depend on any problem parameter, and notation $\tilde{O}(\cdot)$ to hide only absolute constants and log factors. We let $\mathbb{B}_{\mathbf{x}}^{(d)}(r)$ denote the d -dimensional ball centered at \mathbf{x} with radius r ; when it is clear from context, we simply denote it as $\mathbb{B}_{\mathbf{x}}(r)$. We use $\mathcal{P}_{\mathcal{X}}(\cdot)$ to denote projection onto the set \mathcal{X} . Distance and projection are always defined in a Euclidean sense.

2.2 Gradient Descent

The theory of gradient descent often takes its point of departure to be the study of convex minimization where the function is both ℓ -smooth and α -strongly convex:

Definition 1. A differentiable function $f(\cdot)$ is ℓ -smooth (or ℓ -gradient Lipschitz) if:

$$\forall \mathbf{x}_1, \mathbf{x}_2, \quad \|\nabla f(\mathbf{x}_1) - \nabla f(\mathbf{x}_2)\| \leq \ell \|\mathbf{x}_1 - \mathbf{x}_2\|.$$

Definition 2. A twice-differentiable function $f(\cdot)$ is α -strongly convex if $\forall \mathbf{x}, \lambda_{\min}(\nabla^2 f(\mathbf{x})) \geq \alpha$

Such smoothness guarantees imply that the gradient can not change too rapidly, and strong convexity ensures that there is a unique stationary point (and hence a global minimum). Standard analysis using these two properties shows that gradient descent converges linearly to a global optimum \mathbf{x}^* (see e.g. [Bubeck et al., 2015]).

Theorem 1. Assume $f(\cdot)$ is ℓ -smooth and α -strongly convex. For any $\epsilon > 0$, if we run gradient descent with step size $\eta = \frac{1}{\ell}$, iterate \mathbf{x}_t will be ϵ -close to \mathbf{x}^* in iterations:

$$\frac{2\ell}{\alpha} \log \frac{\|\mathbf{x}_0 - \mathbf{x}^*\|}{\epsilon}$$

In a more general setting, we no longer have convexity, let alone strong convexity. Though global optima are difficult to achieve in such a setting, it is possible to analyze convergence to first-order stationary points.

Definition 3. For a differentiable function $f(\cdot)$, we say that \mathbf{x} is a **first-order stationary point** if $\|\nabla f(\mathbf{x})\| = 0$; we also say \mathbf{x} is an ϵ -**first-order stationary point** if $\|\nabla f(\mathbf{x})\| \leq \epsilon$.

Under an ℓ -smoothness assumption, it is well known that by choosing the step size $\eta = \frac{1}{\ell}$, gradient descent converges to first-order stationary points.

Theorem 2 ([Nesterov, 1998]). Assume that the function $f(\cdot)$ is ℓ -smooth. Then, for any $\epsilon > 0$, if we run gradient descent with step size $\eta = \frac{1}{\ell}$ and termination condition $\|\nabla f(\mathbf{x})\| \leq \epsilon$, the output will be ϵ -first-order stationary point, and the algorithm will terminate within the following number of iterations:

$$\frac{\ell(f(\mathbf{x}_0) - f^*)}{\epsilon^2}.$$

Note that the iteration complexity does not depend explicitly on intrinsic dimension; in the literature this is referred to as “dimension-free optimization.”

A first-order stationary point can be either a local minimum or a saddle point or a local maximum. For minimization problems, saddle points and local maxima are undesirable, and we abuse nomenclature to call both of them “saddle points” in this paper. The formal definition is as follows:

Definition 4. For a differentiable function $f(\cdot)$, we say that \mathbf{x} is a **local minimum** if \mathbf{x} is a first-order stationary point, and there exists $\epsilon > 0$ so that for any \mathbf{y} in the ϵ -neighborhood of \mathbf{x} , we have $f(\mathbf{x}) \leq f(\mathbf{y})$; we also say \mathbf{x} is a **saddle point** if \mathbf{x} is a first-order stationary point but not a local minimum. For a twice-differentiable function $f(\cdot)$, we further say a saddle point \mathbf{x} is **strict** (or **non-degenerate**) if $\lambda_{\min}(\nabla^2 f(\mathbf{x})) < 0$.

For a twice-differentiable function $f(\cdot)$, we know a saddle point \mathbf{x} must satisfy $\lambda_{\min}(\nabla^2 f(\mathbf{x})) \leq 0$. Intuitively, for saddle point \mathbf{x} to be strict, we simply rule out the undetermined case $\lambda_{\min}(\nabla^2 f(\mathbf{x})) = 0$, where Hessian information alone is not enough to check whether \mathbf{x} is a local minimum or saddle point. In most non-convex problems, saddle points are undesirable.

To escape from saddle points and find local minima in a general setting, we move both the assumptions and guarantees in Theorem 2 one order higher. In particular, we require the Hessian to be Lipschitz:

Definition 5. A twice-differentiable function $f(\cdot)$ is ρ -**Hessian Lipschitz** if:

$$\forall \mathbf{x}_1, \mathbf{x}_2, \|\nabla^2 f(\mathbf{x}_1) - \nabla^2 f(\mathbf{x}_2)\| \leq \rho \|\mathbf{x}_1 - \mathbf{x}_2\|.$$

That is, Hessian can not change dramatically in terms of spectral norm. We also generalize the definition of first-order stationary point to higher order:

Algorithm 2 Perturbed Gradient Descent: PGD($\mathbf{x}_0, \ell, \rho, \epsilon, c, \delta, \Delta_f$)

$\chi \leftarrow 3 \max\{\log(\frac{d\ell\Delta_f}{c\epsilon^2\delta}), 4\}$, $\eta \leftarrow \frac{c}{\ell}$, $r \leftarrow \frac{\sqrt{c}}{\chi^2} \cdot \frac{\epsilon}{\ell}$, $g_{\text{thres}} \leftarrow \frac{\sqrt{c}}{\chi^2} \cdot \epsilon$, $f_{\text{thres}} \leftarrow \frac{c}{\chi^3} \cdot \sqrt{\frac{\epsilon^3}{\rho}}$, $t_{\text{thres}} \leftarrow \frac{\chi}{c^2} \cdot \frac{\ell}{\sqrt{\rho\epsilon}}$
 $t_{\text{noise}} \leftarrow -t_{\text{thres}} - 1$
for $t = 0, 1, \dots$ **do**
 if $\|\nabla f(\mathbf{x}_t)\| \leq g_{\text{thres}}$ **and** $t - t_{\text{noise}} > t_{\text{thres}}$ **then**
 $\tilde{\mathbf{x}}_t \leftarrow \mathbf{x}_t$, $t_{\text{noise}} \leftarrow t$
 $\mathbf{x}_t \leftarrow \tilde{\mathbf{x}}_t + \xi_t$, ξ_t uniformly $\sim \mathbb{B}_0(r)$
 if $t - t_{\text{noise}} = t_{\text{thres}}$ **and** $f(\mathbf{x}_t) - f(\tilde{\mathbf{x}}_{t_{\text{noise}}}) > -f_{\text{thres}}$ **then**
 return $\tilde{\mathbf{x}}_{t_{\text{noise}}}$
 $\mathbf{x}_{t+1} \leftarrow \mathbf{x}_t - \eta \nabla f(\mathbf{x}_t)$

Definition 6. For a ρ -Hessian Lipschitz function $f(\cdot)$, we say that \mathbf{x} is a **second-order stationary point** if $\|\nabla f(\mathbf{x})\| = 0$ and $\lambda_{\min}(\nabla^2 f(\mathbf{x})) \geq 0$; we also say \mathbf{x} is **ϵ -second-order stationary point** if:

$$\|\nabla f(\mathbf{x})\| \leq \epsilon, \quad \text{and} \quad \lambda_{\min}(\nabla^2 f(\mathbf{x})) \geq -\sqrt{\rho\epsilon}$$

Second-order stationary points are very important in non-convex optimization because when all saddle points are strict, all second-order stationary points are exactly local minima.

Note that the literature sometime defines ϵ -second-order stationary point by two independent error terms; i.e., letting $\|\nabla f(\mathbf{x})\| \leq \epsilon_g$ and $\lambda_{\min}(\nabla^2 f(\mathbf{x})) \geq -\epsilon_H$. We instead follow the convention of Nesterov and Polyak [2006] by choosing $\epsilon_H = \sqrt{\rho\epsilon_g}$ to reflect the natural relations between the gradient and the Hessian. This definition of ϵ -second-order stationary point can also differ by reparametrization (and scaling), e.g. Nesterov and Polyak [2006] use $\epsilon' = \sqrt{\epsilon/\rho}$. We choose our parametrization so that the first requirement of ϵ -second-order stationary point coincides with the requirement of ϵ -first-order stationary point, for a fair comparison of our result with Theorem 2.

3 Main Result

In this section we show that it possible to modify gradient descent in a simple way so that the resulting algorithm will provably converge quickly to a second-order stationary point.

The algorithm that we analyze is a perturbed form of gradient descent (see Algorithm 2). The algorithm is based on gradient descent with step size η . When the norm of the current gradient is small ($\leq g_{\text{thres}}$) (which indicates that the current iterate $\tilde{\mathbf{x}}_t$ is potentially near a saddle point), the algorithm adds a small random perturbation to the gradient. The perturbation is added at most only once every t_{thres} iterations.

To simplify the analysis we choose the perturbation ξ_t to be uniformly sampled from a d -dimensional ball¹. The use of the threshold t_{thres} ensures that the dynamics are mostly those of gradient descent. If the function value does not decrease enough (by f_{thres}) after t_{thres} iterations, the algorithm outputs $\tilde{\mathbf{x}}_{t_{\text{noise}}}$. The analysis in this section shows that under this protocol, the output $\tilde{\mathbf{x}}_{t_{\text{noise}}}$ is necessarily “close” to a second-order stationary point.

We first state the assumptions that we require.

¹Note that uniform sampling from a d -dimensional ball can be done efficiently by sampling $U^{\frac{1}{d}} \times \frac{\mathbf{Y}}{\|\mathbf{Y}\|}$ where $U \sim \text{Uniform}([0, 1])$ and $\mathbf{Y} \sim \mathcal{N}(0, \mathbf{I}_d)$ [Harman and Lacko, 2010].

Assumption A1. Function $f(\cdot)$ is both ℓ -smooth and ρ -Hessian Lipschitz.

The Hessian Lipschitz condition ensures that the function is well-behaved near a saddle point, and the small perturbation we add will suffice to allow the subsequent gradient updates to escape from the saddle point. More formally, we have:

Theorem 3. *Assume that $f(\cdot)$ satisfies A1. Then there exists an absolute constant c_{\max} such that, for any $\delta > 0, \epsilon \leq \frac{\ell^2}{\rho}, \Delta_f \geq f(\mathbf{x}_0) - f^*$, and constant $c \leq c_{\max}$, $\text{PGD}(\mathbf{x}_0, \ell, \rho, \epsilon, c, \delta, \Delta_f)$ will output an ϵ -second-order stationary point, with probability $1 - \delta$, and terminate in the following number of iterations:*

$$O\left(\frac{\ell(f(\mathbf{x}_0) - f^*)}{\epsilon^2} \log^4\left(\frac{d\ell\Delta_f}{\epsilon^2\delta}\right)\right).$$

Strikingly, Theorem 3 shows that perturbed gradient descent finds a second-order stationary point in almost the same amount of time that gradient descent takes to find first-order stationary point. The step size η is chosen as $O(1/\ell)$ which is in accord with classical analyses of convergence to first-order stationary points. Though we state the theorem with a certain choice of parameters for simplicity of presentation, our result holds even if we vary the parameters up to constant factors.

Without loss of generality, we can focus on the case $\epsilon \leq \ell^2/\rho$, as in Theorem 3. This is because in the case $\epsilon > \ell^2/\rho$, standard gradient descent without perturbation—Theorem 2—easily solves the problem (since by A1, we always have $\lambda_{\min}(\nabla^2 f(\mathbf{x})) \geq -\ell \geq -\sqrt{\rho\epsilon}$, which means that all ϵ -second-order stationary points are ϵ -first order stationary points).

We believe that the dependence on at least one $\log d$ factor in the iteration complexity is unavoidable in the non-convex setting, as our result can be directly applied to the principal component analysis problem, for which the best known runtimes (for the power method or Lanczos method) incur a $\log d$ factor. Establishing this formally is still an open question however.

To provide some intuition for Theorem 3, consider an iterate \mathbf{x}_t which is not yet an ϵ -second-order stationary point. By definition, either (1) the gradient $\nabla f(\mathbf{x}_t)$ is large, or (2) the Hessian $\nabla^2 f(\mathbf{x}_t)$ has a significant negative eigenvalue. Traditional analysis works in the first case. The crucial step in the proof of Theorem 3 involves handling the second case: when the gradient is small $\|\nabla f(\mathbf{x}_t)\| \leq g_{\text{thres}}$ and the Hessian has a significant negative eigenvalue $\lambda_{\min}(\nabla^2 f(\tilde{\mathbf{x}}_t)) \leq -\sqrt{\rho\epsilon}$, then adding a perturbation, followed by standard gradient descent for t_{thres} steps, decreases the function value by at least f_{thres} , with high probability. The proof of this fact relies on a novel characterization of geometry around saddle points (see Section 5)

If we are able to make stronger assumptions on the objective function we are able to strengthen our main result. This further analysis is presented in the next section.

3.1 Functions with Strict Saddle Property

In many real applications, objective functions further admit the property that all saddle points are strict [Ge et al., 2015, Sun et al., 2016a,b, Bhojanapalli et al., 2016, Ge et al., 2016]. In this case, all second-order stationary points are local minima and hence convergence to second-order stationary points (Theorem 3) is equivalent to convergence to local minima.

To state this result formally, we introduce a robust version of the strict saddle property [cf. Ge et al., 2015]:

Assumption A2. Function $f(\cdot)$ is (θ, γ, ζ) -strict saddle. That is, for any \mathbf{x} , at least one of following holds:

- $\|\nabla f(\mathbf{x})\| \geq \theta$.
- $\lambda_{\min}(\nabla^2 f(\mathbf{x})) \leq -\gamma$.
- \mathbf{x} is ζ -close to \mathcal{X}^* — the set of local minima.

Intuitively, the strict saddle assumption states that the \mathbb{R}^d space can be divided into three regions: 1) a region where the gradient is large; 2) a region where the Hessian has a significant negative eigenvalue (around saddle point); and 3) the region close to a local minimum. With this assumption, we immediately have the following corollary:

Corollary 4. *Let $f(\cdot)$ satisfy A1 and A2. Then, there exists an absolute constant c_{\max} such that, for any $\delta > 0$, $\Delta_f \geq f(\mathbf{x}_0) - f^*$, constant $c \leq c_{\max}$, and letting $\tilde{\epsilon} = \min(\theta, \gamma^2/\rho)$, $\text{PGD}(\mathbf{x}_0, \ell, \rho, \tilde{\epsilon}, c, \delta, \Delta_f)$ will output a point ζ -close to \mathcal{X}^* , with probability $1 - \delta$, and terminate in the following number of iterations:*

$$O\left(\frac{\ell(f(\mathbf{x}_0) - f^*)}{\tilde{\epsilon}^2} \log^4\left(\frac{d\ell\Delta_f}{\tilde{\epsilon}^2\delta}\right)\right).$$

Corollary 4 shows that by substituting ϵ in Theorem 3 using $\tilde{\epsilon} = \min(\theta, \gamma^2/\rho)$, the output of perturbed gradient descent will be in the ζ -neighborhood of some local minimum.

Note although Corollary 4 only explicitly asserts that the output will lie within some fixed radius ζ from a local minimum. In many real applications, we can further write ζ as a function $\zeta(\cdot)$ of gradient threshold θ , so that when θ decreases, $\zeta(\theta)$ decreases linearly or polynomially depending on θ . Meanwhile, parameter γ is always nondecreasing when θ decreases due to the nature of this strict saddle definition. Therefore, in these cases, the above corollary further gives a convergence rate to a local minimum.

3.2 Functions with Strong Local Structure

The convergence rate in Theorem 3 is polynomial in ϵ , which is similar to that of Theorem 2, but is worse than the rate of Theorem 1 because of the lack of strong convexity. Although global strong convexity does not hold in the non-convex setting that is our focus, in many machine learning problems the objective function may have a favorable local structure in the neighborhood of local minima [Ge et al., 2015, Sun et al., 2016a,b, Sun and Luo, 2016]. Exploiting this property can lead to much faster convergence (linear convergence) to local minima. One such property that ensures such convergence is a local form of smoothness and strong convexity:

Assumption A3.a. In a ζ -neighborhood of the set of local minima \mathcal{X}^* , the function $f(\cdot)$ is α -strongly convex, and β -smooth.

Here we use different letter β to denote the local smoothness parameter (in contrast to the global smoothness parameter ℓ). Note that we always have $\beta \leq \ell$. However, often even local α -strong convexity does not hold. We thus introduce the following relaxation:

Assumption A3.b. In a ζ -neighborhood of the set of local minima \mathcal{X}^* , the function $f(\cdot)$ satisfies a (α, β) -regularity condition if for any \mathbf{x} in this neighborhood:

$$\langle \nabla f(\mathbf{x}), \mathbf{x} - \mathcal{P}_{\mathcal{X}^*}(\mathbf{x}) \rangle \geq \frac{\alpha}{2} \|\mathbf{x} - \mathcal{P}_{\mathcal{X}^*}(\mathbf{x})\|^2 + \frac{1}{2\beta} \|\nabla f(\mathbf{x})\|^2. \quad (1)$$

Algorithm 3 Perturbed Gradient Descent with Local Improvement: PGDli($\mathbf{x}_0, \ell, \rho, \epsilon, c, \delta, \Delta_f, \beta$)

```

 $\mathbf{x}_0 \leftarrow \text{PGD}(\mathbf{x}_0, \ell, \rho, \epsilon, c, \delta, \Delta_f)$ 
for  $t = 0, 1, \dots$  do
     $\mathbf{x}_{t+1} \leftarrow \mathbf{x}_t - \frac{1}{\beta} \nabla f(\mathbf{x}_t)$ 

```

Here $\mathcal{P}_{\mathcal{X}^*}(\cdot)$ is the projection on to the set \mathcal{X}^* . Note (α, β) -regularity condition is more general and is directly implied by standard β -smooth and α -strongly convex conditions. This regularity condition commonly appears in low-rank problems such as matrix sensing and matrix completion, and has been used in Bhojanapalli et al. [2016], Zheng and Lafferty [2016], where local minima form a connected set, and where the Hessian is strictly positive only with respect to directions pointing outside the set of local minima.

Gradient descent naturally exploits local structure very well. In Algorithm 3, we first run Algorithm 2 to output a point within the neighborhood of a local minimum, and then perform standard gradient descent with step size $\frac{1}{\beta}$. We can then prove the following theorem:

Theorem 5. *Let $f(\cdot)$ satisfy A1, A2, and A3.a (or A3.b). Then there exists an absolute constant c_{\max} such that, for any $\delta > 0, \epsilon > 0, \Delta_f \geq f(\mathbf{x}_0) - f^*$, constant $c \leq c_{\max}$, and letting $\tilde{\epsilon} = \min(\theta, \gamma^2/\rho)$, $\text{PGDli}(\mathbf{x}_0, \ell, \rho, \tilde{\epsilon}, c, \delta, \Delta_f, \beta)$ will output a point that is ϵ -close to \mathcal{X}^* , with probability $1 - \delta$, in the following number of iterations:*

$$O\left(\frac{\ell(f(\mathbf{x}_0) - f^*)}{\tilde{\epsilon}^2} \log^4\left(\frac{d\ell\Delta_f}{\tilde{\epsilon}^2\delta}\right) + \frac{\beta}{\alpha} \log \frac{\zeta}{\epsilon}\right).$$

Theorem 5 says that if strong local structure is present, the convergence rate can be boosted to linear convergence ($\log \frac{1}{\epsilon}$). In this theorem we see that sequence of iterations can be decomposed into two phases. In the first phase, perturbed gradient descent finds a ζ -neighborhood by Corollary 4. In the second phase, standard gradient descent takes us from ζ to ϵ -close to a local minimum. Standard gradient descent and Assumption A3.a (or A3.b) make sure that the iterate never steps out of a ζ -neighborhood in this second phase, giving a result similar to Theorem 1 with linear convergence.

Finally, we note our choice of local conditions (Assumption A3.a and A3.b) are not special. The interested reader can refer to Karimi et al. [2016] for other relaxed and alternative notions of convexity, which can also be potentially combined with Assumptions A1 and A2 to yield convergence results of a similar flavor as that of Theorem 5.

4 Example — Matrix Factorization

As a simple example to illustrate how to apply our general theorems to specific non-convex optimization problems, we consider a symmetric low-rank matrix factorization problem, based on the following objective function:

$$\min_{\mathbf{U} \in \mathbb{R}^{d \times r}} f(\mathbf{U}) = \frac{1}{2} \|\mathbf{U}\mathbf{U}^\top - \mathbf{M}^*\|_{\text{F}}^2, \quad (2)$$

where $\mathbf{M}^* \in \mathbb{R}^{d \times d}$. For simplicity, we assume $\text{rank}(\mathbf{M}^*) = r$, and denote $\sigma_1^* := \sigma_1(\mathbf{M}^*)$, $\sigma_r^* := \sigma_r(\mathbf{M}^*)$. Clearly, in this case the global minimum of function value is zero, which is achieved at $\mathbf{V}^* = \mathbf{T}\mathbf{D}^{1/2}$ where $\mathbf{T}\mathbf{D}\mathbf{T}^\top$ is the SVD of the symmetric real matrix \mathbf{M}^* .

The following two lemmas show that the objective function in Eq. (2) satisfies the geometric assumptions A1, A2, and A3.b. Moreover, all local minima are global minima.

Lemma 6. *For any $\Gamma \geq \sigma_1^*$, the function $f(\mathbf{U})$ defined in Eq. (2) is 8Γ -smooth and $12\Gamma^{1/2}$ -Hessian Lipschitz, inside the region $\{\mathbf{U} \mid \|\mathbf{U}\|^2 < \Gamma\}$.*

Lemma 7. *For function $f(\mathbf{U})$ defined in Eq. (2), all local minima are global minima. The set of global minima is $\mathcal{X}^* = \{\mathbf{V}^* \mathbf{R} \mathbf{R}^\top = \mathbf{R}^\top \mathbf{R} = \mathbf{I}\}$. Furthermore, $f(\mathbf{U})$ satisfies:*

1. $(\frac{1}{24}(\sigma_r^*)^{3/2}, \frac{1}{3}\sigma_r^*, \frac{1}{3}(\sigma_r^*)^{1/2})$ -strict saddle property.
2. $(\frac{2}{3}\sigma_r^*, 10\sigma_1^*)$ -regularity condition in $\frac{1}{3}(\sigma_r^*)^{1/2}$ neighborhood of \mathcal{X}^* .

One caveat is that since the objective function is actually a fourth-order polynomial with respect to \mathbf{U} , the smoothness and Hessian Lipschitz parameters from Lemma 6 naturally depend on $\|\mathbf{U}\|$. Fortunately, we can further show that gradient descent (even with perturbation) does not increase $\|\mathbf{U}\|$ beyond $O(\max\{\|\mathbf{U}_0\|, (\sigma_1^*)^{1/2}\})$. Then, applying Theorem 5 gives:

Theorem 8. *There exists an absolute constant c_{\max} such that the following holds. For the objective function in Eq. (2), for any $\delta > 0$ and constant $c \leq c_{\max}$, and for $\Gamma^{1/2} := 2 \max\{\|\mathbf{U}_0\|, 3(\sigma_1^*)^{1/2}\}$, the output of $\text{PGDli}(\mathbf{U}_0, 8\Gamma, 12\Gamma^{1/2}, \frac{(\sigma_r^*)^2}{108\Gamma^{1/2}}, c, \delta, \frac{r\Gamma^2}{2}, 10\sigma_1^*)$, will be ϵ -close to the global minimum set \mathcal{X}^* , with probability $1 - \delta$, after the following number of iterations:*

$$O\left(r\left(\frac{\Gamma}{\sigma_r^*}\right)^4 \log^4\left(\frac{d\Gamma}{\delta\sigma_r^*}\right) + \frac{\sigma_1^*}{\sigma_r^*} \log \frac{\sigma_r^*}{\epsilon}\right).$$

Theorem 8 establishes global convergence of perturbed gradient descent from an arbitrary initial point \mathbf{U}_0 , including exact saddle points. Suppose we initialize at $\mathbf{U}_0 = 0$, then our iteration complexity becomes:

$$O\left(r(\kappa^*)^4 \log^4(d\kappa^*/\delta) + \kappa^* \log(\sigma_r^*/\epsilon)\right),$$

where $\kappa^* = \sigma_1^*/\sigma_r^*$ is the condition number of the matrix \mathbf{M}^* . We see that in the second phase, when convergence occurs inside the local region, we require $O(\kappa^* \log(\sigma_r^*/\epsilon))$ iterations which is the standard local linear rate for gradient descent. In the first phase, to find a neighborhood of the solution, our method requires a number of iterations scaling as $\tilde{O}(r(\kappa^*)^4)$. We suspect that this strong dependence on condition number arises from our generic assumption that the Hessian Lipschitz is uniformly upper bounded; it may well be the case that this dependence can be reduced in the special case of matrix factorization via a finer analysis of the geometric structure of the problem.

5 Proof Sketch for Theorem 3

In this section we will present the key ideas underlying the main result of this paper (Theorem 3). We will first argue the correctness of Theorem 3 given two important intermediate lemmas. Then we turn to the main lemma, which establishes that gradient descent can escape from saddle points quickly. We present full proofs of all these results in Appendix A. Throughout this section, we use $\eta, r, g_{\text{thres}}, f_{\text{thres}}$ and t_{thres} as defined in Algorithm 2.

5.1 Exploiting Large Gradient or Negative Curvature

Recall that an ϵ -second-order stationary point is a point with a small gradient, and where the Hessian does not have a significant negative eigenvalue. Suppose we are currently at an iterate \mathbf{x}_t that is not an ϵ -second-order stationary point; i.e., it does not satisfy the above properties. There are two possibilities:

1. Gradient is large: $\|\nabla f(\mathbf{x}_t)\| \geq g_{\text{thres}}$, or
2. Around saddle point: $\|\nabla f(\mathbf{x}_t)\| \leq g_{\text{thres}}$ and $\lambda_{\min}(\nabla^2 f(\mathbf{x}_t)) \leq -\sqrt{\rho\epsilon}$.

The following two lemmas address these two cases respectively. They guarantee that perturbed gradient descent will decrease the function value in both scenarios.

Lemma 9 (Gradient). *Assume that $f(\cdot)$ satisfies A1. Then for gradient descent with stepsize $\eta < \frac{1}{L}$, we have $f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}_t) - \frac{\eta}{2}\|\nabla f(\mathbf{x}_t)\|^2$.*

Lemma 10 (Saddle). *(informal) Assume that $f(\cdot)$ satisfies A1, If \mathbf{x}_t satisfies $\|\nabla f(\mathbf{x}_t)\| \leq g_{\text{thres}}$ and $\lambda_{\min}(\nabla^2 f(\mathbf{x}_t)) \leq -\sqrt{\rho\epsilon}$, then adding one perturbation step followed by t_{thres} steps of gradient descent, we have $f(\mathbf{x}_{t+t_{\text{thres}}}) - f(\mathbf{x}_t) \leq -f_{\text{thres}}$ with high probability.*

We see that Algorithm 2 is designed so that Lemma 10 can be directly applied. According to these two lemmas, perturbed gradient descent will decrease the function value either in the case of a large gradient, or around strict saddle points. Computing the average decrease per step in function value yields the total iteration complexity. Since Algorithm 2 only terminate when the function value decreases too slowly, this guarantees that the output must be ϵ -second-order stationary point (see Appendix A for formal proofs).

5.2 Main Lemma: Escaping from Saddle Points Quickly

The proof of Lemma 9 is straightforward and follows from traditional analysis. The key technical contribution of this paper is the proof of Lemma 10, which gives a new characterization of the geometry around saddle points.

Consider a point $\tilde{\mathbf{x}}$ that satisfies the the preconditions of Lemma 10 ($\|\nabla f(\tilde{\mathbf{x}})\| \leq g_{\text{thres}}$ and $\lambda_{\min}(\nabla^2 f(\tilde{\mathbf{x}})) \leq -\sqrt{\rho\epsilon}$). After adding the perturbation ($\mathbf{x}_0 = \tilde{\mathbf{x}} + \xi$), we can view \mathbf{x}_0 as coming from a uniform distribution over $\mathbb{B}_{\tilde{\mathbf{x}}}(r)$, which we call the **perturbation ball**. We can divide this perturbation ball $\mathbb{B}_{\tilde{\mathbf{x}}}(r)$ into two disjoint regions: (1) an **escaping region** $\mathcal{X}_{\text{escape}}$ which consists of all the points $\mathbf{x} \in \mathbb{B}_{\tilde{\mathbf{x}}}(r)$ whose function value decreases by at least f_{thres} after t_{thres} steps; (2) a **stuck region** $\mathcal{X}_{\text{stuck}} = \mathbb{B}_{\tilde{\mathbf{x}}}(r) - \mathcal{X}_{\text{escape}}$. Our general proof strategy is to show that $\mathcal{X}_{\text{stuck}}$ consists of a very small proportion of the volume of perturbation ball. After adding a perturbation to $\tilde{\mathbf{x}}$, point \mathbf{x}_0 has a very small chance of falling in $\mathcal{X}_{\text{stuck}}$, and hence will escape from the saddle point efficiently.

Let us consider the nature of $\mathcal{X}_{\text{stuck}}$. For simplicity, let us imagine that $\tilde{\mathbf{x}}$ is an exact saddle point whose Hessian has only one negative eigenvalue, and $d-1$ positive eigenvalues. Let us denote the minimum eigenvalue direction as \mathbf{e}_1 . In this case, if the Hessian remains constant (and we have a quadratic function), the stuck region $\mathcal{X}_{\text{stuck}}$ consists of points \mathbf{x} such that $\mathbf{x} - \tilde{\mathbf{x}}$ has a small \mathbf{e}_1 component. This is a straight band in two dimensions and a flat disk in high dimensions. However, when the Hessian is not constant, the shape of the stuck region is distorted. In two dimensions, it




Figure 1: Perturbation ball in 3D and “thin pancake” shape stuck region




Figure 2: Perturbation ball in 2D and “narrow band” stuck region under gradient flow

forms a “narrow band” as plotted in Figure 2 on top of the gradient flow. In three dimensions, it forms a “thin pancake” as shown in Figure 1.

The major challenge here is to bound the volume of this high-dimensional non-flat “pancake” shaped region $\mathcal{X}_{\text{stuck}}$. A crude approximation of this “pancake” by a flat “disk” loses polynomial factors in the dimensionality, which gives a suboptimal rate. Our proof relies on the following crucial observation: Although we do not know the explicit form of the stuck region, we know it must be very “thin,” therefore it cannot have a large volume. The informal statement of the lemma is as follows:

Lemma 11. *(informal) Suppose $\tilde{\mathbf{x}}$ satisfies the precondition of Lemma 10, and let \mathbf{e}_1 be the smallest eigendirection of $\nabla^2 f(\tilde{\mathbf{x}})$. For any $\delta \in (0, 1/3]$ and any two points $\mathbf{w}, \mathbf{u} \in \mathbb{B}_{\tilde{\mathbf{x}}}(r)$, if $\mathbf{w} - \mathbf{u} = \mu r \mathbf{e}_1$ and $\mu \geq \delta/(2\sqrt{d})$, then at least one of \mathbf{w}, \mathbf{u} is not in the stuck region $\mathcal{X}_{\text{stuck}}$.*

Using this lemma it is not hard to bound the volume of the stuck region: we can draw a straight line along the \mathbf{e}_1 direction which intersects the perturbation ball (shown as purple line segment in Figure 2). For any two points on this line segment that are at least $\delta r/(2\sqrt{d})$ away from each other (shown as red points \mathbf{w}, \mathbf{u} in Figure 2), by Lemma 11, we know at least one of them must not be in $\mathcal{X}_{\text{stuck}}$. This implies if there is one point $\tilde{\mathbf{u}} \in \mathcal{X}_{\text{stuck}}$ on this line segment, then $\mathcal{X}_{\text{stuck}}$ on this line can be at most an interval of length $\delta r/\sqrt{d}$ around $\tilde{\mathbf{u}}$. This establishes the “thickness” of $\mathcal{X}_{\text{stuck}}$ in the \mathbf{e}_1 direction, which is turned into an upper bound on the volume of the stuck region $\mathcal{X}_{\text{stuck}}$ by standard calculus.

6 Conclusion

This paper presents the first (nearly) dimension-free result for gradient descent in a general non-convex setting. We present a general convergence result and show how it can be further strengthened when combined with further structure such as strict saddle conditions and/or local regularity/convexity.

There are still many related open problems. First, in the presence of constraints, it is worthwhile to study whether gradient descent still admits similar sharp convergence results. Another important question is whether similar techniques can be applied to accelerated gradient descent. We hope that this result could serve as a first step towards a more general theory with strong, almost dimension free guarantees for non-convex optimization.

References

- Naman Agarwal, Zeyuan Allen-Zhu, Brian Bullins, Elad Hazan, and Tengyu Ma. Finding approximate local minima for nonconvex optimization in linear time. *arXiv preprint arXiv:1611.01146*, 2016.
- Srinadh Bhojanapalli, Behnam Neyshabur, and Nathan Srebro. Global optimality of local search for low rank matrix recovery. *arXiv preprint arXiv:1605.07221*, 2016.
- Sébastien Bubeck et al. Convex optimization: Algorithms and complexity. *Foundations and Trends® in Machine Learning*, 8(3-4):231–357, 2015.
- Emmanuel J Candes, Xiaodong Li, and Mahdi Soltanolkotabi. Phase retrieval via wirtinger flow: Theory and algorithms. *IEEE Transactions on Information Theory*, 61(4):1985–2007, 2015.
- Yair Carmon and John C Duchi. Gradient descent efficiently finds the cubic-regularized non-convex newton step. *arXiv preprint arXiv:1612.00547*, 2016.
- Yair Carmon, John C Duchi, Oliver Hinder, and Aaron Sidford. Accelerated methods for non-convex optimization. *arXiv preprint arXiv:1611.00756*, 2016.
- Anna Choromanska, Mikael Henaff, Michael Mathieu, Gérard Ben Arous, and Yann LeCun. The loss surface of multilayer networks. *arXiv:1412.0233*, 2014.
- Frank E Curtis, Daniel P Robinson, and Mohammadreza Samadi. A trust region algorithm with a worst-case iteration complexity of $\mathcal{O}(\epsilon^{-3/2})$ for nonconvex optimization. *Mathematical Programming*, pages 1–32, 2014.
- Yann N Dauphin, Razvan Pascanu, Caglar Gulcehre, Kyunghyun Cho, Surya Ganguli, and Yoshua Bengio. Identifying and attacking the saddle point problem in high-dimensional non-convex optimization. In *Advances in Neural Information Processing Systems*, pages 2933–2941, 2014.
- Rong Ge, Furong Huang, Chi Jin, and Yang Yuan. Escaping from saddle points—online stochastic gradient for tensor decomposition. In *COLT*, 2015.
- Rong Ge, Jason D Lee, and Tengyu Ma. Matrix completion has no spurious local minimum. In *Advances in Neural Information Processing Systems*, pages 2973–2981, 2016.
- Radoslav Harman and Vladimír Lacko. On decomposition algorithms for uniform sampling from n-spheres and n-balls. *Journal of Multivariate Analysis*, 101(10):2297–2304, 2010.
- Prateek Jain, Chi Jin, Sham M Kakade, and Praneeth Netrapalli. Computing matrix squareroot via non convex local search. *arXiv preprint arXiv:1507.05854*, 2015.

- Hamed Karimi, Julie Nutini, and Mark Schmidt. Linear convergence of gradient and proximal-gradient methods under the Polyak-Lojasiewicz condition. In *Joint European Conference on Machine Learning and Knowledge Discovery in Databases*, pages 795–811. Springer, 2016.
- Kenji Kawaguchi. Deep learning without poor local minima. In *Advances In Neural Information Processing Systems*, pages 586–594, 2016.
- Jason D Lee, Max Simchowitz, Michael I Jordan, and Benjamin Recht. Gradient descent only converges to minimizers. In *Conference on Learning Theory*, pages 1246–1257, 2016.
- Kfir Y Levy. The power of normalization: Faster evasion of saddle points. *arXiv preprint arXiv:1611.04831*, 2016.
- Yu Nesterov. Introductory lectures on convex programming volume i: Basic course. *Lecture notes*, 1998.
- Yurii Nesterov and Boris T Polyak. Cubic regularization of newton method and its global performance. *Mathematical Programming*, 108(1):177–205, 2006.
- Praneeth Netrapalli, Prateek Jain, and Sujay Sanghavi. Phase retrieval using alternating minimization. In *Advances in Neural Information Processing Systems*, pages 2796–2804, 2013.
- Dohyung Park, Anastasios Kyrillidis, Constantine Caramanis, and Sujay Sanghavi. Non-square matrix sensing without spurious local minima via the burer-monteiro approach. *arXiv preprint arXiv:1609.03240*, 2016.
- Boris T Polyak. Gradient methods for the minimisation of functionals. *USSR Computational Mathematics and Mathematical Physics*, 3(4):864–878, 1963.
- David E Rumelhart, Geoffrey E Hinton, and Ronald J Williams. Learning representations by back-propagating errors. *Cognitive modeling*, 5, 1988.
- Ju Sun, Qing Qu, and John Wright. Complete dictionary recovery over the sphere i: Overview and the geometric picture. *IEEE Transactions on Information Theory*, 2016a.
- Ju Sun, Qing Qu, and John Wright. A geometric analysis of phase retrieval. In *Information Theory (ISIT), 2016 IEEE International Symposium on*, pages 2379–2383. IEEE, 2016b.
- Ruoyu Sun and Zhi-Quan Luo. Guaranteed matrix completion via non-convex factorization. *IEEE Transactions on Information Theory*, 62(11):6535–6579, 2016.
- Qinqing Zheng and John Lafferty. Convergence analysis for rectangular matrix completion using burer-monteiro factorization and gradient descent. *arXiv preprint arXiv:1605.07051*, 2016.

A Detailed Proof of Main Theorem

In this section, we give detailed proof for the main theorem. We will first state two key lemmas that show how the algorithm can make progress when the gradient is large or near a saddle point, and show how the main theorem follows from the two lemmas. Then we will focus on the novel technique in this paper: how to analyze gradient descent near saddle point.

A.1 General Framework

In order to prove the main theorem, we need to show that the algorithm will not be stuck at any point that either has a large gradient or is near a saddle point. This idea is similar to previous works (e.g.[Ge et al., 2015]). We first state a standard Lemma that shows if the current gradient is large, then we make progress in function value.

Lemma 12 (Lemma 9 restated). *Assume $f(\cdot)$ satisfies A1, then for gradient descent with stepsize $\eta < \frac{1}{\ell}$, we have:*

$$f(\mathbf{x}_{t+1}) \leq f(\mathbf{x}_t) - \frac{\eta}{2} \|\nabla f(\mathbf{x}_t)\|^2$$

Proof. By Assumption A1 and its property, we have:

$$\begin{aligned} f(\mathbf{x}_{t+1}) &\leq f(\mathbf{x}_t) + \nabla f(\mathbf{x}_t)^\top (\mathbf{x}_{t+1} - \mathbf{x}_t) + \frac{\ell}{2} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 \\ &= f(\mathbf{x}_t) - \eta \|\nabla f(\mathbf{x}_t)\|^2 + \frac{\eta^2 \ell}{2} \|\nabla f(\mathbf{x}_t)\|^2 \leq f(\mathbf{x}_t) - \frac{\eta}{2} \|\nabla f(\mathbf{x}_t)\|^2 \end{aligned}$$

□

The next lemma says that if we are “close to a saddle points”, i.e., we are at a point where the gradient is small, but the Hessian has a reasonably large negative eigenvalue. This is the main difficulty in the analysis. We show a perturbation followed by small number of standard gradient descent steps can also make the function value decrease with high probability.

Lemma 13 (Lemma 10 formal). *There exist absolute constant c_{\max} , for $f(\cdot)$ satisfies A1, and any $c \leq c_{\max}$, and $\chi \geq 1$. Let $\eta, r, g_{\text{thres}}, f_{\text{thres}}, t_{\text{thres}}$ calculated same way as in Algorithm 2. Then, if $\tilde{\mathbf{x}}_t$ satisfies:*

$$\|\nabla f(\tilde{\mathbf{x}}_t)\| \leq g_{\text{thres}} \quad \text{and} \quad \lambda_{\min}(\nabla^2 f(\tilde{\mathbf{x}}_t)) \leq -\sqrt{\rho\epsilon}$$

Let $\mathbf{x}_t = \tilde{\mathbf{x}}_t + \xi_t$ where ξ_t comes from the uniform distribution over $\mathbb{B}_0(r)$, and let \mathbf{x}_{t+i} be the iterates of gradient descent from \mathbf{x}_t with stepsize η , then with at least probability $1 - \frac{d\ell}{\sqrt{\rho\epsilon}} e^{-\chi}$, we have:

$$f(\mathbf{x}_{t+t_{\text{thres}}}) - f(\tilde{\mathbf{x}}_t) \leq -f_{\text{thres}}$$

The proof of this lemma is deferred to Section A.2. Using this Lemma, we can then prove the main Theorem.

Theorem 3. *There exist absolute constant c_{\max} such that: if $f(\cdot)$ satisfies A1, then for any $\delta > 0, \epsilon \leq \frac{\ell^2}{\rho}, \Delta_f \geq f(\mathbf{x}_0) - f^*$, and constant $c \leq c_{\max}$, with probability $1 - \delta$, the output of PGD($\mathbf{x}_0, \ell, \rho, \epsilon, c, \delta, \Delta_f$) will be ϵ -second order stationary point, and terminate in iterations:*

$$O\left(\frac{\ell(f(\mathbf{x}_0) - f^*)}{\epsilon^2} \log^4\left(\frac{d\ell\Delta_f}{\epsilon^2\delta}\right)\right)$$

Proof. Denote \tilde{c}_{\max} to be the absolute constant allowed in Theorem 13. In this theorem, we let $c_{\max} = \min\{\tilde{c}_{\max}, 1/2\}$, and choose any constant $c \leq c_{\max}$.

In this proof, we will actually achieve some point satisfying following condition:

$$\|\nabla f(\mathbf{x})\| \leq g_{\text{thres}} = \frac{\sqrt{c}}{\chi^2} \cdot \epsilon, \quad \lambda_{\min}(\nabla^2 f(\mathbf{x})) \geq -\sqrt{\rho\epsilon} \quad (3)$$

Since $c \leq 1$, $\chi \geq 1$, we have $\frac{\sqrt{c}}{\chi^2} \leq 1$, which implies any \mathbf{x} satisfy Eq.(3) is also a ϵ -second-order stationary point.

Starting from \mathbf{x}_0 , we know if \mathbf{x}_0 does not satisfy Eq.(3), there are only two possibilities:

1. $\|\nabla f(\mathbf{x}_0)\| > g_{\text{thres}}$: In this case, Algorithm 2 will not add perturbation. By Lemma 12:

$$f(\mathbf{x}_1) - f(\mathbf{x}_0) \leq -\frac{\eta}{2} \cdot g_{\text{thres}}^2 = -\frac{c^2}{2\chi^4} \cdot \frac{\epsilon^2}{\ell}$$

2. $\|\nabla f(\mathbf{x}_0)\| \leq g_{\text{thres}}$: In this case, Algorithm 2 will add a perturbation of radius r , and will perform gradient descent (without perturbations) for the next t_{thres} steps. Algorithm 2 will then check termination condition. If the condition is not met, we must have:

$$f(\mathbf{x}_{t_{\text{thres}}}) - f(\mathbf{x}_0) \leq -f_{\text{thres}} = -\frac{c}{\chi^3} \cdot \sqrt{\frac{\epsilon^3}{\rho}}$$

This means on average every step decreases the function value by

$$\frac{f(\mathbf{x}_{t_{\text{thres}}}) - f(\mathbf{x}_0)}{t_{\text{thres}}} \leq -\frac{c^3}{\chi^4} \cdot \frac{\epsilon^2}{\ell}$$

In case 1, we can repeat this argument for $t = 1$ and in case 2, we can repeat this argument for $t = t_{\text{thres}}$. Hence, we can conclude as long as algorithm 2 has not terminated yet, on average, every step decrease function value by at least $\frac{c^3}{\chi^4} \cdot \frac{\epsilon^2}{\ell}$. However, we clearly can not decrease function value by more than $f(\mathbf{x}_0) - f^*$, where f^* is the function value of global minima. This means algorithm 2 must terminate within the following number of iterations:

$$\frac{f(\mathbf{x}_0) - f^*}{\frac{c^3}{\chi^4} \cdot \frac{\epsilon^2}{\ell}} = \frac{\chi^4}{c^3} \cdot \frac{\ell(f(\mathbf{x}_0) - f^*)}{\epsilon^2} = O\left(\frac{\ell(f(\mathbf{x}_0) - f^*)}{\epsilon^2} \log^4\left(\frac{d\ell\Delta_f}{\epsilon^2\delta}\right)\right)$$

Finally, we would like to ensure when Algorithm 2 terminates, the point it finds is actually an ϵ -second-order stationary point. The algorithm can only terminate when the gradient is small, and the function value does not decrease after a perturbation and t_{thres} iterations. We shall show every time when we add perturbation to iterate $\tilde{\mathbf{x}}_t$, if $\lambda_{\min}(\nabla^2 f(\tilde{\mathbf{x}}_t)) < -\sqrt{\rho\epsilon}$, then we will have $f(\mathbf{x}_{t+t_{\text{thres}}}) - f(\tilde{\mathbf{x}}_t) \leq -f_{\text{thres}}$. Thus, whenever the current point is not an ϵ -second-order stationary point, the algorithm cannot terminate.

According to Algorithm 2, we immediately know $\|\nabla f(\tilde{\mathbf{x}}_t)\| \leq g_{\text{thres}}$ (otherwise we will not add perturbation at time t). By Lemma 13, we know this event happens with probability at least

$1 - \frac{d\ell}{\sqrt{\rho\epsilon}}e^{-\chi}$ each time. On the other hand, during one entire run of Algorithm 2, the number of times we add perturbations is at most:

$$\frac{1}{t_{\text{thres}}} \cdot \frac{\chi^4}{c^3} \cdot \frac{\ell(f(\mathbf{x}_0) - f^*)}{\epsilon^2} = \frac{\chi^3}{c} \frac{\sqrt{\rho\epsilon}(f(\mathbf{x}_0) - f^*)}{\epsilon^2}$$

By union bound, for all these perturbations, with high probability Lemma 13 is satisfied. As a result Algorithm 2 works correctly. The probability of that is at least

$$1 - \frac{d\ell}{\sqrt{\rho\epsilon}}e^{-\chi} \cdot \frac{\chi^3}{c} \frac{\sqrt{\rho\epsilon}(f(\mathbf{x}_0) - f^*)}{\epsilon^2} = 1 - \frac{\chi^3 e^{-\chi}}{c} \cdot \frac{d\ell(f(\mathbf{x}_0) - f^*)}{\epsilon^2}$$

Recall our choice of $\chi = 3 \max\{\log(\frac{d\ell\Delta_f}{c\epsilon^2\delta}), 4\}$. Since $\chi \geq 12$, we have $\chi^3 e^{-\chi} \leq e^{-\chi/3}$, this gives:

$$\frac{\chi^3 e^{-\chi}}{c} \cdot \frac{d\ell(f(\mathbf{x}_0) - f^*)}{\epsilon^2} \leq e^{-\chi/3} \frac{d\ell(f(\mathbf{x}_0) - f^*)}{c\epsilon^2} \leq \delta$$

which finishes the proof. \square

A.2 Main Lemma: Escaping from Saddle Points Quickly

Now we prove the main lemma (Lemma 13), which shows near a saddle point, a small perturbation followed by a small number of gradient descent steps will decrease the function value with high probability. This is the main step where we need new analysis, as the analysis previous works (e.g.[Ge et al., 2015]) do not work when the step size and perturbation do not depend polynomially in dimension d .

Intuitively, after adding a perturbation, the current point of the algorithm comes from a uniform distribution over a d -dimensional ball centered at $\tilde{\mathbf{x}}$, which we call **perturbation ball**. After a small number of gradient steps, some points in this ball (which we call the **escaping region**) will significantly decrease the function; other points (which we call the **stuck region**) does not see a significant decrease in function value. We hope to show that the escaping region constitutes at least $1 - \delta$ fraction of the volume of the perturbation ball.

However, we do not know the exact form of the function near the saddle point, so the escaping region does not have a clean analytic description. Explicitly computing its volume can be very difficult. Our proof rely on a crucial observation: although we do not know the shape of the stuck region, we know the “width” of it must be small, therefore it cannot have a large volume. We will formalize this intuition later in Lemma 15.

The proof of the main lemma requires carefully balancing between different quantities including function value, gradient, parameter space and number of iterations. For clarify, we define following scalar quantities, which serve as the “units” for function value, gradient, parameter space, and time (iterations). We will use these notations throughout the proof.

Let the condition number be the ratio of the smoothness parameter (largest eigenvalue of Hessian) and the negative eigenvalue γ : $\kappa = \ell/\gamma \geq 1$, we define the following units:

$$\begin{aligned} \mathcal{F} &:= \eta\ell \frac{\gamma^3}{\rho^2} \cdot \log^{-3}\left(\frac{d\kappa}{\delta}\right), & \mathcal{G} &:= \sqrt{\eta\ell} \frac{\gamma^2}{\rho} \cdot \log^{-2}\left(\frac{d\kappa}{\delta}\right) \\ \mathcal{S} &:= \sqrt{\eta\ell} \frac{\gamma}{\rho} \cdot \log^{-1}\left(\frac{d\kappa}{\delta}\right), & \mathcal{T} &:= \frac{\log(\frac{d\kappa}{\delta})}{\eta\gamma} \end{aligned}$$

Intuitively, if we plug in our choice of learning rate $\eta\ell = O(1)$ (which we will prove later) and hide the logarithmic dependences, we have $\mathcal{F} = \tilde{O}(\frac{\gamma^3}{\rho^2})$, $\mathcal{G} = \tilde{O}(\frac{\gamma^2}{\rho})$, $\mathcal{S} = \tilde{O}(\frac{\gamma}{\rho})$, which is the only way to correctly describe the units of function value, gradient, parameter space by just γ and ρ . Moreover, these units are closely related, in particular, we know $\sqrt{\frac{\mathcal{F} \cdot \log(\frac{d\kappa}{\delta})}{\gamma}} = \frac{\mathcal{G} \cdot \log(\frac{d\kappa}{\delta})}{\gamma} = \mathcal{S}$.

For simplicity of later proofs, we first restate Lemma 13 into a slightly more general form as follows. Lemma 13 is directly implied following lemma.

Lemma 14 (Lemma 13 restated). *There exists universal constant c_{\max} , for $f(\cdot)$ satisfies A1, for any $\delta \in (0, \frac{d\kappa}{e}]$, suppose we start with point $\tilde{\mathbf{x}}$ satisfying following conditions:*

$$\|\nabla f(\tilde{\mathbf{x}})\| \leq \mathcal{G} \quad \text{and} \quad \lambda_{\min}(\nabla^2 f(\tilde{\mathbf{x}})) \leq -\gamma$$

Let $\mathbf{x}_0 = \tilde{\mathbf{x}} + \xi$ where ξ come from the uniform distribution over ball with radius $\mathcal{S}/(\kappa \cdot \log(\frac{d\kappa}{\delta}))$, and let \mathbf{x}_t be the iterates of gradient descent from \mathbf{x}_0 . Then, when stepsize $\eta \leq c_{\max}/\ell$, with at least probability $1 - \delta$, we have following for any $T \geq \frac{1}{c_{\max}} \mathcal{T}$:

$$f(\mathbf{x}_T) - f(\tilde{\mathbf{x}}) \leq -\mathcal{F}$$

Lemma 14 is almost the same as Lemma 13. It is easy to verify that by substituting $\eta = \frac{e}{\ell}$, $\gamma = \sqrt{\rho e}$ and $\delta = \frac{d\ell}{\sqrt{\rho e}} e^{-\chi}$ into Lemma 14, we immediately obtain Lemma 13.

Now we will formalize the intuition that the “width” of stuck region is small.

Lemma 15 (Lemma 11 restated). *There exists a universal constant c_{\max} , for any $\delta \in (0, \frac{d\kappa}{e}]$, let $f(\cdot), \tilde{\mathbf{x}}$ satisfies the conditions in Lemma 14, and without loss of generality let \mathbf{e}_1 be the minimum eigenvector of $\nabla^2 f(\tilde{\mathbf{x}})$. Consider two gradient descent sequences $\{\mathbf{u}_t\}, \{\mathbf{w}_t\}$ with initial points $\mathbf{u}_0, \mathbf{w}_0$ satisfying: (denote radius $r = \mathcal{S}/(\kappa \cdot \log(\frac{d\kappa}{\delta}))$)*

$$\|\mathbf{u}_0 - \tilde{\mathbf{x}}\| \leq r, \quad \mathbf{w}_0 = \mathbf{u}_0 + \mu \cdot r \cdot \mathbf{e}_1, \quad \mu \in [\delta/(2\sqrt{d}), 1]$$

Then, for any stepsize $\eta \leq c_{\max}/\ell$, and any $T \geq \frac{1}{c_{\max}} \mathcal{T}$, we have:

$$\min\{f(\mathbf{u}_T) - f(\mathbf{u}_0), f(\mathbf{w}_T) - f(\mathbf{w}_0)\} \leq -2.5\mathcal{F}$$

Intuitively, lemma 15 claims for any two points $\mathbf{u}_0, \mathbf{w}_0$ inside the perturbation ball, if $\mathbf{u}_0 - \mathbf{w}_0$ lies in the direction of minimum eigenvector of $\nabla^2 f(\tilde{\mathbf{x}})$, and $\|\mathbf{u}_0 - \mathbf{w}_0\|$ is greater than threshold $\delta r/(2\sqrt{d})$, then at least one of two sequences $\{\mathbf{u}_t\}, \{\mathbf{w}_t\}$ will “efficiently escape saddle point”. In other words, if \mathbf{u}_0 is a point in the stuck region, consider any point \mathbf{w}_0 that is on a straight line along direction of \mathbf{e}_1 . As long as \mathbf{w}_0 is slightly far ($\delta r/\sqrt{d}$) from \mathbf{u}_0 , it must be in the escaping region. This is what we mean by the “width” of the stuck region being small. Now we prove the main Lemma using this observation:

Proof of Lemma 14. By adding perturbation, in worst case we increase function value by:

$$f(\mathbf{x}_0) - f(\tilde{\mathbf{x}}) \leq \nabla f(\tilde{\mathbf{x}})^\top \xi + \frac{\ell}{2} \|\xi\|^2 \leq \mathcal{G} \left(\frac{\mathcal{S}}{\kappa \cdot \log(\frac{d\kappa}{\delta})} \right) + \frac{1}{2} \ell \left(\frac{\mathcal{S}}{\kappa \cdot \log(\frac{d\kappa}{\delta})} \right)^2 \leq \frac{3}{2} \mathcal{F}$$

On the other hand, let radius $r = \frac{\mathcal{S}}{\kappa \cdot \log(\frac{d\kappa}{\delta})}$. We know \mathbf{x}_0 come froms uniform distribution over $\mathbb{B}_{\tilde{\mathbf{x}}}(r)$. Let $\mathcal{X}_{\text{stuck}} \subset \mathbb{B}_{\tilde{\mathbf{x}}}(r)$ denote the set of bad starting points so that if $\mathbf{x}_0 \in \mathcal{X}_{\text{stuck}}$, then

$f(\mathbf{x}_T) - f(\mathbf{x}_0) > -2.5\mathcal{F}$ (thus stuck at a saddle point); otherwise if $\mathbf{x}_0 \in B_{\tilde{\mathbf{x}}}(r) - \mathcal{X}_{\text{stuck}}$, we have $f(\mathbf{x}_T) - f(\mathbf{x}_0) \leq -2.5\mathcal{F}$.

By applying Lemma 15, we know for any $\mathbf{x}_0 \in \mathcal{X}_{\text{stuck}}$, it is guaranteed that $(\mathbf{x}_0 \pm \mu r \mathbf{e}_1) \notin \mathcal{X}_{\text{stuck}}$ where $\mu \in [\frac{\delta}{2\sqrt{d}}, 1]$. Denote $I_{\mathcal{X}_{\text{stuck}}}(\cdot)$ be the indicator function of being inside set $\mathcal{X}_{\text{stuck}}$; and vector $\mathbf{x} = (x^{(1)}, \mathbf{x}^{(-1)})$, where $x^{(1)}$ is the component along \mathbf{e}_1 direction, and $\mathbf{x}^{(-1)}$ is the remaining $d-1$ dimensional vector. Recall $\mathbb{B}^{(d)}(r)$ be d -dimensional ball with radius r ; By calculus, this gives an upper bound on the volume of $\mathcal{X}_{\text{stuck}}$:

$$\begin{aligned} \text{Vol}(\mathcal{X}_{\text{stuck}}) &= \int_{\mathbb{B}_{\tilde{\mathbf{x}}}^{(d)}(r)} d\mathbf{x} \cdot I_{\mathcal{X}_{\text{stuck}}}(\mathbf{x}) \\ &= \int_{\mathbb{B}_{\tilde{\mathbf{x}}}^{(d-1)}(r)} d\mathbf{x}^{(-1)} \int_{\tilde{x}^{(1)} - \sqrt{r^2 - \|\tilde{\mathbf{x}}^{(-1)} - \mathbf{x}^{(-1)}\|^2}}^{\tilde{x}^{(1)} + \sqrt{r^2 - \|\tilde{\mathbf{x}}^{(-1)} - \mathbf{x}^{(-1)}\|^2}} dx^{(1)} \cdot I_{\mathcal{X}_{\text{stuck}}}(\mathbf{x}) \\ &\leq \int_{\mathbb{B}_{\tilde{\mathbf{x}}}^{(d-1)}(r)} d\mathbf{x}^{(-1)} \cdot \left(2 \cdot \frac{\delta}{2\sqrt{d}} r\right) = \text{Vol}(\mathbb{B}_0^{(d-1)}(r)) \times \frac{\delta r}{\sqrt{d}} \end{aligned}$$

Then, we immediately have the ratio:

$$\frac{\text{Vol}(\mathcal{X}_{\text{stuck}})}{\text{Vol}(\mathbb{B}_{\tilde{\mathbf{x}}}^{(d)}(r))} \leq \frac{\frac{\delta r}{\sqrt{d}} \times \text{Vol}(\mathbb{B}_0^{(d-1)}(r))}{\text{Vol}(\mathbb{B}_0^{(d)}(r))} = \frac{\delta}{\sqrt{\pi d}} \frac{\Gamma(\frac{d}{2} + 1)}{\Gamma(\frac{d}{2} + \frac{1}{2})} \leq \frac{\delta}{\sqrt{\pi d}} \cdot \sqrt{\frac{d}{2} + \frac{1}{2}} \leq \delta$$

The second last inequality is by the property of Gamma function that $\frac{\Gamma(x+1)}{\Gamma(x+1/2)} < \sqrt{x + \frac{1}{2}}$ as long as $x \geq 0$. Therefore, with at least probability $1 - \delta$, $\mathbf{x}_0 \notin \mathcal{X}_{\text{stuck}}$. In this case, we have:

$$\begin{aligned} f(\mathbf{x}_T) - f(\tilde{\mathbf{x}}) &= f(\mathbf{x}_T) - f(\mathbf{x}_0) + f(\mathbf{x}_0) - f(0) \\ &\leq -2.5\mathcal{F} + 1.5\mathcal{F} \leq -\mathcal{F} \end{aligned}$$

which finishes the proof. \square

A.3 Bounding the Width of Stuck Region

In order to prove Lemma 15, we do it in two steps:

1. We first show if gradient descent from \mathbf{u}_0 does not decrease function value, then all the iterates must lie within a small ball around \mathbf{u}_0 (Lemma 16).
2. If gradient descent starting from a point \mathbf{u}_0 stuck in a small ball around a saddle point, then gradient descent from \mathbf{w}_0 (moving \mathbf{u}_0 along \mathbf{e}_1 direction for at least a certain distance), will decrease the function value (Lemma 17).

Recall we assumed without loss of generality \mathbf{e}_1 is the minimum eigenvector of $\nabla^2 f(\tilde{\mathbf{x}})$. In this context, we denote $\mathcal{H} := \nabla^2 f(\tilde{\mathbf{x}})$, and for simplicity of calculation, we consider following quadratic approximation:

$$\tilde{f}_{\mathbf{y}}(\mathbf{x}) := f(\mathbf{y}) + \nabla f(\mathbf{y})^\top (\mathbf{x} - \mathbf{y}) + \frac{1}{2} (\mathbf{x} - \mathbf{y})^\top \mathcal{H} (\mathbf{x} - \mathbf{y}) \quad (4)$$

Now we are ready to state two lemmas formally:

Lemma 16. For any constant $\hat{c} \geq 3$, there exists absolute constant c_{\max} : for any $\delta \in (0, \frac{d\kappa}{e}]$, let $f(\cdot), \tilde{\mathbf{x}}$ satisfies the condition in Lemma 14, for any initial point \mathbf{u}_0 with $\|\mathbf{u}_0 - \tilde{\mathbf{x}}\| \leq 2\mathcal{S}/(\kappa \cdot \log(\frac{d\kappa}{\delta}))$, define:

$$T = \min \left\{ \inf_t \left\{ t|\tilde{f}_{\mathbf{u}_0}(\mathbf{u}_t) - f(\mathbf{u}_0) \leq -3\mathcal{F} \right\}, \hat{c}\mathcal{T} \right\}$$

then, for any $\eta \leq c_{\max}/\ell$, we have for all $t < T$ that $\|\mathbf{u}_t - \tilde{\mathbf{x}}\| \leq 100(\mathcal{S} \cdot \hat{c})$.

Lemma 17. There exists absolute constant c_{\max}, \hat{c} such that: for any $\delta \in (0, \frac{d\kappa}{e}]$, let $f(\cdot), \tilde{\mathbf{x}}$ satisfies the condition in Lemma 14, and sequences $\{\mathbf{u}_t\}, \{\mathbf{w}_t\}$ satisfy the conditions in Lemma 15, define:

$$T = \min \left\{ \inf_t \left\{ t|\tilde{f}_{\mathbf{w}_0}(\mathbf{w}_t) - f(\mathbf{w}_0) \leq -3\mathcal{F} \right\}, \hat{c}\mathcal{T} \right\}$$

then, for any $\eta \leq c_{\max}/\ell$, if $\|\mathbf{u}_t - \tilde{\mathbf{x}}\| \leq 100(\mathcal{S} \cdot \hat{c})$ for all $t < T$, we will have $T < \hat{c}\mathcal{T}$.

Note the conclusion $T < \hat{c}\mathcal{T}$ in Lemma 17 equivalently means:

$$\inf_t \left\{ t|\tilde{f}_{\mathbf{w}_0}(\mathbf{w}_t) - f(\mathbf{w}_0) \leq -3\mathcal{F} \right\} < \hat{c}\mathcal{T}$$

That is, for some $T < \hat{c}\mathcal{T}$, $\{\mathbf{w}_t\}$ sequence “escape the saddle point” in the sense of sufficient function value decrease $\tilde{f}_{\mathbf{w}_0}(\mathbf{w}_t) - f(\mathbf{w}_0) \leq -3\mathcal{F}$. Now, we are ready to prove Lemma 15.

Proof of Lemma 15. W.L.O.G, let $\tilde{\mathbf{x}} = 0$ be the origin. Let $(c_{\max}^{(2)}, \hat{c})$ be the absolute constant so that Lemma 17 holds, also let $c_{\max}^{(1)}$ be the absolute constant to make Lemma 16 holds based on our current choice of \hat{c} . We choose $c_{\max} \leq \min\{c_{\max}^{(1)}, c_{\max}^{(2)}\}$ so that our learning rate $\eta \leq c_{\max}/\ell$ is small enough which make both Lemma 16 and Lemma 17 hold. Let $T^* := \hat{c}\mathcal{T}$ and define:

$$T' = \inf_t \left\{ t|\tilde{f}_{\mathbf{u}_0}(\mathbf{u}_t) - f(\mathbf{u}_0) \leq -3\mathcal{F} \right\}$$

Let's consider following two cases:

Case $T' \leq T^*$: In this case, by Lemma 16, we know $\|\mathbf{u}_{T'-1}\| \leq O(\mathcal{S})$, and therefore

$$\|\mathbf{u}_{T'}\| \leq \|\mathbf{u}_{T'-1}\| + \eta\|\nabla f(\mathbf{u}_{T'-1})\| \leq \|\mathbf{u}_{T'-1}\| + \eta\|\nabla f(\tilde{\mathbf{x}})\| + \eta\ell\|\mathbf{u}_{T'-1}\| \leq O(\mathcal{S})$$

By choosing c_{\max} small enough and $\eta \leq c_{\max}/\ell$, this gives:

$$\begin{aligned} f(\mathbf{u}_{T'}) - f(\mathbf{u}_0) &\leq \nabla f(\mathbf{u}_0)^\top (\mathbf{u}_{T'} - \mathbf{u}_0) + \frac{1}{2}(\mathbf{u}_{T'} - \mathbf{u}_0)^\top \nabla^2 f(\mathbf{u}_0)(\mathbf{u}_{T'} - \mathbf{u}_0) + \frac{\rho}{6}\|\mathbf{u}_{T'} - \mathbf{u}_0\|^3 \\ &\leq \tilde{f}_{\mathbf{u}_0}(\mathbf{u}_t) - f(\mathbf{u}_0) + \frac{\rho}{2}\|\mathbf{u}_0 - \tilde{\mathbf{x}}\|\|\mathbf{u}_{T'} - \mathbf{u}_0\|^2 + \frac{\rho}{6}\|\mathbf{u}_{T'} - \mathbf{u}_0\|^3 \\ &\leq -3\mathcal{F} + O(\rho\mathcal{S}^3) = -3\mathcal{F} + O(\sqrt{\eta\ell} \cdot \mathcal{F}) \leq -2.5\mathcal{F} \end{aligned}$$

By choose $c_{\max} \leq \min\{1, \frac{1}{\hat{c}}\}$. We know $\eta < \frac{1}{\hat{c}}$, by Lemma 12, we know gradient descent always decrease function value. Therefore, for any $T \geq \frac{1}{c_{\max}}\mathcal{T} \geq \hat{c}\mathcal{T} = T^* \geq T'$, we have:

$$f(\mathbf{u}_T) - f(\mathbf{u}_0) \leq f(\mathbf{u}_{T^*}) - f(\mathbf{u}_0) \leq f(\mathbf{u}_{T'}) - f(\mathbf{u}_0) \leq -2.5\mathcal{F}$$

Case $T' > T^*$: In this case, by Lemma 16, we know $\|\mathbf{u}_t\| \leq O(\mathcal{S})$ for all $t \leq T^*$. Define

$$T'' = \inf_t \left\{ t \mid \tilde{f}_{\mathbf{w}_0}(\mathbf{w}_t) - f(\mathbf{w}_0) \leq -2\mathcal{F} \right\}$$

By Lemma 17, we immediately have $T'' \leq T^*$. Apply same argument as in first case, we have for all $T \geq \frac{1}{c_{\max}} \mathcal{T}$ that $f(\mathbf{w}_T) - f(\mathbf{w}_0) \leq f(\mathbf{w}_{T^*}) - f(\mathbf{w}_0) \leq -2.5\mathcal{F}$. \square

Next we finish the proof by proving Lemma 16 and Lemma 17.

A.3.1 Proof of Lemma 16

In Lemma 16, we hope to show if the function value did not decrease, then all the iterations must be constrained in a small ball. We do that by analyzing the dynamics of the iterations, and we decompose the d -dimensional space into two subspaces: a subspace \mathcal{S} which is the span of significantly negative eigenvectors of the Hessian and its orthogonal complement.

Recall notation $\mathcal{H} := \nabla^2 f(\tilde{\mathbf{x}})$ and quadratic approximation $\tilde{f}_{\mathbf{y}}(\mathbf{x})$ as defined in Eq.(4). Since $\delta \in (0, \frac{d\kappa}{e}]$, we always have $\log(\frac{d\kappa}{\delta}) \geq 1$. W.L.O.G, set $\mathbf{u}_0 = 0$ to be the origin, by gradient descent update function, we have:

$$\begin{aligned} \mathbf{u}_{t+1} &= \mathbf{u}_t - \eta \nabla f(\mathbf{u}_t) \\ &= \mathbf{u}_t - \eta \nabla f(0) - \eta \left[\int_0^1 \nabla^2 f(\theta \mathbf{u}_t) d\theta \right] \mathbf{u}_t \\ &= \mathbf{u}_t - \eta \nabla f(0) - \eta (\mathcal{H} + \Delta_t) \mathbf{u}_t \\ &= (\mathbf{I} - \eta \mathcal{H} - \eta \Delta_t) \mathbf{u}_t - \eta \nabla f(0) \end{aligned} \tag{5}$$

Here, $\Delta_t := \int_0^1 \nabla^2 f(\theta \mathbf{u}_t) d\theta - \mathcal{H}$. By Hessian Lipschitz, we have $\|\Delta_t\| \leq \rho(\|\mathbf{u}_t\| + \|\tilde{\mathbf{x}}\|)$, and by smoothness of the gradient, we have $\|\nabla f(0)\| \leq \|\nabla f(\tilde{\mathbf{x}})\| + \ell \|\tilde{\mathbf{x}}\| \leq \mathcal{G} + \ell \cdot 2\mathcal{S}/(\kappa \cdot \log(\frac{d\kappa}{\delta})) \leq 3\mathcal{G}$.

We will now compute the projections of \mathbf{u}_t in different eigenspaces of \mathcal{H} . Let \mathcal{S} be the subspace spanned by all eigenvectors of \mathcal{H} whose eigenvalue is less than $-\frac{\gamma}{\hat{c} \log(\frac{d\kappa}{\delta})}$. \mathcal{S}^c denotes the subspace of remaining eigenvectors. Let $\boldsymbol{\alpha}_t$ and $\boldsymbol{\beta}_t$ denote the projections of \mathbf{u}_t onto \mathcal{S} and \mathcal{S}^c respectively i.e., $\boldsymbol{\alpha}_t = \mathcal{P}_{\mathcal{S}} \mathbf{u}_t$, and $\boldsymbol{\beta}_t = \mathcal{P}_{\mathcal{S}^c} \mathbf{u}_t$. We can decompose the update equations Eq.(5) into:

$$\boldsymbol{\alpha}_{t+1} = (\mathbf{I} - \eta \mathcal{H}) \boldsymbol{\alpha}_t - \eta \mathcal{P}_{\mathcal{S}} \Delta_t \mathbf{u}_t - \eta \mathcal{P}_{\mathcal{S}} \nabla f(0) \tag{6}$$

$$\boldsymbol{\beta}_{t+1} = (\mathbf{I} - \eta \mathcal{H}) \boldsymbol{\beta}_t - \eta \mathcal{P}_{\mathcal{S}^c} \Delta_t \mathbf{u}_t - \eta \mathcal{P}_{\mathcal{S}^c} \nabla f(0) \tag{7}$$

By definition of T , we know for all $t < T$:

$$-3\mathcal{F} < \tilde{f}_0(\mathbf{u}_t) - f(0) = \nabla f(0)^\top \mathbf{u}_t - \frac{1}{2} \mathbf{u}_t^\top \mathcal{H} \mathbf{u}_t \leq \nabla f(0)^\top \mathbf{u}_t - \frac{\gamma}{2} \frac{\|\boldsymbol{\alpha}_t\|^2}{\hat{c} \log(\frac{d\kappa}{\delta})} + \frac{1}{2} \boldsymbol{\beta}_t^\top \mathcal{H} \boldsymbol{\beta}_t$$

Combined with the fact $\|\mathbf{u}_t\|^2 = \|\boldsymbol{\alpha}_t\|^2 + \|\boldsymbol{\beta}_t\|^2$, we have:

$$\begin{aligned} \|\mathbf{u}_t\|^2 &\leq \frac{2\hat{c} \log(\frac{d\kappa}{\delta})}{\gamma} \left(3\mathcal{F} + \nabla f(0)^\top \mathbf{u}_t + \frac{1}{2} \boldsymbol{\beta}_t^\top \mathcal{H} \boldsymbol{\beta}_t \right) + \|\boldsymbol{\beta}_t\|^2 \\ &\leq 14 \cdot \max \left\{ \frac{\mathcal{G} \hat{c} \log(\frac{d\kappa}{\delta})}{\gamma} \|\mathbf{u}_t\|, \frac{\mathcal{F} \hat{c} \log(\frac{d\kappa}{\delta})}{\gamma}, \frac{\boldsymbol{\beta}_t^\top \mathcal{H} \boldsymbol{\beta}_t \hat{c} \log(\frac{d\kappa}{\delta})}{\gamma}, \|\boldsymbol{\beta}_t\|^2 \right\} \end{aligned}$$

where last inequality is due to $\|\nabla f(0)\| \leq 3\mathcal{G}$. This gives:

$$\|\mathbf{u}_t\| \leq 14 \cdot \max \left\{ \frac{\mathcal{G} \hat{c} \log(\frac{d\kappa}{\delta})}{\gamma}, \sqrt{\frac{\mathcal{F} \hat{c} \log(\frac{d\kappa}{\delta})}{\gamma}}, \sqrt{\frac{\boldsymbol{\beta}_t^\top \mathcal{H} \boldsymbol{\beta}_t \hat{c} \log(\frac{d\kappa}{\delta})}{\gamma}}, \|\boldsymbol{\beta}_t\| \right\} \quad (8)$$

Now, we use induction to prove that

$$\|\mathbf{u}_t\| \leq 100(\mathcal{S} \cdot \hat{c}) \quad (9)$$

Clearly Eq.(9) is true for $t = 0$ since $\mathbf{u}_0 = 0$. Suppose Eq.(9) is true for all $\tau \leq t$. We will now show that Eq.(9) holds for $t+1 < T$. Note that by the definition of \mathcal{S} , \mathcal{F} and \mathcal{G} , we only need to bound the last two terms of Eq.(8) i.e., $\|\boldsymbol{\beta}_{t+1}\|$ and $\boldsymbol{\beta}_{t+1}^\top \mathcal{H} \boldsymbol{\beta}_{t+1}$.

By update function of $\boldsymbol{\beta}_t$ (Eq.(7)), we have:

$$\boldsymbol{\beta}_{t+1} \leq (\mathbf{I} - \eta \mathcal{H}) \boldsymbol{\beta}_t + \eta \boldsymbol{\delta}_t \quad (10)$$

and the norm of $\boldsymbol{\delta}_t$ is bounded as follows:

$$\begin{aligned} \|\boldsymbol{\delta}_t\| &\leq \|\Delta_t\| \|\mathbf{u}_t\| + \|\nabla f(0)\| \\ &\leq \rho (\|\mathbf{u}_t\| + \|\tilde{\mathbf{x}}\|) \|\mathbf{u}_t\| + \|\nabla f(0)\| \\ &\leq \rho \cdot 100\hat{c}(100\hat{c} + 2/(\kappa \cdot \log(\frac{d\kappa}{\delta}))) \mathcal{S}^2 + \mathcal{G} \\ &= [100\hat{c}(100\hat{c} + 2)\sqrt{\eta\ell} + 1]\mathcal{G} \leq 2\mathcal{G} \end{aligned} \quad (11)$$

The last step follows by choosing small enough constant $c_{\max} \leq \frac{1}{100\hat{c}(100\hat{c}+2)}$ and stepsize $\eta < c_{\max}/\ell$.

Bounding $\|\boldsymbol{\beta}_{t+1}\|$: Combining Eq.(10), Eq.(11) and using the definition of \mathcal{S}^c , we have:

$$\|\boldsymbol{\beta}_{t+1}\| \leq (1 + \frac{\eta\gamma}{\hat{c} \log(\frac{d\kappa}{\delta})}) \|\boldsymbol{\beta}_t\| + 2\eta\mathcal{G}$$

Since $\|\boldsymbol{\beta}_0\| = 0$ and $t+1 \leq T$, by applying above relation recurrively, we have:

$$\|\boldsymbol{\beta}_{t+1}\| \leq \sum_{\tau=0}^t 2(1 + \frac{\eta\gamma}{\hat{c} \log(\frac{d\kappa}{\delta})})^\tau \eta\mathcal{G} \leq 2 \cdot 3 \cdot T\eta\mathcal{G} \leq 6(\mathcal{S} \cdot \hat{c}) \quad (12)$$

The second last inequality is because $T \leq \hat{c}\mathcal{T}$ by definition, so that $(1 + \frac{\eta\gamma}{\hat{c} \log(\frac{d\kappa}{\delta})})^T \leq 3$.

Bounding $\boldsymbol{\beta}_{t+1}^\top \mathcal{H} \boldsymbol{\beta}_{t+1}$: Using Eq.(10), we can also write the update equation as:

$$\boldsymbol{\beta}_t = \sum_{\tau=0}^{t-1} (\mathbf{I} - \eta \mathcal{H})^\tau \boldsymbol{\delta}_\tau$$

Combining with Eq.(11), this gives

$$\begin{aligned}
\beta_{t+1}^\top \mathcal{H} \beta_{t+1} &= \eta^2 \sum_{\tau_1=0}^t \sum_{\tau_2=0}^t \delta_{\tau_1}^\top (\mathbf{I} - \eta \mathcal{H})^{\tau_1} \mathcal{H} (\mathbf{I} - \eta \mathcal{H})^{\tau_2} \delta_{\tau_2} \\
&\leq \eta^2 \sum_{\tau_1=0}^t \sum_{\tau_2=0}^t \|\delta_{\tau_1}\| \|(\mathbf{I} - \eta \mathcal{H})^{\tau_1} \mathcal{H} (\mathbf{I} - \eta \mathcal{H})^{\tau_2}\| \|\delta_{\tau_2}\| \\
&\leq 4\eta^2 \mathcal{G}^2 \sum_{\tau_1=0}^t \sum_{\tau_2=0}^t \|(\mathbf{I} - \eta \mathcal{H})^{\tau_1} \mathcal{H} (\mathbf{I} - \eta \mathcal{H})^{\tau_2}\|
\end{aligned}$$

Let the eigenvalues of \mathcal{H} to be $\{\lambda_i\}$, then for any $\tau_1, \tau_2 \geq 0$, we know the eigenvalues of $(\mathbf{I} - \eta \mathcal{H})^{\tau_1} \mathcal{H} (\mathbf{I} - \eta \mathcal{H})^{\tau_2}$ are $\{\lambda_i(1 - \eta \lambda_i)^{\tau_1 + \tau_2}\}$. Let $g_t(\lambda) := \lambda(1 - \eta \lambda)^t$, and setting its derivative to zero, we obtain:

$$\nabla g_t(\lambda) = (1 - \eta \lambda)^t - t\eta \lambda (1 - \eta \lambda)^{t-1} = 0$$

We see that $\lambda_t^* = \frac{1}{(1+t)\eta}$ is the unique maximizer, and $g_t(\lambda)$ is monotonically increasing in $(-\infty, \lambda_t^*]$. This gives:

$$\|(\mathbf{I} - \eta \mathcal{H})^{\tau_1} \mathcal{H} (\mathbf{I} - \eta \mathcal{H})^{\tau_2}\| = \max_i \lambda_i (1 - \eta \lambda_i)^{\tau_1 + \tau_2} \leq \hat{\lambda} (1 - \eta \hat{\lambda})^{\tau_1 + \tau_2} \leq \frac{1}{(1 + \tau_1 + \tau_2)\eta}$$

where $\hat{\lambda} = \min\{\ell, \lambda_{\tau_1 + \tau_2}^*\}$. Therefore, we have:

$$\begin{aligned}
\beta_{t+1}^\top \mathcal{H} \beta_{t+1} &\leq 4\eta^2 \mathcal{G}^2 \sum_{\tau_1=0}^t \sum_{\tau_2=0}^t \|(\mathbf{I} - \eta \mathcal{H})^{\tau_1} \mathcal{H} (\mathbf{I} - \eta \mathcal{H})^{\tau_2}\| \\
&\leq 4\eta^2 \mathcal{G}^2 \sum_{\tau_1=0}^t \sum_{\tau_2=0}^t \frac{1}{1 + \tau_1 + \tau_2} \leq 8\eta T \mathcal{G}^2 \leq 8\mathcal{S}^2 \gamma \hat{c} \cdot \log^{-1}\left(\frac{d\kappa}{\delta}\right)
\end{aligned} \tag{13}$$

The second last inequality is because by rearrange summation:

$$\sum_{\tau_1=0}^t \sum_{\tau_2=0}^t \frac{1}{1 + \tau_1 + \tau_2} = \sum_{\tau=0}^{2t} \min\{1 + \tau, 2t + 1 - \tau\} \cdot \frac{1}{1 + \tau} \leq 2t + 1 < 2T$$

Finally, substitue Eq.(12) and Eq.(13) into Eq.(8), this gives:

$$\begin{aligned}
\|\mathbf{u}_{t+1}\| &\leq 14 \cdot \max \left\{ \frac{\mathcal{G} \hat{c} \log(\frac{d\kappa}{\delta})}{\gamma}, \sqrt{\frac{\mathcal{F} \hat{c} \log(\frac{d\kappa}{\delta})}{\gamma}}, \sqrt{\frac{\beta_t^\top \mathcal{H} \beta_t \hat{c} \log(\frac{d\kappa}{\delta})}{\gamma}}, \|\beta_t\| \right\} \\
&\leq 100(\mathcal{S} \cdot \hat{c})
\end{aligned}$$

This finishes the induction as well as the proof of the lemma. \square

A.3.2 Proof of Lemma 17

In this Lemma we try to show if all the iterates from \mathbf{u}_0 are constrained in a small ball, iterates from \mathbf{w}_0 must be able to decrease the function value. In order to do that, we keep track of vector \mathbf{v} which is the difference between \mathbf{u} and \mathbf{w} . Similar as before, we also decompose \mathbf{v} into different eigenspaces. However, this time we only care about the projection of \mathbf{v} on the direction \mathbf{e}_1 and its orthogonal subspace.

Again, recall notation $\mathcal{H} := \nabla^2 f(\tilde{\mathbf{x}})$, \mathbf{e}_1 as minimum eigenvector of \mathcal{H} and quadratic approximation $\tilde{f}_{\mathbf{y}}(\mathbf{x})$ as defined in Eq.(4). Since $\delta \in (0, \frac{d\kappa}{e}]$, we always have $\log(\frac{d\kappa}{\delta}) \geq 1$. W.L.O.G, set $\mathbf{u}_0 = 0$ to be the origin. Define $\mathbf{v}_t = \mathbf{w}_t - \mathbf{u}_t$, by assumptions in Lemma 17, we have $\mathbf{v}_0 = \mu[\mathcal{S}/(\kappa \cdot \log(\frac{d\kappa}{\delta}))]\mathbf{e}_1$, $\mu \in [\delta/(2\sqrt{d}), 1]$. Now, consider the update equation for \mathbf{w}_t :

$$\begin{aligned} \mathbf{u}_{t+1} + \mathbf{v}_{t+1} &= \mathbf{w}_{t+1} = \mathbf{w}_t - \eta \nabla f(\mathbf{w}_t) \\ &= \mathbf{u}_t + \mathbf{v}_t - \eta \nabla f(\mathbf{u}_t + \mathbf{v}_t) \\ &= \mathbf{u}_t + \mathbf{v}_t - \eta \nabla f(\mathbf{u}_t) - \eta \left[\int_0^1 \nabla^2 f(\mathbf{u}_t + \theta \mathbf{v}_t) d\theta \right] \mathbf{v}_t \\ &= \mathbf{u}_t + \mathbf{v}_t - \eta \nabla f(\mathbf{u}_t) - \eta(\mathcal{H} + \Delta'_t) \mathbf{v}_t \\ &= \mathbf{u}_t - \eta \nabla f(\mathbf{u}_t) + (\mathbf{I} - \eta \mathcal{H} - \eta \Delta'_t) \mathbf{v}_t \end{aligned}$$

where $\Delta'_t := \int_0^1 \nabla^2 f(\mathbf{u}_t + \theta \mathbf{v}_t) d\theta - \mathcal{H}$. By Hessian Lipschitz, we have $\|\Delta'_t\| \leq \rho(\|\mathbf{u}_t\| + \|\mathbf{v}_t\| + \|\tilde{\mathbf{x}}\|)$. This gives the dynamic for \mathbf{v}_t satisfy:

$$\mathbf{v}_{t+1} = (\mathbf{I} - \eta \mathcal{H} - \eta \Delta'_t) \mathbf{v}_t \quad (14)$$

Since $\|\mathbf{w}_0 - \tilde{\mathbf{x}}\| \leq \|\mathbf{u}_0 - \tilde{\mathbf{x}}\| + \|\mathbf{v}_0\| \leq \mathcal{S}/(\kappa \cdot \log(\frac{d\kappa}{\delta}))$, directly applying Lemma 16, we know $\mathbf{w}_t \leq 100(\mathcal{S} \cdot \hat{c})$ for all $t \leq T$. By condition of Lemma 17, we know $\|\mathbf{u}_t\| \leq 100(\mathcal{S} \cdot \hat{c})$ for all $t < T$. This gives:

$$\|\mathbf{v}_t\| \leq \|\mathbf{u}_t\| + \|\mathbf{w}_t\| \leq 200(\mathcal{S} \cdot \hat{c}) \text{ for all } t < T \quad (15)$$

This in sum gives for $t < T$:

$$\|\Delta'_t\| \leq \rho(\|\mathbf{u}_t\| + \|\mathbf{v}_t\| + \|\tilde{\mathbf{x}}\|) \leq \rho(300\hat{c}\mathcal{S} + \mathcal{S}/(\kappa \cdot \log(\frac{d\kappa}{\delta}))) \leq \rho\mathcal{S}(300\hat{c} + 1)$$

On the other hand, denote ψ_t be the norm of \mathbf{v}_t projected onto \mathbf{e}_1 direction, and φ_t be the norm of \mathbf{v}_t projected onto remaining subspace. Eq.(14) gives us:

$$\begin{aligned} \psi_{t+1} &\geq (1 + \gamma\eta)\psi_t - \mu\sqrt{\psi_t^2 + \varphi_t^2} \\ \varphi_{t+1} &\leq (1 + \gamma\eta)\varphi_t + \mu\sqrt{\psi_t^2 + \varphi_t^2} \end{aligned}$$

where $\mu = \eta\rho\mathcal{S}(300\hat{c} + 1)$. We will now prove via induction that for all $t < T$:

$$\varphi_t \leq 4\mu t \cdot \psi_t \quad (16)$$

By hypothesis of Lemma 17, we know $\varphi_0 = 0$, thus the base case of induction holds. Assume Eq.(16) is true for $\tau \leq t$, For $t + 1 \leq T$, we have:

$$\begin{aligned} 4\mu(t+1)\psi_{t+1} &\geq 4\mu(t+1) \left((1 + \gamma\eta)\psi_t - \mu\sqrt{\psi_t^2 + \varphi_t^2} \right) \\ \varphi_{t+1} &\leq 4\mu t(1 + \gamma\eta)\psi_t + \mu\sqrt{\psi_t^2 + \varphi_t^2} \end{aligned}$$

From above inequalities, we see that we only need to show:

$$(1 + 4\mu(t + 1)) \sqrt{\psi_t^2 + \varphi_t^2} \leq 4(1 + \gamma\eta)\psi_t$$

By choosing $\sqrt{c_{\max}} \leq \frac{1}{300\hat{c}+1} \min\{\frac{1}{2\sqrt{2}}, \frac{1}{4\hat{c}}\}$, and $\eta \leq c_{\max}/\ell$, we have

$$4\mu(t + 1) \leq 4\mu T \leq 4\eta\rho\mathcal{S}(300\hat{c} + 1)\hat{c}\mathcal{T} = 4\sqrt{\eta\ell}(300\hat{c} + 1)\hat{c} \leq 1$$

This gives:

$$4(1 + \gamma\eta)\psi_t \geq 4\psi_t \leq 2\sqrt{2\psi_t^2} \geq (1 + 4\mu(t + 1)) \sqrt{\psi_t^2 + \varphi_t^2}$$

which finishes the induction.

Now, we know $\varphi_t \leq 4\mu t \cdot \psi_t \leq \psi_t$, this gives:

$$\psi_{t+1} \geq (1 + \gamma\eta)\psi_t - \sqrt{2}\mu\psi_t \geq (1 + \frac{\gamma\eta}{2})\psi_t \quad (17)$$

where the last step follows from $\mu = \eta\rho\mathcal{S}(300\hat{c} + 1) \leq \sqrt{c_{\max}}(300\hat{c} + 1)\gamma\eta \cdot \log^{-1}(\frac{d\kappa}{\delta}) < \frac{\gamma\eta}{2\sqrt{2}}$.

Finally, combining Eq.(15) and (17) we have for all $t < T$:

$$\begin{aligned} 200(\mathcal{S} \cdot \hat{c}) &\geq \|\mathbf{v}_t\| \geq \psi_t \geq (1 + \frac{\gamma\eta}{2})^t \psi_0 \\ &= (1 + \frac{\gamma\eta}{2})^t c_0 \frac{\mathcal{S}}{\kappa} \log^{-1}(\frac{d\kappa}{\delta}) \geq (1 + \frac{\gamma\eta}{2})^t \frac{\delta}{2\sqrt{d}} \frac{\mathcal{S}}{\kappa} \log^{-1}(\frac{d\kappa}{\delta}) \end{aligned}$$

This implies:

$$T < \frac{1}{2} \frac{\log(400 \frac{\kappa\sqrt{d}}{\delta} \cdot \hat{c} \log(\frac{d\kappa}{\delta}))}{\log(1 + \frac{\gamma\eta}{2})} \leq \frac{\log(400 \frac{\kappa\sqrt{d}}{\delta} \cdot \hat{c} \log(\frac{d\kappa}{\delta}))}{\gamma\eta} \leq (2 + \log(400\hat{c}))\mathcal{T}$$

The last inequality is due to $\delta \in (0, \frac{d\kappa}{e}]$ we have $\log(\frac{d\kappa}{\delta}) \geq 1$. By choosing constant \hat{c} to be large enough to satisfy $2 + \log(400\hat{c}) \leq \hat{c}$, we will have $T < \hat{c}\mathcal{T}$, which finishes the proof. \square

B Improve Convergence by Local Structure

In this section, we show if the objective function has nice *local structure* (e.g. satisfies Assumptions A3.a or A3.b), then it is possible to combine our analysis with the local analysis in order to get very fast convergence to a local minimum.

In particular, we prove Theorem 5.

Theorem 5. *There exist absolute constant c_{\max} such that: if $f(\cdot)$ satisfies A1, A2, and A3.a (or A3.b), then for any $\delta > 0, \epsilon > 0, \Delta_f \geq f(\mathbf{x}_0) - f^*$, constant $c \leq c_{\max}$, let $\tilde{\epsilon} = \min(\theta, \gamma^2/\rho)$, with probability $1 - \delta$, the output of PGDLi($\mathbf{x}_0, \ell, \rho, \tilde{\epsilon}, c, \delta, \Delta_f, \beta$) will be ϵ -close to \mathcal{X}^* in iterations:*

$$O\left(\frac{\ell(f(\mathbf{x}_0) - f^*)}{\tilde{\epsilon}^2} \log^4\left(\frac{\ell\Delta_f}{\tilde{\epsilon}^2\delta}\right) + \frac{\beta}{\alpha} \log \frac{\zeta}{\epsilon}\right)$$

Proof. Theorem 5 runs PGDli($\mathbf{x}_0, \ell, \rho, \tilde{\epsilon}, c, \delta, \Delta_f, \beta$). According to algorithm 3, we know it calls PGD($\mathbf{x}_0, \ell, \rho, \epsilon, c, \delta, \Delta_f$) first (denote its output as $\hat{\mathbf{x}}$), then run standard gradient descent with learning rate $\frac{1}{\beta}$ starting from $\hat{\mathbf{x}}$.

By Corollary 4, we know $\hat{\mathbf{x}}$ is already in the ζ -neighborhood of \mathcal{X}^* , where \mathcal{X}^* is the set of local minima. Therefore, to prove this theorem, we only need to show prove following two claims:

1. Suppose $\{\mathbf{x}_t\}$ is the sequence of gradient descent starting from $\mathbf{x}_0 = \hat{\mathbf{x}}$ with learning rate $\frac{1}{\beta}$, then \mathbf{x}_t is always in the ζ -neighborhood of \mathcal{X}^* .
2. Local structure (assumption A3.a or A3.b) allows iterates to converge to points ϵ -close to \mathcal{X}^* within $O(\frac{\beta}{\alpha} \log \frac{\zeta}{\epsilon})$ iterations.

We will focus on Assumption A3.b (as we will later see Assumption A3.a is a special case of Assumption A3.b). Assume \mathbf{x}_t is in ζ -neighborhood of \mathcal{X}^* , by gradient updates and the definition of projection, we have:

$$\begin{aligned} \|\mathbf{x}_{t+1} - \mathcal{P}_{\mathcal{X}^*}(\mathbf{x}_{t+1})\|^2 &\leq \|\mathbf{x}_{t+1} - \mathcal{P}_{\mathcal{X}^*}(\mathbf{x}_t)\|^2 = \|\mathbf{x}_t - \eta \nabla f(\mathbf{x}_t) - \mathcal{P}_{\mathcal{X}^*}(\mathbf{x}_t)\|^2 \\ &= \|\mathbf{x}_t - \mathcal{P}_{\mathcal{X}^*}(\mathbf{x}_t)\|^2 - 2\eta \langle \nabla f(\mathbf{x}_t), \mathbf{x}_t - \mathcal{P}_{\mathcal{X}^*}(\mathbf{x}_t) \rangle + \eta^2 \|\nabla f(\mathbf{x}_t)\|^2 \\ &\leq \|\mathbf{x}_t - \mathcal{P}_{\mathcal{X}^*}(\mathbf{x}_t)\|^2 - \eta\alpha \|\mathbf{x}_t - \mathcal{P}_{\mathcal{X}^*}(\mathbf{x}_t)\|^2 + (\eta^2 - \frac{\eta}{\beta}) \|\nabla f(\mathbf{x})\|^2 \\ &\leq (1 - \frac{\alpha}{\beta}) \|\mathbf{x}_t - \mathcal{P}_{\mathcal{X}^*}(\mathbf{x}_t)\|^2 \end{aligned}$$

The second last inequality is due to (α, β) -regularity condition. The last inequality is because of the choice $\eta = \frac{1}{\beta}$.

There are two consequences of this calculation. First, it shows $\|\mathbf{x}_{t+1} - \mathcal{P}_{\mathcal{X}^*}(\mathbf{x}_{t+1})\|^2 \leq \|\mathbf{x}_t - \mathcal{P}_{\mathcal{X}^*}(\mathbf{x}_t)\|^2$. As a result if \mathbf{x}_t in ζ -neighborhood of \mathcal{X}^* , \mathbf{x}_{t+1} is also in this ζ -neighborhood. Since \mathbf{x}_0 is in the ζ -neighborhood by Corollary 4, by induction we know all later iterations are in the same neighborhood.

Now, since we know all the points \mathbf{x}_t are in the neighborhood, the equation also shows linear convergence rate $(1 - \frac{\alpha}{\beta})$. The initial distance is bounded by $\|\mathbf{x}_0 - \mathcal{P}_{\mathcal{X}^*}(\mathbf{x}_0)\| \leq \zeta$, therefore to converge to points ϵ -close to \mathcal{X}^* , we only need the following number of iterations:

$$\frac{\log(\epsilon/\zeta)}{\log(1 - \alpha/\beta)} = O(\frac{\beta}{\alpha} \log \frac{\zeta}{\epsilon}).$$

This finishes the proof under Assumption A3.b.

Finally, we argue assumption A3.a implies A3.b. First, notice that if a function is locally strongly convex, then its local minima are isolated: for any two points $\mathbf{x}, \mathbf{x}' \in \mathcal{X}^*$, the local region $B_{\mathbf{x}}(\zeta)$ and $B_{\mathbf{x}'}(\zeta)$ must be disjoint (otherwise function $f(\mathbf{x})$ is strongly convex in connected domain $B_{\mathbf{x}}(\zeta) \cup B_{\mathbf{x}'}(\zeta)$ but has two distinct local minima, which is impossible). Therefore, W.L.O.G, it suffices to consider one particular disjoint region, with unique local minimum we denote as \mathbf{x}^* , clearly, for all $\mathbf{x} \in B_{\mathbf{x}^*}(\zeta)$ we have $\mathcal{P}_{\mathcal{X}^*}(\mathbf{x}) = \mathbf{x}^*$.

Now by α -strong convexity:

$$f(\mathbf{x}^*) \geq f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{x}^* - \mathbf{x} \rangle + \frac{\alpha}{2} \|\mathbf{x} - \mathbf{x}^*\|^2 \quad (18)$$

On the other hand, for any \mathbf{x} in this ζ -neighborhood, we already proved $\mathbf{x} - \frac{1}{\beta}\nabla f(\mathbf{x})$ also in this ζ -neighborhood. By β -smoothness, we also have:

$$f(\mathbf{x} - \frac{1}{\beta}\nabla f(\mathbf{x})) \leq f(\mathbf{x}) - \frac{1}{2\beta}\|\nabla f(\mathbf{x})\|^2 \quad (19)$$

Combining Eq.(18) and Eq.(19), and using the fact $f(\mathbf{x}^*) \leq f(\mathbf{x} - \frac{1}{\beta}\nabla f(\mathbf{x}))$, we get:

$$\langle \nabla f(\mathbf{x}), \mathbf{x} - \mathbf{x}^* \rangle \geq \frac{\alpha}{2}\|\mathbf{x} - \mathbf{x}^*\|^2 + \frac{1}{2\beta}\|\nabla f(\mathbf{x})\|^2$$

which finishes the proof. \square

C Geometric Structures of Matrix Factorization Problem

In this Section we investigate the global geometric structures of the matrix factorization problem. These properties are summarized in Lemmas 6 and 7. Such structures allow us to apply our main Theorem and get fast convergence (as shown in Theorem 8).

Note that our main results Theorems 3 and 5 are proved for functions $f(\cdot)$ whose input \mathbf{x} is a vector. For the current function in 2, though the input $\mathbf{U} \in \mathbb{R}^{d \times r}$ is a matrix, we can always vectorize it to be a vector in \mathbb{R}^{dr} and apply our results. However, for simplicity of presentation, we still write everything in matrix form (without explicit vectorization), while the reader should keep in mind the operations are same if one vectorizes everything first.

Recall for vectors we use $\|\cdot\|$ to denote the 2-norm, and for matrices we use $\|\cdot\|$ and $\|\cdot\|_F$ to denote spectral norm, and Frobenius norm respectively. Furthermore, we always use $\sigma_i(\cdot)$ to denote the i -th largest singular value of the matrix.

We first show how the geometric properties (Lemma 6 and Lemma 7) imply a fast convergence (Theorem 8).

Theorem 8. *There exists an absolute constant c_{\max} such that the following holds. For matrix factorization (2), for any $\delta > 0$ and constant $c \leq c_{\max}$, let $\Gamma^{1/2} := 2 \max\{\|\mathbf{U}_0\|, 3(\sigma_1^*)^{1/2}\}$, suppose we run PGDLi($\mathbf{U}_0, 8\Gamma, 12\Gamma^{1/2}, \frac{(\sigma_r^*)^2}{108\Gamma^{1/2}}, c, \delta, \frac{r\Gamma^2}{2}, 10\sigma_1^*$), then:*

1. *With probability 1, the iterates satisfy $\|\mathbf{U}_t\| \leq \Gamma^{1/2}$ for every $t \geq 0$.*
2. *With probability $1 - \delta$, the output will be ϵ -close to global minima set \mathcal{X}^* in following iterations:*

$$O\left(r \left(\frac{\Gamma}{\sigma_r^*}\right)^4 \log^4\left(\frac{d\Gamma}{\delta\sigma_r^*}\right) + \frac{\sigma_1^*}{\sigma_r^*} \log \frac{\sigma_r^*}{\epsilon}\right)$$

Proof of Theorem 8. Denote \tilde{c}_{\max} to be the absolute constant allowed in Theorem 5. In this theorem, we let $c_{\max} = \min\{\tilde{c}_{\max}, \frac{1}{2}\}$, and choose any constant $c \leq c_{\max}$.

Theorem 8 consists of two parts. In part 1 we show that the iterations never bring the matrix to a very large norm, while in part 2 we apply our main Theorem to get fast convergence. We will first prove the bound on number of iterations assuming the bound on the norm. We will then proceed to prove part 1.

Part 2: Assume part 1 of the theorem is true i.e., with probability 1, the iterates satisfy $\|\mathbf{U}_t\| \leq \Gamma^{1/2}$ for every $t \geq 0$. In this case, although we are doing unconstrained optimization, we can still use the geometric properties that hold inside this region. .

By Lemma 6 and 7, we know objective function Eq.(2) is 8Γ -smooth, $12\Gamma^{1/2}$ -Lipschitz Hessian, $(\frac{1}{24}(\sigma_r^*)^{3/2}, \frac{1}{3}\sigma_r^*, \frac{1}{3}(\sigma_r^*)^{1/2})$ -strict saddle, and holds $(\frac{2}{3}\sigma_r^*, 10\sigma_1^*)$ -regularity condition in $\frac{1}{3}(\sigma_r^*)^{1/2}$ neighborhood of local minima (also global minima) \mathcal{X}^* . Furthermore, note $f^* = 0$ and recall $\Gamma^{1/2} = 2 \max\{\|\mathbf{U}_0\|, 3(\sigma_1^*)^{1/2}\}$, then, we have:

$$f(\mathbf{U}_0) - f^* = \|\mathbf{U}_0 \mathbf{U}_0^\top - \mathbf{M}^*\|_F^2 \leq 2r \|\mathbf{U}_0 \mathbf{U}_0^\top - \mathbf{M}^*\|^2 \leq \frac{r\Gamma^2}{2}.$$

Thus, we can choose $\Delta_f = \frac{r\Gamma^2}{2}$. Substituting the corresponding parameters from Theorem 5, we know by running $\text{PGDli}(\mathbf{U}_0, 8\Gamma, 12\Gamma^{1/2}, \frac{(\sigma_r^*)^2}{108\Gamma^{1/2}}, c, \delta, \frac{r\Gamma^2}{2}, 10\sigma_1^*)$, with probability $1 - \delta$, the output will be ϵ -close to global minima set \mathcal{X}^* in iterations:

$$O\left(r \left(\frac{\Gamma}{\sigma_r^*}\right)^4 \log^4\left(\frac{d\Gamma}{\delta\sigma_r^*}\right) + \frac{\sigma_1^*}{\sigma_r^*} \log \frac{\sigma_r^*}{\epsilon}\right).$$

Part 1: We will now show part 1 of the theorem. Recall PGDli (Algorithm 3) runs PGD (Algorithm 2) first, and then runs gradient descent within $\frac{1}{3}(\sigma_r^*)^{1/2}$ neighborhood of \mathcal{X}^* . It is easy to verify that $\frac{1}{3}(\sigma_r^*)^{1/2}$ neighborhood of \mathcal{X}^* is a subset of $\{\mathbf{U} \mid \|\mathbf{U}\|^2 \leq \Gamma\}$. Therefore, we only need to show that first phase PGD will not leave the region. Specifically, we now use induction to prove the following for PGD:

1. Suppose at iteration τ we add perturbation, and denote $\tilde{\mathbf{U}}_\tau$ to be the iterate before adding perturbation (i.e., $\mathbf{U}_\tau = \tilde{\mathbf{U}}_\tau + \xi_\tau$, and $\tilde{\mathbf{U}}_\tau = \mathbf{U}_{\tau-1} - \eta \nabla f(\mathbf{U}_{\tau-1})$). Then, $\|\tilde{\mathbf{U}}_\tau\| \leq \frac{1}{2}\Gamma$, and
2. $\|\mathbf{U}_t\| \leq \Gamma$ for all $t \geq 0$.

By choice of parameters of Algorithm 2, we know $\eta = \frac{c}{8\Gamma}$. First, consider gradient descent step without perturbations:

$$\begin{aligned} \|\mathbf{U}_{t+1}\| &= \|\mathbf{U}_t - \eta \nabla f(\mathbf{U}_t)\| = \|\mathbf{U}_t - \eta(\mathbf{U}_t \mathbf{U}_t^\top - \mathbf{M}^*)\mathbf{U}_t\| \\ &\leq \|\mathbf{U}_t - \eta \mathbf{U}_t \mathbf{U}_t^\top \mathbf{U}_t\| + \eta \|\mathbf{M}^* \mathbf{U}_t\| \\ &\leq \max_i [\sigma_i(\mathbf{U}_t) - \eta \sigma_i^3(\mathbf{U}_t)] + \eta \|\mathbf{M}^* \mathbf{U}_t\| \end{aligned}$$

For the first term, we know function $f(t) = t - \eta t^3$ is monotonically increasing in $[0, 1/\sqrt{3\eta}]$. On the other hand, by induction assumption, we also know $\|\mathbf{U}_t\| \leq \Gamma^{1/2} \leq \sqrt{8\Gamma/(3c)} = 1/\sqrt{3\eta}$. Therefore, the max is taken when $i = 1$:

$$\begin{aligned} \|\mathbf{U}_{t+1}\| &\leq \|\mathbf{U}_t\| - \eta \|\mathbf{U}_t\|^3 + \eta \|\mathbf{M}^* \mathbf{U}_t\| \\ &\leq \|\mathbf{U}_t\| - \eta (\|\mathbf{U}_t\|^2 - \sigma_1^*) \|\mathbf{U}_t\|. \end{aligned} \tag{20}$$

We separate our discussion into following cases.

Case $\|\mathbf{U}_t\| > \frac{1}{2}\Gamma^{1/2}$: In this case $\|\mathbf{U}_t\| \geq \max\{\|\mathbf{U}_0\|, 3(\sigma_1^*)^{1/2}\}$. Recall $\Gamma^{1/2} = 2 \max\{\|\mathbf{U}_0\|, 3(\sigma_1^*)^{1/2}\}$. Clearly, $\Gamma \geq 36\sigma_1^*$, we know:

$$\begin{aligned}\|\mathbf{U}_{t+1}\| &\leq \|\mathbf{U}_t\| - \eta(\|\mathbf{U}_t\|^2 - \sigma_1^*)\|\mathbf{U}_t\| \leq \|\mathbf{U}_t\| - \frac{c}{8\Gamma}(\frac{1}{4}\Gamma - \sigma_1^*)\frac{1}{2}\Gamma^{1/2} \\ &\leq \|\mathbf{U}_t\| - \frac{c}{8\Gamma}(\frac{1}{4}\Gamma - \frac{1}{36}\Gamma)\frac{1}{2}\Gamma^{1/2} = \|\mathbf{U}_t\| - \frac{c}{72}\Gamma^{1/2}.\end{aligned}$$

This means that in each iteration, the spectral norm would decrease by at least $\frac{c}{72}\Gamma^{1/2}$.

Case $\|\mathbf{U}_t\| \leq \frac{1}{2}\Gamma^{1/2}$: From (20), we know that as long as $\|\mathbf{U}_t\|^2 \geq \|\mathbf{M}^*\|$, we will always have $\|\mathbf{U}_{t+1}\| \leq \|\mathbf{U}_t\| \leq \frac{1}{2}\Gamma^{1/2}$. For $\|\mathbf{U}_t\|^2 \leq \|\mathbf{M}^*\|$, we have:

$$\begin{aligned}\|\mathbf{U}_{t+1}\| &\leq \|\mathbf{U}_t\| - \eta(\|\mathbf{U}_t\|^2 - \sigma_1^*)\|\mathbf{U}_t\| = \|\mathbf{U}_t\| + \frac{c}{8\Gamma}(\sigma_1^* - \|\mathbf{U}_t\|^2)\|\mathbf{U}_t\| \\ &\leq \|\mathbf{U}_t\| + ((\sigma_1^*)^{1/2} - \|\mathbf{U}_t\|) \times \frac{c}{8\Gamma}((\sigma_1^*)^{1/2} + \|\mathbf{U}_t\|)\|\mathbf{U}_t\| \\ &\leq \|\mathbf{U}_t\| + ((\sigma_1^*)^{1/2} - \|\mathbf{U}_t\|) \times \frac{c\sigma_1^*}{4\Gamma} \leq (\sigma_1^*)^{1/2}\end{aligned}$$

Thus, in this case, we always have $\|\mathbf{U}_{t+1}\| \leq \frac{1}{2}\Gamma^{1/2}$.

In conclusion, if we don't add perturbation in iteration t , we have:

- If $\|\mathbf{U}_t\| > \frac{1}{2}\Gamma^{1/2}$, then $\|\mathbf{U}_{t+1}\| \leq \|\mathbf{U}_t\| - \frac{c}{72}\Gamma^{1/2}$.
- If $\|\mathbf{U}_t\| \leq \frac{1}{2}\Gamma^{1/2}$, then $\|\mathbf{U}_{t+1}\| \leq \frac{1}{2}\Gamma^{1/2}$.

Now consider the iterations where we add perturbation. By the choice of radius of perturbation in Algorithm 2, we increase spectral norm by at most :

$$\|\xi_t\| \leq \|\xi_t\|_F \leq \frac{\sqrt{c}}{\chi^2} \frac{(\sigma_r^*)^2}{108\Gamma^{1/2} \cdot 8\Gamma} \leq \frac{1}{2}\Gamma^{1/2}$$

The first inequality is because $\chi \geq 1$ and $c \leq 1$. That is, if before perturbation we have $\|\tilde{\mathbf{U}}_t\| \leq \frac{1}{2}\Gamma^{1/2}$, then $\|\mathbf{U}_t\| = \|\tilde{\mathbf{U}}_t + \xi_t\| \leq \Gamma^{1/2}$.

On the other hand, according to Algorithm 2, once we add perturbation, we will not add perturbation for next $t_{\text{thres}} = \frac{\chi \cdot 24\Gamma}{c^2 \sigma_r^*} \geq \frac{24}{c^2} \geq \frac{48}{c}$ iterations. Let $T = \min\{\inf_i\{\|\mathbf{U}_{t+i}\| \leq \frac{1}{2}\Gamma^{1/2}\}, t_{\text{thres}}\}$:

$$\|\mathbf{U}_{t+T-1}\| \leq \|\mathbf{U}_t\| - \frac{c}{72}\Gamma^{1/2}(T-1) \leq \Gamma^{1/2}(1 - \frac{c(T-1)}{72})$$

This gives $T \leq \frac{36}{c} < \frac{48}{c} \leq t_{\text{thres}}$. Let $\tau > t$ be the next time when we add perturbation ($\tau \geq t + t_{\text{thres}}$), we immediately know $\|\mathbf{U}_{T+i}\| \leq \frac{1}{2}\Gamma^{1/2}$ for $0 \leq i < \tau - T$ and $\|\tilde{\mathbf{U}}_\tau\| \leq \frac{1}{2}\Gamma^{1/2}$.

Finally, $\|\mathbf{U}_0\| \leq \frac{1}{2}\Gamma^{1/2}$ by definition of Γ , so the initial condition holds. This finishes induction and the proof of the theorem. \square

In the next subsections we prove the geometric structures.

C.1 Smoothness and Hessian Lipschitz

Before we start proofs of lemmas, we first state some properties about gradient and Hessians. The gradient of the objective function $f(\mathbf{U})$ is

$$\nabla f(\mathbf{U}) = 2(\mathbf{U}\mathbf{U}^\top - \mathbf{M}^\star)\mathbf{U}.$$

Furthermore, we have the gradient and Hessian satisfy for any $\mathbf{Z} \in \mathbb{R}^{d \times r}$:

$$\langle \nabla f(\mathbf{U}), \mathbf{Z} \rangle = 2\langle (\mathbf{U}\mathbf{U}^\top - \mathbf{M}^\star)\mathbf{U}, \mathbf{Z} \rangle, \text{ and} \quad (21)$$

$$\nabla^2 f(\mathbf{U})(\mathbf{Z}, \mathbf{Z}) = \|\mathbf{U}\mathbf{Z}^\top + \mathbf{Z}\mathbf{U}^\top\|_F^2 + 2\langle \mathbf{U}\mathbf{U}^\top - \mathbf{M}^\star, \mathbf{Z}\mathbf{Z}^\top \rangle. \quad (22)$$

Lemma 6. For any $\Gamma \geq \sigma_1^\star$, inside the region $\{\mathbf{U} \mid \|\mathbf{U}\|^2 < \Gamma\}$, $f(\mathbf{U})$ defined in Eq.(2) is 8Γ -smooth and $12\Gamma^{1/2}$ -Hessian Lipschitz.

Proof. Denote $\mathcal{D} = \{\mathbf{U} \mid \|\mathbf{U}\|^2 < \Gamma\}$, and recall $\Gamma \geq \sigma_1^\star$.

Smoothness: For any $\mathbf{U}, \mathbf{V} \in \mathcal{D}$, we have:

$$\begin{aligned} \|\nabla f(\mathbf{U}) - \nabla f(\mathbf{V})\|_F &= 2\|(\mathbf{U}\mathbf{U}^\top - \mathbf{M}^\star)\mathbf{U} - (\mathbf{V}\mathbf{V}^\top - \mathbf{M}^\star)\mathbf{V}\|_F \\ &\leq 2\left[\|\mathbf{U}\mathbf{U}^\top\mathbf{U} - \mathbf{V}\mathbf{V}^\top\mathbf{V}\|_F + \|\mathbf{M}^\star(\mathbf{U} - \mathbf{V})\|_F\right] \\ &\leq 2[3 \cdot \Gamma\|\mathbf{U} - \mathbf{V}\|_F + \sigma_1^\star\|\mathbf{U} - \mathbf{V}\|_F] \leq 8\Gamma \cdot \|\mathbf{U} - \mathbf{V}\|_F. \end{aligned}$$

The last line is due to following decomposition and triangle inequality:

$$\mathbf{U}\mathbf{U}^\top\mathbf{U} - \mathbf{V}\mathbf{V}^\top\mathbf{V} = \mathbf{U}\mathbf{U}^\top(\mathbf{U} - \mathbf{V}) + \mathbf{U}(\mathbf{U} - \mathbf{V})^\top\mathbf{V} + (\mathbf{U} - \mathbf{V})\mathbf{V}^\top\mathbf{V}.$$

Hessian Lipschitz: For any $\mathbf{U}, \mathbf{V} \in \mathcal{D}$, and any $\mathbf{Z} \in \mathbb{R}^{d \times r}$, according to Eq.(22), we have:

$$|\nabla^2 f(\mathbf{U})(\mathbf{Z}, \mathbf{Z}) - \nabla^2 f(\mathbf{V})(\mathbf{Z}, \mathbf{Z})| = \underbrace{\|\mathbf{U}\mathbf{Z}^\top + \mathbf{Z}\mathbf{U}^\top\|_F^2 - \|\mathbf{V}\mathbf{Z}^\top + \mathbf{Z}\mathbf{V}^\top\|_F^2}_{\mathfrak{A}} + \underbrace{2\langle \mathbf{U}\mathbf{U}^\top - \mathbf{V}\mathbf{V}^\top, \mathbf{Z}\mathbf{Z}^\top \rangle}_{\mathfrak{B}}.$$

For term \mathfrak{A} , we have:

$$\begin{aligned} \mathfrak{A} &= \langle \mathbf{U}\mathbf{Z}^\top + \mathbf{Z}\mathbf{U}^\top, (\mathbf{U} - \mathbf{V})\mathbf{Z}^\top + \mathbf{Z}(\mathbf{U} - \mathbf{V})^\top \rangle + \langle (\mathbf{U} - \mathbf{V})\mathbf{Z}^\top + \mathbf{Z}(\mathbf{U} - \mathbf{V})^\top, \mathbf{V}\mathbf{Z}^\top + \mathbf{Z}\mathbf{V}^\top \rangle \\ &\leq \|\mathbf{U}\mathbf{Z}^\top + \mathbf{Z}\mathbf{U}^\top\|_F \|(\mathbf{U} - \mathbf{V})\mathbf{Z}^\top + \mathbf{Z}(\mathbf{U} - \mathbf{V})^\top\|_F + \|(\mathbf{U} - \mathbf{V})\mathbf{Z}^\top + \mathbf{Z}(\mathbf{U} - \mathbf{V})^\top\|_F \|\mathbf{V}\mathbf{Z}^\top + \mathbf{Z}\mathbf{V}^\top\|_F \\ &\leq 4\|\mathbf{U}\| \|\mathbf{Z}\|_F^2 \|\mathbf{U} - \mathbf{V}\|_F + 4\|\mathbf{V}\| \|\mathbf{Z}\|_F^2 \|\mathbf{U} - \mathbf{V}\|_F \leq 8\Gamma^{1/2} \|\mathbf{Z}\|_F^2 \|\mathbf{U} - \mathbf{V}\|_F. \end{aligned}$$

For term \mathfrak{B} , we have:

$$\mathfrak{B} \leq 2\|\mathbf{U}\mathbf{U}^\top - \mathbf{V}\mathbf{V}^\top\|_F \|\mathbf{Z}\mathbf{Z}^\top\|_F \leq 4\Gamma^{1/2} \|\mathbf{Z}\|_F^2 \|\mathbf{U} - \mathbf{V}\|_F.$$

The inequality is due to following decomposition and triangle inequality:

$$\mathbf{U}\mathbf{U}^\top - \mathbf{V}\mathbf{V}^\top = \mathbf{U}(\mathbf{U} - \mathbf{V})^\top + (\mathbf{U} - \mathbf{V})\mathbf{V}^\top.$$

Therefore, in sum we have:

$$\begin{aligned} \max_{\mathbf{Z}: \|\mathbf{Z}\|_F \leq 1} |\nabla^2 f(\mathbf{U})(\mathbf{Z}, \mathbf{Z}) - \nabla^2 f(\mathbf{V})(\mathbf{Z}, \mathbf{Z})| &\leq \max_{\mathbf{Z}: \|\mathbf{Z}\|_F \leq 1} 12\Gamma^{1/2} \|\mathbf{Z}\|_F^2 \|\mathbf{U} - \mathbf{V}\|_F \\ &\leq 12\Gamma^{1/2} \|\mathbf{U} - \mathbf{V}\|_F. \end{aligned}$$

□

C.2 Strict-Saddle Property and Local Regularity

Recall the gradient and Hessian of objective function is calculated as in Eq.(21) and Eq.(22). We first prove an elementary inequality regarding to the trace of product of two symmetric PSD matrices. This lemma will be frequently used in the proof.

Lemma 18. *For $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{d \times d}$ both symmetric PSD matrices, we have:*

$$\sigma_{\min}(\mathbf{A})\text{tr}(\mathbf{B}) \leq \text{tr}(\mathbf{AB}) \leq \|\mathbf{A}\|\text{tr}(\mathbf{B})$$

Proof. Let $\mathbf{A} = \mathbf{VDV}^\top$ be the eigendecomposition of \mathbf{A} , where \mathbf{D} is diagonal matrix, and \mathbf{V} is orthogonal matrix. Then we have:

$$\text{tr}(\mathbf{AB}) = \text{tr}(\mathbf{DV}^\top \mathbf{BV}) = \sum_{i=1}^d \mathbf{D}_{ii}(\mathbf{V}^\top \mathbf{BV})_{ii}.$$

Since \mathbf{B} is PSD, we know $\mathbf{V}^\top \mathbf{BV}$ is also PSD, thus the diagonal entries are non-negative. That is, $(\mathbf{V}^\top \mathbf{BV})_{ii} \geq 0$ for all $i = 1, \dots, d$. Finally, the lemma follows from the fact that $\sigma_{\min}(\mathbf{A}) \leq \mathbf{D}_{ii} \leq \|\mathbf{A}\|$ and $\text{tr}(\mathbf{V}^\top \mathbf{BV}) = \text{tr}(\mathbf{BVV}^\top) = \text{tr}(\mathbf{B})$. \square

Now, we are ready to prove Lemma 7.

Lemma 7. *For $f(\mathbf{U})$ defined in Eq.(2), all local minima are global minima. The set of global minima is $\mathcal{X}^* = \{\mathbf{V}^* \mathbf{R} | \mathbf{R} \mathbf{R}^\top = \mathbf{R}^\top \mathbf{R} = \mathbf{I}\}$. Furthermore, $f(\mathbf{U})$ satisfies:*

1. $(\frac{1}{24}(\sigma_r^*)^{3/2}, \frac{1}{3}\sigma_r^*, \frac{1}{3}(\sigma_r^*)^{1/2})$ -strict saddle property, and
2. $(\frac{2}{3}\sigma_r^*, 10\sigma_1^*)$ -regularity condition in $\frac{1}{3}(\sigma_r^*)^{1/2}$ neighborhood of \mathcal{X}^* .

Proof. Let us denote the set $\mathcal{X}^* := \{\mathbf{V}^* \mathbf{R} | \mathbf{R} \mathbf{R}^\top = \mathbf{R}^\top \mathbf{R} = \mathbf{I}\}$, in the end of proof, we will show this set is the set of all local minima (which is also global minima).

Throughout the proof of this lemma, we always focus on the first-order and second-order property for one matrix \mathbf{U} . For simplicity of calculation, when it is clear from the context, we denote $\mathbf{U}^* = \mathcal{P}_{\mathcal{X}^*}(\mathbf{U})$ and $\Delta = \mathbf{U} - \mathcal{P}_{\mathcal{X}^*}(\mathbf{U})$. By definition of \mathcal{X}^* , we know $\mathbf{U}^* = \mathbf{V}^* \mathbf{R}_\mathbf{U}$ and $\Delta = \mathbf{U} - \mathbf{V}^* \mathbf{R}_\mathbf{U}$, where

$$\mathbf{R}_\mathbf{U} = \underset{\mathbf{R}: \mathbf{R} \mathbf{R}^\top = \mathbf{R}^\top \mathbf{R} = \mathbf{I}}{\text{argmin}} \|\mathbf{U} - \mathbf{V}^* \mathbf{R}\|_F^2$$

We first prove following claim, which will be used in many places across this proof:

$$\mathbf{U}^\top \mathbf{U}^* = \mathbf{U}^\top \mathbf{V}^* \mathbf{R}_\mathbf{U} \text{ is a symmetric PSD matrix.} \quad (23)$$

This because by expanding the Frobenius norm, and letting the SVD of $\mathbf{V}^{*\top} \mathbf{U}$ be \mathbf{ADB}^\top , we have:

$$\begin{aligned} \underset{\mathbf{R}: \mathbf{R} \mathbf{R}^\top = \mathbf{R}^\top \mathbf{R} = \mathbf{I}}{\text{argmin}} \|\mathbf{U} - \mathbf{V}^* \mathbf{R}\|_F^2 &= \underset{\mathbf{R}: \mathbf{R} \mathbf{R}^\top = \mathbf{R}^\top \mathbf{R} = \mathbf{I}}{\text{argmin}} -\langle \mathbf{U}, \mathbf{V}^* \mathbf{R} \rangle \\ &= \underset{\mathbf{R}: \mathbf{R} \mathbf{R}^\top = \mathbf{R}^\top \mathbf{R} = \mathbf{I}}{\text{argmin}} -\text{tr}(\mathbf{U}^\top \mathbf{V}^* \mathbf{R}) = \underset{\mathbf{R}: \mathbf{R} \mathbf{R}^\top = \mathbf{R}^\top \mathbf{R} = \mathbf{I}}{\text{argmin}} -\text{tr}(\mathbf{DA}^\top \mathbf{RB}) \end{aligned}$$

Since $\mathbf{A}, \mathbf{B}, \mathbf{R}$ are all orthonormal matrix, we know $\mathbf{A}^\top \mathbf{RB}$ is also orthonormal matrix. Moreover for any orthonormal matrix \mathbf{T} , we have:

$$\text{tr}(\mathbf{DT}) = \sum_i \mathbf{D}_{ii} \mathbf{T}_{ii} \leq \sum_i \mathbf{D}_{ii}$$

The last inequality is because \mathbf{D}_{ii} is singular value thus non-negative, and \mathbf{T} is orthonormal, thus $\mathbf{T}_{ii} \leq 1$. This means the maximum of $\text{tr}(\mathbf{D}\mathbf{T})$ is achieved when $\mathbf{T} = \mathbf{I}$, i.e., the minimum of $-\text{tr}(\mathbf{D}\mathbf{A}^\top \mathbf{R}\mathbf{B})$ is achieved when $\mathbf{R} = \mathbf{A}\mathbf{B}^\top$. Therefore, $\mathbf{U}^\top \mathbf{V}^* \mathbf{R}_\mathbf{U} = \mathbf{B}\mathbf{D}\mathbf{A}^\top \mathbf{A}\mathbf{B}^\top = \mathbf{B}\mathbf{D}\mathbf{B}^\top$ is symmetric PSD matrix.

Strict Saddle Property: In order to show the strict saddle property, we only need to show that for any \mathbf{U} satisfying $\|\nabla f(\mathbf{U})\|_F \leq \frac{1}{24}(\sigma_r^*)^{3/2}$ and $\|\Delta\|_F = \|\mathbf{U} - \mathbf{U}^*\|_F \geq \frac{1}{3}(\sigma_r^*)^{1/2}$, we always have $\sigma_{\min}(\nabla^2 f(\mathbf{U})) \leq -\frac{1}{3}\sigma_r^*$.

Let's consider Hessian $\nabla^2(\mathbf{U})$ in the direction of $\Delta = \mathbf{U} - \mathbf{U}^*$. Clearly, we have:

$$\mathbf{U}\mathbf{U}^\top - \mathbf{M}^* = \mathbf{U}\mathbf{U}^\top - (\mathbf{U} - \Delta)(\mathbf{U} - \Delta)^\top = (\mathbf{U}\Delta^\top + \Delta\mathbf{U}^\top) - \Delta\Delta^\top$$

and by (21):

$$\begin{aligned} \langle \nabla f(\mathbf{U}), \Delta \rangle &= 2\langle (\mathbf{U}\mathbf{U}^\top - \mathbf{M}^*)\mathbf{U}, \Delta \rangle = \langle \mathbf{U}\mathbf{U}^\top - \mathbf{M}^*, \Delta\mathbf{U}^\top + \mathbf{U}\Delta^\top \rangle \\ &= \langle \mathbf{U}\mathbf{U}^\top - \mathbf{M}^*, \mathbf{U}\mathbf{U}^\top - \mathbf{M}^* + \Delta\Delta^\top \rangle \end{aligned}$$

Therefore, by Eq.(22) and above two equalities, we have:

$$\begin{aligned} \nabla^2 f(\mathbf{U})(\Delta, \Delta) &= \|\mathbf{U}\Delta^\top + \Delta\mathbf{U}^\top\|_F^2 + 2\langle \mathbf{U}\mathbf{U}^\top - \mathbf{M}^*, \Delta\Delta^\top \rangle \\ &= \|\mathbf{U}\mathbf{U}^\top - \mathbf{M}^* + \Delta\Delta^\top\|_F^2 + 2\langle \mathbf{U}\mathbf{U}^\top - \mathbf{M}^*, \Delta\Delta^\top \rangle \\ &= \|\Delta\Delta^\top\|_F^2 - 3\|\mathbf{U}\mathbf{U}^\top - \mathbf{M}^*\|_F^2 + 4\langle \mathbf{U}\mathbf{U}^\top - \mathbf{M}^*, \mathbf{U}\mathbf{U}^\top - \mathbf{M}^* + \Delta\Delta^\top \rangle \\ &= \|\Delta\Delta^\top\|_F^2 - 3\|\mathbf{U}\mathbf{U}^\top - \mathbf{M}^*\|_F^2 + 4\langle \nabla f(\mathbf{U}), \Delta \rangle \end{aligned}$$

Consider the first two terms, by expanding, we have:

$$\begin{aligned} 3\|\mathbf{U}\mathbf{U}^\top - \mathbf{M}^*\|_F^2 - \|\Delta\Delta^\top\|_F^2 &= 3\|(\mathbf{U}^*\Delta^\top + \Delta\mathbf{U}^{*\top}) + \Delta\Delta^\top\|_F^2 - \|\Delta\Delta^\top\|_F^2 \\ &= 3 \cdot \text{tr} \left(2\mathbf{U}^{*\top} \mathbf{U}^* \Delta^\top \Delta + 2(\mathbf{U}^{*\top} \Delta)^2 + 4\mathbf{U}^{*\top} \Delta \Delta^\top \Delta + (\Delta^\top \Delta)^2 \right) - \text{tr}((\Delta^\top \Delta)^2) \\ &= \text{tr} \left(6\mathbf{U}^{*\top} \mathbf{U}^* \Delta^\top \Delta + 6(\mathbf{U}^{*\top} \Delta)^2 + 12\mathbf{U}^{*\top} \Delta \Delta^\top \Delta + 2(\Delta^\top \Delta)^2 \right) \\ &= \text{tr}((4\sqrt{3} - 6)\mathbf{U}^{*\top} \mathbf{U}^* \Delta^\top \Delta + (12 - 4\sqrt{3})\mathbf{U}^{*\top} (\mathbf{U}^* + \Delta) \Delta^\top \Delta + 2(\sqrt{3}\mathbf{U}^{*\top} \Delta + \Delta^\top \Delta)^2) \\ &\geq (4\sqrt{3} - 6)\text{tr}(\mathbf{U}^{*\top} \mathbf{U}^* \Delta^\top \Delta) \geq (4\sqrt{3} - 6)\sigma_r^* \|\Delta\|_F^2 \end{aligned}$$

where the second last inequality is because $\mathbf{U}^{*\top} (\mathbf{U}^* + \Delta) \Delta^\top \Delta = \mathbf{U}^{*\top} \mathbf{U} \Delta^\top \Delta$ is the product of two symmetric PSD matrices (thus its trace is non-negative); the last inequality is by Lemma 18.

Finally, in case we have $\|\nabla f(\mathbf{U})\|_F \leq \frac{1}{24}(\sigma_r^*)^{3/2}$ and $\|\Delta\|_F = \|\mathbf{U} - \mathbf{U}^*\|_F \geq \frac{1}{3}(\sigma_r^*)^{1/2}$

$$\begin{aligned} \sigma_{\min}(\nabla^2 f(\mathbf{U})) &\leq \frac{1}{\|\Delta\|_F^2} \nabla^2 f(\mathbf{U})(\Delta, \Delta) \leq -(4\sqrt{3} - 6)\sigma_r^* + 4 \frac{\langle \nabla f(\mathbf{U}), \Delta \rangle}{\|\Delta\|_F^2} \\ &\leq -(4\sqrt{3} - 6)\sigma_r^* + 4 \frac{\|\nabla f(\mathbf{U})\|_F}{\|\Delta\|_F} \leq -(4\sqrt{3} - 6.5)\sigma_r^* \leq -\frac{1}{3}\sigma_r^* \end{aligned}$$

Local Regularity: In $\frac{1}{3}(\sigma_r^*)^{1/2}$ neighborhood of \mathcal{X}^* , by definition, we know,

$$\|\Delta\|_F^2 = \|\mathbf{U} - \mathbf{U}^*\|_F^2 \leq \frac{1}{9}\sigma_r^*.$$

Clearly, by Weyl's inequality, we have $\|\mathbf{U}\| \leq \|\mathbf{U}^\star\| + \|\Delta\| \leq \frac{4}{3}(\sigma_1^\star)^{1/2}$, and $\sigma_r(\mathbf{U}) \geq \sigma_r(\mathbf{U}^\star) - \|\Delta\| \geq \frac{2}{3}(\sigma_r^\star)^{1/2}$. Moreover, since $\mathbf{U}^{\star\top}\mathbf{U}$ is symmetric matrix, we have:

$$\begin{aligned}\sigma_r(\mathbf{U}^{\star\top}\mathbf{U}) &= \frac{1}{2} \left(\sigma_r(\mathbf{U}^\top\mathbf{U}^\star + \mathbf{U}^{\star\top}\mathbf{U}) \right) \\ &\geq \frac{1}{2} \left(\sigma_r(\mathbf{U}^\top\mathbf{U} + \mathbf{U}^{\star\top}\mathbf{U}^\star) - \|(\mathbf{U} - \mathbf{U}^\star)^\top(\mathbf{U} - \mathbf{U}^\star)\| \right) \\ &\geq \frac{1}{2} \left(\sigma_r(\mathbf{U}^\top\mathbf{U}) + \sigma_r(\mathbf{U}^{\star\top}\mathbf{U}^\star) - \|\Delta\|_\text{F}^2 \right) \\ &\geq \frac{1}{2} \left(1 + \frac{4}{9} - \frac{1}{9} \right) \sigma_r^\star = \frac{2}{3} \sigma_r^\star.\end{aligned}$$

At a highlevel, we will prove (α, β) -regularity property (1) by proving that:

1. $\langle \nabla f(\mathbf{x}), \mathbf{x} - \mathcal{P}_{\mathcal{X}^\star}(\mathbf{x}) \rangle \geq \alpha \|\mathbf{x} - \mathcal{P}_{\mathcal{X}^\star}(\mathbf{x})\|^2$, and
2. $\langle \nabla f(\mathbf{x}), \mathbf{x} - \mathcal{P}_{\mathcal{X}^\star}(\mathbf{x}) \rangle \geq \frac{1}{\beta} \|\nabla f(\mathbf{x})\|^2$.

According to (21), we know:

$$\begin{aligned}\langle \nabla f(\mathbf{U}), \mathbf{U} - \mathcal{P}_{\mathcal{X}^\star}(\mathbf{U}) \rangle &= 2\langle (\mathbf{U}\mathbf{U}^\top - \mathbf{M}^\star)\mathbf{U}, \Delta \rangle = 2\langle \mathbf{U}\Delta^\top + \Delta\mathbf{U}^{\star\top}, \Delta\mathbf{U}^\top \rangle \\ &= 2(\text{tr}(\mathbf{U}\Delta^\top\mathbf{U}\Delta^\top) + \text{tr}(\Delta\mathbf{U}^{\star\top}\mathbf{U}\Delta^\top)) \\ &= 2(\|\Delta^\top\mathbf{U}\|_\text{F}^2 + \text{tr}(\mathbf{U}^{\star\top}\mathbf{U}\Delta^\top\Delta)).\end{aligned}\tag{24}$$

The last equality is because $\Delta^\top\mathbf{U}$ is symmetric matrix. Since $\mathbf{U}^{\star\top}\mathbf{U}$ is symmetric PSD matrix, and recall $\sigma_r(\mathbf{U}^{\star\top}\mathbf{U}) \geq \frac{2}{3}\sigma_r^\star$, by Lemma 18 we have:

$$\langle \nabla f(\mathbf{U}), \mathbf{U} - \mathcal{P}_{\mathcal{X}^\star}(\mathbf{U}) \rangle \geq \sigma_r(\mathbf{U}^{\star\top}\mathbf{U})\text{tr}(\Delta^\top\Delta) \geq \frac{2}{3}\sigma_r^\star\|\Delta\|_\text{F}^2.\tag{25}$$

On the other hand, we also have:

$$\begin{aligned}\|\nabla f(\mathbf{U})\|_\text{F}^2 &= 4\langle (\mathbf{U}\mathbf{U}^\top - \mathbf{M}^\star)\mathbf{U}, (\mathbf{U}\mathbf{U}^\top - \mathbf{M}^\star)\mathbf{U} \rangle \\ &= 4\langle (\mathbf{U}\Delta^\top + \Delta\mathbf{U}^{\star\top})\mathbf{U}, (\mathbf{U}\Delta^\top + \Delta\mathbf{U}^{\star\top})\mathbf{U} \rangle \\ &= 4 \left(\underbrace{\text{tr}[(\Delta^\top\mathbf{U}\mathbf{U}^\top\Delta)\mathbf{U}^\top\mathbf{U}]}_{\mathfrak{A}} + 2\text{tr}[\Delta^\top\mathbf{U}\mathbf{U}^\top\mathbf{U}^\star\Delta^\top\mathbf{U}] + \underbrace{\text{tr}(\mathbf{U}^{\star\top}\mathbf{U}\mathbf{U}^\top\mathbf{U}^\star\Delta^\top\Delta)}_{\mathfrak{B}} \right).\end{aligned}$$

For term \mathfrak{A} , by Lemma 18, and $\Delta^\top\mathbf{U}$ being a symmetric matrix, we have:

$$\mathfrak{A} \leq \|\mathbf{U}^\top\mathbf{U}\| \|\Delta^\top\mathbf{U}\|_\text{F}^2 + 2\|\mathbf{U}^\top\mathbf{U}^\star\| \|\Delta^\top\mathbf{U}\|_\text{F}^2 \leq \left(\frac{16}{9} + \frac{8}{3}\right)\sigma_1^\star \|\Delta^\top\mathbf{U}\|_\text{F}^2 \leq 5\sigma_1^\star \|\Delta^\top\mathbf{U}\|_\text{F}^2$$

For term \mathfrak{B} , by Eq.(23) we can denote $\mathbf{C} = \mathbf{U}^{\star\top}\mathbf{U} = \mathbf{U}^\top\mathbf{U}^\star$ which is symmetric PSD matrix, by Lemma 18, we have:

$$\begin{aligned}\mathfrak{B} &= \text{tr}(\mathbf{C}^2\Delta^\top\Delta) = \text{tr}(\mathbf{C}(\mathbf{C}^{1/2}\Delta^\top\Delta\mathbf{C}^{1/2})) \\ &\leq \|\mathbf{C}\| \text{tr}(\mathbf{C}^{1/2}\Delta^\top\Delta\mathbf{C}^{1/2}) = \|\mathbf{C}\| \text{tr}(\mathbf{C}\Delta^\top\Delta) \leq \frac{4}{3}\sigma_1^\star \text{tr}(\mathbf{U}^{\star\top}\mathbf{U}\Delta^\top\Delta).\end{aligned}$$

Combining with (24) we have:

$$\|\nabla f(\mathbf{U})\|_{\text{F}}^2 \leq \sigma_1^*(20\|\Delta^\top \mathbf{U}\|_{\text{F}}^2 + \frac{16}{3}\text{tr}(\mathbf{U}^{\star\top} \mathbf{U} \Delta^\top \Delta)) \leq 10\sigma_1^* \langle \nabla f(\mathbf{U}), \mathbf{U} - \mathcal{P}_{\mathcal{X}^*}(\mathbf{U}) \rangle. \quad (26)$$

Combining (25) and (26), we have:

$$\langle \nabla f(\mathbf{U}), \mathbf{U} - \mathcal{P}_{\mathcal{X}^*}(\mathbf{U}) \rangle \geq \frac{1}{3}\sigma_r^* \|\mathbf{U} - \mathcal{P}_{\mathcal{X}^*}(\mathbf{U})\|_{\text{F}}^2 + \frac{1}{20\sigma_1^*} \|\nabla f(\mathbf{U})\|_{\text{F}}^2.$$

□