

A New Test of Multivariate Nonlinear Causality

Zhidong Bai

KLASMOE and School of Mathematics and Statistics,
Northeast Normal University, China

Yongchang Hui*

School of Mathematics and Statistics,
Xi'an Jiaotong University, China

Zhihui Lv

KLASMOE and School of Mathematics and Statistics,
Northeast Normal University, China

Wing-Keung Wong

Department of Finance, Asia University, Taiwan
Department of Economics, Lingnan University, Hong Kong

Shurong Zheng

KLASMOE and School of Mathematics and Statistics,
Northeast Normal University, China

Zhen-Zhen Zhu

KLASMOE and School of Mathematics and Statistics,
Northeast Normal University, China

*Corresponding author: Yongchang Hui, School of Mathematics and Statistics, Xi'an Jiaotong University. No.28, Xianning West Road, Xi'an, Shaanxi, P.R. China. Tel: (86)-029-82663170, Email: huiyc180@xjtu.edu.cn

A New Test of Multivariate Nonlinear Causality

Abstract The multivariate nonlinear Granger causality developed by Bai et al. (2010) plays an important role in detecting the dynamic interrelationships between two groups of variables. Following the idea of Hiemstra-Jones (HJ) test proposed by Hiemstra and Jones (1994), they attempt to establish a central limit theorem (CLT) of their test statistic by applying the asymptotical property of multivariate U -statistic. However, Bai et al. (2016) revisit the HJ test and find that the test statistic given by HJ is NOT a function of U -statistics which implies that the CLT neither proposed by Hiemstra and Jones (1994) nor the one extended by Bai et al. (2010) is valid for statistical inference. In this paper, we re-estimate the probabilities and reestablish the CLT of the new test statistic. Numerical simulation shows that our new estimates are consistent and our new test performs decent size and power.

Keywords: nonlinear Granger causality, Hiemstra-Jones test, multivariate

1 Introduction

After the pioneering work of Granger (1969), Granger causality tests have been developed into a set of useful methods to detect causal relations between time series in economics and finance. Linear Granger causality tests within the linear autoregressive model class have been developed in many directions, e.g., Hurlin et al. (2001) proposed a procedure for causality tests with panel data, Ghysels et al. (2016) test for Granger causality with mixed frequency data based on the multiple-horizon framework established by Dufour and Renault (1998) and Dufour et al. (2006). Though linear tests of Granger causality have been investigated very deeply, they are limited in their capability to detect nonlinear causality.

The real world is “almost certainly nonlinear” as Granger (1989) notes, so it is more important to test the nonlinear causality. Baek and Brock (1992) develop a nonlinear Granger causality test which is modified by Hiemstra and Jones (1994) later on to study the bivariate nonlinear causal relationship between two series. Among the various tests of nonlinear Granger causality, the Hiemstra-Jones test (hereafter, the HJ test) proposed by Hiemstra and Jones (1994) is the most cited by scholars and the most frequently applied by practitioners in economics and finance. There were over 1100 Google Scholar hits by September 2016, which illustrates its significance in the economics and finance literatures.

Bai et al. (2010) extend the HJ test from bivariate setting to multivariate setting catering to the practical needs that economic and financial factors usually move together and influence others in groups. This extension encourages a large amount of applications. For example, Lam et al. (2012) suggest to use such technics to make better investment decisions. Zheng and Chen (2013) proposed a complete double selection method in identifying external influential factors for a particular stock market. Choudhry et al. (2015) investigates the nonlinear dynamic co-movements between gold returns, stock market returns and stock market volatility during the recent global financial crisis. Choudhry et al. (2016) investigate the relationship between stock market volatility and the business cycle in four major economies US, Canada, Japan and the UK.

However, several works note that counterintuitive results are obtained from the HJ test, Diks and Panchenko (2005, 2006) find that the HJ test is seriously over-rejecting in

simulation studies. In accordance with the evidence presented by Diks and Panchenko (2005, 2006), Bai et al. (2016) reinvestigate the HJ test and reveal some of the underlying reasons for the questionable performance of HJ test. They find that the estimators of the probabilities in the definition are not U -statistics as Hiemstra and Jones (1994) claimed and the central limit theorem of the test statistics is not valid. Bai et al. (2016) propose a set of consistent estimators of the probabilities in the definition of Hiemstra and Jones (1994) and provide a new test statistic with its asymptotic distribution.

Considering the significant importance of the multivariate nonlinear Granger causality test, there is an urgent need to reinvestigate Bai et al. (2010) and extend Bai et al. (2016) to multivariate setting. The remainder of this paper is organized as follows. In Section 2, we simply review the procedure of the multivariate nonlinear Granger causality test (here after BWZ test) extended by Bai et al. (2010). In Section 3, we re-estimate the probabilities in the definition of Bai et al. (2010) and establish the asymptotic distribution of the new test statistics. Simulation results are presented in Section 4. Finally, we provide some concluding remarks in Section 5.

2 The Multivariate Nonlinear Causality Test Extended from HJ Test

Bai et al. (2010) consider two strictly stationary and weakly dependent vector time series processes $X_t = (X_{1,t}, X_{2,t}, \dots, X_{n_1,t})'$, $Y_t = (Y_{1,t}, Y_{2,t}, \dots, Y_{n_2,t})'$. The m_{x_i} -length lead vector of $X_{i,t}$ is defined as $X_{i,t}^{m_{x_i}} \equiv (X_{i,t}, X_{i,t+1}, \dots, X_{i,t+m_{x_i}-1})$, $m_{x_i} = 1, 2, \dots$, $t = 1, 2, \dots$, similarly L_{x_i} -length lag vector of $X_{i,t}$, and L_{y_i} -length lag vector of $Y_{i,t}$ are defined as $X_{i,t-L_{x_i}}^{L_{x_i}} \equiv (X_{i,t-L_{x_i}}, X_{i,t-L_{x_i}+1}, \dots, X_{i,t-1})$, $L_{x_i} = 1, 2, \dots$, $t = L_{x_i} + 1, L_{x_i} + 2, \dots$, $Y_{i,t-L_{y_i}}^{L_{y_i}} \equiv (Y_{i,t-L_{y_i}}, Y_{i,t-L_{y_i}+1}, \dots, Y_{i,t-1})$, $L_{y_i} = 1, 2, \dots$, $t = L_{y_i} + 1, L_{y_i} + 2, \dots$. Denote $M_x = (m_{x_1}, \dots, m_{x_{n_1}})$, $m_x = \max(m_{x_1}, \dots, m_{x_{n_1}})$, $L_x = (L_{x_1}, \dots, L_{x_{n_1}})$, $l_x = \max(L_{x_1}, \dots, L_{x_{n_1}})$, $L_y = (L_{y_1}, \dots, L_{y_{n_2}})$, $l_y = \max(L_{y_1}, \dots, L_{y_{n_2}})$. For given M_x, L_x, L_y, e ,

Bai et al. (2010) define that

$$\begin{aligned} \{\|X_t^{M_x} - X_s^{M_x}\| < e\} &\equiv \{\|X_{i,t}^{m_{x_i}} - X_{i,s}^{m_{x_i}}\| < e, \quad \text{for all } i = 1, 2, \dots, n_1\} \\ \{\|X_{t-L_x}^{L_x} - X_{s-L_x}^{L_x}\| < e\} &\equiv \{\|X_{i,t-L_x}^{L_{x_i}} - X_{i,s-L_x}^{L_{x_i}}\| < e, \quad \text{for all } i = 1, 2, \dots, n_1\} \\ \{\|Y_{t-L_y}^{L_y} - Y_{s-L_y}^{L_y}\| < e\} &\equiv \{\|Y_{i,t-L_y}^{L_{y_i}} - Y_{i,s-L_y}^{L_{y_i}}\| < e, \quad \text{for all } i = 1, 2, \dots, n_2\}, \end{aligned}$$

where $\|\cdot\|$ denotes the maximum norm defined as $\|X - Y\| = \max(|x_1 - y_1|, |x_2 - y_2|, \dots, |x_n - y_n|)$ for any two vectors $X = (x_1, \dots, x_n)$ and $Y = (y_1, \dots, y_n)$.

Definition 2.1. *The vector time series $\{Y_t\}$ does not strictly Granger cause another vector time series $\{X_t\}$ if*

$$\begin{aligned} P\left(\|X_t^{M_x} - X_s^{M_x}\| < e \mid \|X_{t-L_x}^{L_x} - X_{s-L_x}^{L_x}\| < e, \|Y_{t-L_y}^{L_y} - Y_{s-L_y}^{L_y}\| < e\right) \\ = P\left(\|X_t^{M_x} - X_s^{M_x}\| < e \mid \|X_{t-L_x}^{L_x} - X_{s-L_x}^{L_x}\| < e\right), \end{aligned} \quad (1)$$

where $P(\cdot|\cdot)$ denotes conditional probability.

Using the notation

$$\begin{aligned} C_1(M_x + L_x, L_y, e) &\equiv P\left(\|X_{t-L_x}^{M_x+L_x} - X_{s-L_x}^{M_x+L_x}\| < e, \|Y_{t-L_y}^{L_y} - Y_{s-L_y}^{L_y}\| < e\right) \\ C_2(L_x, L_y, e) &\equiv P\left(\|X_{t-L_x}^{L_x} - X_{s-L_x}^{L_x}\| < e, \|Y_{t-L_y}^{L_y} - Y_{s-L_y}^{L_y}\| < e\right) \\ C_3(M_x + L_x, e) &\equiv P\left(\|X_{t-L_x}^{M_x+L_x} - X_{s-L_x}^{M_x+L_x}\| < e\right) \\ C_4(L_x, e) &\equiv P\left(\|X_{t-L_x}^{L_x} - X_{s-L_x}^{L_x}\| < e\right), \end{aligned}$$

Bai, et al. (2010) re-express Equation (1) as

$$\frac{C_1(M_x + L_x, L_y, e)}{C_2(L_x, L_y, e)} = \frac{C_3(M_x + L_x, e)}{C_4(L_x, e)}. \quad (2)$$

For two sets of simultaneous samples $\{x_{i,t}, i = 1, \dots, n_1, t = 1, \dots, T\}$ and $\{y_{i,t}, i =$

$1, \dots, n_2, t = 1, \dots, T\}$, they propose the following test statistic

$$\sqrt{n} \left(\frac{C_1(M_x + L_x, L_y, e, n)}{C_2(L_x, L_y, e, n)} - \frac{C_3(M_x + L_x, e, n)}{C_4(L_x, e, n)} \right), \quad (3)$$

where

$$\begin{aligned} C_1(M_x + L_x, L_y, e, n) &\equiv \frac{2}{n(n-1)} \sum_{t < s} \sum_{i=1}^{n_1} I(x_{i,t-L_{x_i}}^{m_{x_i}+L_{x_i}}, x_{i,s-L_{x_i}}^{m_{x_i}+L_{x_i}}, e) \cdot \prod_{i=1}^{n_2} I(y_{i,t-L_{y_i}}^{L_{y_i}}, y_{i,s-L_{y_i}}^{L_{y_i}}, e), \\ C_2(L_x, L_y, e, n) &\equiv \frac{2}{n(n-1)} \sum_{t < s} \sum_{i=1}^{n_1} I(x_{i,t-L_{x_i}}^{L_{x_i}}, x_{i,s-L_{x_i}}^{L_{x_i}}, e) \cdot \prod_{i=1}^{n_2} I(y_{i,t-L_{y_i}}^{L_{y_i}}, y_{i,s-L_{y_i}}^{L_{y_i}}, e), \\ C_3(M_x + L_x, e, n) &\equiv \frac{2}{n(n-1)} \sum_{t < s} \sum_{i=1}^{n_1} I(x_{i,t-L_{x_i}}^{m_{x_i}+L_{x_i}}, x_{i,s-L_{x_i}}^{m_{x_i}+L_{x_i}}, e), \\ C_4(L_x, e, n) &\equiv \frac{2}{n(n-1)} \sum_{t < s} \sum_{i=1}^{n_1} I(x_{i,t-L_{x_i}}^{L_{x_i}}, x_{i,s-L_{x_i}}^{L_{x_i}}, e), \end{aligned}$$

and

$$I(x, y, e) = \begin{cases} 0, & \text{if } \|x - y\| > e \\ 1, & \text{if } \|x - y\| \leq e \end{cases}.$$

Remark: Following the instruction of Hiemstra and Jones (1994), Bai et al. (2010) take $C_j(*, n)$ s as multivariate U -statistic estimators of their counterparts $C_j(*)$ s and apply the asymptotic property of U -statistic to show the limiting results for the test statistics (3). However the $C_j(*, n)$ s are not U -statistics, because the expectations of the general terms are not the same. Moreover, the $C_j(*)$ s are related to the indices t and s (in fact, related to $|t - s|$ for strongly stationary processes), while the $C_j(*, n)$ s were independent of t and s for summing up over them. Therefore, the $C_j(*, n)$ estimators are neither consistent nor asymptotic normal estimators of their counterparts $C_j(*)$.

3 A New Multivariate Nonlinear Causality Test

We first remind the reader that the pair (s, t) (in fact, $|t - s|$ for strongly stationary processes) in Equation (1) of Definition 2.1 is a key parameter of the probabilities $C_j(*)$. In fact, both Hiemstra and Jones (1994) and Bai et al. (2010) note this, and there is no problem in Equation (1) of Definition 2.1. However, it seems that they overlooked this fact in their proposed estimation of $C_j(*)$. The improper estimators $C_j(*, n)$ thus lead to an invalid asymptotic distribution of the test statistic.

We now begin to state the procedure for our new test. For any given pair (s, t) , we denote

$$\begin{aligned} C_1(M_x + L_x, L_y, e; t, s) &\equiv P \left(\| X_{t-L_x}^{M_x+L_x} - X_{s-L_x}^{M_x+L_x} \| < e, \| Y_{t-L_y}^{L_y} - Y_{s-L_y}^{L_y} \| < e \right) \\ C_2(L_x, L_y, e; t, s) &\equiv P \left(\| X_{t-L_x}^{L_x} - X_{s-L_x}^{L_x} \| < e, \| Y_{t-L_y}^{L_y} - Y_{s-L_y}^{L_y} \| < e \right) \\ C_3(M_x + L_x, e; t, s) &\equiv P \left(\| X_{t-L_x}^{M_x+L_x} - X_{s-L_x}^{M_x+L_x} \| < e \right) \\ C_4(L_x, e; t, s) &\equiv P \left(\| X_{t-L_x}^{L_x} - X_{s-L_x}^{L_x} \| < e \right) \end{aligned}$$

Under the assumption of the stationary, for the given pair (s, t) , if $s - t = l$, we denote $C_1(M_x + L_x, L_y, e; t, s) \equiv C_1(M_x + L_x, L_y, e; t, l)$, which does not depend on t , so we can write $C_1(M_x + L_x, L_y, e; l)$ instead of $C_1(M_x + L_x, L_y, e, t; l)$, the same to the others. So under the assumption of strictly stationary, for each $l > 0$, we examine whether there is nonlinear Granger causality from $\{Y_t\}$ to $\{X_t\}$ by testing the following hypothesis

$$H_0 : \frac{C_1(M_x + L_x, L_y, e; l)}{C_2(L_x, L_y, e; l)} = \frac{C_3(M_x + L_x, e; l)}{C_4(L_x, e; l)}. \quad (4)$$

If we consider two sets of simultaneous samples¹ $\{x_{i,t}, i = 1, \dots, n_1, t = 1, \dots, T\}$ and $\{y_{j,t}, j = 1, \dots, n_2, t = 1, \dots, T\}$, we first provide the consistent estimators of $C_1(M_x +$

¹To implement the test, each series is standardized so that all series share a common standard deviation, and thereby share a common scale parameter.

$L_x, L_y, e; l)$, $C_2(L_x, L_y, e; l)$, $C_3(M_x + L_x, e; l)$ and $C_4(L_x, e; l)$ are

$$\begin{aligned}\hat{C}_1(M_x + L_x, L_y, e; l) &\equiv \frac{1}{n} \sum_{t=L_{xy}+1}^{T-l-m_x+1} \prod_{i=1}^{n_1} I\left(x_{i,t-L_{x_i}}^{m_{x_i}+L_{x_i}}, x_{i,t+l-L_{x_i}}^{m_{x_i}+L_{x_i}}, e\right) \cdot \prod_{j=1}^{n_2} I\left(y_{j,t-L_{y_j}}^{L_{y_j}}, y_{j,t+l-L_{y_j}}^{L_{y_j}}, e\right) \\ \hat{C}_2(L_x, L_y, e; l) &\equiv \frac{1}{n} \sum_{t=L_{xy}+1}^{T-l-m_x+1} \prod_{i=1}^{n_1} I\left(x_{i,t-L_{x_i}}^{L_{x_i}}, x_{i,t+l-L_{x_i}}^{L_{x_i}}, e\right) \cdot \prod_{j=1}^{n_2} I\left(y_{j,t-L_{y_j}}^{L_{y_j}}, y_{j,t+l-L_{y_j}}^{L_{y_j}}, e\right) \\ \hat{C}_3(M_x + L_y, e; l) &\equiv \frac{1}{n} \sum_{t=L_{xy}+1}^{T-l-m_x+1} \prod_{i=1}^{n_1} I\left(x_{i,t-L_{x_i}}^{m_{x_i}+L_{x_i}}, x_{i,t+l-L_{x_i}}^{m_{x_i}+L_{x_i}}, e\right) \\ \hat{C}_4(L_x, e; l) &\equiv \frac{1}{n} \sum_{t=L_{xy}+1}^{T-l-m_x+1} \prod_{i=1}^{n_1} I\left(x_{i,t-L_{x_i}}^{L_{x_i}}, x_{i,t+l-L_{x_i}}^{L_{x_i}}, e\right)\end{aligned}$$

where $L_{xy} = \max(L_x, L_y)$, $n = T - L_{xy} - l - m_x + 1$ and

$$I(x, y, e) \equiv \begin{cases} 0, & \text{if } \|x - y\| > e \\ 1, & \text{if } \|x - y\| \leq e \end{cases}.$$

The consistency of our proposed estimators can be shown straightforwardly and the detail of the proof is omitted. We use a simple numerical study to show that our estimators are consistent whereas those of Bai et al. (2010) are not. Let

$$\begin{pmatrix} X_{1,t} \\ X_{2,t} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a_{1,t-1} \\ a_{2,t-1} \end{pmatrix} + \begin{pmatrix} a_{1,t} \\ a_{2,t} \end{pmatrix},$$

where $\{a_{1,t}\}, \{a_{2,t}\}$ are i.i.d. and mutually independent random variables generated from $N(0, 1)$, while $\{Y_t\}$ could be any stationary sequence. Let $l = 1$, $L_x = L_y = M_x = 1$. We can calculate the exact values of $C_4(L_x, e; l)$, which are 0.2709 and 0.5057, respectively, when $e = 1$ and $e = 1.5$. For simplicity, we denote the true value of $C_4(L_x, e; l)$, the estimate proposed by Bai et al.(2010) and our new estimate as C_4 , \hat{C}_4^{BWZ} and \hat{C}_4 , respectively, in Table 1. Additionally, Table 1 provides the estimated values with their corresponding relative estimation errors in brackets when $T = 1000, 2000$ and 4000 . It is

obvious that \hat{C}_4^{BWZ} is not consistent.

Table 1: $C_4(L_x, e; l)$ and estimated values.

$T =$	$e = 1$			$e = 1.5$		
	C_4	\hat{C}_4	\hat{C}_4^{BWZ}	C_4	\hat{C}_4	\hat{C}_4^{BWZ}
1000	0.2709	0.2497(7.83%)	0.0154(94.32%)	0.5057	0.4774(5.60%)	0.0732(85.53%)
2000	0.2709	0.2639(2.58%)	0.0176(93.50%)	0.5057	0.4847(4.15%)	0.0795(84.28%)
4000	0.2709	0.2692(0.63%)	0.0193(92.88%)	0.5057	0.4909(2.93%)	0.0820(83.78%)

Note: The true value of $C_4(L_x, e; l)$ is denoted C_4 , the BWZ estimate and our new estimate are denoted \hat{C}_4^{BWZ} and \hat{C}_4 , respectively. The relative estimation errors are in the accompanying brackets.

Now, we propose

$$T_n = \sqrt{n} \left(\frac{\hat{C}_1(M_x + L_x, L_y, e, l)}{\hat{C}_2(L_x, L_y, e, l)} - \frac{\hat{C}_3(M_x + L_x, e, l)}{\hat{C}_4(L_x, e, l)} \right) \quad (5)$$

as the test statistic, and we derive the following asymptotic distribution of T_n for the Granger causality test.

Theorem 3.1. *Stationary sequences $\{x_{i,t}, i = 1, \dots, n_1, t = 1, \dots, T\}$ and $\{y_{j,t}, j = 1, \dots, n_2, t = 1, \dots, T\}$ are strong mixing, with mixing coefficients satisfying the conditions of Lemma 1 presented in Appendix, for given values of l, L_x, L_y, M_x and $e > 0$, under the null hypothesis that $\{Y_t\}$ does not strictly Granger cause $\{X_t\}$, then the test statistic is defined in (5)*

$$\sqrt{n} \left(\frac{\hat{C}_1(M_x + L_x, L_y, e, l)}{\hat{C}_2(L_x, L_y, e, l)} - \frac{\hat{C}_3(M_x + L_x, e, l)}{\hat{C}_4(L_x, e, l)} \right) \xrightarrow{d} N(0, \sigma^2(M_x, L_x, L_y, e, l)).$$

The asymptotic variance $\sigma^2(M_x, L_x, L_y, e, l)$ with its consistent estimator $\hat{\sigma}^2(M_x, L_x, L_y, e, l)$ and the proof of theorem 3.1 are given in the Appendix. The hypothesis H_0 defined in (4) is rejected at α if

$$|T_n| / \hat{\sigma}(M_x, L_x, L_y, e, l) > z_{\alpha/2},$$

where $z_{\alpha/2}$ is the up $\alpha/2$ quantile of the standard normal distribution. In this situation, we will conclude that there exists nonlinear Granger causality from $\{Y_t\}$ to $\{X_t\}$.

There are several possible methods to estimate the asymptotic covariance $\sigma^2(M_x, L_x, L_y, e, l)$. A model-based approach uses known laws of $\{X_t\}$ and $\{Y_t\}$ to calculate the expectations in the formula given in the Appendix and simply substitutes $C_j(*), j = 1, 2, 3, 4$ with their corresponding estimates. However, in practice, we can hardly avoid model misspecification and may obtain improper laws of $\{X_t\}$ and $\{Y_t\}$. We suggest the use of bootstrap methods as in the simulation studies we use to test hypothesis H_0 .

4 Simulation

In this section, we perform numerical studies using simulations to illustrate the applicability and superiority of the new multivariate nonlinear Granger causality test developed in Section 3. Let R be the times of rejecting the null hypothesis that Y_t does not strictly Granger cause X_t nonlinearly in 10,000 replications at the α level, and thus, the empirical power is $R/10,000$. In our simulation, the level $\alpha = 0.05$, we standardized the series and chose the same lag length and lead length: $L_x = L_y = M_x = 1$. We set three situations of l and two situations of e : $l = 1, l = 2, l = 3$ and $e = 1, e = 1.5$.

Consider the following model:

$$X_t = \beta Y_{1,t-1} Y_{2,t-1} + \varepsilon_t, \quad (6)$$

where $\{(Y_{1,t}, Y_{2,t})'\}$ are i.i.d. and mutually independent random variables generated from standard normal distribution $N(0, 1)$, $\{\varepsilon_t\}$ is Gaussian white noise generated from $N(0, 0.1)$ and independent of $\{Y_{1,t}\}, \{Y_{2,t}\}$. There is no nonlinear Granger causality from Y_t to X_t when $\beta = 0$, and causality strengthens when β increases.

From the results displayed in Table 2, we conclude first that our test possesses decent size, as we can see when $\beta = 0$ the empirical size are all closed to the test level 0.05 for different settings of parameters and sample size. Second, our test possesses very appropriate power, as we see that empirical power increases as β increases, especially when sample size is 500 the empirical power sharply increase to 1. Further, we find that

Table 2: Test multivariate nonlinear Granger causality form Y_t to X_t

$T = 200$	$e = 1$			$e = 1.5$		
	$l = 1$	$l = 2$	$l = 3$	$l = 1$	$l = 2$	$l = 3$
$\beta = 0$	0.0419	0.0441	0.0432	0.0444	0.0506	0.0438
$\beta = 0.1$	0.1509	0.2341	0.2184	0.3647	0.5209	0.5121
$\beta = 0.2$	0.5629	0.7425	0.7284	0.8352	0.9416	0.9416
$\beta = 0.3$	0.8178	0.9323	0.9267	0.929	0.9810	0.9825
$\beta = 0.4$	0.8994	0.9712	0.9719	0.9512	0.9878	0.9889
$\beta = 0.5$	0.9366	0.9812	0.9808	0.9586	0.9914	0.9915
$\beta = 0.6$	0.9475	0.9870	0.9862	0.9640	0.9918	0.9940
$\beta = 0.7$	0.9574	0.9875	0.9874	0.9664	0.9921	0.9942
$\beta = 0.8$	0.9615	0.9888	0.9882	0.9688	0.9926	0.9953
$\beta = 0.9$	0.9633	0.9896	0.9901	0.9711	0.9923	0.994

$T = 500$	$e = 1$			$e = 1.5$		
	$l = 1$	$l = 2$	$l = 3$	$l = 1$	$l = 2$	$l = 3$
$\beta = 0$	0.0560	0.0560	0.0478	0.0501	0.0529	0.0423
$\beta = 0.1$	0.3969	0.5081	0.5344	0.7835	0.9043	0.9017
$\beta = 0.2$	0.9622	0.9908	0.9900	0.9994	1.0000	1.0000
$\beta = 0.3$	0.9989	1.0000	0.9997	0.9999	1.0000	1.0000
$\beta = 0.4$	0.9998	1.0000	1.0000	1.0000	1.0000	1.0000
$\beta = 0.5$	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
$\beta = 0.6$	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
$\beta = 0.7$	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
$\beta = 0.8$	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
$\beta = 0.9$	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

Note: Y_t and X_t are from the model present in equation (6). $L_x = L_y = M_x = 1$ in our test. Simulation is conducted with the test level $\alpha = 5\%$, and 10,000 replications.

different settings of e may influence the test results. Though the influence is little in our simulation, we still suggest that practitioners choose a couple of different values of e .

5 Conclusion and Remarks

In this paper, we reinvestigate the multivariate nonlinear Granger causality test extended by Bai et al. (2010) which attempt to uncover significant nonlinearities in the dynamic interrelationships between two groups of variables. We find that Bai et al. (2010) as well as Hiemstra and Jones (1994) take the estimators of the probabilities in their definition as U -statistics and establish a CLT of the test statistic by applying the asymptotic property of U -statistics. After revealing that the estimators proposed by Bai et al. (2010) is not U -statistics, we show that their estimators are also not consistent.

The procedure of our new test begins with presenting consistent estimators of probabilities in the definition. Numerical study supports that our estimators are consistent, further our new test possesses admirable properties both in size and power.

There are still amounts of appealing aspects in nonlinear Granger causality test. It is worth noting that Diks and Wolski (2015) extend the test in Diks and Panchenko (2006) which highlight a need for substitutions for the relationship tested in the HJ test.

Appendix

A1: Central Limit Theorems for strong mixing stationary sequence

$\{(Z_t, \mathcal{F}_t), -\infty < t < \infty\}$ is a stochastic process defined on the probability space (Ω, \mathcal{F}, P) . The history and the future of Z_t are σ -algebras $\mathfrak{M}_t^\infty = \{\mathcal{F}_s, s > t\}$ and σ -algebras $\mathfrak{M}_{-\infty}^t = \{\mathcal{F}_s, s < t\}$ respectively.

Let $\{(Z_i, \mathcal{F}_i)\}$ be a stationary sequence with $E(Z_i) = 0$, $E(Z_i^2) < \infty$, and set $S_n^m = \sum_{i=m+1}^{n+m} Z_i$, $\sigma_n^2 = \text{Var}(S_n^m)$. We shall say that the sequence satisfies the central limit theorem if

$$\lim_{n \rightarrow \infty} P\left\{\frac{S_n^m}{\sigma_n} < z\right\} = (2\pi)^{-\frac{1}{2}} \int_{-\infty}^z e^{-\frac{1}{2}u^2} du = \Phi(z) .$$

Definition A1: A stationary process $\{Z_t\}$ is said to be strongly mixing (completely regular) if $\alpha(\tau) = \sup_{A \in \mathfrak{m}_{-\infty}^0, B \in \mathfrak{m}_{\tau}^{\infty}} |P(AB) - P(A)P(B)| \rightarrow 0$ as $\tau \rightarrow \infty$ through positive values.

Lemma A1: Let the stationary sequence $\{Z_i\}$ satisfy the strong mixing condition with mixing coefficient $\alpha(n)$, and let $E|Z_i|^{2+\delta} < \infty$ for some $\delta > 0$. If $\sum_{n=1}^{\infty} \alpha(n)^{\delta/(2+\delta)} < \infty$, then

$$\sigma^2 = E(Z_0^2) + 2 \sum_{j=1}^{\infty} E(Z_0 Z_j) < \infty, \text{ and if } \sigma \neq 0, \text{ then } \lim_{n \rightarrow \infty} P\{\sigma^{-1} n^{-\frac{1}{2}} \sum_{i=1}^n Z_i < z\} = \Phi(z).$$

Readers can be referred to Ibragimov (1971) for a proof and detailed discussion.

A2: Proof of Theorem 3.1

Assume $\{x_{i,1}, x_{i,2}, \dots, x_{i,T}\}$ and $\{y_{j,1}, y_{j,2}, \dots, y_{j,T}\}, i \in \{1, 2, \dots, n_1\}, j \in \{1, 2, \dots, n_2\}$ are both strong mixing stationary sequences whose mixing coefficient satisfying the conditions in Lemma 1. Then the following four sequences

$$\begin{aligned} \{Z_{1t} &= n^{-1} \left(\prod_{i=1}^{n_1} I(x_{i,t-L_{x_i}}^{m_{x_i}+L_{x_i}}, x_{i,t+l-L_{x_i}}^{m_{x_i}+L_{x_i}}, e) \cdot \prod_{j=1}^{n_1} I(y_{j,t-L_{y_j}}^{L_{y_j}}, y_{j,t+l-L_{y_j}}^{L_{y_j}}, e) - C_1(M_x + L_x, L_y, e; l) \right)\}, \\ \{Z_{2t} &= n^{-1} \left(\prod_{i=1}^{n_1} I(x_{i,t-L_{x_i}}^{L_{x_i}}, x_{i,t+l-L_{x_i}}^{L_{x_i}}, e) \cdot \prod_{j=1}^{n_1} I(y_{j,t-L_{y_j}}^{L_{y_j}}, y_{j,t+l-L_{y_j}}^{L_{y_j}}, e) - C_2(L_x, L_y, e; l) \right)\}, \\ \{Z_{3t} &= n^{-1} \left(\prod_{i=1}^{n_1} I(x_{i,t-L_{x_i}}^{L_{x_i}}, x_{i,t+l-L_{x_i}}^{L_{x_i}}, e) \cdot \prod_{j=1}^{n_1} I(y_{j,t-L_{y_j}}^{L_{y_j}}, y_{j,t+l-L_{y_j}}^{L_{y_j}}, e) - C_2(L_x, L_y, e; l) \right)\}, \\ \{Z_{4t} &= n^{-1} \left(\prod_{i=1}^{n_1} I(x_{i,t-L_{x_i}}^{L_{x_i}}, x_{i,t+l-L_{x_i}}^{L_{x_i}}, e) - C_4(L_x, e; l) \right)\}, \\ t &= L_{xy} + 1, \dots, T - l - L_{xy} - m + 1, \end{aligned}$$

satisfy the conditions of Lemma 1, where $n = T - L_{xy} - l - m_x + 1$ and

$$\begin{aligned} C_1(M_x + L_x, L_y, e; t, s) &\equiv P \left(\| X_{t-L_x}^{M_x+L_x} - X_{s-L_x}^{M_x+L_x} \| < e, \| Y_{t-L_y}^{L_y} - Y_{s-L_y}^{L_y} \| < e \right), \\ C_2(L_x, L_y, e; t, s) &\equiv P \left(\| X_{t-L_x}^{L_x} - X_{s-L_x}^{L_x} \| < e, \| Y_{t-L_y}^{L_y} - Y_{s-L_y}^{L_y} \| < e \right) \\ C_3(M_x + L_x, e; t, s) &\equiv P \left(\| X_{t-L_x}^{M_x+L_x} - X_{s-L_x}^{M_x+L_x} \| < e \right) \\ C_4(L_x, e; t, s) &\equiv P \left(\| X_{t-L_x}^{L_x} - X_{s-L_x}^{L_x} \| < e \right). \end{aligned}$$

So $\{Z_{1t}\}$, $\{Z_{2t}\}$, $\{Z_{3t}\}$ and $\{Z_{4t}\}$ satisfy the central limit theorem.

Further, for any real number a_1, a_2, a_3 and a_4 , the sequence $\{Z_t = a_1 Z_{1t} + a_2 Z_{2t} + a_3 Z_{3t} + a_4 Z_{4t}, t = L_{xy}, \dots, T - l - L_{xy} - m + 1\}$ also satisfies the conditions of Lemma 1 which implying that

$$\sqrt{n} \begin{bmatrix} \hat{C}_1(m + L_x, L_y, e; l) - C_1(m + L_x, L_y, e; l) \\ \hat{C}_2(L_x, L_y, e; l) - C_2(L_x, L_y, e; l) \\ \hat{C}_3(m + L_x, e; l) - C_3(m + L_x, e; l) \\ \hat{C}_4(L_x, e; l) - C_4(L_x, e; l) \end{bmatrix} \xrightarrow{d} N(0, \Sigma),$$

where

$$\begin{aligned} \hat{C}_1(M_x + L_x, L_y, e; l) &\equiv \frac{1}{n} \sum_{t=L_{xy}+1}^{T-l-m_x+1} \prod_{i=1}^{n_1} I\left(x_{i,t-L_{x_i}}^{m_{x_i}+L_{x_i}}, x_{i,t+l-L_{x_i}}^{m_{x_i}+L_{x_i}}, e\right) \cdot \prod_{j=1}^{n_2} I\left(y_{j,t-L_{y_j}}^{L_{y_j}}, y_{j,t+l-L_{y_j}}^{L_{y_j}}, e\right) \\ \hat{C}_2(L_x, L_y, e; l) &\equiv \frac{1}{n} \sum_{t=L_{xy}+1}^{T-l-m_x+1} \prod_{i=1}^{n_1} I\left(x_{i,t-L_{x_i}}^{L_{x_i}}, x_{i,t+l-L_{x_i}}^{L_{x_i}}, e\right) \cdot \prod_{j=1}^{n_2} I\left(y_{j,t-L_{y_j}}^{L_{y_j}}, y_{j,t+l-L_{y_j}}^{L_{y_j}}, e\right) \\ \hat{C}_3(M_x + L_y, e; l) &\equiv \frac{1}{n} \sum_{t=L_{xy}+1}^{T-l-m_x+1} \prod_{i=1}^{n_1} I\left(x_{i,t-L_{x_i}}^{m_{x_i}+L_{x_i}}, x_{i,t+l-L_{x_i}}^{m_{x_i}+L_{x_i}}, e\right) \\ \hat{C}_4(L_x, e; l) &\equiv \frac{1}{n} \sum_{t=L_{xy}+1}^{T-l-m_x+1} \prod_{i=1}^{n_1} I\left(x_{i,t-L_{x_i}}^{L_{x_i}}, x_{i,t+l-L_{x_i}}^{L_{x_i}}, e\right) \end{aligned}$$

and Σ is a 4×4 symmetric matrix. Denote

$$h_1(L_x, L_y, M_x, l, k) = I(x_{L_{xy}+1+k-L_x}^{L_x+M_x}, x_{L_{xy}+1+k+l-L_x}^{L_x+M_x}, e),$$

$$h_2(L_x, L_y, l, k) = I(y_{L_{xy}+1+k-L_y}^{L_y}, y_{L_{xy}+1+k+l-L_y}^{L_y}, e),$$

We have

$$\begin{aligned}
\Sigma_{11} &= E \left[\left(h_1(L_x, L_y, M_x, l, 0) h_2(L_x, L_y, l, 0) - C_1(M_x + L_x, L_y, e; l) \right)^2 \right] \\
&+ \sum_{k=1}^{n-1} 2 \left(1 - \frac{k}{n} \right) E \left[\left(h_1(L_x, L_y, M_x, l, 0) h_2(L_x, L_y, l, 0) - C_1(M_x + L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, M_x, l, k) h_2(L_x, L_y, l, k) - C_1(M_x + L_x, L_y, e; l) \right) \right], \\
\Sigma_{12} &= E \left[\left(h_1(L_x, L_y, M_x, l, 0) h_2(L_x, L_y, l, 0) - C_1(M_x + L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, 0, l, 0) h_2(L_x, L_y, l, 0) - C_2(L_x, L_y, e; l) \right) \right] \\
&+ \sum_{k=1}^{n-1} \left(1 - \frac{k}{n} \right) E \left[\left(h_1(L_x, L_y, M_x, l, 0) h_2(L_x, L_y, l, 0) - C_1(M_x + L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, 0, l, k) h_2(L_x, L_y, l, k) - C_2(L_x, L_y, e; l) \right) \right] \\
&+ \sum_{k=1}^{n-1} \left(1 - \frac{k}{n} \right) E \left[\left(h_1(L_x, L_y, M_x, l, k) h_2(L_x, L_y, l, k) - C_1(M_x + L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, 0, l, 0) h_2(L_x, L_y, l, 0) - C_2(L_x, L_y, e; l) \right) \right], \\
\Sigma_{13} &= E \left[\left(h_1(L_x, L_y, M_x, l, 0) h_2(L_x, L_y, l, 0) - C_1(M_x + L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, M_x, l, 0) - C_3(M_x + L_x, e; l) \right) \right] \\
&+ \sum_{k=1}^{n-1} \left(1 - \frac{k}{n} \right) E \left[\left(h_1(L_x, L_y, M_x, l, 0) h_2(L_x, L_y, l, 0) - C_1(M_x + L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, M_x, l, k) - C_3(M_x + L_x, e; l) \right) \right] \\
&+ \sum_{k=1}^{n-1} \left(1 - \frac{k}{n} \right) E \left[\left(h_1(L_x, L_y, M_x, l, k) h_2(L_x, L_y, l, k) - C_1(M_x + L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, M_x, l, 0) - C_3(M_x + L_x, e; l) \right) \right],
\end{aligned}$$

$$\begin{aligned}
\Sigma_{14} &= E \left[\left(h_1(L_x, L_y, M_x, l, 0) h_2(L_x, L_y, l, 0) - C_1(M_x + L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, 0, l, 0) - C_4(L_x, e; l) \right) \right] \\
&+ \sum_{k=1}^{n-1} \left(1 - \frac{k}{n} \right) E \left[\left(h_1(L_x, L_y, M_x, l, 0) h_2(L_x, L_y, l, 0) - C_1(M_x + L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, 0, l, k) - C_4(L_x, e; l) \right) \right] \\
&+ \sum_{k=1}^{n-1} \left(1 - \frac{k}{n} \right) E \left[\left(h_1(L_x, L_y, M_x, l, k) h_2(L_x, L_y, l, k) - C_1(M_x + L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, 0, l, 0) - C_4(L_x, e; l) \right) \right] , \\
\Sigma_{22} &= E \left[\left(h_1(L_x, L_y, 0, l, 0) h_2(L_x, L_y, l, 0) - C_2(L_x, L_y, e; l) \right)^2 \right] \\
&+ \sum_{k=1}^{n-1} 2 \left(1 - \frac{k}{n} \right) E \left[\left(h_1(L_x, L_y, 0, l, 0) h_2(L_x, L_y, l, 0) - C_2(L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, 0, l, k) h_2(L_x, L_y, l, k) - C_2(L_x, L_y, e; l) \right) \right] , \\
\Sigma_{23} &= E \left[\left(h_1(L_x, L_y, 0, l, 0) h_2(L_x, L_y, l, 0) - C_2(L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, M_x, l, 0) - C_3(M_x + L_x, e; l) \right) \right] \\
&+ \sum_{k=1}^{n-1} \left(1 - \frac{k}{n} \right) E \left[\left(h_1(L_x, L_y, 0, l, 0) h_2(L_x, L_y, l, 0) - C_2(L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, M_x, l, k) - C_3(M_x + L_x, e; l) \right) \right] \\
&+ \sum_{k=1}^{n-1} \left(1 - \frac{k}{n} \right) E \left[\left(h_1(L_x, L_y, 0, l, k) h_2(L_x, L_y, l, k) - C_2(L_x, L_y, e; l) \right) \right. \\
&\quad \left. \left(h_1(L_x, L_y, M_x, l, 0) - C_3(M_x + L_x, e; l) \right) \right] ,
\end{aligned}$$

$$\begin{aligned}
\Sigma_{24} &= E \left[(h_1(L_x, L_y, 0, l, 0)h_2(L_x, L_y, l, 0) - C_2(L_x, L_y, e; l)) \right. \\
&\quad \left. (h_1(L_x, L_y, 0, l, 0) - C_4(L_x, e; l)) \right] \\
&+ \sum_{k=1}^{n-1} \left(1 - \frac{k}{n}\right) E \left[(h_1(L_x, L_y, 0, l, 0)h_2(L_x, L_y, l, 0) - C_2(L_x, L_y, e; l)) \right. \\
&\quad \left. (h_1(L_x, L_y, 0, l, k) - C_4(L_x, e; l)) \right] \\
&+ \sum_{k=1}^{n-1} \left(1 - \frac{k}{n}\right) E \left[(h_1(L_x, L_y, 0, l, k)h_2(L_x, L_y, l, k) - C_2(L_x, L_y, e; l)) \right. \\
&\quad \left. (h_1(L_x, L_y, 0, l, 0) - C_4(L_x, e; l)) \right] , \\
\Sigma_{33} &= E \left[(h_1(L_x, L_y, M_x, l, 0) - C_3(M_x + L_x, e; l))^2 \right] \\
&+ \sum_{k=1}^{n-1} 2 \left(1 - \frac{k}{n}\right) E \left[(h_1(L_x, L_y, M_x, l, 0) - C_3(M_x + L_x, e; l)) \right. \\
&\quad \left. (h_1(L_x, L_y, M_x, l, k) - C_3(M_x + L_x, e; l)) \right] , \\
\Sigma_{34} &= E \left[(h_1(L_x, L_y, M_x, l, 0) - C_3(M_x + L_x, e; l)) \right. \\
&\quad \left. (h_1(L_x, L_y, 0, l, 0) - C_4(L_x, e; l)) \right] \\
&+ \sum_{k=1}^{n-1} \left(1 - \frac{k}{n}\right) E \left[(h_1(L_x, L_y, M_x, l, 0) - C_3(M_x + L_x, e; l)) \right. \\
&\quad \left. (h_1(L_x, L_y, 0, l, k) - C_4(L_x, e; l)) \right] \\
&+ \sum_{k=1}^{n-1} \left(1 - \frac{k}{n}\right) E \left[(h_1(L_x, L_y, M_x, l, k) - C_3(M_x + L_x, e; l)) \right. \\
&\quad \left. (h_1(L_x, L_y, 0, l, 0) - C_4(L_x, e; l)) \right] , \\
\Sigma_{44} &= E \left[(h_1(L_x, L_y, 0, l, 0) - C_4(L_x, e; l))^2 \right] \\
&+ \sum_{k=1}^{n-1} 2 \left(1 - \frac{k}{n}\right) E \left[(h_1(L_x, L_y, 0, l, 0) - C_4(L_x, e; l)) \right. \\
&\quad \left. (h_1(L_x, L_y, 0, l, k) - C_4(L_x, e; l)) \right] .
\end{aligned}$$

Under the null hypothesis, applying the delta method (Serfling, 1980), we have

$$\sqrt{n} \left(\frac{\hat{C}_1(M_x + L_x, L_y, e, l)}{\hat{C}_2(L_x, L_y, e, l)} - \frac{\hat{C}_3(M_x + L_x, e, l)}{\hat{C}_4(L_x, e, l)} \right) \xrightarrow{d} N(0, \sigma^2(M_x, L_x, L_y, e, l)),$$

where $\sigma^2(M_x, L_x, L_y, e, l) = \nabla' \Sigma \nabla$, in which

$$\nabla = \left(\frac{1}{C_2(L_x, L_y, e, l)}, -\frac{C_1(M_x + L_x, L_y, e, l)}{C_2^2(L_x, L_y, e, l)}, -\frac{1}{C_4(L_x, e, l)}, \frac{C_3(M_x + L_x, e, l)}{C_4^2(L_x, e, l)} \right)'.$$

An consistent estimator $\hat{\sigma}^2(M_x, L_x, L_y, e, l)$ of the asymptotic variance can be got by replacing all the parts in the sandwich $\nabla' \Sigma \nabla$ by their empirical estimates.

References

- [1] E.G. Baek, W.A. Brok, A general test for nonlinear Granger causality: bivariate model, working paper, Korea Development Institute, University of Wisconsin Madison (1992)
- [2] Z.D. Bai, W.K. Wong, B.Z. Zhang, Multivariate linear and nonlinear causality tests, *Mathematics and Computers in simulation* 81 (2010) 5-17.
- [3] Z.D. Bai, Y.C. Hui, Z.H. Lv, W.K. Wong, Z.Z. Zhu, The Hiemstra-Jones Test Revisited, arXiv:1701.03992 (2016).
- [4] T. Choudhry, S. Hassan, S. Shabi, Relationship between gold and stock markets during the global crisis: Evidence from linear and nonlinear causality tests, *International Review of Finance Analysis* 41 (2015) 247-256.
- [5] T. Choudhry, F. Papadimitriou, S. Shabi, Stock market volatility and business cycle: Evidence from linear and nonlinear causality tests, *Journal of Banking and Finance* 66 (2016) 89-101.
- [6] C. Diks, V. Panchenko, A note on the Hiemstra-Jones test for Granger non-causality, *Studies in Nonlinear Dynamics and Econometrics* 9(2) (2005) Art. No. 4, 1-7.
- [7] C. Diks, V. Panchenko, A new statistic and practical guidelines for nonparametric Granger causality testing, *Journal of Economic Dynamics and Control* 30 (2006) 1647-1669.
- [8] C. Diks, M. Wolski, Nonlinear granger causality: guidelines for multivariate analysis, *Journal of Applied Econometrics* 31 (2016) 1333-1351.
- [9] J.M. Dufour, E. Renault, Short run and long run causality in time series: Theory, *Econometrica* 66 (1998) 1099-1125.

- [10] J.M. Dufour, D. Pelletier, E. Renault, Short run and long run causality in time series: Inference, *Journal of Econometrics* 132 (2006) 337-362.
- [11] E. Ghysels, J.B. Hill, K. Motegi, Testing for Granger causality with mixed frequency data, *Journal of Econometrics* 192(1) (2016) 207-230.
- [12] C.W.J. Granger, Investigating causal relations by econometric models and cross-spectral methods, *Econometrica* 37(3) (1969) 424-438.
- [13] C.W.J. Granger, *Forecasting in business and economics*, Academic Press, 32(1) (1989) 223-226.
- [14] C. Hiemstra, J.D. Jones, Testing for linear and nonlinear Granger causality in the stock price-volume relation, *Journal of Finance* 49(5) (1994) 1639-1664.
- [15] C. Hurlin, B. Venet, Granger causality tests in panel data models with fixed coefficients, *processed Paris University Paris IX* 10 (2001) 1-30.
- [16] I.A. Ibragimov, On the average of real zeroes of random polynomials. I. The coefficients with zero means, *Theory of Probability and its Applications* 16:2 (1971) 228-248.
- [17] N.X. Lam, Twin Deficits Hypothesis and Feldstein-Horioka Puzzle in Vietnam, *International Research Journal of Finance and Economics*, 101 (2012) 169.
- [18] X.L. Zheng, B.M. Chen, Stock market modeling and forecasting: a system adaptation approach, *Lecture Notes in Control and Information Sciences* (2013) 442.