## AN EXPLICIT ERROR TERM IN THEOREM A

NICOLAS BERGERON AND CARLOS MATHEUS

### 1. INTRODUCTION

Recall that Theorem A above ensures the existence of a constant  $\delta > 0$  such that the number N(V) of sLag fibrations with volume  $\leq V$  in a generic twistor family of K3 surfaces is

(1.1) 
$$N(V) = C \cdot V^{20} + O(V^{20-\delta})$$

where C > 0 is the ratio of volumes of two concrete homogenous spaces.

The goal of this appendix is to prove that  $\delta$  can be taken to be  $\left(\frac{4}{697633}\right)^{-}$ :

Theorem 1.1. In the same setting as Theorem A above, one actually has

$$N(V) = C \cdot V^{20} + O_{\varepsilon} \left( V^{\frac{13952050}{697633} + \varepsilon} \right)$$

10050050

for all  $\varepsilon > 0$ .

### 2. REDUCTION OF THEOREM 1.1 TO DYNAMICS IN HOMOGENOUS SPACES

Filip derived his counting formula (1.1) from certain equidistribution results. More precisely, let  $\Lambda_{\mathbb{Z}}$  be a lattice isomorphic to  $H^2(S,\mathbb{Z})$ , where S is a K3 surface. Fix  $P \subset \Lambda_{\mathbb{R}}$  a positive-definite 3-plane. Denote by  $\Lambda_{\mathbb{Z}}^0$  the set of primitive isotropic integral vectors and fix  $e \in \Lambda_{\mathbb{Z}}^0$ . For each  $v \in \Lambda_{\mathbb{R}} = P \oplus P^{\perp}$ , let  $v := (v)_P \oplus (v)_{P^{\perp}}$ with  $(v)_P \in P$  and  $(v)_{P^{\perp}} \in P^{\perp}$ . Consider the orthogonal group  $G := O(\Lambda_{\mathbb{R}})$ , the lattice  $\Gamma := O(\Lambda_{\mathbb{Z}})$  and the maximal compact subgroup  $K := O(P) \times O(P^{\perp})$ of G, and, for a fixed  $e \in \Lambda_{\mathbb{Z}}^0$ , denote by  $H_e := Stab_G(e)$  and  $\Gamma_e = Stab_{\Gamma}(e)$ .

The volumes of the locally homogenous spaces  $X := \Gamma \setminus G$  and  $Y := \Gamma_e \setminus H_e$  are finite. As it is observed in [3, pp. 4], the constants C > 0 and  $\delta > 0$  in (1.1) are the constant described in [3, Theorem 3.1.3]. In particular,

$$C = \frac{\text{Vol } Y}{20 \cdot \text{Vol } X}$$

The constant  $\delta > 0$  is related to the dynamics of a certain one-parameter subgroup  $a_t$  of  $G \simeq SO(3, 19)(\mathbb{R})$ . More concretely, given e and P as above, let e' be the isotropic vector given by

 $e' := (e)_P \oplus -(e)_{P^{\perp}}$  where  $e := (e)_P \oplus (e)_{P^{\perp}}$ 

In this context, we denote by  $\{a_t\}_{t\in\mathbb{R}}\subset G$  the one-parameter subgroup defined as

$$a_t \cdot e = \exp(-t) \cdot e, \quad a_t \cdot e' = \exp(t) \cdot e', \quad a_t|_{(e \oplus e')^{\perp}} = \mathrm{id}$$

Date: January 18, 2022.

It is explained in [3, Subsection 3.6.9] that<sup>1</sup> the quantity  $\delta$  in (1.1) is

(2.1) 
$$\delta = \frac{\delta_0}{d_{l_0} + 1}$$

where  $(d_l)_{l \in \mathbb{N}}$  are the exponents in [3, Proposition 3.5.10 (ii)], and  $\delta_0 > 0$ ,  $l_0 \in \mathbb{N}$  are the constants in the following equidistribution statement in [3, Theorem 4.3.1]:

(2.2) 
$$\int_{Ya_t} w \ d\mu_{Ya_t} = \frac{\text{Vol } Y}{\text{Vol } X} \int_X w \ d\mu_X + O(\|w\|_l \ e^{-\delta_0 t})$$

for all Sobolev scales  $l \ge l_0$  (see [3, §4.2.2] for the definition of the Sobolev norms in this context).

A quick inspection of the proof of [3, Proposition 3.5.10 (ii)] (related to the thickening of K) reveals that the exponents  $d_l$  depend linearly on l. In fact, the constant  $c_1(l)$  in [3, Equation (3.5.15)] gives the power of  $\varepsilon$  associated to the volumes of  $\varepsilon$ -balls at the origin of  $\mathfrak{p}_m \times \mathfrak{n}^+ \times \mathfrak{a}$ , that is,  $c_1(l) = \dim(G) - \dim(K)$  (and, hence,  $c_1(l)$  independs of l). Since the l-th derivative of  $\chi_{\varepsilon}$  is bounded by a multiple of  $\varepsilon^{-c_1(l)-l}$  and it is supported in a  $\varepsilon$ -neighborhood of K, the l-Sobolev norm of  $\chi_{\varepsilon}$  is bounded by a multiple of  $\varepsilon^{-l-c_1(l)/2}$ . Therefore,

(2.3) 
$$d_l := l + \frac{\dim(G) - \dim(K)}{2}$$

## 3. EQUIDISTRIBUTION AND RATES OF MIXING

The constants  $\delta_0 > 0$  and  $l_0 \in \mathbb{N}$  in (2.2) are described in [3, pp. 36] and they are related to the geometry of  $Y \subset X$  and the rate of mixing of  $a_t$ .

3.1. **Injectivity radius.** We denote by inj(x) the local injectivity radius at a point  $x \in X$  and we let  $Y_{\varepsilon} := \{y \in Y : inj(y) \ge \varepsilon\}$ . By [3, Proposition 4.1.3], we know that the arguments of [1, Lemma 11.2] provide a constant p > 0 such that  $\mu_Y(Y \setminus Y_{\varepsilon}) = O(\varepsilon^p)$ . Actually, a close inspection of these arguments (of integration over Siegel sets) reveal that p = 1 in our specific setting (of  $G \simeq SO(3, 19)(\mathbb{R})$ ):

(3.1) 
$$\mu_Y(Y \setminus Y_{\varepsilon}) = O(\varepsilon)$$

3.2. Thickening of Y. Let us fix some parameter 0 < p' < 1 (very close to one in practice) and consider [3, Proposition 4.1.6] (of thickening of Y) where it is constructed a family of smooth versions  $\phi_{\varepsilon}$  of the characteristic function of Y. As it turns out,  $\phi_{\varepsilon}$  is the product of two functions:  $\tau_{\varepsilon}$  is a bump function supported<sup>2</sup> on  $Y_{\varepsilon p'}$  and  $\rho_{\varepsilon}$  is a bump function supported on the  $\varepsilon$ -neighborhood of the identity in a certain Lie group N' of dimension dim $(N') = \dim(X) - \dim(Y)$ .

<sup>&</sup>lt;sup>1</sup>Indeed, [3, pp. 29] says that the optimal choice of  $\delta$  occurs precisely when the terms  $\varepsilon e^{20T} = e^{(20-\delta)T}$  and  $\varepsilon^{-d_{l_0}} e^{(20-\delta_0)T} = e^{(20-\delta_0+\delta d_{l_0})T}$  have the same order in T.

<sup>&</sup>lt;sup>2</sup>In fact, Filip sets p' = 1/2 for his construction of  $\tau_{\varepsilon}$ , but any value of 0 < p' < 1 can be taken here: indeed, the construction of  $\tau_{\varepsilon}$  can be made as soon as the local product structure statement [3, Proposition 4.1.5] holds (and this is the case for any choice of 0 < p' < 1 because  $\varepsilon^{p'} \gg 2\varepsilon$  for all sufficiently small  $\varepsilon > 0$ ).

The bump function  $\rho_{\varepsilon}$  is obtained by rescaling of a fixed smooth bump function on N', so that its *l*-th Sobolev norm satisfies  $\|\rho_{\varepsilon}\|_{l} = O(\varepsilon^{-l - \frac{\dim(X) - \dim(Y)}{2}})$ .

The function  $\tau_{\varepsilon}$  is

$$\tau_{\varepsilon} = \frac{\sum_{y_j \in \mathcal{F}} \beta_{y_j,\varepsilon}}{\sum_{y_i \in \mathcal{G}} \beta_{y_i,\varepsilon}} := \frac{\sum_{y_j \in \mathcal{F}} \beta_{y_j,\varepsilon}}{\beta_{\mathcal{G},\varepsilon}}$$

where  $\{y_k\} \subset Y_{\varepsilon^{p'}}$  is a maximal collection of points such that the balls  $B(y_k, \varepsilon^3) \subset Y$  are mutually disjoint,  $\mathcal{F} = \{y_k\} \cap Y_{4\varepsilon^{p'}}, \mathcal{G} = \{y_k\} \cap Y_{2\varepsilon^{p'}}$ , and the functions  $\beta_{,\varepsilon}$  are translates of a bump function  $\beta_{\varepsilon}$  whose *l*-th Sobolev norm is  $\|\beta_{\varepsilon}\|_l = O(\varepsilon^{-l + \frac{\dim(Y)}{2}})$ .

On one hand, since a ball B of radius  $\varepsilon$  at a point of  $Y_{\varepsilon^{p'}}$  has volume  $O(\varepsilon^{\dim(Y)})$ , the cardinality of  $\mathcal{G} \cap B$  is  $O(\varepsilon^{-2\dim(Y)})$ , the arguments in [1, pp. 1928] imply that the  $L^{\infty}$ -norm of the first l derivatives of  $1/\beta_{\mathcal{G},\varepsilon}$  is  $O(\varepsilon^{-l-2\dim(Y)})$ . On the other hand, the cardinality of  $\mathcal{F}$  is  $O(\varepsilon^{-3\dim(Y)})$  and  $\|\beta_{y_i,\varepsilon}\|_l = \|\beta_{\varepsilon}\|_l$ . It follows that

$$\|\tau_{\varepsilon}\|_{l} = O(\varepsilon^{-l - \frac{9\dim(Y)}{2}}).$$

By inserting these facts into the definition of  $\phi_{\varepsilon}$  in [3, Equation (4.1.7)], we deduce from Sobolev's lemma that

(3.2) 
$$\|\phi_{\varepsilon}\|_{l} = O(\varepsilon^{-2l-4\dim(Y) - \frac{\dim(X)}{2}})$$

for all  $l > \dim(X)/2$ , that is, the constant  $C_l$  in [3, Proposition 4.1.6 (iii)] is

$$C_l := 2l + 4\dim(Y) + \frac{\dim(X)}{2}$$

For later use, notice that  $\phi_{\varepsilon}$  verifies  $\int_X \phi_{\varepsilon} d\mu_X = \text{Vol } Y + O(\text{Vol}(Y \setminus Y_{\varepsilon^{p'}}))$ . By combining this estimate with (3.1), we get

(3.3) 
$$\int_X \phi_{\varepsilon} \ d\mu_X = \operatorname{Vol} Y + O(\varepsilon^{p'})$$

3.3. Wavefront lemma. The proof of Lemma 4.1.10 in [3] says that

$$\int_X w \cdot (\phi_{\varepsilon} \cdot a_t) d\mu_X = \int_Y w(ya_t) d\mu_Y(y) + O(\varepsilon \operatorname{Lip}(w)) + O(\varepsilon^{pp'} ||w||_{L^{\infty}})$$

where p > 0 is the parameter such that  $\mu_Y(Y \setminus Y_{\varepsilon^{p'}}) = O(\varepsilon^{pp'})$ . Therefore, we deduce from (3.1) and Sobolev's lemma that

(3.4) 
$$\int_X w \cdot (\phi_{\varepsilon} \cdot a_t) d\mu_X = \int_Y w(ya_t) d\mu_Y(y) + O(\varepsilon^{p'} ||w||_l)$$

for all  $l > 1 + \dim(X)/2$ .

3.4. **Reduction of equidistribution to rate of mixing.** By following [3, pp. 36], let us compute the constants  $\delta_0 > 0$  and  $l_0 \in \mathbb{N}$  in (2.2) in terms of the following quantitative mixing statement: there exists  $\delta'_0 > 0$  such that

(3.5) 
$$\left| \int_{X} \alpha \cdot (\beta \cdot g) d\mu - \left( \int_{X} \alpha d\mu \right) \left( \int_{X} \beta d\mu \right) \right| = O(\|\alpha\|_{l} \|\beta\|_{l} \|g\|^{-\delta'_{0}})$$

for all  $l \ge l'_0$ . (Here,  $\mu = \mu_X / \text{Vol } X$  is the normalized Haar measure.) For this sake, we observe that (3.5) says that

$$\int_X w \cdot (\phi_{\varepsilon} \cdot a_t) d\mu_X = \frac{1}{\operatorname{Vol} X} \left( \int_X w d\mu_X \right) \left( \int_X \phi_{\varepsilon} d\mu_X \right) + O(\|w\|_l \|\phi_{\varepsilon}\|_l e^{-t\delta_0'})$$
  
for  $l > l_0'$ .

By (3.2) and (3.3), the previous estimate implies

$$\int_X w \cdot (\phi_{\varepsilon} \cdot a_t) d\mu_X = \frac{\operatorname{Vol} Y}{\operatorname{Vol} X} \left( \int_X w d\mu_X \right) + O(\varepsilon^{p'} \|w\|_l) + O(\|w\|_l \varepsilon^{-C_l} e^{-t\delta'_0})$$

for all  $l \ge \max\{l'_0, \lfloor \dim(X)/2 \rfloor + 1\}$ .

By plugging (3.4) into the estimate above, we conclude that

$$\int_{Ya_t} w d\mu_{Ya_t} = \frac{\operatorname{Vol} Y}{\operatorname{Vol} X} \left( \int_X w d\mu_X \right) + O(\varepsilon^{p'} \|w\|_l) + O(\|w\|_l \varepsilon^{-C_l} e^{-t\delta'_0})$$

for all  $l \ge l_0 := \max\{l'_0, \lfloor \dim(X)/2 \rfloor + 2\}$ . By taking  $\varepsilon := e^{-\delta''_0 t}$  and by optimizing<sup>3</sup> the value of  $\delta''_0$ , we obtain that

$$\int_{Ya_t} w d\mu_{Ya_t} = \frac{\operatorname{Vol} Y}{\operatorname{Vol} X} \left( \int_X w d\mu_X \right) + O(\|w\|_{l_0} e^{-t\delta_0})$$

for  $l_0 := \max\{l'_0, \lfloor \dim(X)/2 \rfloor + 2\}$  and  $\delta_0 := \frac{p'}{p' + C_{l_0}} \delta'_0$ .

Since 0 < p' < 1 is an arbitrary parameter, we deduce that (2.2) holds for  $l_0 := \max\{l'_0, |\dim(X)/2| + 2\}$  and any choice of

(3.6) 
$$0 < \delta_0 < \frac{\delta'_0}{1 + 2l_0 + 4\dim(Y) + \frac{\dim(X)}{2}}$$

#### 4. RATES OF MIXING AND REPRESENTATION THEORY

**Definition 4.1.** 1. A *unitary representation*  $\pi$  of G in a (separable) Hilbert space  $\mathcal{H}_{\pi}$  is a morphism  $G \to U(\mathcal{H}_{\pi})$  such that for any  $v \in \mathcal{H}_{\pi}$  the map  $G \to \mathcal{H}_{\pi}$ ;  $g \mapsto \pi(g)v$  is continuous. If this map is smooth one says that v is a  $C^{\infty}$ -vector of  $\pi$ . We denote by  $\mathcal{H}^{\infty}_{\pi}$  the set of  $C^{\infty}$ -vectors of  $\pi$ .

2. Given two vectors  $v, w \in \mathcal{H}_{\pi}$ , we define the *matrix coefficient*  $c_{v,w} : G \to \mathbb{C}$ of  $\pi$  as the continuous map  $g \mapsto \langle \pi(g)v, w \rangle$ . The coefficient  $c_{v,w}$  is said to be *K*-finite if both the vector spaces generated by  $\pi(K) \cdot v$  and  $\pi(K) \cdot w$  are finite dimensional.

3. Let  $p(\pi)$  be the infimum of the set of real numbers  $p \ge 2$  such that all K-finite matrix coefficients of  $\pi$  are in  $L^p(G)$ .

<sup>&</sup>lt;sup>3</sup>I.e., we choose  $\delta_0'' > 0$  so that  $\varepsilon^{p'} = \varepsilon^{-C_{l_0}} e^{-t\delta_0'}$ .

4. Say that a unitary representation  $\sigma$  of G is *weakly contained* in  $\pi$  if any matrix coefficient of  $\sigma$  can be obtained as the limit, with respect to the topology of uniform convergence on compact subsets, of a sequence of matrix coefficients of  $\pi$ .

Given an element  $g = nak \in G$ , we write  $a = e^{H(g)}$ . The Harish-Chandra function is  $\Xi = \Xi_G : G \to \mathbb{R}$  defined by

$$\Xi(g) = \int_{K} e^{-\rho(H(kg^{-1}))} dk$$

where  $\rho$  is half the sum of the positive restricted roots counting multiplicities. The Harish-Chandra function decreases exponentially fast along  $A^+$ ; modulo a polynomial factor of a logarithmic argument, it decreases like  $e^{-\rho(H)}$ .

Let  $d = \dim(K)$  be the dimension of K and fix a basis  $\mathcal{B}$  of the Lie algebra  $\mathfrak{k}$  of K. Given a smooth vector  $v \in \mathcal{H}^{\infty}_{\pi}$  we set

$$S(v) = \sum_{\operatorname{ord}(D) \le \lfloor d/2 \rfloor + 1} ||\pi(D)v||,$$

where D varies among all monomials in elements of B of degree  $\leq \lfloor d/2 \rfloor + 1$  and, if  $X_1, \ldots, X_r$  are elements of B, we have  $\pi(X_1 \cdots X_r) = \pi(X_1) \cdots \pi(X_r)$  and each  $\pi(X_i)$  acts by derivation.

**Proposition 4.2.** For all positive  $\varepsilon$  and  $k \in \mathbb{N}^*$ , there exists a constant  $C = C(\varepsilon, k)$  such that if  $\pi$  is a unitary representation of G with  $p(\pi) \leq 2k$ , then for all  $v, w \in \mathcal{H}^{\infty}_{\pi}$  and for all positive t we have:

(4.1) 
$$|\langle \pi(a_t)v, w \rangle| \le CS(v)S(w)e^{-(p/k-\varepsilon)t}$$

where  $p = \rho(H)$  and H is the infinitesimal generator of the one-parameter subgroup  $(a_t)$ .

*Proof.* Up to replacing  $\pi$  by the tensor product  $\pi^{\otimes k}$  we may suppose that k = 1; see [2, p. 108]. It then follows from [2, Theorem 1] that  $\pi$  is weakly contained in the (right) regular representation  $L^2(G)$ . We are then reduced to prove the proposition in the case where  $\pi$  is the regular representation of G (and k = 1); see the proof of [2, Theorem 2] for more details on this last reduction.

Now consider v and w in  $L^2(G) \cap C^{\infty}(G)$ . The functions

$$\varphi: x \mapsto \sup_{k \in K} |v(xk)| \text{ and } \psi: x \mapsto \sup_{k \in K} |w(xk)|$$

are both positive and K-invariant, and we have:

$$|\langle \pi(a_t)v, w \rangle_{L^2(G)}| \le \int_G \varphi(xa_t)\psi(x)dx = |\langle \pi(a_t)\varphi, \psi \rangle_{L^2(G)}|.$$

Now the Sobolev lemma (see [5, Proposition 2.6]) implies that the  $L^{\infty}$  norms of  $\varphi$  and  $\psi$  can be estimated in terms of their Sobolev norms along K. More precisely: there exists a constant c such that the for all  $x \in G$ ,

$$\varphi(x)^{2} = \sup_{k \in K} |v(xk)|^{2} \le c \sum_{\operatorname{ord}(D) \le \lfloor d/2 \rfloor + 1} ||(\rho(D)v)(x \cdot)||_{L^{2}(K)}.$$

Integrating over G (here we assume for simplicity that the measure of K is 1) one concludes that  $||\varphi||_{L^2(G)} \leq \sqrt{c}S(v)$  and similarly for  $\psi$ . It remains to prove that there exists a constant  $d_{\varepsilon}$  such that if  $\varphi, \psi \in L^2(G)$  are two K-invariant, positive functions of norm 1, then

$$|\langle \pi(a_t)\varphi,\psi\rangle| \le d_{\varepsilon}e^{-(p/k+\varepsilon)t}.$$

First it follows from the computations of [2, pp. 106-107] that

$$\begin{split} |\langle \pi(g)\varphi,\psi\rangle| &= \int_{K} \left( \int_{NA} \varphi(na)\psi(nakg^{-1})e^{2\rho(\log a)}dnda \right) dk \\ &\leq ||\varphi||_{L^{2}(G)} \int_{K} \left( \int_{NA} \psi(naH(kg^{-1}))^{2}e^{2\rho(\log a)}dnda \right)^{1/2} dk \\ &= ||\varphi||_{L^{2}(G)} \cdot ||\psi||_{L^{2}(G)} \int_{K} e^{-\rho(H(kg^{-1}))}dk = ||\varphi||_{L^{2}(G)} \cdot ||\psi||_{L^{2}(G)} \Xi(g) \end{split}$$

Now recall that, up to "polynomial factors of logarithmic arguments", the function  $\Xi(a_t)$  decreases like  $e^{-t\rho(H)} = e^{-pt}$ . The proposition follows.

We shall apply this proposition to the (quasi-)regular representation  $\pi$  of G in the subspace  $L_0^2(\Gamma \setminus G)$  of  $L^2(\Gamma \setminus G)$  that is orthogonal to the space of constant functions. It follows from [4] that  $p(\pi) = 20$ . Proposition 4.2 therefore applies with k = 10. Note that in our case p = 10.

Now let  $\alpha$  and  $\beta$  be two smooth functions in  $L^2(X)$  then

$$\alpha_0 := \alpha - \int_X \alpha d\mu$$
 and  $\beta_0 := \beta - \int_X \beta d\mu \in L^2_0(X)$ 

and we have:

$$\langle \pi(g)\alpha_0,\beta_0\rangle_{L^2_0(X)} = \int_X \alpha \cdot (\beta \cdot g)d\mu - \left(\int_X \alpha d\mu\right) \left(\int_X \beta d\mu\right).$$

From Proposition 4.2 and the fact that  $S(\alpha) \leq ||\alpha||_{|d/2|+1}$  we conclude that

$$\left|\int_{X} \alpha \cdot (\beta \cdot a_{t}) d\mu - \left(\int_{X} \alpha d\mu\right) \left(\int_{X} \beta d\mu\right)\right| = O(\|\alpha\|_{l} \|\beta\|_{l} e^{-t\delta_{0}'})$$

for any  $l \ge l'_0 := \lfloor \dim(K)/2 \rfloor + 1$  and any  $\delta'_0 < 1$ .

# 5. End of proof of Theorem 1.1

The explicit value of  $\delta$  announced in Theorem 1.1 can be easily derived from the discussion above. Indeed, we just saw in Section 4 that  $\delta'_0 = 1-$  and  $l'_0 = \lfloor \dim(K)/2 \rfloor + 1$ . Because  $174 = \dim(K) < \dim(X) = 231$  and  $\dim(Y) = 210$ , we deduce from (3.6) that  $l_0 = \lfloor \dim(X)/2 \rfloor + 2 = 117$  and

$$\delta_0 = \left(\frac{1}{1+2\times 117 + 4\times 210 + \frac{231}{2}}\right)^- = \left(\frac{2}{2381}\right)^-$$

Finally, by inserting these informations into (2.3) and (2.1), we conclude that

$$\delta = \frac{\delta_0}{l_0 + \frac{57}{2} + 1} = \left(\frac{4}{697633}\right)^- \approx (5.7336737224 \dots \times 10^{-6})^-.$$

## REFERENCES

- [1] Y. Benoist and H. Oh, *Effective equidistribution of S-integral points on symmetric varieties*, Ann. Inst. Fourier (Grenoble) 62 (2012), no. 5, 1889–1942.
- M. Cowling, U. Haagerup and R. Howe, Almost L<sup>2</sup> matrix coefficients, J. Reine Angew. Math. 387 (1988), p.97–110.
- [3] S. Filip, *Counting special Lagrangian fibrations in twistor families of K3 surfaces*, Preprint (2016) available at arXiv:161208684.
- [4] J.-S. Li, *The Minimal Decay of Matrix Coefficients for Classical Groups*, in Harmonic Analysis in China Volume 327 of the series Mathematics and Its Applications pp 146-169.
- [5] C. C. Moore, *Exponential decay of correlation coefficients for geodesic flows*, Group representations, ergodic theory, operator algebras, and mathematical physics (Berkeley, Calif., 1984), Math. Sci. Res. Inst. Publ., vol. 6, Springer, New York, 1987, p. 163–181.

NICOLAS BERGERON: UNIVERSITÉ PIERRE ET MARIE CURIE, INSTITUT DE MATHÉMA-TIQUES DE JUSSIEU, CNRS (UMR 7586), 4, PLACE JUSSIEU 75252 PARIS CEDEX 05, FRANCE *E-mail address*: nicolas.bergeron@imj-prg.fr

CARLOS MATHEUS: CMLS, ÉCOLE POLYTECHNIQUE, CNRS (UMR 7640), 91128 PALAISEAU, FRANCE

E-mail address: carlos.matheus@math.cnrs.fr