

Controlled Flavour Changing Neutral Couplings in Two Higgs Doublet Models

Joao M. Alves ^{a,1} Francisco J. Botella ^{b,2}, Gustavo C. Branco ^{a,3},
Fernando Cornet-Gomez ^{b,4}, Miguel Nebot ^{c,5}

^a *Departamento de Física and Centro de Física Teórica de Partículas (CFTP),
Instituto Superior Técnico (IST), U. de Lisboa (UL),
Av. Rovisco Pais, P-1049-001 Lisboa, Portugal.*

^b *Departament de Física Teòrica and IFIC,
Universitat de València-CSIC,
E-46100, Burjassot, Spain.*

^c *Centro de Física Teórica de Partículas (CFTP),
Instituto Superior Técnico (IST), U. de Lisboa (UL),
Av. Rovisco Pais, P-1049-001 Lisboa, Portugal.*

Abstract

We propose a class of Two Higgs Doublet Models where there are Flavour Changing Neutral Currents (FCNC) at tree level, but under control due to the introduction of a discrete symmetry in the full Lagrangian. It is shown that in this class of models, one can have simultaneously FCNC in the up and down sectors, in contrast to the situation encountered in BGL models. The intensity of FCNC is analysed and it is shown that in this class of models one can respect all the strong constraints from experiment without unnatural fine-tuning. It is pointed out that the additional sources of flavour and CP violation are such that they can enhance significantly the generation of the Baryon Asymmetry of the Universe, with respect to the Standard Model.

¹j.magalhaes.alves@tecnico.ulisboa.pt

²Francisco.J.Botella@uv.es

³gbranco@tecnico.ulisboa.pt

⁴Fernando.Cornet@ific.uv.es

⁵miguel.r.nebot.gomez@tecnico.ulisboa.pt

1 Introduction

Recently, there has been a special interest in the scalar sector of the Standard Model (SM) and some of its extensions, motivated by the discovery by both ATLAS [1] and CMS [2], of a particle which can be interpreted as the Higgs Boson of the SM. A central question is whether the Higgs couplings to quarks and leptons are like those in the SM or whether Nature chooses instead a more complex scalar sector. The simplest extension of the scalar sector consists of the addition of another Higgs doublet and the first 2 Higgs Doublet Model (2HDM) was introduced by Lee [3] in order to achieve spontaneous CP violation. The presence of extra symmetries has an important impact on the possibility of generating spontaneous CP violation, an aspect which has been recently reviewed [4, 5]. If no extra symmetries are introduced, the general 2HDM has Flavour Changing Neutral Currents (FCNC) at tree level which have to be suppressed in order to avoid violation of the stringent experimental bounds. The simplest way of avoiding FCNC in the scalar sector is by postulating that quarks of a given charge receive contributions to their mass only from one Higgs doublet. These selective couplings [6, 7] can be easily implemented in a natural way through the introduction of a Z_2 symmetry which leads to Natural Flavour Conservation (NFC) in the scalar sector [6]. Another way of eliminating FCNC at tree level is through the hypothesis of alignment of Yukawa couplings in flavour space [8]. Various studies have been done on the constraints implied by FCNC, in the framework of 2HDM [9–16].

A very interesting approach to the control of FCNC is provided by Branco-Grimus-Lavoura (BGL) models [17], where there are FCNC at tree level but their flavour structure is controlled by the elements of the Cabibbo-Kobayashi-Maskawa (CKM) matrix V , with no other flavour parameters. Although these models were proposed some years before the Minimal Flavour Violation (MFV) hypothesis [18–20] was put forward, BGL models satisfy the MFV hypothesis of having all the flavour structure of New Physics controlled by V [17, 21]. BGL models are renormalisable, since the flavour structure of the Yukawa couplings results from an exact symmetry of the Lagrangian. The general 2HDM contains FCNC, with their flavour structure parametrized by two complex matrices N_d , N_u [22]. These matrices depend on a large number of parameters, in particular on $\mathcal{U}_L^d, \mathcal{U}_L^u, \mathcal{U}_R^d$ and \mathcal{U}_R^u , the unitary matrices which enter in the diagonalization of the down and up quark mass matrices. Having N_d , N_u to depend only on $V = \mathcal{U}_L^{u\dagger} \mathcal{U}_L^d$ in a natural way looks like an impossible task. Yet, this task is accomplished by BGL models, which were first constructed for the quark sector and then generalized to the lepton sector [23]. There are six types of BGL models in the quark sector and six (three) types in the lepton sector for Dirac (Majorana) neutrinos, which can be combined to have a total of 36 (18) BGL models, with different phenomenological implications, which were thoroughly analyzed [24–26]. An interesting feature of BGL models is the fact they contain FCNC either in the up or the down sectors but not in both sectors.

In this paper, we analyze the possibility of generalizing BGL models, having in mind the following questions:

1. Can one have a framework based on 2HDM which keeps the requirement of renormalisability and includes all 36 (18) models in a single model which contain

each one of the BGL models as special cases?

2. Is it possible to have renormalisable 2HDM with controlled but non-vanishing FCNC in both the up and down quark sectors?

The paper is organised as follows. In order to settle the notation, we consider in the next section the general 2HDM with no symmetry introduced, beyond the gauge symmetry. We briefly review BGL models and then propose a more general framework denoted gBGL containing BGL models in special limits. In section 3 we examine Yukawa textures corresponding to gBGL models and show how they contain BGL models as special cases. We also derive in this section Weak Basis (WB) invariant conditions for having gBGL models. In section 4 we propose a convenient parametrisation of gBGL models through the use of WB covariant projectors. In section 5 we describe the scalar potential. In section 6 we analyse the intensity of FCNC in gBGL models, with particular emphasis, in section 7, on models close to BGL models of types b and t . In section 8 the implications of gBGL models for the Baryon Asymmetry of the Universe are discussed and our conclusions are contained in the last section.

2 Generalising BGL models: gBGL

In order to settle the notation, we start by recalling the structure of the Yukawa couplings in the quark sector of a general 2HDM, with no extra symmetry introduced in the Lagrangian:

$$\mathcal{L}_Y = -\bar{Q}_L^0(\Gamma_1\Phi_1 + \Gamma_2\Phi_2)d_R^0 - \bar{Q}_L^0(\Delta_1\tilde{\Phi}_1 + \Delta_2\tilde{\Phi}_2)u_R^0 + \text{h.c.}, \quad (1)$$

with $\tilde{\Phi}_j = i\sigma_2\Phi_j^*$. Electroweak symmetry is spontaneously broken via the vacuum expectation values

$$\langle\Phi_1\rangle = \begin{pmatrix} 0 \\ e^{i\theta_1}v_1/\sqrt{2} \end{pmatrix}, \quad \langle\Phi_2\rangle = \begin{pmatrix} 0 \\ e^{i\theta_2}v_2/\sqrt{2} \end{pmatrix}. \quad (2)$$

Introducing as usual $\theta = \theta_2 - \theta_1$, $v^2 = v_1^2 + v_2^2$, $c_\beta = \cos\beta \equiv v_1/v$, $s_\beta = \sin\beta \equiv v_2/v$ and $t_\beta \equiv \tan\beta$, the Higgs doublets can be rotated into the ‘‘Higgs basis’’ [27–29]

$$\begin{pmatrix} H_1 \\ H_2 \end{pmatrix} = \begin{pmatrix} c_\beta & s_\beta \\ s_\beta & -c_\beta \end{pmatrix} \begin{pmatrix} e^{-i\theta_1}\Phi_1 \\ e^{-i\theta_2}\Phi_2 \end{pmatrix}, \quad (3)$$

where

$$\langle H_1 \rangle = \begin{pmatrix} 0 \\ v/\sqrt{2} \end{pmatrix}, \quad \langle H_2 \rangle = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \quad (4)$$

The Yukawa couplings in eq. (1) read

$$\mathcal{L}_Y = -\frac{\sqrt{2}e^{i\theta_1}}{v}\bar{Q}_L^0(M_d^0H_1 + N_d^0H_2)d_R^0 - \frac{\sqrt{2}e^{-i\theta_1}}{v}\bar{Q}_L^0(M_u^0\tilde{H}_1 + N_u^0\tilde{H}_2)u_R^0 + \text{h.c.}, \quad (5)$$

with the mass matrices M_d^0 , M_u^0 , and the N_d^0 , N_u^0 matrices given by

$$M_d^0 = \frac{v}{\sqrt{2}}(c_\beta \Gamma_1 + e^{i\theta} s_\beta \Gamma_2), \quad N_d^0 = \frac{v}{\sqrt{2}}(s_\beta \Gamma_1 - e^{i\theta} c_\beta \Gamma_2), \quad (6)$$

$$M_u^0 = \frac{v}{\sqrt{2}}(c_\beta \Delta_1 + e^{-i\theta} s_\beta \Delta_2), \quad N_u^0 = \frac{v}{\sqrt{2}}(s_\beta \Delta_1 - e^{-i\theta} c_\beta \Delta_2). \quad (7)$$

It is clear from eqs. (6)–(7) that N_d^0 , N_u^0 are complex matrices containing a huge number of new parameters, including FCNC which have strong experimental constraints. In order to avoid FCNC, Glashow and Weinberg [6] introduced the NFC principle, which constrains the Yukawa couplings so that a quark of a given charge only receives mass from one Higgs doublet. BGL models also control the size of FCNC, while allowing for their appearance at tree level. They have some remarkable features which can be summarized in the following way.

- (i) BGL models are renormalizable, since the pattern of Yukawa couplings are dictated by a symmetry of the full lagrangian.
- (ii) In BGL models, the couplings of the physical neutral scalars to the quark mass eigenstates only depend on V , t_β and the quark masses, with no other parameters.
- (iii) In BGL models there are FCNC either in the up or down sectors, but not in both. It has been shown [30] that if one imposes a flavour symmetry such that the FCNC only depend on V and further assumes that the flavour symmetry is abelian, then BGL models are unique.

Let us recall the symmetries used in order to construct BGL models:

- up-type BGL models (uBGL in the following) are implemented by the symmetry

$$\begin{aligned} Q_{L3} &\mapsto e^{i\tau} Q_{L3}, \\ d_R &\mapsto d_R, \quad \Phi_1 \mapsto \Phi_1, \\ u_{R3} &\mapsto e^{i2\tau} u_{R3}, \quad \Phi_2 \mapsto e^{i\tau} \Phi_2, \end{aligned} \quad (8)$$

- down-type BGL models (dBGL in the following) are implemented by the symmetry

$$\begin{aligned} Q_{L3} &\mapsto e^{i\tau} Q_{L3}, \\ d_{R3} &\mapsto e^{i2\tau} d_{R3}, \quad \Phi_1 \mapsto \Phi_1, \\ u_R &\mapsto u_R, \quad \Phi_2 \mapsto e^{i\tau} \Phi_2, \end{aligned} \quad (9)$$

with $\tau \neq 0, \pi$. It has been shown [23] that if one extends a BGL model to the lepton sector, with Majorana neutrinos and a realistic seesaw mechanism, then $\tau = \pi/2$ and one is led to a \mathbb{Z}_4 symmetry.

In this paper, we address ourselves to the question whether it is possible to generalise BGL models so that the new class of models, called generalised BGL (gBGL), keep

some of the interesting features of BGL models, like renormalizability, but allow for FCNC both in the up and the down sectors. The gBGL models are implemented through a \mathbb{Z}_2 symmetry, where u_R and d_R are even and only one of the scalar doublets and one of the left-handed quark doublets are odd:

$$\begin{aligned} Q_{L_3} &\mapsto -Q_{L_3}, \\ d_R &\mapsto d_R, & \Phi_1 &\mapsto \Phi_1, \\ u_R &\mapsto u_R, & \Phi_2 &\mapsto -\Phi_2. \end{aligned} \tag{10}$$

The above gBGL model includes all BGL models as special cases. Indeed gBGL models have some new parameters and when some of these free parameters are set to zero, one obtains a BGL model and the Lagrangian acquires a larger symmetry, namely \mathbb{Z}_4 ⁶. It is worth emphasizing that gBGL models are implemented through a \mathbb{Z}_2 symmetry, as it is also the case in the Glashow–Weinberg model with NFC. The only difference is that the left-handed quark families transform differently in the two models. In words, one may say that *the principle leading to gBGL constrains the Yukawa couplings so that each line of Γ_j , Δ_j couples only to one Higgs doublet*.

3 Yukawa Textures

Imposing the \mathbb{Z}_2 symmetry in eq. (10), the Yukawa matrices in these models have the general form

$$\begin{aligned} \Gamma_1 &= \begin{pmatrix} \times & \times & \gamma_{13} \\ \times & \times & \gamma_{23} \\ 0 & 0 & 0 \end{pmatrix}, & \Gamma_2 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \gamma_{31} & \gamma_{32} & \times \end{pmatrix}, \\ \Delta_1 &= \begin{pmatrix} \times & \times & \delta_{13} \\ \times & \times & \delta_{23} \\ 0 & 0 & 0 \end{pmatrix}, & \Delta_2 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \delta_{31} & \delta_{32} & \times \end{pmatrix}, \end{aligned} \tag{11}$$

where \times , γ_{ij} and δ_{ij} stand for arbitrary complex parameters. In eq. (11), the γ_{ij} and δ_{ij} entries have been singled out in order to show how gBGL contain BGL models as special cases: it is evident that taking $\gamma_{ij} = 0$, we obtain dBGL models, where

$$\begin{aligned} \Gamma_1 &= \begin{pmatrix} \times & \times & 0 \\ \times & \times & 0 \\ 0 & 0 & 0 \end{pmatrix}, & \Gamma_2 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \times \end{pmatrix}, \\ \Delta_1 &= \begin{pmatrix} \times & \times & \times \\ \times & \times & \times \\ 0 & 0 & 0 \end{pmatrix}, & \Delta_2 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \times & \times & \times \end{pmatrix}, \end{aligned} \tag{12}$$

⁶In the present work, we focus on the generalisation of BGL models in the quark sector and do not address in the detail the inclusion of the leptonic sector.

while taking $\delta_{ij} = 0$ we obtain uBGL models, with

$$\begin{aligned}\Gamma_1 &= \begin{pmatrix} \times & \times & \times \\ \times & \times & \times \\ 0 & 0 & 0 \end{pmatrix}, & \Gamma_2 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \times & \times & \times \end{pmatrix}, \\ \Delta_1 &= \begin{pmatrix} \times & \times & 0 \\ \times & \times & 0 \\ 0 & 0 & 0 \end{pmatrix}, & \Delta_2 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \times \end{pmatrix}.\end{aligned}\tag{13}$$

That is, this class of renormalisable gBGL models includes both up and down type BGL models, as we were looking for. It is also clear that FCNC are present both in the up and in the down sectors: the appearance of FCNC in one sector depends on the Yukawa couplings of that sector alone, without regard to the Yukawa couplings in the other sector. In these gBGL models, both up and down sectors have the FCNC-inducing structure that in BGL models is confined to one and only one sector.

3.1 Weak basis invariant conditions

As a summary of the previous discussion, gBGL models are defined by a \mathbb{Z}_2 symmetry or by the following matrix textures:

$$\begin{aligned}\Gamma_1 &= \begin{pmatrix} \times & \times & \times \\ \times & \times & \times \\ 0 & 0 & 0 \end{pmatrix}, & \Gamma_2 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \times & \times & \times \end{pmatrix}, \\ \Delta_1 &= \begin{pmatrix} \times & \times & \times \\ \times & \times & \times \\ 0 & 0 & 0 \end{pmatrix}, & \Delta_2 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \times & \times & \times \end{pmatrix}.\end{aligned}\tag{14}$$

Obviously, these zero texture structures are valid in a particular set of Weak Basis (WB), the WB where the definition of the symmetry applies – WB transformations are discussed in detail in section 4.1 –. Introducing the projection operator P_3 ,

$$P_3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad P_3 P_3 = P_3, \quad (\mathbf{1} - P_3) P_3 = 0,\tag{15}$$

it is straightforward to check that imposing the textures in eqs. (14) is equivalent to the following definition of gBGL models:

$$\begin{aligned}P_3 \Gamma_1 &= 0, & P_3 \Gamma_2 &= \Gamma_2, \\ P_3 \Delta_1 &= 0, & P_3 \Delta_2 &= \Delta_2.\end{aligned}\tag{16}$$

BGL models, are defined by more relations: in terms of P_3 , uBGL models satisfy

$$\begin{aligned}P_3 \Gamma_1 &= 0, & P_3 \Gamma_2 &= \Gamma_2, \\ P_3 \Delta_1 &= 0, & P_3 \Delta_2 &= \Delta_2, \\ \Delta_1 P_3 &= 0, & \Delta_2 P_3 &= \Delta_2,\end{aligned}\tag{17}$$

while dBGL models satisfy

$$\begin{aligned} P_3 \Gamma_1 &= 0, & P_3 \Gamma_2 &= \Gamma_2, \\ P_3 \Delta_1 &= 0, & P_3 \Delta_2 &= \Delta_2, \\ \Gamma_1 P_3 &= 0, & \Gamma_2 P_3 &= \Gamma_2. \end{aligned} \tag{18}$$

The last two conditions in eqs. (17) and (18) give the block diagonal form of the Yukawa matrices in the corresponding sector (up in uBGL and down in dBGL models), enforcing the absence of FCNC in that sector. From these conditions valid in a set of WB, we can get WB independent matrix conditions for all three types of models. The conditions of interest for gBGL models are

$$\begin{aligned} \Gamma_2^\dagger \Gamma_1 &= 0, & \Gamma_2^\dagger \Delta_1 &= 0, \\ \Delta_2^\dagger \Delta_1 &= 0, & \Delta_2^\dagger \Gamma_1 &= 0. \end{aligned} \tag{19}$$

Notice that eqs. (19) are satisfied trivially in case $\Gamma_1 = \Delta_1 = 0$ or in case $\Gamma_2 = \Delta_2 = 0$, which correspond to 2HDM of types I or X; note, however, that this kind of models are not of the gBGL type. Coming back to eqs. (19), it is straightforward to show that they are necessary conditions for gBGL models since, from eq. (16),

$$\Gamma_2^\dagger \Gamma_1 = (P_3 \Gamma_2)^\dagger \Gamma_1 = (\Gamma_2^\dagger P_3) \Gamma_1 = \Gamma_2^\dagger (P_3 \Gamma_1) = 0, \tag{20}$$

and similarly for the remaining conditions in eq. (19). The sufficiency of these conditions in order to have gBGL models is shown in appendix A, where the relation with 2HDM of type I is also analysed.

4 Parametrisation of gBGL models

It is clear that gBGL models have a great reduction in the number of free parameters, with respect to the general 2HDM. In this section, we use projection operators to suggest some convenient parametrisations of gBGL models.

4.1 Weak basis invariant projectors

Let us recall that under a WB transformation we have

$$Q_L^0 \mapsto Q_L^{0'} = W_L Q_L^0; \quad d_R^0 \mapsto d_R^{0'} = W_{d_R} d_R^0; \quad u_R^0 \mapsto u_R^{0'} = W_{u_R} u_R^0, \tag{21}$$

and

$$\Gamma_i \mapsto \Gamma'_i = W_L^\dagger \Gamma_i W_{d_R}; \quad \Delta_i \mapsto \Delta'_i = W_L^\dagger \Delta_i W_{u_R}, \tag{22}$$

with W_L , W_{d_R} and W_{u_R} unitary matrices.

If we now take the gBGL definition through projectors in eq. (16) for Γ_i and Δ_i , and go to an arbitrary weak basis, we have

$$\begin{aligned} W_L^\dagger P_3 W_L W_L^\dagger \Gamma_1 W_{d_R} &= 0, & W_L^\dagger P_3 W_L W_L^\dagger \Gamma_2 W_{d_R} &= W_L^\dagger \Gamma_2 W_{d_R}, \\ W_L^\dagger P_3 W_L W_L^\dagger \Delta_1 W_{u_R} &= 0, & W_L^\dagger P_3 W_L W_L^\dagger \Delta_2 W_{u_R} &= W_L^\dagger \Delta_2 W_{u_R}. \end{aligned} \tag{23}$$

These equations are valid for any weak basis. Introducing the projector

$$P_3^{[W_L]} = W_L^\dagger P_3 W_L, \quad (24)$$

we have

$$\begin{aligned} P_3^{[W_L]} \Gamma'_1 &= 0, & P_3^{[W_L]} \Gamma'_2 &= \Gamma'_2, \\ P_3^{[W_L]} \Delta'_1 &= 0, & P_3^{[W_L]} \Delta'_2 &= \Delta'_2, \end{aligned} \quad (25)$$

for an arbitrary WB (from now on, we drop the primes). If we choose, for convenience,

$$W_L^\dagger = \mathcal{U}_L^{u\dagger} \mathcal{U}, \quad (26)$$

where $\mathcal{U}_L^{u\dagger}$ appears in the usual bi-diagonalisation $\mathcal{U}_L^{u\dagger} M_u^0 \mathcal{U}_R^u = M_u$ (see section 4.3), then \mathcal{U} is an arbitrary unitary matrix and we have a general WB invariant parametrisation in terms of that arbitrary \mathcal{U} and of $\mathcal{U}_L^{u\dagger}$; eqs. (24) and (25) read

$$P_{\mathcal{U}3}^{[u_L]} = \mathcal{U}_L^u (\mathcal{U} P_3 \mathcal{U}^\dagger) \mathcal{U}_L^{u\dagger}, \quad (27)$$

and

$$\begin{aligned} P_{\mathcal{U}3}^{[u_L]} \Gamma_1 &= 0, & P_{\mathcal{U}3}^{[u_L]} \Gamma_2 &= \Gamma_2, \\ P_{\mathcal{U}3}^{[u_L]} \Delta_1 &= 0, & P_{\mathcal{U}3}^{[u_L]} \Delta_2 &= \Delta_2. \end{aligned} \quad (28)$$

Notice that this \mathcal{U}_L^u dependence reminds us of uBGL models. An alternative parametrisation is obtained taking

$$W_L^\dagger = \mathcal{U}_L^d \mathcal{U}', \quad (29)$$

where in this case $\mathcal{U}_L^{d\dagger}$ comes from the down mass matrix diagonalisation $\mathcal{U}_L^{d\dagger} M_d^0 \mathcal{U}_R^d = M_d$, with

$$P_{\mathcal{U}'3}^{[d_L]} = \mathcal{U}_L^d (\mathcal{U}' P_3 \mathcal{U}'^\dagger) \mathcal{U}_L^{d\dagger}, \quad (30)$$

and

$$P_{\mathcal{U}'3}^{[d_L]} \Gamma_1 = 0, \quad P_{\mathcal{U}'3}^{[d_L]} \Gamma_2 = \Gamma_2, \quad (31)$$

$$P_{\mathcal{U}'3}^{[d_L]} \Delta_1 = 0, \quad P_{\mathcal{U}'3}^{[d_L]} \Delta_2 = \Delta_2, \quad (32)$$

where now the \mathcal{U}_L^d dependence reminds us of dBGL models. Identifying eqs. (26) and (29),

$$W_L^\dagger = \mathcal{U}_L^u \mathcal{U} = \mathcal{U}_L^d \mathcal{U}', \quad (33)$$

and it is straightforward to conclude that one can use equivalently one or the other parametrisation of the same model provided

$$V^\dagger = \mathcal{U}_L^{d\dagger} \mathcal{U}_L^u = \mathcal{U}' \mathcal{U}^\dagger, \quad \text{i.e. } \mathcal{U} \mathcal{U}'^\dagger = V. \quad (34)$$

4.2 Weak basis covariant parametrisation

We have in the general 2HDM

$$N_d^0 = t_\beta M_d^0 - (t_\beta + t_\beta^{-1}) \frac{v}{\sqrt{2}} e^{i\theta} \Gamma_2, \quad (35)$$

$$N_u^0 = t_\beta M_u^0 - (t_\beta + t_\beta^{-1}) \frac{v}{\sqrt{2}} e^{-i\theta} \Delta_2. \quad (36)$$

Assuming the existence of the projection operator $P_{X3}^{[q_L]}$ satisfying

$$P_{X3}^{[q_L]} \Gamma_1 = 0, \quad P_{X3}^{[q_L]} \Gamma_2 = \Gamma_2, \quad (37)$$

$$P_{X3}^{[q_L]} \Delta_1 = 0, \quad P_{X3}^{[q_L]} \Delta_2 = \Delta_2, \quad (38)$$

with X and $P_{X3}^{[q_L]}$ to be specified later,

$$P_{X3}^{[q_L]} M_d^0 = P_{X3}^{[q_L]} \frac{v}{\sqrt{2}} (c_\beta \Gamma_1 + e^{i\theta} s_\beta \Gamma_2) = \frac{ve^{i\theta}}{\sqrt{2}} s_\beta P_{X3}^{[q_L]} \Gamma_2 = \frac{ve^{i\theta}}{\sqrt{2}} s_\beta \Gamma_2, \quad (39)$$

$$P_{X3}^{[q_L]} M_u^0 = P_{X3}^{[q_L]} \frac{v}{\sqrt{2}} (c_\beta \Delta_1 + e^{-i\theta} s_\beta \Delta_2) = \frac{ve^{-i\theta}}{\sqrt{2}} s_\beta P_{X3}^{[q_L]} \Delta_2 = \frac{ve^{-i\theta}}{\sqrt{2}} s_\beta \Delta_2. \quad (40)$$

Equations (35) and (36) can then be rewritten as the general WB covariant gBGL parametrisation

$$N_d^0 = \left[t_\beta \mathbf{1} - (t_\beta + t_\beta^{-1}) P_{X3}^{[q_L]} \right] M_d^0, \quad (41)$$

$$N_u^0 = \left[t_\beta \mathbf{1} - (t_\beta + t_\beta^{-1}) P_{X3}^{[q_L]} \right] M_u^0. \quad (42)$$

We can choose the up parametrisation for $P_{X3}^{[q_L]}$,

$$P_{X3}^{[q_L]} = P_{\mathcal{U}3}^{[u_L]} = \mathcal{U}_L^u (\mathcal{U} P_3 \mathcal{U}^\dagger) \mathcal{U}_L^{u\dagger}, \quad (43)$$

or, equivalently, we can choose the down parametrisation for $P_{X3}^{[q_L]}$,

$$P_{X3}^{[q_L]} = P_{\mathcal{U}'3}^{[d_L]} = \mathcal{U}_L^d (\mathcal{U}' P_3 \mathcal{U}'^\dagger) \mathcal{U}_L^{d\dagger}, \quad (44)$$

with $V = \mathcal{U}\mathcal{U}^\dagger$, to completely define the model. Let us analyse the details of these parametrisations, after we finally rotate the quark fields to the mass basis.

4.3 Parametrisations in the quark mass basis

Quark fields are rotated in the following manner

$$u_L^0 = \mathcal{U}_L^u u_L, \quad u_R^0 = \mathcal{U}_R^u u_R, \quad d_L^0 = \mathcal{U}_L^d d_L, \quad d_R^0 = \mathcal{U}_R^d d_R, \quad (45)$$

with unitary \mathcal{U}_X^q ($q = u, d$, $X = L, R$), such that

$$\mathcal{U}_L^{u\dagger} M_u^0 \mathcal{U}_R^u = M_u = \text{diag}(m_u, m_c, m_t), \quad \mathcal{U}_L^{d\dagger} M_d^0 \mathcal{U}_R^d = M_d = \text{diag}(m_d, m_s, m_b). \quad (46)$$

N_u^0 and N_d^0 are transformed accordingly,

$$N_u^0 \mapsto N_u = \mathcal{U}_L^{u\dagger} N_u^0 \mathcal{U}_R^u, \quad N_d^0 \mapsto N_d = \mathcal{U}_L^{d\dagger} N_d^0 \mathcal{U}_R^d, \quad (47)$$

and, following eqs. (41)–(42),

$$N_d = \left[t_\beta \mathbf{1} - (t_\beta + t_\beta^{-1}) \mathcal{U}_L^{d\dagger} P_{X3}^{[q_L]} \mathcal{U}_L^d \right] M_d, \quad (48)$$

$$N_u = \left[t_\beta \mathbf{1} - (t_\beta + t_\beta^{-1}) \mathcal{U}_L^{u\dagger} P_{X3}^{[q_L]} \mathcal{U}_L^u \right] M_u. \quad (49)$$

If we choose the down parametrisation in eq. (44), eqs. (48)–(49) give

$$N_d = \left[t_\beta \mathbf{1} - (t_\beta + t_\beta^{-1}) \mathcal{U}' P_3 \mathcal{U}'^\dagger \right] M_d, \quad (50)$$

$$N_u = \left[t_\beta \mathbf{1} - (t_\beta + t_\beta^{-1}) V \mathcal{U}' P_3 \mathcal{U}'^\dagger V^\dagger \right] M_u, \quad (51)$$

while, if we choose the up parametrisation in eq. (43), eqs. (48)–(49) give

$$N_d = \left[t_\beta \mathbf{1} - (t_\beta + t_\beta^{-1}) V^\dagger \mathcal{U} P_3 \mathcal{U}^\dagger V \right] M_d, \quad (52)$$

$$N_u = \left[t_\beta \mathbf{1} - (t_\beta + t_\beta^{-1}) \mathcal{U} P_3 \mathcal{U}^\dagger \right] M_u. \quad (53)$$

Equations (50)–(51) and (52)–(53) are of course equivalent, since $V = \mathcal{U} \mathcal{U}'^\dagger$. At this point it is important to discuss a central question: it may appear that there is a large arbitrariness in the definition of the model since there is a completely arbitrary unitary matrix \mathcal{U} involved in eqs. (52)–(53). Nevertheless, there is much less freedom since the quantities involving \mathcal{U} are

$$[\mathcal{U} P_3 \mathcal{U}^\dagger]_{ij} = \mathcal{U}_{i3} \mathcal{U}_{j3}^*, \quad (54)$$

that is, only the elements of the third column of \mathcal{U} , which form a unitary complex vector, are needed to define the model. To stress this fact, we introduce

$$\hat{n}_{[u]i} \equiv \mathcal{U}_{i3}, \quad \text{and} \quad \hat{n}_{[d]i} = \mathcal{U}'_{i3}, \quad \text{with} \quad \hat{n}_{[u]j} = V_{ji} \hat{n}_{[d]i}, \quad (55)$$

where the subindex $[u]$ or $[d]$ specifies the parametrisation under consideration. The matrix elements of N_u and N_d in equations (48) and (49) can be written, explicitly,

$$[N_d]_{ij} = t_\beta \delta_{ij} m_{d_i} - (t_\beta + t_\beta^{-1}) \hat{n}_{[d]i} \hat{n}_{[d]j}^* m_{d_j}, \quad (56)$$

$$[N_u]_{ij} = t_\beta \delta_{ij} m_{u_i} - (t_\beta + t_\beta^{-1}) \hat{n}_{[u]i} \hat{n}_{[u]j}^* m_{u_j}. \quad (57)$$

Since $\hat{n}_{[u]j} = V_{ji} \hat{n}_{[d]i}$, one is free to rewrite eqs. (50)–(51) as

$$[N_d]_{ij} = t_\beta \delta_{ij} m_{d_i} - (t_\beta + t_\beta^{-1}) \hat{n}_{[d]i} \hat{n}_{[d]j}^* m_{d_j}, \quad (58)$$

$$[N_u]_{ij} = t_\beta \delta_{ij} m_{u_i} - (t_\beta + t_\beta^{-1}) V_{ia} V_{jb}^* \hat{n}_{[d]a} \hat{n}_{[d]b}^* m_{u_j}. \quad (59)$$

or eqs. (52)–(53) as

$$[N_d]_{ij} = t_\beta \delta_{ij} m_{d_i} - (t_\beta + t_\beta^{-1}) \hat{n}_{[u]a} \hat{n}_{[u]b}^* V_{ai}^* V_{bj} m_{d_j}, \quad (60)$$

$$[N_u]_{ij} = t_\beta \delta_{ij} m_{u_i} - (t_\beta + t_\beta^{-1}) \hat{n}_{[u]i} \hat{n}_{[u]j}^* m_{u_j}. \quad (61)$$

One can now identify easily all the usual BGL models with this parametrisation. The notation is straightforward, for example, BGL model “s” corresponds to \hat{s} , and so on:

- dBGL models in the down parametrisation,

$$\hat{d}_{[d]} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad \hat{s}_{[d]} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad \hat{b}_{[d]} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad (62)$$

- uBGL models in the up parametrisation,

$$\hat{u}_{[u]} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad \hat{c}_{[u]} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad \hat{t}_{[u]} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad (63)$$

- dBGL models in the up parametrisation,

$$\hat{d}_{[u]} = \begin{pmatrix} V_{ud} \\ V_{cd} \\ V_{td} \end{pmatrix}, \quad \hat{s}_{[u]} = \begin{pmatrix} V_{us} \\ V_{cs} \\ V_{ts} \end{pmatrix}, \quad \hat{b}_{[u]} = \begin{pmatrix} V_{ub} \\ V_{cb} \\ V_{tb} \end{pmatrix}, \quad (64)$$

- uBGL models in the down parametrisation,

$$\hat{u}_{[d]} = \begin{pmatrix} V_{ud}^* \\ V_{us}^* \\ V_{ub}^* \end{pmatrix}, \quad \hat{c}_{[d]} = \begin{pmatrix} V_{cd}^* \\ V_{cs}^* \\ V_{cb}^* \end{pmatrix}, \quad \hat{t}_{[d]} = \begin{pmatrix} V_{td}^* \\ V_{ts}^* \\ V_{tb}^* \end{pmatrix}. \quad (65)$$

It is also possible to give a graphical description of the gBGL class of models, as shown in figure 1. Since a model is defined by a complex unitary vector \hat{n} , $|\hat{n}_1|^2 + |\hat{n}_2|^2 + |\hat{n}_3|^2 = 1$, and $(|\hat{n}_1|, |\hat{n}_2|, |\hat{n}_3|)$ is located on the sphere of unit radius (specifically, on an octant of that sphere). Furthermore, there are two physical complex phases, since one can readily see that $\hat{n}_i \hat{n}_j^*$ is unaffected by a global rephasing of \hat{n} , which can be used to remove one out of the three initial phases in the \hat{n} components. This is illustrated in figure 1(c), where no explicit reference to up, down or other parametrisations is made⁷. Figures 1(a) and 1(b) illustrate the situation for down and up parametrisations, including the usual BGL models in eqs. (62) to (65).


5 The scalar sector

The scalar potential, imposing invariance under $\Phi_2 \mapsto -\Phi_2$ in eq. (10), is


$$V(\Phi_1, \Phi_2) = \mu_{11}^2 \Phi_1^\dagger \Phi_1 + \mu_{22}^2 \Phi_2^\dagger \Phi_2 + \lambda_1 (\Phi_1^\dagger \Phi_1)^2 + \lambda_2 (\Phi_2^\dagger \Phi_2)^2 \\ + 2\lambda_3 (\Phi_1^\dagger \Phi_1)(\Phi_2^\dagger \Phi_2) + 2\lambda_4 (\Phi_1^\dagger \Phi_2)(\Phi_2^\dagger \Phi_1) + [\lambda_5 (\Phi_1^\dagger \Phi_2)^2 + \text{h.c.}]. \quad (66)$$

Obviously, this scalar potential coincides with the one in Glashow-Weinberg model [6], since in both cases a \mathbb{Z}_2 symmetry is introduced. In BGL models where instead


⁷Notice that one can trivially adopt spherical coordinates with, for example, $(|\hat{n}_1|, |\hat{n}_2|, |\hat{n}_3|) = (\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)$.



(a) Down parametrisation.



(b) Up parametrisation.



(c) Generic parametrisation.

Figure 1: gBGL models.

of \mathbb{Z}_2 one uses a larger symmetry, namely \mathbb{Z}_4 , the term in λ_5 is not \mathbb{Z}_4 invariant and therefore cannot be introduced. This leads to a global symmetry which upon spontaneous breaking would lead to a Goldstone boson. This difficulty can be avoided by softly breaking the discrete symmetry through the addition of a term ($m_{12}^2 \Phi_1^\dagger \Phi_2 + \text{h.c.}$). The scalar potential of eq. (66) does not lead to new sources of CP violation. But when the above bilinear term is introduced, one can have either spontaneous [31] or explicit CP breaking in the scalar sector. Expanding

$$\Phi_j = e^{i\theta_j} \begin{pmatrix} \varphi_j^+ \\ (v_j + \rho_j + i\eta_j)/\sqrt{2} \end{pmatrix}, \quad (67)$$

then the rotation to the Higgs basis in eq. (3) identifies the Goldstone boson interpretation of G^\pm and G^0 as the longitudinal degrees of freedom of the W^\pm and Z^0 gauge bosons,

$$H_1 = \begin{pmatrix} G^+ \\ (v + H^0 + iG^0)/\sqrt{2} \end{pmatrix}, \quad H_2 = \begin{pmatrix} H^+ \\ (R^0 + iA)/\sqrt{2} \end{pmatrix}, \quad (68)$$

and

$$\begin{pmatrix} H^0 \\ R^0 \end{pmatrix} = \begin{pmatrix} c_\beta & s_\beta \\ s_\beta & -c_\beta \end{pmatrix} \begin{pmatrix} \rho_1 \\ \rho_2 \end{pmatrix}. \quad (69)$$

The scalar potential in eq. (66) is in general CP conserving in such a way that the physical CP even states do not mix with the CP odd one; then, the physical scalars are a linear combination of ρ_1 and ρ_2 ,

$$\begin{pmatrix} H \\ h \end{pmatrix} = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} \rho_1 \\ \rho_2 \end{pmatrix}, \quad (70)$$

and we will have in general

$$\begin{pmatrix} H^0 \\ R^0 \end{pmatrix} = \begin{pmatrix} c_{\beta\alpha} & s_{\beta\alpha} \\ s_{\beta\alpha} & -c_{\beta\alpha} \end{pmatrix} \begin{pmatrix} H \\ h \end{pmatrix}, \quad (71)$$

where the relevant angle is $\beta - \alpha$ and $c_{\beta\alpha} \equiv \cos(\beta - \alpha)$, $s_{\beta\alpha} \equiv \sin(\beta - \alpha)$. The field A is a physical pseudoscalar. Notice that the Yukawa couplings become

$$\begin{aligned} \mathcal{L}_Y = & -\frac{\sqrt{2}}{v} H^+ \bar{u} (V N_d \gamma_R - N_u^\dagger V \gamma_L) d + \text{h.c.} \\ & -\frac{1}{v} H^0 (\bar{u} M_u u + \bar{d} M_d d) \\ & -\frac{1}{v} R^0 \left[\bar{u} (N_u \gamma_R + N_u^\dagger \gamma_L) u + \bar{d} (N_d \gamma_R + N_d^\dagger \gamma_L) d \right] \\ & +\frac{i}{v} A \left[\bar{u} (N_u \gamma_R - N_u^\dagger \gamma_L) u - \bar{d} (N_d \gamma_R - N_d^\dagger \gamma_L) d \right], \end{aligned} \quad (72)$$

in such a way that, for $h \rightarrow H^0$, h becomes “the standard Higgs”, coupling to fermions like the SM Higgs does. That is, when $s_{\beta\alpha} = 1$ we have h aligned with H^0 with SM couplings.

6 The intensity of FCNC in gBGL

FCNC are extremely constrained by experimental data, consequently we have to worry about their intensity in gBGL models. It is worthwhile to mention that BGL are Minimal Flavour Violating (MFV) models, meaning that FCNC are controlled by the deviation of the CKM matrix V from $\mathbf{1}$. As a result, in the limit $V = \mathbf{1}$, there are no FCNC in BGL models at tree level. However, this is not the case in gBGL models, where FCNC are no longer controlled by the deviations of V from unity. The Yukawa couplings of gBGL models are given, following eqs. (72) and (71), by

$$\mathcal{L}_{h\bar{q}q} = - \sum_{i,j=d,s,b} h\bar{d}_{L_i} Y_{ij}^D d_{R_j} - \sum_{i,j=u,c,t} h\bar{u}_{L_i} Y_{ij}^U u_{R_j} , \quad (73)$$

with

$$\begin{aligned} Y_{ij}^D &= \frac{1}{v} [s_{\beta\alpha}(M_d)_{ij} + c_{\beta\alpha}(N_d)_{ij}] , \\ Y_{ij}^U &= \frac{1}{v} [s_{\beta\alpha}(M_u)_{ij} + c_{\beta\alpha}(N_u)_{ij}] , \end{aligned} \quad (74)$$

with N_q given in eqs. (56)–(57), that generically can be written

$$(N_q)_{ij} = [t_\beta \delta_{ij} - (t_\beta + t_\beta^{-1}) \hat{n}_{[q]i} \hat{n}_{[q]j}^*] m_{q_j} . \quad (75)$$

So, the flavour changing intensities are controlled by N_q with the following factors.

- In an $q_{R_j} \rightarrow q_{L_i}$ and in $q_{L_i} \rightarrow q_{R_j}$ transitions there is a factor m_{q_j}/v . Notice that in $q_{R_i} \rightarrow q_{L_j}$ and in $q_{L_j} \rightarrow q_{R_i}$ transitions the factor is instead m_{q_i}/v . That is, in general, for a $q_j \rightarrow q_i$ vertex, there is a suppression factor given by the heaviest of the two quarks mass, $\max(m_{q_i}, m_{q_j})$. This suppression factor is a very relevant one except, obviously, for any transition where the top quark is present.
- A factor $c_{\beta\alpha}(t_\beta + t_\beta^{-1})$; from perturbative unitarity requirements on the scalar sector, is constrained to be at most one [26, 32–35]. Notice that in the limit $c_{\beta\alpha} \rightarrow 0$, the FCNC associated to the standard Higgs h disappear, and the ones associated to the remaining scalars can be suppressed by making them heavier. Therefore, this is the quantity which, in a global approach, is bounded from above by Higgs mediated FCNC processes.
- Finally, there is the factor $\hat{n}_{[q]i} \hat{n}_{[q]j}^*$, which ranges from 0 to 1/2 (which is only reached when only *one* transition is allowed). We remind that in BGL models, the analogous factor is $V_{iq} V_{jq}^*$. With the naive bounds on $c_{\beta\alpha}(t_\beta + t_\beta^{-1})$ from meson mixings in [26], one can compare and analyse how the suppression factors change in these new gBGL models. We collect the suppression factors for the different transitions in all BGL models in table 1 (with the corresponding power counting in Wolfenstein's parameter λ). One has to compare the maximum value of $|\hat{n}_{[q]i} \hat{n}_{[q]j}^*|_{\max} = 1/2$ with λ , which corresponds to the less suppressed transition in some BGL models. This means that the most stringent constraint on BGL

Model Transition	d	s	b
$u \leftrightarrow c$	$V_{ud}V_{cd}^* \sim \lambda$	$V_{us}V_{cs}^* \sim \lambda$	$V_{ub}V_{cb}^* \sim \lambda^5$
$u \leftrightarrow t$	$V_{ud}V_{td}^* \sim \lambda^3$	$V_{us}V_{ts}^* \sim \lambda^3$	$V_{ub}V_{tb}^* \sim \lambda^3$
$c \leftrightarrow t$	$V_{cd}V_{td}^* \sim \lambda^4$	$V_{cs}V_{ts}^* \sim \lambda^2$	$V_{cb}V_{tb}^* \sim \lambda^2$
Model Transition	u	c	t
$d \leftrightarrow s$	$V_{ud}^*V_{us} \sim \lambda$	$V_{cd}^*V_{cs} \sim \lambda$	$V_{td}^*V_{ts} \sim \lambda^5$
$d \leftrightarrow b$	$V_{ud}^*V_{ub} \sim \lambda^3$	$V_{cd}^*V_{cb} \sim \lambda^3$	$V_{td}^*V_{tb} \sim \lambda^3$
$s \leftrightarrow b$	$V_{us}^*V_{ub} \sim \lambda^4$	$V_{cs}^*V_{cb} \sim \lambda^2$	$V_{ts}^*V_{tb} \sim \lambda^2$

Table 1: CKM suppression in FCNC transitions in BGL models.

models obtained in [26] for $|c_{\beta\alpha}(t_\beta + t_\beta^{-1})|$ should be reduced by a factor 2λ . In this way, taking into account the analysis of [26] and the constraints in [36], one can conclude that the gBGL models are safe, over the entire parameter space, provided

$$|c_{\beta\alpha}(t_\beta + t_\beta^{-1})| \leq 0.02. \quad (76)$$

This constraint arises from $D^0-\bar{D}^0$ mixing and it turns out to be more stringent than the $K^0-\bar{K}^0$ one. It is worth mentioning that in some regions of the parameter space, this constraint will be relaxed. For example, in all BGL models this constraint is much weaker as shown in reference [26], and in some of them $|c_{\beta\alpha}(t_\beta + t_\beta^{-1})|$ can span the entire theoretically allowed parameter space, arriving to values of order 1.

A final remark is related to the absence of the mass suppression factor on $t \leftrightarrow q$ transitions in gBGL models. In this case, the relevant bounds come from rare top decays $t \rightarrow hu, hc$, which give [26]

$$\text{BR}(t \rightarrow hq) = 0.13 \frac{|\hat{n}_{[u]q}\hat{n}_{[u]t}^*|^2}{|V_{tb}|^2} (c_{\beta\alpha}(t_\beta + t_\beta^{-1}))^2. \quad (77)$$

Considering the experimental bounds from ATLAS [37] and CMS [38,39], this yields

$$|c_{\beta\alpha}(t_\beta + t_\beta^{-1})| \leq 0.4, \quad (78)$$

for the maximal value⁸ $|\hat{n}_{[u]q}\hat{n}_{[u]t}^*| = 1/2$.

We conclude that the whole class of gBGL models does not lead to much larger flavour changing transitions than the ones arising in BGL models. This is in spite of having simultaneously FCNC in the up and in the down sectors. In the next sections we discuss other important differences.

⁸The maximal value of $|\hat{n}_{[u]q}\hat{n}_{[u]t}^*|$ cannot be obtained, obviously, for both $t \rightarrow hu$ and $t \rightarrow hc$ simultaneously.

7 Near the top and the bottom models

BGL top and bottom models are the only renormalizable 2HDM that verify the MFV principle in any of the different versions one can find in the literature [21]. We devote this section to analyse in more detail the properties of gBGL models that are “close” to these models, that is, that they depart from the top or bottom models by a “small amount”. Let us recall that the FCNC in the down and the up sectors of the top model are controlled by

$$(N_d)_{ij} = (t_\beta + t_\beta^{-1}) \hat{t}_{[d]i} \hat{t}_{[d]j}^* m_{d_j}, \quad (N_u)_{ij} = (t_\beta + t_\beta^{-1}) \hat{t}_{[u]i} \hat{t}_{[u]j}^* m_{u_j}, \quad (79)$$

and, following eqs. (63)–(65), $\hat{t}_{[u]}$ and $\hat{t}_{[d]} = V^\dagger \hat{t}_{[u]}$, are

$$\hat{t}_{[u]} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad \hat{t}_{[d]} = \begin{pmatrix} V_{td}^* \\ V_{ts}^* \\ V_{tb}^* \end{pmatrix}. \quad (80)$$

It is clear that the BGL top model does not have FCNC in the up sector. Let us now consider small deviations from $\hat{t}_{[d]}$ near the top model parameterised in terms of a complex vector $\vec{\delta} = (\delta_d, \delta_s, \delta_b)$ (with the appropriate normalisation):

$$\hat{t}_{[d]} + \hat{\delta} \hat{t}_{[d]} \sim \begin{pmatrix} V_{td}^*(1 + \delta_d) \\ V_{ts}^*(1 + \delta_s) \\ V_{tb}^*(1 + \delta_b) \end{pmatrix}. \quad (81)$$

The elements of $\hat{t}_{[d]} + \hat{\delta} \hat{t}_{[d]}$ control the flavour structure of the New Physics contributions to $K^0 - \bar{K}^0$, $B_d^0 - \bar{B}_d^0$ and $B_s^0 - \bar{B}_s^0$. In particular, the leading order contributions to the different meson mixings M_{12} have the following form:

$$\begin{aligned} M_{12}[K^0] &\propto (V_{td}^* V_{ts})^2 [1 + 2(\delta_s^* + \delta_d)], \\ M_{12}[B_d^0] &\propto (V_{td}^* V_{tb})^2 [1 + 2(\delta_b^* + \delta_d)], \\ M_{12}[B_s^0] &\propto (V_{ts}^* V_{tb})^2 [1 + 2(\delta_b^* + \delta_s)]. \end{aligned} \quad (82)$$

Therefore, taking into account the phases of the dominant terms $(V_{ti}^* V_{tj})^2$, we can conclude that these “near top” models will give the same contribution to meson mixings provided

$$\text{Re}(\delta_d) \sim \text{Re}(\delta_s) \sim \text{Im}(\delta_s) \leq \mathcal{O}(\lambda^2), \text{ and } \text{Im}(\delta_d) \sim \text{Im}(\delta_b) \leq \mathcal{O}(\lambda^3). \quad (83)$$

Notice that we are not stating that these models do not have any contribution to the meson mixings, the point is, rather, that these models are “like the top BGL” in the sense that they give the same contributions to $K^0 - \bar{K}^0$, $B_d^0 - \bar{B}_d^0$ and $B_s^0 - \bar{B}_s^0$ mixings. The immediate question is then: do these models produce too strong FCNC in the up sector, in particular in $D^0 - \bar{D}^0$ or in top decays? FCNC in the up sector are controlled by

$$\hat{t}_{[u]} + \hat{\delta} \hat{t}_{[u]} = V^\dagger (\hat{t}_{[d]} + \hat{\delta} \hat{t}_{[d]}) \sim \begin{pmatrix} \mathcal{O}(\lambda^5) \\ \delta_b V_{cb} V_{tb}^* \\ 1 + \delta_b \end{pmatrix}. \quad (84)$$

It is clear that $M_{12}[D^0]$ will have a suppression given by

$$(\delta_b \lambda^5 V_{cb} V_{tb}^*)^2 \leq \lambda^{18}, \quad (85)$$

much smaller than any of the contributions in dBGL models.

To analyse $t \rightarrow hc$ we have to compare the maximal value $1/4$ of $|\hat{n}_{[u]q} \hat{n}_{[u]t}^*|^2$ in eq. (77) with the value obtained in the present case, $|(1 + \delta_b) \delta_b V_{cb} V_{tb}^*|^2 \sim \mathcal{O}(\lambda^8)$. The conclusion is evident in the whole parameter space: these models will produce $t \rightarrow hc$ still below the actual experimental bounds. The same conclusion applies to $t \rightarrow hu$. In the next section we will see that these models can depart in a sizable way from the top BGL model in different physical observables.

The bottom dBGL model is specified by \hat{b} (eqs. (62) and (64)):

$$\hat{b}_{[d]} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad \hat{b}_{[u]} = \begin{pmatrix} V_{ub} \\ V_{cb} \\ V_{tb} \end{pmatrix}. \quad (86)$$

This model generates FCNC in the up sector, and in particular it contributes to $D^0 - \bar{D}^0$ mixing but not to $K^0 - \bar{K}^0$, $B_d^0 - \bar{B}_d^0$ and $B_s^0 - \bar{B}_s^0$; gBGL models close to the \hat{b} model that keep its essential properties are obtained with

$$\hat{b}_{[u]} + \delta \hat{b}_{[u]} \sim \begin{pmatrix} V_{ub}(1 + \delta_u) \\ V_{cb}(1 + \delta_c) \\ V_{tb}(1 + \delta_t) \end{pmatrix}, \quad (87)$$

where

$$\delta_u \sim \delta_c \sim \delta_t \leq \mathcal{O}(\lambda^2), \quad (88)$$

as before. To see what happens with the important constraints in the down sector, we show, as before, that $\hat{b}_{[d]} + \delta \hat{b}_{[d]} = V(\hat{b}_{[u]} + \delta \hat{b}_{[u]})$,

$$\hat{b}_{[d]} + \delta \hat{b}_{[d]} \sim \begin{pmatrix} V_{td}^* V_{tb}(\delta_t - \delta_c) + V_{ud}^* V_{ub}(\delta_u - \delta_c) \\ V_{ts}^* V_{tb}(\delta_t - \delta_c) \\ |V_{tb}|^2(1 + \delta_t) \end{pmatrix} \quad (89)$$

is the relevant quantity. With values as in eq. (88), it turns out that the contributions to the mixing in the down sector are much smaller than in any uBGL model. Nevertheless, we will also see, in the next section, that there are important differences with respect to the bottom dBGL model in other observables while considering the same kind of parameter values close to the bottom model.

8 Baryon Asymmetry of the Universe

8.1 The leading gBGL contribution

The presence of additional sources of flavour and CP violation in the gBGL models can enhance the contribution to the Baryonic Asymmetry of the Universe (BAU) with

respect to SM expectations. Having N_u and N_d in addition to M_u and M_d , a weak basis invariant with a non-zero imaginary part already arises [40] at the 4th order^{9,10}:

$$\text{Im} \left(\text{Tr} \left\{ N_d^0 M_d^{0\dagger} M_u^0 M_u^{0\dagger} \right\} \right). \quad (90)$$

Considering N_d^0 in eq. (41),

$$\text{Im} \left(\text{Tr} \left\{ N_d^0 M_d^{0\dagger} M_u^0 M_u^{0\dagger} \right\} \right) = (t_\beta + t_\beta^{-1}) \text{Im} \left(\text{Tr} \left\{ P_{X3}^{[qL]} M_d^0 M_d^{0\dagger} M_u^0 M_u^{0\dagger} \right\} \right). \quad (91)$$

For $P_{X3}^{[qL]} = P_{\mathcal{U}'3}^{[dL]}$, that is, in the down parametrisation, eq. (44),

$$(t_\beta + t_\beta^{-1}) \text{Im} \left(\text{Tr} \left\{ \mathcal{U}_L^d (\mathcal{U}' P_3 \mathcal{U}'^\dagger) \mathcal{U}_L^{d\dagger} \mathcal{U}_L^d M_d M_d^\dagger \mathcal{U}_L^{d\dagger} \mathcal{U}_L^u M_u M_u^\dagger \mathcal{U}_L^{u\dagger} \right\} \right) = \\ (t_\beta + t_\beta^{-1}) \text{Im} \left(\text{Tr} \left\{ (\mathcal{U}' P_3 \mathcal{U}'^\dagger) M_d M_d^\dagger V^\dagger M_u M_u^\dagger V \right\} \right). \quad (92)$$

Notice that, according to eq. (46), $M_u M_u^\dagger = \text{diag}(m_{u_j}^2) = \text{diag}(m_u^2, m_c^2, m_t^2)$, $M_d M_d^\dagger = \text{diag}(m_{d_j}^2) = \text{diag}(m_d^2, m_s^2, m_b^2)$. Following eq. (55),

$$(\mathcal{U}' P_3 \mathcal{U}'^\dagger)_{ij} = \hat{n}_{[d]i} \hat{n}_{[d]j}^*, \quad (93)$$

and thus

$$\text{Tr} \left\{ (\mathcal{U}' P_3 \mathcal{U}'^\dagger) M_d M_d^\dagger V^\dagger M_u M_u^\dagger V \right\} = \sum_{i,j,k} m_{d_j}^2 m_{u_k}^2 \hat{n}_{[d]i} \hat{n}_{[d]j}^* V_{ki} V_{kj}^*, \quad (94)$$

to obtain, finally

$$\text{Im} \left(\text{Tr} \left\{ N_d^0 M_d^{0\dagger} M_u^0 M_u^{0\dagger} \right\} \right) = \frac{i}{2} (t_\beta + t_\beta^{-1}) \sum_{i,j,k} (m_{d_i}^2 - m_{d_j}^2) m_{u_k}^2 \hat{n}_{[d]i} \hat{n}_{[d]j}^* V_{ki} V_{kj}^*. \quad (95)$$

Although the discussion has relied on the use of the down parametrisation (from eq. (92) onwards), completely analogous results are obtained if the up parametrisation is used instead.

In order to estimate the enhancement in the BAU due to $\text{Im} \left(\text{Tr} \left\{ N_d^0 M_d^{0\dagger} M_u^0 M_u^{0\dagger} \right\} \right)$, let us first retain only potentially leading contributions in terms of masses and powers of the Wolfenstein parameter λ [43],

$$\text{Im} \left(\text{Tr} \left\{ N_d^0 M_d^{0\dagger} M_u^0 M_u^{0\dagger} \right\} \right) \simeq \frac{i}{2} (t_\beta + t_\beta^{-1}) \left\{ \hat{n}_{[d]2} \hat{n}_{[d]1}^* m_s^2 [m_c^2 V_{cs} V_{cd}^* + m_t^2 V_{ts} V_{td}^*] \right. \\ \left. + \hat{n}_{[d]3} \hat{n}_{[d]1}^* m_b^2 [m_c^2 V_{cb} V_{cd}^* + m_t^2 V_{tb} V_{td}^*] + \hat{n}_{[d]3} \hat{n}_{[d]2}^* m_b^2 [m_c^2 V_{cb} V_{cs}^* + m_t^2 V_{tb} V_{ts}^*] \right\} \quad (96)$$

⁹To be compared with the 12th order usual one, $-\frac{i}{6} \text{Tr} \left\{ [M_u M_u^\dagger, M_d M_d^\dagger]^3 \right\}$ [41].

¹⁰The rephasing invariance in the Higgs sector [29] imposes that the complete invariant should include the $H_2^\dagger H_1$ coefficient in the Lagrangian: $(\mu_{11}^2 - \mu_{22}^2) s_\beta c_\beta$ [42]. This does not introduce new phases.

that is,

$$\begin{aligned} \text{Im} \left(\text{Tr} \left\{ N_d^0 M_d^{0\dagger} M_u^0 M_u^{0\dagger} \right\} \right) &\sim i(t_\beta + t_\beta^{-1}) \left\{ \hat{n}_{[d]2} \hat{n}_{[d]1}^* m_s^2 [m_c^2 \lambda + m_t^2 \lambda^5] \right. \\ &\quad \left. + \hat{n}_{[d]3} \hat{n}_{[d]1}^* m_b^2 [m_c^2 \lambda^3 + m_t^2 \lambda^3] + \hat{n}_{[d]3} \hat{n}_{[d]2}^* m_b^2 [m_c^2 \lambda^2 + m_t^2 \lambda^2] \right\}. \end{aligned} \quad (97)$$

In the SM, the BAU is proportional to [44, 45]

$$\text{BAU}_{\text{SM}} \sim \frac{m_t^4 m_b^4 m_c^2 m_s^2}{E^{12}} J, \quad (98)$$

where J is the rephasing invariant imaginary part of the CKM quartets [46], $J = \text{Im}(V_{cs}^* V_{ts} V_{tb}^* V_{cb}) \simeq 3 \times 10^{-5}$ and $E \sim 100$ GeV an energy of the order of the electroweak scale one. In eq. (97), we have contributions like the last one, giving

$$\text{BAU}_{\text{gBGL}} \sim (t_\beta + t_\beta^{-1}) \frac{m_t^2 m_b^2}{E^4} \text{Im}(\hat{n}_{[d]3} \hat{n}_{[d]2}^* V_{tb} V_{ts}^*). \quad (99)$$

This simple enhancement estimate with respect to eq. (98), in terms of $\alpha = \arg(\hat{n}_{[d]3} \hat{n}_{[d]2}^* V_{tb} V_{ts}^*)$ and $|\hat{n}_{[d]3} \hat{n}_{[d]2}^*|$ (which has a maximal value of 1/2), is:

$$\begin{aligned} \frac{\text{BAU}_{\text{gBGL}}}{\text{BAU}_{\text{SM}}} &\sim (t_\beta + t_\beta^{-1}) |\hat{n}_{[d]3} \hat{n}_{[d]2}^*| \sin \alpha \frac{|V_{ts}|}{J} \frac{E^8}{m_t^2 m_b^2 m_c^2 m_s^2} \\ &\sim 10^{16} (t_\beta + t_\beta^{-1}) |\hat{n}_{[d]3} \hat{n}_{[d]2}^*| \sin \alpha, \end{aligned} \quad (100)$$

showing that there is margin for substantial enhancement of the BAU with respect to the SM, and with respect to BGL models too [40]. In appendix B it is shown that we do not expect relevant constraints coming from electric dipole moments (EDM).

8.2 The vanishing BGL limits

In BGL models, the previous contribution is vanishing: let us explicitly check this known result [40]. For dBGL models, with \hat{n}_d in eq. (62), only one component is non-vanishing and $\text{Im} \left(\text{Tr} \left\{ N_d^0 M_d^{0\dagger} M_u^0 M_u^{0\dagger} \right\} \right) = 0$. The situation is slightly more involved for uBGL models. From eq. (65), for an uBGL model of type q , $\hat{n}_{[d]i} = V_{qi}^*$. Then, going back to eq. (94),

$$\text{Tr} \left\{ (\mathcal{U}' P_3 \mathcal{U}^{\dagger}) M_d M_d^\dagger V^\dagger M_u M_u^\dagger V \right\} = \sum_{i,j,k} m_{d_j}^2 m_{u_k}^2 V_{qi}^* V_{qj} V_{ki} V_{kj}^*, \quad (\text{uBGL } q). \quad (101)$$

Since, from unitarity of V , $\sum_i V_{qi}^* V_{ki} = \delta_{qk}$, eq. (101) gives

$$\text{Tr} \left\{ (\mathcal{U}' P_3 \mathcal{U}^{\dagger}) M_d M_d^\dagger V^\dagger M_u M_u^\dagger V \right\} = \sum_j m_{d_j}^2 m_{u_q}^2 |V_{qj}|^2, \quad (\text{uBGL } q), \quad (102)$$

and thus $\text{Im} \left(\text{Tr} \left\{ N_d^0 M_d^{0\dagger} M_u^0 M_u^{0\dagger} \right\} \right) = 0$.

8.3 Rephasing invariance

Although we have already mentioned that there are only two physical phases in $\hat{n}_{[d]}$ or $\hat{n}_{[u]}$, it is important to check that the invariant in eq. (95) is, at it should, invariant under individual rephasings of the different quark fields:

$$\begin{aligned} d &\mapsto e^{i\varphi_1} d, & s &\mapsto e^{i\varphi_2} s, & b &\mapsto e^{i\varphi_3} b, \\ V_{qd} &\mapsto e^{i\varphi_1} V_{qd}, & V_{qs} &\mapsto e^{i\varphi_2} V_{qs}, & V_{qb} &\mapsto e^{i\varphi_3} V_{qb}. \end{aligned} \quad (103)$$

The origin of the rephasing of the CKM matrix is straightforward: since $V = \mathcal{U}_L^{u\dagger} \mathcal{U}_L^d$, the rephasing of the down quarks corresponds to

$$\mathcal{U}_L^d \mapsto \mathcal{U}_L^d \begin{pmatrix} e^{i\varphi_1} & 0 & 0 \\ 0 & e^{i\varphi_2} & 0 \\ 0 & 0 & e^{i\varphi_3} \end{pmatrix}. \quad (104)$$

Then, since in eq. (92) $P_{\mathcal{U}'3}^{[d_L]} = \mathcal{U}_L^d (\mathcal{U}' P_3 \mathcal{U}'^\dagger) \mathcal{U}_L^{d\dagger}$,

$$\mathcal{U}' \mapsto \begin{pmatrix} e^{-i\varphi_1} & 0 & 0 \\ 0 & e^{-i\varphi_2} & 0 \\ 0 & 0 & e^{-i\varphi_3} \end{pmatrix} \mathcal{U}' \quad (105)$$

to keep $\mathcal{U}_L^d \mathcal{U}'$ invariant; as a consequence, under the rephasing in eq. (103),

$$\hat{n}_{[d]j} \mapsto e^{-i\varphi_j} \hat{n}_{[d]j}, \quad (106)$$

and eq. (95) is clearly rephasing invariant.

8.4 Enhancements in models near the top and the bottom models

The BAU generated in the top BGL model is proportional to [40]

$$\begin{aligned} &\text{Im} \left(\text{Tr} \left\{ M_d N_d^\dagger M_d M_d^\dagger M_u M_u^\dagger M_d M_d^\dagger \right\} \right) \\ &= -(t_\beta + t_\beta^{-1})(m_b^2 - m_s^2)(m_b^2 - m_d^2)(m_s^2 - m_d^2)(m_c^2 - m_u^2) \text{Im}(V_{cs}^* V_{ts} V_{tb}^* V_{cb}), \end{aligned} \quad (107)$$

in such a way that the ratio to the SM BAU is, for $E \sim 100$ GeV,

$$\frac{\text{BAU}_{\text{BGL-t}}}{\text{BAU}_{\text{SM}}} = (t_\beta + t_\beta^{-1}) \frac{E^4}{m_t^4} \sim 1. \quad (108)$$

Therefore, as far as CP violation is concerned, the top model suffers the same problem as the SM in not being able to generate sufficient BAU. There is no enhancement in the top BGL 2HDM. We can now consider a small departure from the top BGL model as in eq. (81) and apply it to eq. (100). With

$$\hat{n}_{[d]3}^* \sim V_{tb}^*(1 + \delta_b), \quad \hat{n}_{[d]2}^* \sim V_{ts}(1 + \delta_s^*), \quad (109)$$

$$\text{Im} \left(\hat{n}_{[d]2}^* \hat{n}_{[d]3}^* V_{tb} V_{ts}^* \right) \sim |V_{ts}|^2 \text{Im} (\delta_b + \delta_s^*) , \quad (110)$$

and eq. (81) gives

$$\frac{\text{BAU}_{\text{near t}}}{\text{BAU}_{\text{SM}}} = 10^{16} (t_\beta + t_\beta^{-1}) |V_{ts}| \text{Im} (\delta_b + \delta_s^*) , \quad (111)$$

that can produce an enhancement as large as 10^{12} . Even if we are close to a top BGL model we find that, contrary to what happens with top model itself, there can be enough CP violation in this class of models to generate the BAU. In an analogous way, the BAU generated in the bottom BGL model is proportional to

$$\begin{aligned} & \text{Im} \left(\text{Tr} \left\{ M_u N_u^\dagger M_u M_u^\dagger M_d M_d^\dagger M_u M_u^\dagger \right\} \right) \\ &= -(t_\beta + t_\beta^{-1}) (m_t^2 - m_c^2) (m_t^2 - m_u^2) (m_c^2 - m_u^2) (m_s^2 - m_d^2) \text{Im} (V_{cs}^* V_{ts} V_{tb}^* V_{cb}) , \end{aligned} \quad (112)$$

and the ratio to the SM BAU, for $E \sim 100$ GeV, is

$$\frac{\text{BAU}_{\text{BGL-b}}}{\text{BAU}_{\text{SM}}} = (t_\beta + t_\beta^{-1}) \frac{E^4}{m_b^4} \sim 10^5 . \quad (113)$$

In a pure bottom model there is still not enough CP violation – coming from the Yukawa sector – to generate the BAU. But, if we depart from an exact bottom BGL model in the manner explained in eq. (87), with

$$\hat{n}_{[d]3} \sim |V_{tb}|^2 (1 + \delta_t) , \quad \hat{n}_{[d]2}^* \sim V_{ts} V_{tb}^* (\delta_t^* - \delta_c^*) , \quad (114)$$

then

$$\text{Im} \left(\hat{n}_{[d]3} \hat{n}_{[d]2}^* V_{tb} V_{ts}^* \right) \sim |V_{tb}|^4 |V_{ts}|^2 \text{Im} (\delta_t^* - \delta_c^*) , \quad (115)$$

and we obtain, for the models near the bottom BGL one,

$$\frac{\text{BAU}_{\text{near b}}}{\text{BAU}_{\text{SM}}} = 10^{16} (t_\beta + t_\beta^{-1}) |V_{ts}| \text{Im} (\delta_t^* - \delta_c^*) , \quad (116)$$

that is, a potential enhancement of 10^{13} with respect to the SM.

Conclusions

We have analysed the question of FCNC in the framework of 2HDM. In this paper, it has been shown that BGL models can be readily generalized to gBGL models which keep the nice feature of renormalizability, but no longer belong to the Minimal Flavour Violation Framework. Contrary to BGL models, gBGL models contain four extra flavour parameters beyond the CKM matrix V , in spite of the drastic reduction in the number of parameters in gBGL, when compared to those present in the general 2HDM. It has been shown that in gBGL models FCNC are present at tree level, but in a controlled manner, rendering them plausible extensions of the SM. In fact they can have simultaneously FCNC in the up and in the down sector, a difference with BGL models that have FCNC in one of the sectors only.

The flavour structure of Yukawa couplings in gBGL is achieved, in a natural way, through the introduction of a \mathbb{Z}_2 symmetry, at the Lagrangian level. So gBGL models use the same symmetry as proposed by Glashow and Weinberg in NFC models, the only difference lies in the way the quark fields transform under \mathbb{Z}_2 . gBGL models contain BGL models as special cases, in the sense that in the parameter space of gBGL there are regions where one comes close to particular BGL models. Note that in the limit where a BGL is reached, the Lagrangian acquires a larger symmetry, namely a \mathbb{Z}_4 or a $U(1)$ symmetry, dictated by the corresponding neutrino type (either Majorana or Dirac) of the BGL model.

It is well known that the SM does not generate sufficient Baryon Asymmetry of the Universe (BAU) at the electroweak phase transition. One of the reasons for this is the fact that in the SM CP violation is too small. This stems from the fact that the WB invariant [41, 46] controlling CP violation in the SM is of order mass to the 12th. We have analysed in detail how lower order CP odd invariants appear in gBGL models. It turns out that they arise at a much lower mass order which leads to the possibility of generating a much higher BAU, compared to that in the SM.

Acknowledgments

This work is partially supported by Spanish MINECO under grant FPA2015-68318-R and by the Severo Ochoa Excellence Center Project SEV-2014-0398, by Generalitat Valenciana under grant GVPROMETEOII 2014-049 and by Fundação para a Ciência e a Tecnologia (FCT, Portugal) through the projects CERN/FIS-NUC/0010/2015 and CFTP-FCT Unit 777 (UID/FIS/00777/2013) which are partially funded through POCTI (FEDER), COMPETE, QREN and EU. MN acknowledges support from FCT through postdoctoral grant SFRH/BPD/112999/2015.

A Necessary and Sufficient Conditions for gBGL

We complete in this appendix the proof of the sufficient conditions in the following general result: *the WB invariant matrix conditions*

$$\begin{aligned}\Gamma_2^\dagger \Gamma_1 &= 0, & \Gamma_2^\dagger \Delta_1 &= 0, \\ \Delta_2^\dagger \Delta_1 &= 0, & \Delta_2^\dagger \Gamma_1 &= 0,\end{aligned}\tag{19}$$

are the necessary and sufficient conditions to define gBGL models or a type I 2HDM¹¹, provided there are no massless quarks.

It is always possible, in general, to write

$$\Gamma_i = W_{d_i} D_{d_i} U_{d_i}^\dagger, \quad \Delta_i = W_{u_i} D_{u_i} U_{u_i}^\dagger,\tag{117}$$

where W_{d_i} , W_{u_i} , U_{d_i} and U_{u_i} are unitary matrices, and D_{d_i} and D_{u_i} are diagonal ones. From $\Delta_2^\dagger \Delta_1 = 0$ it is straightforward that $[\Delta_1 \Delta_1^\dagger, \Delta_2 \Delta_2^\dagger] = 0$ and thus one can choose

$$W_{u_1} = W_{u_2} = W_u,\tag{118}$$

while from $\Gamma_2^\dagger \Gamma_1 = 0$ it follows that $[\Gamma_1 \Gamma_1^\dagger, \Gamma_2 \Gamma_2^\dagger] = 0$ and therefore we can also choose

$$W_{d_1} = W_{d_2} = W_d.\tag{119}$$

Now, in $\Delta_2^\dagger \Delta_1 = 0$, W_u simplifies away and we have

$$D_{u_2}^\dagger D_{u_1} = 0,\tag{120}$$

and similarly, for $\Gamma_2^\dagger \Gamma_1 = 0$,

$$D_{d_2}^\dagger D_{d_1} = 0.\tag{121}$$

If there are no massless quarks, there are two kinds of solutions for eq. (120),

(a) type I 2HDM

$$D_{u_1} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad D_{u_2} = \begin{pmatrix} u_1 & 0 & 0 \\ 0 & u_2 & 0 \\ 0 & 0 & u_3 \end{pmatrix},\tag{122}$$

with $u_i \neq 0$ in order to have massive up quarks. Notice that interchanging $D_{u_1} \leftrightarrow D_{u_2}$ will give rise to the same model, as explained later.

(b) gBGL

$$D_{u_1} = \begin{pmatrix} u_1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad D_{u_2} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & u_2 & 0 \\ 0 & 0 & u_3 \end{pmatrix},\tag{123}$$

with $u_i \neq 0$ again. As above, exchanging $D_{u_1} \leftrightarrow D_{u_2}$ does not introduce new models. We should also take into account the possibility that $u_1 \neq 0$ is in a different position in the diagonal of D_{u_1} while respecting $D_{u_2}^\dagger D_{u_1} = 0$, which is ensured with a corresponding permutation of the diagonal elements of D_{u_2} .

¹¹Since we are not specifying the leptonic sector, with type I we also refer to type X 2HDM.

Similarly to the up sector, we have different solutions of eq. (121), and we consider three possibilities.

(a) First,

$$D_{d_1} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad D_{d_2} = \begin{pmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{pmatrix}, \quad (124)$$

with $d_i \neq 0$. Notice that eq. (122) together with eq. (124) with interchanged $D_{d_1} \leftrightarrow D_{d_2}$ do not match in order to be a solution of eqs. (19).

(b) Second, D_{d_1} and D_{u_1} have equal rank, and we could consider in general permutations of the diagonal elements, for example

$$D_{d_1} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad D_{d_2} = \begin{pmatrix} d_1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & d_3 \end{pmatrix}. \quad (125)$$

Then, of course, the rank of D_{d_2} is equal to the rank of D_{u_2} .

(c) Third, D_{d_1} and D_{u_1} have different rank (and therefore D_{d_2} and D_{u_2} also have different rank), for example

$$D_{d_1} = \begin{pmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad D_{d_2} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & d_3 \end{pmatrix}. \quad (126)$$

We have to explore now which solutions to eqs. (19) arise from the available possibilities in eqs. (122)–(123) and eqs. (124)–(126).

- $D_{d_1} = 0$ if and only if $D_{u_1} = 0$, which corresponds to a type I or X 2HDM. The proof is simple: if $D_{d_1} = 0$, D_{d_2} has rank 3, and thus Γ_2 has rank 3. Since $\Gamma_2^\dagger \Delta_1 = 0$, all column vectors of Δ_1 are in the null-space of Γ_2^\dagger (they are all non-zero vectors transformed into the zero or null vector), but since $\text{rank}(\Gamma_2^\dagger) = 3$, according to the rank-nullity theorem, the null-space of Γ_2^\dagger has dimension $3 - 3 = 0$, and thus $\Delta_1 = 0$, that is $D_{u_1} = 0$. Eqs. (19) are then trivially verified. Of course, there is also the solution $D_{d_2} = D_{u_2} = 0$, which is completely equivalent with a trivial relabelling of the scalar doublets $\Phi_1 \leftrightarrow \Phi_2$.
- Next we show that concerning eq. (123) and eqs. (125)–(126), the ranks of the Yukawa matrices should match in the following manner: $\text{rank}(D_{d_1}) = \text{rank}(D_{u_1})$ and $\text{rank}(D_{d_2}) = \text{rank}(D_{u_2})$. Consider for definiteness D_{u_1} and D_{u_2} as in eq. (123). First, since $\Delta_2^\dagger \Gamma_1 = 0$ and $\Gamma_2^\dagger \Delta_1 = 0$, with $W = W_u^\dagger W_d$,

$$D_{u_2}^\dagger W D_{d_1} = 0, \quad D_{d_2}^\dagger W^\dagger D_{u_1} = 0. \quad (127)$$

$D_{u_2}^\dagger W$ has rank 2, and thus its null-space has dimension 1; according to the first equation above, if D_{d_1} had rank 2, then the null-space of $D_{u_2}^\dagger W$ would have at least dimension 2 in contradiction with the first statement: consequently, the

rank of D_{d_1} has to be 1. Then $\text{rank}(D_{d_1}) = \text{rank}(D_{u_1}) = 1$ and $\text{rank}(D_{d_2}) = \text{rank}(D_{u_2}) = 2$. That is, the matrices in eq. (125) match the ones in eq. (123) for solutions of eqs. (19) while eqs. (125) and (123) do not match. Finally, relabelling of scalar doublets $\Phi_1 \leftrightarrow \Phi_2$ gives equivalent solutions with $\text{rank}(D_{d_1}) = \text{rank}(D_{u_1}) = 2$ and $\text{rank}(D_{d_2}) = \text{rank}(D_{u_2}) = 1$.

- The following step is to show that the most general case can be taken to be

$$\begin{aligned} D_{u_1} &= \begin{pmatrix} u_1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & D_{u_2} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & u_2 & 0 \\ 0 & 0 & u_3 \end{pmatrix}, \\ D_{d_1} &= \begin{pmatrix} d_1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & D_{d_2} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{pmatrix}. \end{aligned} \quad (128)$$

It is trivial to check that with the insertion of

$$P_{12} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (129)$$

in $\Gamma_i = W_d P_{12} P_{12} D_{d_i} P_{12} P_{12} U_d^\dagger = W'_d (P_{12} D_{d_i} P_{12}) U_d'^\dagger$, one can redefine $W'_d = W_d P_{12}$ and $U'_d = U_d P_{12}$, while P_{12} in $P_{12} D_{d_i} P_{12}$ permutes the first and second elements in the diagonal. This would bring, for example, eq. (125) to the desired form, matching with eq. (123).

- Finally, substituting eq. (128) in $D_{u_1}^\dagger W D_{d_2} = 0$,

$$u_1^* \begin{pmatrix} 0 & d_2 W_{12} & d_3 W_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = 0, \quad (130)$$

we get $W_{12} = W_{13} = 0$, while from $D_{u_2}^\dagger W D_{d_1} = 0$,

$$d_1 \begin{pmatrix} 0 & 0 & 0 \\ u_2^* W_{21} & 0 & 0 \\ u_3^* W_{31} & 0 & 0 \end{pmatrix} = 0, \quad (131)$$

we have $W_{21} = W_{31} = 0$. Then W has the block structure

$$W = W_u^\dagger W_d = e^{i\omega} \begin{pmatrix} 1 & 0 & 0 \\ 0 & W_{22} & W_{23} \\ 0 & W_{32} & W_{33} \end{pmatrix}. \quad (132)$$

One can now complete the proof; going back to eq. (117):

$$\begin{aligned} \Delta_1 &= W_u \begin{pmatrix} u_1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} U_{u_1}^\dagger = W_u \begin{pmatrix} \times & \times & \times \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\ \Delta_2 &= W_u \begin{pmatrix} 0 & 0 & 0 \\ 0 & u_2 & 0 \\ 0 & 0 & u_3 \end{pmatrix} U_{u_2}^\dagger = W_u \begin{pmatrix} 0 & 0 & 0 \\ \times & \times & \times \\ \times & \times & \times \end{pmatrix}, \end{aligned} \quad (133)$$

$$\begin{aligned}
\Gamma_1 &= W_d \begin{pmatrix} d_1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} U_{d_1}^\dagger = W_u W \begin{pmatrix} \times & \times & \times \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = W_u \begin{pmatrix} \times & \times & \times \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\
\Gamma_2 &= W_d \begin{pmatrix} 0 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{pmatrix} U_{d_2}^\dagger = W_u W \begin{pmatrix} 0 & 0 & 0 \\ \times & \times & \times \\ \times & \times & \times \end{pmatrix} = W_u \begin{pmatrix} 0 & 0 & 0 \\ \times & \times & \times \\ \times & \times & \times \end{pmatrix}. \quad (134)
\end{aligned}$$

With a WB transformation given by $Q_L \mapsto W_u Q_L$ we arrive to the equivalent gBGL structures in eqs. (14).

B Electric Dipole Moments

Attending to the explicit form of the N_d and N_u matrices in section 4 and the discussion in section 8, it is clear that gBGL models include new sources of CP violation. Although the dedicated analysis of section 6 addresses the controlled nature of the FCNC, one might still be concerned with the possibility that too large contributions to EDMs are present, induced in particular by the couplings to the charged scalar H^+ or by the (flavour) diagonal couplings to the neutral scalars, present in eqs. (72).

For the latter, let us consider for the moment a generic generation j of the up sector (for the down sector the reasoning is identical); the flavour diagonal couplings in eqs. (72) are controlled by $[M_u]_{jj}$ and $[N_u]_{jj}\gamma_R + [N_u^\dagger]_{jj}\gamma_L$ for the neutral scalars H^0 and R^0 , and by $i[N_u]_{jj}\gamma_R - i[N_u^\dagger]_{jj}\gamma_L$ for the pseudoscalar A . Following eq. (57),

$$[N_u]_{jj} = t_\beta m_{u_j} - (t_\beta + t_\beta^{-1}) \hat{n}_{[u]j} \hat{n}_{[u]j}^* m_{u_j} = (t_\beta - (t_\beta + t_\beta^{-1}) |\hat{n}_{[u]j}|^2) m_{u_j} = [N_u^\dagger]_{jj}, \quad (135)$$

and hence, for the coupling to R^0 ,

$$[N_u]_{jj}\gamma_R + [N_u^\dagger]_{jj}\gamma_L = (t_\beta - (t_\beta + t_\beta^{-1}) |\hat{n}_{[u]j}|^2) m_{u_j}, \quad (136)$$

while for the coupling to A ,

$$i[N_u]_{jj}\gamma_R - i[N_u^\dagger]_{jj}\gamma_L = i(t_\beta - (t_\beta + t_\beta^{-1}) |\hat{n}_{[u]j}|^2) m_{u_j} \gamma_5. \quad (137)$$

In general we have

$$\text{Im}([N_u]_{jj}) = \text{Im}([N_d]_{jj}) = 0 \quad (138)$$

and

$$\text{Im}(\text{Tr}\{N_u M_u^\dagger\}) = \text{Im}(\text{Tr}\{N_d M_d^\dagger\}) = 0. \quad (139)$$

With such couplings, the flavour diagonal quark interactions with the different neutral scalars are not CP violating [47, 48], and no contribution to the EDMs arises. This applies to one loop contributions, to two loop Barr-Zee contributions [49], to two loop contributions to the three gluon Weinberg operator [50] and to effective four fermion operators [51]. Notice that, although H^0 and R^0 are not the physical scalars, this conclusion remains unchanged when they are rotated into the mass eigenstates H and h in eq. (71).

For the charged scalar H^+ , one loop diagrams and contributions to the three gluon operator are not CP violating. Furthermore, the charged Higgs Barr-Zee contributions which are expected to be dominant [52] (see also [53]), are also CP conserving because their proportionality to $\text{Im}([N_u]_{tt}[N_q]_{jj}) = 0$.

References

- [1] **ATLAS** Collaboration, G. Aad *et al.*, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett.* **B716** (2012) 1–29, [1207.7214].
- [2] **CMS** Collaboration, S. Chatrchyan *et al.*, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett.* **B716** (2012) 30–61, [1207.7235].
- [3] T. Lee, *A Theory of Spontaneous T Violation*, *Phys.Rev.* **D8** (1973) 1226–1239.
- [4] G. Branco, P. Ferreira, L. Lavoura, M. Rebelo, M. Sher, *et al.*, *Theory and phenomenology of two-Higgs-doublet models*, *Phys.Rept.* **516** (2012) 1–102, [1106.0034].
- [5] I. P. Ivanov, *Building and testing models with extended Higgs sectors*, 1702.03776.
- [6] S. L. Glashow and S. Weinberg, *Natural Conservation Laws for Neutral Currents*, *Phys.Rev.* **D15** (1977) 1958.
- [7] E. Paschos, *Diagonal Neutral Currents*, *Phys.Rev.* **D15** (1977) 1966.
- [8] A. Pich and P. Tuzon, *Yukawa Alignment in the Two-Higgs-Doublet Model*, *Phys.Rev.* **D80** (2009) 091702, [0908.1554].
- [9] F. Mahmoudi and O. Stal, *Flavor constraints on the two-Higgs-doublet model with general Yukawa couplings*, *Phys. Rev.* **D81** (2010) 035016, [0907.1791].
- [10] A. Crivellin, A. Kokulu, and C. Greub, *Flavor-phenomenology of two-Higgs-doublet models with generic Yukawa structure*, *Phys.Rev.* **D87** (2013), no. 9 094031, [1303.5877].
- [11] R. Gaitán, R. Martinez, and J. H. M. de Oca, *Rare top decay $t \rightarrow c\gamma$ with flavor changing neutral scalar interactions in two Higgs doublet model*, *Phys. Rev.* **D94** (2016), no. 9 094038, [1503.04391].
- [12] B. Altunkaynak, W.-S. Hou, C. Kao, M. Kohda, and B. McCoy, *Flavor Changing Heavy Higgs Interactions at the LHC*, *Phys. Lett.* **B751** (2015) 135–142, [1506.00651].
- [13] A. Arhrib, R. Benbrik, C.-H. Chen, M. Gomez-Bock, and S. Semmlali, *Two-Higgs-doublet type-II and -III models and $t \rightarrow ch$ at the LHC*, *Eur. Phys. J.* **C76** (2016), no. 6 328, [1508.06490].
- [14] C. S. Kim, Y. W. Yoon, and X.-B. Yuan, *Exploring top quark FCNC within 2HDM type III in association with flavor physics*, *JHEP* **12** (2015) 038, [1509.00491].

- [15] T. Enomoto and R. Watanabe, *Flavor constraints on the Two Higgs Doublet Models of Z_2 symmetric and aligned types*, *JHEP* **05** (2016) 002, [1511.05066].
- [16] R. Benbrik, C.-H. Chen, and T. Nomura, *$h, Z \rightarrow \ell_i \bar{\ell}_j$, Δa_μ , $\tau \rightarrow (3\mu, \mu\gamma)$ in generic two-Higgs-doublet models*, *Phys. Rev.* **D93** (2016), no. 9 095004, [1511.08544].
- [17] G. Branco, W. Grimus, and L. Lavoura, *Relating the scalar flavor changing neutral couplings to the CKM matrix*, *Phys.Lett.* **B380** (1996) 119–126, [hep-ph/9601383].
- [18] A. Buras, P. Gambino, M. Gorbahn, S. Jager, and L. Silvestrini, *Universal unitarity triangle and physics beyond the standard model*, *Phys.Lett.* **B500** (2001) 161–167, [hep-ph/0007085].
- [19] G. D’Ambrosio, G. Giudice, G. Isidori, and A. Strumia, *Minimal flavor violation: An Effective field theory approach*, *Nucl.Phys.* **B645** (2002) 155–187, [hep-ph/0207036].
- [20] V. Cirigliano, B. Grinstein, G. Isidori, and M. B. Wise, *Minimal flavor violation in the lepton sector*, *Nucl.Phys.* **B728** (2005) 121–134, [hep-ph/0507001].
- [21] F. Botella, G. Branco, and M. Rebelo, *Minimal Flavour Violation and Multi-Higgs Models*, *Phys.Lett.* **B687** (2010) 194–200, [0911.1753].
- [22] L. Lavoura, *Models of CP violation exclusively via neutral scalar exchange*, *Int.J.Mod.Phys.* **A9** (1994) 1873–1888.
- [23] F. Botella, G. Branco, M. Nebot, and M. Rebelo, *Two-Higgs Leptonic Minimal Flavour Violation*, *JHEP* **1110** (2011) 037, [1102.0520].
- [24] F. Botella, G. Branco, A. Carmona, M. Nebot, L. Pedro, and M. Rebelo, *Physical Constraints on a Class of Two-Higgs Doublet Models with FCNC at tree level*, *JHEP* **1407** (2014) 078, [1401.6147].
- [25] G. Bhattacharyya, D. Das, and A. Kundu, *Feasibility of light scalars in a class of two-Higgs-doublet models and their decay signatures*, *Phys.Rev.* **D89** (2014) 095029, [1402.0364].
- [26] F. J. Botella, G. C. Branco, M. Nebot, and M. N. Rebelo, *Flavour Changing Higgs Couplings in a Class of Two Higgs Doublet Models*, *Eur. Phys. J.* **C76** (2016), no. 3 161, [1508.05101].
- [27] H. Georgi and D. V. Nanopoulos, *Suppression of Flavor Changing Effects From Neutral Spinless Meson Exchange in Gauge Theories*, *Phys. Lett.* **B82** (1979) 95.
- [28] J. F. Donoghue and L. F. Li, *Properties of Charged Higgs Bosons*, *Phys. Rev.* **D19** (1979) 945.

- [29] F. J. Botella and J. P. Silva, *Jarlskog - like invariants for theories with scalars and fermions*, *Phys. Rev.* **D51** (1995) 3870–3875, [[hep-ph/9411288](#)].
- [30] P. M. Ferreira and J. P. Silva, *Abelian symmetries in the two-Higgs-doublet model with fermions*, *Phys. Rev.* **D83** (2011) 065026, [[1012.2874](#)].
- [31] G. C. Branco and M. N. Rebelo, *The Higgs Mass in a Model With Two Scalar Doublets and Spontaneous CP Violation*, *Phys. Lett.* **B160** (1985) 117–120.
- [32] A. G. Akeroyd, A. Arhrib, and E.-M. Naimi, *Note on tree level unitarity in the general two Higgs doublet model*, *Phys. Lett.* **B490** (2000) 119–124, [[hep-ph/0006035](#)].
- [33] I. F. Ginzburg and I. P. Ivanov, *Tree-level unitarity constraints in the most general 2HDM*, *Phys. Rev.* **D72** (2005) 115010, [[hep-ph/0508020](#)].
- [34] B. Grinstein, C. W. Murphy, and P. Uttayarat, *One-loop corrections to the perturbative unitarity bounds in the CP-conserving two-Higgs doublet model with a softly broken \mathbb{Z}_2 symmetry*, *JHEP* **06** (2016) 070, [[1512.04567](#)].
- [35] V. Cacchio, D. Chowdhury, O. Eberhardt, and C. W. Murphy, *Next-to-leading order unitarity fits in Two-Higgs-Doublet models with soft \mathbb{Z}_2 breaking*, *JHEP* **11** (2016) 026, [[1609.01290](#)].
- [36] G. Blankenburg, J. Ellis, and G. Isidori, *Flavour-Changing Decays of a 125 GeV Higgs-like Particle*, *Phys. Lett.* **B712** (2012) 386–390, [[1202.5704](#)].
- [37] **ATLAS** Collaboration, G. Aad *et al.*, *Search for top quark decays $t \rightarrow qH$ with $H \rightarrow \gamma\gamma$ using the ATLAS detector*, *JHEP* **06** (2014) 008, [[1403.6293](#)].
- [38] **CMS** Collaboration, V. Khachatryan *et al.*, *Searches for heavy Higgs bosons in two-Higgs-doublet models and for $t \rightarrow ch$ decay using multilepton and diphoton final states in pp collisions at 8 TeV*, *Phys. Rev.* **D90** (2014) 112013, [[1410.2751](#)].
- [39] **CMS** Collaboration, V. Khachatryan *et al.*, *Search for top quark decays via Higgs-boson-mediated flavor-changing neutral currents in pp collisions at $\sqrt{s} = 8$ TeV*, [1610.04857](#).
- [40] F. Botella, G. Branco, and M. Rebelo, *Invariants and Flavour in the General Two-Higgs Doublet Model*, *Phys. Lett.* **B722** (2013) 76–82, [[1210.8163](#)].
- [41] J. Bernabeu, G. C. Branco, and M. Gronau, *CP restrictions on quark mass matrices*, *Phys. Lett.* **169B** (1986) 243–247.
- [42] S. Davidson and H. E. Haber, *Basis-independent methods for the two-Higgs-doublet model*, *Phys. Rev.* **D72** (2005) 035004, [[hep-ph/0504050](#)]. [[Erratum: Phys. Rev.D72,099902\(2005\)](#)].

- [43] L. Wolfenstein, *Parametrization of the Kobayashi-Maskawa Matrix*, *Phys.Rev.Lett.* **51** (1983) 1945.
- [44] M. Trodden, *Electroweak baryogenesis*, *Rev. Mod. Phys.* **71** (1999) 1463–1500, [[hep-ph/9803479](#)].
- [45] D. E. Morrissey and M. J. Ramsey-Musolf, *Electroweak baryogenesis*, *New J. Phys.* **14** (2012) 125003, [[1206.2942](#)].
- [46] C. Jarlskog, *Commutator of the Quark Mass Matrices in the Standard Electroweak Model and a Measure of Maximal CP Violation*, *Phys. Rev. Lett.* **55** (1985) 1039.
- [47] G. C. Branco, L. Lavoura, and J. P. Silva, *CP Violation, International Series of Monographs on Physics*, *Oxford University Press*.
- [48] M. Nebot and J. P. Silva, *Self-cancellation of a scalar in neutral meson mixing and implications for the LHC*, *Phys. Rev.* **D92** (2015), no. 8 085010, [[1507.07941](#)].
- [49] S. M. Barr and A. Zee, *Electric Dipole Moment of the Electron and of the Neutron*, *Phys. Rev. Lett.* **65** (1990) 21–24. [Erratum: *Phys. Rev. Lett.* 65,2920(1990)].
- [50] S. Weinberg, *Larger Higgs Exchange Terms in the Neutron Electric Dipole Moment*, *Phys. Rev. Lett.* **63** (1989) 2333.
- [51] M. Jung and A. Pich, *Electric Dipole Moments in Two-Higgs-Doublet Models*, *JHEP* **04** (2014) 076, [[1308.6283](#)].
- [52] D. Bowser-Chao, D. Chang, and W.-Y. Keung, *Electron electric dipole moment from CP violation in the charged Higgs sector*, *Phys. Rev. Lett.* **79** (1997) 1988–1991, [[hep-ph/9703435](#)].
- [53] V. Ilisie, *New Barr-Zee contributions to $(\mathbf{g} - \mathbf{2})_\mu$ in two-Higgs-doublet models*, *JHEP* **04** (2015) 077, [[1502.04199](#)].