

Riemannian stochastic quasi-Newton algorithm with variance reduction and its convergence analysis

HiroYuki Kasai*

HiroYuki Sato†

Bamdev Mishra‡

February 7, 2022

Abstract

Stochastic variance reduction algorithms have recently become popular for minimizing the average of a large, but finite number of loss functions. The present paper proposes a Riemannian stochastic quasi-Newton algorithm with variance reduction (R-SQN-VR). The key challenges of averaging, adding, and subtracting multiple gradients are addressed with notions of retraction and vector transport. We present convergence analyses of R-SQN-VR on both non-convex and retraction-convex functions under retraction and vector transport operators. The proposed algorithm is evaluated on the Karcher mean computation on the symmetric positive-definite manifold and the low-rank matrix completion on the Grassmann manifold. In all cases, the proposed algorithm outperforms the state-of-the-art Riemannian batch and stochastic gradient algorithms.

1 Introduction

Let $f : \mathcal{M} \rightarrow \mathbb{R}$ be a smooth real-valued function on a Riemannian manifold \mathcal{M} [1]. The problem under consideration in the present paper is the minimization of the *expected risk* of f for a given model variable $w \in \mathcal{M}$ taken with respect to the distribution of z , i.e., $\min_{w \in \mathcal{M}} f(w)$, where $f(w) = \mathbb{E}_z[f(w; z)] = \int f(w; z) dP(z)$ and z is a random seed representing a single sample or set of samples. When given a set of realizations $\{z_{[n]}\}_{n=1}^N$ of z , we define the loss incurred by the parameter vector w with respect to the n -th sample as $f_n(w) := f(w; z_{[n]})$, and then the *empirical risk* is defined as the average of the sample losses:

$$\min_{w \in \mathcal{M}} \left\{ f(w) := \frac{1}{N} \sum_{n=1}^N f_n(w) \right\}, \quad (1)$$

where N is the total number of the elements. This problem has many applications that include, to name a few, principal component analysis (PCA) and the subspace tracking problem [2] on the Grassmann manifold. The low-rank matrix/tensor completion problem is a promising example of the manifold of fixed-rank matrices/tensors [3, 4]. The linear regression problem is also defined on the manifold of the fixed-rank matrices [5].

*Graduate School of Informatics and Engineering, The University of Electro-Communications, Tokyo, Japan (kasai@is.uec.ac.jp).

†Department of Information and Computer Technology, Tokyo University of Science, Tokyo, Japan (hsato@rs.tus.ac.jp).

‡Core Machine Learning Team, Amazon.com, Bangalore, India (bamdevm@amazon.com.)

Riemannian gradient descent requires the *Riemannian full gradient* estimation, i.e., $\text{grad}f(w) = \sum_{n=1}^N \text{grad}f_n(w)$, for every iteration, where $\text{grad}f_n(w)$ is the *Riemannian stochastic gradient* of $f_n(w)$ on the Riemannian manifold \mathcal{M} for n -th sample. This estimation is computationally heavy when N is extremely large. A popular alternative is *Riemannian stochastic gradient descent* (R-SGD) that extends *stochastic gradient descent* (SGD) in the Euclidean space [6]. Because this uses only one $\text{grad}f_n(w)$, the complexity per iteration is independent of N . However, similarly to SGD [7], R-SGD suffers from a slow convergence due to a *decaying step-size* sequence. *Variance reduction* (VR) methods have been proposed recently to accelerate the convergence of SGD in the Euclidean space [8–12]. One distinguished feature is to calculate a full gradient estimation periodically, and to re-use it to reduce the variance of noisy stochastic gradient. However, because all previously described algorithms are *first-order* algorithms, their convergence speed can be slow because of their poor curvature approximations in *ill-conditioned* problems as seen in Section 4. One promising approach is *second-order* algorithms such as stochastic *quasi-Newton* (QN) methods using Hessian evaluations [13–16]. They achieve faster convergence by exploiting curvature information of the objective function f . Furthermore, addressing these two acceleration techniques, [17] and [18] propose a hybrid algorithm of the stochastic QN method accompanied with the VR method.

Examining the Riemannian manifolds again, many challenges on the QN method have been addressed in deterministic settings [19–21]. The VR method in the Euclidean space has also been extended to Riemannian manifolds, so-called R-SVRG [22, 23]. Nevertheless, the second-order stochastic algorithm with the VR method has not been explored thoroughly for the problem (1). To this end, we propose a Riemannian stochastic QN method based on L-BFGS and the VR method.

Our contributions are four-fold; (i) we propose a novel (and to the best of our knowledge, the first) Riemannian limited-memory QN algorithm with a VR method. (ii) Our convergence analysis deals with both *non-convex* and (*strongly*) *retraction-convex* functions. In this paper, f is said to be strongly retraction-convex when f is (strongly) convex along a curve on \mathcal{M} defined by a retraction R (Assumption 3) while the other functions are called as non-convex functions. (iv) The proposed algorithm and its analyses are considered under computationally efficient *retraction* and *vector transport* operations instead of the more restrictive *exponential mapping* and *parallel translation* operations. This is more challenging than R-SVRG [23], but gives us a big advantage other than computational efficiency, i.e., wider kinds of applicable manifolds. For example, while [23] cannot be applied to the Stiefel and fixed-rank manifolds because these manifolds do not have closed form expressions for parallel translation, our analyses and algorithm can be directly applied to them.

The specific features of the algorithms are two-fold; (i) we update the curvature pair of the QN method every outer loop by exploiting full gradient estimations in the VR method, and thereby capture more precise and stabler curvature information. This avoids additional sweeping of samples required in the Euclidean stochastic QN [16], additional gradient estimations required in the Euclidean online BFGS (oBFGS) [13, 14, 24], or additional sub-sampling of Hessian [16, 17]. (ii) Compared with a simple Riemannian extension of the QN method, a noteworthy advantage of its combination with the VR method is that, as revealed below, frequent transportations of curvature information between different tangent spaces, which are inextricable in such a simple Riemannian extension, can be drastically reduced. This is a special benefit of the Riemannian hybrid algorithm, which does not exist in the Euclidean case [17, 18]. More specifically, the calculations of curvature information and the *second-order modified Riemannian stochastic gradient* are performed uniformly on the tangent space of the

outer loop.

The paper is organized as follows. Section 2 presents details of our proposed R-SQN-VR. Section 3 presents the convergence analyses. In Section 4, numerical comparisons with R-SGD and R-SVRG on two problems are provided with results suggesting the superior performances of R-SQN-VR. The proposed R-SQN-VR is implemented in the Matlab toolbox Manopt [25]. The concrete proofs of theorems and additional experiments are provided as supplementary material.

2 Riemannian stochastic quasi-Newton algorithm with variance reduction (R-SQN-VR)

We assume that the manifold \mathcal{M} is endowed with a Riemannian metric structure, i.e., a smooth inner product $\langle \cdot, \cdot \rangle_w$ of tangent vectors is associated with the tangent space $T_w\mathcal{M}$ for all $w \in \mathcal{M}$ [1]. The *norm* $\|\cdot\|_w$ of a tangent vector is the norm associated with the Riemannian metric. The metric structure allows a systematic framework for optimization over manifolds. Conceptually, the constrained optimization problem (1) is translated into an *unconstrained* problem over \mathcal{M} .

2.1 R-SGD and R-SVRG

R-SGD: Given a starting point $w_0 \in \mathcal{M}$, R-SGD produces a sequence $\{w_t\}$ in \mathcal{M} that converges to a first-order critical point of (1). Specifically, it updates w as

$$w_{t+1} = R_{w_t}(-\alpha_t \text{grad} f_n(w_t, z_t)),$$

where α_t is the step-size, and where $\text{grad} f_n(w_t, z_t)$ is a Riemannian stochastic gradient, which is a tangent vector at $w_t \in \mathcal{M}$. $\text{grad} f_n(w_t, z_t)$ represents an unbiased estimator of the Riemannian full gradient $\text{grad} f(w_t)$, and the expectation of $\text{grad} f_n(w_t, z_t)$ over the choices of z_t is $\text{grad} f(w_t)$, i.e., $\mathbb{E}_{z_t}[\text{grad} f_n(w_t, z_t)] = \text{grad} f(w_t)$. The update moves from w_t in the direction $-\text{grad} f_n(w_t, z_t)$ with a step-size α_t while remaining on \mathcal{M} . This mapping, denoted as $R_w : T_w\mathcal{M} \rightarrow \mathcal{M} : \zeta_w \mapsto R_w(\zeta_w)$, is called *retraction* at w , which maps the tangent bundle $T_w\mathcal{M}$ onto \mathcal{M} with a local rigidity condition that preserves gradients at w . *Exponential mapping* Exp is an instance of the retraction.

R-SVRG: R-SVRG has double loops where a k -th outer loop, called *epoch*, has m_k inner iterations. R-SVRG keeps $\tilde{w}^k \in \mathcal{M}$ after m_{k-1} inner iterations of $(k-1)$ -th epoch, and computes the full Riemannian gradient $\text{grad} f(\tilde{w}^k)$ only for this stored \tilde{w}^k . It also computes the Riemannian stochastic gradient $\text{grad} f_{i_t^k}(\tilde{w}^k)$ for i_t^k -th sample. Then, picking i_t^k -th sample for each t -th inner iteration of k -th epoch at w_t^k , we calculate ξ_t^k , i.e., by modifying $\text{grad} f_{i_t^k}(w_t^k)$ using both $\text{grad} f(\tilde{w}^k)$ and $\text{grad} f_{i_t^k}(\tilde{w}^k)$. Because they belong to different tangent spaces, a simple addition of them is not well-defined because Riemannian manifolds are not vector spaces. Therefore, after $\text{grad} f_{i_t^k}(\tilde{w}^k)$ and $\text{grad} f(\tilde{w}^k)$ are transported to $T_{w_t^k}\mathcal{M}$ by $\mathcal{T}_{\tilde{\eta}_t^k}$, ξ_t^k is set as

$$\xi_t^k = \text{grad} f_{i_t^k}(w_t^k) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f_{i_t^k}(\tilde{w}^k) - \text{grad} f(\tilde{w}^k)),$$

where \mathcal{T} represents *vector transport* from \tilde{w}^k to w_t^k , and $\tilde{\eta}_t^k \in T_{\tilde{w}^k}\mathcal{M}$ satisfies $R_{\tilde{w}^k}(\tilde{\eta}_t^k) = w_t^k$. The vector transport $\mathcal{T} : T\mathcal{M} \oplus T\mathcal{M} \rightarrow T\mathcal{M}, (\eta_w, \xi_w) \mapsto \mathcal{T}_{\eta_w}\xi_w$ is associated with retraction

Algorithm 1 Riemannian stochastic quasi-Newton with variance reduction (R-SQN-VR).

Require: Update frequency m_k , step-size $\alpha_t^k > 0$, memory size L , number of epochs K , and cautious update threshold ϵ .

- 1: Initialize \tilde{w}^0 , and calculate the Riemannian full gradient $\text{grad}f(\tilde{w}^0)$.
 - 2: **for** $k = 0, \dots, K - 1$ **do**
 - 3: Store $w_0^k = \tilde{w}^k$.
 - 4: **for** $t = 0, 1, 2, \dots, m_k - 1$ **do**
 - 5: Choose $i_t^k \in \{1, \dots, N\}$ uniformly at random.
 - 6: Calculate the tangent vector $\tilde{\eta}_t^k$ from \tilde{w}^k to w_t^k by $\tilde{\eta}_t^k = R_{\tilde{w}^k}^{-1}(w_t^k)$.
 - 7: **if** $k > 1$ **then**
 - 8: Transport the stochastic gradient $\text{grad}f_{i_t^k}(w_t^k)$ to $T_{\tilde{w}^k}\mathcal{M}$ by $(\mathcal{T}_{\tilde{\eta}_t^k})^{-1}\text{grad}f_{i_t^k}(w_t^k)$.
 - 9: Calculate $\tilde{\xi}_t^k$ as $\tilde{\xi}_t^k = (\mathcal{T}_{\tilde{\eta}_t^k})^{-1}\text{grad}f_{i_t^k}(w_t^k) - (\text{grad}f_{i_t^k}(\tilde{w}^k) - \text{grad}f(\tilde{w}^k))$.
 - 10: Calculate $\tilde{\mathcal{H}}_t^k \tilde{\xi}_t^k$, transport $\tilde{\mathcal{H}}_t^k \tilde{\xi}_t^k$ back to $T_{w_t^k}\mathcal{M}$ by $\mathcal{T}_{\tilde{\eta}_t^k} \tilde{\mathcal{H}}_t^k \tilde{\xi}_t^k$, and obtain $\mathcal{H}_t^k \xi_t^k$.
 - 11: Update w_{t+1}^k from w_t^k as $w_{t+1}^k = R_{w_t^k}(-\alpha_t^k \mathcal{H}_t^k \xi_t^k)$.
 - 12: **else**
 - 13: Calculate ξ_t^k as $\xi_t^k = \text{grad}f_{i_t^k}(w_t^k) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad}f_{i_t^k}(\tilde{w}^k) - \text{grad}f(\tilde{w}^k))$.
 - 14: Update w_{t+1}^k from w_t^k as $w_{t+1}^k = R_{w_t^k}(-\alpha_t^k \xi_t^k)$.
 - 15: **end if**
 - 16: **end for**
 - 17: Option I: $\tilde{w}^{k+1} = g_{m_k}(w_1^k, \dots, w_{m_k}^k)$ (or $\tilde{w}^{k+1} = w_t^k$ for randomly chosen $t \in \{1, \dots, m_k\}$).
 - 18: Option II: $\tilde{w}^{k+1} = w_{m_k}^k$.
 - 19: Calculate the Riemannian full gradient $\text{grad}f(\tilde{w}^{k+1})$.
 - 20: Calculate the tangent vector η_k from \tilde{w}^k to \tilde{w}^{k+1} by $\eta_k = R_{\tilde{w}^k}^{-1}(\tilde{w}^{k+1})$.
 - 21: Compute $s_k^{k+1} = \mathcal{T}_{\eta_k} \eta_k$, and $y_k^{k+1} = \kappa_k^{-1} \text{grad}f(\tilde{w}^{k+1}) - \mathcal{T}_{\eta_k} \text{grad}f(\tilde{w}^k)$ where $\kappa_k = \|\eta_k\|_{\tilde{w}^k} / \|\mathcal{T}_{R_{\eta_k}} \eta_k\|_{\tilde{w}^k}$.
 - 22: **if** $\langle y_k^{k+1}, s_k^{k+1} \rangle_{\tilde{w}^{k+1}} \geq \epsilon \|s_k^{k+1}\|_{\tilde{w}^{k+1}}^2$ **then**
 - 23: Discard pair (s_{k-L}^k, y_{k-L}^k) when $k > L$, and store pair (s_k^{k+1}, y_k^{k+1}) .
 - 24: **end if**
 - 25: Transport $\{(s_j^k, y_j^k)\}_{j=k-\tau+1}^{k-1} \in T_{\tilde{w}^k}\mathcal{M}$ to $\{(s_j^{k+1}, y_j^{k+1})\}_{j=k-\tau+1}^{k-1} \in T_{\tilde{w}^{k+1}}\mathcal{M}$ by \mathcal{T}_{η_k} .
 - 26: **end for**
 - 27: Option III: output $w_{\text{sol}} = \tilde{w}^K$
 - 28: Option IV: output $w_{\text{sol}} = w_t^k$ for randomly chosen $t \in \{1, \dots, m_k\}$ and $k \in \{1, \dots, K\}$.
-

R and all $\xi_w, \zeta_w \in \mathcal{T}_w \mathcal{M}$. It holds that (i) $\mathcal{T}_{\eta_w} \xi_w \in \mathcal{T}_{R(\eta_w)} \mathcal{M}$, (ii) $\mathcal{T}_{0_w} \xi_w = \xi_w$, and (iii) \mathcal{T}_{η_w} is a linear map. *Parallel translation* P is an instance of the vector transport. Consequently, the final update is defined as $w_{t+1}^k = R_{w_t^k}(-\alpha_t^k \xi_t^k)$.

2.2 Proposed R-SQN-VR

We propose a Riemannian stochastic QN method accompanied with a VR method (R-SQN-VR). A straightforward extension is to update the modified stochastic gradient ξ_t^k by premultiplying a linear *inverse Hessian approximation operator* \mathcal{H}_t^k at w_t^k as

$$w_{t+1}^k = R_{w_t^k}(-\alpha_t^k \mathcal{H}_t^k \xi_t^k),$$

where $\mathcal{H}_t^k := \mathcal{T}_{\tilde{\eta}_t^k} \circ \tilde{\mathcal{H}}^k \circ (\mathcal{T}_{\tilde{\eta}_t^k})^{-1}$ by denoting the inverse Hessian approximation at \tilde{w}^k simply as $\tilde{\mathcal{H}}^k$. Here, \mathcal{T} is an isometric vector transport explained in Section 3. \mathcal{H}_t^k should be positive definite, i.e., $\mathcal{H}_t^k \succ 0$ and is close to the Hessian of f , i.e., $\text{Hess}f(w_t^k)$. It is noteworthy that $\tilde{\mathcal{H}}^k$ is calculated only every outer epoch, and remains to be used for \mathcal{H}_t^k throughout the corresponding k -th epoch.

Curvature pair (s_k^{k+1}, y_k^{k+1}) : This paper particularly addresses the operator $\tilde{\mathcal{H}}^k$ used in L-BFGS intended for a large-scale data. Thus, let s_k^{k+1} and y_k^{k+1} be the variable variation and the gradient variation at $T_{\tilde{w}^{k+1}} \mathcal{M}$, respectively, where the superscript expresses explicitly that they belong to $T_{\tilde{w}^{k+1}} \mathcal{M}$. It should be noted that the curvature pair (s_k^{k+1}, y_k^{k+1}) is calculated at the new $T_{\tilde{w}^{k+1}} \mathcal{M}$ just after k -th epoch finished. Furthermore, after the epoch index k is incremented, the curvature pair must be used only at $T_{\tilde{w}^k} \mathcal{M}$ because the calculation of $\tilde{\mathcal{H}}^k$ is performed only at $T_{\tilde{w}^k} \mathcal{M}$.

The variable variation s_k^{k+1} is calculated from the difference between \tilde{w}^{k+1} and \tilde{w}^k . This is represented by the tangent vector η_k from \tilde{w}^k to \tilde{w}^{k+1} , which is calculated using the inverse of the retraction $R_{\tilde{w}^k}^{-1}(\tilde{w}^{k+1})$. Since η_k belongs to the $T_{\tilde{w}^k} \mathcal{M}$, transporting this onto $T_{\tilde{w}^{k+1}} \mathcal{M}$ yields

$$s_k^{k+1} = \mathcal{T}_{\eta_k} \eta_k (= \mathcal{T}_{\eta_k} R_{\tilde{w}^k}^{-1}(\tilde{w}^{k+1})). \quad (2)$$

The gradient variation y_k^{k+1} is calculated from the difference between the new full gradient $\text{grad}f(\tilde{w}^{k+1}) \in T_{\tilde{w}^{k+1}} \mathcal{M}$ and the previous, but transported $\mathcal{T}_{\eta_k} \text{grad}f(\tilde{w}^k) \in T_{\tilde{w}^k} \mathcal{M}$ [21] as

$$y_k^{k+1} = \kappa_k^{-1} \text{grad}f(\tilde{w}^{k+1}) - \mathcal{T}_{\eta_k} \text{grad}f(\tilde{w}^k), \quad (3)$$

where $\kappa_k > 0$ is explained in Section 3.

Inverse Hessian approximation operator $\tilde{\mathcal{H}}^k$: $\tilde{\mathcal{H}}^k$ is calculated using the past curvature pairs. More specifically, $\tilde{\mathcal{H}}^k$ is updated as $\tilde{\mathcal{H}}^{k+1} = (\check{\mathcal{V}}^k)^{\flat} \check{\mathcal{H}}_k \check{\mathcal{V}}^k + \rho_k s_k s_k^{\flat}$, where $\check{\mathcal{H}}_k = \mathcal{T}_{\eta_k} \circ \tilde{\mathcal{H}}^k \circ \mathcal{T}_{\eta_k}^{-1}$, $\rho_k = 1/\langle y_k, s_k \rangle$, $\check{\mathcal{V}}^k = \text{id} - \rho_k y_k s_k^{\flat}$ with identity mapping id [21]. Therein, a^{\flat} denotes the flat of $a \in T_w \mathcal{M}$, i.e., $a^{\flat} : T_w \mathcal{M} \rightarrow \mathbb{R} : v \rightarrow \langle a, v \rangle_w$. Thus, $\tilde{\mathcal{H}}^k$ depends on $\tilde{\mathcal{H}}^{k-1}$ and (s_{k-1}, y_{k-1}) , and similarly $\tilde{\mathcal{H}}^{k-1}$ depends on $\tilde{\mathcal{H}}^{k-2}$ and (s_{k-2}, y_{k-2}) . Proceeding recursively, $\tilde{\mathcal{H}}^k$ is a function of the initial $\tilde{\mathcal{H}}^0$ and all previous k curvature pairs $\{(s_j, y_j)\}_{j=0}^{k-1}$. Meanwhile, L-BFGS restricts use to the most recent L pairs $\{(s_j, y_j)\}_{j=k-L}^{k-1}$ since (s_j, y_j) with $j < k-L$ are likely to have little curvature information. Based on this idea, L-BFGS performs L updates by the initial $\tilde{\mathcal{H}}^0$. We use the k pairs $\{(s_j, y_j)\}_{j=0}^{k-1}$ when $k < L$.

Now, we consider the final calculation of $\tilde{\mathcal{H}}^k$ used for \mathcal{H}_t^k in the inner iterations of k -th outer epoch using the L most recent curvature pairs. Here, since this calculation is executed

at $T_{\tilde{w}^k}\mathcal{M}$ and a Riemannian manifold is in general not a vector space, all the L curvature pairs must be located at $T_{\tilde{w}^k}\mathcal{M}$. To this end, just after the curvature pair is calculated in (2) and (3), the past $(L-1)$ pairs of $\{(s_j^k, y_j^k)\}_{j=k-L+1}^{k-1} \in T_{\tilde{w}^k}\mathcal{M}$ are transported into $T_{\tilde{w}^{k+1}}\mathcal{M}$ by the same vector transport \mathcal{T}_{η_k} used when calculating s_k^{k+1} and y_k^{k+1} . It should be emphasized that this transport is necessary only for every outer epoch instead of every inner loop, and results in drastic reduction of computational complexity in comparison with the straightforward extension of the Euclidean stochastic L-BFGS [24] into the manifold setting. Consequently, the update is defined as

$$\begin{aligned}\tilde{\mathcal{H}}^k &= ((\check{\mathcal{V}}_{k-1}^k)^{\flat} \cdots (\check{\mathcal{V}}_{k-L}^k)^{\flat}) \check{\mathcal{H}}_0^k (\check{\mathcal{V}}_{k-L}^k \cdots \check{\mathcal{V}}_{k-1}^k) \\ &\quad + \rho_{k-2} (\check{\mathcal{V}}_{k-1}^k)^{\flat} s_{k-2}^k (s_{k-2}^k)^{\flat} (\check{\mathcal{V}}_{k-1}^k) + \rho_{k-1} s_{k-1}^k (s_{k-1}^k)^{\flat},\end{aligned}$$

where $\check{\mathcal{V}}_j^k = \text{id} - \rho_j y_j^k (s_j^k)^{\flat}$, and $\check{\mathcal{H}}_0^k$ is the initial inverse Hessian approximation. id is the identity mapping. Because $\check{\mathcal{H}}_0^k$ is not necessarily $\check{\mathcal{H}}^{k-L}$, and because it is any positive definite self-adjoint operator, we use $\check{\mathcal{H}}_0^k = \langle s_{k-1}^k, y_{k-1}^k \rangle_{\tilde{w}^k} / \langle y_{k-1}^k, y_{k-1}^k \rangle_{\tilde{w}^k} \text{id}$ similar to the Euclidean case. The practical update of $\tilde{\mathcal{H}}^k$ uses *two-loop recursion* algorithm [26] in Algorithm A.1 of the supplementary material.

Cautious update: Euclidean L-BFGS fails on non-convex problems because the Hessian approximation has eigenvalues that are away from zero and are not uniformly bounded above. To circumvent this issue, *cautious update* has been proposed in the Euclidean space [27]. By following this, we skip the update of the curvature pair when the following condition is not satisfied;

$$\langle y_k^{k+1}, s_k^{k+1} \rangle_{\tilde{w}^{k+1}} \geq \epsilon \|s_k^{k+1}\|_{\tilde{w}^{k+1}}^2, \quad (4)$$

where $\epsilon > 0$ is a predefined constant parameter. According to this update, the positive definiteness of $\tilde{\mathcal{H}}^k$ is guaranteed as far as $\tilde{\mathcal{H}}^{k-1}$ is positive definite.

Second-order modified stochastic gradient $\mathcal{H}_t^k \xi_t^k$: R-SVRG transports $\text{grad} f(\tilde{w}^k)$ and $\text{grad} f_{i_t^k}(\tilde{w}^k)$ at $T_{\tilde{w}^k}\mathcal{M}$ into $T_{w_t^k}\mathcal{M}$ to add them to $\text{grad} f_{i_t^k}(w_t^k)$ at $T_{w_t^k}\mathcal{M}$. If we follow the same strategy, we must also transport L pairs of $\{(s_j^k, y_j^k)\}_{j=k-L}^{k-1} \in T_{\tilde{w}^k}\mathcal{M}$ into the current $T_{w_t^k}\mathcal{M}$ at every inner iteration. Addressing this problem and the fact that both the full gradient and the curvature pairs belong to the same tangent space $T_{\tilde{w}^k}\mathcal{M}$, we transport $\text{grad} f_{i_t^k}(w_t^k)$ from $T_{w_t^k}\mathcal{M}$ into $T_{\tilde{w}^k}\mathcal{M}$, and complete all the calculations on $T_{\tilde{w}^k}\mathcal{M}$. More specifically, after transporting $\text{grad} f_{i_t^k}(w_t^k)$ as $(\mathcal{T}_{\tilde{\eta}_t^k})^{-1} \text{grad} f_{i_t^k}(w_t^k)$ from w_t^k to \tilde{w}^k using $\tilde{\eta}_t^k (= R_{\tilde{w}^k}^{-1}(w_t^k))$, the modified stochastic gradient $\tilde{\xi}_t^k \in T_{\tilde{w}^k}\mathcal{M}$ is computed as

$$\tilde{\xi}_t^k = (\mathcal{T}_{\tilde{\eta}_t^k})^{-1} \text{grad} f_{i_t^k}(w_t^k) - (\text{grad} f_{i_t^k}(\tilde{w}^k) - \text{grad} f(\tilde{w}^k)).$$

After calculating $\tilde{\mathcal{H}}_t^k \tilde{\xi}_t^k \in T_{\tilde{w}^k}\mathcal{M}$ using the two-loop recursion algorithm, we obtain $\mathcal{H}_t^k \xi_t^k \in T_{w_t^k}\mathcal{M}$ by transporting $\tilde{\mathcal{H}}_t^k \tilde{\xi}_t^k$ to $T_{w_t^k}\mathcal{M}$ by $\mathcal{T}_{\tilde{\eta}_t^k} \tilde{\mathcal{H}}_t^k \tilde{\xi}_t^k$. Finally, we update w_{t+1}^k from w_t^k as $w_{t+1}^k = R_{w_t^k}(-\alpha_t^k \mathcal{H}_t^k \xi_t^k)$. It should be noted that, although $-\xi_t^k$ is not generally guaranteed as a descent direction, $\mathbb{E}_{i_t^k}[-\xi_t^k] = -\text{grad} f(w_t^k)$ is a descent direction. Furthermore, the positive definiteness of \mathcal{H}_t^k yields that $-\mathcal{H}_t^k \xi_t^k$ is an average descent direction due to $\mathbb{E}_{i_t^k}[-\mathcal{H}_t^k \xi_t^k] = -\mathcal{H}_t^k \text{grad} f(w_t^k)$.

3 Convergence analysis

This section presents convergence analyses on both non-convex and retraction-convex functions under retraction and vector transport operations.. The concrete proofs are in the supplementary file.

Assumption 1. We assume below [21];

(1.1) The objective function f and its components f_1, \dots, f_N are twice continuously differentiable.

(1.2) For a sequence $\{w_t^k\}$ generated by Algorithm 1, there exists a compact and connected set $K \subset \mathcal{M}$ such that $w_t^k \in K$ for all $k, t \geq 0$. Also, for each $k \geq 1$, there exists a totally retractive neighborhood Θ_k of \tilde{w}^k such that w_t^k stays in Θ_k for any $t \geq 0$, where the ρ -totally retractive neighborhood Θ of w is a set such that for all $z \in \Theta$, $\Theta \subset R_z(\mathbb{B}(0_z, \rho))$, and $R_z(\cdot)$ is a diffeomorphism on $\mathbb{B}(0_z, \rho)$, which is the ball in $T_w\mathcal{M}$ with center 0_z and radius ρ , where 0_z is the zero vector in $T_z\mathcal{M}$. Furthermore, suppose that there exists $I > 0$ such that $\inf_{k \geq 1} \{\sup_{z \in \Theta_k} \|R_{\tilde{w}^k}^{-1}(z)\|_{\tilde{w}^k}\} \geq I$.

(1.3) The sequence $\{w_t^k\}$ continuously remains in ρ -totally retractive neighborhood Θ of critical point w^* and f is retraction-smooth with respect to retraction R in Θ . Here, f is said to be retraction-smooth in Θ if $f(R_w(t\eta_w))$ for all $w \in \mathcal{M}$, i.e., there exists a constant $0 < \Lambda$ such that $\frac{d^2 f(R_w(t\eta_w))}{dt^2} \leq \Lambda$, for all $w \in \Theta$, all $\|\eta_w\|_w = 1$, and all t such that $R_w(\tau\eta_w) \in \Theta$ for all $\tau \in [0, t]$.

(1.4) The vector transport \mathcal{T} is isometric on \mathcal{M} . It satisfies $\langle \mathcal{T}_{\xi_w} \eta_w, \mathcal{T}_{\xi_w} \zeta_w \rangle_{R_w(\xi_w)} = \langle \eta_w, \zeta_w \rangle_w$ for any $w \in \mathcal{M}$ and $\xi_w, \eta_w, \zeta_w \in T_w\mathcal{M}$.

(1.5) There exists a constant c_0 such that the vector transport \mathcal{T} satisfies the following conditions for all $w, z \in \mathcal{U}$, which is some neighborhood of an arbitrary point $\bar{w} \in \mathcal{M}$: $\|\mathcal{T}_{\eta_w} - \mathcal{T}_{R_{\eta_w}}\| \leq c_0 \|\eta_w\|_w$, $\|\mathcal{T}_{\eta_w}^{-1} - \mathcal{T}_{R_{\eta_w}}^{-1}\| \leq c_0 \|\eta_w\|_w$, where \mathcal{T}_R denotes the differentiated retraction, i.e., $\mathcal{T}_{R_{\zeta_w}} \xi_w = DR_w(\zeta_w)[\xi_w]$ with $\xi_w \in T_w\mathcal{M}$, and $\eta_w = R_w^{-1}(z)$.

(1.6) Riemannian stochastic gradient is bounded as $\mathbb{E}_{i_t^k} [\|\text{grad} f_{i_t^k}(w_t^k)\|_{w_t^k}^2] < C^2$ as [14–16].

Essential inequalities. We briefly summarize essential inequalities. They are detailed in the supplementary material. For all $w, z \in \mathcal{U}$, which is a neighborhood of \bar{w} , the difference between the parallel translation and the vector transport is given with a constant θ as (Lemma C.14)

$$\|\mathcal{T}_\eta \xi - P_\eta \xi\|_z \leq \theta \|\xi\|_w \|\eta\|_w, \quad (5)$$

where $\xi, \eta \in T_w\mathcal{M}$ and $R_w(\eta) = z$. Similarly, as for the difference between the exponential mapping and the retraction, there exist $\tau_1 > 0$, $\tau_2 > 0$ for all $w \in \mathcal{U}$ and all small length of $\xi \in T_w\mathcal{M}$ such that (Lemma C.15)

$$\tau_1 \text{dist}(w, R_w(\xi)) \leq \|\xi\|_w \leq \tau_2 \text{dist}(w, R_w(\xi)). \quad (6)$$

Then, the variance of ξ_t^k is upper bounded by (Lemma D.9)

$$\mathbb{E}_{i_t^k} [\|\xi_t^k\|_{w_t^k}^2] \leq 4(\beta^2 + \tau_2^2 C^2 \theta^2) (7(\text{dist}(w_t^k, w^*))^2 + 4(\text{dist}(\tilde{w}^k, w^*))^2), \quad (7)$$

where C is the constant of Assumption 1, β is a Lipschitz constant, and θ is the constant in (5). Finally, there exist $0 < \gamma < \Gamma$ such that (Proposition C.7 for non-convex functions and Proposition D.6 for retraction-convex functions)

$$\gamma \text{id} \preceq \mathcal{H}_t^k \preceq \text{Fid}, \quad (8)$$

where the $A \preceq B$ with $A, B \in \mathbb{R}^{n \times n}$ means that $B - A$ is positive semidefinite.

Now, we first present a global convergence analysis to a critical point starting from any initialization point, which is common in a non-convex setting with additional but mild assumptions;

Assumption 2. *We assume that f is bounded below by a scalar f_{\inf} , and a decaying step-size sequence $\{\alpha_t^k\}$ satisfies $\sum \alpha_t^k = \infty$ and $\sum (\alpha_t^k)^2 < \infty$. Additionally, since Θ is compact, all continuous functions on Θ can be bounded. Therefore, there exists $S > 0$ such that for all $w \in \Theta$ and $n \in N$, we have $\|\text{grad}f(w)\|_w \leq S$ and $\|\text{grad}f_n(w)\|_w \leq S$.*

Theorem 3.1 (Global convergence analysis on non-convex functions). *Let \mathcal{M} be a Riemannian manifold and $w^* \in \mathcal{M}$ be a non-degenerate local minimizer of f . Consider Algorithm 1 and suppose Assumptions 1 and 2, and that the mapping $w \mapsto \|\text{grad}f(w)\|_w^2$ has the positive real number that the largest eigenvalue of its Riemannian Hessian is bounded for all $w \in \mathcal{M}$. Then, we have $\lim_{k \rightarrow \infty} \mathbb{E}[\|\text{grad}f(w_t^k)\|_{w_t^k}^2] = 0$.*

We next present a global convergence rate analysis. This requires an strict selection of a fixed step size satisfying the condition below, but, instead, provides a convergence *rate* under it.

Theorem 3.2 (Global convergence rate analysis on non-convex functions). *Let \mathcal{M} be a Riemannian manifold and $w^* \in \mathcal{M}$ be a non-degenerate local minimizer of f . Consider Algorithm 1 with Option II and IV, and suppose Assumption 1. Let the constants θ in (5), τ_1 and τ_2 in (6), and β , and C in (7). Λ is the constant Assumption 1.3, and γ_{nc} and Γ_{nc} are the constants γ and Γ in (8). Set $\nu = \frac{\sqrt{\beta^2 + \tau_2^2 C^2 \theta^2} \Gamma_{nc} \tau_1}{N^{a_1/2}} \zeta^{1-a_2}$ and $\alpha_t^k = \alpha = \frac{\mu_0 \tau_1}{\sqrt{\beta^2 + \tau_2^2 C^2 \theta^2} N^{a_1} \Gamma_{nc} \zeta^{a_2}}$, where $0 < a_1 < 1$, and $0 < a_2 < 2$. Given sufficiently small $\mu_0 \in (0, 1)$, suppose that $\varrho > 0$ is chosen such that $\frac{\sqrt{\beta^2 + \tau_2^2 C^2 \theta^2}}{\Lambda \Gamma_{nc}} \gamma_{nc} \left(1 - \frac{\varrho \Gamma_{nc}}{\mu_0 \gamma_{nc} \tau_1}\right) > \frac{2\mu_0(e-1)}{\zeta^{2-a_2} \tau_1} + \frac{\mu_0 \tau_1}{N^{a_1} \zeta^{a_2}} + \frac{4\mu_0^2(e-1)}{N^{\frac{3a_1}{2}} \zeta^{a_2} (2\tau_1 + 1)}$ holds. Set $m = \lfloor \frac{N^{3a_1/2}}{5\mu_0 \zeta^{1-a_2} \tau_1 (2\tau_1 + 1)} \rfloor$ and $T = mK$. Then, we have*

$$\mathbb{E}[\|\text{grad}f(w_{\text{sol}})\|^2] \leq \frac{\sqrt{\beta^2 + \tau_2^2 C^2 \theta^2} N^{a_1} \zeta^{a_2} [f(w^0) - f(w^*)]}{T \varrho}. \quad (9)$$

The total number of gradient evaluations is $\mathcal{O}(N^{a_1}/\epsilon)$ to obtain an ϵ -solution. The proof is given by extending those of [12, 15, 23].

As a final analysis, we present a local convergence rate in neighborhood of a local minimum by introducing additionally a *local* assumption for retraction-convexity below. This is also very common and standard in manifold optimization.

Assumption 3. *We assume that the objective function f is strongly retraction-convex with respect to R in Θ . Here, f is said to be strongly retraction-convex in Θ if $f(R_w(t\eta_w))$ for all $w \in \mathcal{M}$ and $\eta_w \in T_w \mathcal{M}$ is strongly convex, i.e., there exists a constant $0 < \lambda$ such that $\lambda \leq \frac{d^2 f(R_w(t\eta_w))}{dt^2}$, for all $w \in \Theta$, all $\|\eta_w\|_w = 1$, and all t such that $R_w(t\eta_w) \in \Theta$ for all $\tau \in [0, t]$. Additionally, the vector transport \mathcal{T} satisfies the locking condition, which is defined as*

$$\mathcal{T}_{\eta_w} \xi_w = \kappa \mathcal{T}_{R_{\eta_w}} \xi_w, \text{ where } \kappa = \frac{\|\xi_w\|_w}{\|\mathcal{T}_{R_{\eta_w}} \xi_w\|_{R_w(\eta_w)}}, \quad (10)$$

for all $\eta_w, \xi_w \in T_w \mathcal{M}$ and all $w \in \mathcal{M}$.

It should be noted that, if we extend this local assumption to the entire manifold, as R-SVRG [23], our rate below directly results in the global rate. However, such a global assumption is fairly restrictive in terms of what cost functions and manifolds can be considered, and hence, the standard manifold literature mostly focuses on local rate analysis. For example, R-SVRG [22] does not show a global rate on retraction-convex functions.

Theorem 3.3 (Local convergence rate analysis on retraction-convex functions). *Let \mathcal{M} be a Riemannian manifold and $w^* \in \mathcal{M}$ be a non-degenerate local minimizer of f . Suppose Assumption 1 holds. Λ and λ are constants in Assumption 1 and 3, respectively. Let the constants θ in (5), τ_1 and τ_2 in (6), and β , and C in (7). γ_c and Γ_c are the constants in (8). Let α be a positive number satisfying $\lambda\tau_1^2 > 2\alpha(\lambda^2\tau_1^2 - 14\alpha\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2))$ and $\gamma_c\lambda^2\tau_1^2 > 14\alpha\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2)$. It then follows that for any sequence $\{\tilde{w}^k\}$ generated by Algorithm 1 with Option I under a fixed step size $\alpha_t^k := \alpha$ and $m_k := m$ converging to w^* , there exists $0 < K_{th} < K$ such that for all $k > K_{th}$,*

$$\mathbb{E}[(\text{dist}(\tilde{w}^{k+1}, w^*))^2] \leq \frac{2(\Lambda\tau_2^2 + 16m\alpha^2\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2))}{m\alpha(\gamma_c\lambda^2\tau_1^2 - 14\alpha\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2))} \mathbb{E}[(\text{dist}(\tilde{w}^k, w^*))^2]. \quad (11)$$

The proof structure is different from that of [22, 23] due to the way of bounding of $\mathbb{E}[\|\xi_t^k\|^2]$ and the existence of \mathcal{H}_t^k . Additionally, comparing (11) with that of R-SVRG [22, 23], we notice the rate degradation. To the best of our knowledge, no theoretical rate result that is better than or equals to that of SVRG [8] has been also given in the Euclidean SQN-VR [17]. Thus, this issue is a common area of research in both the Euclidean and Riemannian settings to further improve the theoretical rate. However, it should be emphasized that R-SQN-VR shows much better performances than R-SVRG, especially on a ill-conditioned problem, as shown later in Figure 1.

4 Numerical comparisons

This section compares R-SQN-VR with R-SGD with a decaying step-size sequence and R-SVRG with a fixed step size. The decaying step-size sequence is $\alpha_k = \alpha(1 + \alpha\lfloor k/m_k \rfloor)^{-1}$, where $\lfloor \cdot \rfloor$ denotes the floor function. As references, we also compare them with two Riemannian batch methods, i.e., R-SD, which is the steepest descent algorithm on Riemannian manifolds with backtracking line search [1], and R-L-BFGS, which is the Riemannian L-BFGS with strong wolfe condition [20, 28]. All experiments are executed in Matlab on a 4.0 GHz Intel Core i7 PC with 16 GB RAM, and are stopped when the gradient norm gets below 10^{-8} or when they reach a predefined maximum iteration. All results except R-SD and R-L-BFGS are the *best-tuned* results from multiple choices of step sizes α and a fixed $\varsigma = 10^{-3}$. This paper addresses the Karcher mean computation problem of symmetric positive-definite (SPD) manifold, and the low-rank matrix completion (MC) problem on the Grassmann manifold. The details of the problems and manifolds are in the supplementary file.


Karcher mean problem on SPD manifold. The first comparison is the Karcher mean problem on SPD matrices [28]. All experiments use the batch size fixed to 1 and $L = 4$, and are initialized randomly and are stopped when the number of iterations reaches 10 for R-SVRG and R-SQN-VR, and 60 for others. α are tuned from $\{10^{-5}, \dots, 10^{-1}\}$. m_k and the batch size are $3N$ and 1, respectively. Figures 1(a) and (b) show the results of the optimality gap when $N = 500$ with $d = 3$ (**Case KM-1**) and the larger size case with $N = 1500$ (**Case KM-2**), respectively. These results reveal that R-SQN-VR outperforms others.

MC problem on Grassmann manifold. We first consider a synthetic dataset. The proposed algorithm is also compared with Grouse [2], a state-of-the-art stochastic gradient algorithm on the Grassmann manifold. Algorithms are initialized randomly as [29]. α are tuned from $\{10^{-3}, 5 \times 10^{-3}, \dots, 10^{-2}, 5 \times 10^{-2}\}$ for R-SGD, R-SVRG and R-SQN-VR, and $\{1, 10, 100\}$ for Grouse. We set explicitly the condition number, denoted as CN, of the matrix, which represents the ratio of the maximal and the minimal singular values of the matrix. We also set the over-sampling ratio (OS) for the number of known entries. The Gaussian noise is also added with the noise level σ as suggested in [29]. m_k and the batch size are set to $5N$ and 50, respectively. The maximum number of the outer iterations to stop is 100 for R-SVRG and R-SQN-VR, and $100(m_k + 1)$ for the others. This experiment evaluates the projection-based vector transport and the QR-decomposition-based retraction, which do not satisfy the locking condition, but is computationally efficient. The baseline problem instance (**Case MC-S1**) is the case of $N = 5000$, $d = 200$, rank $r = 5$, $L = 10$, $OS = 8$, $\sigma = 10^{-10}$ and $CN = 50$. Additionally, changing some parameters of those in **Case MC-S1**, we evaluate the lower-sampling case with $OS = 4$ (**Case MC-S2**), the ill-conditioning case with $CN = 100$ (**Case MC-S3**), the higher noise case with $\sigma = 10^{-6}$ (**Case MC-S4**), and the higher rank case with $r = 10$ (**Case MC-S5**). The results of the MSE on test set Φ , which is different from the training set Ω , are shown in Figures 1(c)-(h), respectively. This gives the prediction accuracy of missing elements. From the figures, we confirm the superior performance of R-SQN-VR. **Case MC-S6** for different memory sizes L reveals that the larger size does not always show better results, which is also noticed in [15]. Finally, we compare the algorithms on a real-world dataset, the MovieLens-1M dataset¹. It contains a million ratings for 3952 movies (N) of 6040 users (d). We further randomly split this set into 80/10/10 percent data out of the entire data as train/validation/test partitions. α is chosen from $\{10^{-5}, 5 \times 10^{-5}, \dots, 10^{-2}, 5 \times 10^{-2}\}$, the batch size is 50, $r = 10$, and $L = 10$. The algorithms are terminated when the MSE on the validation set starts to increase or the number of the outer iteration reaches 100. Figure 1(i) shows the result except Grouse, which faces issues with convergence on this set (**Case MC-R**). R-SQN-VR shows much faster convergences than others.


5 Conclusions

We have proposed a Riemannian stochastic quasi-Newton algorithm with variance reduction (R-SQN-VR) on manifolds that is well suited for finite-sum minimization problems. We presented a rigorous convergence analysis for taking the Hessian approximation into a variance reduction stochastic setting on a manifold. Our proposed algorithm makes the explicit use of retraction and vector transport operators on manifolds, which makes the proposed algorithm appealing on a wider number of manifolds. The numerical comparisons show the benefits of our proposed algorithm on a number of applications.


¹<http://grouplens.org/datasets/movielens/>




(a) Case KM-1: small size.




(b) Case KM-2: large size.




(c) Case MC-S1: baseline.




(d) Case MC-S2: low sampling.




(e) Case MC-S3: ill-conditioning.




(f) Case MC-S4: noisy data.



(g) Case MC-S5: higher rank.



(h) Case MC-S6: memory sizes.



(i) Case MC-R: MovieLens-1M.

Figure 1: Performance evaluations on Karcher mean (KM) problem and low-rank MC problem.

References

- [1] P.-A. Absil, R. Mahony, and R. Sepulchre. *Optimization Algorithms on Matrix Manifolds*. Princeton University Press, 2008.
- [2] L. Balzano, R. Nowak, and B. Recht. Online identification and tracking of subspaces from highly incomplete information. In *Allerton*, pages 704–711, 2010.
- [3] B. Mishra and R. Sepulchre. R3MC: A Riemannian three-factor algorithm for low-rank matrix completion. In *IEEE CDC*, pages 1137–1142, 2014.
- [4] H. Kasai and B. Mishra. Low-rank tensor completion: a Riemannian manifold preconditioning approach. In *ICML*, 2016.
- [5] G. Meyer, S. Bonnabel, and R. Sepulchre. Linear regression under fixed-rank constraints: A Riemannian approach. In *ICML*, 2011.
- [6] S. Bonnabel. Stochastic gradient descent on Riemannian manifolds. *IEEE Trans. on Automatic Control*, 58(9):2217–2229, 2013.
- [7] H. Robbins and S. Monro. A stochastic approximation method. *Ann. Math. Statistics*, pages 400–407, 1951.
- [8] R. Johnson and T. Zhang. Accelerating stochastic gradient descent using predictive variance reduction. In *NIPS*, pages 315–323, 2013.
- [9] N. L. Roux, M. Schmidt, and F. R. Bach. A stochastic gradient method with an exponential convergence rate for finite training sets. In *NIPS*, pages 2663–2671, 2012.
- [10] S. Shalev-Shwartz and T. Zhang. Stochastic dual coordinate ascent methods for regularized loss minimization. *JMLR*, 14:567–599, 2013.
- [11] A. Defazio, F. Bach, and S. Lacoste-Julien. SAGA: A fast incremental gradient method with support for non-strongly convex composite objectives. In *NIPS*, 2014.
- [12] S. J. Reddi, A. Hefny, S. Sra, B. Póczos, and A. Smola. Stochastic variance reduction for nonconvex optimization. In *ICML*, 2016.
- [13] N. N. Schraudolph, J. Yu, and S. Gunter. A stochastic quasi-Newton method for online convex optimization. In *AISTATS*, 2007.
- [14] A. Mokhtari and A. Ribeiro. RES: Regularized stochastic BFGS algorithm. *IEEE Trans. on Signal Process.*, 62(23):6089–6104, 2014.
- [15] X. Wang, S. Ma, D. Goldfarb, and W. Liu. Stochastic quasi-Newton methods for non-convex stochastic optimization. *arXiv preprint arXiv:1607.0123*, 2016.
- [16] R. H. Byrd, S. L. Hansen, J. Nocedal, and Y. Singer. A stochastic quasi-Newton method for large-scale optimization. *SIAM J. Optim.*, 26(2), 2016.
- [17] P. Moritz, R. Nishihara, and M. I. Jordan. A linearly-convergent stochastic L-BFGS algorithm. In *AISTATS*, pages 249–258, 2016.

- [18] R. Kolte, M. Erdogdu, and A. Ozgur. Accelerating SVRG via second-order information,. In *OPT2015*, 2015.
- [19] D. Gabay. Minimizing a differentiable function over a differential manifold. *Journal of Optimization Theory and Applications*, 37(2):177–219, 1982.
- [20] W. Ring and B. Wirth. Optimization methods on Riemannian manifolds and their application to shape space. *SIAM J. Optim.*, 22(2):596–627, 2012.
- [21] W. Huang, K. A. Gallivan, and P.-A. Absil. A Broyden class of quasi-Newton methods for Riemannian optimization. *SIAM J. Optim.*, 25(3):1660–1685, 2015.
- [22] H. Sato, H. Kasai, and B. Mishra. Riemannian stochastic variance reduced gradient. *arXiv preprint: arXiv:1702.05594*, 2017.
- [23] H. Zhang, S. J. Reddi, and S. Sra. Fast stochastic optimization on Riemannian manifolds. In *NIPS*, 2016.
- [24] A. Mokhtari and A. Ribeiro. Global convergence of online limited memory BFGS. *JMLR*, 16:3151–3181, 2015.
- [25] N. Boumal, B. Mishra, P.-A. Absil, and R. Sepulchre. Manopt: a Matlab toolbox for optimization on manifolds. *JMLR*, 15(1):1455–1459, 2014.
- [26] J. Nocedal and Wright S.J. *Numerical Optimization*. Springer, New York, USA, 2006.
- [27] D. Li and M. Fukushima. On the global convergence of BFGS method for nonconvex unconstrained optimization. *SIAM J. Optim.*, 11(4):1054–1064, 2011.
- [28] X. Yuana, P.-A. Huang, W. Absil, and K. A. Gallivan. A Riemannian limited-memory BFGS algorithm for computing the matrix geometric mean. In *ICCS*, 2016.
- [29] D. Kressner, M. Steinlechner, and B. Vandereycken. Low-rank tensor completion by Riemannian optimization. *BIT Numer. Math.*, 54(2):447–468, 2014.
- [30] B. Jeuris, R. Vandebril, and B. Vandereycken. A survey and comparison of contemporary algorithms for computing the matrix geometric mean. *ETNA*, 2012.
- [31] H Karcher. Riemannian center of mass and mollifier smoothing. *Comm. Pure Appl. Math.*, 30(5):509–541, 1977.
- [32] S. Jayasumana, R. Hartley, M. Salzmann, H. Li, and M. Harandi. Kernel methods on Riemannian manifolds with Gaussian RBF kernels. *IEEE Trans. Pattern Anal. Mach. Intell.*, 37(12), 2015.
- [33] N. Boumal and P.-A. Absil. Low-rank matrix completion via preconditioned optimization on the Grassmann manifold. *Linear Algebra and its Applications*, 475:200–239, 2015.
- [34] L. Bottou, F. Curtis, and J. Nocedal. Optimization mehtods for large-scale machine learning. *arXiv preprint: arXiv:1606.04838*, 2016.
- [35] H. Zhang and S. Sra. First-order methods for geodesically convex optimization. In *COLT*, 2016.

- [36] W. Huang, P.-A. Absil, and K. A. Gallivan. A Riemannian symmetric rank-one trust-region method. *Math. Program., Ser. A*, 150:179–216, 2015.
- [37] S. Boyd and L. Vandenberghe. *Convex Optimization*. Cambridge University Press, 2004.

A Problems and manifolds in numerical comparison

This section gives a brief explanation of the problems and the manifolds that are evaluated in the numerical comparisons in Section 4.

A.1 SPD manifold and Karcher mean problem

SPD manifold \mathcal{S}_{++}^d . Let \mathcal{S}_{++}^d be the manifold of $d \times d$ SPD matrices. If we endow \mathcal{S}_{++}^d with the Riemannian metric defined by

$$\langle \xi_{\mathbf{X}}, \eta_{\mathbf{X}} \rangle_{\mathbf{X}} = \text{trace}(\xi_{\mathbf{X}} \mathbf{X}^{-1} \eta_{\mathbf{X}} \mathbf{X}^{-1})$$

at $\mathbf{X} \in \mathcal{S}_{++}^d$, the SPD manifold \mathcal{S}_{++}^d becomes a Riemannian manifold. The explicit formula for the exponential mapping is given by

$$\text{Exp}_{\mathbf{X}}(\xi_{\mathbf{X}}) = \mathbf{X}^{1/2} \exp(\mathbf{X}^{-1/2} \xi_{\mathbf{X}} \mathbf{X}^{-1/2}) \mathbf{X}^{1/2}$$

for any $\xi_{\mathbf{X}} \in T_{\mathbf{X}} \mathcal{S}_{++}^d$ and $\mathbf{X} \in \mathcal{S}_{++}^d$. On the other hand, $R_{\mathbf{X}}(\xi_{\mathbf{X}}) = \mathbf{X} + \xi_{\mathbf{X}} + \frac{1}{2} \xi_{\mathbf{X}} \mathbf{X}^{-1} \xi_{\mathbf{X}}$ proposed in [30] is a retraction, which is symmetric positive-definite for all $\xi_{\mathbf{X}} \in T_{\mathbf{X}} \mathcal{S}_{++}^d$ and $\mathbf{X} \in \mathcal{S}_{++}^d$. The parallel translation on \mathcal{S}_{++}^d along $\eta_{\mathbf{X}}$ is given by

$$P_{\eta_{\mathbf{X}}}(\xi_{\mathbf{X}}) = \mathbf{X}^{1/2} \mathbf{Y} \mathbf{X}^{-1/2} \xi_{\mathbf{X}} \mathbf{X}^{-1/2} \mathbf{Y} \mathbf{X}^{1/2},$$

where $\mathbf{Y} = \exp(\mathbf{X}^{-1/2} \eta_{\mathbf{X}} \mathbf{X}^{-1/2} / 2)$. A more efficient algorithm that constructs an isometric vector transport is proposed based on a field of orthonormal tangent bases [28] while satisfying the locking condition (10). We use it in this experiment, and the details are in [21, 28]. The logarithm map of \mathbf{Y} at \mathbf{X} is given by

$$\text{Log}_{\mathbf{X}}(\mathbf{Y}) = \mathbf{X}^{1/2} \log(\mathbf{X}^{-1/2} \mathbf{Y} \mathbf{X}^{-1/2}) \mathbf{X}^{1/2} = \log(\mathbf{Y} \mathbf{X}^{-1}) \mathbf{X}.$$

Karcher mean problem on \mathcal{S}_{++}^d . The Karcher mean is introduced as a notion of *mean* on Riemannian manifolds by Karcher [31]. It generalizes the notion of an “average” on a manifold. Given N points on \mathcal{S}_{++}^d with matrix representations $\mathbf{Q}_1, \dots, \mathbf{Q}_N$, the Karcher mean is defined as the solution to the problem

$$\min_{\mathbf{X} \in \mathcal{S}_{++}^d} \frac{1}{N} \sum_{n=1}^N (\text{dist}(\mathbf{X}, \mathbf{Q}_n))^2,$$

$\min_{\mathbf{X} \in \mathcal{S}_{++}^d} \frac{1}{N} \sum_{n=1}^N (\text{dist}(\mathbf{X}, \mathbf{Q}_n))^2$, where $\text{dist}(p, q) = \|\log(p^{-1/2} q p^{-1/2})\|_F$ represents the distance along the corresponding geodesic between the elements on \mathcal{S}_{++}^d with respect to the affine-invariant metric. The gradient of the loss function is computed as $\frac{2}{N} \sum_{n=1}^N -\log(\mathbf{Q}_n \mathbf{X}^{-1}) \mathbf{X}$. The Karcher mean on \mathcal{S}_{++}^d is frequently used for computer vision problems, such as visual object categorization and pose categorization [32]. Since recursive calculations are needed with each visual image, stochastic gradient algorithms become an appealing choice for large datasets.

A.2 Grassmann manifold and MC problem

Grassmann manifold $\text{Gr}(r, d)$. A point on the Grassmann manifold is an equivalence class represented by a $d \times r$ orthogonal matrix \mathbf{U} with orthonormal columns, i.e., $\mathbf{U}^T \mathbf{U} = \mathbf{I}$. Two orthogonal matrices express the same element on the Grassmann manifold if they are related by right multiplication of an $r \times r$ orthogonal matrix $\mathbf{O} \in \mathcal{O}(r)$. Equivalently, an element of $\text{Gr}(r, d)$ is identified with a set of $d \times r$ orthogonal matrices $[\mathbf{U}] := \{\mathbf{U}\mathbf{O} : \mathbf{O} \in \mathcal{O}(r)\}$. That is, $\text{Gr}(r, d) := \text{St}(r, d)/\mathcal{O}(r)$, where $\text{St}(r, d)$ is the *Stiefel manifold* that is the set of matrices of size $d \times r$ with orthonormal columns. The Grassmann manifold has the structure of a Riemannian quotient manifold [1, Section 3.4].

The exponential mapping for the Grassmann manifold from $\mathbf{U}(0) := \mathbf{U} \in \text{Gr}(r, d)$ in the direction of $\xi \in T_{\mathbf{U}(0)}\text{Gr}(r, d)$ is given in a closed form as [1, Section 5.4]

$$\mathbf{U}(t) = [\mathbf{U}(0)\mathbf{V} \ \mathbf{W}] \begin{bmatrix} \cos t\Sigma \\ \sin t\Sigma \end{bmatrix} \mathbf{V}^T,$$

where $\xi = \mathbf{W}\Sigma\mathbf{V}^T$ is the singular value decomposition (SVD) of ξ with rank r . The $\sin(\cdot)$ and $\cos(\cdot)$ operations are performed only on the diagonal entries. The parallel translation of $\zeta \in T_{\mathbf{U}(0)}\text{Gr}(r, d)$ on the Grassmann manifold along $\gamma(t)$ with $\dot{\gamma}(0) = \mathbf{W}\Sigma\mathbf{V}^T$ is given in a closed form by

$$\zeta(t) = \left([\mathbf{U}(0)\mathbf{V} \ \mathbf{W}] \begin{bmatrix} -\sin t\Sigma \\ \cos t\Sigma \end{bmatrix} \mathbf{W}^T + (\mathbf{I} - \mathbf{W}\mathbf{W}^T) \right) \zeta.$$

The logarithm map of $\mathbf{U}(t)$ at $\mathbf{U}(0)$ on the Grassmann manifold is given by

$$\xi = \text{Log}_{\mathbf{U}(0)}(\mathbf{U}(t)) = \mathbf{W} \arctan(\Sigma) \mathbf{V}^T,$$

where $\mathbf{W}\Sigma\mathbf{V}^T$ is the SVD of $(\mathbf{U}(t) - \mathbf{U}(0)\mathbf{U}(0)^T\mathbf{U}(t))(\mathbf{U}(0)^T\mathbf{U}(t))^{-1}$ with rank r . Furthermore, a popular retraction is

$$R_{\mathbf{U}(0)}(\xi) = \text{qf}(\mathbf{U}(0) + t\xi) \quad (= \mathbf{U}(t))$$

which extracts the orthonormal factor based on QR decomposition, and a popular vector transport uses an orthogonal projection of $t\xi$ to the horizontal space at $\mathbf{U}(t)$, i.e., $(\mathbf{I} - \mathbf{U}(t)\mathbf{U}(t)^T)t\xi$ [1].

Matrix completion problem. The matrix completion problem is completing an incomplete matrix \mathbf{X} , say of size $d \times N$, from a small number of entries by assuming that the latent structure of the matrix is low-rank. If Ω is the set of known indices in \mathbf{X} , the rank- r matrix completion problem amounts to solving

$$\min_{\mathbf{U}, \mathbf{A}} \|\mathcal{P}_\Omega(\mathbf{U}\mathbf{A}) - \mathcal{P}_\Omega(\mathbf{X})\|_F^2,$$

where $\mathbf{U} \in \mathbb{R}^{d \times r}$, $\mathbf{A} \in \mathbb{R}^{r \times N}$, and the operator \mathcal{P}_Ω acts as $\mathcal{P}_\Omega(\mathbf{X}_{ij}) = \mathbf{X}_{ij}$ if $(i, j) \in \Omega$ and $\mathcal{P}_\Omega(\mathbf{X}_{ij}) = 0$ otherwise. Partitioning $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_N]$, the previous problem is equivalent to

$$\min_{\mathbf{U} \in \mathbb{R}^{d \times r}, \mathbf{a}_n \in \mathbb{R}^r} \frac{1}{N} \sum_{n=1}^N \|\mathcal{P}_{\Omega_n}(\mathbf{U}\mathbf{a}_n) - \mathcal{P}_{\Omega_n}(\mathbf{x}_n)\|_2^2,$$

where $\mathbf{x}_n \in \mathbb{R}^d$ and the operator \mathcal{P}_{Ω_n} is the sampling operator for the n -th column. Given \mathbf{U} , \mathbf{a}_n admits a closed form solution. Consequently, the problem only depends on the column space of \mathbf{U} and is on $\text{Gr}(r, d)$ [33].

B Two-loop Hessian inverse updating algorithm

The section summarizes the Riemannian two-loop Hessian inverse updating algorithm in Algorithm A.1. This is an straightforward extension of that in the Euclidean space explained in [26, Section 7.2].

Algorithm A.1 Hessian inverse updating

Require: Pair-updating counter t , memory depth τ , correction pairs $\{s_u^k, y_u^k\}_{u=k-\tau}^{k-1}$, gradient p .

- 1: $p_0 = p$.
- 2: $\mathcal{H}_k^0 = \chi_k \text{id} = \frac{\langle s_t^k, y_t^k \rangle}{\langle y_t^k, y_t^k \rangle} \text{id}$.
- 3: **for** $u = 0, 1, 2, \dots, \tau - 1$ **do**
- 4: $\rho_{k-u} = 1 / \langle s_{k-u-1}^k, y_{k-u-1}^k \rangle$.
- 5: $\alpha_u = \rho_{k-u-1} \langle s_{k-u-1}^k, p_u \rangle$.
- 6: $p_{u+1} = p_u - \alpha_u y_{k-u-1}^k$.
- 7: **end for**
- 8: $q_0 = \mathcal{H}_k^0 p_\tau$.
- 9: **for** $u = 0, 1, 2, \dots, \tau - 1$ **do**
- 10: $\beta_u = \rho_{k-\tau+u} \langle y_{k-\tau+u}^k, q_u \rangle$.
- 11: $q_{u+1} = q_u + (\alpha_{\tau-u-1} - \beta_u) s_{k-\tau+u}^k$.
- 12: **end for**
- 13: $q = q_\tau$.

C Proofs of convergence analysis on non-convex functions

This section presents the proof of the global convergence analysis on non-convex functions. Hereinafter, we use $\mathbb{E}[\cdot]$ to express expectation with respect to the joint distribution of all random variables. For example, w_t is determined by the realizations of the independent random variables $\{i_1, i_2, \dots, i_{t-1}\}$, the total expectation of $f(w_t)$ for any $t \in \mathbb{N}$ can be taken as $\mathbb{E}[f(w_t)] = \mathbb{E}_{i_1} \mathbb{E}_{i_2} \dots \mathbb{E}_{i_{t-1}}[f(w_t)]$. We also use $\mathbb{E}_{i_t}[\cdot]$ to denote an expected value taken with respect to the distribution of the random variable i_t . In addition, we omit the subscript \tilde{w}^k for a Riemannian metric $\langle \cdot, \cdot \rangle_{\tilde{w}^k}$ when the tangent space to be considered is clear.

C.1 Preliminary lemmas

This subsection first states some preliminary lemmas.

The literature [1] generalizes a Taylor's theorem to Riemannian manifolds. However, it addresses the exponential mapping instead of the retraction. Therefore, [21] applies Taylor's theorem on the retraction by newly introducing a function along a curve on the manifold. Here, we denote $f(R_{w_t^k}(\tau \eta_k / \|\eta_k\|_{w_t^k}))$ for a twice continuously differentiable objective function. From Taylor's theorem, we obtain below;

Lemma C.1 (In Lemma 3.2 in [21]). *Under Assumptions 1.1, 1.2, and 1.3, there exists Λ such that*

$$f(w_{t+1}^k) - f(w_t^k) \leq \langle \text{grad} f(w_t^k), \alpha_t^k \eta_k \rangle_{w_t^k} + \frac{1}{2} \Lambda (\alpha_t^k \|\eta_k\|_{w_t^k})^2. \quad (\text{A.1})$$

Proof. From Taylor's theorem, we have

$$\begin{aligned} & f(w_{t+1}^k) - f(w_t^k) \\ &= f(R_{w_t^k}(\alpha_t^k \eta_k)) - f(R_{w_t^k}(0)) \\ &= \left. \frac{d}{d\tau} f(R_{w_t^k}(\tau \eta_k / \|\eta_k\|_{w_t^k})) \right|_{\tau=0} \cdot \alpha_t^k \|\eta_k\|_{w_t^k} + \frac{1}{2} \frac{d^2}{d\tau^2} f(R_{w_t^k}(\tau \eta_k / \|\eta_k\|_{w_t^k})) \Big|_{\tau=p} \cdot (\alpha_t^k \|\eta_k\|_{w_t^k})^2 \\ &= \langle \text{grad} f(w_t^k), \alpha_t^k \eta_k \rangle_{w_t^k} + \frac{1}{2} \frac{d^2}{d\tau^2} f(R_{w_t^k}(\tau \eta_k / \|\eta_k\|_{w_t^k})) \Big|_{\tau=p} \cdot (\alpha_t^k \|\eta_k\|_{w_t^k})^2 \\ &\leq \langle \text{grad} f(w_t^k), \alpha_t^k \eta_k \rangle_{w_t^k} + \frac{1}{2} \Lambda (\alpha_t^k \|\eta_k\|_{w_t^k})^2, \end{aligned}$$

where $0 \leq p \leq \alpha_t^k \|\eta_k\|_{w_t^k}$, and Λ is the constant in Assumption 1.3. This completes the proof. \square

Lemma C.2. *Suppose Assumption 1 holds. Then there exists a constant $0 < v$ for all k such that*

$$v \leq \frac{\langle y_k, y_k \rangle}{\langle s_k, y_k \rangle}. \quad (\text{A.2})$$

Proof. The claim for the case $s_k = 0$ is obvious. Assume that $s_k \neq 0$ below. This is given by applying Cauchy-Schwarz inequality to the condition (4) recursively. More specifically, (4) yields that

$$\epsilon \|s_k\|^2 \leq \langle y_k, s_k \rangle \leq \|y_k\| \|s_k\|,$$

and considering the most left and right terms, we obtain

$$\|s_k\| \leq \frac{1}{\epsilon} \|y_k\|.$$

Substituting this into the above equation yields

$$\langle s_k, y_k \rangle \leq \|s_k\| \|y_k\| \leq \frac{1}{\epsilon} \|y_k\|^2.$$

Consequently, we obtain

$$\frac{\|y_k\|^2}{\langle s_k, y_k \rangle} \geq \epsilon \quad (=v).$$

This completes the claim by denoting ϵ as v . \square

Lemma C.3. *Suppose Assumption 1 holds. There exists a constant $0 < \Upsilon_{nc}$ for all k such that*

$$\frac{\langle y_k, y_k \rangle}{\langle s_k, y_k \rangle} \leq \Upsilon_{nc}. \quad (\text{A.3})$$

Proof. Most part of this proof is given in Lemma 3.9 of [21]. But, [21] uses the strongly retraction-convexity assumption for the final part. Therefore, we include that of [21] for completeness, and describe the final part with a slight modifications. This proof is also included for the subsequent analysis.

Define $y_k^P = \text{grad}f(\tilde{w}^{k+1}) - P_{\gamma_k}^{1 \leftarrow 0} \text{grad}f(\tilde{w}^k)$, where $\gamma_k(t) = R_{\tilde{w}^k}(t\eta_k)$, i.e., the retraction curve connecting \tilde{w}^k and \tilde{w}^{k+1} , and P_{γ_k} is the parallel translation along $\gamma_k(t)$. We have $\|P_{\gamma_k}^{1 \leftarrow 0} y_k^P - \bar{H}_k \eta_k\| \leq b_0 \|\eta_k\|^2 = b_0 \|s_k\|^2$, where $\bar{H}_k = \int_0^1 P_{\gamma_k}^{0 \leftarrow t} \text{Hess}f(\gamma_k(t)) P_{\gamma_k}^{t \leftarrow 0} dt$ and $b_0 > 0$. It follows that

$$\begin{aligned} \|y_k\| &\leq \|y_k - y_k^P\| + \|y_k^P\| \\ &= \|y_k - y_k^P\| + \|P_{\gamma_k}^{0 \leftarrow 1} y_k^P\| \\ &\leq \|y_k - y_k^P\| + \|P_{\gamma_k}^{0 \leftarrow 1} y_k^P - \bar{H}_k \eta_k\| + \|\bar{H}_k \eta_k\| \\ &\leq \|\text{grad}f(\tilde{w}^{k+1})/\kappa_k - \mathcal{T}_{\eta_k} \text{grad}f(\tilde{w}^k) - \text{grad}f(\tilde{w}^{k+1}) + P_{\gamma_k}^{0 \leftarrow 1} \text{grad}f(\tilde{w}^k)\| \\ &\quad + \|\bar{H}_k \eta_k\| + b_0 \|s_k\|^2 \\ &\leq \|\text{grad}f(\tilde{w}^{k+1})/\kappa_k - \text{grad}f(\tilde{w}^{k+1})\| + \|P_{\gamma_k}^{0 \leftarrow 1} \text{grad}f(\tilde{w}^k) - \mathcal{T}_{\eta_k} \text{grad}f(\tilde{w}^k)\| \\ &\quad + \|\bar{H}_k \eta_k\| + b_0 \|s_k\|^2 \\ &\leq b_1 \|s_k\| \|\text{grad}f(\tilde{w}^{k+1})\| + b_2 \|s_k\| \|\text{grad}f(\tilde{w}^k)\| + b_3 \|s_k\| + b_0 \|s_k\|^2 \\ &\leq b_4 \|s_k\|, \end{aligned} \quad (\text{A.4})$$

where b_1, b_2, b_3 , and $b_4 > 0$. Here, we directly obtain the following fact from (4) in the cautious update as

$$\frac{\|s_k\|^2}{\langle y_k, s_k \rangle} \leq \frac{1}{\epsilon}. \quad (\text{A.5})$$

Therefore, we finally obtain the upper bound of $\frac{\langle y_k, y_k \rangle}{\langle s_k, y_k \rangle}$ from (A.4) and (A.5) as

$$\frac{\langle y_k, y_k \rangle}{\langle s_k, y_k \rangle} = \frac{\|s_k\|^2}{\langle s_k, y_k \rangle} \cdot \frac{\|y_k\|^2}{\|s_k\|^2} \leq \frac{b_4^2}{\epsilon} \quad (= \Upsilon_{nc}). \quad (\text{A.6})$$

Denoting b_4^2/ϵ as Υ_{nc} , this completes the proof. \square

Remark C.4. From the proof of Lemma C.3, if the parallel translation is used for vector transport, i.e., $\mathcal{T} = P$, the first two terms in (A.4) are equal to zero, and the upper bound Υ_{nc} in (A.3) can get smaller than that of the case in the vector transport.

C.2 Eigenvalue bounds of \mathcal{H}_t^k on non-convex functions

This subsection presents Proposition C.7, which is an essential proposition that bounds the eigenvalues of \mathcal{H}_t^k at w_t^k , i.e., $\mathcal{H}_t^k := \mathcal{T}_{\tilde{\eta}_t^k} \circ \tilde{\mathcal{H}}^k \circ (\mathcal{T}_{\tilde{\eta}_t^k})^{-1}$. To this end, we particularly use the *Hessian approximation operator* $\tilde{\mathcal{B}}^k = (\tilde{\mathcal{H}}^k)^{-1}$ as opposed to $\tilde{\mathcal{H}}^k$. Since mentioned in the algorithm description, we consider the curvature information for $\tilde{\mathcal{H}}^k$ at \tilde{w}^k , i.e., every outer epoch, and reuse this $\tilde{\mathcal{H}}^k$ in the calculation of the second-order modified stochastic gradient $\mathcal{H}_t^k \xi_t^k$ at w_t^k . Thereby, the way of the proof consists of two steps as follows;

1. We first address the bounds of $\tilde{\mathcal{H}}^k$ at \tilde{w}^k . The main task of the proof is to bound the Hessian operator $\tilde{\mathcal{B}}^k = (\tilde{\mathcal{H}}^k)^{-1}$.
2. Next, we bound \mathcal{H}_t^k at w_t^k based on the bounds of $\tilde{\mathcal{H}}^k$ at \tilde{w}^k .

It should be noted that, in this subsection, the curvature pair $\{s_j^k, y_j^k\}_{j=k-L}^{k-1} \in T_{\tilde{w}^k} \mathcal{M}$ is simply notated as $\{s_j, y_j\}_{j=k-L}^{k-1}$.

First, we attempt to bound $\text{trace}(\hat{\tilde{\mathcal{B}}})$ in order to bound the eigenvalues of $\tilde{\mathcal{H}}^k$, where a *hat* denotes the coordinate expression of the operator. The basic structure of the proof follows stochastic L-BFGS methods in the Euclidean space, e.g., [15, 16, 24]. Nevertheless, some special treatments considering the Riemannian setting and the lemmas earlier are required. It should be noted that $\text{trace}(\hat{\tilde{\mathcal{B}}})$ does not depend on the chosen basis.

Lemma C.5 (Bounds of trace of $\tilde{\mathcal{B}}^k$). *Consider the recursion of $\tilde{\mathcal{B}}_u^k$ as*

$$\tilde{\mathcal{B}}_{u+1}^k = \tilde{\mathcal{B}}_u^k - \frac{\tilde{\mathcal{B}}_u^k s_{k-\tau+u} (\tilde{\mathcal{B}}_u^k s_{k-\tau+u})^b}{(\tilde{\mathcal{B}}_u^k s_{k-\tau+u})^b s_{k-\tau+u}} + \frac{y_{k-\tau+u}^b}{y_{k-\tau+u}^b s_{k-\tau+u}}, \quad (\text{A.7})$$

where $\tilde{\mathcal{B}}_u^k = \mathcal{T}_{\eta_k} \tilde{\mathcal{B}}_u^k (\mathcal{T}_{\eta_k})^{-1}$ for $u = 0, \dots, \tau - 1$. The Hessian approximation at k -th outer epoch is $\tilde{\mathcal{B}}^k = \tilde{\mathcal{B}}_\tau^k$ when $u = \tau - 1$. Then, consider the Hessian approximation $\tilde{\mathcal{B}}^k = \tilde{\mathcal{B}}_\tau^k$ in (A.7) with $\tilde{\mathcal{B}}_0^k = \gamma_k^{-1} \text{id}$. If Assumption 1 holds, the $\text{trace}(\hat{\tilde{\mathcal{B}}})$ in a coordinate expression of $\tilde{\mathcal{B}}^k$ is uniformly upper bounded for all $k \geq 1$ as

$$\text{trace}(\hat{\tilde{\mathcal{B}}}) \leq (M + \tau) \Upsilon_{nc}, \quad (\text{A.8})$$

where M is the dimension of \mathcal{M} . Here, a *hat* expression represents the coordinate expression of an operator.

Proof. The proof can be completed parallel to the Euclidean case [17]. We use a *hat* symbol in order to represent the coordinate expression of the operator $\tilde{\mathcal{B}}_{u+1}^k$ and $\tilde{\mathcal{B}}_u^k$ in update formula (A.7). Because \mathcal{T} is an isometric vector transport, \mathcal{T}_{η_k} is invertible for all k . Accordingly, $\text{trace}(\hat{\tilde{\mathcal{B}}})$ and $\det(\hat{\tilde{\mathcal{B}}})$ can be reformulated as

$$\text{trace}(\hat{\tilde{\mathcal{B}}}) = \text{trace}(\hat{\mathcal{T}}_{\eta_k} \hat{\tilde{\mathcal{B}}} \hat{\mathcal{T}}_{\eta_k}^{-1}) = \text{trace}(\hat{\tilde{\mathcal{B}}}), \quad (\text{A.9})$$

$$(\text{A.10})$$

We first consider the trace lower bound of $\text{trace}(\hat{\mathcal{B}}_\tau^k)$ from (A.9) and (A.7) as

$$\begin{aligned}\text{trace}(\hat{\mathcal{B}}_{u+1}^k) &= \text{trace}(\hat{\mathcal{B}}_u^k) - \frac{\langle \hat{\mathcal{B}}_u^k \hat{s}_{k-\tau+u}, \hat{\mathcal{B}}_u^k \hat{s}_{k-\tau+u} \rangle}{\langle \hat{\mathcal{B}}_u^k \hat{s}_{k-\tau+u}, \hat{s}_{k-\tau+u} \rangle} + \frac{\langle y_{k-\tau+u}, y_{k-\tau+u} \rangle}{\langle y_{k-\tau+u}, s_{k-\tau+u} \rangle} \\ &= \text{trace}(\hat{\mathcal{B}}_u^k) - \frac{\|\hat{\mathcal{B}}_u^k \hat{s}_{k-\tau+u}\|^2}{\langle \hat{\mathcal{B}}_u^k \hat{s}_{k-\tau+u}, \hat{s}_{k-\tau+u} \rangle} + \frac{\|y_{k-\tau+u}\|^2}{\langle y_{k-\tau+u}, s_{k-\tau+u} \rangle},\end{aligned}$$

where we use the same notation $\langle \cdot, \cdot \rangle$ for the inner product with respect to local coordinates corresponding to the Riemannian metric. Here, the positive definiteness of $\hat{\mathcal{B}}_u^k$ guarantees the negativity of the second term. Therefore, the bound of the third term yields from Lemma C.3 as

$$\text{trace}(\hat{\mathcal{B}}_{u+1}^k) \leq \text{trace}(\hat{\mathcal{B}}_u^k) + \Upsilon_{nc}.$$

By calculating recursively this for $u = 0, \dots, \tau - 1$, we can conclude that

$$\text{trace}(\hat{\mathcal{B}}_u^k) \leq \text{trace}(\hat{\mathcal{B}}_0^k) + u\Upsilon_{nc}.$$

All that is left is to bound $\text{trace}(\hat{\mathcal{B}}_0^k)$. For this purpose, we consider the definition $\hat{\mathcal{B}}_0^k = \text{id}/\chi_k$, where, as a common choice in L-BFGS in the Euclidean, $\chi_k = \frac{\langle s_k, y_k \rangle}{\langle y_k, y_k \rangle}$, and we obtain

$$\text{trace}(\hat{\mathcal{B}}_0^k) = \text{trace}\left(\frac{\mathbf{I}}{\chi_k}\right) = \frac{M}{\chi_k} = M \frac{\langle y_k, y_k \rangle}{\langle s_k, y_k \rangle} \leq M\Upsilon_{nc}.$$

Consequently, we obtain

$$\text{trace}(\hat{\mathcal{B}}_u^k) \leq (M + u)\Upsilon_{nc}. \quad (\text{A.11})$$

Plugging $u = \tau$ into (A.11) yields the claim (A.8).

Thus, this completes the proof. \square

Now we prove the main lemma for Proposition C.7.

Lemma C.6 (Bounds of $\tilde{\mathcal{H}}^k$). *Suppose the constant $0 < \gamma_{nc} < \Gamma_{nc} < \infty$. If Assumption 1 holds, the eigenvalues of $\tilde{\mathcal{H}}^k$ is bounded by γ_{nc} and Γ_{nc} for all $k \geq 1$ as*

$$\gamma_{nc}\text{id} \preceq \tilde{\mathcal{H}}^k \preceq \Gamma_{nc}\text{id},$$

where γ_{nc} and Γ_{nc} are some positive constants.

Proof. We first state the lower bound part. The proof is obtained as parallel to the Euclidean case [24]. The sum of its eigenvalues of $\hat{\mathcal{B}}^k$ corresponds to the bounds on the trace. Here, we denote π_i as the i -th largest eigenvalue of the operator matrix $\hat{\mathcal{B}}^k$ for $1 \leq i \leq M$. From (A.8) in Lemma C.5, the sum of the eigenvalues of $\hat{\mathcal{B}}^k$ satisfies below

$$\sum_{i=1}^M \pi_i = \text{trace}(\hat{\mathcal{B}}^k) \leq (M + \tau)\Upsilon_{nc}. \quad (\text{A.12})$$

Because all the eigenvalues are positive due to the positive definiteness of $\hat{\mathcal{B}}^k$, it is obvious that every eigenvalue is less than the upper bound of the sum of all of the eigenvalues. Consequently, we obtain $\lambda_i \leq (M + \tau)\Upsilon_{nc}$ for all i , and finally obtain $\hat{\mathcal{B}}^k \preceq (M + \tau)\Upsilon_{nc}\text{id}$. The bounds in (A.12) implies that its inverse is the bound for the eigenvalues of $\hat{\mathcal{H}}^k = (\hat{\mathcal{B}}^k)^{-1}$ as

$$(\gamma_{nc}\text{id}) = \frac{1}{(M + \tau)\Upsilon_{nc}}\text{id} \preceq \tilde{\mathcal{H}}^k. \quad (\text{A.13})$$

Denoting $\frac{1}{(M + \tau)\Upsilon_{nc}}$ as γ_{nc} , we obtain the lower bound of the claim.

Next, we present the proof for the upper bound part by referring [15]. $\tilde{\mathcal{H}}^k$ is defined as

$$\tilde{\mathcal{H}}_u^k = (\text{id} - \rho_k y_k s_k^\flat)^\flat \tilde{\mathcal{H}}_{u-1}^k (\text{id} - \rho_k y_k s_k^\flat) + \rho_k s_k s_k^\flat, \quad (\text{A.14})$$

where $\tilde{\mathcal{H}}_{u-1}^k = \mathcal{T}_{\eta_k} \circ \tilde{\mathcal{H}}_{u-1}^k \circ \mathcal{T}_{\eta_k}^{-1}$, and $\rho_k = 1/\langle y_k, s_k \rangle$ [21]. Therefore, the coordinate representation of $\tilde{\mathcal{H}}_u^k$ is

$$\begin{aligned} \hat{\mathcal{H}}_u^k &= (\text{id} - \rho_k y_k s_k^T)^T \hat{\mathcal{H}}_{u-1}^k (\text{id} - \rho_k y_k s_k^T) + \rho_k s_k s_k^T, \\ &= \hat{\mathcal{H}}_{u-1}^k - \rho_k (\hat{\mathcal{H}}_{u-1}^k y_k s_k^T + s_k y^T \hat{\mathcal{H}}_{u-1}^k) + \rho_k s_k s_k^T + \rho_k^2 s_k y_k^T \hat{\mathcal{H}}_{u-1}^k y_k s_k^T. \end{aligned} \quad (\text{A.15})$$

Here, noticing below from the fact \mathcal{T} is isometric in Assumption 1.4,

$$\|\tilde{\mathcal{H}}_{u-1}^k\| = \|\mathcal{T}_{\eta_k} \circ \tilde{\mathcal{H}}_{u-1}^k \circ \mathcal{T}_{\eta_k}^{-1}\| = \|\tilde{\mathcal{H}}_{u-1}^k\|, \quad (\text{A.16})$$

we obtain below from (A.5) and (A.6),

$$\begin{aligned} \|\hat{\mathcal{H}}_u^k\| &\leq \|\hat{\mathcal{H}}_{u-1}^k\| + \frac{2\|\hat{\mathcal{H}}_{u-1}^k\|\|y_k\|\|s_k\|}{\langle s_k, y_k \rangle} + \frac{\|s_k\|^2}{\langle s_k, y_k \rangle} + \frac{\|s_k\|^2}{\langle s_k, y_k \rangle} \frac{\|\hat{\mathcal{H}}_{u-1}^k\|\|y_k\|^2}{\langle s_k, y_k \rangle} \\ &= \|\hat{\mathcal{H}}_{u-1}^k\| + 2\|\hat{\mathcal{H}}_{u-1}^k\| \left[\frac{\|y_k\|^2}{\langle s_k, y_k \rangle} \cdot \frac{\|s_k\|^2}{\langle s_k, y_k \rangle} \right]^{1/2} + \frac{\|s_k\|^2}{\langle s_k, y_k \rangle} + \frac{\|s_k\|^2}{\langle s_k, y_k \rangle} \frac{\|\hat{\mathcal{H}}_{u-1}^k\|\|y_k\|^2}{\langle s_k, y_k \rangle} \\ &= \left(1 + \frac{b_4}{\epsilon}\right)^2 \|\hat{\mathcal{H}}_{u-1}^k\| + \frac{1}{\epsilon} \\ &= q\|\hat{\mathcal{H}}_{u-1}^k\| + \frac{1}{\epsilon}, \end{aligned} \quad (\text{A.17})$$

where we denote $(1 + b_4/\epsilon)^2$ as q for simplicity. Because we consider the definition $\hat{\mathcal{B}}_0^k = \text{id}/\chi_k$, where, as a common choice in L-BFGS in the Euclidean, $\chi_k = \frac{\langle s_k, y_k \rangle}{\langle y_k, y_k \rangle}$, we obtain $\|\hat{\mathcal{H}}_0^k\|$ as

$$\|\hat{\mathcal{H}}_0^k\| = \|\chi_k\| = \frac{\langle s_k, y_k \rangle}{\|y_k\|^2} \leq \frac{1}{\epsilon},$$

where the last inequality uses (A.2) in Lemma C.2. Then, it follows that

$$\|\hat{\mathcal{H}}_1^k\| \leq q\|\hat{\mathcal{H}}_0^k\| + \frac{1}{\epsilon} \leq \frac{1}{\epsilon}(q + 1).$$

By recurrence relation, we calculate $\|\hat{\mathcal{H}}_u^k\|$ from $\|\hat{\mathcal{H}}_1^k\|$ as

$$\begin{aligned}\|\hat{\mathcal{H}}_u^k\| &\leq \left(\frac{1}{\epsilon}(q+1) - \frac{1}{\epsilon(1-q)}\right)q^{u-1} + \frac{1}{\epsilon(1-q)} \\ &= \frac{q^{u+1} - 1}{\epsilon(q-1)}.\end{aligned}$$

Consequently, plugging $u = \tau$ and considering $\hat{\mathcal{H}}_\tau^k = \hat{\mathcal{H}}^k$ and $\lambda_{\max}(\hat{\mathcal{H}}_\tau^k) = \|\hat{\mathcal{H}}_\tau^k\|$, we obtain below;

$$\tilde{\mathcal{H}}^k \preceq \frac{(1+b_4/\epsilon)^{2(\tau+1)} - 1}{\epsilon((1+b_4/\epsilon)^2 - 1)} \text{id} \quad (= \Gamma_{nc} \text{id}). \quad (\text{A.18})$$

Denoting the upper bound $\frac{(1+b_4/\epsilon)^{2(\tau+1)} - 1}{\epsilon((1+b_4/\epsilon)^2 - 1)}$ as Γ_{nc} , we obtain the upper bound part of the claim. This completes the proof. \square

Finally we present Proposition C.7.

Proposition C.7 (Bounds of \mathcal{H}_t^k on non-convex functions). *Consider the operator $\mathcal{H}_t^k := \mathcal{T}_{\tilde{\eta}_t^k} \circ \tilde{\mathcal{H}}^k \circ (\mathcal{T}_{\tilde{\eta}_t^k})^{-1}$. If Assumption 1 holds, the range of eigenvalues of \mathcal{H}_t^k is bounded by γ_{nc} and Γ_{nc} for all $k \geq 1, t \geq 1$, i.e.,*

$$\gamma_{nc} \text{id} \preceq \mathcal{H}_t^k \preceq \Gamma_{nc} \text{id}, \quad (\text{A.19})$$

where γ_{nc} and Γ_{nc} are some positive constants.

Proof. Considering $\mathcal{H}_t^k := \mathcal{T}_{\tilde{\eta}_t^k} \circ \tilde{\mathcal{H}}^k \circ (\mathcal{T}_{\tilde{\eta}_t^k})^{-1}$, where $\tilde{\eta}_t^k = R_{\tilde{w}^k}^{-1}(w_t^k)$, since $\mathcal{T}_{\tilde{\eta}_t^k}$ is a liner transformation operator, we can conclude that the eigenvalues of \mathcal{H}_t^k and $\tilde{\mathcal{H}}^k$ are identical. Actually, let hat expressions be representation matrices with some bases of $T_{w_t^k} \mathcal{M}$ and $T_{\tilde{w}^k} \mathcal{M}$, we have the relation below;

$$\begin{aligned}\det(\lambda \text{id} - \mathcal{H}_t^k) &= \det(\lambda \text{id} - \hat{\mathcal{T}}_{\tilde{\eta}_t^k} \hat{\mathcal{H}}^k (\hat{\mathcal{T}}_{\tilde{\eta}_t^k})^{-1}) \\ &= \det(\hat{\mathcal{T}}_{S_{\tilde{\eta}_t^k}} (\lambda \text{id} - \hat{\mathcal{H}}^k) (\hat{\mathcal{T}}_{\tilde{\eta}_t^k})^{-1}) \\ &= \det(\hat{\mathcal{T}}_{S_{\tilde{\eta}_t^k}}) \det(\lambda \text{id} - \hat{\mathcal{H}}^k) \det((\hat{\mathcal{T}}_{\tilde{\eta}_t^k})^{-1}) \\ &= \det(\hat{\mathcal{T}}_{S_{\tilde{\eta}_t^k}}) \det(\lambda \text{id} - \hat{\mathcal{H}}^k) \det(\hat{\mathcal{T}}_{\tilde{\eta}_t^k})^{-1} \\ &= \det(\lambda \text{id} - \hat{\mathcal{H}}^k).\end{aligned}$$

Therefore, Lemma C.6 directly yields the claim. This completes the proof. \square

C.3 Proof of global convergence analysis (Theorem 3.1)

This subsection shows the global convergence analysis of the proposed R-SQN-VR. This analysis partially extends the expectation-based analysis of SGD in the Euclidean space [34] into the proposed algorithm.

C.3.1 Essential lemmas

We first obtain the following lemma from (A.1) in Lemma C.1. Subsequently, $\mathbb{E}_{i_t^k}[f(w_{t+1}^k)]$ is a meaningful quantity because w_{t+1}^k depends on i_t^k through the update in Algorithm 1.

Lemma C.8. *Under Lemma C.1, the iterates of Algorithm 1 satisfy the following inequality for all $k \in \mathbb{N}$:*

$$\mathbb{E}_{i_t^k}[f(w_{t+1}^k)] - f(w_t^k) \leq -\alpha_t^k \langle \text{grad} f(w_t^k), \mathbb{E}_{i_t^k}[\mathcal{H}_t^k \xi_t^k] \rangle_{w_t^k} + \frac{1}{2}(\alpha_t^k)^2 \Lambda \mathbb{E}_{i_t^k}[\|\mathcal{H}_t^k \xi_t^k\|_{w_t^k}^2]. \quad (\text{A.20})$$

Proof. When $w_{t+1}^k = R_{w_t^k}(-\alpha_t^k \mathcal{H}_t^k \xi_t^k)$, substituting $-\mathcal{H}_t^k \xi_t^k$ into η_k , the iterates generated by Algorithm 1 satisfy from (A.1) in Lemma C.1

$$\begin{aligned} f(w_{t+1}^k) - f(w_t^k) &\leq \langle \text{grad} f(w_t^k), -\alpha_t^k \mathcal{H}_t^k \xi_t^k \rangle_{w_t^k} + \frac{1}{2} \Lambda \|\alpha_t^k \mathcal{H}_t^k \xi_t^k\|_{w_t^k}^2 \\ &= -\alpha_t^k \langle \text{grad} f(w_t^k), \mathcal{H}_t^k \xi_t^k \rangle_{w_t^k} + \frac{1}{2}(\alpha_t^k)^2 \Lambda \|\mathcal{H}_t^k \xi_t^k\|_{w_t^k}^2. \end{aligned} \quad (\text{A.21})$$

Taking expectations in the inequalities above with respect to the distribution of i_t^k , and noting that w_{t+1}^k , but not w_t^k , depends on i_t^k , we obtain the desired bound. \square

This lemma shows that, regardless of how Algorithm 1 arrived at w_t^k , the expected decrease in the objective function yielded by the k -th step is bounded above by a quantity involving: (i) the *expected directional derivative* of f at w_t^k along $-\mathcal{H}_t^k \xi_t^k$ and (ii) the *second moment* of $\mathcal{H}_t^k \xi_t^k$.

Next, we derive the following lemma;

Lemma C.9. *Under Assumptions 1 and 2, the sequence of average function $f(w_t^k)$ satisfies*

$$\mathbb{E}[f(w_{t+1}^k)] \leq f(w_t^k) - \alpha_t^k \gamma_{nc} \|\text{grad} f(w_t^k)\|_{w_t^k}^2 + \frac{9\Lambda(\alpha_t^k)^2 \Gamma_{nc}^2 S^2}{2}. \quad (\text{A.22})$$

Proof. Taking expectation (A.20) in Lemma C.8 with regard to w_t^k considering that \mathcal{H}_t^k is deterministic when w_t^k is given, we write

$$\begin{aligned} &\mathbb{E}[f(w_{t+1}^k)] \\ &\leq f(w_t^k) - \alpha_t^k \langle \text{grad} f(w_t^k), \mathcal{H}_t^k \mathbb{E}_{i_t^k}[\xi_t^k] \rangle_{w_t^k} + \frac{(\alpha_t^k)^2 \Lambda}{2} \mathbb{E}_{i_t^k}[\|\mathcal{H}_t^k \xi_t^k\|_{w_t^k}^2] \\ &\leq f(w_t^k) - \alpha_t^k \langle \text{grad} f(w_t^k), \mathcal{H}_t^k \text{grad} f(w_t^k) \rangle_{w_t^k} + \frac{(\alpha_t^k)^2 \Lambda}{2} \mathbb{E}_{i_t^k}[\|\mathcal{H}_t^k \xi_t^k\|_{w_t^k}^2] \\ &\leq f(w_t^k) - \alpha_t^k \langle \text{grad} f(w_t^k), \mathcal{H}_t^k \text{grad} f(w_t^k) \rangle_{w_t^k} + \frac{(\alpha_t^k)^2 \Lambda}{2} \mathbb{E}_{i_t^k}[\Gamma_{nc}^2 \|\xi_t^k\|_{w_t^k}^2] \\ &\leq f(w_t^k) - \alpha_t^k \gamma_{nc} \|\text{grad} f(w_t^k)\|_{w_t^k}^2 + \frac{9\Lambda(\alpha_t^k)^2 \Gamma_{nc}^2 S^2}{2}, \end{aligned} \quad (\text{A.23})$$

where the second inequality is obtained from $\mathbb{E}_{i_t^k}[\xi_t^k] = \text{grad} f(w_t^k)$ because ξ_t^k is an unbiased estimate of $\text{grad} f(w_t^k)$. The last inequality comes from Assumption 2 since

$$\begin{aligned} \|\xi_t^k\|_{w_t^k} &= \|\text{grad} f_{i_t^k}(w_t^k) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f_{i_t^k}(\tilde{w}^k)) + \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f(\tilde{w}^k))\|_{w_t^k} \\ &\leq S + S + S = 3S, \end{aligned} \quad (\text{A.24})$$

where $\tilde{\eta}_t^k \in T_{\tilde{w}^k} \mathcal{M}$ satisfies $R_{\tilde{w}^k}(\tilde{\eta}_t^k) = w_t^k$. This completes the proof. \square

Proposition C.10. *Under Assumptions 1 and 2, suppose that Algorithm 1 is run with a step-size sequence satisfying Assumption 2. Then, we have*

$$\mathbb{E} \left[\sum_{k=1}^K \sum_{t=1}^{m_k} \alpha_t^k \|\text{grad} f(w_t^k)\|_{w_t^k}^2 \right] < \infty. \quad (\text{A.25})$$

Proof. Taking the total expectation of (A.22) in Lemma C.9 yields

$$\mathbb{E}[f(w_{t+1}^k)] - \mathbb{E}[f(w_t^k)] \leq -\alpha_t^k \gamma_{nc} \mathbb{E}[\|\text{grad} f(w_t^k)\|_{w_t^k}^2] + \frac{9\Lambda(\alpha_t^k)^2 \Gamma_{nc}^2 S^2}{2}.$$

Summing both sides of this inequality for $\{w_1^1, \dots, w_{m_1}^1, \dots, w_1^{K-1}, \dots, w_{m_{K-1}}^{K-1}, w_1^K, \dots, w_{m_K}^K\}$ gives

$$\begin{aligned} f_{\inf} - f(w_1^1) &\leq \mathbb{E}[f(w_{t+1}^k)] - f(w_1^1) \\ &\leq -\gamma_{nc} \sum_{k=1}^K \sum_{t=1}^{m_k} \alpha_t^k \mathbb{E}[\|\text{grad} f(w_t^k)\|_{w_t^k}^2] + \frac{9\Lambda \Gamma_{nc}^2 S^2}{2} \sum_{k=1}^K \sum_{t=1}^{m_k} (\alpha_t^k)^2. \end{aligned}$$

Dividing by γ_{nc} and rearranging the terms, we obtain

$$\sum_{k=1}^K \sum_{t=1}^{m_k} \alpha_t^k \mathbb{E}[\|\text{grad} f(w_t^k)\|_{w_t^k}^2] \leq \frac{(f(w_1^1) - f_{\inf})}{\gamma_{nc}} + \frac{9\Lambda \Gamma_{nc}^2 S^2}{2\gamma_{nc}} \sum_{k=1}^K \sum_{t=1}^{m_k} (\alpha_t^k)^2.$$

The second condition of the decaying step-size sequence in Assumption 2, i.e., $\sum (\alpha_t^k)^2 < \infty$, implies that the right-hand side of this inequality converges to a *finite limit* when K increases. This completes the proof. \square

Then, we obtain the following proposition by taking (A.25) into account with the first condition of Assumption 2.

Proposition C.11. *Under Assumptions 1 and 2, suppose that Algorithm 1 is run with a step-size sequence satisfying Assumption 2. Then, we have*

$$\liminf_{k \rightarrow \infty} \mathbb{E}[\|\text{grad} f(w_t^k)\|_{w_t^k}^2] = 0. \quad (\text{A.26})$$

Proof. The proof is by contradiction. Assume that (A.26) does not hold. Then, there exists $\delta > 0$ such that $\mathbb{E}[\|\text{grad} f(w_t^k)\|_{w_t^k}^2] > \delta$ for all k sufficiently large, say, $k > N$. We have

$$\mathbb{E} \left[\sum_{k=1}^{\infty} \sum_{t=1}^{m_k} \alpha_t^k \|\text{grad} f(w_t^k)\|_{w_t^k}^2 \right] \geq \sum_{k=N}^{\infty} \sum_{t=1}^{m_k} \alpha_t^k \mathbb{E}[\|\text{grad} f(w_t^k)\|_{w_t^k}^2] > \delta \sum_{k=N}^{\infty} \sum_{t=1}^{m_k} \alpha_t^k = \infty.$$

This contradicts (A.25). \square

This implies that, for the R-SQN-VR with decaying step-sizes sequence, the expected gradient norms cannot stay bounded away from zero.

C.3.2 Main proof of Theorem 3.1

Theorem. 3.1. *Let \mathcal{M} be a Riemannian manifold and $w^* \in \mathcal{M}$ be a non-degenerate local minimizer of f . Consider Algorithm 1 and suppose Assumptions 1 and 2, and that the mapping $w \mapsto \|\text{grad}f(w)\|_w^2$ has the positive real number that the largest eigenvalue of its Riemannian Hessian is bounded for all $w \in \mathcal{M}$. Then, we have*

$$\lim_{k \rightarrow \infty} \mathbb{E}[\|\text{grad}f(w_t^k)\|_{w_t^k}^2] = 0.$$

Proof. We define $h(w)$ as $h(w) := \|\text{grad}f(w)\|_{w_t^k}^2$ and let Λ_h be the absolute value of the eigenvalue with the largest magnitude of the Hessian of h . Then, from Taylor's theorem, we obtain

$$h(w_{t+1}^k) - h(w_t^k) \leq -2\alpha_t^k \langle \text{grad}h(w_t^k), \text{Hess}f(w_t^k)[\mathcal{H}_t^k \xi_t^k] \rangle_{w_t^k} + \frac{1}{2}(\alpha_t^k)^2 \Lambda_h \|\mathcal{H}_t^k \xi_t^k\|_{w_t^k}^2.$$

Taking the expectation with respect to the distribution of i_t^k , we obtain below;

$$\begin{aligned} & \mathbb{E}_{i_t^k}[h(w_{t+1}^k)] - h(w_t^k) \\ & \leq -2\alpha_t^k \langle \text{grad}f(w_t^k), \mathbb{E}_{i_t^k}[\text{Hess}f(w_t^k)[\mathcal{H}_t^k \xi_t^k]] \rangle_{w_t^k} + \frac{1}{2}(\alpha_t^k)^2 \Lambda_h \mathbb{E}_{i_t^k}[\|\mathcal{H}_t^k \xi_t^k\|_{w_t^k}^2] \\ & = -2\alpha_t^k \langle \text{grad}f(w_t^k), \text{Hess}f(w_t^k)[\mathcal{H}_t^k \mathbb{E}_{i_t^k}[\xi_t^k]] \rangle_{w_t^k} + \frac{1}{2}(\alpha_t^k)^2 \Lambda_h \mathbb{E}_{i_t^k}[\|\mathcal{H}_t^k \xi_t^k\|_{w_t^k}^2] \\ & \leq 2\alpha_t^k \|\text{grad}f(w_t^k)\|_{w_t^k} \|\text{Hess}f(w_t^k)[\mathcal{H}_t^k \text{grad}f(w_t^k)]\|_{w_t^k} + \frac{1}{2}(\alpha_t^k)^2 \Lambda_h \Gamma_{nc}^2 \mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2] \\ & \leq 2\alpha_t^k \Lambda \Gamma_{nc} \|\text{grad}f(w_t^k)\|_{w_t^k}^2 + \frac{9}{2}(\alpha_t^k)^2 \Lambda_h S^2 \Gamma_{nc}^2, \end{aligned}$$

where the last inequality comes from $\|\text{Hess}f(w)[\mathcal{H}_t^k \text{grad}f(w_t^k)]\|_{w_t^k} \leq \Lambda \|\mathcal{H}_t^k \text{grad}f(w_t^k)\|_{w_t^k} \leq \Lambda \Gamma_{nc} \|\text{grad}f(w_t^k)\|_{w_t^k}$.

Taking the total expectation simply yields

$$\mathbb{E}[h(w_{t+1}^k)] - \mathbb{E}[h(w_t^k)] \leq 2\alpha_t^k \Lambda \Gamma_{nc} \mathbb{E}[\|\text{grad}f(w_t^k)\|_{w_t^k}^2] + \frac{9}{2}(\alpha_t^k)^2 \Lambda_h S^2 \Gamma_{nc}^2. \quad (\text{A.27})$$

Recall that Proposition C.10 establishes that the first component of this bound is the term of a convergent sum. The second component of this bound is also the term of a convergent sum since $\sum_{k=1}^{\infty} \sum_{t=1}^{m_k} (\alpha_t^k)^2$ converges. This means that again the result of Proposition C.10 can be applied. Therefore, the right-hand side of (A.27) is the term of a convergent sum. Let us now define $S_K^+ = \sum_{k=1}^K \sum_{t=1}^{m_k} \max(0, \mathbb{E}[h(w_{t+1}^k)] - \mathbb{E}[h(w_t^k)])$, and $S_K^- = \sum_{k=1}^K \sum_{t=1}^{m_k} \max(0, \mathbb{E}[h(w_t^k)] - \mathbb{E}[h(w_{t+1}^k)])$.

Since the bound (A.27) is positive and forms a convergent sum, the nondecreasing sequence S_K^+ is upper bounded and therefore converges. Since, for any $K \in \mathbb{N}$, one has $\mathbb{E}[h(w_K)] = h(w_1) + S_K^+ - S_K^- \geq 0$, the nondecreasing sequence S_K^- is upper bounded and therefore also converges. Therefore $\mathbb{E}[h(w_K)]$ converges. Consequently, this implies that this limit must be zero from Proposition C.11. This completes the proof. \square

C.4 Proof of global convergence rate analysis (Theorem 3.2)

The global convergence rate analysis on non-convex functions in the Euclidean SVRG is proposed in [12]. Its further extensions into the stochastic L-BFGS setting and the Riemannian setting are proposed in [15] and [23], respectively. The proof in this subsection mainly follows that in [12] by integrating its two extensions in [15, 23]. Besides that, the special and careful treatments for the retraction and the vector transport operations are particularly taken in the proof. The results for the exponential mapping and the parallel translation are given in the corresponding corollaries as a special case.

C.4.1 Preliminary lemmas

We first present some essential lemmas.

Lemma C.12 (Lemma 6 in [35]). *If a , b , and c are the side lengths of a geodesic triangle in an Alexandrov space with curvature lower-bounded by κ , and A is the angle between sides b and c , then*

$$a^2 \leq \frac{\sqrt{|\kappa|}c}{\tanh(\sqrt{|\kappa|}c)}b^2 + c^2 - 2bc \cos(A).$$

Lemma C.13 (In the proof of Lemma 3.9 in [21]). *Under Assumptions 1.1 and 1.2, there exists a constant $\beta > 0$ such that*

$$\|P_\gamma^{w \leftarrow z}(\text{grad} f(z)) - \text{grad} f(w)\|_w \leq \beta \text{dist}(z, w), \quad (\text{A.28})$$

where w and z are in Θ in Assumption 1.2 and γ is a curve $\gamma(t) := R_z(\tau\eta)$ for $\eta \in T_z\mathcal{M}$ defined by a retraction R on \mathcal{M} . $P_\gamma^{w \leftarrow z}(\cdot)$ is a parallel translation operator along the curve γ from z to w .

Note that the curve γ in this lemma is not necessarily the geodesic. The relation (A.28) is a generalization of the Lipschitz continuity condition. In addition, we specifically use β_0 when the curve is geodesic.

Lemma C.14 (Lemma 3.5 in [21]). *Let $\mathcal{T} \in C^0$ be a vector transport associated with the same retraction R as that of the parallel translation $P \in C^\infty$. Under Assumption 1.5, for any $\bar{w} \in \mathcal{M}$ there exists a constant $\theta > 0$ and a neighborhood \mathcal{U} of \bar{w} such that for all $w, z \in \mathcal{U}$,*

$$\|\mathcal{T}_\eta \xi - P_\eta \xi\|_z \leq \theta \|\xi\|_w \|\eta\|_w, \quad (\text{A.29})$$

where $\xi, \eta \in T_w\mathcal{M}$ and $R_w(\eta) = z$.

Modifying slightly Lemma 3 in [36], we obtain the following lemma.

Lemma C.15 (Lemma 3 in [36]). *Let \mathcal{M} be a Riemannian manifold endowed with retraction R and let $\bar{w} \in \mathcal{M}$. Then there exist $\tau_1 > 0$, $\tau_2 > 0$ and δ_{τ_1, τ_2} such that for all w in a sufficiently small neighborhood of \bar{w} and all $\xi \in T_w\mathcal{M}$ with $\|\xi\|_w \leq \delta_{\tau_1, \tau_2}$, the inequalities*

$$\tau_1 \text{dist}(w, R_w(\xi)) \leq \|\xi\|_w \leq \tau_2 \text{dist}(w, R_w(\xi)) \quad (\text{A.30})$$

hold.

C.4.2 Essential propositions

This subsection first presents an essential lemma about the bound of $\mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2]$, where the vector transport is carefully handled to give the lemma. Next, an important proposition C.18 is presented by extending [12, 15, 23]. It should be noted that we carefully treat the difference between the exponential case and the retraction case for Proposition C.18.

Lemma C.16. *Suppose Assumptions 1.1, 1.2, 1.3, 1.4, 1.5, and 1.7, which guarantee Lemmas C.13, C.14, and C.15 for $\bar{w} = w^*$. Let $\beta > 0$ be a constant such that*

$$\|P_\gamma^{w \leftarrow z}(\text{grad} f_n(z)) - \text{grad} f_n(w)\|_w \leq \beta \text{dist}(z, w), \quad w, z \in \Theta, \quad n = 1, 2, \dots, N. \quad (\text{A.31})$$

The existence of such β is guaranteed by Lemma C.13. Then, the upper bound of the variance of $\mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2]$ is given by

$$\mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2] \leq 4(\beta^2 + \tau_2^2 C^2 \theta^2)(\text{dist}(w_t^k, \tilde{w}^k))^2 + 2\|\text{grad} f(w_t^k)\|_{w_t^k}^2. \quad (\text{A.32})$$

Proof. The proof is partially similar to that of Lemma 5.8 in [22]. We first consider

$$\mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2] = \mathbb{E}_{i_t^k}[\|\text{grad} f_{i_t^k}(w_t^k) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f_{i_t^k}(\tilde{w}^k)) + \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f(\tilde{w}^k))\|_{w_t^k}^2]. \quad (\text{A.33})$$

The first and second terms in (A.33) is

$$\mathbb{E}_{i_t^k}[\|\text{grad} f_{i_t^k}(w_t^k) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f_{i_t^k}(\tilde{w}^k))\|_{w_t^k}^2] = \|\text{grad} f(w_t^k) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f(\tilde{w}^k))\|_{w_t^k}^2,$$

which is equivalent to

$$\mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f(\tilde{w}^k)) = \text{grad} f(w_t^k) - \mathbb{E}_{i_t^k}[\|\text{grad} f_{i_t^k}(w_t^k) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f_{i_t^k}(\tilde{w}^k))\|_{w_t^k}^2].$$

Plugging this into the third term of (A.33) yields

$$\begin{aligned} \mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2] &= \mathbb{E}_{i_t^k}[\|\text{grad} f_{i_t^k}(w_t^k) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f_{i_t^k}(\tilde{w}^k)) \\ &\quad - \mathbb{E}_{i_t^k}[\|\text{grad} f_{i_t^k}(w_t^k) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f_{i_t^k}(\tilde{w}^k))\|_{w_t^k}^2] + \text{grad} f(w_t^k)\|_{w_t^k}^2] \\ &\leq 2\mathbb{E}_{i_t^k}[\|\text{grad} f_{i_t^k}(w_t^k) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f_{i_t^k}(\tilde{w}^k)) \\ &\quad - \mathbb{E}_{i_t^k}[\|\text{grad} f_{i_t^k}(w_t^k) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f_{i_t^k}(\tilde{w}^k))\|_{w_t^k}^2] + 2\|\text{grad} f(w_t^k)\|_{w_t^k}^2] \\ &\leq 2\mathbb{E}_{i_t^k}[\|\text{grad} f_{i_t^k}(w_t^k) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f_{i_t^k}(\tilde{w}^k))\|_{w_t^k}^2] + 2\|\text{grad} f(w_t^k)\|_{w_t^k}^2, \end{aligned} \quad (\text{A.34})$$

where the second inequality comes from the fact that, for arbitrary random vector z on arbitrary tangent space, $\mathbb{E}[\|z - \mathbb{E}[z]\|^2] = \mathbb{E}[\|z\|^2] - \|\mathbb{E}[z]\|^2 \leq \mathbb{E}[\|z\|^2]$. Now, the first term in

the right-hand side is upper-bounded by the distance between \tilde{w}^k and w_t^k as

$$\begin{aligned}
& \mathbb{E}_{i_t^k} [\|\text{grad} f_{i_t^k}(w_t^k) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f_{i_t^k}(\tilde{w}^k))\|_{w_t^k}^2] \\
= & \mathbb{E}_{i_t^k} \left[\|\text{grad} f_{i_t^k}(w_t^k) - P^{w_t^k \leftarrow \tilde{w}^k}(\text{grad} f_{i_t^k}(\tilde{w}^k)) \right. \\
& \left. + P^{w_t^k \leftarrow \tilde{w}^k}(\text{grad} f_{i_t^k}(\tilde{w}^k)) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f_{i_t^k}(\tilde{w}^k))\|_{w_t^k}^2 \right] \\
\leq & 2\mathbb{E}_{i_t^k} \left[\|\text{grad} f_{i_t^k}(w_t^k) - P^{w_t^k \leftarrow \tilde{w}^k}(\text{grad} f_{i_t^k}(\tilde{w}^k))\|_{w_t^k}^2 \right] \\
& + 2\mathbb{E}_{i_t^k} \left[\|P^{w_t^k \leftarrow \tilde{w}^k}(\text{grad} f_{i_t^k}(\tilde{w}^k)) - \mathcal{T}_{\tilde{\eta}_t^k}(\text{grad} f_{i_t^k}(\tilde{w}^k))\|_{w_t^k}^2 \right] \\
\stackrel{(A.29)}{\leq} & 2\beta^2(\text{dist}(w_t^k, \tilde{w}^k))^2 + 2\mathbb{E}_{i_t^k} [\theta^2 \|\tilde{\eta}_t^k\|_{w_t^k}^2 \|\text{grad} f_{i_t^k}(w_t^k)\|_{w_t^k}^2] \\
= & 2\beta^2(\text{dist}(w_t^k, \tilde{w}^k))^2 + 2\theta^2 \|\tilde{\eta}_t^k\|_{w_t^k}^2 \mathbb{E}_{i_t^k} [\|\text{grad} f_{i_t^k}(w_t^k)\|_{w_t^k}^2] \\
\leq & 2\beta^2(\text{dist}(w_t^k, \tilde{w}^k))^2 + 2C^2\theta^2 \|\tilde{\eta}_t^k\|_{w_t^k}^2 \\
\stackrel{(A.30)}{\leq} & 2(\beta^2 + \tau_2^2 C^2 \theta^2)(\text{dist}(w_t^k, \tilde{w}^k))^2, \tag{A.35}
\end{aligned}$$

where the first inequality uses $\|a + b\| \leq 2\|a\|^2 + 2\|b\|^2$ for vector a and b , and the second inequality uses (A.29) in Lemma C.14. The third inequality uses Assumption 1.7, and the last inequality uses (A.30) in Lemma C.15. Substituting this into (A.34) yields the claimed statement. This complete the proof. \square

We obtain the counterpart result of Lemma C.16 for the parallel translation and the exponential mapping.

Corollary C.17. *Suppose Assumptions 1.1, 1.2, 1.3, 1.4, and 1.5 which guarantee Lemmas C.13 for $\bar{w} = w^*$. Consider $\mathcal{T} = P$ and $R = \text{Exp}$, i.e., the parallel translation and the exponential mapping case. Let $\beta > 0$ be a constant such that*

$$\|P_\gamma^{w \leftarrow z}(\text{grad} f_n(z)) - \text{grad} f_n(w)\|_w \leq \beta \text{dist}(z, w), \quad w, z \in \Theta, \quad n = 1, 2, \dots, N.$$

The existence of such β_0 is guaranteed by Lemma C.13. Then, the upper bound of the variance of $\mathbb{E}_{i_t^k} [\|\xi_t^k\|_{w_t^k}^2]$ is given by

$$\mathbb{E}_{i_t^k} [\|\xi_t^k\|_{w_t^k}^2] \leq 2\beta_0^2(\text{dist}(w_t^k, \tilde{w}^k))^2 + 2\|\text{grad} f(w_t^k)\|_{w_t^k}^2.$$

Proposition C.18. *Let \mathcal{M} be a Riemannian manifold and $w^* \in \mathcal{M}$ be a non-degenerate local minimizer of f (i.e., $\text{grad} f(w^*) = 0$ and the Hessian $\text{Hess} f(w^*)$ of f at w^* is positive definite). Suppose Assumption 1 holds. Assume also that w_t^k and w_{t+1}^k are sufficiently close to each other such that $\langle R_{w_t^k}^{-1}(w_{t+1}^k), \text{Exp}_{w_t^k}^{-1}(\tilde{w}^k) \rangle_{w_t^k} \leq \langle \text{Exp}_{w_t^k}^{-1}(w_{t+1}^k), \text{Exp}_{w_t^k}^{-1}(\tilde{w}^k) \rangle_{w_t^k} / \phi$ for some positive constant ϕ . Let the constants θ be in (5), τ_2 in (6), and β and C in (7). Let Λ be the constant in Assumption 1.3, and γ_{nc} and Γ_{nc} in (8). For $c_t^k, c_{t+1}^k, \nu_t > 0$, we set*

$$c_t^k = c_{t+1}^k (1 + \phi \alpha_t^k \nu_t + 4\zeta(\alpha_t^k)^2 \frac{\Gamma_{nc}^2}{\tau_1^2} (\beta^2 + \tau_2^2 C^2 \theta^2)) + 2(\alpha_t^k)^2 \Lambda \Gamma_{nc}^2 (\beta^2 + \tau_2^2 C^2 \theta^2). \tag{A.36}$$

We also define

$$\Delta_t := \alpha_t^k \left(\gamma_{nc} - \frac{\phi c_{t+1}^k \Gamma_{nc}^2}{\nu_t} - \alpha_t^k \Lambda \Gamma_{nc}^2 - 2c_{t+1}^k \zeta \alpha_t^k \frac{\Gamma_{nc}^2}{\tau_1^2} \right). \tag{A.37}$$

Let α_t^k, ν_t and c_{t+1}^k be defined such that $\Delta_t > 0$. It then follows that for any sequence $\{\tilde{w}_t^k\}$ generated by Algorithm 1 with option II and with a fixed step-size $\alpha_t^k := \alpha$ and $m_k := m$ converging to w^* , the expected squared norm of the Riemannian gradient, $\text{grad}f(w_t^k)$, satisfies the following bound as

$$\mathbb{E}[\|\text{grad}f(w_t^k)\|^2] \leq \frac{V_t^k - V_{t+1}^k}{\Delta_t}, \quad (\text{A.38})$$

where $V_t^k := \mathbb{E}[f(w_t^k) + c_t^k(\text{dist}(\tilde{w}_t^k, w_t^k))^2]$ for $0 \leq k \leq K - 1$.

Proof. We first obtain the following from (A.23) in Lemma C.9 as

$$\begin{aligned} \mathbb{E}[f(w_{t+1}^k)] &\leq f(w_t^k) - \alpha_t^k \langle \text{grad}f(w_t^k), \mathcal{H}_t^k \text{grad}f(w_t^k) \rangle_{w_t^k} + \frac{(\alpha_t^k)^2 \Lambda}{2} \mathbb{E}_{i_t^k}[\Gamma_{nc}^2 \|\xi_t^k\|_{w_t^k}^2] \\ &\leq f(w_t^k) - \alpha_t^k \gamma_{nc} \|\text{grad}f(w_t^k)\|_{w_t^k}^2 + \frac{(\alpha_t^k)^2 \Lambda \Gamma_{nc}^2}{2} \mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2]. \end{aligned} \quad (\text{A.39})$$

Next, we bound the expected squared distance between \tilde{w}_t^k and w_{t+1}^k , i.e., $\mathbb{E}[(\text{dist}(\tilde{w}_t^k, w_{t+1}^k))^2]$, from Lemma C.12 as

$$\begin{aligned} &\mathbb{E}[(\text{dist}(\tilde{w}_t^k, w_{t+1}^k))^2] \\ &\leq \mathbb{E}[\zeta(\text{dist}(w_t^k, w_{t+1}^k))^2 + (\text{dist}(w_t^k, \tilde{w}_t^k))^2 - 2\langle \text{Exp}_{w_t^k}^{-1}(w_{t+1}^k), \text{Exp}_{w_t^k}^{-1}(\tilde{w}_t^k) \rangle_{w_t^k}] \\ &\leq \mathbb{E}[\zeta(\text{dist}(w_t^k, w_{t+1}^k))^2 + (\text{dist}(w_t^k, \tilde{w}_t^k))^2 - 2\phi\langle R_{w_t^k}^{-1}(w_{t+1}^k), \text{Exp}_{w_t^k}^{-1}(\tilde{w}_t^k) \rangle_{w_t^k}] \\ (\text{A.30}) \quad &\leq \mathbb{E}\left[\zeta \frac{\| -\alpha_t^k \mathcal{H}_t^k \xi_t^k \|_{w_t^k}^2}{\tau_1^2}\right] + (\text{dist}(w_t^k, \tilde{w}_t^k))^2 - 2\phi\langle -\alpha_t^k \mathcal{H}_t^k \xi_t^k, \text{Exp}_{w_t^k}^{-1}(\tilde{w}_t^k) \rangle_{w_t^k} \\ &\leq \zeta(\alpha_t^k)^2 \frac{\Gamma_{nc}^2}{\tau_1^2} \mathbb{E}[\mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2]] + \mathbb{E}[(\text{dist}(w_t^k, \tilde{w}_t^k))^2] \\ &\quad + \phi\alpha_t^k \mathbb{E}\left[\frac{1}{\nu_t} \Gamma_{nc}^2 \|\text{grad}f(w_t^k)\|_{w_t^k}^2 + \nu_t \|\text{Exp}_{w_t^k}^{-1}(\tilde{w}_t^k)\|_{w_t^k}^2\right] \\ &= \zeta(\alpha_t^k)^2 \frac{\Gamma_{nc}^2}{\tau_1^2} \mathbb{E}[\mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2]] + \mathbb{E}[(\text{dist}(w_t^k, \tilde{w}_t^k))^2] \\ &\quad + \phi\alpha_t^k \mathbb{E}\left[\frac{1}{\nu_t} \Gamma_{nc}^2 \|\text{grad}f(w_t^k)\|_{w_t^k}^2\right] + \phi\alpha_t^k \mathbb{E}\left[\nu_t \|\text{Exp}_{w_t^k}^{-1}(\tilde{w}_t^k)\|_{w_t^k}^2\right] \\ &= \zeta(\alpha_t^k)^2 \frac{\Gamma_{nc}^2}{\tau_1^2} \mathbb{E}[\mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2]] + \frac{\phi\alpha_t^k \Gamma_{nc}^2}{\nu_t} \mathbb{E}[\|\text{grad}f(w_t^k)\|_{w_t^k}^2] + (1 + \phi\alpha_t^k \nu_t) \mathbb{E}[(\text{dist}(w_t^k, \tilde{w}_t^k))^2], \end{aligned} \quad (\text{A.40})$$

where the second inequality uses (A.30) in Lemma C.15 and $\mathbb{E}_{i_t^k}[-\mathcal{H}_t^k \xi_t^k] = -\mathcal{H}_t^k \text{grad}f(w_t^k)$.

The third inequality uses the relation $2\langle a, b \rangle \leq \frac{1}{\nu_t} \|a\|^2 + \nu_t \|b\|^2$.

Now, we introduce the following function defined as

$$V_t^k := \mathbb{E}[f(w_t^k) + c_t^k(\text{dist}(\tilde{w}_t^k, w_t^k))^2]. \quad (\text{A.41})$$

This function measures how far the current parameter w_t^k is from \tilde{w}_t^k and the objective function

value. Then, V_{t+1}^k is calculated from Lemma C.16 as

$$\begin{aligned}
& V_{t+1}^k \\
&= \mathbb{E}[f(w_{t+1}^k) + c_{t+1}^k (\text{dist}(\tilde{w}^k, w_{t+1}^k))^2] \\
&\stackrel{(A.39), (A.40)}{\leq} \mathbb{E}[f(w_t^k) - \alpha_t^k \gamma_{nc} \|\text{grad} f(w_t^k)\|_{w_t^k}^2 + \frac{(\alpha_t^k)^2 \Lambda \Gamma_{nc}^2}{2} \mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2] \\
&\quad + c_{t+1}^k \left[\zeta (\alpha_t^k)^2 \frac{\Gamma_{nc}^2}{\tau_1^2} \mathbb{E}[\mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2]] + \frac{\phi \alpha_t^k \Gamma_{nc}^2}{\nu_t} \mathbb{E}[\|\text{grad} f(w_t^k)\|_{w_t^k}^2] \right. \\
&\quad \left. + (1 + \phi \alpha_t^k \nu_t) \mathbb{E}[(\text{dist}(w_t^k, \tilde{w}^k))^2] \right] \\
&= \mathbb{E} \left[f(w_t^k) - \alpha_t^k \left(\gamma_{nc} - \frac{\phi c_{t+1}^k \Gamma_{nc}^2}{\nu_t} \right) \|\text{grad} f(w_t^k)\|_{w_t^k}^2 \right] + c_{t+1}^k (1 + \phi \alpha_t^k \nu_t) \mathbb{E}[(\text{dist}(w_t^k, \tilde{w}^k))^2] \\
&\quad + \left(\frac{(\alpha_t^k)^2 \Lambda \Gamma_{nc}^2}{2} + c_{t+1}^k \zeta (\alpha_t^k)^2 \frac{\Gamma_{nc}^2}{\tau_1^2} \right) \mathbb{E}[\mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2]] \\
&\stackrel{(A.32)}{\leq} \mathbb{E} \left[f(w_t^k) - \alpha_t^k \left(\gamma_{nc} - \frac{\phi c_{t+1}^k \Gamma_{nc}^2}{\nu_t} \right) \|\text{grad} f(w_t^k)\|_{w_t^k}^2 \right] + c_{t+1}^k (1 + \phi \alpha_t^k \nu_t) \mathbb{E}[(\text{dist}(w_t^k, \tilde{w}^k))^2] \\
&\quad + \left(\frac{(\alpha_t^k)^2 \Lambda \Gamma_{nc}^2}{2} + c_{t+1}^k \zeta (\alpha_t^k)^2 \frac{\Gamma_{nc}^2}{\tau_1^2} \right) \\
&\quad \times \left(4(\beta^2 + \tau_2^2 C^2 \theta^2) \mathbb{E}[(\text{dist}(w_t^k, \tilde{w}^k))^2] + 2 \mathbb{E}[\|\text{grad} f(w_t^k)\|_{w_t^k}^2] \right) \\
&= \mathbb{E} \left[f(w_t^k) - \alpha_t^k \left(\gamma_{nc} - \frac{\phi c_{t+1}^k \Gamma_{nc}^2}{\nu_t} - \alpha_t^k \Lambda \Gamma_{nc}^2 - 2c_{t+1}^k \zeta \alpha_t^k \frac{\Gamma_{nc}^2}{\tau_1^2} \right) \|\text{grad} f(w_t^k)\|_{w_t^k}^2 \right] \\
&\quad + (c_{t+1}^k (1 + \phi \alpha_t^k \nu_t + 4\zeta (\alpha_t^k)^2 \frac{\Gamma_{nc}^2}{\tau_1^2} (\beta^2 + \tau_2^2 C^2 \theta^2)) + 2(\alpha_t^k)^2 \Lambda \Gamma_{nc}^2 (\beta^2 + \tau_2^2 C^2 \theta^2)) \\
&\quad \times \mathbb{E}[(\text{dist}(w_t^k, \tilde{w}^k))^2] \\
&= V_t^k - \alpha_t^k \left(\gamma_{nc} - \frac{\phi c_{t+1}^k \Gamma_{nc}^2}{\nu} - \alpha_t^k \Lambda \Gamma_{nc}^2 - 2c_{t+1}^k \zeta \alpha_t^k \frac{\Gamma_{nc}^2}{\tau_1^2} \right) \mathbb{E}[\|\text{grad} f(w_t^k)\|_{w_t^k}^2] \\
&= V_t^k - \Delta_t \mathbb{E}[\|\text{grad} f(w_t^k)\|_{w_t^k}^2]. \tag{A.42}
\end{aligned}$$

Rearranging the above yields the claim. This completes the proof. \square

We obtain the counterpart result of Proposition C.18 for the parallel translation and the exponential mapping.

Corollary C.19. *Let \mathcal{M} be a Riemannian manifold and $w^* \in \mathcal{M}$ be a non-degenerate local minimizer of f . Consider Algorithm 1 with option II and with $\mathcal{T} = P$ and $R = \text{Exp}$, i.e., the parallel translation and the exponential mapping case. Suppose Assumption 1 holds. Let the constants θ be in (5), τ_2 in (6), and β_0 , and C in (7). Λ is the constant in Assumption 1.3, and γ_{nc} and Γ_{nc} are the constants γ and Γ in (8). For $c_t^k, c_{t+1}^k, \nu_t > 0$, we set*

$$c_t^k = c_{t+1}^k (1 + \nu_t \alpha_t^k + 2\zeta (\alpha_t^k)^2 \Gamma_{nc}^2 \beta_0^2) + 2(\alpha_t^k)^2 \Lambda \Gamma_{nc}^2 \beta_0^2. \tag{A.43}$$

We also define

$$\Delta_t := \alpha_t^k \left(\gamma_{nc} - \frac{c_{t+1}^k \Gamma_{nc}^2}{\nu_t} - \alpha_t^k \Lambda \Gamma_{nc}^2 - 2c_{t+1}^k \zeta \alpha_t^k \Gamma_{nc}^2 \right). \tag{A.44}$$

Let α_t^k, ν_t and c_{t+1}^k be defined such that $\Delta_t > 0$. It then follows that, for any sequence $\{\tilde{w}_t^k\}$ generated by Algorithm 1 with a fixed step-size $\alpha_t^k := \alpha$ and $m_k := m$ converging to w^* , the expected squared Riemannian gradient satisfies the following bound as

$$\mathbb{E}[\|\text{grad}f(w_t^k)\|^2] \leq \frac{V_t^k - V_{t+1}^k}{\Delta_t}, \quad (\text{A.45})$$

where $V_t^k := \mathbb{E}[f(w_t^k) + c_t^k(\text{dist}(\tilde{w}_t^k, w_t^k))^2]$ for $0 \leq k \leq K - 1$.

The following proposition is very similar to Theorem 2 in [12].

Proposition C.20 (Theorem 2 in [12]). *Let \mathcal{M} be a Riemannian manifold and $w^* \in \mathcal{M}$ be a non-degenerate local minimizer of f . Consider Algorithm 1 with option II and IV, and suppose Assumption 1 holds. Let the constants θ be in (5), τ_1 and τ_2 in (6), and β , and C in (7). Λ is the constant in Assumption 1.3, and γ_{nc} and Γ_{nc} are the constants γ and Γ in (8). Let $c_m = 0$, $\alpha_t^k = \alpha > 0$, $\nu_t = \nu > 0$, and c_t^k is defined as (A.36) such that Δ_t defined in (A.37) satisfies $\Delta_t > 0$ for $0 \leq t \leq m - 1$. Define $\delta_t := \min_t \Delta_t$. Let T be mK . It then follows that, for the output w_{sol} of Algorithm 1, we have*

$$\mathbb{E}[\|\text{grad}f(w_{\text{sol}})\|^2] \leq \frac{f(w^0) - f(w^*)}{T\delta_t}. \quad (\text{A.46})$$

Proof. Because the proof is identical to those in [12, 15, 23], we omit it. The complete proof is therein. The sketch of the proof is as follows; we first telescoping the sum of (A.45) from $t = 0$ to $t = m - 1$ by introducing δ_t , then estimate its upper bound from the difference between V_0^s and V_0^m defined in (A.41). After showing that this difference is equivalent to the expected difference between $f(\tilde{w}^k)$ and $f(\tilde{w}^{k+1})$, summing up from $k = 0$ to $k = K - 1$, we obtain the desired claim. \square

C.4.3 Main proof of Theorem 3.2

We finally present the proof of Theorem 3.2 based on the extensions of results in [12, 15, 23].

Theorem. 3.2. *Let \mathcal{M} be a Riemannian manifold and $w^* \in \mathcal{M}$ be a non-degenerate local minimizer of f . Consider Algorithm 1 with option II and IV, and suppose Assumption 1 holds. Let the constants θ in (5), τ_1 and τ_2 in (6), and β , and C in (7). Λ is the constant in Assumption 1.3, and γ_{nc} and Γ_{nc} are the constants γ and Γ in (8). Set $\nu = \frac{\sqrt{\beta^2 + \tau_2^2 C^2 \theta^2} \Gamma_{nc}}{N^{a_1/2} \tau_1} \zeta^{1-a_2}$ and $\alpha_t^k = \alpha = \frac{\mu_0 \tau_1}{\sqrt{\beta^2 + \tau_2^2 C^2 \theta^2} N^{a_1} \Gamma_{nc} \zeta^{a_2}}$, where $0 < a_1 < 1$, and $0 < a_2 < 2$. Given sufficiently small $\mu_0 \in (0, 1)$, suppose that $\varrho > 0$ is chosen such that*

$$\frac{\sqrt{\beta^2 + \tau_2^2 C^2 \theta^2}}{\Lambda \Gamma_{nc}} \gamma_{nc} \left(1 - \frac{\varrho \Gamma_{nc}}{\mu_0 \gamma_{nc}}\right) > \frac{2\phi \mu_0 (e-1) \tau_1}{\zeta^{2-a_2}} + \frac{\mu_0 \tau_1}{N^{a_1} \zeta^{a_2}} + \frac{4\mu_0^2 (e-1)}{N^{\frac{3a_1}{2}} \zeta^{a_2} \tau_1} \quad (\text{A.47})$$

holds. Set $m = \lfloor \frac{N^{3a_1/2}}{5\phi_1 \mu_0 \zeta^{1-2a_2}} \rfloor$ and $T = mK$. Then, we have

$$\mathbb{E}[\|\text{grad}f(w_a)\|^2] \leq \frac{\sqrt{\beta^2 + \tau_2^2 C^2 \theta^2} N^{a_1} \zeta^{a_2} [f(w^0) - f(w^*)]}{T\varrho}. \quad (\text{A.48})$$

Proof. From (A.37) in Proposition C.18, we need to consider the upper bound of c_t^k defined in (A.36). To this end, the upper bound of c_0^k is first derived. Denoting, for simplicity, $\varphi = \phi\alpha\nu + 4\zeta\alpha^2\frac{\Gamma_{nc}^2}{\tau_1^2}(\beta^2 + \tau_2^2 C^2\theta^2)$ and $\omega = \tau_2 C\theta$, we first consider the bound of φ as

$$\begin{aligned}\varphi &= \phi\alpha\nu + 4\zeta\alpha^2\frac{\Gamma_{nc}^2}{\tau_1^2}(\beta^2 + \omega^2) \\ &= \frac{\phi\mu_0\zeta^{1-2a_2}}{N^{\frac{3a_1}{2}}} + \frac{4\mu_0^2\zeta^{1-2a_2}}{N^{2a_1}} \\ &= \frac{\phi\mu_0\zeta^{1-2a_2}}{N^{\frac{3a_1}{2}}} \left(1 + \frac{4\mu_0}{\phi N^{\frac{1}{2a_1}}}\right).\end{aligned}$$

Consequently, we obtain the bound of φ as $\varphi \in \left(\frac{\phi\mu_0\zeta^{1-2a_2}}{N^{3a_1/2}}, 5\frac{\phi\mu_0\zeta^{1-2a_2}}{N^{3a_1/2}}\right)$.

Then, we consider the recurrence relation $c_t^k = c_{t+1}^2(1 + \varphi) + 2\alpha^2\Lambda\frac{\Gamma_{nc}^2}{\tau_1^2}(\beta^2 + \omega^2)$ as

$$\begin{aligned}c_0^k &= \left(c_m^k - \frac{2\alpha^2\Lambda\Gamma_{nc}^2(\beta^2 + \omega^2)}{-\varphi\tau_1^2}\right)(1 + \varphi)^m + \frac{2\alpha^2\Lambda\Gamma_{nc}^2(\beta^2 + \omega^2)}{-\varphi\tau_1^2} \\ &= 2\alpha^2\Lambda\frac{\Gamma_{nc}^2}{\tau_1^2}(\beta^2 + \omega^2)\frac{(1 + \varphi)^m - 1}{\varphi} \\ &= 2\frac{\mu_0^2\Lambda}{N^{2a_1}\zeta^{2a_2}}\frac{(1 + \varphi)^m - 1}{\varphi} \\ &\leq 2\frac{\mu_0\Lambda}{N^{\frac{a_1}{2}}\zeta}((1 + \varphi)^m - 1) \\ &\leq 2\frac{\mu_0\Lambda}{N^{\frac{a_1}{2}}\zeta}(e - 1),\end{aligned}\tag{A.49}$$

where the second equality uses $c_m^k = 0$, and the first inequality uses the lower bound of φ derived above. Regarding the last inequality, because $m = \lfloor \frac{N^{3a_1/2}}{5\phi\mu_0\zeta^{1-2a_2}} \rfloor$, $\varphi \leq 1/m$. In addition, noting that $\lim_{r \rightarrow \infty} (1 + 1/r)^r = e$ for $r > 0$ where e is the Euler's number, we used the relation $(1 + \varphi)^m < e$.

Now, we attempt to estimate the lower bound of δ_t , i.e., $\min_t \Delta_t$.

$$\begin{aligned}\delta_t &= \min_t \Delta_t \\ &= \min_t \alpha \left(\gamma_{nc} - \frac{\phi c_{t+1}^k \Gamma_{nc}^2}{\nu} - \alpha \Lambda \Gamma_{nc}^2 - 2c_{t+1}^k \zeta \alpha \frac{\Gamma_{nc}^2}{\tau_1^2} \right) \\ &\geq \alpha \left(\gamma_{nc} - \frac{\phi c_0^k \Gamma_{nc}^2}{\nu} - \alpha \Lambda \Gamma_{nc}^2 - 2c_0^k \zeta \alpha \frac{\Gamma_{nc}^2}{\tau_1^2} \right) \\ &\stackrel{(A.49)}{\geq} \alpha \left(\gamma_{nc} - \frac{2\phi\mu_0(e-1)}{\zeta^{2-a_2}} \frac{\Lambda}{\sqrt{\beta^2 + \omega^2}} \Gamma_{nc} \tau_1 - \frac{\mu_0}{N^{a_1}\zeta^{a_2}} \frac{\Lambda}{\sqrt{\beta^2 + \omega^2}} \Gamma_{nc} \tau_1 \right. \\ &\quad \left. - \frac{4\mu_0^2(e-1)}{N^{\frac{3a_1}{2}}\zeta^{a_2}} \frac{\Lambda}{\sqrt{\beta^2 + \omega^2}} \frac{\Gamma_{nc}}{\tau_1} \right) \\ &\stackrel{(A.47)}{\geq} \frac{\varrho}{\sqrt{\beta^2 + \omega^2} N^{a_1} \zeta^{a_2}},\end{aligned}\tag{A.50}$$

where the second inequality uses (A.49) for the second and the fourth terms. Substituting (A.50) into (A.46) in Proposition C.20 completes the proof. \square

Corollary C.21. *Suppose the same assumptions and conditions as those of Theorem 3.2. Then, the total number of gradient evaluations in Algorithm 1 is $\mathcal{O}(N^{a_1}/\epsilon)$ to obtain an ϵ -solution.*

Proof. The total number of gradient evaluations is equals to $(N + m)K$. Comparing the left term in (A.48) with ϵ , we obtain $K \approx \mathcal{O}(N^{a_1}/(m\epsilon))$. Additionally, $m = \lfloor \frac{N^{3a_1/2}}{5\mu_0\zeta^{1-2a_2}} \rfloor \approx \mathcal{O}(N^{3a_1/2})$. Consequently, it results in that the total number of gradient evaluations is $\mathcal{O}(N^{a_1}/\epsilon)$. \square

The obtained complexity is the same as that of R-SVRG [23] in terms of the total number of samples, N .

We obtain the corresponding result of Theorem 3.2 when the parallel translation and the exponential mapping are used.

Corollary C.22. *Let \mathcal{M} be a Riemannian manifold and $w^* \in \mathcal{M}$ be a non-degenerate local minimizer of f . Consider Algorithm 1 with option II and IV and with $\mathcal{T} = P$ and $R = \text{Exp}$, i.e., the parallel translation and the exponential mapping case. Let the constant β_0 be β in (7). Λ is the constant in Assumption 1.3, and γ_{nc} and Γ_{nc} are the constants γ and Γ in (8). Set $\nu = \frac{\beta_0\Gamma_{nc}\zeta^{1-2a_2}}{N^{a_1/2}}$ and $\alpha_t^k = \alpha = \frac{\mu_0}{\beta_0 N^{a_1}\Gamma_{nc}\zeta^{a_2}}$, where $0 < a_1 < 1$, and $0 < a_2 < 2$. Given sufficiently small $\mu_0 \in (0, 1)$, suppose that $\varrho > 0$ is chosen such that*

$$\frac{\beta_0}{\Lambda\Gamma_{nc}}\gamma_{nc}\left(1 - \frac{\varrho\Gamma_{nc}}{\mu_0\gamma_{nc}}\right) > \frac{\mu_0(e-1)}{\zeta^{2-a_2}} + \frac{\mu_0}{N^{a_1}\zeta^{a_2}} + \frac{2\mu_0^2(e-1)}{N^{3a_1/2}\zeta^{a_2}}$$

holds. Set $m = \lfloor \frac{N^{3a_1/2}}{3\mu_0\zeta^{1-2a_2}} \rfloor$, and $T = mK$. Then, we have

$$\mathbb{E}[\|\text{grad}f(w_a)\|^2] \leq \frac{\beta_0 N^{a_1} \zeta^{a_2} [f(w^0) - f(w^*)]}{T\varrho}.$$

Proof. The proof is similar to that of Theorem 3.2. Therefore, we omit it. However, it should be noted that we follow Corollary C.17 to bound $\mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2]$ instead of using Lemma C.16. \square

D Proof of convergence analysis on retraction-convex functions

This subsection presents a local convergence rate analysis in neighborhood of a local minimum for retraction-convex functions. This *local* setting is very common and standard in manifold optimization.

D.1 Preliminary lemmas

This subsection first states some essential lemmas. Since f is strongly retraction-convex on Θ by Assumption 3, there exist constants $0 < \lambda$ such that $\lambda \leq \frac{d^2 f(R_{w_t^k}(\tau \eta_k / \|\eta_k\|_{w_t^k}))}{d\tau^2}$ for all $t \in [0, \alpha_t^k \|\eta_k\|_{w_t^k}]$. From Taylor's theorem, we obtain below;

Lemma D.1 (In Lemma 3.2 in [21]). *Under Assumptions 1.1, 1.2, and Assumption 3, there exists λ such that*

$$f(w_{t+1}^k) - f(w_t^k) \geq \langle \text{grad} f(w_t^k), \alpha_t^k \eta_k \rangle_{w_t^k} + \frac{1}{2} \lambda (\alpha_t^k \|\eta_k\|_{w_t^k})^2. \quad (\text{A.51})$$

Proof. From Taylor's theorem, we have

$$\begin{aligned} & f(w_{t+1}^k) - f(w_t^k) \\ &= f(R_{w_t^k}(\alpha_t^k \eta_k)) - f(R_{w_t^k}(0)) \\ &= \frac{d}{d\tau} f(R_{w_t^k}(\tau \eta_k / \|\eta_k\|_{w_t^k})) \Big|_{\tau=0} \cdot \alpha_t^k \|\eta_k\|_{w_t^k} + \frac{1}{2} \frac{d^2}{d\tau^2} f(R_{w_t^k}(\tau \eta_k / \|\eta_k\|_{w_t^k})) \Big|_{\tau=p} \cdot (\alpha_t^k \|\eta_k\|_{w_t^k})^2 \\ &= \langle \text{grad} f(w_t^k), \alpha_t^k \eta_k \rangle_{w_t^k} + \frac{1}{2} \frac{d^2}{d\tau^2} f(R_{w_t^k}(\tau \eta_k / \|\eta_k\|_{w_t^k})) \Big|_{\tau=p} \cdot (\alpha_t^k \|\eta_k\|_{w_t^k})^2 \\ &\geq \langle \text{grad} f(w_t^k), \alpha_t^k \eta_k \rangle_{w_t^k} + \frac{1}{2} \lambda (\alpha_t^k \|\eta_k\|_{w_t^k})^2, \end{aligned}$$

where $0 \leq p \leq \alpha_t^k \|\eta_k\|_{w_t^k}$, and the inequality uses Assumption 3. This yields (A.51). This completes the proof. \square

Lemma D.2 (Lemma 3.3 in [21]). *Under Assumptions 1.1, 1.2, 1.3, 1.4, 1.6, and Assumption 3, there exist two constants $0 < \lambda < \Lambda$ such that*

$$\lambda \leq \frac{\langle s_k, y_k \rangle}{\langle s_k, s_k \rangle} \leq \Lambda \quad (\text{A.52})$$

for all k .

Proof. From Lemma D.1, the proof of this lemma is given, but we omit it. The reader can see the complete proof in Lemma 3.3 in [21]. \square

Lemma D.3 (Lemma 3.9 in [21]). *Suppose Assumption 1 and Assumption 3 hold, there exists a constant $0 < \Upsilon_c$ for all k such that*

$$\frac{\langle y_k, y_k \rangle}{\langle s_k, y_k \rangle} \leq \Upsilon_c. \quad (\text{A.53})$$

Proof. (Lemma 3.9 in [21]) The complete proof is given in [21]. It is also stated in Lemma C.3 for completeness. (A.4) yields $\|y_k\| \leq b_4\|s_k\|$ derived in Lemma C.3, where $b_4 > 0$. Therefore, by Lemma D.2, we have

$$\frac{\langle y_k, y_k \rangle}{\langle s_k, y_k \rangle} \leq \frac{\langle y_k, y_k \rangle}{\lambda \langle s_k, s_k \rangle} \leq \frac{b_4^2}{\lambda} \quad (= \Upsilon_c). \quad (\text{A.54})$$

Denoting b_4^2/λ as Υ_c , this completes the proof. \square

D.2 Eigenvalue bounds of \mathcal{H}_t^k on retraction-convex functions

Now, we attempt to bound $\text{trace}(\hat{\tilde{\mathcal{B}}})$ and $\det(\hat{\tilde{\mathcal{B}}})$ in order to bound the eigenvalues of $\tilde{\mathcal{H}}^k$, where a *hat* denotes the coordinate expression of the operator. The bound of $\text{trace}(\hat{\tilde{\mathcal{B}}})$ is identical to that of the non-convex case in Lemma C.5. Therefore, we concentrate on the bound of $\det(\hat{\tilde{\mathcal{B}}})$. As the same as Lemma C.5, the proof follows stochastic L-BFGS methods in the Euclidean space, e.g., [16, 24]. Similarly to Section C.2, it should be noted that $\text{trace}(\hat{\tilde{\mathcal{B}}})$ and $\det(\hat{\tilde{\mathcal{B}}})$ do not depend on the chosen basis.

Lemma D.4 (Bounds of trace and determinant of $\tilde{\mathcal{B}}^k$). *Consider the recursion of $\tilde{\mathcal{B}}_u^k$ defined in (A.7). If Assumption 1 and Assumption 3 hold, the $\text{trace}(\hat{\tilde{\mathcal{B}}^k})$ in a coordinate expression of $\tilde{\mathcal{B}}^k$ is uniformly upper bounded for all $k \geq 1$,*

$$\text{trace}(\hat{\tilde{\mathcal{B}}^k}) \leq (M + \tau)\Upsilon_c. \quad (\text{A.55})$$

where M is the dimension of \mathcal{M} . Similarly, if Assumption 1 and Assumption 3 hold, the $\det(\hat{\tilde{\mathcal{B}}^k})$ in a coordinate expression of $\tilde{\mathcal{B}}^k$ is uniformly lower bounded for all k ,

$$\det(\hat{\tilde{\mathcal{B}}^k}) \geq v^M \left[\frac{\lambda}{(M + \tau)\Upsilon_c} \right]^\tau. \quad (\text{A.56})$$

Here, a *hat* expression represents the coordinate expression of an operator.

Proof. The proof can be completed parallel to the Euclidean case [17]. As mentioned, the proof for the bound of $\text{trace}(\hat{\tilde{\mathcal{B}}})$ is given in Lemma C.5, we address only $\det(\hat{\tilde{\mathcal{B}}^k})$.

Because \mathcal{T} is an isometry vector transport, \mathcal{T}_{η_k} is invertible for all k . Accordingly, $\det(\hat{\tilde{\mathcal{B}}^k})$ can be reformulated as

$$\det(\hat{\tilde{\mathcal{B}}^k}) = \det(\hat{\mathcal{T}_{\eta_k} \tilde{\mathcal{B}}^k \mathcal{T}_{\eta_k}^{-1}}) = \det(\hat{\tilde{\mathcal{B}}^k}). \quad (\text{A.57})$$

We consider the determinant lower bound of $\det(\hat{\mathcal{B}}_{k,\tau})$ from (A.7) as

$$\begin{aligned}
\det(\hat{\mathcal{B}}_{u+1}^k) &= \det(\hat{\mathcal{B}}_u^k) \det \left(\mathbf{I} - \frac{s_{k-\tau+u}(\hat{\mathcal{B}}_u^k s_{k-\tau+u})^T}{\langle \hat{\mathcal{B}}_u^k s_{k-\tau+u}, s_{k-\tau+u} \rangle} + \frac{(\hat{\mathcal{B}}_u^k)^{-1} y_{k-\tau+u} y_{k-\tau+u}^T}{\langle y_{k-\tau+u}, s_{k-\tau+u} \rangle} \right) \\
&= \det(\hat{\mathcal{B}}_u^k) \det \left(\frac{(\hat{\mathcal{B}}_u^k s_{k-\tau+u})^T}{\langle \hat{\mathcal{B}}_u^k s_{k-\tau+u}, s_{k-\tau+u} \rangle} (\hat{\mathcal{B}}_u^k)^{-1} y_{k-\tau+u} \right) \\
&= \det(\hat{\mathcal{B}}_u^k) \frac{\langle s_{k-\tau+u}, y_{k-\tau+u} \rangle}{\langle \hat{\mathcal{B}}_u^k s_{k-\tau+u}, s_{k-\tau+u} \rangle} \\
&= \det(\hat{\mathcal{B}}_u^k) \frac{\langle s_{k-\tau+u}, y_{k-\tau+u} \rangle}{\|s_{k-\tau+u}\|^2} \frac{\|s_{k-\tau+u}\|^2}{\langle \hat{\mathcal{B}}_u^k s_{k-\tau+u}, s_{k-\tau+u} \rangle} \\
&\geq \det(\hat{\mathcal{B}}_u^k) \frac{\lambda}{\lambda_{\max}(\hat{\mathcal{B}}_u^k)} \\
&\geq \det(\hat{\mathcal{B}}_u^k) \frac{\lambda}{\text{trace}(\hat{\mathcal{B}}_u^k)} \\
&\geq \det(\hat{\mathcal{B}}_u^k) \frac{\lambda}{(M + \tau) \Upsilon_c}. \tag{A.58}
\end{aligned}$$

Regarding the second equality, we obtain it from the formula $\det(\mathbf{I} + \mathbf{u}_1 \mathbf{v}_1^T + \mathbf{u}_2 \mathbf{v}_2^T) = (1 + \mathbf{u}_1^T \mathbf{v}_1)(1 + \mathbf{u}_2^T \mathbf{v}_2) - (\mathbf{u}_1^T \mathbf{v}_2)(\mathbf{u}_2^T \mathbf{v}_1)$ by setting $\mathbf{u}_1 = -s_{k-\tau+u}$, $\mathbf{v}_1 = \hat{\mathcal{B}}_u^k s_{k-\tau+u} / \langle \hat{\mathcal{B}}_u^k s_{k-\tau+u}, s_{k-\tau+u} \rangle$, $\mathbf{u}_2 = (\hat{\mathcal{B}}_u^k)^{-1} y_{k-\tau+u}$, and $\mathbf{v}_2 = y_{k-\tau+u} / \langle s_{k-\tau+u}, y_{k-\tau+u} \rangle$. The first inequality follows from (A.52) in Lemma D.2 and the fact $\langle \hat{\mathcal{B}}_u^k s_{k-\tau+u}, s_{k-\tau+u} \rangle \leq \lambda_{\max}(\hat{\mathcal{B}}_u^k) \|s_{k-\tau+u}\|^2$. Actually, we use the fact the trace of a positive definite matrix bounds its maximal eigenvalue for the second inequality. The last inequality follows (A.11). Then, applying (A.57), (A.58) turns to be

$$\det(\hat{\mathcal{B}}_{u+1}^k) \geq \det(\hat{\mathcal{B}}_u^k) \frac{\lambda}{(M + \tau) \Upsilon_c}. \tag{A.59}$$

Applying (A.59) recursively from $u = 0$ to $u = \tau - 1$, we obtain that

$$\det(\hat{\mathcal{B}}_\tau^k) \geq \left[\frac{\lambda}{(M + \tau) \Upsilon_c} \right]^\tau \det(\hat{\mathcal{B}}_0^k).$$

To bound the determinant of $\hat{\mathcal{B}}_0^k$, considering $\hat{\mathcal{B}}_0^k = \text{id} / \chi_k$ as above and Lemma C.2, we can rewrite for $k \geq 1$ as

$$\det(\hat{\mathcal{B}}_0^k) = \det \left(\frac{\mathbf{I}}{\chi_k} \right) = \frac{1}{\chi_k^M} = \left(\frac{\langle y_k, y_k \rangle}{\langle s_k, y_k \rangle} \right)^M \geq v^M,$$

where v is defined in Lemma C.2. Consequently, we obtain as

$$\det(\hat{\mathcal{B}}_\tau^k) \geq v^M \left[\frac{\lambda}{(M + \tau) \Upsilon_c} \right]^\tau.$$

Thus, this yields (A.56), and these complete the proof. \square

Now we prove the main lemma for Proposition D.6.

Lemma D.5. *If Assumption 1 and Assumption 3 hold, the eigenvalues of $\tilde{\mathcal{H}}^k$ is bounded by γ_c and Γ_c with $0 < \gamma_c < \Gamma_c < \infty$ for all $k \geq 1$ as*

$$\gamma_c \text{id} \preceq \tilde{\mathcal{H}}^k \preceq \Gamma_c \text{id}.$$

Proof. The proof is obtained as parallel to the Euclidean case [24]. The lower part is identical to the proof that is given in Lemma C.5. Regarding the upper bound, because the determinant of a matrix is the product of its eigenvalues, the lower bound in (A.56) bounds the product of the eigenvalues of $\hat{\mathcal{B}}^k$ from below. This means that $\prod_{i=1}^M \lambda_i \geq \frac{\lambda^\tau v^M}{[(M+\tau)\Upsilon_c]^\tau}$. Thus, we have below for any given eigenvalue of $\hat{\mathcal{B}}^k$, say λ_j ,

$$\lambda_j \geq \frac{1}{\prod_{k=1, k \neq j}^M \lambda_k} \cdot \frac{\lambda^\tau v^M}{[(M+\tau)\Upsilon_c]^\tau}. \quad (\text{A.60})$$

Considering that $(M+\tau)\Upsilon_c$ is an upper bound for the eigenvalues of $\hat{\mathcal{B}}^k$, $[(M+\tau)\Upsilon_c]^{M-1}$ gives the upper bound of the product of the $(M-1)$ eigenvalues $\prod_{k=1, k \neq j}^M \lambda_k$.

As a result, we obtain that any eigenvalues of $\hat{\mathcal{B}}^k$ is lower bounded as

$$\lambda_j \geq \frac{1}{[(M+\tau)\Upsilon_c]^{M-1}} \cdot \frac{\lambda^\tau v^M}{[(M+\tau)\Upsilon_c]^\tau} = \frac{\lambda^\tau v^M}{[(M+\tau)\Upsilon_c]^{M+\tau-1}}. \quad (\text{A.61})$$

Consequently, we finally obtain $\frac{\lambda^\tau v^M}{[(M+\tau)\Upsilon_c]^{M+\tau-1}} \text{id} \preceq \hat{\mathcal{B}}^k$.

Now, we obtain the claim. The bounds in (A.55) and (A.61) imply that their inverses are bounds for the eigenvalues of $\hat{\mathcal{H}}^k = (\hat{\mathcal{B}}^k)^{-1}$ as

$$(\gamma_c \text{id} =) \frac{1}{(M+\tau)\Upsilon_c} \text{id} \preceq \tilde{\mathcal{H}}^k \preceq \frac{[(M+\tau)\Upsilon_c]^{M+\tau-1}}{\lambda^\tau v^M} \text{id} \quad (= \Gamma_c \text{id}). \quad (\text{A.62})$$

Denoting $\frac{1}{(M+\tau)\Upsilon_c}$ as γ_c as in Lemma C.6, and $\frac{[(M+\tau)\Upsilon_c]^{M+\tau-1}}{\lambda^\tau v^M}$ as Γ_c , we obtain the claim. This completes the proof. \square

Finally we give Proposition D.6 for retraction-convex functions.

Proposition D.6 (Bounds of \mathcal{H}_t^k for retraction-convex functions). *Consider the operator $\tilde{\mathcal{H}}^k := \mathcal{T}_{\tilde{\eta}_t^k} \circ \tilde{\mathcal{H}}^k \circ (\mathcal{T}_{\tilde{\eta}_t^k})^{-1}$. Define the constant $0 < \gamma_c < \Gamma_c < \infty$. If Assumption 1 holds, the range of eigenvalues of \mathcal{H}_t^k is bounded by γ_c and Γ_c for all $k \geq 1, t \geq 1$, i.e.,*

$$\gamma_c \text{id} \preceq \mathcal{H}_t^k \preceq \Gamma_c \text{id}. \quad (\text{A.63})$$

Proof. The proof is identical to that of Proposition C.7. \square

Remark D.7. We discuss the obtained bounds of \mathcal{H}_t^k by comparing the retraction-convex case in Proposition D.6 with the non-convex case in Proposition C.7. The lower bound of \mathcal{H}_t^k in the convex case is $\gamma_c = 1/((M+\tau)\Upsilon_c) = \lambda/((M+\tau)b_4^2)$ from (A.53) and (A.62). The non-convex case is $\gamma_{nc} = 1/((M+\tau)\Upsilon_{nc}) = \epsilon/((M+\tau)b_4^2)$ from (A.6) and (A.13). In terms of ϵ and λ , γ_c is $\mathcal{O}(\lambda)$ and γ_{nc} is $\mathcal{O}(\epsilon)$. Assuming λ of the strongly retraction-convex functions

much larger than ϵ because it is generally set to very small value [27], we conclude $\gamma_c > \gamma_{nc}$. Meanwhile, the upper bound of \mathcal{H}_t^k in the convex case is $\Gamma_c = \frac{[(M+\tau)\Upsilon_c]^{M+\tau-1}}{\lambda^\tau v^M} = \frac{[(M+\tau)b_4^2]^{M+\tau-1}}{\lambda^{M+2\tau-1}\epsilon^M}$ from (A.53) and (A.62). The non-convex case is $\Gamma_{nc} = \frac{(1+b_4/\epsilon)^{2(\tau+1)-1}}{\epsilon((1+b_4/\epsilon)^2-1)}$ from (A.18). With respect to ϵ and λ , Γ_c is $\mathcal{O}(1/(\lambda^{M+2\tau-1}\epsilon^M))$ and Γ_{nc} is $\mathcal{O}(1/\epsilon^{2\tau})$. Similarly to the lower bound mentioned above, we conclude $\Gamma_c < \Gamma_{nc}$. Consequently, the range of the bounds of \mathcal{H}_t^k on strongly retraction-convex functions is smaller than that on non-convex functions.

D.3 Proof of local convergence rate analysis (Theorem 3.3)

This subsection first introduces some essential lemmas. Then, the main proof of Theorem 3.3 is given. This section also derives at the end a corollary about the analysis when the using exponential mapping and the parallel translation that are special cases of the retraction and the vector transport.

D.3.1 Essential lemmas

We first introduce a property of the Karcher mean on a general Riemannian manifold.

Lemma D.8 (Lemma C.2 in [22]). *Let w_1, \dots, w_m be points on a Riemannian manifold \mathcal{M} and let w be the Karcher mean of the m points. For an arbitrary point p on \mathcal{M} , we have*

$$(\text{dist}(p, w))^2 \leq \frac{4}{m} \sum_{i=1}^m (\text{dist}(p, w_i))^2.$$

We now bound the variance of ξ_t^k as follows.

Lemma D.9 (Lemma 5.8 in [22]). *Suppose Assumptions 1.1, 1.2, 1.4, 1.5, and 1.7, which guarantee Lemmas C.13, C.14, and C.15 for $\bar{w} = w^*$. Let $\beta > 0$ be a constant such that*

$$\|P_\gamma^{w \leftarrow z}(\text{grad} f_n(z)) - \text{grad} f_n(w)\|_w \leq \beta \text{dist}(z, w), \quad w, z \in \Theta, \quad n = 1, 2, \dots, N.$$

The existence of such β is guaranteed by Lemma C.13. The upper bound of the variance of ξ_t^k is given by

$$\mathbb{E}_{i_t^k} [\|\xi_t^k\|_{w_t^k}^2] \leq 4(\beta^2 + \tau_2^2 C^2 \theta^2)(7(\text{dist}(w_t^k, w^*))^2 + 4(\text{dist}(\tilde{w}^k, w^*))^2), \quad (\text{A.64})$$

where the constant θ corresponds to that in Lemma C.14, C is the constant of Assumption 1, and $\tau_2 > 0$ appears in (A.30).

We also have the following corollary of the previous lemma with the case $R = \text{Exp}$ and $\mathcal{T} = P$.

Corollary D.10 (Corollary 5.1 in [22]). *Consider Algorithm 1 with $\mathcal{T} = P$ and $R = \text{Exp}$, i.e., the parallel translation and the exponential mapping case. When each $\text{grad} f_n$ is β_0 -Lipschitz continuously differentiable, the upper bound of the variance of ξ_t^k is given by*

$$\mathbb{E}_{i_t^k} [\|\xi_t^k\|_{w_t^k}^2] \leq \beta_0^2(14(\text{dist}(w_t^k, w^*))^2 + 8\text{dist}(\tilde{w}^k, w^*))^2). \quad (\text{A.65})$$

Next, we show the lemma that finds a lower bound for $\|\text{grad} f(w_t^k)\|_{w_t^k}$ with respect to the error $f(w_t^k) - f(w^*)$, which is a standard derivation in the Euclidean space. See, e.g., [37]. We extend this into manifolds.

Lemma D.11. *Let $w \in \mathcal{M}$ and z be in a totally retractive neighborhood of w . It holds that*

$$2\lambda(f(w) - f(z)) \leq \|\text{grad}f(w)\|_w^2. \quad (\text{A.66})$$

Proof. Let $\zeta = R_w^{-1}(z)$. Using (A.51) in Lemma D.1, which is equivalent to the strong convexity of $\text{grad}f$, we obtain

$$\begin{aligned} f(z) &\geq f(w) + \langle \text{grad}f(w), \zeta \rangle_w + \frac{\lambda}{2} \|\zeta\|_w^2 \\ &\geq f(w) + \min_{\xi \in T_w \mathcal{M}} \left(\langle \text{grad}f(w), \xi \rangle_w + \frac{\lambda}{2} \|\xi\|_w^2 \right) \\ &\geq f(w) - \frac{1}{2\lambda} \|\text{grad}f(w)\|_w^2. \end{aligned}$$

Rearranging this inequality completes the proof. \square

D.3.2 Main proof of Theorem 3.3

Theorem 3.3. *Let \mathcal{M} be a Riemannian manifold and $w^* \in \mathcal{M}$ be a non-degenerate local minimizer of f (i.e., $\text{grad}f(w^*) = 0$ and the Hessian $\text{Hess}f(w^*)$ of f at w^* is positive definite). Suppose Assumptions 1 and 3 hold. Let the constants β, θ , and C be in Lemma D.9, and τ_1 and τ_2 be in Lemma C.15. Λ and λ are the constants in Lemmas C.1 and D.1, respectively. γ_c and Γ_c are the constants in Proposition D.6. Let α be a positive number satisfying $\lambda\tau_1^2 > 2\alpha(\lambda^2\tau_1^2 - 14\alpha\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2))$ and $\gamma_c\lambda^2\tau_1^2 > 14\alpha\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2)$. It then follows that for any sequence $\{\tilde{w}^k\}$ generated by Algorithm 1 with Option II under a fixed step-size $\alpha_t^k := \alpha$ and $m_k := m$ converging to w^* , there exists $0 < K_{th} < K$ such that for all $k > K_{th}$,*

$$\mathbb{E}[(\text{dist}(\tilde{w}^{k+1}, w^*))^2] \leq \frac{2(\Lambda\tau_2^2 + 16m\alpha^2\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2))}{m\alpha(\gamma_c\lambda^2\tau_1^2 - 14\alpha\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2))} \mathbb{E}[(\text{dist}(\tilde{w}^{k-1}, w^*))^2].$$

Proof. Using (A.1) in Lemma C.1, which is equivalent to the Lipschitz continuity of $\text{grad}f$ from Assumptions 1, we obtain

$$f(w_{t+1}^k) - f(w_t^k) \leq \langle \text{grad}f(w_t^k), -\alpha\mathcal{H}_t^k \xi_t^k \rangle_{w_t^k} + \frac{1}{2}\Lambda(-\alpha\|\mathcal{H}_t^k \xi_t^k\|_{w_t^k})^2.$$

Taking expectation with regard to i_t^k , this becomes

$$\begin{aligned} \mathbb{E}_{i_t^k}[f(w_{t+1}^k)] - f(w_t^k) &\leq \mathbb{E}_{i_t^k}[\langle \text{grad}f(w_t^k), -\alpha\mathcal{H}_t^k \xi_t^k \rangle_{w_t^k} + \frac{1}{2}\alpha^2\Lambda\|\mathcal{H}_t^k \xi_t^k\|_{w_t^k}^2] \\ &\leq -\alpha\langle \text{grad}f(w_t^k), \mathbb{E}_{i_t^k}[\mathcal{H}_t^k \xi_t^k] \rangle_{w_t^k} + \frac{1}{2}\alpha^2\Lambda\mathbb{E}_{i_t^k}[\|\mathcal{H}_t^k \xi_t^k\|_{w_t^k}^2] \\ &\leq -\alpha\langle \text{grad}f(w_t^k), \mathcal{H}_t^k \text{grad}f(w_t^k) \rangle_{w_t^k} + \frac{1}{2}\alpha^2\Lambda\mathbb{E}_{i_t^k}[\|\mathcal{H}_t^k \xi_t^k\|_{w_t^k}^2] \\ &\leq -\alpha\gamma_c\|\text{grad}f(w_t^k)\|_{w_t^k}^2 + \frac{1}{2}\alpha^2\Lambda\Gamma_c^2\mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2]. \end{aligned} \quad (\text{A.67})$$

where the third inequality used the fact that $\mathbb{E}_{i_t^k}[\mathcal{H}_t^k \xi_t^k] = \mathcal{H}_t^k \text{grad}f(w_t^k)$. The last inequality used the bound of \mathcal{H}_t^k in Proposition D.6.

From Lemma D.11, (A.67) yields

$$\mathbb{E}_{i_t^k}[f(w_{t+1}^k)] - f(w_t^k) \leq -2\alpha\gamma_c\lambda(f(w_t^k) - f(w^*)) + \frac{1}{2}\alpha^2\Lambda\Gamma_c^2\mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2]. \quad (\text{A.68})$$

Using (A.51) in Lemma D.1 with $\text{grad}f(w^*) = 0$, and using Lemma C.15, we obtain

$$f(w_t^k) - f(w^*) \geq \frac{\lambda}{2}\|R_{w^*}^{-1}(w_t^k)\|_{w^*}^2 \geq \frac{\lambda\tau_1^2}{2}(\text{dist}(w_t^k, w^*))^2. \quad (\text{A.69})$$

Plugging (A.69) and the bound of $\mathbb{E}_{i_t^k}[\|\xi_k^k\|^2]$ in (A.64) in Lemma D.9 into (A.68) yields

$$\begin{aligned} \mathbb{E}_{i_t^k}[f(w_{t+1}^k)] - f(w_t^k) &\leq -\alpha\gamma_c\lambda^2\tau_1^2(\text{dist}(w_t^k, w^*))^2 + \frac{1}{2}\alpha^2\Lambda\Gamma_c^2\mathbb{E}_{i_t^k}[\|\xi_t^k\|_{w_t^k}^2] \\ &\leq -\alpha\gamma_c\lambda^2\tau_1^2(\text{dist}(w_t^k, w^*))^2 \\ &\quad + \frac{1}{2}\alpha^2\Lambda\Gamma_c^2\{4(\beta^2 + \tau_2^2C^2\theta^2)(7(\text{dist}(w_t^k, w^*))^2 + 4(\text{dist}(\tilde{w}^k, w^*))^2)\} \\ &\leq (-\alpha\gamma_c\lambda^2\tau_1^2 + 14\alpha^2\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2))(\text{dist}(w_t^k, w^*))^2 \\ &\quad + 8\alpha^2\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2)(\text{dist}(\tilde{w}^k, w^*))^2. \end{aligned}$$

Taking expectations over all random variables, we obtain below by further summing over $t = 0, \dots, m-1$ of the inner loop on k -th epoch

$$\begin{aligned} \mathbb{E}[f(w_m^k) - f(w_0^k)] &\leq -(\alpha\gamma_c\lambda^2\tau_1^2 - 14\alpha^2\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2)) \sum_{t=0}^{m-1} \mathbb{E}[(\text{dist}(w_t^k, w^*))^2] \\ &\quad + 8m\alpha^2\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2)\mathbb{E}[(\text{dist}(\tilde{w}^k, w^*))^2]. \end{aligned} \quad (\text{A.70})$$

Here, considering the difference with the solution w^* in terms of the cost function value, we obtain

$$\begin{aligned} \mathbb{E}[f(w_m^k) - f(w_0^k)] &= \mathbb{E}[f(w_m^k) - f(w^*) - (f(w_0^k) - f(w^*))] \\ &\geq \frac{1}{2}\mathbb{E}[\lambda\tau_1^2(\text{dist}(w_m^k, w^*))^2 - \Lambda\tau_2^2(\text{dist}(w_0^k, w^*))^2]. \end{aligned}$$

Plugging the above into (A.70) yields

$$\begin{aligned} &\mathbb{E}[\lambda\tau_1^2(\text{dist}(w_m^k, w^*))^2 - \Lambda\tau_2^2(\text{dist}(w_0^k, w^*))^2] \\ &\leq -2\alpha(\gamma_c\lambda^2\tau_1^2 - 14\alpha\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2)) \sum_{t=0}^{m-1} \mathbb{E}[(\text{dist}(w_t^k, w^*))^2] \\ &\quad + 16m\alpha^2\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2)\mathbb{E}[(\text{dist}(\tilde{w}^k, w^*))^2]. \end{aligned}$$

Rearranging this gives

$$\begin{aligned} &2\alpha(\gamma_c\lambda^2\tau_1^2 - 14\alpha\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2)) \sum_{t=0}^{m-1} \mathbb{E}[(\text{dist}(w_t^k, w^*))^2] \\ &\leq \mathbb{E}[\Lambda\tau_2^2(\text{dist}(w_0^k, w^*))^2] - \mathbb{E}[\lambda\tau_1^2(\text{dist}(w_m^k, w^*))^2] \\ &\quad + 16m\alpha^2\Lambda\Gamma_c^2(\beta^2 + \tau_2^2C^2\theta^2)\mathbb{E}[(\text{dist}(\tilde{w}^k, w^*))^2]. \end{aligned} \quad (\text{A.71})$$

Now, addressing option I in Algorithm 1, which uses $\tilde{w}^{k+1} = g_{m_k}(w_1^k, \dots, w_m^k)$, we derive below from Lemma D.8 as

$$\begin{aligned}
& \frac{m}{4} 2\alpha(\gamma_c \lambda^2 \tau_1^2 - 14\alpha \Lambda \Gamma_c^2(\beta^2 + \tau_2^2 C^2 \theta^2)) \mathbb{E}[(\text{dist}(\tilde{w}^{k+1}, w^*))^2] \\
& \leq 2\alpha(\gamma_c \lambda^2 \tau_1^2 - 14\alpha \Lambda \Gamma_c^2(\beta^2 + \tau_2^2 C^2 \theta^2)) \\
& \quad \times \mathbb{E} \left[\sum_{t=0}^{m-1} (\text{dist}(w_t^k, w^*))^2 + (\text{dist}(w_m^k, w^*))^2 - (\text{dist}(w_0^k, w^*))^2 \right] \\
& \stackrel{(A.71)}{\leq} \mathbb{E}[\Lambda \tau_2^2 (\text{dist}(w_0^k, w^*))^2] - \mathbb{E}[\lambda \tau_1^2 (\text{dist}(w_m^k, w^*))^2] \\
& \quad + 16m\alpha^2 \Lambda \Gamma_c^2(\beta^2 + \tau_2^2 C^2 \theta^2) \mathbb{E}[(\text{dist}(\tilde{w}^k, w^*))^2] \\
& \quad + 2\alpha(\gamma_c \lambda^2 \tau_1^2 - 14\alpha \Lambda \Gamma_c^2(\beta^2 + \tau_2^2 C^2 \theta^2)) \mathbb{E}[(\text{dist}(w_m^k, w^*))^2 - (\text{dist}(w_0^k, w^*))^2] \\
& \leq (\Lambda \tau_2^2 - 2\alpha(\gamma_c \lambda^2 \tau_1^2 - 14\alpha \Lambda \Gamma_c^2(\beta^2 + \tau_2^2 C^2 \theta^2))) \mathbb{E}[(\text{dist}(w_0^k, w^*))^2] \\
& \quad + 16m\alpha^2 \Lambda \Gamma_c^2(\beta^2 + \tau_2^2 C^2 \theta^2) \mathbb{E}[(\text{dist}(\tilde{w}^k, w^*))^2] \\
& \quad - (\lambda \tau_1^2 - 2\alpha(\gamma_c \lambda^2 \tau_1^2 - 14\alpha \Lambda \Gamma_c^2(\beta^2 + \tau_2^2 C^2 \theta^2))) \mathbb{E}[(\text{dist}(w_m^k, w^*))^2].
\end{aligned}$$

Combining the relation $\Lambda \tau_2^2 > \lambda \tau_1^2$ and the assumption $\lambda \tau_1^2 > 2\alpha(\gamma_c \lambda^2 \tau_1^2 - 14\alpha \Lambda \Gamma_c^2(\beta^2 + \tau_2^2 C^2 \theta^2))$, since $w_0^k = \tilde{w}^k$, we obtain

$$\begin{aligned}
& \frac{m}{4} 2\alpha(\gamma_c \lambda^2 \tau_1^2 - 14\alpha \Lambda \Gamma_c^2(\beta^2 + \tau_2^2 C^2 \theta^2)) \mathbb{E}[(\text{dist}(\tilde{w}^{k+1}, w^*))^2] \\
& \leq (\Lambda \tau_2^2 + 16m\alpha^2 \Lambda \Gamma_c^2(\beta^2 + \tau_2^2 C^2 \theta^2)) \mathbb{E}[(\text{dist}(\tilde{w}^k, w^*))^2].
\end{aligned}$$

Finally, we obtain

$$\mathbb{E}[(\text{dist}(\tilde{w}^{k+1}, w^*))^2] \leq \frac{2(\Lambda \tau_2^2 + 16m\alpha^2 \Lambda \Gamma_c^2(\beta^2 + \tau_2^2 C^2 \theta^2))}{m\alpha(\gamma_c \lambda^2 \tau_1^2 - 14\alpha \Lambda \Gamma_c^2(\beta^2 + \tau_2^2 C^2 \theta^2))} \mathbb{E}[(\text{dist}(\tilde{w}^k, w^*))^2]. \quad (\text{A.72})$$

This completes the proof. \square

Remark D.12. From the proof of Lemma D.3, if we adopt the parallel translation as the vector transport, i.e., $\mathcal{T} = P$, the first two terms in (A.4) are equal to zero, and Υ_c in (A.53) gets smaller than that of the case of vector transport. This leads to a smaller Γ_c and a larger γ_c in Proposition D.6. Then, the smaller Γ_c and the larger γ_c leads to a smaller coefficient in (A.72) of Theorem 3.3. Consequently, the parallel translation can result in a faster local convergence rate.

We obtain the following corollary of the previous theorem with the case $R = \text{Exp}$ and $\mathcal{T} = P$.

Corollary D.13. Consider Algorithm 1 with $\mathcal{T} = P$ and $R = \text{Exp}$, i.e., the parallel translation and the exponential mapping case. Let \mathcal{M} be a Riemannian manifold and $w^* \in \mathcal{M}$ be a non-degenerate local minimizer of f (i.e., $\text{grad}f(w^*) = 0$ and the Hessian $\text{Hess}f(w^*)$ of f at w^* is positive definite). Suppose Assumptions 1 and 3 hold. Let the constants θ , and C in Lemma D.9. β_0 is the constant in Corollary D.10. Λ and λ are the constants in Lemmas C.1 and D.1, respectively. γ_c and Γ_c are the constants in Proposition D.6. Let α be a positive number satisfying $\lambda > 2\alpha(\gamma_c \lambda^2 - 7\alpha \Lambda \Gamma_c^2 \beta_0^2)$ and $\gamma_c \lambda^2 \tau_1^2 > 14\alpha \Lambda \Gamma_c^2(\beta^2 + \tau_2^2 C^2 \theta^2)$. It then follows that for any sequence $\{\tilde{w}^k\}$ generated by Algorithm 1 with Option II under a fixed

step-size $\alpha_t^k := \alpha$ and $m_k := m$ converging to w^* , there exists $0 < K_{th} < K$ such that for all $k > K_{th}$,

$$\mathbb{E}[(\text{dist}(\tilde{w}^{k+1}, w^*))^2] \leq \frac{2(\Lambda + 8m\alpha^2\Lambda\Gamma_c^2\beta_0^2)}{m\alpha(\gamma_c\lambda^2 - 7\alpha\Lambda\Gamma_c^2\beta_0^2)}\mathbb{E}[(\text{dist}(\tilde{w}^k, w^*))^2] \quad (\text{A.73})$$

Proof. The proof is given similarly to Theorem 3.3. We use Corollary D.10, and also set as $\theta = 0$ in Lemma C.14, and as $\tau_1 = \tau_2 = 1$ in Lemma C.15. \square

E Additional numerical experiments

In this section, we show additional numerical experiments which do not appear in the main text.

E.1 Matrix completion problem on synthetic datasets

E.1.1 Additional results

This section shows the results of six problem instances. Due to the page limitations, we only show the loss on a test set Φ , which is different from the training set Ω . The loss on the test set demonstrates the convergence speed to a good prediction accuracy of missing entries.


Case MC-S1: We first show the results of the comparison when the number of samples $N = 5000$, the dimension $d = 200$, the memory size $L = 10$, the oversampling ratio (OS) is 8, and the condition number (CN) is 50. We also add Gaussian noise $\sigma = 10^{-10}$. Figures A.1 show the results of 4 runs except the result shown in the main text, which corresponds to "run 1." They show superior performances than other algorithms.

Case MC-S2: influence on low sampling. We look into problem instances from scarcely sampled data, e.g. OS is 4. Other conditions are the same as **Case MC-S1**. From Figures A.2, we can find that the proposed algorithm gives much better and stabler performances against other algorithms.


Case MC-S3: influence on ill-conditioning. We consider the problem instances with higher condition number (CN) 100. Other conditions are the same as **Case MC-S1**. Figures A.3 show the superior performances of the proposed algorithm against other algorithms.

Case MC-S4: influence on higher noise. We consider noisy problem instances, where $\sigma = 10^{-6}$. Other conditions are the same as **Case MC-S1**. Figures A.4 show that the convergent MSE values are much higher than the other cases. Then, we can see the superior performance of the proposed R-SQN-VR against other algorithms.


Case MC-S5: influence on higher rank. We consider problem instances with higher rank, where $r = 10$. Other conditions are the same as **Case MC-S1**. From Figures A.5, the proposed R-SQN-VR still shows the superior performances against other algorithms. Grouse indicates the faster decrease of the MSE at the begging of the iterations. However, the convergent MSE values are much higher than those of others.




(a) run 2



(b) run 3




(c) run 4




(d) run 5


Figure A.1: Performance evaluations on low-rank MC problem (**Case MC-S1: baseline.**).




run 2



(b) run 3




(c) run 4




(d) run 5


Figure A.2: Performance evaluations on low-rank MC problem (**Case MC-S2: low sampling.**).




(a) run 2



(b) run 3




(c) run 4




(d) run 5


Figure A.3: Performance evaluations on low-rank MC problem (**Case MC-S3: ill-conditioning**).




(a) run 2



(b) run 3




(c) run 4




(d) run 5


Figure A.4: Performance evaluations on low-rank MC problem (**Case MC-S4: noisy data.**).




(a) run 2



(b) run 3



(c) run 4



(d) run 5

Figure A.5: Performance evaluations on low-rank MC problem (**Case MC-S5: higher rank.**).

E.1.2 Processing time experiments

The results in terms of the processing time is presented.

Case MC-S7: Comparison in terms of processing time. Because one major concern of second-order algorithms is, in general, higher computational processing load than first-order algorithms, we additionally show the results in terms of the processing time. This evaluation addresses only R-SGD, R-SVRG and R-SQN-VR because the code structures of them are similar whereas the batch-based algorithms, i.e., R-SD and R-L-BFGS, have completely different implementations. Figures A.6 (a)-(e) show the results of the relationship between test MSE and the processing time [sec]. From the figures, as expected, R-SGD gains much faster speed in comparison with the results in terms of iteration than other algorithms. However, it should be noted that R-SGD suffers from the problem that it heavily decreases the convergence speed around the solution as reported in the literature. Comparing R-SQN-VR with R-SVRG, R-SQN-VR still gives better performance although R-SQN-VR requires one more additional vector transport of a gradient in each inner iteration and L vector transports of the curvature pairs at every outer epoch than R-SVRG does. Overall, R-SQN-VR outperforms R-SGD and R-SVRG in terms of the processing time. Consequently, we also have confirmed the effectiveness of the proposed R-SQN-VR from the viewpoint of processing time.




Figure A.6: Performance evaluations on low-rank MC problem (Case MC-S7).

Finally, Figure A.7 shows the results when the memory size of L is changed in R-SQN-VR. Comparing the results with Figure 1 (h), the lower size cases improved their results very slightly, but we do not observe a big advantage of lower memory sizes in terms of processing load. From these results of both the convergence speed and the processing load, we cannot conclude which size of L is the best. This should be left to a future research topic.





Figure A.7: Performance evaluations on low-rank MC problem (processing time) (**Case MC-S6: different memory sizes**).

E.2 Matrix completion problem on MovieLens 1M dataset


Figures A.8 and A.9 show the results of the cases of $r = 10$ (**MC-R1: lower rank**) and $r = 20$ (**MC-R2: higher rank**). They show the convergence plots of the training error on Ω and the test error on Φ for all the five runs when rank $r = 10$ and $r = 20$, respectively. They show that the proposed R-SQN-VR give good performances on other algorithms in all runs.



(a-1) run 1

(a-2) run 2


(a-3) run 3



(a-4) run 4

(a-5) run 5


(a) MSE on train set Ω



(b-1) run 1

(b-2) run 2

(b-3) run 3




(b-4) run 4

(b-5) run 5

(b) MSE on test set Φ


Figure A.8: Performance evaluations on low-rank MC problem (**MC-R1: lower rank**).



(a-1) run 1

(a-2) run 2


(a-3) run 3



(a-4) run 4

(a-5) run 5


(a) MSE on train set Ω



(b-1) run 1

(b-2) run 2

(b-3) run 3



(b-4) run 4

(b-5) run 5

(b) MSE on test set Φ

Figure A.9: Performance evaluations on low-rank MC problem (**MC-R2: higher rank**).