

Resilience: A Criterion for Learning in the Presence of Arbitrary Outliers

Jacob Steinhardt^{*}

Moses Charikar[†]

Gregory Valiant[‡]

JSTEINHA@STANFORD.EDU

MOSES@STANFORD.EDU

VALIANT@STANFORD.EDU

Abstract

We introduce a criterion, *resilience*, which allows properties of a dataset (such as its mean or best low rank approximation) to be robustly computed, even in the presence of a large fraction of arbitrary additional data. Resilience is a weaker condition than most other properties considered so far in the literature, and yet enables robust estimation in a broader variety of settings. We provide new information-theoretic results on robust distribution learning, robust estimation of stochastic block models, and robust mean estimation under bounded k th moments. We also provide new algorithmic results on robust distribution learning, as well as robust mean estimation in ℓ_p -norms. Among our proof techniques is a method for pruning a high-dimensional distribution with bounded 1st moments to a stable “core” with bounded 2nd moments, which may be of independent interest.

Keywords: robust learning, outliers, stochastic block models, p -norm estimation, low rank approximation

1. Introduction

What are the fundamental properties that allow one to robustly learn from a dataset, even if some fraction of that dataset consists of arbitrarily corrupted data? While much work has been done in the setting of noisy data, or for restricted families of outliers, it is only recently that provable algorithms for learning in the presence of a large fraction of arbitrary (and potentially adversarial) data have been formulated in high-dimensional settings (Klivans et al., 2009; Xu et al., 2010; Diakonikolas et al., 2016a; Lai et al., 2016; Steinhardt et al., 2016; Charikar et al., 2017). In this work, we formulate a conceptually simple criterion that a dataset can satisfy—*resilience*—which guarantees that properties such as the mean of that dataset can be robustly estimated even if a large fraction of additional arbitrary data is inserted.

To illustrate our setting, consider the following game between Alice (the adversary) and Bob. First, a set $S \subseteq \mathbb{R}^d$ of $(1 - \epsilon)n$ points is given to Alice. Alice then adds ϵn additional points to S to create a new set $\tilde{S}$, and passes $\tilde{S}$ to Bob. Bob wishes to output a parameter $\hat{\mu}$ that is as close as possible to the mean μ of the points in the original set S , with error measured according to some norm $\|\hat{\mu} - \mu\|$. The question is: how well can Bob do, assuming that Alice is an adversary with knowledge of Bob’s algorithm?

The above game models mean estimation in the presence of arbitrary outliers; one can easily consider other problems as well (e.g. regression) but we will focus on mean estimation in this paper.

With no assumptions on S , Bob will clearly incur arbitrarily large error in the worst case—Alice can add points arbitrarily far away from the true mean μ , and Bob has no way of telling whether those points actually belong to S or were added by Alice. A first pass assumption is to suppose that S has diameter at most ρ ; then by discarding points that are very far away from most other points, Bob can obtain error $\mathcal{O}(\epsilon\rho)$. However, in most high-dimensional settings, the diameter ρ grows polynomially with the dimension

^{*} JS was supported by a Fannie & John Hertz Foundation Fellowship, an NSF Graduate Research Fellowship, and a Future of Life Institute grant.

[†] MC was supported by NSF grants CCF-1617577, CCF-1302518 and a Simons Investigator Award.

[‡] GV was supported by NSF CAREER Award CCF-1351108 and a Sloan Research Fellowship.




Figure 1: Illustration of the robust mean estimation setting. First, a set of points (blue) is given to Alice, who adds an ϵ fraction of adversarially chosen points (red). Bob’s goal is to output the mean of the original set (indicated in green).

d (e.g. the d -dimensional hypercube has ℓ_2 -diameter $\Theta(\sqrt{d})$). Subtler criteria are therefore needed to obtain dimension-independent bounds in most settings of interest.

Recently, [Diakonikolas et al. \(2016a\)](#) showed that Bob can incur ℓ_2 -error $\mathcal{O}(\epsilon\sqrt{\log(1/\epsilon)})$ when the points in S are drawn from a d -dimensional Gaussian, while [Lai et al. \(2016\)](#) concurrently showed that Bob can incur ℓ_2 -error $\mathcal{O}(\sqrt{\epsilon}\log(d))$ if the points in S are drawn from a distribution with bounded 4th moments. Since then, a considerable amount of additional work has studied high-dimensional estimation in the presence of adversaries, which we discuss in detail below. However, in general, both Bob’s strategy and its analysis tend to be quite complex, and specialized to particular distributional assumptions. This raises the question—is it possible to formulate a general and simple-to-understand criterion for the set S under which Bob has a (possibly inefficient) strategy for incurring small error?

In this paper, we provide such a criterion; we identify an assumption—*resilience*—on the set S , under which Bob has a straightforward exponential-time algorithm for estimating μ accurately. This yields new information-theoretic bounds for a number of robust learning problems, including robust learning of stochastic block models, of discrete distributions, and of distributions with bounded k th moments. We also identify additional assumptions under which Bob has an efficient (polynomial-time) strategy for estimating μ , which yields an efficient algorithm for robust learning of discrete distributions, as well as for robust mean estimation in ℓ_p -norms.

The resilience condition is essentially that the mean of every large subset of S must be close to the mean of all of S . More formally, for a norm $\|\cdot\|$, our criterion is as follows:

Definition 1 (Resilience) *A set of points $\{x_i\}_{i \in S}$ lying in $\mathbb{R}^d$ is (σ, ϵ) -resilient in a norm $\|\cdot\|$ around a point μ if, for all subsets $T \subseteq S$ of size at least $(1 - \epsilon)|S|$,*

$$\left\| \frac{1}{|T|} \sum_{i \in T} (x_i - \mu) \right\| \leq \sigma. \quad (1)$$

More generally, a distribution p is said to be (σ, ϵ) -resilient if $\|\mathbb{E}[x - \mu \mid E]\| \leq \sigma$ for every event E of probability at least $1 - \epsilon$.

In the definition above, μ need not equal the mean of S ; this distinction is useful in statistical settings where the sample mean of a finite set of points differs slightly from the true mean. However, (1) implies that μ differs from the mean of S by at most σ .

Importantly, Definition 1 is satisfied with high probability by a finite sample in many settings. For instance, samples from a distribution with k th moments bounded by σ will be $(\mathcal{O}(\sigma\epsilon^{1-1/k}), \epsilon)$ -resilient in the ℓ_2 -norm with high probability. Resilience also holds with high probability under many other natural distributional assumptions, discussed in more detail in Sections 1.1, 6, and 7.

Assuming that the original set S is (σ, ϵ) -resilient, Bob’s strategy is actually quite simple—find *any* large (σ, ϵ) -resilient subset S' of the corrupted set $\tilde{S}$, and output the mean of S' . By pigeonhole, S' and S have large intersection, and hence by (1) must have similar means. We establish this formally in Section 2.

Pleasingly, resilience reduces the question of whether Bob can win the game to a purely algorithmic question—that of finding any large resilient set. Rather than wondering whether it is even information-theoretically possible to estimate μ , we can instead focus on efficiently finding resilient subsets of $\tilde{S}$.

We provide one such algorithm in Section 4, assuming that the norm $\|\cdot\|$ is *strongly convex* and that we can approximately solve a certain generalized eigenvalue problem in the dual norm. When specialized to the ℓ_1 -norm, our general algorithm yields an efficient procedure for robust learning of discrete distributions.

In the remainder of this section, we will outline our main results, starting with information-theoretic results and then moving on to algorithmic results. In Section 1.1, we show that resilience is indeed information-theoretically sufficient for robust mean estimation. In Section 1.2, we then provide finite-sample bounds showing that resilience holds with high probability for i.i.d. samples from a distribution.

In Section 1.3, we turn our attention to algorithmic bounds. We identify a property—bounded variance in the dual norm—under which efficient algorithms exist. We then show that, as long as the norm is strongly convex, every resilient set has a large subset with bounded variance, thus enabling efficient algorithms. This connection between resilience and bounded variance is the most technically non-trivial component of our results, and may be of independent interest.

Both our information-theoretic and algorithmic bounds yield new results in concrete settings, which we discuss in the corresponding subsections. In Section 1.4, we also discuss an extension of resilience to low-rank matrix approximation, which enables us to derive new bounds in that setting as well. In Section 1.5 we outline the rest of the paper and point to technical highlights, and in Section 1.6 we discuss related work.

1.1. Information-Theoretic Sufficiency

First, we show that resilience is indeed information-theoretically sufficient for robust recovery of the mean μ . In what follows, we use $\sigma_*(\epsilon)$ to denote the smallest σ such that S is (σ, ϵ) -resilient.

Proposition 2 *Suppose that $\tilde{S} = \{x_1, \dots, x_n\}$ contains a set S of size $(1 - \epsilon)n$ that is resilient around μ (where S and μ are both unknown). Then if $\epsilon < \frac{1}{2}$, it is possible to recover a $\hat{\mu}$ such that $\|\hat{\mu} - \mu\| \leq 2\sigma_*(\frac{\epsilon}{1-\epsilon})$.*

More generally, if $|S| \geq \alpha n$ (even if $\alpha < \frac{1}{2}$), it is possible to output a (random) $\hat{\mu}$ such that $\|\hat{\mu} - \mu\| \leq \frac{16}{\alpha}\sigma_(\frac{\alpha}{4})$ with probability at least $\frac{\alpha}{2}$.*

The first part says that robustness to an ϵ fraction of outliers depends on resilience to a $\frac{\epsilon}{1-\epsilon}$ fraction of deletions. Thus, Bob has a good strategy as long as $\sigma_*(\frac{\epsilon}{1-\epsilon})$ is small.

The second part, which is more surprising, says that Bob has a good strategy *even if the majority of $\tilde{S}$ is controlled by Alice*. Here one cannot hope for recovery in the usual sense, because if $\alpha = \frac{1}{2}$ (i.e., Alice controls half the points) then Alice can make $\tilde{S}$ the disjoint union of two identical copies of S (one of which is shifted by a large amount) and Bob has no way of determining which of the two copies is the true S . Nevertheless, in this situation Bob can still identify S (and hence μ) with probability $\frac{1}{2}$; more generally, the second part of Proposition 2 says that if $|S| = \alpha|\tilde{S}|$ then Bob can identify μ with probability at least $\frac{\alpha}{2}$.

The fact that estimation is possible even when $\alpha < \frac{1}{2}$ was first established by Steinhardt et al. (2016) in a crowdsourcing setting, and later by Charikar et al. (2017) in a number of settings including mean

estimation. Apart from being interesting due to its unexpectedness, estimation in this regime has immediate implications for robust estimation of mixtures of distributions (by considering each mixture component in turn as the “good” set S) or of planted substructures in random graphs. We refer the reader to [Charikar et al. \(2017\)](#) for a full elaboration of this point.

The proof of Proposition 2, given in detail in Section 2, is a pigeonhole argument. For the $\epsilon < \frac{1}{2}$ case, we simply search for any large resilient set S' and output its mean; then S and S' must have large overlap, and by resilience their means must both be close to the mean of their intersection, and hence to each other.

For the general case where $|S| = \alpha|\tilde{S}|$ (possibly with $\alpha < \frac{1}{2}$), a similar pigeonhole argument applies but we now need to consider a covering of $\tilde{S}$ by $\frac{2}{\alpha}$ approximately disjoint sets $S'_1, \dots, S'_{2/\alpha}$. We can show that the true set S must overlap at least one of these sets by a decent amount, and so outputting the mean of one of these sets at random gives a good approximation to the mean of S with probability $\frac{\alpha}{2}$.

1.2. Finite-Sample Concentration

While Proposition 2 provides a deterministic condition under which robust mean estimation is possible, we would also like a way of checking that resilience holds with high probability given samples $x_1, \dots, x_n$ from a distribution p . First, we provide an alternate characterization of resilience which says that a distribution is resilient if it has *thin tails* in every direction:

Lemma 3 *Given a norm $\|\cdot\|$, define the dual norm $\|v\|_* = \sup_{\|x\| \leq 1} \langle v, x \rangle$. For a fixed vector v , let $\tau_\epsilon(v)$ denote the ϵ -quantile of $\langle x - \mu, v \rangle$: $\mathbb{P}_{x \sim p}[\langle x - \mu, v \rangle \geq \tau_\epsilon(v)] = \epsilon$. Then, p is (σ, ϵ) -resilient around its mean μ if and only if*

$$\mathbb{E}_p[\langle x - \mu, v \rangle \mid \langle x - \mu, v \rangle \geq \tau_\epsilon(v)] \leq \frac{1 - \epsilon}{\epsilon} \sigma \text{ whenever } \|v\|_* \leq 1. \quad (2)$$

In other words, if we project onto any unit vector v in the dual norm, the ϵ -tail of $x - \mu$ must have mean at most $\frac{1 - \epsilon}{\epsilon} \sigma$. Thus, for instance, a distribution with variance at most σ_0^2 along every unit vector would have $\sigma = \mathcal{O}(\sigma_0 \sqrt{\epsilon})$. Note that Lemma 3 requires μ to be the mean, rather than an arbitrary vector as before.

We next provide a meta-result establishing that resilience of a population distribution p very generically transfers to a finite set of samples from that distribution. The number of samples necessary depends on two quantities B and $\log M$ that will be defined in detail later; for now we note that they are ways of measuring the effective dimension of the space.

Proposition 4 *Suppose that a distribution p is (σ, ϵ) -resilient around its mean μ with $\epsilon < \frac{1}{2}$. Let B be such that $\mathbb{P}[\|x - \mu\| \geq B] \leq \epsilon/2$. Also let M be the covering number of the unit ball in the dual norm $\|\cdot\|_*$.*

Then, given n samples $x_1, \dots, x_n \sim p$, with probability $1 - \delta - \exp(-\epsilon n/6)$ there is a subset T of $(1 - \epsilon)n$ of the x_i such that T is (σ', ϵ) -resilient with $\sigma' = \mathcal{O}\left(\sigma \cdot \left(1 + \sqrt{\frac{\log(M/\delta)}{\epsilon^2 n}} + \frac{(B/\sigma) \log(M/\delta)}{n}\right)\right)$.

Note that Proposition 4 only guarantees resilience on a $(1 - \epsilon)n$ -element subset of the x_i , rather than all of $x_1, \dots, x_n$. From the perspective of robust estimation, this is sufficient, as we can simply regard the remaining ϵn points as part of the “bad” points controlled by Alice. This weaker requirement seems to be actually necessary to achieve Proposition 4, and was also exploited in [Charikar et al. \(2017\)](#) to yield improved bounds for a graph partitioning problem. There has been a great deal of recent interest in showing how to “prune” samples to achieve faster rates in random matrix settings ([Guédon and Vershynin, 2014](#); [Le et al., 2015](#); [Rebrova and Tikhomirov, 2015](#); [Rebrova and Vershynin, 2016](#)), and we think the general investigation of such pruning results is likely to be fruitful.

We remark that the sample complexity in Proposition 4 is suboptimal in many cases, requiring roughly $d^{1.5}$ samples when d samples would suffice. At the end of the next subsection we discuss a tighter but more specialized bound based on spectral graph sparsification.

Applications. Propositions 2 and 4 together give us a powerful tool for deriving information-theoretic robust recovery results: one needs simply establish resilience for the population distribution p , then use Proposition 4 to obtain finite sample bounds and Proposition 2 to obtain robust recovery guarantees. We do this in three illustrative settings: ℓ_2 mean estimation, learning discrete distributions, and stochastic block models. We outline the results below; formal statements and proofs are deferred to Section 7.

Mean estimation in ℓ_2 -norm. Suppose that a distribution on $\mathbb{R}^d$ has bounded k th moments: $\mathbb{E}_{x \sim p}[|\langle x - \mu, v \rangle|^k]^{1/k} \leq \sigma \|v\|_2$ for all v for some $k \geq 2$. Then p is $(\mathcal{O}(\sigma \epsilon^{1-1/k}), \epsilon)$ -resilient in the ℓ_2 -norm. Propositions 4 and 2 then imply that, given $n \geq \frac{d^{1.5}}{\epsilon} + \frac{d}{\epsilon^2}$ samples from p , and an ϵ -fraction of corruptions, it is possible to recover the mean to ℓ_2 -error $\mathcal{O}(\sigma \epsilon^{1-1/k})$. Moreover, if only an α -fraction of points are good, the mean can be recovered to error $\mathcal{O}(\sigma \alpha^{-1/k})$ with probability $\Omega(\alpha)$.

The $d^{1.5}/\epsilon$ term in the sample complexity is likely loose, and we believe the true dependence on d is at most $d \log(d)$. This looseness comes from Proposition 4, which uses a naïve covering argument and could potentially be improved with more sophisticated tools. Nevertheless, it is interesting that resilience holds long before the empirical k th moments concentrate, which would require $d^{k/2}$ samples.

Distribution learning. Suppose that we are given k -tuples of independent samples from a discrete distribution: $p = \pi^k$, where π is a distribution on $\{1, \dots, m\}$. By taking the empirical average of the k samples from π , we can treat a sample from p as an element in the m -dimensional simplex Δ_m . This distribution turns out to be resilient in the ℓ_1 -norm with $\sigma = \mathcal{O}(\epsilon \sqrt{\log(1/\epsilon)/k})$, which allows us to estimate p in the ℓ_1 -norm (i.e., total variation norm) and recover $\hat{\pi}$ such that $\|\hat{\pi} - \pi\|_{TV} = \mathcal{O}(\epsilon \sqrt{\log(1/\epsilon)/k})$. This reveals a pleasing “error correction” property: if we are given k samples at a time, either all or none of which are good, then our error is $\sqrt{k}$ times smaller than if we only observe the samples individually.

Stochastic block models. Finally, we consider the *semi-random stochastic block model* studied in Charikar et al. (2017) (described in detail in Section 7.3). For a graph on n vertices, this model posits a subset S of αn “good” vertices, which are connected to each other with probability $\frac{a}{n}$ and to the other (“bad”) vertices with probability $\frac{b}{n}$ (where $b < a$); the connections among the bad vertices can be arbitrary. The goal is to recover the set S .

We think of each row of the adjacency matrix as a vector in $\{0, 1\}^n$, and show that for the good vertices these vectors are resilient in a truncated ℓ_1 -norm $\|x\|$, defined as the sum of the αn largest coordinates of x (in absolute value). In this case, we have $\sigma = \mathcal{O}(\alpha \sqrt{a \log(2/\alpha)})$ (this requires a separate argument from Proposition 4 to get tight bounds). Applying Proposition 2, we find that we are able to recover (with probability $\frac{\alpha}{2}$) a set $\hat{S}$ with

$$\frac{1}{\alpha n} |S \Delta \hat{S}| = \mathcal{O} \left(\frac{a \log(2/\alpha)}{(a-b)^2 \alpha^2} \right). \quad (3)$$

In particular, we get non-trivial guarantees as long as $\frac{(a-b)^2}{a} \gg \frac{\log(2/\alpha)}{\alpha^2}$. Charikar et al. (2017) derive a weaker (but computationally efficient) bound when $\frac{(a-b)^2}{a} \gg \frac{\log(2/\alpha)}{\alpha^3}$, and remark on the similarity to the famous *Kesten-Stigum threshold* $\frac{(a-b)^2}{a} \gg \frac{1}{\alpha^2}$, which is the conjectured threshold for computationally efficient recovery in the classical stochastic block model (see Decelle et al. (2011) for the conjecture, and Mossel et al. (2013); Massoulié (2014) for a proof in the two-block case). Our information-theoretic upper bound matches the Kesten-Stigum threshold up to a $\log(2/\alpha)$ factor. We conjecture that this upper bound is tight; some evidence for this is given in Steinhardt (2017), which provides a nearly matching information-theoretic lower bound when $a = 1$, $b = \frac{1}{2}$.

1.3. Strong Convexity, Second Moments, and Efficient Algorithms

Most existing algorithmic results on robust mean estimation rely on analyzing the empirical covariance of the data in some way (see, e.g., [Lai et al. \(2016\)](#); [Diakonikolas et al. \(2016a\)](#); [Balakrishnan et al. \(2017\)](#)). In this section we establish connections between bounded covariance and resilience, and show that in a very general sense, bounded covariance is indeed sufficient to enable robust mean estimation.

Given a norm $\|\cdot\|$, we say that a set of points $x_1, \dots, x_n$ has *variance bounded by σ_0^2* in that norm if $\frac{1}{n} \sum_{i=1}^n \langle x_i - \mu, v \rangle^2 \leq \sigma_0^2 \|v\|_*^2$ (recall $\|\cdot\|_*$ denotes the dual norm). Since this implies a tail bound along every direction, it is easy to see (c.f. [Lemma 3](#)) that a set with variance bounded by σ_0^2 is $(\mathcal{O}(\sigma_0 \sqrt{\epsilon}), \epsilon)$ -resilient around its mean for all $\epsilon < \frac{1}{2}$. Therefore, bounded variance implies resilience.

An important result is that the converse is also true, *provided the norm is strongly convex*. We say that a norm $\|\cdot\|$ is γ -strongly convex if $\|x + y\|^2 + \|x - y\|^2 \geq 2(\|x\|^2 + \gamma\|y\|^2)$ for all $x, y \in \mathbb{R}^d$.¹ As an example, the ℓ_p -norm is $(p - 1)$ -strongly convex for $p \in (1, 2]$. For strongly convex norms, we show that any resilient set has a highly resilient “core” with bounded variance:

Theorem 5 *If S is $(\sigma, \frac{1}{2})$ -resilient in a γ -strongly convex norm $\|\cdot\|$, then S contains a set S_0 of size at least $\frac{1}{2}|S|$ with bounded variance: $\frac{1}{|S_0|} \sum_{i \in S_0} \langle x_i - \mu, v \rangle^2 \leq \frac{288\sigma^2}{\gamma} \|v\|_*^2$ for all v .*

Using [Lemma 3](#), we can show that $(\sigma, \frac{1}{2})$ -resilience is equivalent to having bounded 1st moments in every direction; [Theorem 5](#) can thus be interpreted as saying that any set with bounded 1st moments can be pruned to have bounded 2nd moments.

We found this result quite striking—the fact that [Theorem 5](#) can hold with no dimension-dependent factors is far from obvious. In fact, if we replace 2nd moments with 3rd moments or take a non-strongly-convex norm then the analog of [Theorem 5](#) is false: we incur polynomial factors in the dimension even if S is the standard basis of $\mathbb{R}^d$ (see [Section A](#) for details). The proof of [Theorem 5](#) involves minimax duality and Khintchine’s inequality. We can also strengthen [Theorem 5](#) to yield S_0 of size $(1 - \epsilon)|S|$. The proofs of both results are given in [Section 3](#) and may be of independent interest.

Algorithmic results. Given points with bounded variance, we establish algorithmic results assuming that one can solve the “generalized eigenvalue” problem $\max_{\|v\|_* \leq 1} v^\top A v$ up to some multiplicative accuracy κ . Specifically, we make the following assumption:

Assumption 6 (κ -Approximability) *There is a convex set $\mathcal{P}$ of PSD matrices such that*

$$\sup_{\|v\|_* \leq 1} v^\top A v \leq \sup_{M \in \mathcal{P}} \langle A, M \rangle \leq \kappa \sup_{\|v\|_* \leq 1} v^\top A v \quad (4)$$

for every PSD matrix A . Moreover, it is possible to optimize linear functions over $\mathcal{P}$ in polynomial time.

A result of [Nesterov \(1998\)](#) implies that this is true with $\kappa = \mathcal{O}(1)$ if $\|\cdot\|_*$ is any “quadratically convex” norm, which includes the ℓ_q -norms for $q \in [2, \infty]$. Also, while we do not use it in this paper, one can sometimes exploit weaker versions of [Assumption 6](#) that only require $\sup_{M \in \mathcal{P}} \langle A, M \rangle$ to be small for certain matrices A ; see for instance [Li \(2017\)](#), which obtains an algorithm for robust sparse mean estimation even though [Assumption 6](#) (as well as strong convexity) fails to hold in that setting.

Our main algorithmic result is the following:

Theorem 7 *Suppose that $x_1, \dots, x_n$ contains a subset S of size $(1 - \epsilon)n$ whose variance around its mean μ is bounded by σ_0^2 in the norm $\|\cdot\|$. Also suppose that [Assumption 6](#) holds for the dual norm $\|\cdot\|_*$. Then, if $\epsilon \leq \frac{1}{4}$, there is a polynomial-time algorithm whose output satisfies $\|\hat{\mu} - \mu\| = \mathcal{O}(\sigma_0 \sqrt{\kappa \epsilon})$.*

1. In the language of Banach space theory, this is also referred to as having bounded co-type.

If, in addition, $\|\cdot\|$ is γ -strongly convex, then even if S only has size αn there is a polynomial-time algorithm such that $\|\hat{\mu} - \mu\| = \mathcal{O}\left(\frac{\sqrt{\kappa\sigma_0}}{\sqrt{\gamma\alpha}}\right)$ with probability $\Omega(\alpha)$.

This is essentially a more restrictive, but computationally efficient version of Proposition 2. We note that for the ℓ_2 -norm, the algorithm can be implemented as an SVD (singular value decomposition) combined with a filtering step; for more general norms, the SVD is replaced with a semidefinite program.

In the small- ϵ regime, Theorem 7 is in line with existing results which typically achieve errors of $\mathcal{O}(\sqrt{\epsilon})$ in specific norms (see for instance a concurrent result of Diakonikolas et al. (2017a)). While several papers achieve stronger rates of $\mathcal{O}(\epsilon^{3/4})$ (Lai et al., 2016) or $\tilde{\mathcal{O}}(\epsilon)$ (Diakonikolas et al., 2016a; Balakrishnan et al., 2017), these stronger results rely crucially on specific distributional assumptions such as Gaussianity. At the time of writing of this paper, no results obtained rates better than $\mathcal{O}(\sqrt{\epsilon})$ for any general class of distributions (even under strong assumptions such as sub-Gaussianity). After initial publication of this paper, Kothari and Steinhardt (2017) surpassed $\sqrt{\epsilon}$ and obtained rates of $\epsilon^{1-\gamma}$ for any $\gamma > 0$, for distributions satisfying the Poincaré isoperimetric inequality.

In the small- α regime, Theorem 7 generalizes the mean estimation results of Charikar et al. (2017) to norms beyond the ℓ_2 -norm. That paper achieves a better rate of $1/\sqrt{\alpha}$ (versus the $1/\alpha$ rate given here); we think it is likely possible to achieve the $1/\sqrt{\alpha}$ rate in general, but leave this for future work.

Applications. Because Assumption 6 holds for ℓ_p -norms, we can perform robust estimation in ℓ_p -norms for any $p \in [1, 2]$, as long as the data have bounded variance in the dual ℓ_q -norm (where $\frac{1}{p} + \frac{1}{q} = 1$); see Corollary 27 for a formal statement. This is the first efficient algorithm for performing robust mean estimation in any ℓ_p -norm with $p \neq 2$. The ℓ_1 -norm in particular is often a more meaningful metric than the ℓ_2 -norm in discrete settings, allowing us to improve on existing results.

Indeed, as in the previous section, suppose we are given k -tuples of samples from a discrete distribution π on $\{1, \dots, m\}$. Applying Theorem 7 with the ℓ_1 -norm yields an algorithm recovering a $\hat{\pi}$ with $\|\hat{\pi} - \pi\|_{TV} = \tilde{\mathcal{O}}(\sqrt{\epsilon/k})$.² In contrast, bounds using the ℓ_2 -norm would only yield $\|\hat{\pi} - \pi\|_2 = \mathcal{O}(\sqrt{\epsilon\pi_{\max}/k})$, which is substantially weaker when the maximum probability $\pi_{\max}$ is large. Our result has a similar flavor to that of Diakonikolas et al. (2016a) on robustly estimating binary product distributions, for which directly applying ℓ_2 mean estimation was also insufficient. We discuss our bounds in more detail in Section 8.

Finite-sample bound. To get the best sample complexity for the applications above, we provide an additional finite-sample bound focused on showing that a set of points has bounded variance. This is a simple but useful generalization of Proposition B.1 of Charikar et al. (2017); it shows that in a very generic sense, given d samples from a distribution on $\mathbb{R}^d$ with bounded population variance, we can find a subset of samples with bounded variance with high probability. It involves pruning the samples in a non-trivial way based on ideas from graph sparsification (Batson et al., 2012). The formal statement is given in Section 6.2.

1.4. Low-Rank Recovery

Finally, to illustrate that the idea behind resilience is quite general and not restricted to mean estimation, we also provide results on recovering a rank- k approximation to the data in the presence of arbitrary outliers. Given a set of points $[x_i]_{i \in S}$, let X_S be the matrix whose columns are the x_i . Our goal is to obtain a low-rank matrix P such that the operator norm $\|(I - P)X_S\|_2$ is not much larger than $\sigma_{k+1}(X_S)$, where σ_{k+1} denotes the $k + 1$ st singular value; we wish to do this even if S is corrupted to a set $\tilde{S}$ by adding arbitrary outliers.

As before, we start by formulating an appropriate resilience criterion:

2. The $\tilde{\mathcal{O}}$ notation suppresses log factors in m and ϵ ; the dependence on m can likely be removed with a more careful analysis.

Definition 8 (Rank-resilience) A set of points $[x_i]_{i \in S}$ in $\mathbb{R}^d$ is δ -rank-resilient if for all subsets T of size at least $(1 - \delta)|S|$, we have $\text{col}(X_T) = \text{col}(X_S)$ and $\|X_T^\dagger X_S\|_2 \leq 2$, where † is the pseudoinverse and col denotes column space.

Rank-resilience says that the variation in X should be sufficiently spread out: there should not be a direction of variation that is concentrated in only a δ -fraction of the points. Under rank-resilience, we can perform efficient rank- k recovery even in the presence of a δ -fraction of arbitrary data:

Theorem 9 Let $\delta \leq \frac{1}{3}$. If a set of n points contains a set S of size $(1 - \delta)n$ that is δ -rank-resilient, then it is possible to efficiently recover a matrix P of rank at most $15k$ such that $\|(I - P)X_S\|_2 = \mathcal{O}(\sigma_{k+1}(X_S))$.

The power of Theorem 9 comes from the fact that the error depends on σ_{k+1} rather than e.g. σ_2 , which is what previous results yielded. This distinction is crucial in practice, since most data have a few (but more than one) large singular values followed by many small singular values. Note that in contrast to Theorem 7, Theorem 9 only holds when S is relatively large: at least $(1 - \delta)n \geq \frac{2}{3}n$ in size.

1.5. Summary, Technical Highlights, and Roadmap

In summary, we have provided a deterministic condition on a set of points that enables robust mean estimation, and provided finite-sample bounds showing that this condition holds with high probability in many concrete settings. This yields new results for distribution learning, stochastic block models, mean estimation under bounded moments, and mean estimation in ℓ_p norms. We also provided an extension of our condition that yields results for robust low-rank recovery.

Beyond the results themselves, the following technical aspects of our work may be particularly interesting: The proof of Proposition 2 (establishing that resilience is indeed sufficient for robust estimation), while simple, is a nice pigeonhole argument that we found to be conceptually illuminating.

In addition, the proof of Theorem 5, on pruning resilient sets to obtain sets with bounded variance, exploits strong convexity in a non-trivial way in conjunction with minimax duality; we think it reveals a fairly non-obvious geometric structure in resilient sets, and also shows how the ability to prune points can yield sets with meaningfully stronger properties.

Finally, in the proof of our algorithmic result (Theorem 7), we establish an interesting generalization of the inequality $\sum_{i,j} X_{ij}^2 \leq \text{rank}(X) \cdot \|X\|_2^2$, which holds not just for the ℓ_2 -norm but for any strongly convex norm. This is given as Lemma 18.

Roadmap. The rest of the paper is organized as follows. In Section 2, we prove our information-theoretic recovery result for resilient sets (Proposition 2). In Section 3, we prove Theorem 5 establishing that all resilient sets in strongly convex norms contain large subsets with bounded variance; we also prove a more precise version of Theorem 5 in Section 3.1. In Section 4, we prove our algorithmic results, warming up with the ℓ_2 -norm (Section 4.1) and then moving to general norms (Section 4.2). In Section 5, we prove our results on rank- k recovery. In Section 6, we present and prove the finite-sample bounds discussed in Section 1.2. In Sections 7 and 8, we provide applications of our information-theoretic and algorithmic results, respectively.

1.6. Related Work

A number of authors have recently studied robust estimation and learning in high-dimensional settings: Lai et al. (2016) study mean and covariance estimation, while Diakonikolas et al. (2016a) focus on estimating Gaussian and binary product distributions, as well as mixtures thereof; note that this implies mean/covariance estimation of the corresponding distributions. Charikar et al. (2017) recently showed that robust estimation is possible even when the fraction α of “good” data is less than $\frac{1}{2}$. We refer to these papers

for an overview of the broader robust estimation literature; since those papers, a number of additional results have also been published: [Diakonikolas et al. \(2017a\)](#) provide a case study of various robust estimation methods in a genomic setting, [Balakrishnan et al. \(2017\)](#) study sparse mean estimation, and others have studied problems including regression, Bayes nets, planted clique, and several other settings ([Diakonikolas et al., 2016b,c, 2017b,c](#); [Kane et al., 2017](#); [Meister and Valiant, 2017](#)).

Special cases of the resilience criterion are implicit in some of these earlier works; for instance, ℓ_2 -resilience appears in equation (9) in [Diakonikolas et al. \(2016a\)](#), and resilience in a sparsity-inducing norm appears in Theorem 4.5 of [Li \(2017\)](#). However, these conditions typically appear concurrently with other stronger conditions, and the general sufficiency of resilience for information-theoretic recovery appears to be unappreciated (for instance, [Li \(2017\)](#), despite already having implicitly established resilience, proves its information-theoretic results via reduction to a tournament lemma from [Diakonikolas et al. \(2016a\)](#)).

Low rank estimation was studied by [Lai et al. \(2016\)](#), but their bounds depend on the maximum eigenvalue $\|\Sigma\|_2$ of the covariance matrix, while our bound provides robust recovery guarantees in terms of lower singular values of Σ . (Some work, such as [Diakonikolas et al. \(2016a\)](#), shows how to estimate all of Σ in e.g. Frobenius norm, but appears to require the samples to be drawn from a Gaussian.)

2. Resilience and Robustness: Information-Theoretic Sufficiency

Recall the definition of resilience: S is (σ, ϵ) -resilient if $\|\frac{1}{|T|} \sum_{i \in T} (x_i - \mu)\| \leq \sigma$ whenever $T \subseteq S$ and $|T| \geq (1 - \epsilon)|S|$. Here we establish Proposition 2 showing that, if we ignore computational efficiency, resilience leads directly to an algorithm for robust mean estimation.

Proof (Proposition 2) We prove Proposition 2 via a constructive (albeit exponential-time) algorithm. To prove the first part, suppose that the true set S is $(\sigma, \frac{\epsilon}{1-\epsilon})$ -resilient around μ , and let S' be any set of size $(1 - \epsilon)n$ that is $(\sigma, \frac{\epsilon}{1-\epsilon})$ -resilient (around some potentially different vector μ'). We claim that μ' is sufficiently close to μ .

Indeed, let $T = S \cap S'$, which by the pigeonhole principle has size at least $(1 - 2\epsilon)n = \frac{1-2\epsilon}{1-\epsilon}|S| = (1 - \frac{\epsilon}{1-\epsilon})|S|$. Therefore, by the definition of resilience,

$$\left\| \frac{1}{|T|} \sum_{i \in T} (x_i - \mu) \right\| \leq \sigma. \quad (5)$$

But by the same argument, $\|\frac{1}{|T|} \sum_{i \in T} (x_i - \mu')\| \leq \sigma$ as well. By the triangle inequality, $\|\mu - \mu'\| \leq 2\sigma$, which completes the first part of the proposition.

For the second part, we need the following simple lemma relating ϵ -resilience to $(1 - \epsilon)$ -resilience:

Lemma 10 *For any $0 < \epsilon < 1$, a distribution/set is (σ, ϵ) -resilient around its mean μ if and only if it is $(\frac{1-\epsilon}{\epsilon}\sigma, 1 - \epsilon)$ -resilient. More generally, even if μ is not the mean then the distribution/set is $(\frac{2-\epsilon}{\epsilon}\sigma, 1 - \epsilon)$ -resilient. In other words, if $\|\frac{1}{|T|} \sum_{i \in T} (x_i - \mu)\| \leq \sigma$ for all sets T of size at least $(1 - \epsilon)n$, then $\|\frac{1}{|T'|} \sum_{i \in T'} (x_i - \mu)\| \leq \frac{2-\epsilon}{\epsilon}\sigma$ for all sets T' of size at least ϵn .*

Given Lemma 10, the second part of Proposition 2 is similar to the first part, but requires us to consider multiple resilient sets S_i rather than a single S' . Suppose S is $(\sigma, \frac{\alpha}{4})$ -resilient around μ —and thus also $(\frac{8}{\alpha}\sigma, 1 - \frac{\alpha}{4})$ -resilient by Lemma 10—and let $S_1, \dots, S_m$ be a maximal collection of subsets of $[n]$ such that:

1. $|S_j| \geq \frac{\alpha}{2}n$ for all j .
2. S_j is $(\frac{8}{\alpha}\sigma, 1 - \frac{\alpha}{2})$ -resilient around some point μ_j .
3. $S_j \cap S_{j'} = \emptyset$ for all $j \neq j'$.

Clearly $m \leq \frac{2}{\alpha}$. We claim that at least one of the μ_j is close to μ . By maximality of the collection $\{S_j\}_{j=1}^m$, it must be that $S_0 = S \setminus (S_1 \cup \dots \cup S_m)$ cannot be added to the collection. First suppose that $|S_0| \geq \frac{\alpha}{2}n$. Then S_0 is $(\frac{8}{\alpha}\sigma, 1 - \frac{\alpha}{2})$ -resilient (because any subset of $\frac{\alpha}{2}|S_0|$ points in S_0 is a subset of at least $\frac{\alpha}{4}|S|$ points in S). But this contradicts the maximality of $\{S_j\}_{j=1}^m$, so we must have $|S_0| < \frac{\alpha}{2}n$.

Now, this implies that $|S \cap (S_1 \cup \dots \cup S_m)| \geq \frac{\alpha}{2}n$, so by pigeonhole we must have $|S \cap S_j| \geq \frac{\alpha}{2}|S_j|$ for some j . Letting $T = S \cap S_j$ as before, we find that $|T| \geq \frac{\alpha}{2}|S_j| \geq \frac{\alpha}{4}|S|$ and hence by resilience of S_j and S we have $\|\mu - \mu_j\| \leq 2 \cdot (\frac{8}{\alpha}\sigma) = \frac{16}{\alpha}\sigma$. If we output one of the μ_j at random, we are then within the desired distance of μ with probability $\frac{1}{m} \geq \frac{\alpha}{2}$. ■

3. Powering up Resilience: Finding a Core with Bounded Variance

In this section we prove Theorem 5, which says that for strongly convex norms, every resilient set contains a core with bounded variance. Recall that this is important for enabling algorithmic applications that depend on a bounded variance condition.

First recall the definition of resilience (Definition 1): a set S is (σ, ϵ) -resilient if for every set $T \subseteq S$ of size $(1 - \epsilon)|S|$, we have $\|\frac{1}{|T|} \sum_{i \in T} (x_i - \mu)\| \leq \sigma$. For $\epsilon = \frac{1}{2}$, we observe that resilience in a norm is equivalent to having bounded first moments in the dual norm:

Lemma 11 *Suppose that S is $(\sigma, \frac{1}{2})$ -resilient in a norm $\|\cdot\|$, and let $\|\cdot\|_*$ be the dual norm. Then S has 1st moments bounded by 3σ : $\frac{1}{|S|} \sum_{i \in S} |\langle x_i - \mu, v \rangle| \leq 3\sigma \|v\|_*$ for all $v \in \mathbb{R}^d$.*

Conversely, if S has 1st moments bounded by σ , it is $(2\sigma, \frac{1}{2})$ -resilient.

The proof is routine and can be found in Section C. Supposing a set has bounded 1st moments, we will show that it has a large core with bounded second moments. This next result is *not* routine:

Proposition 12 *Let S be any set with 1st moments bounded by σ . Then if the norm $\|\cdot\|$ is γ -strongly convex, there exists a core S_0 of size at least $\frac{1}{2}|S|$ with variance bounded by $\frac{32\sigma^2}{\gamma}$. That is, $\frac{1}{|S_0|} \sum_{i \in S_0} |\langle x_i - \mu, v \rangle|^2 \leq \frac{32\sigma^2}{\gamma} \|v\|_*^2$ for all $v \in \mathbb{R}^d$.*

The assumptions seem necessary: i.e., such a core does not exist when $\|\cdot\|$ is the ℓ_p -norm with $p > 2$ (which is a non-strongly-convex norm), or with bounded 3rd moments for $p = 2$ (see Section A). The proof of Proposition 12 uses minimax duality and Khintchine's inequality (Khintchine, 1923). Note that Lemma 11 and Proposition 12 together imply Theorem 5.

Proof (Proposition 12) Without loss of generality take $\mu = 0$ and suppose that $S = [n]$. We can pose the problem of finding a resilient core as an integer program:

$$\min_{c \in \{0,1\}^n, \|c\|_1 \geq \frac{n}{2}} \max_{\|v\|_* \leq 1} \frac{1}{n} \sum_{i=1}^n c_i |\langle x_i, v \rangle|^2. \quad (6)$$

Here the variable c_i indicates whether the point i lies in the core S_0 . By taking a continuous relaxation and applying a standard duality argument, we obtain the following:

Lemma 13 *Suppose that for all m and all vectors $v_1, \dots, v_m$ satisfying $\sum_{j=1}^m \|v_j\|_*^2 \leq 1$, we have*

$$\frac{1}{n} \sum_{i=1}^n \sqrt{\sum_{j=1}^m |\langle x_i, v_j \rangle|^2} \leq B. \quad (7)$$

Then the value of (6) is at most $8B^2$.

The proof is straightforward and deferred to Section D. Now, to bound (7), let $s_1, \dots, s_m \in \{-1, +1\}$ be i.i.d. random sign variables. We have

$$\frac{1}{n} \sum_{i=1}^n \sqrt{\sum_{j=1}^m |\langle x_i, v_j \rangle|^2} \stackrel{(i)}{\leq} \mathbb{E}_{s_{1:m}} \left[\frac{\sqrt{2}}{n} \sum_{i=1}^n \left| \sum_{j=1}^m s_j \langle x_i, v_j \rangle \right| \right] \quad (8)$$

$$= \mathbb{E}_{s_{1:m}} \left[\frac{\sqrt{2}}{n} \sum_{i=1}^n \left| \left\langle x_i, \sum_{j=1}^m s_j v_j \right\rangle \right| \right] \quad (9)$$

$$\stackrel{(ii)}{\leq} \mathbb{E}_{s_{1:m}} \left[\sqrt{2} \sigma \left\| \sum_{j=1}^m s_j v_j \right\|_* \right] \quad (10)$$

$$\leq \sqrt{2} \sigma \mathbb{E}_{s_{1:m}} \left[\left\| \sum_{j=1}^m s_j v_j \right\|_*^2 \right]^{\frac{1}{2}}. \quad (11)$$

Here (i) is Khintchine's inequality (Haagerup, 1981) and (ii) is the assumed first moment bound. It remains to bound (11). The key is the following inequality asserting that the dual norm $\|\cdot\|_*$ is strongly smooth whenever $\|\cdot\|$ is strongly convex (c.f. Lemma 17 of Shalev-Shwartz (2007)):

Lemma 14 *If $\|\cdot\|$ is γ -strongly convex, then $\|\cdot\|_*$ is $(1/\gamma)$ -strongly smooth: $\frac{1}{2}(\|v+w\|_*^2 + \|v-w\|_*^2) \leq \|v\|_*^2 + (1/\gamma)\|w\|_*^2$.*

Applying Lemma 14 inductively to $\mathbb{E}_{s_{1:m}} \left[\left\| \sum_{j=1}^m s_j v_j \right\|_*^2 \right]$, we obtain

$$\mathbb{E}_{s_{1:m}} \left[\left\| \sum_{j=1}^m s_j v_j \right\|_*^2 \right] \leq \frac{1}{\gamma} \sum_{j=1}^m \|v_j\|_*^2 \leq \frac{1}{\gamma}. \quad (12)$$

Combining with (11), we have the bound $B \leq \sigma \sqrt{2/\gamma}$, which yields the desired result. $\blacksquare$

3.1. Finding Resilient Cores when $\alpha \approx 1$

Lemma 11 together with Proposition 12 show that a $(\sigma, \frac{1}{2})$ -resilient set has a core with bounded 2nd moments. One piece of looseness is that Proposition 12 only exploits resilience for $\epsilon = \frac{1}{2}$, and hence is not sensitive to the degree of (σ, ϵ) -resilience as $\epsilon \rightarrow 0$. In particular, it only yields a core S_0 of size $\frac{1}{2}|S|$, while we might hope to find a much larger core of size $(1-\epsilon)|S|$ for some small ϵ .

Here we tighten Proposition 12 to make use of finer-grained resilience information. Recall that we let $\sigma_*(\epsilon)$ denote the resilience over sets of size $(1-\epsilon)|S|$. For a given ϵ , our goal is to construct a core S_0 of size $(1-\epsilon)|S|$ with small second moments. The following key quantity will tell us how small the second moments can be:

$$\tilde{\sigma}_*(\epsilon) \stackrel{\text{def}}{=} \sqrt{\int_{\epsilon/2}^{1/2} u^{-2} \sigma_*(u)^2 du}. \quad (13)$$

The following proposition, proved in Section E, says that $\tilde{\sigma}_*$ controls the 2nd moments of S_0 :

Proposition 15 *Let S be any resilient set in a γ -strongly-convex norm. Then for any $\epsilon \leq \frac{1}{2}$, there exists a core S_0 of size $(1-\epsilon)|S|$ with variance bounded by $\mathcal{O}(\tilde{\sigma}_*^2(\epsilon)/\gamma)$.*

The proof is similar to Proposition 12, but requires more careful bookkeeping.

To interpret $\tilde{\sigma}_*$, suppose that $\sigma_*(\epsilon) = \sigma\epsilon^{1-1/r}$ for some $r \in [1, 2]$, which roughly corresponds to having bounded r th moments. Then $\tilde{\sigma}_*^2(\epsilon) = \sigma^2 \int_{\epsilon/2}^{1/2} u^{-2/r} du \leq \frac{\sigma^2}{2/r-1} \left(\frac{2}{\epsilon}\right)^{2/r-1}$. If $r = 1$ then a core of size $(1 - \epsilon)|S|$ might require second moments as large as $\frac{\sigma^2}{\epsilon}$; on the other hand, as $r \rightarrow 2$ the second moments can be almost as small as σ^2 . In general, $\tilde{\sigma}_*(\epsilon)$ is $\mathcal{O}(\sigma\epsilon^{1/2-1/r})$ if $r \in [1, 2]$, is $\mathcal{O}(\sigma\sqrt{\log(1/\epsilon)})$ if $r = 2$, and is $\mathcal{O}(\sigma)$ if $r > 2$.

4. Efficient Recovery Algorithms

We now turn our attention to the question of efficient algorithms. The main point of this section is to prove Theorem 7, which yields efficient robust mean estimation for a general class of norms.

4.1. Warm-Up: Recovery in ℓ_2 -norm

We first prove a warm-up to Theorem 7 which focuses on the ℓ_2 -norm. Our warm-up result is:

Proposition 16 *Let $x_1, \dots, x_n \in \mathbb{R}^d$, and let S be a subset of size αn with bounded variance in the ℓ_2 -norm: $\lambda_{\max}(\frac{1}{|S|} \sum_{i \in S} (x_i - \mu)(x_i - \mu)^\top) \leq \sigma^2$, where μ is the mean of S . Then there is an efficient randomized algorithm (Algorithm 1) which with probability $\Omega(\alpha)$ outputs a parameter $\hat{\mu}$ such that $\|\mu - \hat{\mu}\|_2 = \mathcal{O}(\frac{\sigma}{\alpha})$. Moreover, if $\alpha = 1 - \epsilon \geq \frac{3}{4}$ then $\|\mu - \hat{\mu}\|_2 = \mathcal{O}(\sigma\sqrt{\epsilon})$ with probability 1.*

At the heart of Algorithm 1 is the following optimization problem:

$$\begin{aligned} & \underset{W \in \mathbb{R}^{n \times n}}{\text{minimize}} && \|X - XW\|_2^2 \\ & \text{subject to} && 0 \leq W_{ji} \leq \frac{1}{\alpha n} \quad \forall i, j, \quad \sum_j W_{ji} = 1 \quad \forall i. \end{aligned} \tag{14}$$

Here $X \in \mathbb{R}^{d \times n}$ is the data matrix $[x_1 \cdots x_n]$ and $\|X - XW\|_2$ is the operator norm (maximum singular value) of $X - XW$. Note that (14) can be expressed as a semidefinite program; however, it can actually be solved more efficiently than this, via a singular value decomposition (see Section F for details).

The idea behind (14) is to re-construct each x_i as an average of αn other x_j . Note that by assumption we can always re-construct each element of S using the mean of S , and have small error. Intuitively, any element that cannot be re-constructed well must not lie in S , and can be safely removed. We do a soft form of removal by maintaining weights c_i on the points x_i (initially all 1), and downweighting points with high reconstruction error. We also maintain an active set $\mathcal{A}$ of points with $c_i \geq \frac{1}{2}$.

Informally, Algorithm 1 for estimating μ takes the following form:

1. Solve the optimization problem (14).
2. If the optimum is $\gg \sigma^2 n$, then find the columns of X that are responsible for the optimum being large, and downweight them.
3. Otherwise, if the optimum is $\mathcal{O}(\sigma^2 n)$, then take a low rank approximation W_0 to W , and return a randomly chosen column of XW_0 .

The hope in step 3 is that the low rank projection XW_0 will be close to μ for the columns belonging to S . The choice of operator norm is crucial: it means we can actually expect XW to be close to X (on the order of $\sigma\sqrt{n}$). In contrast, the Frobenius norm would scale as $\sigma\sqrt{nd}$.

Algorithm 1 Algorithm for recovering the mean of a set with bounded variance in ℓ_2 -norm.

- 1: Initialize $c_i = 1$ for all $i = 1, \dots, n$ and $\mathcal{A} = \{1, \dots, n\}$.
- 2: Let $Y \in \mathbb{R}^{d \times d}$ and $W \in \mathbb{R}^{A \times A}$ be the maximizer/minimizer of the saddle point problem

$$\max_{\substack{Y \succeq 0, \\ \text{tr}(Y) \leq 1}} \min_{\substack{0 \leq W_{ji} \leq \frac{4-\alpha}{\alpha(2+\alpha)n}, \\ \sum_j W_{ji} = 1}} \sum_{i \in \mathcal{A}} c_i (x_i - X_{\mathcal{A}} w_i)^\top Y (x_i - X_{\mathcal{A}} w_i). \quad (15)$$

- 3: Let $\tau_i^* = (x_i - X_{\mathcal{A}} w_i)^\top Y (x_i - X_{\mathcal{A}} w_i)$.
 - 4: **if** $\sum_{i \in \mathcal{A}} c_i \tau_i^* > 4n\sigma^2$ **then**
 - 5: For $i \in \mathcal{A}$, replace c_i with $\left(1 - \frac{\tau_i^*}{\tau_{\max}}\right) c_i$, where $\tau_{\max} = \max_{i \in \mathcal{A}} \tau_i^*$.
 - 6: For all i with $c_i < \frac{1}{2}$, remove i from $\mathcal{A}$.
 - 7: Go back to line 2.
 - 8: **end if**
 - 9: Let W_1 be the result of zeroing out all singular values of W that are greater than 0.9.
 - 10: Let $Z = X_{\mathcal{A}} W_0$, where $W_0 = (W - W_1)(I - W_1)^{-1}$.
 - 11: **if** $\text{rank}(Z) = 1$ **then**
 - 12: Output the average of the columns of $X_{\mathcal{A}}$.
 - 13: **else**
 - 14: Output a column of Z at random.
 - 15: **end if**
-

Finally, we note that $\|X - XW\|_2^2$ is equal to

$$\|X - XW\|_2^2 = \max_{Y \succeq 0, \text{tr}(Y) \leq 1} \sum_{i=1}^n (x_i - Xw_i)^\top Y (x_i - Xw_i), \quad (16)$$

which is the form we use in Algorithm 1.

Proof (Proposition 16) We need to show two things: (1) that the outlier removal step removes many more outliers than good points, and (2) that many columns of XW_0 are close to μ .

Outlier removal. To analyze the outlier removal step (step 2 above, or lines 5-6 of Algorithm 1), we make use of the following general lemma:

Lemma 17 *For any scalars τ_i and a , suppose that $\sum_{i \in \mathcal{A}} c_i \tau_i \geq 4a$ while $\sum_{i \in S \cap \mathcal{A}} c_i \tau_i \leq \alpha a$. Then the following invariants are preserved by lines 5-6 of Algorithm 1: (i) $\sum_{i \in S} (1 - c_i) \leq \frac{\alpha}{4} \sum_{i=1}^n (1 - c_i)$, and (ii) $|S \cap \mathcal{A}| \geq \frac{\alpha(2+\alpha)}{4-\alpha} n$.*

Lemma 17 says that we downweight points within S at least 4 times slower than we do overall (property i), and in particular we never remove too many points from S (property ii). This type of lemma is not new (cf. Lemma 4.5 of Charikar et al. (2017)) but for completeness we prove it in Section G.

We will show that we can take $a = n\sigma^2$ in Lemma 17, or in other words that $\sum_{i \in S \cap \mathcal{A}} c_i \tau_i^* \leq \alpha n\sigma^2$. Let $\tau_i(w) = (x_i - X_{\mathcal{A}} w)^\top (x_i - X_{\mathcal{A}} w)$, and note that $\tau_i^* = \tau_i(w_i) = \min\{\tau_i(w) \mid 0 \leq w_j \leq \frac{1}{\alpha n}, \sum_j w_j = 1\}$. This is because for a fixed Y , each of the w_i are optimized independently.

We can therefore bound τ_i^* by substituting any feasible $\hat{w}_i$. We will choose $\hat{W}_{ji} = \frac{\mathbb{I}[j \in S \cap \mathcal{A}]}{|S \cap \mathcal{A}|}$, in which case $X_{\mathcal{A}}\hat{w}_i = \hat{\mu}$, where $\hat{\mu}$ is the average of x_j over $S \cap \mathcal{A}$. Then we have

$$\sum_{i \in S \cap \mathcal{A}} c_i \tau_i^* \leq \sum_{i \in S \cap \mathcal{A}} c_i \tau_i(\hat{w}_i) \quad (17)$$

$$\leq \sum_{i \in S \cap \mathcal{A}} c_i (x_i - \hat{\mu})^\top Y (x_i - \hat{\mu}) \quad (18)$$

$$\stackrel{(i)}{\leq} \sum_{i \in S \cap \mathcal{A}} c_i (x_i - \mu)^\top Y (x_i - \mu) \quad (19)$$

$$\leq \sum_{i \in S} (x_i - \mu)^\top Y (x_i - \mu) \leq \alpha n \sigma^2 \text{tr}(Y) \leq \alpha n \sigma^2 \quad (20)$$

as desired; (i) is because the covariance around the mean ($\hat{\mu}$) is smaller than around any other point (μ).

Analyzing XW_0 . By Lemma 17, we will eventually exit the if statement and obtain $Z = X_{\mathcal{A}}W_0$. It therefore remains to analyze Z ; we will show in particular that $\|Z_{\mathcal{A} \cap S} - \mu \mathbb{1}^\top\|_F$ is small, where the subscript indicates restricting to the columns in $S \cap \mathcal{A}$. At a high level, it suffices to show that W_0 has low rank (so that Frobenius norm is close to spectral norm) and that XW_0 and X are close in spectral norm (note that X and $\mu \mathbb{1}^\top$ are close by assumption).

To bound $\text{rank}(W_0)$, note that the constraints in (15) imply that $\|W\|_F^2 \leq \frac{4-\alpha}{\alpha(2+\alpha)}$, and so at most $\frac{4-\alpha}{0.81\alpha(2+\alpha)}$ singular values of W can be greater than 0.9. Importantly, at most 1 singular value can be greater than 0.9 if $\alpha \geq \frac{3}{4}$, and at most $\mathcal{O}(\frac{1}{\alpha})$ can be in general. Therefore, $\text{rank}(W_0) \leq \mathcal{O}(\frac{1}{\alpha})$.

Next, we show that $X_{\mathcal{A}}$ and Z are close in operator norm. Indeed, $X_{\mathcal{A}} - Z = X_{\mathcal{A}}(I - W_0) = X_{\mathcal{A}}(I - W)(I - W_1)^{-1}$, hence:

$$\|X_{\mathcal{A}} - Z\|_2 = \|X_{\mathcal{A}}(I - W)(I - W_1)^{-1}\|_2 \quad (21)$$

$$\leq \|X_{\mathcal{A}}(I - W)\|_2 \|(I - W_1)^{-1}\|_2 \quad (22)$$

$$\stackrel{(i)}{\leq} 10 \|X_{\mathcal{A}}(I - W)\|_2 \quad (23)$$

$$\stackrel{(ii)}{\leq} 10\sqrt{2} \|X_{\mathcal{A}}(I - W) \text{diag}(c_{\mathcal{A}})^{1/2}\|_2 \stackrel{(iii)}{\leq} 20\sqrt{2}n\sigma. \quad (24)$$

Here (i) is because all singular values of W_1 are less than 0.9, (ii) is because $\text{diag}(c_{\mathcal{A}})^{1/2} \succeq \frac{1}{\sqrt{2}}I$, and (iii) is by the condition in the if statement (line 4 of Algorithm 1), since the sum on line 4 is equal to $\|X_{\mathcal{A}}(I - W) \text{diag}(c_{\mathcal{A}})^{1/2}\|_2^2$.

Combining the previous two observations, we have

$$\sum_{i \in S \cap \mathcal{A}} \|z_i - \mu\|_2^2 \leq (\text{rank}(Z) + 1) \|[z_i - \mu]_{i \in S \cap \mathcal{A}}\|_2^2 \quad (25)$$

$$\leq (\text{rank}(Z) + 1) (\|[z_i - x_i]_{i \in S \cap \mathcal{A}}\|_2 + \|[x_i - \mu]_{i \in S \cap \mathcal{A}}\|_2)^2 \quad (26)$$

$$\stackrel{(i)}{\leq} (\text{rank}(Z) + 1) \left(20\sqrt{2}n\sigma + \sqrt{\alpha n}\sigma\right)^2 = \mathcal{O}\left(\frac{\sigma^2}{\alpha}n\right). \quad (27)$$

Here (i) uses the preceding bound on $\|X_{\mathcal{A}} - Z\|_2$, together with the 2nd moment bound $\|[x_i - \mu]_{i \in S}\|_2 \leq \sqrt{\alpha n}\sigma$. Note that $\text{rank}(Z) \leq \text{rank}(W_0) = \mathcal{O}(\frac{1}{\alpha})$.

Since $|S \cap \mathcal{A}| = \Omega(\alpha n)$ by Lemma 17, the average value of $\|z_i - \mu\|_2^2$ over $S \cap \mathcal{A}$ is $\mathcal{O}(\frac{\sigma^2}{\alpha^2})$, and hence with probability at least $\frac{|S \cap \mathcal{A}|}{2|\mathcal{A}|} = \Omega(\alpha)$, a randomly chosen z_i will be within distance $\mathcal{O}(\frac{\sigma}{\alpha})$ of μ , which completes the first part of Proposition 16.

For the second part, when $\alpha = 1 - \epsilon \geq \frac{3}{4}$, recall that we have $\text{rank}(W_0) = 1$, and that $W_0 = (W - W_1)(I - W_1)^{-1}$. One can then verify that $\mathbb{1}^\top W_0 = \mathbb{1}^\top$. Therefore, $W_0 = u\mathbb{1}^\top$ for some u . Letting $\tilde{\mu} = X_{\mathcal{A}}u$, we have $\|X_{\mathcal{A}} - \tilde{\mu}\mathbb{1}^\top\|_2 \leq 20\sqrt{2n}\sigma$ by (24). In particular, $\mathcal{A}$ is resilient (around its mean) with $\sigma(\epsilon) \leq 20\sigma\sqrt{\frac{2\epsilon}{1-\epsilon}} \leq 40\sigma\sqrt{\epsilon}$ for $\epsilon \leq \frac{1}{2}$. Thus by the proof of Proposition 2 and the fact that $|\mathcal{A}| \geq |S \cap \mathcal{A}| \geq \frac{\alpha(2+\alpha)}{4-\alpha}n \geq (1 - \frac{5}{3}(1-\alpha))n$, the mean of $\mathcal{A}$ is within $\mathcal{O}(\sigma\sqrt{1-\alpha})$ of μ , as desired. ■

4.2. General Case

We are now ready to prove our general algorithmic result, Theorem 7 from Section 1.3. For convenience we recall Theorem 7 here:

Theorem *Suppose that $x_1, \dots, x_n$ contains a subset S of size $(1-\epsilon)n$ whose variance around its mean μ is bounded by σ^2 in the norm $\|\cdot\|$. Also suppose that Assumption 6 (κ -approximability) holds for the dual norm $\|\cdot\|_*$. Then, if $\epsilon \leq \frac{1}{4}$, there is a polynomial-time algorithm whose output satisfies $\|\hat{\mu} - \mu\| = \mathcal{O}(\sigma\sqrt{\kappa\epsilon})$.*

If, in addition, $\|\cdot\|$ is γ -strongly convex, then even if S only has size αn there is a polynomial-time algorithm such that $\|\hat{\mu} - \mu\| = \mathcal{O}(\frac{\sqrt{\kappa}\sigma}{\sqrt{\gamma\alpha}})$ with probability $\Omega(\alpha)$.

Recall that bounded variance means that $\frac{1}{|S|} \sum_{i \in S} \langle x_i - \mu, v \rangle^2 \leq \sigma^2 \|v\|_*^2$ for all $v \in \mathbb{R}^d$. There are two equivalent conditions to bounded variance that will be useful. The first is $\sup_{\|v\|_* \leq 1} v^\top \Sigma v \leq \sigma^2$ for all $v \in \mathbb{R}^d$, where $\Sigma = \frac{1}{|S|} \sum_{i \in S} (x_i - \mu)(x_i - \mu)^\top$; this is useful because Assumption 6 allows us to κ -approximate this supremum for any given Σ .

The second equivalent condition re-interprets σ in terms of a matrix norm. Let $\|\cdot\|_\psi$ denote the norm $\|\cdot\|$ above, and for a matrix M define the induced $2 \rightarrow \psi$ -norm $\|M\|_{2 \rightarrow \psi}$ as $\sup_{\|u\|_2 \leq 1} \|Mu\|_\psi$. Then the set S has variance at most σ^2 if and only if $\|[x_i - \mu]_{i \in S}\|_{2 \rightarrow \psi} \leq \sqrt{|S|}\sigma$. This will be useful because induced norms satisfy helpful compositional properties such as $\|AB\|_{2 \rightarrow \psi} \leq \|A\|_{2 \rightarrow \psi} \|B\|_2$.

The algorithm establishing Theorem 7 is almost identical to Algorithm 1, with two changes. The first change is that on line 4, the quantity $4n\sigma^2$ is replaced with $4\kappa n\sigma^2$, where κ is the approximation factor in Assumption 6. The second change is that in the optimization (15), the constraint $Y \succeq 0, \text{tr}(Y) \leq 1$ is replaced with $Y \in \mathcal{P}$, where $\mathcal{P}$ is the feasible set in Assumption 6. In other words, the only difference is that rather than finding the maximum eigenvalue, we κ -approximate the $2 \rightarrow \psi$ norm using Assumption 6. We therefore end up solving the saddle point problem

$$\max_{Y \in \mathcal{P}} \min_{\substack{0 \leq W_{ji} \leq \frac{4-\alpha}{\alpha(2+\alpha)n}, \\ \sum_j W_{ji} = 1}} \sum_{i \in \mathcal{A}} c_i (x_i - X_{\mathcal{A}} w_i)^\top Y (x_i - X_{\mathcal{A}} w_i). \quad (28)$$

Standard optimization algorithms such as Frank-Wolfe allow us to solve (28) to any given precision with a polynomial number of calls to the linear optimization oracle guaranteed by Assumption 6.

While Algorithm 1 essentially minimizes the quantity $\|X - XW\|_2^2$, this new algorithm can be thought of as minimizing $\|X - XW\|_{2 \rightarrow \psi}^2$. However, for general norms computing the $2 \rightarrow \psi$ norm is NP-hard, and so we rely on a κ -approximate solution by optimizing over $\mathcal{P}$. We are now ready to prove Theorem 7.

Proof (Theorem 7) The proof is similar to Proposition 16, so we only provide a sketch of the differences. First, the condition of Lemma 17 still holds, now with a equal to $\kappa n \sigma^2$ rather than $n \sigma^2$ due to the approximation ratio κ . (This is why we needed to change line 4.)

Next, we need to modify equations (21-24) to hold for the $2 \rightarrow \psi$ norm rather than operator norm:

$$\|X_{\mathcal{A}} - Z\|_{2 \rightarrow \psi} = \|X_{\mathcal{A}}(I - W)(I - W_1)^{-1}\|_{2 \rightarrow \psi} \quad (29)$$

$$\stackrel{(i)}{\leq} \|X_{\mathcal{A}}(I - W)\|_{2 \rightarrow \psi} \|(I - W_1)^{-1}\|_{2 \rightarrow 2} \quad (30)$$

$$\leq 10 \|X_{\mathcal{A}}(I - W)\|_{2 \rightarrow \psi} \quad (31)$$

$$\leq 10\sqrt{2} \|X_{\mathcal{A}}(I - W) \text{diag}(c_{\mathcal{A}})^{1/2}\|_{2 \rightarrow \psi} \leq 20\sqrt{2\kappa n} \sigma. \quad (32)$$

Here (i) is from the general fact $\|AB\|_{2 \rightarrow \psi} \leq \|A\|_{2 \rightarrow \psi} \|B\|_{2 \rightarrow 2}$, and the rest of the inequalities follow for the same reasons as in (21-24).

We next need to modify equations (25-27). This can be done with the following inequality:

Lemma 18 *For any matrix $A = [a_1 \cdots a_n]$ of rank r and any γ -strongly convex norm $\|\cdot\|_{\psi}$, we have $\sum_{i=1}^n \|a_i\|_{\psi}^2 \leq \frac{r}{\gamma} \|A\|_{2 \rightarrow \psi}^2$.*

This generalizes the inequality $\|A\|_F^2 \leq \text{rank}(A) \cdot \|A\|_2^2$. Using Lemma 18 (proved below), we have

$$\sum_{i \in S \cap \mathcal{A}} \|z_i - \mu\|_{\psi}^2 \leq \frac{\text{rank}(Z)+1}{\gamma} \|[z_i - \mu]_{i \in S \cap \mathcal{A}}\|_{2 \rightarrow \psi}^2 \quad (33)$$

$$\leq \frac{\text{rank}(Z)+1}{\gamma} (\|[z_i - x_i]_{i \in S \cap \mathcal{A}}\|_{2 \rightarrow \psi} + \|[x_i - \mu]_{i \in S \cap \mathcal{A}}\|_{2 \rightarrow \psi})^2 \quad (34)$$

$$= \mathcal{O}\left(\frac{\kappa \sigma^2 n}{\alpha \gamma}\right). \quad (35)$$

The inequalities again follow for the same reasons as before. If we choose z_i at random, with probability $\Omega(\alpha)$ we will output a z_i with $\|z_i - \mu\|_{\psi} = \mathcal{O}\left(\frac{\sigma \sqrt{\kappa}}{\alpha \sqrt{\gamma}}\right)$. This completes the first part of the proposition.

For the second part, by the same reasoning as before we obtain $\tilde{\mu}$ with $\|X_{\mathcal{A}} - \tilde{\mu} \mathbb{1}^T\|_{2 \rightarrow \psi} = \mathcal{O}(\sqrt{n\kappa}\sigma)$, which implies that $\mathcal{A}$ is resilient with $\sigma_*(\epsilon) = \mathcal{O}(\sigma \sqrt{\kappa\epsilon})$ for $\epsilon \leq \frac{1}{2}$. The mean of $\mathcal{A}$ will therefore be within distance $\mathcal{O}(\sigma \sqrt{\kappa\epsilon})$ of μ as before, which completes the proof. ■

We finish by proving Lemma 18.

Proof (Lemma 18) Let $s \in \{-1, +1\}^n$ be a uniformly random sign vector. We will compare $\mathbb{E}_s[\|As\|_{\psi}^2]$ in two directions. Let P be the projection onto the span of A . On the one hand, we have $\|As\|_{\psi}^2 = \|APs\|_{\psi}^2 \leq \|A\|_{2 \rightarrow \psi}^2 \|Ps\|_2^2$, and hence $\mathbb{E}_s[\|As\|_{\psi}^2] \leq \mathbb{E}_s[\|Ps\|_2^2] \|A\|_{2 \rightarrow \psi}^2 = \text{rank}(A) \|A\|_{2 \rightarrow \psi}^2$. On the other hand, similarly to (12) we have $\mathbb{E}_s[\|As\|_{\psi}^2] \geq (1/\gamma) \sum_{i=1}^n \|a_i\|_{\psi}^2$ by the strong convexity of the norm $\|\cdot\|_{\psi}$. Combining these yields the desired result. ■

5. Robust Low-Rank Recovery

In this section we present results on rank- k recovery. We first justify the definition of rank-resilience (Definition 8) by showing that it is information-theoretically sufficient for (approximately) recovering the best rank- k subspace. Then, we provide an algorithm showing that this subspace can be recovered efficiently.

5.1. Information-Theoretic Sufficiency

Let $X_S = [x_i]_{i \in S}$. Recall that δ -rank-resilience asks that $\text{col}(X_T) = \text{col}(X_S)$ and $\|X_T^\dagger X_S\|_2 \leq 2$ for $|T| \geq (1 - \delta)|S|$. This is justified by the following:

Proposition 19 *Let $S \subseteq [n]$ be a set of points of size $(1 - \delta)n$ that is $\frac{\delta}{1-\delta}$ -rank-resilient. Then it is possible to output a rank- k projection matrix P such that $\|(I - P)X_S\|_2 \leq 2\sigma_{k+1}(X_S)$.*

Proof Find the $\frac{\delta}{1-\delta}$ -rank-resilient set S' of size $(1 - \delta)n$ such that $\sigma_{k+1}(X_{S'})$ is smallest, and let P be the projection onto the top k singular vectors of $X_{S'}$. Then we have $\|(I - P)X_{S'}\|_2 = \sigma_{k+1}(X_{S'}) \leq \sigma_{k+1}(X_S)$. Moreover, if we let $T = S \cap S'$, we have $\|(I - P)X_T\|_2 \leq \|(I - P)X_{S'}\|_2 \leq \sigma_{k+1}(X_S)$ as well. By pigeonhole, $|T| \geq (1 - 2\delta)n = (1 - \frac{\delta}{1-\delta})|S|$. Therefore $\text{col}(X_T) = \text{col}(X_S)$, and $\|(I - P)X_S\|_2 = \|(I - P)X_T X_T^\dagger X_S\|_2 \leq \|(I - P)X_T\|_2 \|X_T^\dagger X_S\|_2 \leq 2\sigma_{k+1}(X_S)$ as claimed. ■

5.2. Efficient Recovery

We now address the question of efficient recovery, by proving Theorem 9, whose statement we recall here (with a few details added). For convenience let $\sigma \stackrel{\text{def}}{=} \sigma_{k+1}(X_S)/\sqrt{|S|}$.

Theorem *Let $\delta \leq \frac{1}{3}$. If a set of n points contains a subset S of size $(1 - \delta)n$ that is δ -rank-resilient, then there is an efficient algorithm (Algorithm 2) for recovering a matrix P of rank at most $15k$ such that $\|(I - P)X_S\|_2 = \mathcal{O}(\sigma\sqrt{|S|}) = \mathcal{O}(\sigma_{k+1}(X_S))$.*

Algorithm 2 is quite similar to Algorithm 1, but we include it for completeness. The proof is also similar to the proof of Proposition 16, but there are enough differences that we provide details below.

Algorithm 2 Algorithm for recovering a rank- k subspace.

- 1: Initialize $c_i = 1$ for all $i = 1, \dots, n$ and $\mathcal{A} = \{1, \dots, n\}$. Set $\lambda = \frac{(1-\delta)n\sigma^2}{k}$.
- 2: Let $Y \in \mathbb{R}^{d \times d}$ and $Q \in \mathbb{R}^{\mathcal{A} \times \mathcal{A}}$ be the maximizer/minimizer of the saddle point problem

$$\max_{\substack{Y \succeq 0, \\ \text{tr}(Y) \leq 1}} \min_{Q \in \mathbb{R}^{n \times n}} \sum_{i \in \mathcal{A}} c_i [(x_i - X_{\mathcal{A}} q_i)^\top Y (x_i - X_{\mathcal{A}} q_i) + \lambda \|q_i\|_2^2]. \quad (36)$$

- 3: Let $\tau_i^* = (x_i - X_{\mathcal{A}} q_i)^\top Y (x_i - X_{\mathcal{A}} q_i) + \lambda \|q_i\|_2^2$.
 - 4: **if** $\sum_{i \in \mathcal{A}} c_i \tau_i^* > 8n\sigma^2$ **then**
 - 5: For $i \in \mathcal{A}$, replace c_i with $(1 - \frac{\tau_i^*}{\tau_{\max}})c_i$, where $\tau_{\max} = \max_{i \in \mathcal{A}} \tau_i^*$.
 - 6: For all i with $c_i < \frac{1}{2}$, remove i from $\mathcal{A}$.
 - 7: Go back to line 3.
 - 8: **end if**
 - 9: Let Q_1 be the result of zeroing out all singular values of Q greater than 0.9.
 - 10: Output $P = X_{\mathcal{A}} Q_0 X_{\mathcal{A}}^\dagger$, where $Q_0 = (Q - Q_1)(I - Q_1)^{-1}$.
-

Proof (Theorem 9) First, we show that Lemma 17 holds with $a = 2n\sigma^2$. As before, the q_i can be optimized independently for each i , so $\sum_{i \in S \cap \mathcal{A}} c_i \tau_i^* \leq \sum_{i \in S \cap \mathcal{A}} c_i \tau_i(\hat{q}_i)$ for any matrix $\hat{Q}$. Take $\hat{Q}$ to be the projection onto the top k (right) singular vectors of $X_{S \cap \mathcal{A}}$. Then we have $\sum_{i \in S \cap \mathcal{A}} \lambda \|\hat{q}_i\|_2^2 \leq \lambda \|\hat{Q}\|_F^2 = \lambda k$. We also have $\sum_{i \in S \cap \mathcal{A}} c_i (x_i - X_{\mathcal{A}} \hat{q}_i)^\top Y (x_i - X_{\mathcal{A}} \hat{q}_i) \leq \sigma_{k+1}(X_{S \cap \mathcal{A}})^2 \leq \sigma^2 |S| \leq (1 - \delta)n\sigma^2$. Together these imply

that $\sum_{i \in S \cap \mathcal{A}} c_i \tau_i(\hat{q}_i) \leq (1 - \delta)n\sigma^2 + \lambda k = 2(1 - \delta)n\sigma^2$, so we can indeed take $a = 2n\sigma^2$. Consequently, $|S \cap \mathcal{A}| \geq \frac{(1-\delta)(3-\delta)}{3+\delta}n \geq (1 - \frac{5\delta}{3})n$.

Now, when we get to line 9, we have $\|X_{\mathcal{A}}(I - Q) \text{diag}(c_{\mathcal{A}})^{1/2}\|_2^2 + \lambda \|Q\|_F^2 \leq 8n\sigma^2$, and in particular each of the two terms individually is bounded by $8n\sigma^2$. Therefore, $\|Q\|_F^2 \leq \frac{8n\sigma^2}{\lambda} = \frac{8k}{1-\delta}$, and so Q_0 will have rank at most $\frac{8k}{0.81(1-\delta)} \leq 15k$. Moreover, $\|X_{\mathcal{A}}(I - Q)\|_2 \leq \sqrt{2}\|X_{\mathcal{A}}(I - Q) \text{diag}(c_{\mathcal{A}})^{1/2}\|_2 \leq 4\sqrt{n}\sigma$. We then have

$$\|(I - P)X_{\mathcal{A}}\|_2 = \|(I - X_{\mathcal{A}}Q_0X_{\mathcal{A}}^\dagger)X_{\mathcal{A}}\|_2 \quad (37)$$

$$= \|X_{\mathcal{A}}(I - Q_0)X_{\mathcal{A}}^\dagger X_{\mathcal{A}}\|_2 \quad (38)$$

$$\leq \|X_{\mathcal{A}}(I - Q_0)\|_2 \quad (39)$$

$$= \|X_{\mathcal{A}}(I - Q)(I - Q_1)^{-1}\|_2 \quad (40)$$

$$\leq 10\|X_{\mathcal{A}}(I - Q)\|_2 \leq 40\sqrt{n}\sigma = \mathcal{O}\left(\sigma\sqrt{|S|}\right). \quad (41)$$

By the same argument as Proposition 19, since $|S \cap \mathcal{A}| \geq \frac{1-5\delta/3}{1-\delta}|S| \geq (1 - \delta)|S|$, it follows that $\|(I - P)X_S\|_2 \leq 2\|(I - P)X_{S \cap \mathcal{A}}\|_2 = \mathcal{O}\left(\sigma\sqrt{|S|}\right)$ as well, which completes the proof. $\blacksquare$

6. Finite-Sample Concentration

In this section we provide two general finite-sample concentration results that establish resilience with high probability. The first holds for arbitrary resilient distributions but has suboptimal sample complexity, while the latter is specialized to distributions with bounded variance and has near-optimal sample complexity.

6.1. Concentration for Resilient Distributions

Our first result, stated as Proposition 4 in Section 1.2, applies to any (σ, ϵ) -resilient distribution p ; recall that p is (σ, ϵ) -resilient iff $\|\mathbb{E}[x \mid E] - \mu\| \leq \sigma$ for any event E of probability $1 - \epsilon$.

We define the *covering number* of the unit ball in a norm $\|\cdot\|_*$ to be the minimum M for which there are vectors $v_1, \dots, v_M$, each with $\|v_j\|_* \leq 1$, such that $\max_{j=1}^M \langle x, v_j \rangle \geq \frac{1}{2} \sup_{\|v\|_* \leq 1} \langle x, v \rangle$ for all vectors $x \in \mathbb{R}^d$. Note that $\log M$ is a measure of the effective dimension of the unit ball, i.e. $\log M = \Theta(d)$ if $\|\cdot\|_*$ is the ℓ_∞ or ℓ_2 norm, while $\log M = \Theta(\log d)$ for the ℓ_1 norm.

We recall Proposition 4 for convenience:

Proposition *Suppose that a distribution p is (σ, ϵ) -resilient around its mean μ with $\epsilon < \frac{1}{2}$. Let B be such that $\mathbb{P}[\|x - \mu\| \geq B] \leq \epsilon/2$. Also let M be the covering number of the unit ball in the dual norm $\|\cdot\|_*$.*

Then, given n samples $x_1, \dots, x_n \sim p$, with probability $1 - \delta - \exp(-\epsilon n/6)$ there is a subset T of $(1 - \epsilon)n$ of the x_i such that T is (σ', ϵ) -resilient with $\sigma' = \mathcal{O}\left(\sigma\left(1 + \sqrt{\frac{\log(M/\delta)}{\epsilon^2 n}} + \frac{(B/\sigma)\log(M/\delta)}{n}\right)\right)$.

Proof Let p' be the distribution of samples from p conditioned on $\|x - \mu\| \leq B$. Note that p' is $(\sigma, \frac{\epsilon}{2})$ -resilient since every event with probability $1 - \epsilon/2$ in p' is an event of probability $(1 - \epsilon/2)^2 \geq 1 - \epsilon$ in p . Moreover, with probability $1 - \exp(-\epsilon n/6)$, at least $(1 - \epsilon)n$ of the samples from p will come from p' . Therefore, we can focus on establishing resilience of the $n' = (1 - \epsilon)n$ samples from p' .

With a slight abuse of notation, let $x_1, \dots, x_{n'}$ be the samples from p' . Then to check resilience we need to bound $\|\frac{1}{|T|} \sum_{i \in T} (x_i - \mu)\|$ for all sets T of size at least $(1 - \epsilon)n'$. We will first use the covering $v_1, \dots, v_M$ to obtain

$$\left\| \frac{1}{|T|} \sum_{i \in T} (x_i - \mu) \right\| \leq 2 \max_{j=1}^M \frac{1}{|T|} \sum_{i \in T} \langle x_i - \mu, v_j \rangle. \quad (42)$$

The idea will be to analyze the sum over $\langle x_i - \mu, v_j \rangle$ for a fixed v_j and then union bound over the M possibilities. For a fixed v_j , we will split the sum into two components: those with small magnitude (roughly σ/ϵ) and those with large magnitude (between σ/ϵ and B). We can then bound the former with Hoeffding's inequality, and using resilience we will be able to upper-bound the second moment of the latter, after which we can use Bernstein's inequality.

More formally, let $\tau = \frac{1-\epsilon}{\epsilon/4} \sigma$ and define

$$y_i = \langle x_i - \mu, v_j \rangle \mathbb{I}[|\langle x_i - \mu, v_j \rangle| < \tau], \quad (43)$$

$$z_i = \langle x_i - \mu, v_j \rangle \mathbb{I}[|\langle x_i - \mu, v_j \rangle| \geq \tau]. \quad (44)$$

Clearly $y_i + z_i = \langle x_i - \mu, v_j \rangle$. Also, we have $|y_i| \leq \tau$ almost surely, and $|z_i| \leq B$ almost surely (because $x_i \sim p'$ and hence $\langle x_i - \mu, v_j \rangle \leq \|x_i - \mu\| \leq B$). The threshold τ is chosen so that z_i is non-zero with probability at most $\epsilon/2$ under p (see Lemma 3).

Now, for any set T of size at least $(1 - \epsilon)n'$, we have

$$\frac{1}{|T|} \sum_{i \in T} \langle x_i - \mu, v_j \rangle = \frac{1}{|T|} \sum_{i \in T} y_i + z_i \quad (45)$$

$$\leq \left| \frac{1}{|T|} \sum_{i \in T} y_i \right| + \frac{1}{|T|} \sum_{i \in T} |z_i| \quad (46)$$

$$\leq \left| \frac{1}{|T|} \sum_{i=1}^{n'} y_i \right| + \left| \frac{1}{|T|} \sum_{i \notin T} y_i \right| + \frac{1}{|T|} \sum_{i=1}^{n'} |z_i| \quad (47)$$

$$\leq \frac{1}{1-\epsilon} \left| \frac{1}{n'} \sum_{i=1}^{n'} y_i \right| + \frac{\epsilon}{1-\epsilon} \tau + \frac{1}{(1-\epsilon)n'} \sum_{i=1}^{n'} |z_i|. \quad (48)$$

The last step uses the fact that $|y_i| \leq \tau$ for all i . It thus suffices to bound $|\frac{1}{n'} \sum_{i=1}^{n'} y_i|$ as well as $\frac{1}{n'} \sum_{i=1}^{n'} |z_i|$.

For the y_i term, note that by resilience $\|\mathbb{E}[y_i]\| \leq \sigma$ (since y_i is sampled from p conditioned on $|\langle x_i - \mu, v_j \rangle| < \tau$ and $\|x_i - \mu\| \leq B$, which each occur with probability at least $1 - \epsilon/2$). Then by Hoeffding's inequality, $|\frac{1}{n'} \sum_{i=1}^{n'} y_i| = \mathcal{O}(\sigma + \tau \sqrt{\log(2/\delta)/n'})$ with probability $1 - \delta$.

For the z_i term, we note that $\mathbb{E}[|z_i|] = \mathbb{E}[\max(z_i, 0)] + \mathbb{E}[\max(-z_i, 0)]$. Let τ' be the ϵ -quantile of $\langle x_i - \mu, v_j \rangle$ under p , which by Lemma 3 is at most τ . Then we have

$$\mathbb{E}_p[\max(z_i, 0)] = \mathbb{E}_p[\langle x_i - \mu, v_j \rangle \mathbb{I}[\langle x_i - \mu, v_j \rangle \geq \tau]] \quad (49)$$

$$\leq \mathbb{E}_p[\langle x_i - \mu, v_j \rangle \mathbb{I}[\langle x_i - \mu, v_j \rangle \geq \tau']] \quad (50)$$

$$\stackrel{(i)}{\leq} \epsilon \cdot \frac{1-\epsilon}{\epsilon} \sigma = (1-\epsilon)\sigma, \quad (51)$$

where (i) is again Lemma 3. Then we have $\mathbb{E}_{p'}[\max(z_i, 0)] \leq \frac{1}{1-\epsilon} \mathbb{E}_p[\max(z_i, 0)] \leq \sigma$, and hence $\mathbb{E}_{p'}[|z_i|] \leq 2\sigma$ (as $\mathbb{E}[\max(-z_i, 0)] \leq \sigma$ by the same argument as above). Since $|z_i| \leq B$, we then have $\mathbb{E}[|z_i|^2] \leq 2B\sigma$.

Therefore, by Bernstein's inequality, with probability $1 - \delta$ we have

$$\frac{1}{n'} \sum_{i=1}^{n'} |z_i| \leq \mathcal{O}\left(\sigma + \sqrt{\frac{\sigma B \log(2/\delta)}{n'}} + \frac{B \log(2/\delta)}{n'}\right) = \mathcal{O}\left(\sigma + \frac{B \log(2/\delta)}{n'}\right). \quad (52)$$

Taking a union bound over the v_j for both y and z , and plugging back into (48), we get that $\frac{1}{|T|} \sum_{i \in T} \langle x_i - \mu, v_j \rangle \leq \mathcal{O}\left(\sigma + \frac{\sigma}{\epsilon} \sqrt{\frac{\log(2M/\delta)}{n}} + \frac{B \log(2M/\delta)}{n}\right)$ for all T and v_j with probability $1 - \delta$. Plugging back into (42), we get that $\left\| \frac{1}{|T|} \sum_{i \in T} (x_i - \mu) \right\| \leq \mathcal{O}\left(\sigma + \frac{\sigma}{\epsilon} \sqrt{\frac{\log(2M/\delta)}{n}} + \frac{B \log(2M/\delta)}{n}\right)$, as was to be shown. $\blacksquare$

6.2. Concentration Under Bounded Covariance

In this section we state a stronger but more restrictive finite-sample bound giving conditions under which samples have bounded variance. It is a straightforward extension of Proposition B.1 of Charikar et al. (2017), so we defer the proof to Section H.

Proposition 20 *Suppose that a distribution p has bounded variance in a norm $\|\cdot\|_*$: $\mathbb{E}_{x \sim p}[\langle x - \mu, v \rangle^2] \leq \sigma^2 \|v\|_*^2$ for all $v \in \mathbb{R}^d$. Then, given n samples $x_1, \dots, x_n \sim p$, with probability $1 - \exp(-\epsilon^2 n/16)$ there is a subset T of $(1 - \epsilon)n$ of the points such that*

$$\frac{1}{|T|} \sum_{i \in T} \langle x_i - \mu, v \rangle^2 \leq (\sigma')^2 \|v\|_*^2 \text{ for all } v \in \mathbb{R}^d, \quad (53)$$

where $(\sigma')^2 = \frac{4\sigma^2}{\epsilon} \left(1 + \frac{d}{(1-\epsilon)n}\right)$.

Proposition 20 says that whenever a distribution p on $\mathbb{R}^d$ has bounded variance, if $n \geq d$ samples x_i are drawn from p then some large subset of the samples will have bounded variance as well.

7. Information-Theoretic Applications

Using Proposition 2 together with our finite-sample bounds, we establish information-theoretic robust recovery results in several cases of interest.

7.1. Distribution Learning

First we consider the case where we are given k -tuples of independent samples from a distribution π on $\{1, \dots, m\}$. By taking the empirical distribution of the k samples, we get an element of the simplex Δ_m ; for instance if $m = 5$ and $k = 3$ and a given triple of samples is $(2, 4, 2)$, then we produce the vector $(0, \frac{2}{3}, 0, \frac{1}{3}, 0) \in \Delta_5$. For general π and k , we will denote the resulting distribution over Δ_m as $F(\pi, k)$, and think of both π and samples $x \sim F(\pi, k)$ as elements of $\mathbb{R}^m$.

We first establish finite-sample concentration for $F(\pi, k)$:

Proposition 21 *Let $p = F(\pi, k)$, where π is a distribution on $\{1, \dots, m\}$. Then:*

- *Given m samples $x_1, \dots, x_m$ from p , with probability $1 - \exp(-\Omega(m))$ there is a subset T of $\frac{3m}{4}$ of the samples such that $\frac{1}{|T|} \sum_{i \in T} \langle x_i - \pi, v \rangle^2 \leq \mathcal{O}\left(\frac{1}{k}\right) \|v\|_\infty^2$ for all $v \in \mathbb{R}^d$.*

- Given $\frac{m\sqrt{k}}{\epsilon} + \frac{m}{\epsilon^2}$ samples from p , with probability $1 - \exp(-\Omega(m))$ there is a subset of $(1 - \epsilon)m$ of the samples that is (σ, ϵ) -resilient in the ℓ_1 -norm, with $\sigma = \mathcal{O}(\epsilon\sqrt{\log(1/\epsilon)/k})$.

As a corollary, we immediately get robust recovery bounds for π . For instance, we have:

Corollary 22 Suppose $\epsilon > 0$ is sufficiently small. Then, given $n = \Omega(\frac{m\sqrt{k}}{\epsilon} + \frac{m}{\epsilon^2})$ samples $x_1, \dots, x_n$, $(1 - \epsilon)n$ of which are drawn from π^k , with probability $1 - \exp(-\Omega(m))$ it is information-theoretically possible to recover π with total variational error $\mathcal{O}(\epsilon\sqrt{\log(1/\epsilon)/k})$.

We omit the proof of Corollary 22, as it follows straightforwardly from Proposition 21 and Proposition 2.

Proof (Proposition 21) Consider a sample $y \sim \pi$, interpreted as an indicator vector in $\{0, 1\}^m$. First, we observe that for any vector $v \in \{-1, +1\}^m$, $\mathbb{E}[\langle y, v \rangle^2] = \sum_{j=1}^m \pi_j v_j^2 = 1$, and so certainly $\mathbb{E}[\langle y - \pi, v \rangle^2] \leq 1$. Therefore, if x is an average of k independent samples from π , $\mathbb{E}[\langle x - \pi, v \rangle^2] \leq 1/k$ for all v with $\|v\|_\infty \leq 1$. Invoking Proposition 20 then yields the first part of Proposition 21.

We next analyze the moment generating function $\mathbb{E}[\exp(\lambda^\top (y - \pi))]$ in order to get a tail bound on $y - \pi$. Provided $\|\lambda\|_\infty \leq 1$, the moment generating function is bounded as

$$\mathbb{E}[\exp(\lambda^\top (y - \pi))] = \exp(-\lambda^\top \pi) \sum_{j=1}^m \pi_j \exp(\lambda_j) \quad (54)$$

$$\leq \exp(-\lambda^\top \pi) \sum_{j=1}^m \pi_j (1 + \lambda_j + \lambda_j^2) \quad (55)$$

$$\leq \exp(-\lambda^\top \pi) \exp\left(\sum_{j=1}^m \pi_j (\lambda_j + \lambda_j^2)\right) = \exp\left(\sum_{j=1}^m \pi_j \lambda_j^2\right). \quad (56)$$

In particular, for any sign vector v and $c \in [0, 1]$ we have $\mathbb{E}[\exp(cv^\top (y - \pi))] \leq \exp(c^2 \sum_{j=1}^m \pi_j) = \exp(c^2)$. For an average x of k independent samples from π , we therefore have $\mathbb{E}[\exp(cv^\top (x - \pi))] \leq \exp(c^2/k)$ for $c \in [0, k]$. But note that $v^\top (x - \pi) \leq 2$, and so $\mathbb{E}[\exp(cv^\top (x - \pi))] \leq \exp(2c^2/k)$ for all $c \geq k$ as well. Hence $\mathbb{E}[\exp(cv^\top (x - \pi))] \leq \exp(2c^2/k)$ for all $c \geq 0$.

Now, let E be any event of probability ϵ . We have

$$\mathbb{E}[v^\top (x - \pi) \mid E] \leq \frac{1}{c} \log(\mathbb{E}[\exp(cv^\top (x - \pi)) \mid E]) \quad (57)$$

$$\leq \frac{1}{c} \log\left(\frac{1}{\epsilon} \mathbb{E}[\exp(cv^\top (x - \pi))]\right) \quad (58)$$

$$\leq \frac{1}{c} \log\left(\frac{1}{\epsilon} \exp(2c^2/k)\right) \quad (59)$$

$$= \frac{\log(1/\epsilon) + 2c^2/k}{c}. \quad (60)$$

Optimizing c yields $\mathbb{E}[v^\top (x - \pi) \mid E] \leq \sqrt{8 \log(1/\epsilon)/k}$, and so (by Lemma 10) p is (σ, ϵ) -resilient around its mean in the ℓ_1 -norm, with $\sigma = \frac{\epsilon}{1-\epsilon} \sqrt{8 \log(1/\epsilon)/k} = \mathcal{O}(\epsilon\sqrt{\log(1/\epsilon)/k})$.

The second part of Proposition 21 now follows immediately upon invoking Proposition 4, with $M = 2^d$ (for the 2^d sign vectors), $B = 2$ and $\sigma = \mathcal{O}(\epsilon\sqrt{\log(1/\epsilon)/k})$. $\blacksquare$

7.2. Bounded k th Moments

We next give results for distributions with bounded k th moments.

Proposition 23 *Suppose that a distribution p on $\mathbb{R}^d$ has bounded k th moments: $\mathbb{E}[|\langle x - \mu, v \rangle|^k]^{1/k} \leq \sigma_k \|v\|_2$ for all $v \in \mathbb{R}^d$ and some $k \geq 2$. Then, given $n \geq \frac{d^{1.5}}{\epsilon} + \frac{d}{\epsilon^2}$ samples $x_1, \dots, x_n \sim p$, with probability $1 - \exp(-\Omega(d))$ there is a set T of $(1 - \epsilon)n$ of the x_i that is $(\mathcal{O}(\sigma_k \epsilon^{1-1/k}), \epsilon)$ -resilient in the ℓ_2 -norm.*

Proof We will invoke Proposition 4. Let U be the uniform distribution on the unit sphere. First note that

$$\mathbb{E}_x[\|x - \mu\|_2^k] = \mathbb{E}_x[(d \mathbb{E}_{v \sim U}[|\langle x - \mu, v \rangle|^2])^{k/2}] \quad (61)$$

$$\leq d^{k/2} \mathbb{E}_{x,v}[|\langle x - \mu, v \rangle|^k] \quad (62)$$

$$\leq d^{k/2} \sigma_k^k. \quad (63)$$

Therefore, by Chebyshev's inequality, $\mathbb{P}[\|x - \mu\|_2 \geq (\epsilon/2)^{-1/k} \sigma_k \sqrt{d}] \leq \epsilon/2$. We can therefore take $B = \mathcal{O}(\epsilon^{-1/k} \sigma_k \sqrt{d})$ in Proposition 4. Similarly, we can take $\sigma = \mathcal{O}(\epsilon^{-1/k} \sigma_k)$ in Proposition 4, and there is a covering of the ℓ_2 ball with 9^d elements, so $\log(M) = \mathcal{O}(d)$. Together, these imply that with probability $1 - \exp(-d)$ there is a set T of size $(1 - \epsilon)n$ that is $(\mathcal{O}(\sigma_k \epsilon^{1-1/k}), \epsilon)$ -resilient, as was to be shown. ■

As before, we obtain the following immediate corollary:

Corollary 24 *Let p be a distribution on $\mathbb{R}^d$ with k th moments bounded by σ_k , for $k \geq 2$, and suppose that $\epsilon > 0$ is sufficiently small. Then, given $n = \Omega(\frac{d^{1.5}}{\epsilon} + \frac{d}{\epsilon^2})$ samples $x_1, \dots, x_n$, $(1 - \epsilon)n$ of which are drawn from p , with probability $1 - \exp(-\Omega(d))$ it is possible to recover $\hat{\mu}$ within ℓ_2 -distance $\mathcal{O}(\epsilon^{1-1/k} \sigma_k)$ of the mean of p .*

7.3. Stochastic Block Model

Our final set of information-theoretic results is for the semi-random stochastic block model from Charikar et al. (2017). In this model, we consider a graph G on n vertices, with an unknown set S of αn “good” vertices. For simplicity we assume the graph is a directed graph. For $i, j \in S$, i is connected to j with probability $\frac{a}{n}$, and for $i \in S$, $j \notin S$, i is connected to j with probability $\frac{b}{n}$, where $b < a$. For $i \notin S$ the edges are allowed to be arbitrary. If we let $A \in \{0, 1\}^{n \times n}$ be the adjacency matrix of G , then the rows in S are jointly independent samples from a distribution p on $\{0, 1\}^n$. The mean of this distribution is $\frac{a}{n}$ for the coordinates lying in S , and $\frac{b}{n}$ for the coordinates outside of S . We will apply our robust mean estimation results to this distribution, and use this to robustly recover the set S itself. We let $x_1, \dots, x_{\alpha n}$ denote the αn samples from p corresponding to the elements of S .

Lemma 25 *Suppose that $x_1, \dots, x_{\alpha n}$ are drawn from a stochastic block model with parameters (α, a, b) . Take the norm $\|x\| = \max_{|J|=\alpha n} \|x_J\|_1$, which is the maximum ℓ_1 -norm over any αn coordinates of x . Define $\mu_j = \frac{a}{n}$ if $j \in S$ and $\frac{b}{n}$ if $j \notin S$. Then, with probability $1 - \exp(-\Omega(\alpha n))$, the x_i are $(\sigma, \alpha/2)$ -resilient under $\|\cdot\|$ around μ with parameter*

$$\sigma = \mathcal{O}\left(\alpha \sqrt{a \log(2/\alpha)} + \log(2/\alpha)\right). \quad (64)$$

Proof Note that we can express $\|x\|$ as $\max_{v \in \mathcal{V}} \langle x, v \rangle$, where $\mathcal{V}$ is the set of αn -sparse $\{0, +1, -1\}$ vectors; in particular, $|\mathcal{V}| = \binom{n}{\alpha n} 2^{\alpha n}$. By Lemma 10 and the definition of resilience, σ is equal to

$$\frac{\alpha/2}{1 - \alpha/2} \max_{T \subseteq \{1, \dots, \alpha n\}, |T| = \frac{1}{2} \alpha^2 n} \max_{v \in \mathcal{V}} \left\langle \frac{1}{|T|} \sum_{i \in T} x_i - \mu, v \right\rangle. \quad (65)$$

We will union bound over all T and v . For a fixed T and v , the inner expression is equal to $\frac{2}{\alpha^2 n} \sum_{i \in T} \sum_{j=1}^n v_j (x_{ij} - \mu_j)$. Note that all of the $v_j (x_{ij} - \mu_j)$ are independent, zero-mean random variables with variance at most $\frac{a}{n} v_j^2$ and are bounded in $[-1, 1]$. By Bernstein's inequality, we have

$$\mathbb{P} \left[\sum_{i \in T} \sum_{j=1}^n v_j (x_{ij} - \mu_j) \geq t \right] \leq \exp \left(-\frac{1}{2} \frac{t^2}{\sum_{i,j} \frac{a}{n} v_j^2 + \frac{1}{3} t} \right) \quad (66)$$

$$= \exp \left(-\frac{t^2}{\alpha^3 a n + \frac{2}{3} t} \right), \quad (67)$$

Now, if we want our overall union bound to hold with probability $1 - \delta$, we need to set the term in (67) to be at most $\delta / \left[\binom{\alpha n}{\alpha^2 n/2} \binom{n}{\alpha n} 2^{\alpha n} \right]$, so $\frac{t^2}{\alpha^3 a n + \frac{2}{3} t} = \log(1/\delta) + \mathcal{O}(\alpha n \log(2/\alpha)) = \mathcal{O}(\alpha n \log(2/\alpha))$ (since we will take $\delta = \exp(-\Theta(\alpha n))$). Hence we can take $t = \mathcal{O}(\sqrt{\alpha^4 a n^2 \log(2/\alpha)} + \alpha n \log(2/\alpha))$. Dividing through by $\frac{1}{2} \alpha^2 n$ and multiplying by $\frac{\alpha/2}{1-\alpha/2}$, we get

$$\sigma = \mathcal{O} \left(\alpha \sqrt{a \log(2/\alpha)} + \log(2/\alpha) \right), \quad (68)$$

as was to be shown. $\blacksquare$

Using Lemma 25, we obtain our result on robust recovery of the set S :

Corollary 26 *Under the semi-random stochastic block model with parameters α , a , and b , it is information-theoretically possible to obtain a set $\hat{S}$ satisfying*

$$\frac{1}{\alpha n} |\hat{S} \triangle S| = \mathcal{O} \left(\sqrt{\frac{a \log(2/\alpha)}{\alpha^2 (a-b)^2}} \right). \quad (69)$$

with probability $\Omega(\alpha)$.

Proof By Lemma 25 and Proposition 2, we can recover a $\hat{\mu}$ that with probability $\Omega(\alpha)$ satisfies

$$\|\hat{\mu} - \mu\| = \mathcal{O} \left(\sqrt{a \log(2/\alpha)} + \frac{\log(2/\alpha)}{\alpha} \right). \quad (70)$$

Note that $\mu_j = \frac{a}{n}$ if $j \in S$ and $\frac{b}{n}$ if $j \notin S$, where $b < a$. We will accordingly define $\hat{S}$ to be the set of coordinates j such that $\hat{\mu}_j \geq \frac{a+b}{2n}$. We then have that $\|\hat{\mu} - \mu\| = \Omega(\frac{a-b}{n} \min(|\hat{S} \triangle S|, \alpha n))$, and hence $\frac{1}{\alpha n} |\hat{S} \triangle S| = \mathcal{O}(\frac{1}{\alpha(a-b)} \|\hat{\mu} - \mu\|)$ whenever the right-hand-side is at most 1. Using (70), we have that

$$\frac{1}{\alpha n} |\hat{S} \triangle S| = \mathcal{O} \left(\frac{1}{\alpha(a-b)} \left(\sqrt{a \log(2/\alpha)} + \frac{\log(2/\alpha)}{\alpha} \right) \right) \quad (71)$$

$$= \mathcal{O} \left(\sqrt{\frac{a \log(2/\alpha)}{\alpha^2 (a-b)^2}} + \frac{\log(2/\alpha)}{\alpha^2 (a-b)} \right) \leq \mathcal{O} \left(\sqrt{\frac{a \log(2/\alpha)}{\alpha^2 (a-b)^2}} + \frac{a \log(2/\alpha)}{\alpha^2 (a-b)^2} \right) \quad (72)$$

Here the last step simply multiplies the second term by $\frac{a}{a-b}$, which is at least 1. Note that the first term in (72) dominates whenever the bound is meaningful, which yields the desired result. $\blacksquare$

8. Algorithmic Applications

We end by presenting applications of our algorithmic result Theorem 7. First, we show that Theorem 7 implies a deterministic recovery result for sets that are resilient in the ℓ_p -norm, for any $p \in (1, 2]$:

Corollary 27 *Suppose that a set of points $x_1, \dots, x_n$ contains a set S of size αn that is $(\sigma, \frac{1}{2})$ -resilient around its mean μ in ℓ_p -norm, with $p \in (1, 2]$. Then there is an efficient algorithm whose output $\hat{\mu}$ satisfies $\|\mu - \hat{\mu}\|_p = \mathcal{O}\left(\frac{\sigma}{\alpha^{p-1}}\right)$ with probability $\Omega(\alpha)$.*

Moreover, if $\alpha = 1 - \epsilon \geq 0.87$ and S is $(\sigma\sqrt{\delta}, \delta)$ -resilient for $\delta \in [\frac{\epsilon}{2}, \frac{1}{2}]$, the same algorithm has $\|\mu - \hat{\mu}\|_p = \tilde{\mathcal{O}}\left(\sigma\sqrt{\frac{\epsilon}{p-1}}\right)$ with probability 1. In particular, this holds if S has variance bounded by σ^2 .

Proof First, we observe that the ℓ_p -norm is $(p-1)$ -strongly convex for $p \in (1, 2]$ (c.f. Lemma 17 of Shalev-Shwartz (2007)). Therefore, if S has the assumed level of resilience, then we can find a core S_0 of size $\frac{\alpha}{2}$ with second moments bounded by $\mathcal{O}(\frac{\sigma^2}{p-1})$ by Proposition 12. Moreover, if $\alpha = 1 - \epsilon \geq 0.87$, then we can find a core of size $(1 - \epsilon)^2 n \geq \frac{3}{4}n$ by Proposition 15. In this latter case, the core will have second moments bounded by $\mathcal{O}(\frac{\sigma^2 \log(1/\epsilon)}{p-1})$.

In each case, we can apply Theorem 7 to this bounded-variance core. We need to check that Assumption 6 holds. This is due to the following result, which follows from Theorem 3 of Nesterov (1998):

Theorem 28 (Nesterov) *Let $\|\cdot\|_q$ denote the ℓ_q -norm for $q \geq 2$. For a matrix $A \in \mathbb{R}^{d \times d}$, let $f(A) = \max_{\|v\|_q \leq 1} v^\top A v$, and let $g(A) = \max_{Y \succeq 0, \|\text{diag}(Y)\|_{q/2} \leq 1} \langle A, Y \rangle$. Then $f(A) \leq g(A) \leq \frac{\pi}{2} f(A)$.*

In particular, for the ℓ_p -norm with $p \in [1, 2]$, the dual norm is the ℓ_q -norm with $q = \frac{p}{p-1} \in [2, \infty]$. Therefore, Theorem 28 asserts that Assumption 6 holds for $\mathcal{P} = \{Y \mid Y \succeq 0, \|\text{diag}(Y)\|_{q/2} \leq 1\}$ and $\kappa = \frac{\pi}{2} = \mathcal{O}(1)$.

Now, applying Theorem 7 for general α , this yields a recovery guarantee of $\|\hat{\mu} - \mu\|_p = \mathcal{O}(\frac{\sigma}{\alpha^{p-1}})$ with probability $\Omega(\alpha)$, while for $\alpha = 1 - \epsilon \geq 0.87$ this yields a recovery guarantee of $\|\hat{\mu} - \mu\|_p = \mathcal{O}(\frac{\sigma}{\sqrt{p-1}} \sqrt{\epsilon \log(1/\epsilon)})$ with probability 1, as claimed. ■

We next specialize Corollary 27 to the distribution learning case and provide finite-sample bounds.

Corollary 29 *For the distribution learning problem from Section 7.1, suppose that we are given n samples, of which αn are independent k -tuples from a distribution π on $\{1, \dots, m\}$, and the remaining samples are arbitrary. Then:*

- *If $\alpha n \geq m$, with probability $1 - \exp(-\Omega(m))$ there is a randomized algorithm outputting a $\hat{\pi}$ satisfying $\|\hat{\pi} - \pi\|_{TV} = \tilde{\mathcal{O}}(\frac{1}{\alpha\sqrt{k}})$ with probability $\Omega(\alpha)$.*
- *If $\alpha = 1 - \epsilon$, where ϵ is sufficiently small, and $n \geq \frac{m\sqrt{k}}{\epsilon} + \frac{m}{\epsilon^2}$, then with probability $1 - \exp(-\Omega(m))$ there is a deterministic algorithm outputting a $\hat{\pi}$ satisfying $\|\hat{\pi} - \pi\|_{TV} = \tilde{\mathcal{O}}(\sqrt{\epsilon/k})$.*

In both of the above bounds, the $\tilde{\mathcal{O}}$ notation hides polylog factors in both ϵ and m .

Proof By Proposition 21, in the first case there will with high probability be a set of size $\frac{3}{4}\alpha n$ with bounded variance under the ℓ_1 -norm. Since the ℓ_1 -norm and $\ell_{1+\frac{1}{\log(m)}}$ -norm are within constant factors of each other, the set will also have bounded variance in the $\ell_{1+\frac{1}{\log(m)}}$ -norm, which is $\Omega(\frac{1}{\log(m)})$ -strongly convex. We can then invoke Corollary 27 to obtain error $\mathcal{O}(\frac{\sigma \log(m)}{\alpha})$.

In the second case, Proposition 21 says that we can remove $\epsilon|S|/2$ points from S to make the remaining set $(\mathcal{O}(\epsilon\sqrt{\log(1/\epsilon)/k}), \epsilon/2)$ -resilient. This implies that it is also $(\mathcal{O}(\delta\sqrt{\log(1/\epsilon)/k}), \delta)$ -resilient for all $\delta \in [\frac{\epsilon}{2}, \frac{1}{2}]$, and so by Proposition 15 it has a core of size $(1 - \epsilon)(1 - \epsilon/2)|S| = (1 - \mathcal{O}(\epsilon))n$ with variance at most $\frac{\log(m)\log(1/\epsilon)}{k}$ (we again need to pass from ℓ_1 to $\ell_{1+\log(m)}$ to apply Proposition 15, which is why we incur the $\log(m)$ factor).

Applying Theorem 7 as in the second part of Corollary 27 then yields the desired error bound of $\mathcal{O}(\sqrt{\epsilon \log(1/\epsilon) \log(m)/k}) = \tilde{\mathcal{O}}(\sqrt{\epsilon/k})$. ■

Appendix A. Counterexample to Analogues of Proposition 12

In this section we show:

Proposition 30 *Let $S = \{e_1, \dots, e_n\}$, where the e_i are the standard basis in $\mathbb{R}^n$. Then:*

1. *S is $(2n^{1/p-1}, \frac{1}{2})$ -resilient around 0 in ℓ_p -norm, for all $p \in [1, \infty]$.*
2. *Any subset T of $\frac{n}{2}$ elements of S has $\max_{\|v\|_q \leq 1} \frac{1}{|T|} \sum_{i \in T} |\langle e_i, v \rangle|^k = (n/2)^{\max(-1, k(\frac{1}{p}-1))}$.*

Here we let $\|\cdot\|_q$ denote the dual norm to $\|\cdot\|_p$, i.e. q is such that $\frac{1}{p} + \frac{1}{q} = 1$.

Proposition 30 implies in particular that for large n , an analog of Proposition 12 can only hold if $k(1/p - 1) \geq -1$, which implies that $k \leq \frac{p}{p-1}$. For $k = 2$, this implies $p \leq 2$ (so the norm is strongly convex), and for $p = 2$ this implies $k \leq 2$. We do not know whether an analog of Proposition 12 holds for e.g. $k = 3$, $p = 1.5$.

Proof (Proposition 30) For the first part, note that the mean of any subset of $\frac{n}{2}$ elements of S will have all coordinates lying in $[0, \frac{2}{n}]$. Therefore, the ℓ_p -norm of the mean is at most $\frac{2}{n} \cdot n^{1/p} = 2n^{1/p-1}$, as claimed.

For the second part, assume without loss of generality that $T = \{e_1, \dots, e_{n/2}\}$. The optimization then reduces to $\max_{\|v\|_q \leq 1} \frac{2}{n} \|v\|_k^k$, where $v \in \mathbb{R}^{n/2}$. For the optimal v , it is clear that $\|v\|_k = 1$ if $k \geq q$, and $\|v\|_k = (n/2)^{\frac{1}{k} - \frac{1}{q}}$ if $k < q$, whence $\frac{2}{n} \|v\|_k^k = \frac{2}{n} \cdot (n/2)^{\max(0, 1 - \frac{k}{q})} = (n/2)^{\max(-1, -\frac{k}{q})} = (n/2)^{\max(-1, -k(1 - \frac{1}{p}))}$, as claimed. ■

Appendix B. Basic Properties of Resilience

Here we prove Lemmas 3 and 10. Recall that the first states that resilience around the mean μ is equivalent to bounded tails, while the latter relates ϵ -resilience to $(1 - \epsilon)$ -resilience.

Proof (Lemma 10) Note that for a distribution p , resilience around μ can be computed as $\sigma_*(\epsilon) = \max\{\|\mathbb{E}[X | E] - \mu\| \mid E \text{ has probability } 1 - \epsilon\}$. When μ is the mean, then $\mu = \mathbb{E}[X]$. We can then write

$$= \|\mathbb{E}[X | E] - \mathbb{E}[X]\| \tag{73}$$

$$= \|\mathbb{E}[X | E] - \mathbb{P}[E]\mathbb{E}[X | E] - \mathbb{P}[\neg E]\mathbb{E}[X | \neg E]\| \tag{74}$$

$$= (1 - \mathbb{P}[E])\|\mathbb{E}[X | E] - \mathbb{E}[X | \neg E]\|. \tag{75}$$

From (75), it is clear that $\sigma_*(1 - \epsilon) = \frac{1-\epsilon}{\epsilon}\sigma_*(\epsilon)$, as (75) is invariant under replacing E with $\neg E$, except that $1 - \mathbb{P}[E]$ becomes $\mathbb{P}[E]$.

More generally, if μ is not the mean then we still have $\|\mu - \mathbb{E}[X]\| = \sigma_*(0) \leq \sigma_*(\epsilon)$ by resilience. Then if E is an event with probability ϵ , we have

$$\|\mathbb{E}[X | E] - \mu\| \leq \|\mathbb{E}[X | E] - \mathbb{E}[X]\| + \|\mathbb{E}[X] - \mu\| \quad (76)$$

$$\leq \frac{1-\epsilon}{\epsilon} \|\mathbb{E}[X | \neg E] - \mathbb{E}[X]\| + \sigma_*(\epsilon) \quad (77)$$

$$\leq \frac{1-\epsilon}{\epsilon} (\|\mathbb{E}[X | \neg E] - \mu\| + \|\mu - \mathbb{E}[X]\|) + \sigma_*(\epsilon) \quad (78)$$

$$\leq \frac{1-\epsilon}{\epsilon} \cdot 2\sigma_*(\epsilon) + \sigma_*(\epsilon) = \frac{2-\epsilon}{\epsilon} \sigma_*(\epsilon). \quad (79)$$

Therefore, even when μ is not the mean we have $\sigma_*(1-\epsilon) \leq \frac{2-\epsilon}{\epsilon} \sigma_*(\epsilon)$, which completes the proof. $\blacksquare$

Proof (Lemma 3) We note that

$$\sigma_*(\epsilon) \stackrel{(i)}{=} \frac{1-\epsilon}{\epsilon} \sigma_*(1-\epsilon) \quad (80)$$

$$\stackrel{(ii)}{=} \frac{1-\epsilon}{\epsilon} \max_E \{\|\mathbb{E}[X - \mu | E]\| \mid \mathbb{P}[E] \geq \epsilon\} \quad (81)$$

$$\stackrel{(iii)}{=} \frac{1-\epsilon}{\epsilon} \max_{E,v} \{\langle \mathbb{E}[X - \mu | E], v \rangle \mid \mathbb{P}[E] \geq \epsilon, \|v\|_* \leq 1\} \quad (82)$$

$$\stackrel{(iv)}{=} \frac{1-\epsilon}{\epsilon} \max_{E,v} \{\mathbb{E}[\langle X - \mu, v \rangle \mid E] \mid \mathbb{P}[E] \geq \epsilon, \|v\|_* \leq 1\} \quad (83)$$

$$\stackrel{(v)}{=} \frac{1-\epsilon}{\epsilon} \max_v \{\mathbb{E}[\langle X - \mu, v \rangle \mid \langle X - \mu, v \rangle \geq \tau_\epsilon(v)] \mid \|v\|_* \leq 1\}, \quad (84)$$

as was to be shown. Here (i) is Lemma 10, (ii) is the definition of resilience, and (iii) is the definition of the dual norm. Then (iv) uses linearity of expectation, while (v) explicitly maximizes $\mathbb{E}[\langle X - \mu, v \rangle \mid E]$ over events E with probability at least ϵ . $\blacksquare$

Appendix C. Proof of Lemma 11

For a fixed v , let $S_+ = \{i \mid \langle x_i - \mu, v \rangle > 0\}$, and define S_- similarly. Either S_+ or S_- must have size at least $\frac{1}{2}|S|$, without loss of generality assume it is S_+ . Then we have

$$\frac{1}{|S|} \sum_{i \in S} |\langle x_i - \mu, v \rangle| = \frac{1}{|S|} \left(2 \sum_{i \in S_+} \langle x_i - \mu, v \rangle - \sum_{i \in S} \langle x_i - \mu, v \rangle \right) \quad (85)$$

$$\leq \frac{1}{|S|} \left(2 \sum_{i \in S_+} (x_i - \mu) \|v\|_* + \left\| \sum_{i \in S} (x_i - \mu) \right\| \|v\|_* \right) \quad (86)$$

$$\leq \left(2 \frac{|S_+|}{|S|} + 1 \right) \sigma \|v\|_* \leq 3\sigma \|v\|_*, \quad (87)$$

which completes the first part. In the other direction, for any set T , we have

$$\left\| \frac{1}{|T|} \sum_{i \in T} (x_i - \mu) \right\| = \max_{\|v\|_* = 1} \frac{1}{|T|} \sum_{i \in T} \langle x_i - \mu, v \rangle \quad (88)$$

$$\leq \max_{\|v\|_* = 1} \frac{1}{|T|} \sum_{i \in S} |\langle x_i - \mu, v \rangle| \leq \frac{|S|}{|T|} \sigma \leq 2\sigma, \quad (89)$$

as was to be shown. ■

Appendix D. Proof of Lemma 13

We start by taking a continuous relaxation of (6), asking for weights $c_i \in [0, 1]$ rather than $\{0, 1\}$:

$$\min_{c \in [0, 1]^n, \|c\|_1 \geq \frac{3n}{4}} \max_{\|v\|_* \leq 1} \frac{1}{n} \sum_{i=1}^n c_i |\langle x_i, v \rangle|^2. \quad (90)$$

Note that we strengthened the inequality to $\|c\|_1 \geq \frac{3n}{4}$, whereas in (6) it was $\|c\|_1 \geq \frac{n}{2}$. Given any solution $c_{1:n}$ to (90), we can obtain a solution c' to (6) by letting $c'_i = \mathbb{I}[c_i \geq \frac{1}{2}]$. Then $c'_i \in \{0, 1\}$ and $\|c'\|_1 \geq \frac{n}{2}$. Moreover, $c'_i \leq 2c_i$, so $\frac{1}{n} \sum_{i=1}^n c'_i |\langle x_i, v \rangle|^2 \leq \frac{2}{n} \sum_{i=1}^n c_i |\langle x_i, v \rangle|^2$ for all v . Therefore, the value of (6) is at most twice the value of (90).

Now, by the minimax theorem, we can swap the min and max in (90) in exchange for replacing the single vector v with a distribution over vectors v_j , thus obtaining that (90) is equal to

$$\lim_{m \rightarrow \infty} \max_{\substack{\alpha_1 + \dots + \alpha_m \leq 1 \\ \alpha \geq 0, \|v_j\|_* \leq 1}} \min_{\substack{c \in [0, 1]^n \\ \|c\|_1 \geq \frac{3n}{4}}} \frac{1}{n} \sum_{i=1}^n c_i \sum_{j=1}^m \alpha_j |\langle x_i, v_j \rangle|^2. \quad (91)$$

By letting $v'_j = \alpha_j v_j$, the above is equivalent to optimizing over v_j satisfying $\sum_j \|v_j\|_*^2 \leq 1$:

$$\lim_{m \rightarrow \infty} \max_{\|v_1\|_*^2 + \dots + \|v_m\|_*^2 \leq 1} \min_{\substack{c \in [0, 1]^n \\ \|c\|_1 \geq \frac{3n}{4}}} \frac{1}{n} \sum_{i=1}^n c_i \sum_{j=1}^m |\langle x_i, v_j \rangle|^2. \quad (92)$$

For any $v_1, \dots, v_m$, we will find c such that the above sum is bounded. Indeed, define $B(v_{1:m})$ to be $\frac{1}{n} \sum_{i=1}^n \sqrt{\sum_{j=1}^m |\langle x_i, v_j \rangle|^2}$. Then take $c_i = \mathbb{I}[\sum_{j=1}^m |\langle x_i, v_j \rangle|^2 < 16B^2]$, which has $\|c\|_1 \geq \frac{3n}{4}$ by Markov's inequality, and for which $\sum_i c_i \sum_j |\langle x_i, v_j \rangle|^2 \leq 4B(v_{1:m})^2$.

Therefore, the value of (90) is bounded by $\max_{m, v_{1:m}} 4B(v_{1:m})^2$, and so the value of (6) is bounded by $\max_{m, v_{1:m}} 8B(v_{1:m})^2$, which yields the desired result. ■

Appendix E. Proof of Proposition 15

The proof mirrors that of Proposition 12, but with more careful bookkeeping. Instead of (90), we consider the minimax problem

$$\min_{\substack{c \in [0, 1]^n \\ \|c\|_1 \geq (1-\epsilon/2)n}} \max_{\|v\|_* \leq 1} \frac{1}{n} \sum_{i=1}^n c_i |\langle x_i, v \rangle|^2. \quad (93)$$

The only difference is that $\frac{3}{4}n$ has been replaced with $(1 - \epsilon/2)n$ in the constraint on $\|c\|_1$. We then end up needing to bound

$$\max_{\sum_j \|v_j\|_*^2 \leq 1} \frac{1}{n} \sum_{k=1}^{(1-\frac{\epsilon}{2})n} \sum_{j=1}^m |\langle x_{i_k}, v_j \rangle|^2. \quad (94)$$

Here we suppose that the indices i are arranged in order $i_1, i_2, \dots, i_n$ such that the output of the inner sum is monotonically increasing in i_k (in other words, the outer sum over k is over the $(1 - \frac{\epsilon}{2})n$ smallest values of the inner sum, which corresponds to the optimal choice of c). Now we have

$$\frac{1}{n} \sum_{k=1}^{(1-\frac{\epsilon}{2})n} \sum_{j=1}^m |\langle x_{i_k}, v_j \rangle|^2 \stackrel{(i)}{\leq} \frac{1}{n} \sum_{k=1}^{(1-\frac{\epsilon}{2})n} \left(\frac{1}{n-k+1} \sum_{l=k}^n \sqrt{\sum_{j=1}^m |\langle x_{i_l}, v_j \rangle|^2} \right)^2 \quad (95)$$

$$\stackrel{(ii)}{\leq} \frac{2}{n} \sum_{k=1}^{(1-\frac{\epsilon}{2})n} \left(\frac{1}{n-k+1} \mathbb{E}_s \left[\sum_{l=k}^n \left| \langle x_{i_l}, \sum_{j=1}^m s_j v_j \rangle \right| \right] \right)^2 \quad (96)$$

$$\stackrel{(iii)}{\leq} \frac{2}{n} \sum_{k=1}^{(1-\frac{\epsilon}{2})n} \left(\frac{3n \min(\sigma_*(\frac{n-k+1}{n}), \sigma_*(\frac{1}{2}))}{n-k+1} \mathbb{E}_s \left[\left\| \sum_{j=1}^m s_j v_j \right\|_* \right] \right)^2 \quad (97)$$

$$\leq \frac{18}{n\gamma} \sum_{k=1}^{(1-\frac{\epsilon}{2})n} \left(\frac{\min(\sigma_*(\frac{n-k+1}{n}), \sigma_*(\frac{1}{2}))}{\frac{n-k+1}{n}} \right)^2. \quad (98)$$

Here (i) bounds the k th smallest term by the average of the $n - k + 1$ largest terms, (ii) is Khinchine's inequality, and (iii) is the following lemma which we prove later:

Lemma 31 *If S is resilient around μ and $T \subseteq S$ has size at most $\epsilon|S|$, then $\sum_{i \in T} |\langle x_i - \mu, v \rangle| \leq 3|S| \|v\|_* \cdot \min(\sigma_*(\epsilon), \sigma_*(\frac{1}{2}))$.*

Continuing, we can bound (98) by

$$\frac{18}{\gamma} \int_0^{1-\frac{\epsilon}{2}} \left(\frac{\min(\sigma_*(1-u), \sigma_*(\frac{1}{2}))}{1-u-1/n} \right)^2 du \leq \frac{18}{\gamma} \left(\frac{1}{\frac{\epsilon}{2} - \frac{1}{n}} \right)^2 \int_{\epsilon/2}^1 \left(\frac{\min(\sigma_*(u), \sigma_*(\frac{1}{2}))}{u} \right)^2 du \quad (99)$$

$$\leq \mathcal{O}(1) \cdot \frac{1}{\gamma} \cdot \left(\int_{\epsilon/2}^{1/2} u^{-2} \sigma_*(u)^2 du + 2\sigma_*(\frac{1}{2})^2 \right) \quad (100)$$

$$= \mathcal{O} \left(\frac{1}{\gamma} \cdot (\tilde{\sigma}_*(\epsilon)^2 + \sigma_*(\frac{1}{2})^2) \right). \quad (101)$$

Since the $\tilde{\sigma}_*(\epsilon)$ term dominates, this completes the bound on (93), from which the remainder of the proof follows identically to Proposition 12. $\blacksquare$

We end this section by proving Lemma 31.

Proof (Lemma 31) Note that the $\sigma_*(\frac{1}{2})$ part of the bound was already established in Lemma 11, so it suffices to show the $\sigma_*(\epsilon)$ part. Let $\hat{\mu}$ be the mean of S . Note that we must have $\|\mu - \hat{\mu}\| \leq \sigma_*(\epsilon)$ by definition. We can thus replace μ with $\hat{\mu}$ in the statement of Lemma 31 at the cost of changing the left-hand-side by at most

$|T|\sigma_*(\epsilon)\|v\|_*$. Now let T_+ be the elements of T with $\langle x_i - \hat{\mu}, v \rangle > 0$ and define T_- similarly. We have

$$\sum_{i \in T} |\langle x_i - \mu, v \rangle| = \sum_{i \in T_+} \langle x_i - \mu, v \rangle - \sum_{i \in T_-} \langle x_i - \mu, v \rangle \quad (102)$$

$$= - \sum_{i \in S \setminus T_+} \langle x_i - \mu, v \rangle + \sum_{i \in S \setminus T_-} \langle x_i - \mu, v \rangle \quad (103)$$

$$\leq \left\| \sum_{i \in S \setminus T_+} (x_i - \mu) \right\| \|v\|_* + \left\| \sum_{i \in S \setminus T_-} (x_i - \mu) \right\| \|v\|_* \quad (104)$$

$$\leq 2|S|\sigma_*(\epsilon)\|v\|_*, \quad (105)$$

which yields the desired result. $\blacksquare$

Appendix F. Fast Algorithm for ℓ_2 Mean Estimation

Recall that in Section 4.1 we wanted to solve the problem

$$\min_W \|X - XW\|_2 \text{ subject to } 0 \leq W_{ij} \leq \frac{1}{\alpha n}, \mathbb{1}^\top W = \mathbb{1}^\top. \quad (106)$$

An examination of the proof of Proposition 16 reveals that it actually suffices to replace the constraint $0 \leq W_{ij} \leq \frac{1}{\alpha n}$ with the Frobenius norm constraint $\|W\|_F^2 \leq \frac{1}{\alpha}$. We will show how to solve this modified problem efficiently (via a singular value decomposition and linear-time post-processing). For $r = \frac{1}{\alpha}$, we want to solve

$$\min_W \|X - XW\|_2 \text{ subject to } \mathbb{1}^\top W = \mathbb{1}^\top, \|W\|_F^2 \leq r. \quad (107)$$

Let $P = \frac{1}{n} \mathbb{1} \mathbb{1}^\top$. Then given any W , we can pass to the matrix $W' = P + (I - P)W$, which will always satisfy $\mathbb{1}^\top W' = \mathbb{1}^\top$. Therefore, it suffices to minimize

$$\|X - XW'\|_2^2 = \|X - X(P + (I - P)W)\|_2^2 = \|X(I - P) - X(I - P)W\|_2^2 \quad (108)$$

with a constraint on

$$\|W'\|_F^2 = \|P + (I - P)W\|_F^2 = \|P\|_F^2 + \|(I - P)W\|_F^2. \quad (109)$$

Now, at the optimum we have $W = (I - P)W$, and for all W we have $\|(I - P)W\|_F^2 \leq \|W\|_F^2$, so we can replace $\|(I - P)W\|_F^2$ with $\|W\|_F^2$ without changing the optimization. In addition, P is a constant and $\|P\|_F^2 = 1$, so the constraint boils down to $\|W\|_F^2 \leq r - 1$. Letting $X' = X(I - P)$, it therefore suffices to solve

$$\min_W \|X' - X'W\|_2^2 \text{ subject to } \|W\|_F^2 \leq r - 1. \quad (110)$$

This problem is unitarily invariant and so can be solved efficiently via an SVD. In particular let $X' = U\Lambda V^\top$. The optimal W lies in the span of $V^\top$, so after an appropriate change of variables it suffices to minimize $\|\Lambda - \Lambda W\|_2^2$ subject to $\|W\|_F^2 \leq r - 1$. We can also check that the optimal W is diagonal in this basis, so we need only solve the vector minimization

$$\min_w \max_j \lambda_j^2 (1 - w_j)^2 \text{ subject to } \sum_j w_j^2 \leq r - 1, \quad (111)$$

which can be done in linear time.

Appendix G. Proof of Lemma 17

Assuming that (i) and (ii) hold prior to line 5 of the algorithm, we need to show that they continue to hold.

By assumption, we have $\sum_{i \in S \cap \mathcal{A}} c_i \tau_i \leq \alpha a$, while $\sum_{i \in \mathcal{A}} c_i \tau_i \geq 4a$. But note that the amount that $\sum_{i \in S} c_i$ decreases in line 5 is proportional to $\sum_{i \in S \cap \mathcal{A}} c_i \tau_i$, while the amount that $\sum_{i=1}^n c_i$ decreases is proportional to $\sum_{i \in \mathcal{A}} c_i \tau_i$. Therefore, the former decreases at most $\frac{\alpha}{4}$ times as fast as the latter, meaning that (i) is preserved.

Since (i) is preserved, we have $\sum_{i \in S} (1 - c_i) \leq \frac{\alpha}{4} (\sum_{i \in S} (1 - c_i) + \sum_{i \notin S} (1 - c_i))$. Re-arranging yields $\sum_{i \in S} (1 - c_i) \leq \frac{\alpha}{4-\alpha} \sum_{i \notin S} (1 - c_i) \leq \frac{\alpha(1-\alpha)}{4-\alpha} n$. In particular, $\#\{i \mid c_i \leq \frac{1}{2}\} \leq \frac{2\alpha(1-\alpha)}{4-\alpha} n$, so at most $\frac{2\alpha(1-\alpha)}{4-\alpha} n$ elements of S have been removed from $\mathcal{A}$ in total. This implies $|S \cap \mathcal{A}| \geq \frac{\alpha(2+\alpha)}{4-\alpha} n$, and so (ii) is preserved as well. ■

Appendix H. Proof of Proposition 20

This is a corollary of Proposition B.1 of Charikar et al. (2017), which states that given a distribution with covariance Σ_0 , where $\Sigma_0 \preceq \sigma_0^2 I$, with probability $1 - \exp(-\epsilon^2 n/16)$ it is possible to find a set T of $(1 - \epsilon)n$ points such that $\frac{1}{|T|} \sum_{i \in T} (x_i - \mu)(x_i - \mu)^\top \preceq (\sigma'_0)^2 I$, where σ'_0 and σ_0 have the same relation as σ' and σ above. Applying this proposition directly to p will not give us our result, so we instead apply it to a transformed version of p .

In particular, let Σ be the covariance of p , and given a sample $x_i \sim p$, let $y_i = \Sigma^{-1/2}(x_i - \mu)$.³ Then the y_i have identity covariance and mean 0, and so Proposition B.1 of Charikar et al. (2017) implies that we can find $(1 - \epsilon)n$ of the y_i such that $\frac{1}{|T|} \sum_{i \in T} y_i y_i^\top \preceq \frac{4}{\epsilon} \left(1 + \frac{d}{(1-\epsilon)n}\right) I$. In terms of the x_i , after multiplying by $\Sigma^{1/2}$ on the left and right, we obtain

$$\frac{1}{|T|} \sum_{i \in T} (x_i - \mu)(x_i - \mu)^\top \preceq \frac{4}{\epsilon} \left(1 + \frac{d}{(1-\epsilon)n}\right) \Sigma, \quad (112)$$

which implies that

$$\frac{1}{|T|} \sum_{i \in T} \langle x_i - \mu, v \rangle^2 \leq \frac{4}{\epsilon} \left(1 + \frac{d}{(1-\epsilon)n}\right) v^\top \Sigma v \quad (113)$$

$$= \frac{4}{\epsilon} \left(1 + \frac{d}{(1-\epsilon)n}\right) \mathbb{E}_{x \sim p} [\langle x - \mu, v \rangle^2] \quad (114)$$

$$\leq \frac{4}{\epsilon} \left(1 + \frac{d}{(1-\epsilon)n}\right) \sigma^2 \|v\|_*^2 = (\sigma')^2 \|v\|_*^2, \quad (115)$$

as was to be shown.

References

S. Balakrishnan, S. S. Du, J. Li, and A. Singh. Computationally efficient robust sparse estimation in high dimensions. In *Conference on Learning Theory (COLT)*, pages 169–212, 2017.

3. In the case that Σ is not invertible, we need to add a small multiple of the identity matrix before inverting, but we omit this part of the argument for brevity.

- J. Batson, D. A. Spielman, and N. Srivastava. Twice-Ramanujan sparsifiers. *SIAM Journal on Computing*, 41(6):1704–1721, 2012.
- M. Charikar, J. Steinhardt, and G. Valiant. Learning from untrusted data. In *Symposium on Theory of Computing (STOC)*, 2017.
- A. Decelle, F. Krzakala, C. Moore, and L. Zdeborová. Asymptotic analysis of the stochastic block model for modular networks and its algorithmic applications. *Physical Review E*, 84(6), 2011.
- I. Diakonikolas, G. Kamath, D. Kane, J. Li, A. Moitra, and A. Stewart. Robust estimators in high dimensions without the computational intractability. In *Foundations of Computer Science (FOCS)*, 2016a.
- I. Diakonikolas, D. Kane, and A. Stewart. Robust learning of fixed-structure Bayesian networks. *arXiv*, 2016b.
- I. Diakonikolas, D. M. Kane, and A. Stewart. Statistical query lower bounds for robust estimation of high-dimensional Gaussians and Gaussian mixtures. *arXiv*, 2016c.
- I. Diakonikolas, G. Kamath, D. Kane, J. Li, A. Moitra, and A. Stewart. Being robust (in high dimensions) can be practical. *arXiv*, 2017a.
- I. Diakonikolas, G. Kamath, D. M. Kane, J. Li, A. Moitra, and A. Stewart. Robustly learning a Gaussian: Getting optimal error, efficiently. *arXiv*, 2017b.
- I. Diakonikolas, D. M. Kane, and A. Stewart. Learning geometric concepts with nasty noise. *arXiv*, 2017c.
- O. Guédon and R. Vershynin. Community detection in sparse networks via Grothendieck’s inequality. *arXiv*, 2014.
- U. Haagerup. The best constants in the khintchine inequality. *Studia Mathematica*, 70(3):231–283, 1981.
- D. Kane, S. Karmalkar, and E. Price. Robust polynomial regression up to the information theoretic limit. *arXiv*, 2017.
- A. Khintchine. Über dyadische brüche. *Mathematische Zeitschrift*, 18:109–116, 1923.
- A. R. Klivans, P. M. Long, and R. A. Servedio. Learning halfspaces with malicious noise. *Journal of Machine Learning Research (JMLR)*, 10:2715–2740, 2009.
- P. Kothari and J. Steinhardt. Better agnostic clustering via tensor norms. *arXiv*, 2017.
- K. A. Lai, A. B. Rao, and S. Vempala. Agnostic estimation of mean and covariance. In *Foundations of Computer Science (FOCS)*, 2016.
- C. M. Le, E. Levina, and R. Vershynin. Concentration and regularization of random graphs. *arXiv*, 2015.
- J. Li. Robust sparse estimation tasks in high dimensions. *arXiv*, 2017.
- L. Massoulié. Community detection thresholds and the weak Ramanujan property. In *Symposium on Theory of Computing (STOC)*, pages 694–703, 2014.
- M. Meister and G. Valiant. A data prism: Semi-verified learning in the small-alpha regime. *arXiv*, 2017.

- E. Mossel, J. Neeman, and A. Sly. A proof of the block model threshold conjecture. *arXiv*, 2013.
- Y. Nesterov. Semidefinite relaxation and nonconvex quadratic optimization. *Optimization methods and software*, 9:141–160, 1998.
- E. Rebrova and K. Tikhomirov. Coverings of random ellipsoids, and invertibility of matrices with iid heavy-tailed entries. *arXiv*, 2015.
- E. Rebrova and R. Vershynin. Norms of random matrices: local and global problems. *arXiv*, 2016.
- S. Shalev-Shwartz. *Online Learning: Theory, Algorithms, and Applications*. PhD thesis, The Hebrew University of Jerusalem, 2007.
- J. Steinhardt. Does robustness imply tractability? A lower bound for planted clique in the semi-random model. *arXiv*, 2017.
- J. Steinhardt, G. Valiant, and M. Charikar. Avoiding imposters and delinquents: Adversarial crowdsourcing and peer prediction. In *Advances in Neural Information Processing Systems (NIPS)*, 2016.
- H. Xu, C. Caramanis, and S. Mannor. Principal component analysis with contaminated data: The high dimensional case. *arXiv*, 2010.