

Radiative decay widths of ground and excited states of vector charmonium and bottomonium

Hluf Negash^a and Shashank Bhatnagar^b

July 21, 2022

^aDepartment of Physics, Samara University, Samara, Ethiopia

^bDepartment of Physics, University Institute of Sciences, Chandigarh University, Mohali, India

Abstract

In this work we study the radiative decay widths of vector quarkonia for the process of $J/\psi(nS) \rightarrow \eta_c(nS)\gamma$ and $\Upsilon(nS) \rightarrow \eta_b(nS)\gamma$ (for principal quantum numbers $n = 1, 2, 3$) in the framework of Bethe-Salpeter equation under the covariant instantaneous ansatz using a 4×4 form of BSE. The parameters of the framework were determined by a fit to the mass spectrum of ground states of pseudoscalar and vector quarkonia, such as; η_c , η_b , J/ψ and Υ . These input parameters so fixed were found to give good agreements with data on mass spectra of ground and excited states of pseudoscalar and vector quarkonia, leptonic decay constants of pseudoscalar and vector quarkonia, two photon decays and two gluon decays of pseudoscalar quarkonia in our recent paper. With these input parameters so fixed, in the present work, the radiative decay widths of ground (1S) and excited (2S, 3S) states of heavy vector quarkonia (J/Ψ and Υ) are calculated and found to be in reasonable agreement with data and other models.

1 Introduction

Studies on mass spectra and decays of heavy quarkonia ($c\bar{c}$, and $b\bar{b}$) have become a hot topic in recent years, due to observation of many new states at various high energy accelerators at BABAR,

Belle, CLEO and BES-III collaborations [1, 2, 3, 28]. All this has opened up new challenges in theoretical understanding of heavy hadrons and provide an important tool for exploring the structure of these simplest bound states in QCD and for studying the non-perturbative (long distance) behavior of strong interactions.

As regards the dynamical framework, to investigate these properties is concerned, many non-perturbative approaches, such as Lattice QCD [5], Chiral perturbation theory [6], heavy quark effective theory [7], QCD sum rules [8], N.R.QCD [9], dynamical-equation based approaches like Schwinger-Dyson equation and Bethe-Salpeter equation (BSE) [10, 11, 12, 13, 14, 15], and potential models [16] have been proposed to deal with the long distance property of QCD.

Bethe-Salpeter equation (BSE) [11, 15, 17, 18, 19] is a conventional approach in dealing with relativistic bound state problems. From the solutions, we can obtain useful information about the inner structure of hadrons, which is also crucial in treating hadronic decays. The BSE framework which is firmly rooted in field theory, provides a realistic description for analyzing hadrons as composite objects. Despite its drawback of having to input model-dependent kernel, these studies have become an interesting topic in recent years, since calculations have shown that BSE framework using phenomenological potentials can give satisfactory results as more and more data are being accumulated. The BSE is frequently adopted as starting point of QCD inspired models, due to the fact that this equation has a firm base in quantum field theory.

In a recent work [20, 21], we employed a 4×4 representation for two-body ($q\bar{q}$) BS amplitude for calculating both the mass spectra as well as the transition amplitudes for various processes. However, the price one to pay in this approach is to solve a coupled set of Salpeter equations for both pseudoscalar and vector quarkonia. However, in [21], we explicitly showed that these coupled Salpeter equations can indeed get decoupled in the heavy-quark approximation, leading to mass spectral equations with analytical solutions for both masses, as well as eigenfunctions for all the ground and excited states of pseudoscalar and vector $c\bar{c}$ and $b\bar{b}$ systems in an approximate harmonic oscillator basis. These analytical forms of eigen functions for ground and excited states so obtained were used to evaluate the transition amplitudes for different processes in [21]. Thus in [21], we had calculated the mass spectrum, weak decay constants, two photon decay widths and two gluon decay widths of ground (1S) and radially excited (2S, 3S,...) states of pseudoscalar

charmonium and bottomonium such as η_c and η_b , as well as the mass spectrum and leptonic decay constants of ground state (1S), excited (2S, 1D, 3S, 2D, 4S and 3D) states of vector charmonium and bottomonium such as J/ψ and Υ , using this formulation of 4×4 Bethe-Salpeter equation under covariant Instantaneous Ansatz (CIA). Our results were in good agreement with data (where ever available) and other models. However, in all the above processes, the quark anti-quark loop involved a single hadron-quark vertex, which was simple to handle.

However for the transitions such as $V \rightarrow P + \gamma$, the process requires calculation of triangle quark-loop diagram involving two hadron-quark vertices and is difficult to evaluate in BSE-CIA [11]. However, the problems involving quark-loop diagrams with two or more hadron-quark vertices can be handled effectively in 4×4 representation of BSE [22, 23]. Thus, in the present work, we will precisely apply an instantaneous formalism employing 4×4 BSE under CIA for transitions involving the process, $V \rightarrow P\gamma$. This is an extension of our earlier work in [21], where we were involved in calculations of processes that involve one quark loop with a single hadron-quark vertex such as leptonic decays of P and V quarkonia, and the two-photon and two-gluon decays of P-quarkonia.

This paper is organized as follows. In Sec. **2**, we will formulate the hadronic process $V \rightarrow P + \gamma$ in the framework of 4×4 BSE under CIA and calculate its radiative decay width. The numerical results for radiative decay widths of the processes are worked out. Section **3** is reserved for discussion and conclusions.

2 Electromagnetic transition of $V \rightarrow P + \gamma$

The lowest order, Feynman diagrams for the process, $V \rightarrow P\gamma$, are given in Fig.1, where $V(1^{--})$ and $P(0^{-+})$ are vector and pseudoscalar quarkonia respectively. The second diagram is obtained from the first one by reversing the directions of internal fermionic lines. The electromagnetic transition amplitude M_{fi} , corresponding to Fig.1a, and 1b is given by the one-loop momentum integral as in Ref. [23, 24, 25]:

$$M_{fi} = -ie_q \int \frac{d^4q}{(2\pi)^4} \text{Tr}[\bar{\Psi}_P(P', q') \not{\epsilon} \Psi_V(P, q) S_F^{-1}(-p_2) + \bar{\Psi}_P(P', q') S_F^{-1}(p_1) \Psi_V(P, q) \not{\epsilon}] \quad (1)$$




Figure 1: Electromagnetic transitions for the process $V \rightarrow P\gamma$ via quark triangle-loop diagrams.

where Ψ_P and Ψ_V are the 4D BS wave functions of pseudoscalar and vector quarkonia involved in the process, and are expressed as, $\Psi_V(P, q) = S_F(p_1)\Gamma(\hat{q})S_F(-p_2)$ and $\bar{\Psi}_P(P', q') = S_F(-p'_2)\Gamma(\hat{q}')S_F(p'_1)$. e_q is the electric charge of quarks. ϵ_μ is the polarization vector of the emitted photon. We have used the momentum relations: $P = p_1 + p_2$, $P' = p'_1 + p'_2$, $k = P - P'$, and $p_2 = p'_2$. However for second diagram, the last momentum relation is $p_1 = p'_1$, while the other relations are the same as in first diagram. Here, the second term (coming from Fig.1b) in Eq. (1) gives the same contribution as the first term (coming from Fig.1a), and hence we can write M_{fi} as:

$$M_{fi} = -2ie_q \int \frac{d^4q}{(2\pi)^4} Tr[\bar{\Psi}_P(P', q')\not{\epsilon}\Psi_V(P, q)S_F^{-1}(-p_2)] \quad (2)$$

with an overall multiplicative factor of 2.

Thus, one can write the instantaneous Bethe-Salpeter form for the electromagnetic transition amplitude between two bound states as (see Ref. [26]):

$$M_{fi} = -\frac{2e_q}{M} \int \frac{d^3\hat{q}}{(2\pi)^3} Tr\{\not{P}\bar{\psi}_P^{++}(\hat{q}')\not{\epsilon}\psi_V^{++}(\hat{q}) - \not{P}\bar{\psi}_P^{--}(\hat{q}')\not{\epsilon}\psi_V^{--}(\hat{q})\} \quad (3)$$

where, we resolve the amplitude corresponding to the first term (i.e. Fig.1a) in the previous equation into $++$, and $--$ components, where the dominant contribution comes from the $++$ components, while the contribution from the $--$ components is less than 1 percent [27]. Thus we can to a good approximation, write, the total amplitude for the process $V \rightarrow P\gamma$ in terms of only the $++$ components as:

$$M_{fi} = -\frac{2e_q}{M} \int \frac{d^3\hat{q}}{(2\pi)^3} Tr\{\not{P}\bar{\psi}_P^{++}(\hat{q}')\not{\epsilon}\psi_V^{++}(\hat{q})\} \quad (4)$$

where, $\hat{q}' = \hat{q} + \frac{1}{2}\hat{P}'$ and M is the mass of the initial quarkonia. The relativistic BS wave function of $P(0^{-+})$ meson bound state is taken as (see Ref. [20, 21]):

$$\psi_P(\hat{q}') = N'_P[M' + \not{P}' + \frac{\not{q}'\not{P}'}{m}]\gamma_5\phi_P(\hat{q}') \quad (5)$$

where P' is the momentum, N'_P is the BS normalization of the final state P-quarkonia and M' is the mass of the final state P-quarkonia.

The relativistic BS wave function of $V(1^{--})$ bound state can ultimately be written as (see Ref. [21]):

$$\psi_V(\hat{q}) = N_V[M\not{\epsilon} + \hat{q}\cdot\epsilon\frac{M}{m} + \not{\epsilon}\not{P} + \frac{\not{P}\hat{q}\cdot\epsilon}{m} - \frac{\not{P}\not{\epsilon}\not{\hat{q}}}{m}]\phi_V(\hat{q}) \quad (6)$$

Here, ϵ is the polarization vector of the vector quarkonia and N_V is the BS normalizer of the initial state quarkonia. The projected wave function for positive energy, $\psi^{++}(\hat{q})$ is obtained by the operation of projection operators on $\psi_P(\hat{q}')$ and $\psi_V(\hat{q})$ respectively as:

$$\begin{aligned} \psi_P^{++}(\hat{q}') &= \Lambda_1^+(\hat{q}')\frac{\not{P}'}{M'}\psi_P(\hat{q}')\frac{\not{P}'}{M'}\Lambda_2^+(\hat{q}') \\ \psi_V^{++}(\hat{q}) &= \Lambda_1^+(\hat{q})\frac{\not{P}}{M}\psi_V(\hat{q})\frac{\not{P}}{M}\Lambda_2^+(\hat{q}) \end{aligned} \quad (7)$$

where $\omega^2 = m^2 + \hat{q}^2$, $\omega'^2 = m^2 + \hat{q}'^2$, $\Lambda_{1,2}^+(\hat{q}) = \frac{1}{2\omega}[\frac{\not{P}}{M}\omega \pm m \pm \not{\hat{q}}]$, $\Lambda_{1,2}^+(\hat{q}') = \frac{1}{2\omega'}[\frac{\not{P}'}{M'}\omega' \pm m \pm \not{\hat{q}}']$, are called as projection operators. The relativistic positive energy wave function of equal mass pseudoscalar quarkonia in the center of mass system can be express as:

$$\psi_P^{++}(\hat{q}') = \frac{N'_P\phi_P(\hat{q}')}{2}\gamma_5\left\{M'\left[\frac{m^2}{\omega'^2} + \frac{m}{\omega'} - \frac{\hat{q}'^2}{m\omega'}\right] - \not{P}'\left[\frac{m^2}{\omega'^2} + \frac{m}{\omega'}\right] + \not{q}'\not{P}'\left[\frac{m}{\omega'^2} + \frac{1}{\omega'}\right]\right\} \quad (8)$$

And the relativistic positive energy wave function of equal mass vector quarkonia, in the center of mass system can be written as:

$$\begin{aligned} \psi_V^{++}(\hat{q}) &= \frac{N_V\phi_V(\hat{q})}{2}\left\{M\left[\frac{m^2}{\omega^2} - \frac{m}{\omega} + \frac{\hat{q}^2}{m\omega}\right]\not{\epsilon} + M\hat{q}\cdot\epsilon\left[\frac{m}{\omega^2} - \frac{1}{\omega}\right] + \left[\frac{m^2}{\omega^2} - \frac{m}{\omega}\right]\not{\epsilon}\not{P} \right. \\ &\quad \left. + \hat{q}\cdot\epsilon\left[\frac{m}{\omega^2} - \frac{1}{\omega}\right]\not{P} - \left[\frac{m}{\omega^2} - \frac{1}{\omega}\right]\not{P}\not{\hat{q}} + \frac{\hat{q}\cdot\epsilon}{m\omega}\not{\hat{q}}\not{P} - \frac{M\hat{q}\cdot\epsilon}{m\omega}\not{\hat{q}}\right\} \end{aligned} \quad (9)$$

The conjugate of $\psi_P^{++}(\hat{q}')$ is evaluated from $\bar{\psi}_P^{++}(\hat{q}') = \gamma^0(\psi_P^{++})^+\gamma^0$, after a sequence of steps is expressed as:

$$\bar{\psi}_P^{++}(\hat{q}') = -\frac{N'_P\phi_P(\hat{q}')}{2}\gamma_5\left\{M'\left[\frac{m^2}{\omega'^2} + \frac{m}{\omega'} - \frac{\hat{q}'^2}{m\omega'}\right] - \not{P}'\left[\frac{m^2}{\omega'^2} + \frac{m}{\omega'}\right] + \not{P}'\not{q}'\left[\frac{m}{\omega'^2} + \frac{1}{\omega'}\right]\right\} \quad (10)$$

Now, if we evaluate the total transition amplitude using equations Eq. (4), Eq. (9) and Eq. (10), we get,

$$M_{fi} = -\frac{e_q N'_P N_V}{2M} \int \frac{d^3 \hat{q}}{(2\pi)^3} \phi_P(\hat{q}') \phi_V(\hat{q}) [TR] \quad (11)$$

where

$$\begin{aligned} [TR] = & Tr \left\{ -\gamma_5 \not{P} \not{P}' \left[\frac{m^2}{\omega'^2} + \frac{m}{\omega'} \right] \left[M \left[\frac{m^2}{\omega^2} - \frac{m}{\omega} + \frac{\hat{q}^2}{m\omega} \right] \not{\epsilon} - \left[\frac{m}{\omega^2} - \frac{1}{\omega} \right] \not{\epsilon} \not{P} \not{\hat{q}} - \frac{M \hat{q} \cdot \epsilon}{m\omega} \not{\epsilon} \not{\hat{q}} \right] \right. \\ & \left. + \gamma_5 \not{P} \not{P}' \not{\hat{q}}' \left[\frac{m}{\omega'^2} + \frac{1}{\omega'} \right] \left[M \hat{q} \cdot \epsilon \left[\frac{m}{\omega^2} - \frac{1}{\omega} \right] \not{\epsilon} + \left[\frac{m^2}{\omega^2} - \frac{m}{\omega} \right] \not{\epsilon} \not{P} + \frac{\hat{q} \cdot \epsilon}{m\omega} \not{\epsilon} \not{\hat{q}} \not{P} \right] \right\} \end{aligned} \quad (12)$$

Evaluating trace over the gamma matrices, one can obtain the expression:

$$[TR] = -4M \epsilon_{\mu\nu\alpha\beta} P_\mu P'_\nu \epsilon_\alpha \epsilon_\beta \left[\frac{m}{\omega'^2} + \frac{1}{\omega'} \right] \left[\frac{m^3}{\omega^2} - \frac{m^2}{\omega} + \left(\frac{m}{\omega^2} - \frac{1}{\omega} \right) \hat{q} \cdot \hat{q}' \right] \quad (13)$$

where, P and P' are momenta of the initial (V) and final (P) quarkonia respectively. One can rewrite the transition amplitude after evaluation of the gamma matrices as:

$$M_{fi} = 2e_q N'_P N_V \int \frac{d^3 \hat{q}}{(2\pi)^3} \phi_P(\hat{q}') \phi_V(\hat{q}) \left[\frac{m}{\omega'^2} + \frac{1}{\omega'} \right] \left[\frac{m^3}{\omega^2} - \frac{m^2}{\omega} + \left(\frac{m}{\omega^2} - \frac{1}{\omega} \right) \hat{q} \cdot \hat{q}' \right] \epsilon_{\mu\nu\alpha\beta} P_\mu P'_\nu \epsilon_\alpha \epsilon_\beta \quad (14)$$

The decay width for $V \rightarrow P\gamma$, can be expressed as (see Ref.[25]):

$$\Gamma_{V \rightarrow P\gamma} = \frac{|\vec{P}'|}{8\pi M^2} |M|^2; |\vec{P}'| = \frac{1}{2M} [M^2 - M'^2] \quad (15)$$

From Equ. (14) for total amplitude, M_{tot} , one can obtain:

$$\begin{aligned} |M|^2 &= |F_V(\hat{q})|^2 (\epsilon_{\mu\nu\alpha\beta} P_\mu P'_\nu \epsilon_\alpha \epsilon_\beta)^2 \\ &= (M^4 + 10M^2 M'^2 + M'^4) |F_V(\hat{q})|^2 \end{aligned} \quad (16)$$

where

$$|F_V(\hat{q})|^4 = 2e_q N'_P N_V \int \frac{d^3 \hat{q}}{(2\pi)^3} \phi_P(\hat{q}') \phi_V(\hat{q}) \left[\frac{m}{\omega'^2} + \frac{1}{\omega'} \right] \left[\frac{m^3}{\omega^2} - \frac{m^2}{\omega} + \left(\frac{m}{\omega^2} - \frac{1}{\omega} \right) \hat{q} \cdot \hat{q}' \right] \quad (17)$$

where we have used $\hat{q}' = \hat{q}$, due to the nearly degenerate masses of P and V mesons corresponding to the transitions between states of same radial quantum numbers. One can obtain the expression of the decay rate of heavy quarkonia for the process, $V \rightarrow P\gamma$, as:

$$\Gamma_{V \rightarrow P\gamma} = \frac{1}{64\pi} \left[M^3 + \frac{M'^4}{M} - M'^2 M - \frac{M'^6}{M^3} \right]^3 |F_V(\hat{q})|^2 \quad (18)$$

M' and M are the masses of the Pseudoscalar heavy quarkonia (η_c, η_b) and Vector heavy quarkonia ($J/\psi, \Upsilon$) respectively. In the expression for $|f(\hat{q})|^2$, N_V and N'_P are the BS normalizers for heavy vector and pseudoscalar quarkonia, respectively, which are given in a simple form as in Ref.[21]:

$$N_V = \left[16mM_V \int \frac{d^3\hat{q}}{(2\pi)^3} \frac{\hat{q}^2}{\omega^3} \phi_V^2(\hat{q}) \right]^{-1/2} \quad (19)$$

and

$$N'_P = \left[\frac{16M_P}{m} \int \frac{d^3\hat{q}'}{(2\pi)^3} \frac{\hat{q}'^2}{\omega'} \phi_P^2(\hat{q}') \right]^{-1/2} \quad (20)$$

The ground state (1S) wave function and the radial wave functions for the 2S and 3S excitations for initial heavy vector quarkonia are written as in Ref.[21]:

$$\begin{aligned} \phi_V(1S, \hat{q}) &\approx \frac{1}{\beta_V^{3/2}} e^{-\frac{\hat{q}^2}{2\beta_V^2}}, \\ \phi_V(2S, \hat{q}) &\approx \frac{1}{\beta_V^{7/2}} (3\beta_V^2 - 2\hat{q}^2) e^{-\frac{\hat{q}^2}{2\beta_V^2}}, \\ \phi_V(3S, \hat{q}) &\approx \frac{1}{\beta_V^{11/2}} (15\beta_V^4 - 20\beta_V^2\hat{q}^2 + 4\hat{q}^4) e^{-\frac{\hat{q}^2}{2\beta_V^2}} \end{aligned} \quad (21)$$

where, $\beta_V = \left(2 \frac{m\omega_{q\bar{q}}^2}{\sqrt{1+2A_0(N+\frac{3}{2})}} \right)^{\frac{1}{4}}$, for the heavy vector quarkonia, which is dependent on the input kernel and contains the dynamical information of the initial heavy vector quarkonia.

The ground state (1S) wave function and the radial wave functions for the 2S and 3S excitations for final heavy pseudoscalar quarkonia are written as in Ref.[21]:

$$\begin{aligned} \phi_P(1S, \hat{q}') &\approx \frac{1}{\beta_P^{3/2}} e^{-\frac{\hat{q}'^2}{2\beta_P^2}}, \\ \phi_P(2S, \hat{q}') &\approx \frac{1}{\beta_P^{7/2}} (3\beta_P^2 - 2\hat{q}'^2) e^{-\frac{\hat{q}'^2}{2\beta_P^2}}, \\ \phi_P(3S, \hat{q}') &\approx \frac{1}{\beta_P^{11/2}} (15\beta_P^4 - 20\beta_P^2\hat{q}'^2 + 4\hat{q}'^4) e^{-\frac{\hat{q}'^2}{2\beta_P^2}} \end{aligned} \quad (22)$$

where $\hat{q}' = \hat{q} + \frac{1}{2}|\vec{P}'|$ and $|\vec{P}'| = \frac{M_V}{2}(1 - \frac{M_P^2}{M_V^2})$ and $\beta_P = \left(4 \frac{m\omega_{q\bar{q}}^2}{\sqrt{1+2A_0(N+\frac{3}{2})}} \right)^{\frac{1}{4}}$, for the heavy pseudoscalar quarkonia, which is dependent on the input kernel and contains the dynamical information of the final heavy vector quarkonia.

Numerical results

We use the same input parameters of our model fixed from the mass spectrum: $C_0 = 0.21$, $\omega_0 =$

0.15GeV , QCD length scale $\Lambda = 0.20\text{GeV}$, $A_0 = 0.01$ and the input quark masses, $m_c = 1.49\text{GeV}$ and $m_b = 5.07\text{GeV}$ to calculate the decay widths for the process $V- \rightarrow P + \gamma$ studied above. The results of radiative decay widths of our model are listed in Table 1.

Transition	Our work	Expt.[28]	LFM[29]	PM[30]
$\Gamma_{J/\psi(1S) \rightarrow \eta_c(1S)\gamma}$	2.0054	1.5687 ± 0.011	1.67 ± 0.05	1.8
$\Gamma_{\psi(2S) \rightarrow \eta_c(2S)\gamma}$	0.5709	0.2093 ± 0.002		0.4
$\Gamma_{\psi(3S) \rightarrow \eta_c(3S)\gamma}$	0.2984			
$\Gamma_{\Upsilon(1S) \rightarrow \eta_b(1S)\gamma}$	0.3387		$0.043^{+0.09}_{-0.03}$	0.001
$\Gamma_{\Upsilon(2S) \rightarrow \eta_b(2S)\gamma}$	0.1053			0.0002
$\Gamma_{\Upsilon(3S) \rightarrow \eta_b(3S)\gamma}$	0.0781			

Table 1: Radiative decay widths of equal mass heavy vector quarkonium of ground state (1S) and radially excited states (2S, 3S) in present calculation (BSE-CIA) along with their decay widths in other models and experimental data (all values are in units of Kev).

3 Discussions and conclusion

We have employed a 3D reduction of BSE (with a 4×4 representation for two-body ($q\bar{q}$) BS amplitude) under Covariant Instantaneous Ansatz (CIA) for deriving the algebraic forms of the electromagnetic transition amplitude equation, in Eq.(1), leading to decay width for ground and radially excited states of vector (J/Ψ , and Υ) quarkonia.

The decay width calculated in this BSE framework for (1S, 2S, 3S) states of J/ψ and Υ are shown in Table 1. This numerical calculation have been done using Mathematica. We first fit our parameters to the ground state masses of η_c , η_b , J/ψ and Υ . These input parameters were found to be $C_0 = 0.130$, $\omega_0 = 0.100$ GeV, $\Lambda = 0.200$ GeV and $A_0 = 0.010$. With input quark masses $m_c = 1.485$ GeV and $m_b = 5.070$ GeV we obtained the best fit to these ground state masses. The same set of parameters above was used to calculate the masses of all the other (excited) states of η_c , η_b , J/ψ and Υ , as well as the leptonic decay constants of these states. Two-photon as well as two gluon decay widths of η_c , and η_b was further studied (see our recent paper [21]).

The results obtained for decay width of ground and radially excited states of J/ψ and Υ are somewhat on the higher side in comparison to central values of data for these states, which might be probably due to the absence of the negative energy part employed in Eq.(1), where we have considered only the positive energy part (as in Ref. [27] employed earlier for heavy mesons). However, from the Table, a wide range of variation of radiative decay widths of $J/\Psi(1S)$, and $\Psi(2S)$ states in different models can be observed. However, the important thing is that our radiative decay width values for vector quarkonia, show a marked decrease as one goes from $1S$ to $3S$ state for J/ψ and Υ , which is again in conformity with data and other models. We have also given our predictions for radiative decay widths of $\Psi(3S)$, $\Upsilon(1S)$, $\Upsilon(2S)$, and $\Upsilon(3S)$ states, for which data is currently not yet available. The aim of doing this study was to mainly test our analytic forms of wave functions obtained as solutions of mass spectral equations in an approximate harmonic oscillator basis obtained analytically from the 4×4 BSE as our starting point (that had so far given good predictions for leptonic decays of P and V quarkonia, and the two-photon, and the two-gluon decays of P-quarkonia [21]), to the single photon radiative decays of V-quarkonia. This would in turn lead to validation of our approach, which provides a much deeper insight than the purely numerical calculations in 4×4 BSE approach that are prevalent in the literature.

We are not aware of any other BSE framework, involving 4×4 BS amplitude, and with all the Dirac structures incorporated (in fact many works use only the leading Dirac structure), that treats the mass spectral problem analytically, and uses the wave functions derived analytically from this equation for calculation of various transitions. To the best of our knowledge, all the other 4×4 BSE approaches treat this problem numerically after obtaining the coupled set of equations..

References

- [1] B.Auger *et al.*(BaBar collaboration), Phys. Rev. Lett. 103, 161801 (2009).
- [2] K.F.Chen *et al.*(Belle collaboration), Phys. Rev. D82, 091106(R) (2010).
- [3] K.W.Edwards *et al.*(CLEO collaboration), Phys. Rev.Lett. 86, 30 (2001).
- [4] K.A.Olive *et al.*, (Particle Data Group), Chin. Phys. **C38**, 090001 (2014).
- [5] C.McNielle, C.T.H.Davies, E.Follana, K.Hornbostel, G.P.Lepage (HPQCD Collaboration), arxiv: 1207.0994[hep-lat] (2012).
- [6] J.Gasser, H.Leutwyler, Ann. Phys. 158, 142 (1984).
- [7] M.Neubert, Phys. Rep. 245, 259 (1994).
- [8] E.Veli Veliev, K.Azizi, H.Sundu, N.Aksit, J.Phys. G39, 015002(2012).
- [9] G.T.Bodwin, E.Braaten, G.P.Lepage, Phys. Rev. D51, 1125 (1995).
- [10] C.H.L.Smith, Ann. Phys. (N.Y.) 53, 521 (1969).
- [11] A. N. Mitra, B. M. Sodermark, Nucl. Phys. **A695**, 328 (2001).
- [12] R. Alkofer, L. W. Smekel, Phys. Rep.**353**, 281 (2001).
- [13] C-H. Chang, G. L. Wang, Sci. China G53, 2005 (2010).
- [14] R. Ricken, M. Koll, D. Merten, B. Metsch, H. Petry, Eur. Phys. J. A9, 221 (2000).
- [15] A.N.Mitra, S.Bhatnagar, Intl. J. Mod. Phys. A07, 121 (1992).
- [16] S.Patel, P.C.Vinodkumar, S.Bhatnagar, Chinese Phys. C40 (2016)(In press); arXiv:1504.01103[hep-ph].
- [17] A. N. Mitra, Proc. Indian Natl. Sci. Acad. 65, 527 (1999).
- [18] S.Bhatnagar, D.S.Kulshreshtha, A.N.Mitra, Physics Letters B 263, 485 (1991).
- [19] S.Bhatnagar, J.Mahecha, Y.Mengesha, Phys. Rev. D90, 014034 (2014).

- [20] H.Negash, S.Bhatnagar, Intl. J. Mod. Phys. E24, 1550030 (2015).
- [21] H.Negash, S.Bhatnagar, Intl. J. Mod. Phys. E25,1650059 (2016); arXiv:1508.06131v2 [hep-ph] (2016).
- [22] CHANG Chao-Hsi, CHEN Jiao-Kai, and WANG Guo-Li, Commun. Theor. Phys. (Beijing, China) 46 (2006) pp. 467-480.
- [23] C-H. Chang, J-K. Chen, G-L. Wang: [arXiv:hep-th/0312250] (2003).
- [24] N. Nimai Singh and A. N. Mitra, Phys. Re. D 38, 5 (1988).
- [25] E.Mengesha, S.Bhatnagar, Intl. J. Mod. Phys. E21, 1250084 (2012).
- [26] Tian-Hong Wang, and and Guo-Li Wang, arXiv:1006.3363v4 [hep-ph] 7 Feb 2011.
- [27] Tianhong Wang, Guo-Li Wang, Yue Jiang, and Wan-Li Ju, arXiv:1205.5725v2 [hep-ph] 7 Sep 2012.
- [28] K.A.Olive *et al.*, (Particle Data Group), Chin. Phys. **C38**, 090001 (2014).
- [29] Ho-Meoyng Choi, arXiv:hep-ph/0701263v2 28 Mar 2007.
- [30] Stanley F. Radford, Wayne W. Repko, Phys. Re. D 75, 074031 (2007).