

On Consistency of Graph-based Semi-supervised Learning

Chengan Du

Center for Outcomes Research
and Evaluation
Yale University
New Haven, Connecticut
chengan.du@yale.edu

Yunpeng Zhao

School of Mathematical and
Natural Sciences
Arizona State University
Tempe, Arizona
yunpeng.zhao@asu.edu

Feng Wang

School of Mathematical and
Natural Sciences
Arizona State University
Tempe, Arizona
fwang25@asu.edu

Abstract—Graph-based semi-supervised learning is one of the most popular methods in machine learning. Some of its theoretical properties such as bounds for the generalization error and the convergence of the graph Laplacian regularizer have been studied in computer science and statistics literature. However, a fundamental statistical property – *consistency*¹ – has not been proved.

In this article, we study the consistency problem under a non-parametric framework. We obtain the following two results: 1) We prove that graph-based semi-supervised learning on the test data is consistent in the case that the estimated scores are enforced to be equal to the observed responses for the labeled data (the hard criterion). The sample size of unlabeled data are allowed to grow at a slower rate than the size of the labeled data in this result. 2) We give a counterexample demonstrating that the estimator can be inconsistent for the case when the estimated scores are not required to be equal to the observed responses (the soft criterion), where a tuning parameter is used to balance the loss function and the graph Laplacian regularizer. These somewhat surprising theoretical findings are supported by numerical studies on both synthetic and real datasets.

Moreover, numerical studies show that the hard criterion constantly outperforms the soft criterion even when the sample size of unlabeled data is smaller than the size of labeled data. This suggests that practitioners can safely choose the hard criterion without the burden of selecting the tuning parameter in the soft criterion.

Index Terms—semi-supervised learning, consistency, graph Laplacian

I. INTRODUCTION

Semi-supervised learning is a class of machine learning methods in the middle ground between *supervised learning*, where all training data are labeled, and *unsupervised learning*, where no training data are labeled. Specifically, in addition to the labeled training data $X_1, \dots, X_n$, there exist unlabeled inputs $X_{n+1}, \dots, X_{n+m}$. Under certain assumptions on the geometric structure of the input data, such

as the cluster assumption or the low-dimensional manifold assumption [1], the use of both labeled and unlabeled data can achieve better prediction accuracy than supervised learning, which only uses labeled inputs $X_1, \dots, X_n$.

Semi-supervised learning has become popular because the acquisition of unlabeled data is relatively inexpensive. A large number of methods have been developed under the framework of semi-supervised learning. For example, [2] proposed that the combination of labeled and unlabeled data will improve the prediction accuracy under the assumption of mixture models. The self-training method [3] and the co-training method [4] were then applied to semi-supervised learning when mixture models are not assumed. Reference [5] described an approach to semi-supervised clustering based on hidden Markov random fields (HMRFs) that can combine multiple approaches in a unified probabilistic framework. Reference [6] proposed a probabilistic framework for semi-supervised learning incorporating a K-means-type clustering algorithm (HMRF-Kmeans). Reference [7] proposed the transductive support vector machines (TSVMs) that used the idea of transductive learning by including unlabeled data in the computation of the margin. Reference [8] used a convex relaxation of the optimization problem called semi-definite programming as a different approaches to the TSVMs.

In this article, we focus on a particular semi-supervised method – graph-based semi-supervised learning. In this method, the geometric structure of the input data are represented by a weighted graph $\mathbf{G} = (V, E)$, where nodes $V = \{v_1, \dots, v_{n+m}\}$ represent the inputs $X_1, \dots, X_{n+m}$ and edges E represent the similarities between them. The similarities are given in an $n + m$ by $n + m$ symmetric similarity matrix (or called *kernel matrix*), $\mathbf{W} = [w_{ij}]$, where $0 \leq w_{ij} \leq 1$. The larger w_{ij} implies that X_i and X_j are more similar. Furthermore, let $Y_1, \dots, Y_n$ be the responses of the labeled data.

Reference [9] proposed the following graph-based learn-

¹Throughout the paper, *consistency* is used as a statistical term referring to an asymptotic property – that is, the prediction by the algorithm can identify the underlying truth with unlimited data. This is not to be confused with the existence of solutions in an equation system, which is a term used in algebra.

ing method,

$$\begin{aligned} \min_{\mathbf{f}=(f_1, \dots, f_{n+m})^T} & \sum_{i=1}^{n+m} \sum_{j=1}^{n+m} w_{ij} (f_i - f_j)^2 \\ \text{subject to } & f_i = Y_i, i = 1, \dots, n. \end{aligned} \quad (1)$$

We call the solution estimated scores. The objective function (1) (hereafter referred to as “hard criterion”), requires all estimated scores to be exactly the same as the responses for the labeled data. Reference [10] relaxed this requirement by proposing a soft version (hereafter referred to as “soft criterion”). We follow an equivalent form given in [11],

$$\begin{aligned} \min_{\mathbf{f}=(f_1, \dots, f_{n+m})^T} & \sum_{i=1}^n (Y_i - f_i)^2 \\ & + \frac{\lambda}{2} \sum_{i=1}^{n+m} \sum_{j=1}^{n+m} w_{ij} (f_i - f_j)^2. \end{aligned} \quad (2)$$

The soft criterion belongs to the “loss+penalty” paradigm: it searches for the minimizer $\hat{\mathbf{f}}$, which improves the smoothness of $\hat{\mathbf{f}}$ by a penalty-based similarity matrix while causing a training error.

These two criteria are closely related: when $\lambda = 0$ the soft criterion is equivalent to the hard criterion.

Remark 1. *The tuning parameter λ being 0 in the soft criterion (2) is understood in the following sense: the squared loss has an infinite weight and thereby f_i is enforced to be Y_i for all the labeled data. But the penalty term $\sum_{i=1}^{n+m} \sum_{j=1}^{n+m} w_{ij} (f_i - f_j)^2$ still plays a crucial role when it has no conflict with the hard constraints on the labeled data – that is, it builds a connection between f_i ’s on the labeled and unlabeled data. More precisely, the solution of Eq. (2) goes to the solution of Eq. (1) as $\lambda \rightarrow 0$ (see page 203 of [1] and Proposition II.1 for a detailed explanation).*

References [12], [13] also proposed different variants of graph-based learning methods. We only focus on Eq. (1) and (2) in this article.

The theoretical properties of graph-based learning have been studied in computer science and statistics literature. Reference [14] derived the limit of the Laplacian regularizer when the sample size of unlabeled data goes to infinity. Reference [15] considered the convergence of Laplacian regularizer on Riemannian manifolds. Reference [16] reinterpreted the graph Laplacian as a measure of intrinsic distances between inputs on a manifold and reformulated the problem as a functional optimization in a reproducing kernel Hilbert space. Reference [17] pointed out that the hard criterion can yield a completely noninformative solution when the size of unlabeled data goes to infinity and labeled data are finite – that is, the solution can give a perfect fit on the labeled data but remains as 0 on the unlabeled data. Reference [18] obtained the asymptotic mean squared error of a different version of graph-based learning criterion.

Reference [13] gave a bound of the generalization error for a slightly different version of objective function (2). Reference [19] studied the theoretical properties of ℓ_p -based Laplacian regularization – in particular the phase transition of p for an informative solution.

However, no result is available in the literature on a very fundamental question – the consistency of graph-based learning – which is the main focus of this article. Specifically, we want to answer the question of *under what conditions $\hat{f}_i$ will converge to $\mathbb{E}[Y_i|X_i]$ on unlabeled data*, where $\mathbb{E}[Y_i|X_i]$ is the true probability of a positive label given X_i if responses are binary, and $\mathbb{E}[Y_i|X_i]$ is the regression function on X_i if responses are continuous. For simplicity, we will always call $\mathbb{E}[Y_i|X_i]$ as regression function.

Most of the literatures on graph-based semi-supervised learning considered a “functional version” of Eq. (1) and (2). Previous works used a functional optimization problem with the optimizer $\hat{f}(x)$ being a function, as an approximation of the original problem with the optimizer $\hat{\mathbf{f}}$ being a vector. The behavior of the limit of graph Laplacian and the solution $\hat{f}(x)$ were studied in this context.

Instead of adopting this framework, we use a more direct approach. We focus on the original problem and study the relations of $\hat{f}_i$ and $\mathbb{E}[Y_i|X_i]$ directly under the general non-parametric setting. Our approach essentially belongs to the framework of transductive learning, which focuses on the prediction on the given unlabeled data $X_{n+1}, \dots, X_{n+m}$, not the general mapping from inputs to responses. By establishing a link between the optimizer of Eq. (1) and the Nadaraya-Watson estimator [20], [21] for kernel regression, we prove the consistency of the hard criterion. Unlike [17], our result requires the sample size of labeled data goes infinity, which is natural in asymptotic theory of statistics. The result also allows the size of unlabeled data goes to infinity. On the other hand, we show that the soft criterion is inconsistent for sufficiently large λ . To the best of our knowledge, this is the first result that explicitly distinguishes the hard criterion and the soft criterion of graph-based learning from a theoretical perspective.

The main results are stated in Section II and proved in Section IV. We give a toy example in Section III to give more intuition into the somewhat surprising theoretical findings. The results are further supported by numerical studies on synthetic and real datasets in Section V. Moreover, numerical studies also show that the hard criterion constantly outperforms the soft criterion even when the sample size of unlabeled data is smaller than the size of labeled data. This suggests that practitioners can safely choose the hard criterion without the burden of selecting the tuning parameter in the soft criterion.

II. MAIN RESULTS

Let $(X_1, Y_1), \dots, (X_{n+m}, Y_{n+m})$ be independently and identically distributed pairs. Each X_i is a d -dimensional vector and $\mathbf{Y} = (Y_1, \dots, Y_{n+m})^T$ are binary responses labeled as 1 and 0 (the classification case) or continuous responses (the regression case). The last m responses are unobserved.

We now give the solution of the soft version (2). Recall that W is the similarity matrix. Let $\mathbf{D} = \text{diag}(d_1, \dots, d_{n+m})$ where $d_i = \sum_{j=1}^{n+m} w_{ij}$, and $\mathbf{L} = \mathbf{D} - \mathbf{W}$ being the unnormalized graph Laplacian (see [22] for details). Soft criterion (2) can be written in matrix form

$$\min_{\mathbf{f}} (\mathbf{f} - \mathbf{Y})^T \mathbf{V} (\mathbf{f} - \mathbf{Y}) + \lambda \mathbf{f}^T \mathbf{L} \mathbf{f}, \quad (3)$$

where $\mathbf{V}$ is an $n+m$ by $n+m$ matrix defined as

$$\mathbf{V} = \begin{pmatrix} \mathbf{I}_n & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{pmatrix}.$$

By taking the derivative of Eq. (3) with respect to $\mathbf{f}$ and setting equal to zero, we obtain the following solution:

$$\hat{\mathbf{f}} = (\mathbf{V} + \lambda \mathbf{L})^{-1} \begin{pmatrix} \mathbf{Y}_n \\ \mathbf{0} \end{pmatrix}.$$

where $\mathbf{Y}_n = (Y_1, \dots, Y_n)^T$.

Our objective is to determine the estimated scores on the unlabeled data, i.e., $\hat{\mathbf{f}}_{(n+1):(n+m)} = (\hat{f}_{n+1}, \dots, \hat{f}_{n+m})^T$. In order to obtain an explicit form for $\hat{\mathbf{f}}_{(n+1):(n+m)}$, we use a formula for inverse of a block matrix: for any non-singular square matrix

$$\mathbf{A} = \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{pmatrix},$$

$$\mathbf{A}^{-1} = \begin{pmatrix} (\mathbf{A}_{11} - \mathbf{A}_{12} \mathbf{A}_{22}^{-1} \mathbf{A}_{21})^{-1} & -(\mathbf{A}_{22} - \mathbf{A}_{21} \mathbf{A}_{11}^{-1} \mathbf{A}_{12})^{-1} \mathbf{A}_{21} \mathbf{A}_{11}^{-1} \\ -(\mathbf{A}_{11} - \mathbf{A}_{12} \mathbf{A}_{22}^{-1} \mathbf{A}_{21})^{-1} \mathbf{A}_{12} \mathbf{A}_{22}^{-1} & (\mathbf{A}_{22} - \mathbf{A}_{21} \mathbf{A}_{11}^{-1} \mathbf{A}_{12})^{-1} \end{pmatrix}.$$

Write $\mathbf{D}$ and $\mathbf{W}$ as 2×2 block matrices,

$$\mathbf{D} = \begin{pmatrix} \mathbf{D}_{11} & \mathbf{D}_{12} \\ \mathbf{D}_{21} & \mathbf{D}_{22} \end{pmatrix}, \mathbf{W} = \begin{pmatrix} \mathbf{W}_{11} & \mathbf{W}_{12} \\ \mathbf{W}_{21} & \mathbf{W}_{22} \end{pmatrix}.$$

By the formula above,

$$\begin{aligned} \hat{\mathbf{f}}_{(n+1):(n+m)} &= \\ &(\mathbf{D}_{22} - \mathbf{W}_{22} - \lambda \mathbf{W}_{21} (\mathbf{I}_n + \lambda \mathbf{D}_{11} - \lambda \mathbf{W}_{11})^{-1} \mathbf{W}_{12})^{-1} \\ &\cdot \mathbf{W}_{21} (\mathbf{I}_n + \lambda \mathbf{D}_{11} - \lambda \mathbf{W}_{11})^{-1} \mathbf{Y}_n. \end{aligned} \quad (4)$$

By letting $\lambda = 0$, we obtain

$$\hat{\mathbf{f}}_{(n+1):(n+m)} = (\mathbf{D}_{22} - \mathbf{W}_{22})^{-1} \mathbf{W}_{21} \mathbf{Y}_n. \quad (5)$$

Some literatures such as [11] used Lagrange multipliers to the constrained optimization problem and obtained the

same solution for the hard criterion (1). Therefore, we have essentially proved the following proposition:

Proposition II.1. *The solution of the soft criterion (2) at $\lambda = 0$ is equivalent to the solution of the hard criterion.*

It is worth noting that the time complexities of computing Eq. (5) and (4) are respectively $O(m^3)$ and $O((m+n)^3)$ when using Gaussian elimination for solving systems of linear equations. Therefore, it is more efficient to solve the hard criterion, which is another advantage of the hard criterion in addition to theoretical considerations.

The form of Eq. (5) is closely related to the Nadaraya-Watson estimator [20], [21] for kernel regression, which is

$$\hat{q}_{n+a} = \frac{\sum_{i=1}^n w_{n+a,i} Y_i}{\sum_{k=1}^n w_{n+a,k}}, \quad a = 1, \dots, m. \quad (6)$$

The Nadaraya-Watson estimator is well studied under the non-parametric framework. We can construct $\mathbf{W}$ by a kernel function – that is, let $w_{ij} = K((X_i - X_j)/h_n)$, where K is a nonnegative function on $\mathbb{R}^d$, and h_n is a positive constant controlling the bandwidth of the kernel. Let $q(X) = \mathbb{E}[Y|X]$ be the true regression function. The consistency of Nadaraya-Watson estimator was first proved by [20], [21]. Many other researchers such as [23] and [24] studied its asymptotic properties under different assumptions. Here, we follow the result in [25]. If $h_n \rightarrow 0$, $nh_n^d \rightarrow \infty$ as $n \rightarrow \infty$, and K satisfies:

- (i) K is bounded by $k^* < \infty$;
- (ii) The support of K is compact;
- (iii) $K \geq \beta I_B$ for some $\beta > 0$ and some closed ball B centered at the origin and having positive radius δ ,

then $\hat{q}_{n+a}$ converges to $q(X_{n+a})$ in probability for $a = 1, \dots, m$.

By establishing a connection between the solution of the hard criterion and Nadaraya-Watson estimator, we prove the following main theorem:

Theorem II.1. *Suppose that*

$(X_1, Y_1), \dots, (X_{n+m}, Y_{n+m})$ are independently and identically distributed with Y_i being bounded; h_n and K satisfy the above conditions. Further, we assume that the density function $\phi(\cdot)$ of X_1 has a compact support $\mathcal{X}$. And for every inner point x in $\mathcal{X}$,

$$\phi(x) \geq s^* > 0. \quad (7)$$

Then, for $m = o(nh_n^d)$, $\hat{f}_{n+a}$ given in Eq. (4) converges to $q(X_{n+a})$ in probability, for $a = 1, \dots, m$.

The proof will be given in Section IV.

Theorem II.1 establishes the consistency of the hard criterion under the standard non-parametric framework with two additional assumptions. First, both labeled data and unlabeled data are allowed to grow but the size of unlabeled

data m grows slower than the size of labeled data n . We conjecture that when m grows faster than n , the graph-based semi-supervised learning may not be consistent based on the simulation studies in Section V although it still outperforms the soft criterion. Reference [17] also suggested that the method may not work well when m grows too fast. Second, we assume the density function of the difference of two independent inputs is strictly positive near the origin, which is a mild technical condition valid for commonly used density functions.

We now consider the soft criterion ($\lambda \neq 0$).

Proposition II.2. *Suppose that $(X_1, Y_1), \dots, (X_{n+m}, Y_{n+m})$ are independently and identically distributed with Y_i being bounded. Furthermore, suppose $\mathbf{W}$ represents a connected graph. Then for sufficiently large λ , the soft criterion (2) is inconsistent.*

Proof. Consider another extreme case of the soft criterion (2), $\lambda = \infty$. When $\mathbf{W}$ represents a connected graph, the objective function becomes

$$\min_{\mathbf{f}=(f_1, \dots, f_n)^T} \sum_{i=1}^n (Y_i - f_i)^2 \quad (8)$$

subject to $f_i = f_j, 1 \leq i, j \leq n+m$.

It is simple to verify that the solution of Eq. (8), denoted by $\hat{\mathbf{f}}(\infty)$, is given by

$$\hat{f}_{n+a}(\infty) = \frac{1}{n} \sum_{i=1}^n Y_i, \quad a = 1, \dots, m.$$

By the law of large numbers,

$$\lim_{n \rightarrow \infty} \hat{f}_{n+a}(\infty) = \mathbb{E}[q(X_1)] \text{ almost surely.}$$

Clearly, $\mathbb{E}[q(X_1)] \neq q(X_{n+a})$ since the right-hand side is a random variable. This implies that for sufficiently large λ , the soft criterion is inconsistent. $\square$

This counterexample in fact suggests more general results than its first look. We prove that under $\lambda \rightarrow \infty$, the soft criterion predicts the same label (an extremely inaccurate prediction). On the contrary, the hard criterion ($\lambda = 0$) gives a consistent prediction. Note that Eq. (5) is a continuous function of λ , so the prediction cannot suddenly jump from consistent to extremely inaccurate. This suggests in general the soft criterion is inconsistent for all or a wide range of nonzero λ .

III. A TOY EXAMPLE

Theorem II.1 and Proposition II.2 provide somewhat surprising insights about the graph-based semi-supervised learning. At a first glance, the hard criterion makes an impractical assumption that requires the responses to be noiseless, while the soft criterion seems to be a more natural choice. According to our theoretical result, the hard criterion

is however consistent under the standard non-parametric framework where the responses on training data are allowed to be random and noisy by default. Below we provide a toy example² that further illustrates the rationale behind the hard criterion.

Consider the case of $X_1, \dots, X_{n+m}$ being the same constant. Thus, $Y_1, \dots, Y_{n+m}$ become independently and identically distributed random variables. We still assume that the first n responses are observed and the last m are to be predicted. Let $\mathbf{W}$ be the Gaussian radial basis function (RBF) kernel, that is,

$$w_{ij} = \exp\left(-\frac{\|X_i - X_j\|^2}{\sigma^2}\right), \text{ for } 1 \leq i, j \leq m+n.$$

Then $w_{ij} \equiv 1$ for $1 \leq i, j \leq m+n$. Thus,

$$\mathbf{D}_{22} - \mathbf{W}_{22} = \begin{pmatrix} m+n-1 & -1 & \cdots & -1 \\ -1 & m+n-1 & \cdots & -1 \\ \vdots & \vdots & \ddots & \vdots \\ -1 & -1 & \cdots & m+n-1 \end{pmatrix}_{m \times m}.$$

It is easy to verify that³

$$(\mathbf{D}_{22} - \mathbf{W}_{22})^{-1} = \begin{pmatrix} \frac{n+1}{n(m+n)} & \frac{1}{n(m+n)} & \cdots & \frac{1}{n(m+n)} \\ \frac{1}{n(m+n)} & \frac{n+1}{n(m+n)} & \cdots & \frac{1}{n(m+n)} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{n(m+n)} & \frac{1}{n(m+n)} & \cdots & \frac{n+1}{n(m+n)} \end{pmatrix}_{m \times m}.$$

Thus, from Eq. (5) the solution of the hard criterion is $\hat{\mathbf{f}}_i \equiv \frac{1}{n} \sum_{j=1}^n Y_j$ for $i = n+1, \dots, n+m$, and $\hat{\mathbf{f}}_i = Y_i$ for $i = 1, \dots, n$. This is in fact the best solution one can expect: for labeled data we simply used the observed responses and for unlabeled data we used the mean of the observed responses to do prediction (since there is no other information available). This example shows in the transductive learning setup it is not an issue that the hard criterion did not smoothen the labeled data (there is no need to do prediction on labeled data in the first place), while the prediction on unlabeled data is indeed based on a weighted average of the observed responses.

IV. PROOF OF THE MAIN THEOREM

We give the proof of Theorem II.1 in this section.

Recall that

$$\hat{\mathbf{f}}_{(n+1):(n+m)} = (\mathbf{D}_{22} - \mathbf{W}_{22})^{-1} \mathbf{W}_{21} \mathbf{Y}_n.$$

²This example is motivated by comments from an anonymous reviewer.

³Note that the size of $\mathbf{D}_{22} - \mathbf{W}_{22}$ is $m \times m$ but not $(m+n) \times (m+n)$ and this matrix is invertible.

We first focus on $(\mathbf{D}_{22} - \mathbf{W}_{22})^{-1}$. Clearly,

$$(\mathbf{D}_{22} - \mathbf{W}_{22})^{-1} = (\mathbf{I}_m - \mathbf{D}_{22}^{-1}\mathbf{W}_{22})^{-1}\mathbf{D}_{22}^{-1}.$$

For any positive integer l , define

$$\mathbf{S}_l = \mathbf{D}_{22}^{-1}\mathbf{W}_{22} + (\mathbf{D}_{22}^{-1}\mathbf{W}_{22})^2 + (\mathbf{D}_{22}^{-1}\mathbf{W}_{22})^3 + \dots + (\mathbf{D}_{22}^{-1}\mathbf{W}_{22})^l.$$

Our goal is to prove that the limit of $\mathbf{S}_l$ exists with probability approaching 1, and thus we have

$$(\mathbf{I}_m - \mathbf{D}_{22}^{-1}\mathbf{W}_{22})^{-1} = \mathbf{I}_m + \lim_{l \rightarrow \infty} \mathbf{S}_l$$

with probability approaching 1 [26].

By definition,

$$\mathbf{D}_{22} = \begin{pmatrix} d_{n+1,n+1} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & d_{n+m,n+m} \end{pmatrix},$$

$$\mathbf{W}_{22} = \begin{pmatrix} w_{n+1,n+1} & \cdots & w_{n+1,n+m} \\ \vdots & \ddots & \vdots \\ w_{n+m,n+1} & \cdots & w_{n+m,n+m} \end{pmatrix},$$

where

$$d_{n+a,n+a} = \sum_{k=1}^{n+m} w_{n+a,k}, \quad w_{n+a,i} = K \left(\frac{X_i - X_{n+a}}{h_n} \right),$$

for $1 \leq a \leq m, 1 \leq i \leq n+m$. Thus we have

$$\mathbf{D}_{22}^{-1}\mathbf{W}_{22} = \begin{pmatrix} \frac{w_{n+1,n+1}}{d_{n+1,n+1}} & \cdots & \frac{w_{n+1,n+m}}{d_{n+1,n+1}} \\ \vdots & \ddots & \vdots \\ \frac{w_{n+m,n+1}}{d_{n+m,n+1}} & \cdots & \frac{w_{n+m,n+m}}{d_{n+m,n+m}} \end{pmatrix}.$$

Define $p(X_{n+a}) = \mathbb{P}(\|X_i - X_{n+a}\| \leq \delta h_n \mid X_{n+a})$. Since $h_n \rightarrow 0$, there exist $n_0 \in \mathbb{N}$ such that $\delta h_n \leq c$ holds for every $n > n_0$. Then by Eq. (7) and the definition of multiple integral, with probability 1.

$$\lim_{n \rightarrow \infty} \frac{p(X_{n+a})}{V_d(\delta h_n)} = \phi(X_{n+a}) \geq s^*,$$

where $V_d(\delta h_n)$ denotes the volume of a d -dimensional ball with radius δh_n . Thus, for sufficiently large n ,

$$p(X_{n+a}) \geq \frac{1}{2}s^*V_d(\delta h_n) = sh_n^d,$$

where s is a constant only related to s^* and δ .

Since $nh_n^d \rightarrow \infty$, the above inequality implies $np(X_{n+a}) \rightarrow \infty$. On the other side, $p(X_{n+a}) \rightarrow 0$ since $h_n \rightarrow 0$.

Further,

$$\begin{aligned} & \text{Var}(I\{\|X_i - X_{n+a}\| \leq \delta h_n\} \mid X_{n+a}) \\ &= p(X_{n+a})(1 - p(X_{n+a})). \end{aligned}$$

By Chebyshev's Inequality, for any $0 < \epsilon < 1/2$, since $nh_n^d \rightarrow \infty$,

$$\begin{aligned} & \mathbb{P} \left(\left| \frac{\sum_{i=1}^n I\{\|X_i - X_{n+a}\| \leq \delta h_n\}}{np(X_{n+a})} - 1 \right| \geq \epsilon \mid X_{n+a} \right) \\ &= \mathbb{P} \left(\left| \frac{1}{n} \sum_{i=1}^n I\{\|X_i - X_{n+a}\| \leq \delta h_n\} - p(X_{n+a}) \right| \geq \epsilon p(X_{n+a}) \mid X_{n+a} \right) \\ &\leq \frac{p(X_{n+a})(1 - p(X_{n+a}))}{n\epsilon^2 p(X_{n+a})^2} \leq \frac{1}{\epsilon^2 p(X_{n+a})n} \leq \frac{1}{\epsilon^2 snh_n^d}. \end{aligned} \quad (9)$$

Therefore,

$$\begin{aligned} & \mathbb{P} \left(\left| \frac{\sum_{i=1}^n I\{\|X_i - X_{n+a}\| \leq \delta h_n\}}{np(X_{n+a})} - 1 \right| \geq \epsilon \right) \\ &\leq \frac{1}{\epsilon^2 snh_n^d} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

This further implies

$$\frac{\sum_{i=1}^n I\{\|X_i - X_{n+a}\| \leq \delta h_n\}}{np(X_{n+a})} \rightarrow 1 \quad \text{in probability.}$$

We now continue to study the property of $\mathbf{D}_{22}^{-1}\mathbf{W}_{22}$. Consider each element $(\mathbf{D}_{22}^{-1}\mathbf{W}_{22})_{ab}$ of this matrix. For $1 \leq a, b \leq m$,

$$\begin{aligned} & (\mathbf{D}_{22}^{-1}\mathbf{W}_{22})_{ab} = \frac{w_{n+a,n+b}}{d_{n+a,n+a}} \\ &= K \left(\frac{X_{n+b} - X_{n+a}}{h_n} \right) / \sum_{i=1}^{n+m} K \left(\frac{X_i - X_{n+a}}{h_n} \right) \\ &\leq \frac{k^*}{\beta \sum_{i=1}^n I\{\|X_i - X_{n+a}\| \leq \delta h_n\}}, \end{aligned}$$

by condition (i) and (iii). For simplicity of notation, let

$$\Phi_n(a) = \frac{\sum_{i=1}^n I\{\|X_i - X_{n+a}\| \leq \delta h_n\}}{np(X_{n+a})},$$

where Φ_n is a nonnegative function depending on n . By Eq. (9), we have

$$\mathbb{P}(0 \leq \Phi_n(a) \leq 1 - \epsilon) \leq \mathbb{P}(|\Phi_n(a) - 1| \geq \epsilon) \leq \frac{1}{\epsilon^2 snh_n^d},$$

which implies

$$\begin{aligned}\mathbb{P}\left(\min_{1 \leq a \leq m} \Phi_n(a) \leq 1 - \epsilon\right) &= \mathbb{P}\left(\bigcup_{a=1}^m \{\Phi_n(a) \leq 1 - \epsilon\}\right) \\ &\leq \sum_{a=1}^m \mathbb{P}(\Phi_n(a) \leq 1 - \epsilon) \leq \frac{m}{\epsilon^2 s n h_n^d},\end{aligned}$$

and

$$\begin{aligned}\mathbb{P}\left(\max_{1 \leq a \leq m} \frac{k^*}{\beta \Phi_n(a) n p(X_{n+a})} \leq \frac{k^*}{\beta(1 - \epsilon) n p(X_{n+a})}\right) \\ \geq 1 - \frac{m}{\epsilon^2 s n h_n^d}.\end{aligned}$$

Since $\frac{m}{\epsilon^2 s n h_n^d} \rightarrow 0$, we have

$$\begin{aligned}\mathbb{P}\left(\max_{1 \leq a, b \leq m} (\mathbf{D}_{22}^{-1} \mathbf{W}_{22})_{ab} \leq \max_{1 \leq a \leq m} \frac{k^*}{\beta \Phi_n(a) n p(X_{n+a})}\right. \\ \left. \leq M \frac{1}{n h_n^d}\right) \rightarrow 1, \quad \text{as } n \rightarrow \infty, \quad (10)\end{aligned}$$

where $M = \frac{2k^*}{s\beta} > \frac{k^*}{(1-\epsilon)s\beta}$. Note that M is a constant independent with n and m .

For the sake of simplicity, we say a matrix $\mathbf{A}$ has *tiny elements*, if

$$\|\mathbf{A}\|_{\max} \leq M \frac{1}{n h_n^d},$$

with probability approaching 1, where $\|\mathbf{A}\|_{\max} = \max_{ij} \mathbf{A}_{ij}$. $(\mathbf{A})_i$ denotes the i -th row of $\mathbf{A}$. Thus, $\mathbf{D}_{22}^{-1} \mathbf{W}_{22}$ has tiny elements by Eq. (10). Moreover,

$$\begin{aligned}\|(\mathbf{D}_{22}^{-1} \mathbf{W}_{22})^2\|_{\max} &= \|(\mathbf{D}_{22}^{-1} \mathbf{W}_{22})(\mathbf{D}_{22}^{-1} \mathbf{W}_{22})\|_{\max} \\ &\leq (M \frac{1}{n h_n^d})^2 m = \frac{M}{n h_n^d} (\frac{mM}{n h_n^d})\end{aligned}$$

holds with probability approaching 1. By induction,

$$\begin{aligned}\|(\mathbf{D}_{22}^{-1} \mathbf{W}_{22})^l\|_{\max} &= \|(\mathbf{D}_{22}^{-1} \mathbf{W}_{22})(\mathbf{D}_{22}^{-1} \mathbf{W}_{22})^{l-1}\|_{\max} \\ &\leq \frac{M}{n h_n^d} (\frac{mM}{n h_n^d})^{l-1},\end{aligned}$$

with probability approaching 1. Therefore,

$$\begin{aligned}\|\mathbf{S}_l\|_{\max} &= \|\mathbf{D}_{22}^{-1} \mathbf{W}_{22} + \cdots + (\mathbf{D}_{22}^{-1} \mathbf{W}_{22})^l\|_{\max} \\ &\leq \|\mathbf{D}_{22}^{-1} \mathbf{W}_{22}\|_{\max} + \cdots + \|(\mathbf{D}_{22}^{-1} \mathbf{W}_{22})^l\|_{\max} \\ &\leq \frac{M}{n h_n^d} \left(1 + \cdots + (\frac{mM}{n h_n^d})^{l-1}\right)\end{aligned}$$

with probability approaching 1.

$$\begin{aligned}\lim_{l \rightarrow \infty} \|\mathbf{S}_l\|_{\max} &\leq \lim_{l \rightarrow \infty} \frac{M}{n h_n^d} \left(1 + \cdots + (\frac{mM}{n h_n^d})^{l-1}\right) \\ &\leq \frac{M}{n h_n^d} / (1 - \frac{mM}{n h_n^d}) \leq \frac{2M}{n h_n^d}\end{aligned}$$

with probability approaching 1.

Thus, $\mathbf{S} \triangleq \lim_{l \rightarrow \infty} \mathbf{S}_l$ exists with probability approaching 1 since $\lim_{l \rightarrow \infty} \|\mathbf{S}_l\|_{\max} < \infty$, and $\mathbf{S}$ also has tiny elements. Therefore,

$$\begin{aligned}(\mathbf{D}_{22} - \mathbf{W}_{22})^{-1} &= (\mathbf{I}_m - \mathbf{D}_{22}^{-1} \mathbf{W}_{22})^{-1} \mathbf{D}_{22}^{-1} \\ &= (\mathbf{I}_m + \mathbf{S}) \mathbf{D}_{22}^{-1},\end{aligned}$$

with probability approaching 1.

We now go back to the solution of the hard criterion of graph-based semi-supervised learning,

$$\begin{aligned}\hat{\mathbf{f}}_{(n+1):(n+m)} &= (\mathbf{D}_{22} - \mathbf{W}_{22})^{-1} \mathbf{W}_{21} \mathbf{Y}_n \\ &= (\mathbf{I}_m + \mathbf{S}) \mathbf{D}_{22}^{-1} \mathbf{W}_{21} \mathbf{Y}_n \\ &= \mathbf{D}_{22}^{-1} \mathbf{W}_{21} \mathbf{Y}_n + \mathbf{S} \mathbf{D}_{22}^{-1} \mathbf{W}_{21} \mathbf{Y}_n,\end{aligned}$$

with probability approaching 1. For $1 \leq a \leq m$, $\hat{f}_{(n+a)}$ equals to the a th row of $(\mathbf{D}_{22} - \mathbf{W}_{22})^{-1} \mathbf{W}_{21} \mathbf{Y}_n$, i.e.,

$$\begin{aligned}\hat{f}_{(n+a)} &= ((\mathbf{D}_{22} - \mathbf{W}_{22})^{-1} \mathbf{W}_{21} \mathbf{Y}_n)_a \\ &= \sum_{i=1}^n \frac{w_{i,n+a}}{d_{n+a,n+a}} Y_i + (\mathbf{S})_a \mathbf{D}_{22}^{-1} \mathbf{W}_{21} \mathbf{Y}_n,\end{aligned}$$

with probability approaching 1, where $(\mathbf{S})_a$ denotes the a th row of $\mathbf{S}$.

By assumption, Y_i s are bounded. Without loss of generality, assume $\|Y_n\|_{\max} \leq 1$. For $1 \leq a \leq m$, define

$$g_{(n+a)} = \sum_{i=1}^n Y_i \left(\frac{w_{i,n+a}}{\sum_{k=1}^n w_{k,n+a}} - \frac{w_{i,n+a}}{d_{n+a,n+a}} \right).$$

We have

$$\begin{aligned}|g_{(n+a)}| &\leq \sum_{i=1}^n \|Y_n\|_{\max} \left(\frac{w_{i,n+a}}{\sum_{k=1}^n w_{k,n+a}} - \frac{w_{i,n+a}}{d_{n+a,n+a}} \right) \\ &= \frac{\sum_{i=1}^n w_{i,n+a}}{\sum_{k=1}^n w_{k,n+a}} - \frac{\sum_{i=1}^n w_{i,n+a}}{\sum_{k=1}^{n+m} w_{k,n+a}} \\ &= \frac{\sum_{k=n+1}^{n+m} w_{k,n+a}}{d_{n+a,n+a}} \\ &\leq \frac{m k^*}{\beta \Phi_n(a) n p(X_{n+a})} \leq \frac{mM}{n h_n^d} \rightarrow 0,\end{aligned}$$

with probability approaching 1 as $n \rightarrow \infty$. This implies

$$g_{(n+a)} \rightarrow 0 \text{ in probability,}$$

since for any $\epsilon > 0$ we can find $m, n \in \mathbb{N}$ such that $\frac{mM}{n h_n^d} \leq \epsilon$ and

$$\mathbb{P}(|g_{(n+a)}| \leq \epsilon) \geq \mathbb{P}\left(|g_{(n+a)}| \leq \frac{mM}{n h_n^d}\right) \rightarrow 1.$$

Finally, for each $1 \leq a \leq m$,

$$\begin{aligned}\hat{f}_{(n+a)} &= \sum_{i=1}^n \frac{w_{i,n+a}}{d_{n+a,n+a}} Y_i + (\mathbf{S})_a \mathbf{D}_{22}^{-1} \mathbf{W}_{21} \mathbf{Y}_n \\ &= \sum_{i=1}^n \frac{w_{i,n+a}}{\sum_{k=1}^n w_{k,n+a}} Y_i + (\mathbf{S})_a \mathbf{D}_{22}^{-1} \mathbf{W}_{21} \mathbf{Y}_n \\ &\quad - g_{(n+a)},\end{aligned}$$

Since $\mathbf{S}$ has tiny elements,

$$\|(\mathbf{S})_a \mathbf{D}_{22}^{-1} \mathbf{W}_{21} \mathbf{Y}_n\| \leq \frac{mM}{nh_n^d} \rightarrow 0$$

with probability approaching 1, which implies $(\mathbf{S})_a \mathbf{D}_{22}^{-1} \mathbf{W}_{21} \mathbf{Y}_n \rightarrow 0$ in probability. The theorem then holds by the consistency of Nadaraya-Watson estimator.

V. EXPERIMENTS

A. Synthetic Data

In this sub-section, we compare the performance of the hard and soft criteria with different tuning parameters under a linear and non-linear model.

The inputs $X_1, \dots, X_{n+m}$ are generated independently from a truncated multivariate normal distribution. Specifically, let $\tilde{X}_i$ follow a p -dimensional multivariate normal with the mean $\mu = (0.5, \dots, 0.5)$ and the variance-covariance matrix

$$\begin{pmatrix} 0.1 & 0.05 & 0.05 & \dots & 0.05 \\ 0.05 & 0.1 & 0.05 & \dots & 0.05 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0.05 & 0.05 & 0.05 & \dots & 0.1 \end{pmatrix}.$$

We set $p = 5$. For $i = 1, \dots, n+m$ and $k = 1, \dots, p$, let $X_{ik} = \tilde{X}_{ik}$ if $\tilde{X}_{ik} \in [0, 1]$ and $\tilde{X}_{ik} = 0$ otherwise, where X_{ik} and $\tilde{X}_{ik}$ are the k -th component of X_i and $\tilde{X}_i$, respectively.

Let $\mathbf{W}$ be the Gaussian radial basis function (RBF) kernel with $\sigma = h_n = (\log n/n)^{1/5}$. Note that $\mathbf{W}$ has compact support since X_i are truncated and the choice of h_n satisfies the condition in Theorem II.1.

We consider two models in this study. In Model 1, the responses Y_i follow a logistic regression with

$$\begin{aligned}\text{logit } q(X_i) &= -1.35 + 2X_{i1} - X_{i2} + X_{i3} - X_{i4} \\ &\quad + 2X_{i5},\end{aligned}\tag{11}$$

for $i = 1, \dots, m+n$. Model 2 uses a non-linear logit function,

$$\begin{aligned}\text{logit } q(X_i) &= -1.35 + 2X_{i1} - X_{i2} + X_{i3} - X_{i4} \\ &\quad + 2X_{i5} + X_{i1}X_{i3} + X_{i2}X_{i4},\end{aligned}$$

for $i = 1, \dots, m+n$.

We compare the performance of graph-based learning methods with four different tuning parameters, $\lambda =$

0, 0.01, 0.1 and 5. Performance is measured by the root mean squared error (RMSE) on the unlabeled data:

$$\sqrt{\frac{1}{m} \sum_{a=1}^m (q(X_{n+a}) - \hat{q}_{n+a})^2}.$$

Each simulation is repeated 1000 times and the average RMSEs are reported.

Figure 1 shows the RMSEs under Model 1 when the sample size of unlabeled data m is fixed as 30 and the sample size of labeled data $n = 10, 30, 50, 100, 200, 300, 500, 800, 1000$ and 1500. As n increases, the RMSEs of all methods decrease as expected. More importantly, the RMSE increases as λ increases. In particular, the hard criterion always outperforms the soft criterion, which is in line with our theoretical results.

Figure 2 shows the RMSEs under Model 1 when n is fixed as 100 and $m = 30, 60, 100, 300, 500$ and 1000. As before, the RMSE increases as λ increases. Moreover, the RMSEs of all methods increase as m increases, which suggests that the hard criterion may not be consistent when m grows faster than n although the hard criterion still performs constantly better. For a non-linear logit function, Figure 3 and 4 show the same patterns as in Figure 1 and 2, which further supports our theoretical results.




Fig. 1. Average RMSEs for $m = 30$ under Model 1

B. The Columbia Object Image Library Data

We test the performance of the hard and soft criteria on the Columbia object image library dataset compiled by [1], which is listed in the Chapter 21 of the book as a benchmark. The dataset contains color images of 24 different objects taken from 72 different angles. These subjects were classified into six classes and the authors randomly discarded 38 images of each class, leaving 250 each, i.e., 1500 samples in total. The author also created a binary version of this data, which groups the first three and last three together,




Fig. 2. Average RMSEs for $n = 100$ under Model 1




Fig. 3. Average RMSEs for $m = 30$ under Model 2




Fig. 4. Average RMSEs for $n = 100$ under Model 2

respectively, leaving two classes. We use this binary dataset to test the hard and soft criteria. The inputs X were created from 16×16 pixels of each image. We use the Gaussian RBF kernel as $\mathbf{W}$ with σ^2 being the median of squared distances between each pair of inputs.

When responses are binary, the RMSE cannot be used to measure the performance of classification algorithms in real dataset because the true probability $\mathbb{E}[Y_i|X_i]$ is unknown (for continuous responses, the root mean square prediction error can be used). Instead, we use the area under the receiver operating characteristic curve (AUC) to measure the performance. The receiver operating characteristic (ROC) curve is obtained via plotting the sensitivity (true positive rate) versus $1 - \text{specificity}$ (false positive rate) of the classification.

We vary the ratio between the training/labeled and test/unlabeled sets and prepare the data sets in the following way: in the first setting, we randomly split the data into five subsets of approximately equal size. We then use each subset as the test set and the rest four as the training set. In this way, every part of the data has the chance to be predicted in the experiment. We repeat the above procedure 100 times and thus the reported results are based on the average of 500 experiments. In each experiment, we compare the criteria on seven different tuning parameters, $\lambda = 0, 0.01, 0.05, 0.1, 0.5, 1$ and 5 .

In the second and the third settings, we follow the same procedure as above but use only 20% and 10% of the data as labeled samples, respectively. Specifically, we randomly split the data into five subsets of approximately equal size and use one subset as the training set and the rest four as the test set in the second setting. In the third setting, we split the data into ten subsets and use one as the training and nine as the test. We repeat the above procedure 100 times in each setting. The reported results are thus based on the average of 500 experiments as before in the second setting but are the average of 1000 experiments in the last setting since we split the data into ten subsets under this scenario.

According to Figure 5 the hard criterion ($\lambda = 0$) constantly gives the best AUC on all combinations of the scales of the labeled and unlabeled data. The AUC decreases as λ increases in general, although the difference between the AUCs for $\lambda = 1$ and $\lambda = 5$ are negligible. Moreover, the AUC decrease as the proportion of labeled data decreases as expected. The pattern is consistent with that of RMSEs for the synthetic data and the theoretical results.

VI. SUMMARY

In this article, we proved the consistency of graph-based semi-supervised learning when the tuning parameter of the graph Laplacian is zero (the hard criterion) and showed that the method can be inconsistent when the tuning parameter is nonzero (the soft criterion). Moreover, the numerical studies also suggest that the hard criterion outperforms the




Fig. 5. Average AUCs for the COIL data

soft criterion in terms of RMSE and AUC. These results provide a better understanding about the statistical properties of graph-based semi-supervised learning. It suggests that practitioners can safely choose the hard criterion ($\lambda = 0$) with no need for tuning λ in the soft criterion.

For future work, we plan to investigate the theoretical properties of other indicators of prediction accuracy such as AUC and MCC (Matthews correlation coefficient) in more depth. The asymptotic properties of these indicators in the setting of semi-supervised learning remains unknown. Moreover, we would also like to investigate the behavior of graph-based semi-supervised learning when the unlabeled data grow faster than the label data. The numerical results suggest that the hard criterion may not be consistent when the size of labeled data grows faster than the size of unlabeled data although the hard criterion still performs constantly better. A theoretical comparison between the two criteria is intriguing under this scenario.

REFERENCES

- [1] O. Chapelle, B. Schölkopf, and A. Zien, *Semi-supervised Learning*. The MIT Press, 2006.
- [2] J. Ratsaby and S. S. Venkatesh, "Learning from a mixture of labeled and unlabeled examples with parametric side information," In *Proceedings of COLT '95 Proceedings of the eighth annual conference on Computational learning theory*, pp. 412–417, 1995.
- [3] C. Rosenberg, M. Hebert, and H. Schneiderman, "Semi-supervised self-training of object detection models," In *Proceedings of Seventh IEEE Workshop on Applications of Computer Vision*, 2005.
- [4] R. Jones, "Learning to extract entities from labeled and unlabeled text," *PhD Thesis*, 2005.
- [5] Y. Zhang, M. Brady, and S. Smith, "Hidden markov random field model and segmentation of brain mr images," *IEEE Transactions on Medical Imaging*, vol. 20(1), pp. 45–57, 2001.
- [6] S. Basu, A. Banerjee, and R. J. Mooney, "Semi-supervised clustering by seeding," In *Proceedings of the International Conference on Machine Learning*, pp. 19–26, 2002.
- [7] V. Vapnik, *Statistical Learning Theory*. Wiley, New York, 1998.
- [8] T. D. Bie and N. Cristianini, "Convex methods for transduction," In *Advances in Neural Information Processing Systems*, vol. 16, pp. 73–80, 2004.
- [9] X. Zhu, Z. Ghahramani, and J. Lafferty, "Semi-supervised learning using Gaussian Fields and Harmonic Functions," *ICML*, no. 118, 2003.
- [10] O. Delalleau, Y. Bengio, and N. L. Roux, "Efficient non-parametric function induction in semi-supervised learning," In *Artificial Intelligence and Statistics*, 2005.
- [11] X. Zhu and A. B. Goldberg, *Introduction to Semi-supervised Learning*, Morgan & Claypool Publishers, 2009.
- [12] D. Zhou, O. Bousquet, T. N. Lal, J. Weston, and B. Schölkopf, "Learning with local and global consistency," in S. Thrun, L. Saul, and B. Schölkopf, editors, *MIT Press, Cambridge, MA*, 2004.
- [13] M. Belkin, I. Matveeva, and P. Niyogi, "Regularization and semi-supervised learning on large graphs," In *Proceedings of the Seventeenth Annual Conference on Computational Learning Theory*, pp. 624–638, Banff, Canada, 2004.
- [14] O. Bosquet, O. Chapelle, and M. Hein, "Measure based regularization," *NIPS*, vol. 16, 2004.
- [15] M. Hein, "Uniform convergence of adaptive graph-based regularization," *COLT: Learning Theory*, pp. 50–64, 2006.
- [16] M. Belkin, P. Niyogi, and S. Sindhvani, "Manifold Regularization: A geometric framework for learning from labeled and unlabeled examples," *JMLR*, vol. 7, pp. 2399–2434, 2006.
- [17] B. Nadler, N. Srebro, and X. Zhou, "Semi-supervised learning with the Graph Laplacian: The limit of infinite unlabelled data," *NIPS*, 2009.
- [18] J. Lafferty and L. Wasserman, "Statistical analysis of semi-supervised regression," *NIPS*, vol. 20, 2008.
- [19] A. E. Alaoui, X. Cheng, A. Ramdas, M. J. Wainwright, and M. I. Jordan, "Asymptotic behavior of ℓ_p -based Laplacian regularization in semi-supervised learning," in *29th Annual Conference on Learning Theory*, vol. 49. Columbia University, New York, New York, USA: Proceedings of Machine Learning Research, 23–26 Jun 2016, pp. 879–906.
- [20] E. A. Nadaraya, "On estimating regression," *Theor. Probability Appl.*, vol. 9, pp. 141–142, 1964.
- [21] G. S. Watson, "Smooth regression analysis," *Sankhyā, Series A*, vol. 26, pp. 359–372, 1964.
- [22] M. E. J. Newman, *Networks: An introduction*. Oxford University Press, 2010.
- [23] L. P. Devroye, "The uniform convergence of the Nadaraya-Watson regression function estimate," *The Canadian Journal of Statistics*, vol. 6, pp. 179–191, 1978.
- [24] Z. Cai, "Weighted Nadaraya Watson regression estimation," *Statistics & Probability Letters*, vol. 51, pp. 307–318, 2001.
- [25] L. P. Devroye and T. J. Wagner, "Distribution-free consistency results in nonparametric discrimination and regression function estimation," *Annals of Statistics*, vol. 8, no. 2, pp. 231–239, 1980.
- [26] D. Werner, *Functional Analysis (in German)*. Springer Verlag, 2005.