

On the Apparent Ambiguities in the Post-Newtonian Expansion for Binary Systems

Rafael A. Porto¹ and Ira Z. Rothstein²

¹ *ICTP South American Institute for Fundamental Research
Rua Dr. Bento Teobaldo Ferraz 271, 01140-070 São Paulo, SP Brazil*

² *Department of Physics and Astronomy
Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

We discuss the source of the apparent ambiguities arising in the calculation of the dynamics of binary black holes within the Post-Newtonian framework. Divergences appear in both the near and far zone calculations, and may be of either ultraviolet (UV) or infrared (IR) nature. The effective field theory (EFT) formalism elucidates the origin of the singularities which may introduce apparent ambiguities. In particular, the only (physical) ‘ambiguity parameters’ that necessitate a matching calculation correspond to unknown finite size effects, which first appear at fifth Post-Newtonian (5PN) order for non-spinning bodies. We demonstrate that the ambiguities linked to IR divergences in the near zone, that plague the recent derivations of the binding energy at 4PN order, both in the ADM and ‘Fokker-action’ approach, can be resolved by implementing the so-called *zero-bin* subtraction in the EFT framework. The procedure yields ambiguity-free results without the need of additional information beyond the PN expansion.

1 Introduction

The calculation of high accuracy Post-Newtonian (PN) templates for binary inspirals (for extensive reviews see [1–3]) plays a key role in the data analysis program in this nascent era of gravitational wave (GW) science [4–8]. The PN formalism is a systematic expansion in the relative velocity, v , of the constituents during the early stages of the inspiral, for which $v/c \ll 1$. The expansion parameter also serves to separate out the relevant length scales in the problem: The orbital radius r and the wavelength of the radiation $\lambda_{\text{rad}} \sim r/v$. In addition, there is a third scale, namely the typical radius of the constituents, R , which is, in general, independent of r and v . The starting point of a perturbative approach corresponds to a multipole expansion where extended objects, such as neutron stars or black holes, are treated as localized sources. In principle, this is logically independent of the PN expansion, and it is valid even in the relativistic limit, as long as the typical wavelength (or frequency) of the perturbation, λ , obeys $\lambda \gg R$. For a bound state of compact objects ($R \sim 2G_N m$)¹, the virial theorem relates the size of the bodies to the orbital radius, $R/r \sim v^2$, and therefore all relevant scales in the problem are organized in powers of v .

In the PN formalism, after the constituents are replaced by point-like sources, the calculations are then separated into *regions*, the near (or potential) zone entailing (instantaneous) modes of the gravitational field varying on the scale of the bound state radius, and the far (or radiation) zone, where modes propagate ($p^0 \sim |\mathbf{p}|$) with the typical wavelength of the GW emission. The calculations are performed in the two regions—in principle—independently. A *matching* procedure is required to read off the relevant parameters entering in the GW amplitude and phase in the radiation region (multipole moments) in terms of quantities from the near zone.

While the full computation within general relativity (GR)—using extended bodies and without separating into zones—is devoid of UV divergences, there may be IR singularities, which are due to the long range nature of the Newtonian potential. For the computation of the GW amplitude, the IR sensitivity has two sources: The unknown value of the phase as the GW enters the detectors’ band, and the resulting phase shift associated with scattering off of a long range potential. The former is absorbed into an initial conditions and the latter cancels out in the total radiated power.² On the other hand, the approximation scheme in which we split the calculation into regions generates new singular integrals. These additional divergences may be either of IR or UV origin. The latter appear as a consequence of the point-particle approximation, whereas the former may include spurious poles which are not present in the full theory calculations.

The effective field theory (EFT) framework (NRGR) [9] is particularly useful to, not only organize the computation systematically, but also to elucidate the origin of IR and UV divergences, and consistently eliminate them. For reviews of the EFT approach see [10–15]. In the EFT the near/far zone separation is accomplished at the level of the action, by splitting the gravitational field into potential and radiation modes, such that each term scales homogeneously in v . Since all observables are computed in terms of Feynman diagrams, whose vertices are fixed by the effective action, it is a simple matter to determine which ones contribute at any given order in the expansion parameter.

Near zone calculations, which involve the potential field, may contain UV divergences due to the point-particle approximation, and can be removed by *counter-terms* that include all possible terms in the action compatible with the symmetries of the problem. In this sense, the removal of classical divergences is identical to renormalization in quantum field theories [16,17]. It can be shown that, up to order

¹For objects other than black holes, the compactness: $c \equiv R/(2G_N m)$ depends on the equation of state. Therefore, while finite size effects will be organized in powers of v , there could be large coefficients, e.g. c^5 for neutron stars, that may enhance the naive power counting.

²Let us emphasize a cancellation occurs for physical observables, and may not be the case for gauge-dependent, unphysical, quantities.

$(R/r)^5 \sim v^{10}$ (or 5PN), the only unknown parameters (in addition to G_N) are the masses of constituents, for non-rotating bodies. This is known as the effacement theorem [18]. In the EFT formalism, this result follows from the fact that the first term –beyond minimal coupling– that one can write down in the worldline effective action is proportional to the square of the electric and magnetic components of the Weyl tensor [9]. The coefficients are known as the Love numbers [12, 19]. (For spinning objects [20], on the other hand, finite size effects appear already at 2PN order [21–23]. We will only deal with non-rotating bodies in what follows.) Thus, from the point of view of renormalization, there can be no physically relevant divergences until 5PN order, or in other words, no ‘ambiguity parameters’ are needed until this order. At 5PN order, after the divergence is subtracted away, the remaining finite part is chosen such that it matches the physical value of the Love number for the underlying constituent. Moreover, the divergence must be UV in origin, since it arises from the short-distance region of the Feynman integral, which in the full theory is cut off by the size of the constituents. While the subtraction scheme is arbitrary, the physical results are not, and the independence from the choice of renormalization scale translates into the renormalization group evolution of the coefficients of the effective theory [9, 15].

In the far zone we also encounter UV divergences as a consequence of the multipole expansion in which the binary (as a whole) is also replaced by a point-like source. In that case, a counter-term is needed which is proportional to the (time-dependent) multipole moments of the binary [24]. However, the previously alluded to matching condition takes care of the ambiguity in the remaining finite part of the (renormalized) multipoles, since these are obtained in terms of masses, positions, velocities, etc., of the constituents. (Because of the non-linearities of GR, the binding energy of the system also contributes to the source multipoles.) On the other hand, because the effective theory is constructed to agree with the full theory in the IR, IR divergences in the far zone will be identical to the ones in the complete GR computation, and consequently cancel out [24, 25].

The near zone, on the other hand, may introduce spurious IR divergences if the split into regions is not handled with care. In principle, IR singularities cannot develop with potential modes. This would signal a region of momenta in the Feynman integrals which is much softer than what is allowed –by definition– in the near zone. In other words, the near region should be cut off where the far zone begins. Hence, the IR divergences in the computation of the binding potential cannot have a physical origin. The solution to this problem is to remove from the integrals in the near zone the contribution from Fourier modes with wavelengths larger than r . The procedure goes by the name of the *zero-bin* subtraction [26]. The purpose of this note is to demonstrate how to implement the latter in the PN framework, without the need of ambiguity parameters. The logic we are about to describe holds to all orders in the PN expansion. As an explicit example we will consider the conservative binding energy at 4PN order, where the spurious IR divergences in the near zone first appear [27–33]. An illustrative example –the derivation of the Lamb shift in an EFT framework– is presented in full in a companion paper, where similar considerations apply [34].

2 The zero-bin subtraction

The calculation of the binding energy in full-fledged GR is finite, therefore the IR/UV divergences appear as a result of the approximation scheme. The above mentioned IR singularities that arises in near zone calculations is a signal of double-counting, since modes which are thought to be confined to the bound state are overlapping with the radiation region. Therefore, a consistent methodology must ensure that spurious IR divergences are absent. In the EFT formalism such a methodology entails including “zero-bin” subtractions [26]. We sketch the basic steps here, see [26] for more details.

The binding energy in NRGR is determined by convolutions of Green’s functions of potential and radiation modes. Hence, if we denote by V_{full} the full gravitational potential, we have

$$V_{\text{full}} = V_{\text{pot}} + V_{\text{rad}} , \quad (2.1)$$

where V_{pot} is the potential mode contribution in NRGR, and V_{rad} is the conservative part of the tail corrections in the radiation theory. In principle, at a given n PN order we will have a result of the form

$$V_{\text{pot}}^{(n)}(\mathbf{q}) = \frac{A_{\text{pot}}^{(n)}}{\epsilon_{\text{UV}}} + \frac{B_{\text{pot}}^{(n)}}{\epsilon_{\text{IR}}} + f_{\text{pot}}^{(n)}(\mathbf{q}, c_i^{(n)}, \mu), \quad (2.2)$$

with $\mathbf{q} \sim 1/r$, and $\epsilon_{\text{UV/IR}} \equiv (d-4)_{\text{UV/IR}}$ in dimensional regularization (dim. reg.), which we use to regularize the divergent integrals. The factors of μ appear also from dim. reg. since, in $d \neq 4$, Newton's constant acquires an extra mass scale proportional to μ^{d-4} . The $A_{\text{pot}}^{(n)}$ term can be either set to zero by a coordinate transformation for $n \leq 4$; or removed by a counter-term, $(c_i^{(n)})_{\text{c.t.}} \propto 1/\epsilon_{\text{UV}}$, using the coefficients of the effective action (beyond minimal coupling) for $n \geq 5$. The finite part, encoded in $c_i^{(n)}(\mu)$, accounts for the finite size effects starting at 5PN order. The μ -independence of physical results leads to a renormalization group evolution for these coefficients [9, 15]. By explicit calculation it is known that $B^{(n)} = 0$ for $n \leq 3$. On the other hand, the far zone contribution to the conservative part of the potential will be of the form³

$$V_{\text{rad}}^{(n)}(\omega) = \frac{A_{\text{rad}}^{(n)}}{\epsilon_{\text{UV}}} + \frac{B_{\text{rad}}^{(n)}}{\epsilon_{\text{IR}}} + f_{\text{rad}}^{(n)}(\omega, \mu), \quad (2.3)$$

with $\omega \sim v/r$. For our purposes we will take $B_{\text{rad}}^{(n)} = 0$, since IR divergences do not appear in the conservative part of the radiation sector. From the computation of the tail effect we find $A_{\text{rad}}^{(n)} \neq 0$ for $n \geq 4$, and $A_{\text{rad}}^{(n)} = -B_{\text{pot}}^{(n)}$. This relationship is key in the cancellation of the spurious IR/UV divergences in the far/near zones from the conservative sector [35].

The role of the zero-bin subtraction is to remove the IR divergent contribution from the potential region, and transform it into the required UV pole to cancel the one arising from the far zone. In other words, we must shift

$$V_{\text{pot}} \rightarrow V_{\text{pot}} - V_{\text{zero-bin}}, \quad (2.4)$$

where $V_{\text{zero-bin}}$ corresponds to an asymptotic expansion of the integral around the region responsible for the IR singularity. This procedure removes the double counting induced by the overlap between the IR-sensitive part of V_{pot} and the contribution from V_{rad} . The zero-bin part may involve a scaleless integral in dim. reg. These are usually set to zero, however, the procedure is more subtle for integrals which are simultaneously IR and UV divergent. That is because we need to isolate the UV and IR divergences, which otherwise could be incorrectly removed. For example, consider the following scaleless integral in the limit $d \rightarrow 4$,

$$I = \int \frac{d^d k}{k^4}, \quad (2.5)$$

which is clearly both IR and UV sensitive. One can easily manipulate the integrand and re-write it as IR and UV divergent parts, with M some generic scale,

$$I = \int \frac{d^d k}{k^2 - M^2} \frac{1}{k^2} - \int \frac{d^d k}{k^2 - M^2} \frac{M^2}{k^4}. \quad (2.6)$$

To regularize the integral, in the first term we use a UV regularization, $\epsilon_{\text{UV}} < 0$, whereas in the second we chose $\epsilon_{\text{IR}} > 0$, leading to

$$I \propto \left(\frac{1}{\epsilon_{\text{UV}}} - \frac{1}{\epsilon_{\text{IR}}} \right). \quad (2.7)$$

The zero-bin is often of this form, i.e.

$$V_{\text{zero-bin}}^{(n)} = B_{\text{pot}}^{(n)} \left(\frac{1}{\epsilon_{\text{IR}}} - \frac{1}{\epsilon_{\text{UV}}} \right), \quad (2.8)$$

³In principle we also need to incorporate the running of the multipole moments, e.g. $I^{ij}(\mu)$, see [24]. However, these effects—which do not introduce ambiguities—will enter at much higher orders. We suppress their contribution here for simplicity.

such that the IR part is precisely what we need to cancel the pole from the potential region. At the same time, the remaining UV pole readily removes the divergence from the far zone. Hence, after the zero-bin is incorporated, we have

$$V_{\text{pot}}^{(n)} - V_{\text{zero-bin}}^{(n)} + V_{\text{rad}}^{(n)} \rightarrow V_{\text{tot}}^{(n)}(\mathbf{q}, \omega, c_i^{(n)}(\mu), \mu), \quad (2.9)$$

where the $c_i^{(n)}(\mu)$ are the only parameters which require a matching computation beyond the PN framework, first entering at 5PN order according to the effacement theorem [9, 15, 18]. For the case $n \leq 4$, the final expression is not only finite, but also all the $\log \mu$ terms –associated to either IR or UV poles– cancel out (or are removed by a coordinate transformation). This is the situation we encounter at 4PN order, we describe below. We point the reader to [26] for a more elaborate description of the zero-bin procedure.

3 The conservative dynamics at 4PN order

The complete conservative dynamics of a gravitationally bound (non-spinning) two-body system has been recently determined at 4PN order in [27–33]. However, the calculations thus far have been plagued by ambiguity parameters. Unlike previous ones at 3PN order due to UV singularities, which may be removed by coordinate transformations [1] (or field redefinitions in the EFT language [9, 15]), on this occasion the ambiguities are due to the scheme-dependence introduced in [27–33] to handle the presence of IR divergences. Hence, the exact form of the analytic contribution to the gravitational binding energy at leading order in the symmetric mass-ratio, $\nu = \frac{m_1 m_2}{(m_1 + m_2)^2}$, has been derived after a comparison with gravitational self-force results was made [29, 32], see also [36]. However, one should not have to rely on calculations outside the scope of the PN expansion to remove spurious divergences which are due to the approximation scheme. (If for no other reason than it requires unnecessary extra work.) In fact, this issue will re-emerge at higher PN orders, involving IR-sensitive contributions at $\mathcal{O}(\nu^k)$, with $k \geq 2$, for which self-force computations may not be available. The zero-bin subtraction [26], on the other hand, removes the need of ambiguity parameters altogether, yielding IR-safe quantities. We illustrate the procedure for the calculation of the 4PN gravitational potential below.

3.1 The ambiguities

In [27, 28], where the calculation was performed using dim. reg. in ADM gauge, the regularized gravitational potential contains an IR pole,⁴

$$\int dt V_{4\text{PN}}^{JS}(t, \mu) = -\frac{G_N^2 M}{5} \int \frac{d\omega}{2\pi} \omega^6 I^{ij}(-\omega) I^{ij}(\omega) \left(\frac{1}{\epsilon_{\text{IR}}} - 2 \log(\mu r) \right) + \text{local/finite}, \quad (3.1)$$

We may attempt to remove the IR divergence from the near zone calculation, by subtracting it away. However, this is not only misguided, it is also totally arbitrary since one could subtract either $1/\epsilon_{\text{IR}}$ or $1/\epsilon_{\text{IR}} + \text{constant}$. In general, different subtraction schemes, or even choices of regulators other than dim. reg., e.g. graviton-mass, momentum cut-off, Hadamard, etc., would produce a non-unique result. Therefore, the final answer would be ambiguous and the 4PN near zone Hamiltonian would depend on a regularization scale, μ , as well as an undetermined dimensionless constant, C , through (the factor of $2/5$ is a convention)

$$V_{(C)}^{JS} \equiv \frac{2}{5} C G_N^2 M \left(I^{ij(3)}(t) \right)^2. \quad (3.2)$$

The same type of ambiguities (in principle more than one) were introduced in the computation of the Fokker-action in harmonic gauge [31, 32]. The existence of these extra parameter(s) was also discussed in

⁴As it turns out, the derivation in [27, 28] uses a combination of regularization methods before matching the calculation into the far zone later on in [29]. This exacerbates the ambiguous nature of the computation.

more detail in [30]. In the ADM and Fokker-action approaches, the value of C is fixed after importing knowledge from an independent calculation. Since the ambiguity shows up at leading order in ν , the constant C (in principle more than one) was fixed by comparison with existent self-force calculations [29,32].

On the other end, the contribution to the gravitational potential at 4PN order from radiation-reaction was computed within the EFT approach in [35],

$$\int dt V_{\text{tail}}(\mu) = \frac{G_N^2 M}{5} \int \frac{d\omega}{2\pi} \omega^6 I^{ij}(-\omega) I^{ij}(\omega) \left[\frac{1}{\epsilon_{\text{UV}}} + \log \frac{\omega^2}{\mu^2} + \text{finite} \right]. \quad (3.3)$$

We notice that, instead of IR issues, we encounter a UV divergence. The alert reader will immediately realize that the IR and UV near/far zone poles in (3.1) and (3.3) have –up to a crucial sign– the same coefficients [35], provided the IR divergence from [27,28] is reproduced in NRGR (see below).⁵ In our language, an expression like (3.2) would arise if we were to use two different subtraction scales for the IR and UV poles (i.e. μ_{IR} and μ_{UV}), yielding an ambiguity parameter, $C \propto \log(\mu_{\text{UV}}/\mu_{\text{IR}})$ [35]; however, this would be inconsistent.⁶ We show in what follows how to unify the treatment of the potential and radiation singularities in the EFT formalism.

3.2 Infrared divergences in the EFT framework

While the UV pole in the conservative sector from the tail effect identified in [35] was expected, the IR divergences in the potential region have not been yet fully isolated. Hence, we must first identify the terms from the near zone which yield IR poles at 4PN order. There are many diagrams which contribute to the gravitational potential at this order [39,40]. One such term is depicted in fig. 1. Following the representation in [40], the integral may be written as

$$\text{fig. 1} = m_1^2 m_2^3 \int_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3} \frac{\text{Num}(\mathbf{k}_i, \mathbf{p})}{\mathbf{k}_2^2 \mathbf{k}_3^2 \mathbf{k}_{12}^2 \mathbf{k}_{13}^2 \mathbf{k}_{23}^2 \mathbf{p}_2^2 \mathbf{p}_3^2}, \quad (3.4)$$

where $\int_{\mathbf{k}_i} \equiv \int \frac{d^{d-1} \mathbf{k}_i}{(2\pi)^{d-1}}$, $\mathbf{p}_i \equiv \mathbf{p} - \mathbf{k}_i$ and $\mathbf{k}_{ij} \equiv \mathbf{k}_i - \mathbf{k}_j$. The numerator, $\text{Num}(\mathbf{k}_i, \mathbf{p})$, depends on the loop momenta, \mathbf{k}_i , as well as the Fourier variable, \mathbf{p} , related to the relative distance, $\mathbf{r} \equiv \mathbf{x}_1 - \mathbf{x}_2$. In principle this topology contributes at $\mathcal{O}(G_N^4)$, and naively it would have entered also at 3PN order. However, the ϕ^3 vertex introduces an extra factor of v^2 [41]. Therefore, the diagram in fig. 1 enters at order $G_N^4 v^2$, namely 4PN. The calculation can be reduced to a series of master integrals, in particular one such integral will be of the form

$$\mathcal{M}(\mathbf{p}) = \int_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3} \frac{1}{\mathbf{k}_1^2 \mathbf{k}_2^2 \mathbf{k}_3^2 \mathbf{d}_2^2 \mathbf{d}_3^2}, \quad (3.5)$$

with $\mathbf{d}_2 = \mathbf{p} + \mathbf{k}_1 - \mathbf{k}_2$, $\mathbf{d}_3 = \mathbf{p} + \mathbf{k}_1 - \mathbf{k}_2 - \mathbf{k}_3$, so that the contribution to the gravitational potential is given by,

$$V_{\text{fig.1}} = c_{\mathcal{M}} m_1^2 m_2^3 \int_{\mathbf{p}} \mathbf{p}^2 \mathcal{M}(\mathbf{p}) e^{i\mathbf{p} \cdot \mathbf{r}} + \dots, \quad (3.6)$$

with some numerical pre-factor, $c_{\mathcal{M}} \propto G_N^4$.

The master integral may be computed in d -dimensions and is IR divergent for $d \rightarrow 4$. The relevant IR divergent part is given by⁷

$$\mathcal{M} \propto |\mathbf{p}|^{-1 + \frac{3}{2}\epsilon_{\text{IR}}} \Gamma[\epsilon_{\text{IR}}] \dots. \quad (3.7)$$

⁵As emphasized in [35], this must be the case since the factors for the pole and logarithm are related in dim. reg., and the coefficient for the latter is fixed by the long-distance contribution to the binding energy, first obtained in [37] and re-derived in [35] (see also [38]).

⁶The dim. reg. prescription must be universally applied in both regions (regardless of whether we have IR or UV divergent integrals) with the d -dimensional G_N determined from the Einstein-Hilbert action.

⁷See [42] for the complete expression which includes various Γ -functions.




Figure 1: A contribution in NRGR to the gravitational potential from the near zone at 4PN order. We use the decomposition introduced in [41] in terms of $(\phi, \mathbf{A}_i, \gamma_{ij})$ fields. As described in [40], the integral may be cast as a three-loop self-energy diagram in gauge theory.

The origin of this divergence can be traced down to the standard formula,

$$I[n_1, n_2] \equiv \int_{\mathbf{k}} \frac{1}{[[\mathbf{k}^2]^{n_1} (\mathbf{k} + \mathbf{p})^2]^{n_2}} \quad (3.8)$$

$$= \frac{\Gamma[n_1 + n_2 - (d-1)/2] \Gamma[(d-1)/2 - n_2] \Gamma[(d-1)/2 - n_1]}{\Gamma[n_1] \Gamma[n_2] \Gamma[(d-1) - n_1 - n_2]} (\mathbf{p}^2)^{(d-1)/2 - n_1 - n_2},$$

which can be used repeatedly (three times) to factorize the integral. The IR divergence is manifest by the fact that at some point a term may become singular for $d < 4$.

3.3 Removing the double-counting

The implementation of the zero-bin subtraction for the master integral in (3.5) is relatively straightforward. Because it can be factorized, the IR pole can be simply isolated from (see (3.8))

$$I \left[\frac{3-\epsilon}{2}, \frac{1-\epsilon}{2} \right] \propto |\mathbf{p}|^{-1+\frac{3}{2}\epsilon} \Gamma[\epsilon], \quad (3.9)$$

which contains the expected $\propto 1/\epsilon_{\text{IR}}$ coming from the region $\mathbf{k} \ll \mathbf{p}$, which must be subtracted away. Hence, expanding the factor of $(\mathbf{k} + \mathbf{p})$, we find

$$I_{\text{zero-bin}}[n_1, n_2] = \int_{\mathbf{k}} \frac{1}{[\mathbf{k}^2]^{n_1} [\mathbf{p}^2]^{n_2}} \xrightarrow{(n_1 \rightarrow 3/2, n_2 \rightarrow 1/2)} |\mathbf{p}|^{-1} \int_{\mathbf{k}} \frac{1}{\mathbf{k}^3} \propto |\mathbf{p}|^{-1} \left(\frac{1}{\epsilon_{\text{UV}}} - \frac{1}{\epsilon_{\text{IR}}} \right). \quad (3.10)$$

Notice the zero-bin subtraction transformed the IR pole into a UV singularity. This example illustrates how the procedure works at the level of the Feynman diagrams.

Plugging the expression for the master integral in (3.6), including the zero-bin subtraction, we get a term,

$$V_{\text{fig. 1}} \rightarrow \frac{G_N^4 m_1^2 m_2^3 v^2}{r^4} \left(-\frac{1}{\epsilon_{\text{UV}}} + 2 \log \mu r \right) + \dots, \quad (3.11)$$

where the IR pole turned into a UV divergence. Using the equations of motion (EOM), we have

$$G_N^2 (m_1 + m_2) (I^{ij(3)})^2 \xrightarrow{\text{EOM}} \frac{G_N^4 (m_1^2 m_2^3 + m_2^2 m_1^3) v^2}{r^3} (1 + \mathcal{O}(v^2)), \quad (3.12)$$

such that the contribution from fig. 1 to the gravitational binding potential has the correct structure to cancel the UV divergence from the conservative part of the tail effect (and $\log \mu$ factor), as advertised [35]. The procedure continues by implementing the zero-bin subtraction in all the IR-sensitive master integrals that enter at $\mathcal{O}(G_N^4 v^2)$.

4 Concluding remarks

The gravitational binding energy contains a logarithmic contribution, $E_{\log} \propto v^8 \log v$, in GR. When performing PN calculations, using all the present methodologies, one splits the problem up into near and far zone, which leads to both IR and UV logarithmically divergent integrals. The sum of these results reproduces the aforementioned logarithm, but at the cost of introducing a set of spurious new poles. While the UV divergences are well understood, the presence of IR divergences is a new development in the PN framework, which first entered scene at 4PN order for non-rotating bodies.⁸ The IR sensitivity in the near region has led to the introduction of ambiguity parameters in the methodology implemented in [27–33]. This occurs in any framework in which the IR/UV divergences, from the near/far zone contributions to the gravitational potential, are treated independently. That is what happens in the computations in [27–32], which removed the IR singularity in the near zone prior to appending the contribution from the far region, obtained independently in [43, 44]. However, by implementing the zero-bin subtraction there are no ambiguity parameters at any stage of the calculation. The procedure transforms the near zone IR divergence into a UV pole which readily cancels the one from the far region. We illustrated the zero-bin subtraction in the calculation of the conservative dynamics at 4PN order using dim. reg., in which case it simply replaces $\epsilon_{\text{IR}} \rightarrow \epsilon_{\text{UV}}$, by subtracting from the near region calculation a series of scaleless integrals. This justifies the claims in [35] that the sum of the near/far region IR/UV poles in dim. reg. must cancel each other out. However, the formalism is more general, and may be also implemented for other regularization schemes, where a cancellation of IR and UV divergences would be far less transparent to implement, including the ones in [27–33].⁹ The virtue of this procedure is precisely the independence from the regulator and subsequent lack of ambiguities to all orders in the PN expansion. The zero-bin subtraction not only provides finite results, it also uniquely fixes the analytic (local) contribution to the potential without the need of extra matching conditions beyond the PN framework. An explicit example is described in [34].

Acknowledgments

We thank the participants of the workshop ‘Analytic Methods in General Relativity’ held at ICTP-SAIFR,¹⁰ (supported by the São Paulo Research Foundation (FAPESP) grant 2016/01343-7) for fruitful interactions, in particular to Luc Blanchet, Guillaume Faye and Gerhard Schäfer. We thank also Aneesh Manohar for discussions. R.A.P. thanks the theory group at DESY (Hamburg) for hospitality while this paper was being completed. The work of R.A.P. is supported by the Simons Foundation and FAPESP Young Investigator Awards, grants 2014/25212-3 and 2014/10748-5. I.Z.R. is supported by the National Science Foundation under grant NSF-1407744.

⁸Up until now, the computations within the EFT approach had not encountered IR divergences in the potential region, and the only IR poles in the radiation theory arose from the tail contribution to the one-point function (radiative multipole moments), previously mentioned. There are other UV divergences in the radiation sector, which are responsible for the renormalization group evolution for the multipole moments (or can be consistently set to zero in dim. reg.), e.g. [15].

⁹See [26] for a discussion of the zero-bin subtraction also in the confines of a cut-off regulator.

¹⁰<http://www.ictcp-saifr.org/gr2016>

References

- [1] L. Blanchet, “Gravitational Radiation from Post-Newtonian Sources and Inspiralling Compact Binaries,” *Living Rev.Rel.* **17** (2014) 2, [arXiv:1310.1528 \[gr-qc\]](#).
- [2] A. Buonanno and B. S. Sathyaprakash, “Sources of Gravitational Waves: Theory and Observations,” [arXiv:1410.7832 \[gr-qc\]](#).
- [3] A. Le Tiec, “The Overlap of Numerical Relativity, Perturbation Theory and Post-Newtonian Theory in the Binary Black Hole Problem,” *Int. J. Mod. Phys.* **D23** no. 10, (2014) 1430022, [arXiv:1408.5505 \[gr-qc\]](#).
- [4] **Virgo, LIGO Scientific** Collaboration, J. Abadie *et al.*, “Predictions for the Rates of Compact Binary Coalescences Observable by Ground-based Gravitational-wave Detectors,” *Class. Quant. Grav.* **27** (2010) 173001, [arXiv:1003.2480 \[astro-ph.HE\]](#).
- [5] **Virgo, LIGO Scientific** Collaboration, B. P. Abbott *et al.*, “Observation of Gravitational Waves from a Binary Black Hole Merger,” *Phys. Rev. Lett.* **116** no. 6, (2016) 061102, [arXiv:1602.03837 \[gr-qc\]](#).
- [6] **Virgo, LIGO Scientific** Collaboration, B. P. Abbott *et al.*, “Astrophysical Implications of the Binary Black-Hole Merger GW150914,” *Astrophys. J.* **818** no. 2, (2016) L22, [arXiv:1602.03846 \[astro-ph.HE\]](#).
- [7] **Virgo, LIGO Scientific** Collaboration, B. P. Abbott *et al.*, “Binary Black Hole Mergers in the first Advanced LIGO Observing Run,” *Phys. Rev.* **X6** no. 4, (2016) 041015, [arXiv:1606.04856 \[gr-qc\]](#).
- [8] A. Sesana, “Prospects for Multiband Gravitational-Wave Astronomy after GW150914,” *Phys. Rev. Lett.* **116** no. 23, (2016) 231102, [arXiv:1602.06951 \[gr-qc\]](#).
- [9] W. D. Goldberger and I. Z. Rothstein, “An Effective field theory of gravity for extended objects,” *Phys.Rev.* **D73** (2006) 104029, [arXiv:hep-th/0409156 \[hep-th\]](#).
- [10] W. D. Goldberger, “Les Houches lectures on effective field theories and gravitational radiation,” 2007. [arXiv:hep-ph/0701129 \[hep-ph\]](#).
- [11] R. A. Porto and R. Sturani, “Scalar gravity: Post-Newtonian corrections via an effective field theory approach,” in *Les Houches Summer School - Session 86: Particle Physics and Cosmology: The Fabric of Spacetime Les Houches, France, July 31-August 25, 2006*. 2007. [arXiv:gr-qc/0701105 \[gr-qc\]](#).
- [12] I. Z. Rothstein, “Progress in effective field theory approach to the binary inspiral problem,” *Gen. Rel. Grav.* **46** (2014) 1726.
- [13] V. Cardoso and R. A. Porto, “Analytic approximations, perturbation theory, effective field theory methods and their applications,” *Gen. Rel. Grav.* **46** (2014) 1682, [arXiv:1401.2193 \[gr-qc\]](#).
- [14] S. Foffa and R. Sturani, “Effective field theory methods to model compact binaries,” *Class. Quant. Grav.* **31** no. 4, (2014) 043001, [arXiv:1309.3474 \[gr-qc\]](#).
- [15] R. A. Porto, “The effective field theorist’s approach to gravitational dynamics,” *Phys. Rept.* **633** (2016) 1–104, [arXiv:1601.04914 \[hep-th\]](#).

- [16] W. D. Goldberger and M. B. Wise, “Renormalization group flows for brane couplings,” *Phys. Rev.* **D65** (2002) 025011, [arXiv:hep-th/0104170](#) [hep-th].
- [17] I. Z. Rothstein, “TASI lectures on effective field theories,” [arXiv:hep-ph/0308266](#) [hep-ph].
- [18] T. Damour, “THE PROBLEM OF MOTION IN NEWTONIAN AND EINSTEINIAN GRAVITY,” in *300 Years of Gravity: A Conference to Mark the 300th Anniversary of the Publication of Newton’s Principia Cambridge, England, June 29-July 4, 1987*. 1986.
- [19] R. A. Porto, “The Tune of Love and the Nature(ness) of Spacetime,” [arXiv:1606.08895](#) [gr-qc].
- [20] R. A. Porto, “Post-Newtonian corrections to the motion of spinning bodies in NRGR,” *Phys.Rev.* **D73** (2006) 104031, [arXiv:gr-qc/0511061](#) [gr-qc].
- [21] R. A. Porto and I. Z. Rothstein, “The Hyperfine Einstein-Infeld-Hoffmann potential,” *Phys.Rev.Lett.* **97** (2006) 021101, [arXiv:gr-qc/0604099](#) [gr-qc].
- [22] R. A. Porto and I. Z. Rothstein, “Spin(1)Spin(2) Effects in the Motion of Inspiralling Compact Binaries at Third Order in the Post-Newtonian Expansion,” *Phys.Rev.* **D78** (2008) 044012, [arXiv:0802.0720](#) [gr-qc].
- [23] R. A. Porto and I. Z. Rothstein, “Next to Leading Order Spin(1)Spin(1) Effects in the Motion of Inspiralling Compact Binaries,” *Phys.Rev.* **D78** (2008) 044013, [arXiv:0804.0260](#) [gr-qc].
- [24] W. D. Goldberger and A. Ross, “Gravitational radiative corrections from effective field theory,” *Phys. Rev.* **D81** (2010) 124015, [arXiv:0912.4254](#) [gr-qc].
- [25] R. A. Porto, A. Ross, and I. Z. Rothstein, “Spin induced multipole moments for the gravitational wave amplitude from binary inspirals to 2.5 Post-Newtonian order,” *JCAP* **1209** (2012) 028, [arXiv:1203.2962](#) [gr-qc].
- [26] A. V. Manohar and I. W. Stewart, “The Zero-Bin and Mode Factorization in Quantum Field Theory,” *Phys. Rev.* **D76** (2007) 074002, [arXiv:hep-ph/0605001](#) [hep-ph].
- [27] P. Jaranowski and G. Schafer, “Dimensional regularization of local singularities in the 4th post-Newtonian two-point-mass Hamiltonian,” *Phys. Rev.* **D87** (2013) 081503, [arXiv:1303.3225](#) [gr-qc].
- [28] P. Jaranowski and G. Schafer, “Derivation of the local-in-time fourth post-Newtonian ADM Hamiltonian for spinless compact binaries,” [arXiv:1508.01016](#) [gr-qc].
- [29] T. Damour, P. Jaranowski, and G. Schafer, “Nonlocal-in-time action for the fourth post-Newtonian conservative dynamics of two-body systems,” *Phys. Rev.* **D89** no. 6, (2014) 064058, [arXiv:1401.4548](#) [gr-qc].
- [30] T. Damour, P. Jaranowski, and G. Schafer, “On the conservative dynamics of two-body systems at the fourth post-Newtonian approximation of general relativity,” [arXiv:1601.01283](#) [gr-qc].
- [31] L. Bernard, L. Blanchet, A. Bohe, G. Faye, and S. Marsat, “Fokker action of non-spinning compact binaries at the fourth post-Newtonian approximation,” [arXiv:1512.02876](#) [gr-qc].
- [32] L. Bernard, L. Blanchet, A. Bohe, G. Faye, and S. Marsat, “Energy and periastron advance of compact binaries on circular orbits at the fourth post-Newtonian order,” *Phys. Rev.* **D95** (2017) 044026, [arXiv:1610.07934](#) [gr-qc].

- [33] T. Damour and P. Jaranowski, “On the four-loop static contribution to the gravitational interaction potential of two point masses,” [arXiv:1701.02645](#) [[gr-qc](#)].
- [34] R. A. Porto, “The Lamb shift and the gravitational binding energy for binary black holes,” *Companion paper* .
- [35] C. R. Galley, A. K. Leibovich, R. A. Porto, and A. Ross, “Tail effect in gravitational radiation reaction: Time nonlocality and renormalization group evolution,” *Phys. Rev.* **D93** (2016) 124010, [arXiv:1511.07379](#) [[gr-qc](#)].
- [36] L. Blanchet and A. Le Tiec, “First Law of Compact Binary Mechanics with Gravitational-Wave Tails,” [arXiv:1702.06839](#) [[gr-qc](#)].
- [37] L. Blanchet, S. L. Detweiler, A. Le Tiec, and B. F. Whiting, “High-Order Post-Newtonian Fit of the Gravitational Self-Force for Circular Orbits in the Schwarzschild Geometry,” *Phys. Rev.* **D81** (2010) 084033, [arXiv:1002.0726](#) [[gr-qc](#)].
- [38] W. D. Goldberger, A. Ross, and I. Z. Rothstein, “Black hole mass dynamics and renormalization group evolution,” *Phys. Rev.* **D89** no. 12, (2014) 124033, [arXiv:1211.6095](#) [[hep-th](#)].
- [39] S. Foffa and R. Sturani, “Dynamics of the gravitational two-body problem at fourth post-Newtonian order and at quadratic order in the Newton constant,” *Phys. Rev.* **D87** no. 6, (2013) 064011, [arXiv:1206.7087](#) [[gr-qc](#)].
- [40] S. Foffa, P. Mastrolia, R. Sturani, and C. Sturm, “Effective field theory approach to the gravitational two-body dynamics, at fourth post-Newtonian order and quintic in the Newton constant,” [arXiv:1612.00482](#) [[gr-qc](#)].
- [41] B. Kol and M. Smolkin, “Non-Relativistic Gravitation: From Newton to Einstein and Back,” *Class. Quant. Grav.* **25** (2008) 145011, [arXiv:0712.4116](#) [[hep-th](#)].
- [42] T. Gehrmann, G. Heinrich, T. Huber, and C. Studerus, “Master integrals for massless three-loop form-factors: One-loop and two-loop insertions,” *Phys. Lett.* **B640** (2006) 252–259, [arXiv:hep-ph/0607185](#) [[hep-ph](#)].
- [43] L. Blanchet and T. Damour, “Hereditary effects in gravitational radiation,” *Phys. Rev.* **D46** (1992) 4304–4319.
- [44] L. Blanchet, “Time asymmetric structure of gravitational radiation,” *Phys. Rev.* **D47** (1993) 4392–4420.