

Pentaquark photoproduction

César Fernández-Ramírez

Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Ciudad de México 04510, Mexico

E-mail: `cesar.fernandez@nucleares.unam.mx`

Astrid N. Hiller Bin

Departamento de Física Teórica and IFIC, Centro Mixto Universidad de Valencia-CSIC, Institutos de Investigación de Paterna, E-46071 Valencia, Spain

Alessandro Pilloni

Theory Center, Thomas Jefferson National Accelerator Facility, Newport News, VA 23606, USA

Abstract. We present results and suggestions on how to confirm the existence and resonant nature of the $P_c(4450)$ detected at LHCb through photoproduction experiments. We find that this narrow structure might have escaped detection in past experiments and use those to give a constraint for the upper limit of the branching ratio/coupling to the $J/\psi p$ channel.

1. Introduction

One of the most remarkable properties of Quantum Chromodynamics (QCD) is the color confinement phenomenon that makes hadrons to be built out of color singlets of quarks. The simplest way to build a color neutral baryon (half-integer spin) state is out of three quarks (qqq). However, QCD tells us that there are other ways to build color neutral baryons, for example: hybrid baryons with gluonic components ($qqqg$, $qqqqg$, $\dots$), pentaquarks ($qqqq\bar{q}$), heptaquarks ($qqqqq\bar{q}\bar{q}$) and so on. Finding complete multiplets of baryons beyond the qqq picture is one of the main goals of hadron spectroscopy and will provide insight on the dynamics of QCD in the hadronic energy regime. The recent discovery of two structures compatible with pentaquark states in the $\Lambda_b^0 \rightarrow J/\psi K^- p$ decay by LHCb [1] together with the observation of a compatible structure in $\Lambda_b^0 \rightarrow J/\psi \pi^- p$ [2], has triggered a lot of research activities aimed to explain them. Each signal has been associated to a pentaquark candidate, $P_c(4380)$ and $P_c(4450)$, with opposite parity assignments, and one state having spin $3/2$ and the other $5/2$ [1, 3]. The $P_c(4380)$ signal is less apparent and the properties of this state (mass and width) could depend on the amplitude analysis performed, especially for the $K^- p$ background [3, 4]. However, the $P_c(4450)$ signal is very narrow and its mass and width are not expected to change by refinements in the amplitude analysis. Hence, its study has been the focus of many research groups. Several explanations for this signal have been proposed: a kinematical effect [5], a molecular state [6] and a compact QCD (pentaquark) state [7].

One of the best approaches to gain insight on the nature of $P_c(4450)$ consists on detecting the state in a different reaction. In [8] it was proposed to use $J/\psi p$ photoproduction. From the LHCb Λ_b^0 decay data we know that the $P_c(4450)$ state couples strongly to the $J/\psi p$, and the photon energy can be tuned to excite this particular state. This experiment has just been approved to run at Hall C in Jefferson Lab [9] and results are expected in the next years. Hence, as a first step it is necessary to develop the amplitudes that will be needed to provide realistic predictions for the planning of the experiment and later to analyze the experimental data. In the next section we sketch the main features of the J/ψ photoproduction model that we developed with these purposes in mind. The detailed description of the model and the corresponding results can be found in [10].

2. Pentaquark photoproduction model

The differential cross section for the $\gamma p \rightarrow J/\psi p$ reaction can be written

$$\frac{d\sigma}{d\cos\theta} = \frac{4\pi\alpha}{32\pi s} \frac{p_f}{p_i} \frac{1}{4} \sum_{\lambda_\gamma, \lambda_p, \lambda_\psi, \lambda_{p'}} |\langle \lambda_\psi \lambda_{p'} | T | \lambda_\gamma \lambda_p \rangle|^2, \quad (1)$$

where p_i is the incoming momentum, p_f is the outgoing momentum, θ is the scattering angle, s is the standard Mandelstam variable, α is the electromagnetic fine structure constant, and λ_ψ , $\lambda_{p'}$, λ_γ and λ_p are the J/ψ , outgoing proton, photon and incoming proton helicities respectively. All the kinematical variables are defined in the center of mass reference system.

The amplitude $\langle \lambda_\psi \lambda_{p'} | T | \lambda_\gamma \lambda_p \rangle$ is constructed as the addition of two pieces: (i) the pentaquark resonance as a Breit–Wigner amplitude

$$\langle \lambda_\psi \lambda_{p'} | T_r | \lambda_\gamma \lambda_p \rangle = \frac{\langle \lambda_\psi \lambda_{p'} | T_{\text{dec}} | \lambda_r \rangle \langle \lambda_r | T_{\text{em}}^\dagger | \lambda_\gamma \lambda_p \rangle}{M_r^2 - W^2 - i\Gamma_r M_r}, \quad (2)$$

where M_r , Γ_r and λ_r are the $P_c(4450)$ mass, width and helicity respectively, $W = \sqrt{s}$ is the invariant mass, T_{dec} stands for the strong vertex $pJ/\psi P_c$, and T_{em} stands for the electromagnetic vertex $\gamma p P_c$; and (ii) a nonresonant background described through an affective Pomeron exchange that is constrained by existing high-energy data [11]

$$\langle \lambda_\psi \lambda_{p'} | T_P | \lambda_\gamma \lambda_p \rangle = iA \left(\frac{s - s_t}{s_0} \right)^{\alpha(t)} e^{b_0(t - t_{\min})} \delta_{\lambda_p \lambda_{p'}} \delta_{\lambda_\psi \lambda_\gamma}, \quad (3)$$

where t is the standard Mandelstam variable, $t_{\min}$ is the minimum t for a given s , $s_0 = 1 \text{ GeV}^2$ is the hadron physics scale, $\alpha(t) = \alpha_0 + \alpha' t$ is the Pomeron Regge trajectory whose parameters α_0 and α' are fitted to the data, and A , b_0 and t_0 are also fitted to the data. The electromagnetic vertex can be related to the helicity amplitudes $A_{1/2}$ and $A_{3/2}$ through

$$\langle \lambda_\gamma \lambda_p | T_{\text{em}} | \lambda_r \rangle = \frac{W}{M_r} \sqrt{\frac{8M_N M_r \bar{p}_i}{4\pi\alpha}} \sqrt{\frac{\bar{p}_i}{p_i}} A_{1/2, 3/2}, \quad (4)$$

where M_N is the nucleon mass and $\bar{p}_i$ is the momentum p_i evaluated at the resonance mass M_r . We do not know the helicity amplitudes of the pentaquark, hence, we need to make assumptions to describe this piece of the model. We assume the electromagnetic and strong vertices to be related by vector meson dominance

$$\langle \lambda_\gamma \lambda_p | T_{\text{em}} | \lambda_r \rangle = \frac{\sqrt{4\pi\alpha} f_\psi}{M_\psi} \langle \lambda_\psi = \lambda_\gamma, \lambda_p | T_{\text{dec}} | \lambda_r \rangle, \quad (5)$$

where M_ψ and $f_\psi = 280$ MeV are the J/ψ mass and decay constant respectively, and

$$\langle \lambda_\psi \lambda_{p'} | T_{\text{dec}} | \lambda_r \rangle = g_{\lambda_\psi \lambda_{p'}} d_{\lambda_r, \lambda_\psi - \lambda_{p'}}^{J_r}(\cos \theta), \quad (6)$$

where $d_{\lambda_r, \lambda_\psi - \lambda_{p'}}^{J_r}(\cos \theta)$ are the Wigner d -functions with J_r the resonance spin, and $g_{\lambda_\psi, \lambda_{p'}}$ are the helicity couplings. We assume the three helicity couplings to be equal, $g = g_{\lambda_\psi, \lambda_{p'}}$. Hence the partial decay width $\Gamma_{\psi p}$ reads

$$\Gamma_{\psi p} = \mathcal{B}_{\psi p} \Gamma_r = \frac{\bar{p}_f}{32\pi^2 M_r^2} \frac{1}{2J_r + 1} \sum_{\lambda_R} \int d\Omega |\langle \lambda_\psi \lambda_{p'} | T_{\text{dec}} | \lambda_R \rangle|^2 = \frac{\bar{p}_f}{8\pi M_r^2} \frac{6g^2}{2J_r + 1}, \quad (7)$$

where $\mathcal{B}_{\psi p}$ is the $P_c(4450) \rightarrow J/\psi p$ branching ratio and $\bar{p}_f$ the p_f momentum at the resonance peak. Since there are no data in the resonance peak region we can only determine an upper limit for $\mathcal{B}_{\psi p}$ within a certain confidence level (CL) to ensure compatibility with the available data.

3. Results

We fit the available data on the $\gamma p \rightarrow J/\psi p$ reaction for $|t| \leq 1.5$ GeV² [11] and we compute the uncertainties in the parameters up to 68% and 95% CL using the bootstrap technique [12]. We also incorporate the uncertainty of the pentaquark mass and width through a Monte Carlo. In this way we can propagate exactly the experimental uncertainties to our predictions. Figure 1 shows the differential cross section $d\sigma/dt$ at $t = t_{\text{min}}$ in the pentaquark energy region for the $J_r^P = 3/2^-$ and $5/2^+$ spin-parity assignments. The pentaquark peak is very pronounced, which implies that, if the LHCb signal is a true resonance and does not couple too weakly to the photon (as expected from vector meson dominance), it will be visible in the proposed photoproduction experiment at Jefferson Lab [9]. We found that data allow for the existence of a pentaquark whose $J/\psi p$ branching ratio has to be below 30% for $J^P = 3/2^-$ and below 17% for $J^P = 5/2^+$ at a 95% CL. The values of the parameters, their uncertainties, and the full extent of the analysis can be found in [10].




Figure 1. Differential cross section in the $P_c(4450)$ peak region at $t = t_{\text{min}}$ for the $J_r = 3/2^-$ (left plot) and the $J_r = 5/2^+$ (right plot) spin-parity assignments.

4. Final remarks

The signals found in the $\Lambda_b^0 \rightarrow J/\psi K^- p$ decay by the LHCb collaboration [1] indicate the possible existence of a hidden charm pentaquark with a mass of 4450 MeV. One of the most

promising ways to independently confirm the pentaquark nature of the signal consists on trying to photoproduce the state using the $\gamma p \rightarrow J/\psi p$ reaction. The recent approval by Jefferson Lab's PAC44 of this very experiment at Hall C [9] and the development of photoproduction models to analyze the data [8, 10] make us believe that in the near future we will get a clear answer. The codes for the model in [10] can be downloaded and run online from the Joint Physics Analysis Center (JPAC) web page [13]. If the $P_c(4450)$ state is also detected in J/ψ photoproduction, it would likely be a compact state. Then, new experiments aimed to hunt the other members of this hidden-charm pentaquark multiplet can begin [14] as well as experiments to map the Q^2 dependence of the pentaquark electromagnetic form factor and provide insight into its internal structure.

Acknowledgments

This work is part of the efforts of the Joint Physics Analysis Center (JPAC) collaboration. CF-R work is supported in part by research grant IA101717 from PAPIIT-DGAPA (UNAM), by CONACYT (Mexico) grant No. 251817, and by Red Temática CONACYT de Física en Altas Energías (Red FAE, Mexico). ANHB research is supported in part by the Spanish Ministerio de Economía y Competitividad (MINECO), European FEDER funds under contract No. FIS2014-51948-C2-2-P, SEV-2014-0398, and Generalitat Valenciana under Contract PROMETEOII/2014/0068. AP research is supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under contract DE-AC05-06OR23177. CF-R thanks the organizers for their invitation to the conference and their warm hospitality at Cocoyoc.

References

- [1] Aaij R *et al* (LHCb collaboration) 2015 *Phys. Rev. Lett.* **115** 072001
Aaij R *et al* (LHCb collaboration) 2016 *Phys. Rev. Lett.* **117** 082002
- [2] Aaij R *et al* (LHCb collaboration) 2016 *Phys. Rev. Lett.* **117** 082003
- [3] Patrignani C *et al* (Particle Data Group) 2016 *Chin. Phys. C* **40** 100001
- [4] Fernández-Ramírez C *et al* 2016 *Phys. Rev. D* **93** 034029
- [5] Meißner UG and Oller JA 2015 *Phys. Lett. B* **751** 59
Guo FK, Meißner UG, Wang W and Yang Z 2015 *Phys. Rev. D* **92** 071502
Guo FK, Meißner UG, Nieves J and Yang Z, 2016 *Eur. Phys. J. A* **52** 318
- [6] Chen R, Liu X, Li Q and Zhu SL *Phys. Rev. Lett.* **115**, 132002 (2015)
Chen HX *et al* *Phys. Rev. Lett.* **115**, 172001 (2015)
Roca L, Nieves J and Oset E 2015 *Phys. Rev. D* **92** 094003
He J 2016 *Phys. Lett. B* **753** 547
Lü QF and Dong YB 2016 *Phys. Rev. D* **93** 074020
- [7] Maiani L, Polosa AD and Riquer V 2015 *Phys. Lett. B* **749** 289
Lebed RF 2015 *Phys. Lett. B* **749** 454
Mironov A and Morozov A 2015 *JETP Lett.* **102** 271
Esposito A, Pilloni A and Polosa AD 2017 *Phys. Rep.* **668** 1
- [8] Wang Q, Liu XH and Zhao Q 2015 *Phys. Rev. D* **92** 034022
Kubarovsky V and Voloshin MB 2015 *Phys. Rev. D* **92** 031502
Karliner M and Rosner JL 2016 *Phys. Lett. B* **752** 329
- [9] Mezziani ZE *et al* (Hall C collaboration) 2016 *A Search for the LHCb Charmed "Pentaquark" using Photoproduction of J/ψ at Threshold in Hall C at Jefferson Lab*, Preprint 1609.00676 [hep-ex]
- [10] Hiller Blin AN *et al* 2016 *Phys. Rev. D* **94** 034002
- [11] Camerini U *et al* 1975 *Phys. Rev. Lett.* **35** 483
Ritson DM 1976 *AIP Conf. Proc.* **30** 75
Anderson RL 1976 *Madison Conf.* 102 Preprint SLAC-PUB-1741
Chekanov S *et al* (ZEUS collaboration) 2002 *Eur. Phys. J. C* **24** 345
Aktas A *et al* (H1 collaboration) 2006 *Eur. Phys. J. C* **46** 585
- [12] Landay J *et al* 2017 *Phys. Rev. C* **95** 015203
- [13] Mathieu V 2016 *AIP Conf. Proc.* **1735** 070004; <http://www.indiana.edu/~jpac/>
- [14] Bijker R 2017 *J. Phys.: Conf. Ser.* submitted
Bijker R and Fernández-Ramírez C 2017 *Phys. Rev. C* submitted