

The structure of pentaquarks Ω_c^0 in the chiral quark model

Gang Yang, Jialun Ping*

¹*Department of Physics and Jiangsu Key Laboratory for Numerical Simulation of Large Scale Complex Systems, Nanjing Normal University, Nanjing 210023, P. R. China*

Recently, the experimental results of LHCb Collaboration suggested the existence of five new excited states of Ω_c^0 , $\Omega_c(3000)^0$, $\Omega_c(3050)^0$, $\Omega_c(3066)^0$, $\Omega_c(3090)^0$ and $\Omega_c(3119)^0$, the quantum numbers of these new particles are not determined now. To understand the nature of the states, a dynamical calculation of 5-quark systems with quantum numbers $IJ^P = 0(\frac{1}{2})^-$, $0(\frac{3}{2})^-$ and $0(\frac{5}{2})^-$ is performed in the framework of chiral quark model with the help of gaussian expansion method. The results show the $\Xi\bar{D}$, $\Xi_c\bar{K}$ and $\Xi_c^*\bar{K}$ are possible the candidates of these new particles. The distances between quark pairs suggest that the nature of pentaquark states.

PACS numbers: 14.20.Lq, 14.40.Lb, 12.39.Jh

I. INTRODUCTION

Recently, CERN announced an exceptional new discovery that was made by the LHCb, which unveiled five new states all at once [1]. Each of the five particles were found to be the excited states of Ω_c^0 , a particle with three quarks, *css*. These particle states are named, according to the standard convention, $\Omega_c(3000)^0$, $\Omega_c(3050)^0$, $\Omega_c(3066)^0$, $\Omega_c(3090)^0$ and $\Omega_c(3119)^0$. Just after the announcement, the theoretical interpretations were proposed. S. S. Agaev *et al.* interpreted two of these excited charmed baryons ($\Omega_c(3066)^0$ and $\Omega_c(3119)^0$) as the first radial excitation with $(2S, 1/2^+)$ and $(2S, 3/2^+)$, respectively in QCD sum rules [2]. The same conclusion is proposed by H. X. Chen *et al.* [3] in studying the decay properties of P -wave charmed baryons from light-cone QCD sum rules, besides they also suggest that one of these Ω_c^0 states ($\Omega_c(3000)^0$, $\Omega_c(3050)^0$ or $\Omega_c(3066)^0$) as a $J^P = 1/2^-$ state, the rest two states is with $J^P = 3/2^-$ and $J^P = 5/2^-$. In Ref. [4], Karliner and Rosner suggested that the parity was negative for all of the five states, two $J^P = 1/2^-$ states ($\Omega_c(3000)^0$ and $\Omega_c(3050)^0$), two $J^P = 3/2^-$ states ($\Omega_c(3066)^0$ and $\Omega_c(3090)^0$), and the last one is $\Omega_c(3119)^0$ $J^P = 5/2^-$. These exciting announcements and the theoretical work along with the pentaquarks P_c^+ discovered also by the LHCb Collaboration in 2015 [5] have bring us lots of peculiar understanding to the world of microcosmic particles.

The quantum numbers of these new particles are not determined for the moment, and the explanation of them as the excited states of q^3 baryon is reasonable. However, the possibility of the multi-quark candidates of these excited states cannot be excluded. The ground states of Ω_c have been observed experimentally, $\Omega_c(2695)^0$ with $J^P = \frac{1}{2}^+$ and $\Omega_c(2770)^0$ with $J^P = \frac{3}{2}^+$. The excited energies of the newly reported states with respect to the ground states are 230-424 MeV, which are enough to excite light quark-antiquark pair from the vacuum. From

the masses of Ξ_c baryon and K meson, 2468 MeV and 495 MeV, we have the threshold for $\Xi_c - \bar{K}$ state around 2963 MeV. It is expected that the 5- q components will play a role in these Ω_c 's. In Ref. [27], spectrum of low-lying pentaquark states with strangeness $S = -3$ and negative parity is studied in three kinds of constituent quark models. The results indicate that the lowest energy state Ω^* is around 1.8 GeV, which is about 200 MeV lower than predictions of various quenched three quarks models, and the energy cost to excite ground state of Ω to a 5-quark state is less than that to an orbital excitation.

The interesting in pentaquark is revived after the observation of the exotic hadrons, $P_c^+(4380)$ and $P_c^+(4450)$ in the decay of Λ_b^0 , $\Lambda_b^0 \rightarrow J/\psi K^- p$ by the LHCb Collaboration lately [5], there are a lots of theoretical calculations have been performed to investigate these two exotic states [6–16], even though the $\Theta^+(1540)$ pentaquark was reported by several experimental groups [17–19] in 2003 and has been denied by JLab with more higher precision results [20] (LEPS Collaboration still insisted on the existence of pentaquark $\Theta^+(1540)$ [21]). Besides, it is shown that there should be notable five-quark components in the baryon resonances [22–24]. In addition, the valence-sea quark mixing (Fock space expansion) model ($q^3 + q^3 q\bar{q}$) of nucleon ground state had been used to explain the mysterious proton spin structure well [25]. Such a sea quark excitation model had also been used to show that the $q^3 q\bar{q}$ excitation is more favorable than the p -wave excitation in q^3 configuration for $1/2^-$ baryons [26].

Quark model is the most common approach to multi-quark system. With the recent experimental data on multi-quark states and the development of quark model, It is expected to perform a serious calculation of multi-quark state in the framework of quark model. In the present work, the chiral quark model (ChQM) is employed to study the pentaquark states Ω_c^0 . To find the structure of the pentaquark states, a general, powerful method of few-body system, gaussian expansion method (GEM) [28] is used to do the calculation. The GEM has been successfully applied to many few-body systems, light nuclei, hyper-nuclei, hadron physics and so on [28]. It suits for both of compact multi-quark systems and loosely bound molecular states. In this approach, the

*jlping@njnu.edu.cn, corresponding author

four relative orbital motions of the system are expanded by gaussians with various widths. By taking into account of all the possible couplings for color-flavor-spin degrees of freedom, the structure of the system determined by its dynamics can be found.

The structure of the paper is as follows. In section II the quark model, wave-functions and calculation method is presented. Section III is devoted to the calculated results and discussions. A brief summary is given in the last section.

II. MODEL AND WAVE FUNCTION

The chiral quark model has achieved a success both in describing the hadron spectra and hadron-hadron in-

teraction. In this model, the constituent quark and antiquark interact with each other through the Goldstone boson exchange and the effective one-gluon-exchange, in addition to the phenomenological color confinement. Besides, the scalar nonet (the extension of chiral partner σ -meson) exchange are also introduced. The details of the model can be found in Ref.[29]. So the Hamiltonian in the present calculation takes the form

$$H = \sum_{i=1}^n \left(m_i + \frac{p_i^2}{2m_i} \right) - T_{CM} + \sum_{j>i=1}^n [V_{CON}(\mathbf{r}_{ij}) + V_{OGE}(\mathbf{r}_{ij}) + V_\chi(\mathbf{r}_{ij}) + V_s(\mathbf{r}_{ij})], \quad (1)$$

$$V_{CON}(\mathbf{r}_{ij}) = \boldsymbol{\lambda}_i^c \cdot \boldsymbol{\lambda}_j^c [-a_c(1 - e^{-\mu_c r_{ij}}) + \Delta],$$

$$V_{OGE}(\mathbf{r}_{ij}) = \frac{1}{4} \alpha_s \boldsymbol{\lambda}_i^c \cdot \boldsymbol{\lambda}_j^c \left[\frac{1}{r_{ij}} - \frac{1}{6m_i m_j} \boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j \frac{e^{-r_{ij}/r_0(\mu)}}{r_{ij} r_0^2(\mu)} \right], \quad r_0(\mu) = \hat{r}_0/\mu,$$

$$V_\chi(\mathbf{r}_{ij}) = v_\pi(\mathbf{r}_{ij}) \sum_{a=1}^3 \boldsymbol{\lambda}_i^a \cdot \boldsymbol{\lambda}_j^a + v_K(\mathbf{r}_{ij}) \sum_{a=4}^7 \boldsymbol{\lambda}_i^a \cdot \boldsymbol{\lambda}_j^a + v_\eta(\mathbf{r}_{ij}) [\boldsymbol{\lambda}_i^8 \cdot \boldsymbol{\lambda}_j^8 \cos \theta_P - \boldsymbol{\lambda}_i^0 \cdot \boldsymbol{\lambda}_j^0 \sin \theta_P], \quad (2)$$

$$v_\chi(\mathbf{r}_{ij}) = \frac{g_{ch}^2}{4\pi} \frac{m_\chi^2}{12m_i m_j} \frac{\Lambda_\chi^2}{\Lambda_\chi^2 - m_\chi^2} m_\chi \left[Y(m_\chi r_{ij}) - \frac{\Lambda_\chi^3}{m_\chi^3} Y(\Lambda_\chi r_{ij}) \right] \boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j, \quad \chi = \pi, K, \eta,$$

$$V_s(\mathbf{r}_{ij}) = v_\sigma(\mathbf{r}_{ij})(\boldsymbol{\lambda}_i^0 \cdot \boldsymbol{\lambda}_j^0) + v_{a_0}(\mathbf{r}_{ij}) \sum_{a=1}^3 \boldsymbol{\lambda}_i^a \cdot \boldsymbol{\lambda}_j^a + v_\kappa(\mathbf{r}_{ij}) \sum_{a=4}^7 \boldsymbol{\lambda}_i^a \cdot \boldsymbol{\lambda}_j^a + v_{f_0}(\mathbf{r}_{ij})(\boldsymbol{\lambda}_i^8 \cdot \boldsymbol{\lambda}_j^8), \quad (3)$$

$$v_s(\mathbf{r}_{ij}) = -\frac{g_{ch}^2}{4\pi} \frac{\Lambda_s^2}{\Lambda_s^2 - m_s^2} m_s \left[Y(m_s r_{ij}) - \frac{\Lambda_s}{m_s} Y(\Lambda_s r_{ij}) \right], \quad s = \sigma, a_0, \kappa, f_0.$$

All the symbols take their usual meanings. μ is the reduced mass of two interacting quarks. To simplify the calculation, only the central parts of the interactions are employed in the present work to consider the ground state of multi-quark system. The model parameters are fixed by fitting the spectrum of baryons and mesons and their values are listed in Table I, the calculated masses of baryons and mesons are shown in Table II. There are two sets of parameters are given, the fixed quark-gluon coupling constant is used in the set I, the set II has the running coupling constants which are given as

$$\alpha_s = \frac{\alpha_0}{\ln((\mu^2 + \mu_0^2)/\Lambda_0^2)}.$$

It is worth to mention that the above quark-quark interaction is assumed to be universal according to the "Casimir scaling" [30], it can be applied to the multi-

quark system directly. The possible multi-body interaction in the multi-quark system is not considered, although it may give different spectra of multi-quark states [31].

From Table II, we can see that the masses of P -wave Ω_c 's are higher than 3200 MeV although the mass of P -wave nucleon is close the experimental value (for the set I). The parameters of set II is used to check the dependence of the results on the model parameters. The results show that the P -wave baryons have rather large masses, comparing with the experimental data. So it is still difficult to have a good description of the negative parity states of baryons in the quark model. In the following, we use set I parameters to study the 5-quark states.

The wavefunctions for the system are constructed just as the way in Ref. [6]. Here only the wavefunctions of each degree of freedom for five-quark system and parts of the sub-clusters of three-quark and quark-antiquark

TABLE I: Quark model parameters. The masses of mesons take their experimental values. $m_\pi = 0.7 \text{ fm}^{-1}$, $m_K = 2.51 \text{ fm}^{-1}$, $m_\pi = 2.77 \text{ fm}^{-1}$.

		set I	set II
Quark mass	$m_u=m_d \text{ (MeV)}$	313	313
	$m_s \text{ (MeV)}$	555	555
	$m_c \text{ (MeV)}$	1752	1752
Goldstone boson	$\Lambda_\pi = \Lambda_\sigma \text{ (fm}^{-1}\text{)}$	4.20	4.20
	$\Lambda_K = \Lambda_\eta \text{ (fm}^{-1}\text{)}$	5.20	5.20
	$\theta_P(^{\circ})$	-15	-15
	$g_{ch}^2/(4\pi)$	0.54	0.54
SU(3)	$m_s \text{ (fm}^{-1}\text{)}$	4.97	4.97
Scalar nonet $s = \sigma, a_0, \kappa, f_0$	$\Lambda_s \text{ (fm}^{-1}\text{)}$	5.20	5.20
	$m_\sigma \text{ (fm}^{-1}\text{)}$	3.42	3.42
Confinement	$a_c \text{ (MeV)}$	180	184.08
	$\mu_c \text{ (fm}^{-1}\text{)}$	0.645	0.634
	$\Delta \text{ (MeV)}$	55.5	40.249
OGE		$\alpha_0 = 1.293$	
	α_s	0.69	$\Lambda_0 = 1.5585 \text{ fm}^{-1}$
			$\mu_0 = 621.5 \text{ MeV}$
	$\hat{r}_0 \text{ (MeV fm)}$	28.170	43.882

TABLE II: Masses of baryon and meson in ChQM (unit: MeV).

P	$N(939)$	$\Delta(1232)$	$\Omega(1672)$	$\Lambda(1116)$	$\Sigma(1189)$	$\Xi(1315)$
set I						
+	936	1208	1643	1154	1173	1362
-	1575	1625	2203	1772	1777	1981
set II						
+	939	1231	1671	1187	1209	1408
-	1661	1716	2301	1889	1895	2098
	$\Sigma^*(1383)$	$\Xi^*(1532)$	$\Omega_c(2695)$	$\Omega_c(2765)$	$\Xi_c(2467)$	$\Xi_c^*(2645)$
set I						
+	1342	1488	2675	2748	2541	2603
-	1805	1999	3257	3282	3086	3093
set II						
+	1393	1539	2748	2818	2629	2727
-	1928	2119	3378	3389	3145	3166
set I						
P	$\pi(140)$	$\rho(775)$	$\eta(548)$	$\omega(782)$	$K(495)$	$K^*(892)$
-	93	800	611	705	326	965
	$\eta'(958)$	$\phi(1019)$	$D^0(1865)$	$D^*(2007)$		
-	914	1056	1842	2043		

are listed. One need to notice that there are many different ways to construct the wave-functions of the system. However, it makes no difference by choosing any one configuration if all the possible coupling are considered.

For the Ω_c^0 with quark content $ssc q \bar{q}$, $q = u, d, s$ in flavor SU(3) case, there are two types of separation, one is $(qss)\bar{q}c$ and the other is $(ssc)\bar{q}q$. The flavor wavefunc-

tions for the sub-clusters constructed are shown below.

$$\begin{aligned}
B_{00}^1 &= ssc, & B_{00}^2 &= sss, \\
B_{\frac{1}{2}, \frac{1}{2}}^1 &= \frac{1}{\sqrt{6}}(sus + uss - 2ssu), \\
B_{\frac{1}{2}, -\frac{1}{2}}^1 &= \frac{1}{\sqrt{6}}(sds + dss - 2ssd), \\
B_{\frac{1}{2}, \frac{1}{2}}^2 &= \frac{1}{\sqrt{2}}(us - su)s, \\
B_{\frac{1}{2}, -\frac{1}{2}}^2 &= \frac{1}{\sqrt{2}}(ds - sd)s, \\
B_{\frac{1}{2}, \frac{1}{2}}^3 &= \frac{1}{\sqrt{3}}(ssu + sus + uss), \\
B_{\frac{1}{2}, -\frac{1}{2}}^3 &= \frac{1}{\sqrt{3}}(ssd + sds + dss), \\
B_{\frac{1}{2}, \frac{1}{2}}^4 &= \frac{1}{\sqrt{2}}(us + su)c, \\
B_{\frac{1}{2}, -\frac{1}{2}}^4 &= \frac{1}{\sqrt{2}}(ds + sd)c, \\
B_{\frac{1}{2}, \frac{1}{2}}^5 &= \frac{1}{\sqrt{2}}(us - su)c, \\
B_{\frac{1}{2}, -\frac{1}{2}}^5 &= \frac{1}{\sqrt{2}}(ds - sd)c,
\end{aligned} \tag{4}$$

$$\begin{aligned}
M_{\frac{1}{2}, \frac{1}{2}}^1 &= \bar{d}c, & M_{\frac{1}{2}, -\frac{1}{2}}^1 &= -\bar{u}c, \\
M_{\frac{1}{2}, \frac{1}{2}}^2 &= \bar{d}s, & M_{\frac{1}{2}, -\frac{1}{2}}^2 &= -\bar{u}s, \\
M_{00}^1 &= \frac{1}{\sqrt{2}}(\bar{u}u + \bar{d}d), & M_{00}^2 &= \bar{s}s, & M_{00}^3 &= \bar{s}c.
\end{aligned} \tag{5}$$

The flavor wavefunctions for 5-quark system with isospin $I = 0$ are obtained by the following couplings,

$$\begin{aligned}
\chi_1^f &= \sqrt{\frac{1}{2}}(B_{\frac{1}{2}, \frac{1}{2}}^1 M_{\frac{1}{2}, -\frac{1}{2}}^1 - B_{\frac{1}{2}, -\frac{1}{2}}^1 M_{\frac{1}{2}, \frac{1}{2}}^1), \\
\chi_2^f &= \sqrt{\frac{1}{2}}(B_{\frac{1}{2}, \frac{1}{2}}^2 M_{\frac{1}{2}, -\frac{1}{2}}^2 - B_{\frac{1}{2}, -\frac{1}{2}}^2 M_{\frac{1}{2}, \frac{1}{2}}^2), \\
\chi_3^f &= \sqrt{\frac{1}{2}}(B_{\frac{1}{2}, \frac{1}{2}}^3 M_{\frac{1}{2}, -\frac{1}{2}}^3 - B_{\frac{1}{2}, -\frac{1}{2}}^3 M_{\frac{1}{2}, \frac{1}{2}}^3), \\
\chi_4^f &= \sqrt{\frac{1}{2}}(B_{\frac{1}{2}, \frac{1}{2}}^4 M_{\frac{1}{2}, -\frac{1}{2}}^4 - B_{\frac{1}{2}, -\frac{1}{2}}^4 M_{\frac{1}{2}, \frac{1}{2}}^4), \\
\chi_5^f &= \sqrt{\frac{1}{2}}(B_{\frac{1}{2}, \frac{1}{2}}^5 M_{\frac{1}{2}, -\frac{1}{2}}^5 - B_{\frac{1}{2}, -\frac{1}{2}}^5 M_{\frac{1}{2}, \frac{1}{2}}^5), \\
\chi_6^f &= B_{00}^1 M_{00}^1, \quad \chi_7^f = B_{00}^1 M_{00}^2, \quad \chi_8^f = B_{00}^2 M_{00}^3.
\end{aligned} \tag{6}$$

In a similar way, the spin and color wavefunctions for 5-quark system can be constructed, which are the same as the expressions of Ref. [6]. Here we only give the expressions of 5-quark system, the wavefunctions for the sub-clusters can be found in Ref. [6].

$$\begin{aligned}
\chi_{\frac{1}{2}, \frac{1}{2}}^{\sigma 1}(5) &= \sqrt{\frac{1}{6}}\chi_{\frac{3}{2}, -\frac{1}{2}}^{\sigma}(3)\chi_{11}^{\sigma} - \sqrt{\frac{1}{3}}\chi_{\frac{3}{2}, \frac{1}{2}}^{\sigma}(3)\chi_{10}^{\sigma} \\
&\quad + \sqrt{\frac{1}{2}}\chi_{\frac{3}{2}, \frac{3}{2}}^{\sigma}(3)\chi_{1-1}^{\sigma}
\end{aligned}$$

$$\begin{aligned}
\chi_{\frac{1}{2},\frac{1}{2}}^{\sigma_2}(5) &= \sqrt{\frac{1}{3}}\chi_{\frac{1}{2},\frac{1}{2}}^{\sigma_1}(3)\chi_{10}^\sigma - \sqrt{\frac{2}{3}}\chi_{\frac{1}{2},-\frac{1}{2}}^{\sigma_1}(3)\chi_{11}^\sigma \\
\chi_{\frac{1}{2},\frac{1}{2}}^{\sigma_3}(5) &= \sqrt{\frac{1}{3}}\chi_{\frac{1}{2},\frac{1}{2}}^{\sigma_2}(3)\chi_{10}^\sigma - \sqrt{\frac{2}{3}}\chi_{\frac{1}{2},-\frac{1}{2}}^{\sigma_2}(3)\chi_{11}^\sigma \\
\chi_{\frac{1}{2},\frac{1}{2}}^{\sigma_4}(5) &= \chi_{\frac{1}{2},\frac{1}{2}}^{\sigma_1}(3)\chi_{00}^\sigma \\
\chi_{\frac{1}{2},\frac{1}{2}}^{\sigma_5}(5) &= \chi_{\frac{1}{2},\frac{1}{2}}^{\sigma_2}(3)\chi_{00}^\sigma \\
\chi_{\frac{3}{2},\frac{3}{2}}^{\sigma_1}(5) &= \sqrt{\frac{3}{5}}\chi_{\frac{3}{2},\frac{3}{2}}^{\sigma_2}(3)\chi_{10}^\sigma - \sqrt{\frac{2}{5}}\chi_{\frac{3}{2},\frac{1}{2}}^{\sigma_2}(3)\chi_{11}^\sigma \\
\chi_{\frac{3}{2},\frac{3}{2}}^{\sigma_2}(5) &= \chi_{\frac{3}{2},\frac{3}{2}}^{\sigma_1}(3)\chi_{00}^\sigma \\
\chi_{\frac{3}{2},\frac{3}{2}}^{\sigma_3}(5) &= \chi_{\frac{1}{2},\frac{1}{2}}^{\sigma_1}(3)\chi_{11}^\sigma \\
\chi_{\frac{3}{2},\frac{3}{2}}^{\sigma_4}(5) &= \chi_{\frac{1}{2},\frac{1}{2}}^{\sigma_2}(3)\chi_{11}^\sigma \\
\chi_{\frac{5}{2},\frac{5}{2}}^{\sigma_1}(5) &= \chi_{\frac{3}{2},\frac{3}{2}}^{\sigma_1}(3)\chi_{11}^\sigma
\end{aligned} \tag{7}$$

$$\begin{aligned}
\chi_1^c &= \frac{1}{\sqrt{18}}(rgb - rbg + gbr - grb + brg - bgr) \\
&\quad (\bar{r}r + \bar{g}g + \bar{b}b), \\
\chi_k^c &= \frac{1}{\sqrt{8}}(\chi_{3,1}^k\chi_{2,8} - \chi_{3,2}^k\chi_{2,7} - \chi_{3,3}^k\chi_{2,6} + \chi_{3,4}^k\chi_{2,5} \\
&\quad + \chi_{3,5}^k\chi_{2,4} - \chi_{3,6}^k\chi_{2,3} - \chi_{3,7}^k\chi_{2,2} + \chi_{3,8}^k\chi_{2,2}), \tag{8}
\end{aligned}$$

with $k = 2, 3$. For the color part, both the color singlet channels ($k = 1$) and the hidden color channels ($k = 2, 3$), are considered here to have an economic way to describe multi-quark system [6].

For the orbital wavefunctions, there are four relative motions for 5-body system. In the present work, the orbital wavefunctions for each relative motion of the system are determined by the dynamics of the system, The orbital wavefunctions for this purpose is written as follows,

$$\psi_{LM_L} = [[[\phi_{n_1 l_1}(\boldsymbol{\rho})\phi_{n_2 l_2}(\boldsymbol{\lambda})]_l \phi_{n_3 l_3}(\mathbf{r})]_{l'} \phi_{n_4 l_4}(\mathbf{R})]_{LM_L}$$

where the Jacobi coordinates are defined as,

$$\begin{aligned}
\boldsymbol{\rho} &= \mathbf{x}_1 - \mathbf{x}_2, \\
\boldsymbol{\lambda} &= \mathbf{x}_3 - \left(\frac{m_1 \mathbf{x}_1 + m_2 \mathbf{x}_2}{m_1 + m_2}\right), \\
\mathbf{r} &= \mathbf{x}_4 - \mathbf{x}_5, \\
\mathbf{R} &= \left(\frac{m_1 \mathbf{x}_1 + m_2 \mathbf{x}_2 + m_3 \mathbf{x}_3}{m_1 + m_2 + m_3}\right) - \left(\frac{m_4 \mathbf{x}_4 + m_5 \mathbf{x}_5}{m_4 + m_5}\right).
\end{aligned} \tag{10}$$

To find the orbital wavefunctions, the gaussian expansion method (GEM) is employed, i.e., every ϕ is expanded by gaussians with various sizes [28]

$$\phi_{nlm}(\mathbf{r}) = \sum_{n=1}^{n_{max}} c_n N_{nl} r^l e^{-(r/r_n)^2} Y_{lm}(\hat{\mathbf{r}}), \tag{11}$$

where N_{nl} is the normalization constant,

$$N_{nl} = \left[\frac{2^{l+2} (2\nu_n)^{l+\frac{3}{2}}}{\sqrt{\pi} (2l+1)} \right]^{\frac{1}{2}}. \tag{12}$$

The size parameters of gaussians r_n are taken as the geometric progression numbers

$$r_n = r_1 a^{n-1}. \tag{13}$$

c_n is the variational parameters, which is determined by the dynamics of the system.

Finally, the complete channel wave function for the 5-quark system is written as

$$\Psi_{JM,i,j,k,n} = \mathcal{A} \left[[\chi_S^{\sigma_i}(5) \psi_L]_{JM_J} \chi_j^f \chi_k^c \right] \tag{14}$$

where \mathcal{A} is the antisymmetry operator of the system. In the flavor SU(3) case, it has six terms for the system with three identical particles and it can be reduced to three terms, as follows, due to the symmetry between first two particles has been considered when constructing the wavefunctions of the 3-quark clusters. For the two types of separations, 1-(uss)($\bar{u}c$) + (dss)($\bar{d}c$), (sss)($\bar{s}c$), 2-(ssc)($\bar{u}u + \bar{d}d$), (ssc)($\bar{s}s$), we have the following anti-symmetric operators,

$$\mathcal{A}_1 = 1 - (13) - (23), \tag{15}$$

$$\mathcal{A}_2 = 1 - (15) - (25). \tag{16}$$

The eigen-energy of the system is obtained by solving the following eigen-equation

$$H\Psi_{JM} = E\Psi_{JM}, \tag{17}$$

by using variational principle. The eigen functions Ψ_{JM} are the linear combination of the above channel wavefunctions Eq.(14).

In evaluating the matrix elements of hamiltonian, the calculation is rather complicated, if the orbital angular momenta of relative motions of system are not all zero. Here a useful method named the infinitesimally-shifted gaussian are used [28]. In this method, the spherical harmonic function is absorbed into the shifted gaussians,

$$\phi_{nlm}(\mathbf{r}) = N_{nl} \lim_{\varepsilon \rightarrow 0} \frac{1}{(\nu\varepsilon)^l} \sum_{k=1}^{k_{max}} C_{lm,k} e^{-\nu_n(\mathbf{r}-\varepsilon\mathbf{D}_{lm,k})^2},$$

the calculation becomes easy with no tedious angular-momentum algebra required.

III. RESULTS AND DISCUSSIONS

In the present calculation, we are interested in the low-lying states of $ussc\bar{u}$, $dssc\bar{d}$ pentaquark system, so all the orbital angular momenta are set to 0. Then the parity of five-quark system with one antiquark is negative. In this way, the total angular momentum J can take values 1/2, 3/2 and 5/2. The possible channels under the consideration are listed in Tables III-V.

First, the single channel calculations are performed. The eigen-energies of each states with different quantum numbers are shown in Tables VI-X, where the eigen-energies of the states are shown in column 2, along with

TABLE III: The channels with $IJ^P = 0\frac{1}{2}^-$.

index	$\chi_{1/2}^{\sigma_i}$	χ_j^f	χ_k^c	physical channel
1	$i = 1$	$j = 3$	$k = 1$	$\Xi^* \bar{D}^*$
2	$i = 1$	$j = 3$	$k = 3$	
3	$i = 1$	$j = 4$	$k = 1$	$\Xi_c^* \bar{K}^*$
4	$i = 1$	$j = 4, 5$	$k = 2, 3$	
5	$i = 1$	$j = 6$	$k = 1$	$\Omega_c^* \omega$
6	$i = 1$	$j = 6$	$k = 3$	
7	$i = 2, 3$	$j = 1, 2$	$k = 1$	$\Xi \bar{D}^*$
8	$i = 2, 3$	$j = 1, 2$	$k = 2, 3$	
9	$i = 2, 3$	$j = 4, 5$	$k = 1$	$\Xi_c \bar{K}^*$
10	$i = 2, 3$	$j = 4, 5$	$k = 2, 3$	
11	$i = 2$	$j = 6$	$k = 1$	$\Omega_c \omega$
12	$i = 2, 3$	$j = 6$	$k = 2, 3$	
13	$i = 4, 5$	$j = 1, 2$	$k = 1$	$\Xi \bar{D}$
14	$i = 4, 5$	$j = 1, 2$	$k = 2, 3$	
15	$i = 4, 5$	$j = 4, 5$	$k = 1$	$\Xi_c \bar{K}$
16	$i = 4, 5$	$j = 4, 5$	$k = 2, 3$	
17	$i = 4$	$j = 6$	$k = 1$	$\Omega_c \eta$
18	$i = 4, 5$	$j = 6$	$k = 2, 3$	

the theoretical thresholds in column 3 and experimental thresholds in column 5, column 4 gives the binding energies, the difference between the eigen-energies and the theoretical thresholds, $E_B = E - E_{th}^{Theo}$. The corrected energies of the states (column 6), which are obtained by taking the sum of experimental thresholds and the binding energies. Namely, and $E' = E_B + E_{th}^{Exp}$.

Secondly, the three types of channel coupling calculations are performed. The first is the channel coupling between color-singlet and hidden-color channels with the same flavor-spin structures. The second is the coupling among all color-singlet channels with different flavor-spin structures and the last is the full coupling, including all channels for given J^P . Table XI gives the spacial configurations of the states by calculating the distances between any two quarks or quark and antiquark in the full channel coupling calculation.

In the following we analyze the results in detail.

(a) $J^P = \frac{1}{2}^-$: The single channel calculations show that there are weak attractions for the most channels, the exceptions are $\Omega_c \eta, \Omega_c \omega, \Omega_c^* \omega$ and $\Xi_c \bar{K}$. The coupling to hidden-color channels helps a little, increasing the attraction a few MeVs and pushing $\Omega_c^* \omega$ and $\Xi_c \bar{K}$ below the corresponding thresholds. So the resonances can be formed. Most of the states have higher masses compared with that of the five new excited states of Ω_c . For $\Xi \bar{D}$, the second lowest state, it has the energy 3156 MeV, which is close to the highest Ω_c , 3119 MeV. The lowest state $\Xi_c \bar{K}$ has the energy 2949 MeV with the help of hidden-color channel coupling, which is a little smaller than the mass of the lowest excited state of Ω_c , 3000 MeV.

TABLE IV: The channels with $IJ^P = 0\frac{3}{2}^-$.

index	$\chi_{3/2}^{\sigma_i}$	χ_j^f	χ_k^c	physical channel
1	$i = 1$	$j = 3$	$k = 1$	$\Xi^* \bar{D}^*$
2	$i = 1$	$j = 3$	$k = 3$	
3	$i = 1$	$j = 4$	$k = 1$	$\Xi_c^* \bar{K}^*$
4	$i = 1$	$j = 4, 5$	$k = 2, 3$	
5	$i = 1$	$j = 6$	$k = 1$	$\Omega_c^* \omega$
6	$i = 1$	$j = 6$	$k = 3$	
7	$i = 2$	$j = 3$	$k = 1$	$\Xi^* \bar{D}$
8	$i = 2$	$j = 3$	$k = 3$	
9	$i = 2$	$j = 4$	$k = 1$	$\Xi_c^* \bar{K}$
10	$i = 2$	$j = 4, 5$	$k = 2, 3$	
11	$i = 2$	$j = 6$	$k = 1$	$\Omega_c^* \eta$
12	$i = 2$	$j = 6$	$k = 3$	
13	$i = 3, 4$	$j = 1, 2$	$k = 1$	$\Xi \bar{D}^*$
14	$i = 3, 4$	$j = 1, 2$	$k = 2, 3$	
15	$i = 3, 4$	$j = 4, 5$	$k = 1$	$\Xi_c \bar{K}^*$
16	$i = 3, 4$	$j = 4, 5$	$k = 2, 3$	
17	$i = 3$	$j = 6$	$k = 1$	$\Omega_c \omega$
18	$i = 3, 4$	$j = 6$	$k = 2, 3$	

TABLE V: The channels with $IJ^P = 0\frac{5}{2}^-$.

index	$\chi_{5/2}^{\sigma_i}$	χ_j^f	χ_k^c	physical channel
1	$i = 1$	$j = 3$	$k = 1$	$\Xi^* \bar{D}^*$
2	$i = 1$	$j = 3$	$k = 3$	
3	$i = 1$	$j = 4$	$k = 1$	$\Xi_c^* \bar{K}^*$
4	$i = 1$	$j = 4, 5$	$k = 2, 3$	
5	$i = 1$	$j = 6$	$k = 1$	$\Omega_c^* \omega$
6	$i = 1$	$j = 6$	$k = 3$	

The situation changes a lot after coupling all the color-singlet channels, the lowest energy we obtained is 2865 MeV. And the full channel-coupling calculation decreases the lowest energy further to 2769 MeV. Table VII shows the six lowest eigen-energies in the full-channel calculation. E' denotes the corrected energy,

$$E' = E_{th}^{Exp}(\Xi_c \bar{K}) - E_{th}^{Theo}(\Xi_c \bar{K}) + E.$$

In this way, there are many Ω_c pentaquark states with $IJ^P = 0\frac{1}{2}^-$ in the quark model. To assign these states to the excited Ω_c states announced by LHCb, further work is needed. The problem has to be solved is how to correct the eigen-energies from the full channel-coupling calculation.

One interesting state is $\Omega_c^* \omega$, the hidden-color channel has lower energy than the colorless one. It is a possible good resonance because of its color structure, although it has a rather high energy, 3497 MeV.

(b) $J^P = \frac{3}{2}^-$: We have similar results with that of $J^P = \frac{1}{2}^-$. Four channels, $\Omega_c^* \omega, \Xi_c^* \bar{K}, \Xi_c \bar{K}^*$ and $\Omega_c \omega$, have no attraction in single channel calculations. and the hidden-color channel-coupling induces a very weak

TABLE VI: The lowest eigen-energies of the $udc\bar{c}u$ system with $J^P = \frac{1}{2}^-$ (unit: MeV). The percentages of color-singlet (S) and hidden-color (H) channels are also given.

Channel	E	E_{th}^{Theo}	E_B	E_{th}^{Exp}	E'
1	3526	3531	-5	3539($\Xi^* \bar{D}^*$)	3534
2	4016				
1+2	3525		-6		3533
percentage(S;H): 99.8%; 0.2%					
3	3566	3568	-2	3537($\Xi_c^* \bar{K}^*$)	3535
4	3616				
3+4	3564		-4		3533
percentage(S;H): 96.3%; 3.7%					
5	3453	3453	0	3548($\Omega_c^* \omega$)	3453
6	3404				
5+6	3402		-51		3497
percentage(S;H): 0.2%; 99.8%					
7	3374	3405	-31	3322($\Xi \bar{D}^*$)	3291
8	3672				
7+8	3373		-32		3290
percentage(S;H): 99.8%; 0.2%					
9	3495	3506	-11	3359($\Xi_c \bar{K}^*$)	3348
10	3613				
9+10	3472		-34		3325
percentage(S;H): 85.2%; 14.8%					
11	3380	3380	0	3477($\Omega_c \omega$)	3477
12	3608				
11+12	3380				
13	3175	3204	-29	3185($\Xi \bar{D}$)	3156
14	3811				
13+14	3175		-29		3156
percentage(S;H): 100.0%; 0.0%					
15	2867	2867	0	2961($\Xi_c \bar{K}$)	2961
16	3807				
15+16	2855		-12		2949
percentage(S;H): 96.7%; 3.3%					
17	3286	3286	0	3243($\Omega_c \eta$)	3243
18	3828				
17+18	3286				
mixed (singlet)	2771	2867	-96	2961($\Xi_c \bar{K}$)	2865
mixed (full)	2675	2867	-192	2961($\Xi_c \bar{K}$)	2769

attraction for $\Xi_c^* \bar{K}$. But, it introduces a large attraction for $\Xi_c \bar{K}^*$, -158 MeV, a good candidate of color structure resonance to be confirmed.

All color-singlet channel-coupling calculation gives a very weak bound state with energy 3138 MeV after correction. The full channel-coupling lowered the energy further to 3067 MeV. Table IX shows the four lowest eigen-energies in the full-channel coupling calculation. After correction, their energies are below 3.2 GeV.

(c) $J^P = \frac{5}{2}^-$: Only one channel, $\Xi^* \bar{D}^*$, has attractive in the single channel calculation. Coupling to the hidden-

TABLE VII: The eigen-energies of full channel-coupling calculation below 3.2 GeV with $IJ^P = 0\frac{1}{2}^-$. (unit: MeV).

index	1	2	3	4	5	6
E	2675	2867	2873	2882	2901	2937
E'	2769	2961	2967	2976	2995	3031

TABLE VIII: The lowest eigen-energies of the $udc\bar{c}u$ system with $\frac{3}{2}^-$ (unit: MeV).

Channel	E	E_{th}^{Theo}	E_B	E_{th}^{Exp}	E'
1	3521	3531	-10	3539($\Xi^* \bar{D}^*$)	3529
2	4026				
1+2	3521		-10		3529
percentage(S;H): 100.0%; 0.0%					
3	3565	3568	-3	3537($\Xi_c^* \bar{K}^*$)	3534
4	3617				
3+4	3562		-6		3531
percentage(S;H): 94.0%; 6.0%					
5	3453	3453	0	3548($\Omega_c^* \omega$)	3548
6	3477				
5+6	3453				
7	3309	3330	-21	3397($\Xi^* \bar{D}$)	3376
8	4145				
7+8	3309		-21		3376
percentage(S;H): 100.0%; 0.0%					
9	2929	2929	0	3139($\Xi_c \bar{K}$)	3139
10	3782				
9+10	2728		-1		3138
percentage(S;H): 99.7%; 0.3%					
11	3359	3359	0	3314($\Omega_c^* \eta$)	3314
12	3763				
11+12	3359				
13	3388	3405	-17	3322($\Xi \bar{D}^*$)	3305
14	3705				
13+14	3388		-17		3305
percentage(S;H): 100.0%; 0.0%					
15	3506	3506	0	3359($\Xi_c \bar{K}^*$)	3359
16	3656				
15+16	3348		-158		3201
percentage(S;H): 57.7%; 42.3%					
17	3380	3380	0	3477($\Omega_c \omega$)	3477
18	3588				
17+18	3380				
mixed (singlet)	2928	2929	-1	3139($\Xi_c \bar{K}$)	3138
mixed (full)	2857	2929	-72	3139($\Xi_c \bar{K}$)	3067

color channels, an additional channel, $\Xi_c^* \bar{K}^*$, is induced out an attraction. Channel-couplings, color-singlet and full, do not produce any bound state. The D -wave Ξ - \bar{D} and/or Ξ_c - \bar{K} scattering phase shift calculation is needed to check that the resonances, $\Xi^* \bar{D}^*$ and $\Xi_c^* \bar{K}^*$, can survive or not after the coupling.

TABLE IX: The eigen-energies of full channel-coupling calculation below 3.2 GeV with $IJ^P = 0\frac{3}{2}^-$. (unit: MeV).

index	1	2	3	4
E	2857	2931	2940	2956
E'	3067	3141	3150	3166

TABLE X: The lowest eigen-energies of the $ssc\bar{u}u + ssc\bar{d}d$ system with $\frac{5}{2}^-$ (unit: MeV).

Channel	E	E_{th}^{Theo}	E_B	E_{th}^{Exp}	E'
1	3508	3531	-23	3539($\Xi^* \bar{D}^*$)	3516
2	4042				
1+2	3507		-24		3515
percentage(S;H): 99.8%; 0.2%					
3	3568	3568	0	3537($\Xi_c^* \bar{K}^*$)	3537
4	3646				
3+4	3532		-36		3501
percentage(S;H): 80.0%; 20.0%					
5	3453	3453	0	3548($\Omega_c^* \omega$)	3548
6	3563				
5+6	3453				
mixed (singlet)	3453				
mixed (full)	3453				

TABLE XI: Distances between quarks, q is for u, d quark and Q is for c quark (unit: fm).

J^P	Channel	r_{qq}	r_{qQ}	$r_{q\bar{q}}$	$r_{Q\bar{q}}$
$\frac{1}{2}^-$	$\Omega_c^0(2769)$	1.3	1.1	1.4	1.2
$\frac{3}{2}^-$	$\Omega_c^0(3067)$	1.4	1.1	1.2	1.4

Table XI gives the distances between quarks for two states, $\Omega_c^0(2769)$ and $\Omega_c^0(3067)$. All the quark-pairs have similar distances and all are smaller than 1.5 fm. So these two states are compact ones.

IV. SUMMARY

In the framework of the chiral quark model, the 5-quark systems with quark contents $ssc\bar{u}u$, $ssc\bar{d}d$ are investigated by means of Gaussian expansion method. The

calculation shows that there are several resonance states for $I(J^P) = 0(\frac{1}{2}^-)$, $0(\frac{3}{2}^-)$ below 3.2 GeV. $\Xi\bar{D}$, $\Xi_c\bar{K}$ and $\Xi_c^*\bar{K}$ are possible the candidates of the newly announced excited states of Ω_c^0 by LHCb Collaboration. In the present calculation, the masses of the lowest states with quantum numbers $IJ^P = 0\frac{1}{2}^-$ and $IJ^P = 0\frac{3}{2}^-$ are 2769 MeV and 3067 MeV, respectively. And the distances between quark pairs suggest these two states are compact states or pentaquark structures. It manifests the effects of hidden-color channels. So it is interesting to identify the states experimentally. In this work, in fact we cannot identify the excited states of Ω_c^0 reported by LHCb Collaboration with the pentaquarks we calculated. We want stress that the P -wave q^3 baryon will mix strongly with the S -wave pentaquark. The unquenched quark model, including the high Fock components, study of Ω_c is needed to clarify the situation.

In the present calculation, the internal structures of the sub-clusters are not fixed, the structure of a 5-quark system is determined by the dynamics of the system, because all the possible coupling are included except the high orbital angular momentum. The further work of considering the high orbital angular momenta along with the spin-orbit and tensor interactions is expected.

Pentaquark involves two subcluster, q^3 and $q\bar{q}$. If the two subclusters are colorless, they are corresponding to baryon and meson. To describe baryon and meson simultaneously in quark model with one set of parameters is still difficult. It is main source of the uncertainty of the model calculation of pentaquark. Unquenched quark model may be a solution for the unified description of baryon and meson, since the $q\bar{q}$ cluster is always involved.

Multiquark states are ideal place to develop the quark model. Because the model approach is a phenomenological one, its development depends on the accumulated experimental data. We hope that the model description of the multiquark states will be improved with the accumulation of the experimental data on multiquark state,

Acknowledgments

The work is supported partly by the National Natural Science Foundation of China under Grant Nos. 11535005, 11175088, and 11205091.

[1] R. Aaij et al. [LHCb Collaboration], arXiv:1703.04639 [hep-ex].
[2] S. S. Agaev, K. Azizi and H. Sundu, arXiv:1703.07091 [hep-ex].
[3] H. X. Chen, Q. Mao, W. Chen, A. Hosaka, X. Liu and S. L. Zhu, arXiv:1703.07703 [hep-ph].
[4] M. Karliner and J. L. Rosner, arXiv:1703.07774 [hep-ph].
[5] R. Aaij et al. [LHCb Collaboration], Phys. Rev. Lett. **115**, 072001 (2015).
[6] G. Yang, J. L. Ping and F. Wang, Phys. Rev. D **95**,

014010 (2017).
[7] H. X. Huang, C. R. Deng, J. L. Ping and F. Wang, Eur. Phys. J. C **76**, 624 (2016).
[8] R. Chen, X. Liu, X. Q. Li and S. L. Zhu, Phys. Rev. Lett. **115**, 132002 (2015).
[9] J. He, Phys. Lett. B **753**, 547 (2016).
[10] H. X. Chen, W. Chen, X. Liu, T. G. Steel and S. L. Zhu, Phys. Rev. Lett. **115**, 172001 (2015).
[11] Z. G. Wang, Eur. Phys. J. C **76**, 70 (2016).
[12] L. Roca, J. Nieves and E. Oset, Phys. Rev. D **92**, 094003

- (2015).
- [13] R. L. Zhu, C. F. Qiao, Phys. Lett. B **756**, 259 (2016).
 - [14] X. H. Liu, Q. Wang and Q. Zhao, Phys. Lett. B **757**, 231 (2016).
 - [15] F. K. Guo, U. G. Meißner, W. Wang and Z. Yang, Phys. Rev. D **92** 071502 (2015).
 - [16] H. X. Chen, W. Chen, X. Liu and S. L. Zhu, Phys. Rep. **639**, 1 (2016).
 - [17] T. Nakano *et al.* [LEPS Collaboration], Phys. Rev. Lett. **91**, 012002 (2003).
 - [18] V. V. Barmin *et al.* [DIANA Collaboration], Yad. Fiz. **66**, 1763 (2003) [Phys. At. Nucl. **66**, 1715 (2003)].
 - [19] S. Stepanyan *et al.* [CLAS Collaboration], Phys. Rev. Lett. **91**, 252001 (2003).
 - [20] M. Battaglieri *et al.* [CLAS Collaboration], Phys. Rev. Lett. **96**, 042001 (2006) and references therein.
 - [21] T. Nakano *et al.* [LEPS Collaboration], Phys. Rev. C **79**, 025210 (2009).
 - [22] R. Bijker and E. Santopinto, Phys. Rev. C **80**, 065210 (2009).
 - [23] E. Santopinto and R. Bijker, Phys. Rev. C **82**, 062202(R) (2010).
 - [24] C. S. An and B. S. Zou, Eur. Phys. J. A **39**, 195 (2009).
 - [25] D. Qing, X.S. Chen and F. Wang, Phys. Rev. C **57**, R31 (1998), D **58**, 114032(998).
 - [26] B. S. Zou, Chin. Phys. C (High Ener. Phys. Nucl. Phys.), **33**, 1113 (2009).
 - [27] S. G. Yuan, C. S. An, K. W. Wei, B. S. Zou and H. S. Xu, Phys. Rev. C **87**, 025205 (2013).
 - [28] E. Hiyama, Y. Kino and M. Kamimura, Prog. Part. Nucl. Phys. **51**, 223 (2003).
 - [29] J. Vijande, F. Fernandez and A. Valcarce, J. Phys. G **31**, 481 (2005).
 - [30] G. S. Bali, Phys. Rev. D **62**, 114503 (2000).
 - [31] J.-M. Richard, Phys. Rev. C **81**, 015205 (2010); M. W. Paris, Phys. Rev. Lett. **95**, 202002 (2005); C. R. Deng, J. L. Ping, F. Wang and T. Goldman, Phys. Rev. D **82**, 074001 (2010).
 - [32] M. Harvey, Nucl. Phys. A **352**, 301 (1981); 326 (1981)
 - [33] F. Wang, Prog. Phys. **9**, 297 (1989) (in Chinese).
 - [34] J. Vijande, A. Valcarce and N. Barnea, Phys. Rev. D **79**, 074010 (2009); J. Vijande and A. Valcarce, Phys. Rev. C **80**, 035204 (2009); D. Janc and M. Rosina, Few Body Syst. **35**, 175 (2004).