

# Properties of Low-Lying Charmonium States in a Phenomenological Approach

Praveen P D'Souza <sup>1,2</sup> and Antony Prakash Monteiro <sup>1\*</sup>

<sup>1</sup>*P. G. Department of Physics, St Philomena college Darbe, Puttur 574 202, India*

K. B. Vijaya Kumar <sup>2</sup>

<sup>2</sup>*Department of Physics, Mangalore University,  
Mangalagangothri P.O., Mangalore - 574199, INDIA*

(Dated: March 11, 2022)

We investigate the spectrum and decay rates of low lying charmonium states within the framework of the non relativistic quark model by employing a Coulomb like potential from the perturbative one gluon exchange and the linear confining potential along with the potential derived from instanton vacuum to account for the hyperfine mass splitting of charmonium states in variational approach. We predict radiative E1, M1, two-photon, leptonic and two-gluon decay rates of low lying charmonium states. An overall agreement is obtained with the experimental masses and decay widths.

## I. INTRODUCTION

Charmonia are bound states of a charm and an anticharm quark ( $c\bar{c}$ ), and represent an important testing ground for the properties of the strong interaction. There has been a great progress in the observation of the charmonium states from the past few years. The discovery of the first charmonium state  $J/\psi$ [1, 2] has revolutionized the field of hadron spectroscopy. This led to a clear understanding of the prevailing theory of particle physics. Several quarkonium states have been observed after the discovery of the charmonium state  $J/\psi$  at BNL and SLAC[3]. The first observation of a singlet ground state of charmonium  $\eta_c$  was done by Mark II and Crystal Ball experiments in the radiative decays of  $J/\psi$  and  $\psi'$ [3]. The discoveries of conventional states  $h_c(1P)$ ,  $h_c(2P)$ ,  $\chi_c(1P)$ ,  $\chi_c(2P)$ ,  $\eta_c(1S)$  and the observation of the exotic states like X(3872), X(3915), Y(4260), Z(3930) at Belle, BaBar, LHC, BESIII, CLEO, etc have created a renewed interest in quarkonium physics[3]. These new observations have given a deeper understanding of the charmonium physics and have unraveled many mysteries[4]. Charmonium system is a powerful tool for the study of forces between quarks in QCD in non-perturbative regime. Studies of charmonia production can improve our understanding of heavy quark production and the formation of bound states.

The exploration and understanding of the substructure of hadrons, presented in terms of quarks and gluons by quantum chromodynamics(QCD), has led to a considerable progress in the study of charmonium states. Though there have been numerous calculations of charmonium spectra from first principles such as Lattice QCD[5, 6] and NRQCD which provide rigorous theoretical implications for the experimental observations, quark model calculations provide more intuitive insights and supply important phenomenological guidance towards their study.

The quark antiquark potential cannot be obtained from the first principles of QCD. Therefore, one has to use potential models to explain the observed hadronic properties. The QCD inspired potential models have been playing an important role in investigating heavy quarkonium. Most of the quark potential models [7–18] have common ingredients in the non relativistic quark models(NRQM). Recently it is shown that the NRQM with instanton induced interaction(III) explains  $^1S_0$  and  $^3S_1$  the nucleon- nucleon potential[19].

The NRQM formalism provides a systematic treatment of the perturbative and non perturbative components of QCD at hadronic scale [20, 21]. These QCD inspired potential models use a short range part motivated by perturbative QCD (Coulomb like or one gluon exchange potential) [20, 22–30] and a phenomenological long range part accounting for confinement (i.e. linear, logarithmic or quadratic potential)[31–38]. Our Model uses the NRQM formalism for the study of low lying charmonium states using a Hamiltonian which has the heavy quark potential from the instanton vacuum depending on  $r$ , the inter quark distance. The heavy quark potential derived from the instanton ensemble rises linearly as the relative distance between the quark and antiquark increases, then it gets saturated. As the quark and the antiquark distance increases the central potential turns out to be Coulomb like potential. Therefore to study the mass spectra of the quarkonia we have added Coulombic type potential to central instanton potential. This can be understood as a non perturbative contribution to the perturbative potential from instanton vacuum at large inter quark distances greater than the instanton size. At large distance the instanton vacuum cannot explain quark confinement, hence in our phenomenological potential model we have included a linear confinement potential. Taking into consideration the above factors, we have developed a nonrelativistic potential model to obtain a reliable estimate of the masses of the low lying charmonium states and their decay widths.

The paper is organized in 4 sections. In sec. II we briefly review the theoretical background for the non rel-

---

\* aprakashmonteiro@gmail.com

ativistic model. In sec. III we present the results and discussions. In sec. IV we draw up conclusions.

## II. THEORETICAL BACKGROUND

### A. The Model

In a potential model approach the entire dynamics of quarks in a meson is governed by a Hamiltonian which is composed of a kinetic energy term  $K$  and a potential energy term  $V$ , that is,

$$H = K + V. \quad (1)$$

The kinetic energy  $K$  is given by,

$$K = M + \frac{p^2}{2\mu} \quad (2)$$

Here  $p$  is the relative momentum,  $\mu = \frac{m_Q m_{\bar{Q}}}{m_Q + m_{\bar{Q}}}$  is the reduced mass of the  $Q\bar{Q}$  system, where  $m_Q$  and  $m_{\bar{Q}}$  are the masses of the individual quark and anti quark respectively and  $M$  is the total mass of quark and antiquark[39].

The potential energy  $V$  is the sum of the heavy-quark potential  $V_{Q\bar{Q}}(\vec{r})$ , confining potential  $V_{conf}(\vec{r})$  and Coulomb potential  $V_{coul}(\vec{r})$ , that is

$$V(\vec{r}) = V_{Q\bar{Q}}(\vec{r}) + V_{coul}(\vec{r}) + V_{conf}(\vec{r}) \quad (3)$$

The heavy-quark potential  $V_{Q\bar{Q}}(\vec{r})$  is,

$$V_{Q\bar{Q}}(\vec{r}) = V_C(\vec{r}) + V_{SD}(\vec{r}). \quad (4)$$

Here  $V_C(\vec{r})$  and  $V_{SD}(\vec{r})$  are central and spin dependent potentials due to instanton vacuum respectively[40].

$V_C(\vec{r})$  is given by the following expression

$$V_C(\vec{r}) \simeq \frac{4\pi\bar{\rho}^3}{R^4 N_c} \left( 1.345 \frac{r^2}{\bar{\rho}^2} - 0.501 \frac{r^4}{\bar{\rho}^4} \right) \quad (5)$$

Here,  $\rho = \frac{1}{3}$  fm the average size of the instanton,  $\bar{R} = 1$  fm the average separation between instantons and number of colors  $N_c$  is 3.

The spin-spin interaction  $V_{SS}(\vec{r})$ , the spin-orbit coupling term  $V_{LS}(\vec{r})$  and the tensor part  $V_T(\vec{r})$  contribute to the spin dependent potential;

$$\begin{aligned} V_{SS}(\vec{r}) &= \frac{1}{3m_Q^2} \nabla^2 V_C(\vec{r}); \quad V_{LS}(\vec{r}) = \frac{1}{2m_Q^2} \frac{1}{r} \frac{dV_C(\vec{r})}{dr}; \\ V_T(\vec{r}) &= \frac{1}{3m_Q^2} \left( \frac{1}{r} \frac{dV_C(\vec{r})}{dr} - \frac{d^2 V_C(\vec{r})}{dr^2} \right). \end{aligned} \quad (6)$$

The coulomb-like (perturbative) one gluon exchange part of the potential is given by

$$V_{coul}(\vec{r}) = \frac{-4\alpha_s}{3r} \quad (7)$$

with the strong coupling constant  $\alpha_s$  and inter quark distance  $r$ .

The confinement term represents the non perturbative effect of QCD which includes the spin-independent linear confinement term[39]

$$V_{conf}(\vec{r}) = - \left[ \frac{3}{4} V_0 + \frac{3}{4} cr \right] F_1 \cdot F_2 \quad (8)$$

where  $c$  and  $V_0$  are constants.  $F$  is related to the Gell-Mann matrix,  $F_1 = \frac{\lambda_1}{2}$  and  $F_2 = \frac{\lambda_2^*}{2}$  and  $F_1 \cdot F_2 = \frac{-4}{3}$  for the mesons.

### B. Radiative Transitions

The study of radiative transitions can help in understanding the theory of strong interaction in the nonperturbative regime of QCD. The non-relativistic treatment adopted for the study of charmonium systems allows us to apply the usual multi-pole expansion in electrodynamics to compute the transition between the quarkonia states with the emission of a photon. The lowest order of multi-pole expansion tends to dominate the transition. The resulting transitions are the magnetic dipole  $M_1$  and electric dipole  $E_1$  transitions. The electric and magnetic dipole operators are decomposed by a multi-pole expansion into components with definite spherical tensor ranks.

In the phenomenological potential model the expression used for the radiative decay width does not directly influence the structure of the potential model Hamiltonian. Thus the transition probabilities are influenced by the chosen Hamiltonian only through the chosen wave function. In a  $M_1$  transition only the spin of the quarkonium state is changed, while the parity and the orbital angular momentum remain unchanged. The expression for the decay width of a spin flip  $M_1$  transition between heavy quarkonium states depend on the radial matrix. The different radial matrix elements can be obtained from the corresponding spin-flip magnetic moment operator. The  $M_1$  decay widths of charmonium are calculated using an assumption that the effective confining interaction is purely scalar. In electric dipole  $E_1$  transition, the parity of the states changes while spin remains unchanged. The one gluon exchange contribution survives in the  $E_1$  transition, whereas the contribution from the scalar confining interaction term vanishes.

In calculating the radiative decay widths, we have assumed that in the non relativistic limit, the dipole radial matrix elements are independent of  $J$ , i.e all states within the same angular momentum multiplet have the same wave function [41]. Radiative transitions could play an important role in the discovery and identification of charmonium states. They are sensitive to the internal structure of states, in particular to  $^3L_L - ^1L_L$  mixing for states with  $J = L$ . For our study of low lying charmonium states we have taken  $E_1$  and  $M_1$  radiative transitions

since the other order transitions contribute a little to the radiative decays. The partial width for an  $E_1$  radiative transition is given by,

$$\Gamma(i \rightarrow f + \gamma) = \frac{4\alpha e_c^2}{3} (2J_f + 1) S_{if}^E k_0^3 |\mathcal{E}_{if}|^2 \quad (9)$$

where  $k_0 = m_i - m_f$  is the energy of the emitted photon,  $\alpha$  is the fine structure constant.  $e_c = 2/3$  is the charge of the  $c$  quark in units of  $|e|$ ,  $m_i$  and  $m_f$  are the masses of initial and final mesons,  $S_{if}^E = \max(L_i, L_f) \left\{ \begin{matrix} J_i & 1 & J_f \\ L_f & S & L_i \end{matrix} \right\}^2$  is the statistical factor,  $J_i$  and  $J_f$  are the total angular momenta of initial and final mesons,  $L_i$  and  $L_f$  are the orbital angular momenta of initial and final mesons and  $S$  is the spin of the initial meson. The radial overlap integral which has the dimension of length is,

$$\mathcal{E}_{if} = \frac{3}{k_0} \int_0^\infty r^3 R_{nl}(r) R'_{nl}(r) dr \left[ \frac{k_0 r}{2} j_0 \left( \frac{k_0 r}{2} \right) - j_1 \left( \frac{k_0 r}{2} \right) \right] \quad (10)$$

with  $R_{nl}(r)$  and  $R'_{nl}(r)$  as the normalized radial wave functions for the corresponding states and  $j_0$  and  $j_1$  are spherical Bessel functions.

The  $M_1$  transitions between S-wave  $c\bar{c}$  states are given in the non relativistic approximation by [42–48],

$$\Gamma(i \rightarrow f + \gamma) = \frac{4\alpha e_c^2}{3m_c^2} \frac{2J_f + 1}{2L_i + 1} \delta_{L_i L_f} \delta_{S_i S_f} k_0^3 |M_{if}(r)|^2 \quad (11)$$

Here  $M_{if}$  is the radial overlap integral which has the dimension of length,

$$M_{if} = \int_0^\infty 4\pi r^3 R_{nl}(r) j_0(kr/2) R'_{nl}(r) dr. \quad (12)$$

In the overlap integral for unit operator between the coordinate wave functions of the initial and the final meson states,  $j_0(kr/2)$  is the spherical Bessel function,  $m_c$  is the mass of charm quark,  $J_f$  is the total angular momentum of final meson state,  $L_i$  is the orbital angular momentum of the initial meson state.

### C. Annihilation Decays

The annihilation decays of charmonium states into gluons and light quarks make significant contributions to the total decay width of the states. The annihilation decays allow us to determine wave function at the origin. The annihilation decays of some  $c\bar{c}$  states into photons can be used as a tool for the production and identification of the resonances.

#### 1. Two Photon Decays

Two-photon branching fraction for the charmonium provides a probe for the strong coupling constant at the

charmonium scale via the two-photon decay width. This can be utilized as a sensitive test for the corrections for the non-relativistic approximation in the potential models or in the effective field theories such as non relativistic QCD (NRQCD). The two-photon decays of P-wave charmonia are helpful for better understanding the nature of inter-quark forces and decay mechanisms.

The  $q\bar{q}$  quark pair in charge conjugation even states with  $J \neq 1$  can annihilate into two photons. The expressions for the decay rates of  $n^1S_0$ ,  $n^3P_0$  and  $n^3P_2$  states into two photons with the first order QCD radiative corrections are given by [49].

$$\Gamma(n^1S_0 \rightarrow \gamma\gamma) = \frac{3e_c^4 \alpha^2}{m_c^2} |R_{nS}(0)|^2 \left( 1 - \frac{3.4\alpha_s}{\pi} \right) \quad (13)$$

$$\Gamma(n^3P_0 \rightarrow \gamma\gamma) = \frac{27e_c^4 \alpha^2}{m_c^4} |R'_{nP}(0)|^2 \left( 1 + \frac{0.2\alpha_s}{\pi} \right) \quad (14)$$

$$\Gamma(n^3P_2 \rightarrow \gamma\gamma) = \frac{36e_c^4 \alpha^2}{5m_c^4} |R'_{nP}(0)|^2 \left( 1 - \frac{16\alpha_s}{\pi} \right) \quad (15)$$

The two photon decay widths of P wave charmonium states depend on the derivative of the radial wave function at the origin.

#### 2. Two Gluon Decays

The even states in charge conjugation of quarkonium with  $J \neq 1$  can annihilate into two gluons, much in the same way as they decay into two photons. The charmonium states  $^1S_0$ ,  $^3P_0$ ,  $^3P_2$  and  $^1D_2$  can decay into two gluons, which account for a substantial portion of the hadronic decays for states below  $c\bar{c}$  threshold. The two gluon decay widths are given by [49, 50].

$$\Gamma(n^1S_0 \rightarrow 2g) = \frac{2\alpha_s^2}{3m_c^2} |R_{nS}(0)|^2 \left( 1 + \frac{4.8\alpha_s}{\pi} \right) \quad (16)$$

$$\Gamma(n^3P_0 \rightarrow 2g) = \frac{6\alpha_s^2}{m_c^4} |R'_{nP}(0)|^2 \left( 1 + \frac{9.5\alpha_s}{\pi} \right) \quad (17)$$

$$\Gamma(n^3P_2 \rightarrow 2g) = \frac{8\alpha_s^2}{5m_c^4} |R'_{nP}(0)|^2 \left( 1 - \frac{2.2\alpha_s}{\pi} \right) \quad (18)$$

$$\Gamma(n^1D_2 \rightarrow 2g) = \frac{2\alpha_s^2}{3\pi m_c^6} |R''_{nD}(0)|^2 \quad (19)$$

It is natural that in the non-relativistic potential model of charmonium, the ratio of the two-photon and two-gluon widths of the charmonium decays does not depend on the wave function and slowly grows with increase of the charmonium mass because of the proportionality to  $\frac{1}{\alpha_s^2}$ . According to QCD, the decay of charmonium is due to the annihilation of  $c\bar{c}$  pair. The mass of  $c\bar{c}$  pair is large and the annihilations of  $c\bar{c}$  into gluons are perturbative, so the two-gluon decay mode is dominant in the charmonium. The two gluon decay widths are sensitive to the behavior of the  $q\bar{q}$  wave function and its derivatives near the origin.

### 3. Leptonic Decays

The vector mesons decay leptonically through interaction with the electromagnetic current. The leptonic decay width is proportional to the average value of the squared charge and the squared wave function at the origin. Which gives the probability that the quark and anti-quark will interact with the photon at the origin of their relative coordinates and the mass of the vector mesons. The quark-antiquark assignments for the vector mesons, as well as the fractional values for the quark charges, may be experimented from the values of their leptonic decay widths.

The decay of vector meson into charged leptons proceeds through the virtual photon ( $q\bar{q} \rightarrow l^+l^-$  where  $l = e^-, \mu^-, \tau^-$ ). The  $^3S_1$  and  $^3D_1$  states have quantum numbers of a virtual photon,  $J^{PC} = 1^{--}$  and can annihilate into lepton pairs through one photon.

The leptonic decay width of the vector meson ( $^3S_1$  charmonium) including the first order radiative QCD correction is given by [49, 50]

$$\Gamma(n \ ^3S_1 \rightarrow e^+e^-) = \frac{4\alpha^2 e_c^2 |R_{nS}(0)|^2}{M_{nS}^2} \left(1 - \frac{16\alpha_s}{3\pi}\right) \quad (20)$$

where  $\alpha \approx \frac{1}{137}$  is the fine structure constant,  $M_{nS}$  is the mass of the decaying charmonium state and  $e_c = 2/3$  is the charge of the charm quark in units of the electron charge. For D wave charmonium states the leptonic decay width with leading order QCD correction is given by

$$\Gamma(n \ ^3D_1 \rightarrow e^+e^-) = \frac{25\alpha^2 e_c^2 |R''_{nD}(0)|^2}{2m_c^4 M_{nD}^2} \left(1 - \frac{16\alpha_s}{3\pi}\right) \quad (21)$$

where  $M_{nD}$  is the mass of the decaying charmonium state. The leptonic partial widths are an exploration of the compactness of the quarkonium system and provide important information supplementary to level spacings. The quark-antiquark assignments for the vector mesons, as well as the fractional values for the quark charges, may be experimented from the values of their leptonic decay widths.

### III. RESULTS AND DISCUSSIONS

In our work, we have used the three-dimensional harmonic oscillator wave function which has been extensively used in atomic and nuclear physics is used as the trial wave function for obtaining the  $QQ$  mass spectrum.

$$\psi_{nlm}(r, \theta, \phi) = N \left(\frac{r}{b}\right)^l L_n^{l+1/2}\left(\frac{r}{b}\right) \exp\left(-\frac{r^2}{2b^2}\right) Y_{lm}(\theta, \phi) \quad (22)$$

where  $|N|$  is the normalizing constant given by

$$|N|^2 = \frac{2\alpha^3 n!}{\sqrt{\pi}} \frac{2^{(2(n+l)+1)}}{(2n+2l+1)!} (n+l)! \quad (23)$$

and  $L_n^{l+1/2}(x)$  are the associated Laguerre polynomials. The harmonic oscillator wave function allows the separation of motion of the center of mass and has been used to study the spectra of baryons and mesons [51, 52]. If the basic states are harmonic oscillator wave functions it is rather easy to evaluate the matrix elements of a few body systems such as mesons or baryons. In the harmonic oscillator wave function  $b$  is treated as a variational parameter, which is determined for each state by minimizing the expectation value of the Hamiltonian. The obtained  $b$  value is used in the harmonic oscillator wave function to find the mass spectrum [53]. We have two important parameters characterizing the dilute instanton liquid; the average size of the instanton  $\rho = \frac{1}{3}$  fm, the values of  $\rho$  is less effective in the spin-dependent parts of the potential. The average separation between instantons is  $\bar{R} = 1$  fm [54, 55]. The strength of each part of the potential becomes stronger when smaller value of  $\bar{R}$  is employed. The instanton density is given as  $N/V \simeq (200 \text{ MeV})^4$  and number of colors  $N_C$  is 3. Other parameters in our potential model are, the coupling constant  $\alpha_s$ , the charm quark mass  $m_c$ , the confinement strength  $c$  and a constant  $V_0$ . The confinement strength  $c$  is fixed by the stability condition for variation of mass of the vector meson against the size parameter  $b$ . The mass of the charm quark  $m_c$  and constant  $V_0$  were fixed so as to reproduce the ground state masses. We start with a set of reasonable values of  $m_c$  and  $V_0$ . We used the following set of parameters in our work.

$$\begin{aligned} m_c &= 1475 \text{ MeV}; & \alpha_s &= 0.3 \\ c &= 260 \text{ MeV fm}^{-1}; & V_0 &= -125 \text{ MeV}; \end{aligned}$$

It should be noted that, for harmonic oscillator wave function  $|\Psi(0)|^2 \propto \frac{1}{b^3}$ , which is required to estimate the leptonic and two photon and two gluon decay widths. The  $\alpha_s$  quoted by the latest PDG is 0.1182(12)[56]. The value of  $\alpha_s$  used in the present investigation is 0.3.

Table I lists the low lying charmonium states in comparison with experimental data and other theoretical models. Our predictions for the masses agree with PDG data

within a few MeV. The model correctly reproduces the mass spectrum of charmonium states. The mass of singlet state  $\eta_c(1S)$  is found to be 2984 MeV which is in good

TABLE I. Mass spectrum (MeV).

$n^{2S+1}L_J$	Name	$J^{PC}$	Present Work MeV	$M_{exp}$ MeV	[10] MeV	[57] MeV	[58] MeV	[59] MeV	[60] MeV
$1^1S_0$	$\eta_c(1S)$	$0^{-+}$	2984	$2983.6 \pm 0.7$	2970	2981	2981.7	2990.4	2990
$2^1S_0$	$\eta_c(2S)$	$0^{-+}$	3640	$3639.2 \pm 0.11$	3620	3635	3619.2	3646.5	3643
$3^1S_0$	$\eta_c(3S)$	$0^{-+}$	4061	....	4060	3989	4052.5	4071.9	4054
$1^3S_1$	$J/\psi$	$1^{--}$	3097	$3096.916 \pm 0.011$	3100	3096	3096.92	3085.1	3096
$2^3S_1$	$\psi(2S)$	$1^{--}$	3687	$3686.108 \pm 0.018$	3680	3685	3686.1	3682.1	3703
$3^3S_1$	$\psi(3S)$	$1^{--}$	4039	$4039 \pm 1$	4100	4039	4102	4100.2	4097
$1^1P_1$	$h_c(1P)$	$1^{+-}$	3525	$3525.38 \pm 0.11$	3520	3525	3523.7	3514.6	3515
$2^1P_1$	$h_c(2P)$	$1^{+-}$	3927	....	3960	3926	3963.2	3944.6	3956
$3^1P_1$	$h_c(3P)$	$1^{+-}$	4337	....	....	4337	....	4333.9	4278
$1^3P_0$	$\chi_{c0}(1P)$	$0^{++}$	3414	$3414.75 \pm 0.31$	3440	3413	3415.2	3351.6	3452
$2^3P_0$	$\chi_{c0}(2P)$	$0^{++}$	3916	$3915 \pm 3 \pm 2$	3920	3870	3864.3	3835.7	3909
$3^3P_0$	$\chi_{c0}(3P)$	$0^{++}$	4303	....	....	4301	....	4216.7	4242
$1^3P_1$	$\chi_{c1}(1P)$	$1^{++}$	3510	$3510.66 \pm 0.07$	3510	3511	3510.6	3500.4	3452
$2^3P_1$	$\chi_{c1}(2P)$	$1^{++}$	3872	3872	3950	3906	3950.0	3933.5	3947
$3^3P_1$	$\chi_{c1}(3P)$	$1^{++}$	4312	....	....	4319	....	4317.9	4272
$1^3P_2$	$\chi_{c2}(1P)$	$2^{++}$	3555	$3556.20 \pm 0.09$	3550	3555	3556.2	3551.4	3532
$2^3P_2$	$\chi_{c2}(2P)$	$2^{++}$	3929	$3927.2 \pm 2.6$	3980	3949	3992.3	3979.8	3969
$3^3P_2$	$\chi_{c2}(3P)$	$2^{++}$	4042	....	4010	4041	....	4383.4	4043
$1^1D_2$	$\eta_{c2}(1D)$	$2^{-+}$	3812	....	3840	3807	3822.3	3807.3	3812
$2^1D_2$	$\eta_{c2}(2D)$	$2^{-+}$	4198	....	4210	4196	4196.9	4173.7	4166
$1^3D_1$	$\psi_1(1D)$	$1^{--}$	3779	3778	3820	3783	3789.4	3785.3	3796
$2^3D_1$	$\psi_1(2D)$	$1^{--}$	4192	$4191 \pm 5$	4190	4159	4159.2	4150.4	4153
$1^3D_2$	$\psi_2(1D)$	$2^{--}$	3823	3823	3840	3795	3822.1	3807.7	3810
$2^3D_2$	$\psi_2(2D)$	$2^{--}$	4195	....	4210	4190	4195.8	4173.7	4160
$1^3D_3$	$\psi_2(1D)$	$3^{--}$	3845	....	....	....	3844.8	3814.6	....
$2^3D_3$	$\psi_2(2D)$	$3^{--}$	4220	....	....	....	4218.9	4182.9	....

agreement with the experimentally measured mass value  $2983.6 \pm 0.7 \text{ MeV}$  [3]. Lattice QCD calculations predict a mass of  $2985(1) \text{ MeV}$  [61] for  $\eta_c(1S)$  state. The mass of the spin triplet state  $J/\psi(1S)$  calculated in our model is  $3097 \text{ MeV}$  which is in good agreement with the experimental value  $3096.96 \pm 0.011 \text{ MeV}$  [3] and the values of other theoretical models [10],[57],[58],[59],[60]. The lattice QCD calculations predict a mass of  $3099 \pm 1 \text{ MeV}$  for  $J/\psi(1S)$  state. The masses of radially excited Charmonium state  $\eta_c(2S)$  and its triplet partner  $\psi(2S)$  calculated in our model are in good agreement with both experimental value and with other theoretical models. The lattice QCD calculations predict a slightly less mass value for  $\eta_c(2S)$ , ( $M(\eta_c(2S)) = 3612 \pm 9 \text{ MeV}$ ) and for  $\psi(2S)$  ( $M_{\psi(2S)} = 3653 \pm 12 \text{ MeV}$ ) [61]. It is clear from the table, that the mass of  $\eta_c(3S)$  and its triplet partner calculated in our model is reasonably in good agreement with the experimental and with other model values. The lattice QCD calculations predict a mass of  $4074 \pm 20 \text{ MeV}$  for  $\eta_c(3S)$  and  $4099 \pm 24 \text{ MeV}$  for  $\psi(3S)$  [61].

The mass of spin-singlet P-wave charmonium  $h_c(1P)$  is in good agreement with experiment and other theoretical models. However the lattice QCD predicts a mass of  $3506 \pm 6 \text{ MeV}$  for  $h_c(1P)$  state which is slightly below the experimental value [61]. The spin triplet states  $\chi_{cJ}$  are in good agreement with both experiment and other phenomenological models. The lattice QCD obtains the masses  $20 - 25 \text{ MeV}$  smaller than the experimental val-

ues for these states [61]. We have also predicted masses of low lying  $D$  wave states which is reasonably in good agreement with available experimental data [56] and with other theoretical models [58],[59],[60].

The spin-orbit and tensor potentials in eqn(6) are responsible for the splitting of the charmonium levels. The hyperfine splitting of  $1S$  state obtained in our model  $\Delta M(1^3S_1 - 1^1S_0)$  is  $113 \text{ MeV}$ . The hyperfine mass splitting calculated in our model agrees with the experimental value  $\Delta M(1^3S_1 - 1^1S_0) = 113.2 \pm 0.7 \text{ MeV}$  [3] and lattice QCD results  $\Delta M(1^3S_1 - 1^1S_0) = 114 \pm 1 \text{ MeV}$ . The hyperfine mass splitting of  $2S$  states  $\Delta M(2^3S_1 - 2^1S_0) = 47 \text{ MeV}$  is in good agreement with the  $\Delta M(2^3S_1 - 2^1S_0) = 47 \pm 1 \text{ MeV}$  [3]. The lattice QCD predicts a  $2S$  hyperfine splittings of  $57.9 \pm 2.0 \text{ MeV}$  [62] which is slightly higher than the experimental value. For  $1^3P_1 - 1^3P_0$  splitting, we obtain  $\Delta M(1^3P_1 - 1^3P_0) = 96 \text{ MeV}$  which is good agreement with the experimental value  $\Delta M(1^3P_1 - 1^3P_0) = 95.5 \pm 0.8 \text{ MeV}$  [3]. The Lattice QCD calculations predict a  $1P$  splitting of  $68.4 \pm 5.0 + 11.8 - 3.0 \text{ MeV}$  [63] which is reasonably in good agreement with our calculations. For  $1^3P_2 - 1^3P_1$  splitting, we obtain  $\Delta M(1^3P_2 - 1^3P_1) = 45 \text{ MeV}$ , which is in good agreement with the experimental data  $\Delta M(1^3P_2 - 1^3P_1) = 45.7 \pm 0.2 \text{ MeV}$  [3]. The lattice QCD calculations predict  $\Delta M(1^3P_2 - 1^3P_1) = 31.4 \pm 8.4 \text{ MeV}$  [63], which is rather a low value compared to experimental value.

The ratio between the two hyperfine structures,

TABLE II. Spin averaged mass splittings (MeV)

Mass Splittings	Present Work	Exp	Lattice QCD [63]	GI[66]	[67]	[68]
$M(1^3S_1 - 1^1S_0)$	113	$113.2 \pm 0.7$	$114 \pm 1$	113	114	118
$M(2^3S_1 - 2^1S_0)$	47	$47 \pm 1$	$57.9 \pm 2$	53	44	50
$M(1^3P_2 - 1^3P_1)$	45	$45.7 \pm 0.2$	$31.4 \pm 8.4$	40	36	44
$M(1^3P_1 - 1^3P_0)$	96	$95.5 \pm 0.8$	68.4	65	101	77
$M(1^1P_1 - 1^3P)$	0.6	0.9	-1.4			
$M(1^1P_1 - 1\bar{S})$	456.25	458.5	$448.8 \pm 29$			
$M(1^3P_0 - 1\bar{S})$	345.25	347.1	$419.8 \pm 47$			
$M(1^3P_1 - 1\bar{S})$	441.25	442.9	$448.8 \pm 34$			
$M(1^3P_2 - 1\bar{S})$	486.25	488.6	$448.8 \pm 29$			
$M(1\bar{P} - 1\bar{S})$	455.75	457.9	457.9			
$M(2\bar{P} - 2\bar{S})$	237.91	....	$462 \pm 72$			
$M(2\bar{S} - 1\bar{S})$	606.75	595	$671 \pm 21$			
$M(2\bar{P} - 1\bar{P})$	388.66	....	$675 \pm 76$			

$\frac{\Delta M(1^3P_2 - 1^3P_1)}{\Delta M(1^3P_1 - 1^3P_0)}$  sheds light on the nature of the confinement. Our estimate of the ratio is  $\frac{\Delta M(1^3P_2 - 1^3P_1)}{\Delta M(1^3P_1 - 1^3P_0)} = 0.46$ , and that from the experiment is 0.48[63, 64]. Another interesting quantity is the P-state hyperfine splitting  $\Delta M(1^1P_1 - 1^3P)$ , where,

$$M(1^3P) = \frac{5M(1^3P_2) + 3M(1^3P_1) + M(1^3P_0)}{9} \quad (24)$$

is the center of gravity of the P-wave system. The P-state hyperfine splitting should be much smaller than S-state hyperfine splitting, since the P state wave function is zero at the origin. Our estimate of the P-state hyperfine splitting is  $\Delta M(1^1P_1 - 1^3P) = 0.6$  MeV and the experimental value is  $\Delta M(1^1P_1 - 1^3P) = 0.9$  MeV [3]. The lattice QCD calculations obtain P-state hyperfine splitting of -1.4 MeV.

Radiative decays of excited charmonium states are powerful tool to study the internal structure of the mesons. The possible  $E1$  decay modes have been listed in Table III. Most of the predictions for  $E1$  transitions are in qualitative agreement with other theoretical models. However, there are some differences in the predictions due to differences in phase space arising from different mass predictions and also from the wave function effects. We find our results are compatible with other theoretical model values for most of the channels.

The M1 transition rates of charmonium states have been calculated using eqn(11). Allowed M1 transitions correspond to triplet-singlet transition between S-wave states and between P-state of the same n quantum number, while hindered M1 transitions are either triplet-singlet or singlet-triplet transitions between S-wave states of different quantum numbers. In order to calculate decay rates of hindered transitions, we need to include relativistic corrections, viz, modification of the nonrelativistic wave functions, relativistic modification

The spin averaged masses are defined by,

$$M(n\bar{S}) = \frac{3M(n^3S_1) + M(n^1S_0)}{4} \quad (25)$$

$$M(n\bar{P}) = \frac{3M(n^1P_1) + 5M(n^3P_2) + 3M(n^3P_1) + M(n^3P_0)}{12} \quad (26)$$

with  $n=1,2,3,\dots$  the radial quantum numbers. The spin averaged masses calculated in our model are  $M(1\bar{S})=3068.75$  MeV,  $M(2\bar{S})=3675.25$  MeV,  $M(1\bar{P})=3524.5$  MeV and  $M(2\bar{P})=3913.16$  MeV and the experimental value of spin averaged masses are 2984.3, 3638.56 MeV, 3525.3 MeV and 3929 MeV respectively [63] [65]. The calculated spin averaged mass splittings are listed in Table II.

of the electromagnetic transition operator, and finite-size corrections. In addition to these, there are additional corrections arising from the quark anomalous magnetic moment.

Corrections to the wave function that give contribution to the transition amplitude are of two categories: (1) higher order potential corrections, which are distinguished as (a) the zero recoil effect and (b)recoil effects of the final state meson, and (2) color octet effects. The color octet effects have not been included in potential model formulation and are not considered so far in radiative transitions. The spherical Bessel function  $j_0(k_0r/2)$ , introduced in eqn(11), takes into account the so called finite size effect(equivalently, re-summing the multipole-expanded magnetic amplitude to all orders). For small value of  $k_0$ ,  $j_0(k_0r/2) \rightarrow 1$ , the transitions with  $n' = n$  have dominant contribution to the matrix elements, though the corresponding partial decay widths are suppressed by smaller  $k_0^3$  factors. For a large value of photon energy (k), transitions with  $n \neq n'$  have domi-

TABLE III. E1 Transition rates

E1 Transition $i \rightarrow f$	k (MeV)	Present Work $\Gamma$ ( $i \rightarrow f$ )(keV)	$\Gamma_{Expt}$ (keV)	[66] <sup>a</sup> (keV)	[66] <sup>b</sup> (keV)	[44] (keV)	[59] (keV)	[58] (keV)
$1^3P_0 \rightarrow 1^3S_1 + \gamma$	317	114.74	119.5 $\pm$ 8	152	114	120	97	139.3
$1^3P_1 \rightarrow 1^3S_1 + \gamma$	413	297.61	295.8 $\pm$ 13	314	239	241	330	38.4
$1^3P_1 \rightarrow 1^1S_0 + \gamma$	526	486.26		498	352	482	465	546.4
$1^3P_2 \rightarrow 1^3S_1 + \gamma$	458	350.58	384.2 $\pm$ 16	424	313	315	421	319.4
$2^3S_1 \rightarrow 1^3P_0 + \gamma$	273	55.61		63	26	47	34	25.2
$2^1S_0 \rightarrow 1^1P_1 + \gamma$	115	40.26		49	36	35.1	72	17.4
$2^3S_1 \rightarrow 1^3P_1 + \gamma$	177	25.45	28.0 $\pm$ 1.2	54	29	42.8	48	29.1
$2^3S_1 \rightarrow 1^3P_2 + \gamma$	132	23.9	26.6 $\pm$ 1.1	38	24	30.1	43	26.5
$1^3D_1 \rightarrow 1^3P_0 + \gamma$	365	200.66	199.3 $\pm$ 25	403	213	299	367	243.9
$1^3D_1 \rightarrow 1^3P_1 + \gamma$	269	75.34	79.2 $\pm$ 16	125	77	99	146	104.9
$1^3D_1 \rightarrow 1^3P_2 + \gamma$	224	3.70	3.88	< 24.6	4.9	3.3	6.8	1.9
$1^3D_2 \rightarrow 1^3P_1 + \gamma$	313	245.35		307	268	313	321	256.7
$1^3D_2 \rightarrow 1^3P_2 + \gamma$	268	80.64		64	66	69.5	79	61.8
$1^1D_2 \rightarrow 1^1P_1 + \gamma$	287	310.52		339	344	389	398	
$1^3D_3 \rightarrow 1^3P_2 + \gamma$	290	246.79		272	296	402	340	

<sup>a</sup>Non relativistic quark model<sup>b</sup>Relativistic quark model

TABLE IV. M1 Transition rates

( $M_1$ )Transition $i \rightarrow f$	k (MeV)	Present Work $\Gamma$ ( $i \rightarrow f$ )(keV)	$\Gamma_{Expt}$ (keV)	[66] <sup>c</sup> (keV)	[66] <sup>d</sup> (keV)	[59] <sup>c</sup> (keV)	[59] <sup>d</sup> (keV)
$1^3S_1 \rightarrow 1^1S_0 + \gamma$	113	2.38	1.58 $\pm$ 0.37	2.9	2.4	1.5	2.2
$2^3S_1 \rightarrow 2^1S_0 + \gamma$	47	0.17	0.143 $\pm$ 0.027	0.21	0.17	3.1	3.8
$2^3S_1 \rightarrow 1^1S_0 + \gamma$	703	3.96	0.97 $\pm$ 0.027	4.6	9.6	6.1	6.9
$2^3S_1 \rightarrow 1^3S_1 + \gamma$	590	5.30		7.9	5.6	0.70	0.71
$3^3S_1 \rightarrow 2^1S_0 + \gamma$	399	1.12		0.61	2.6	3.2	3.7
$3^1S_0 \rightarrow 2^3S_1 + \gamma$	374	1.42		1.3	0.84	1.7	1.6
$3^1S_0 \rightarrow 1^3S_1 + \gamma$	964	6.51		6.3	6.9	5.9	6.5
$2^1P_1 \rightarrow 1^3P_2 + \gamma$	372	0.23		0.071	0.11	1.3	1.2
$2^1P_1 \rightarrow 1^3P_1 + \gamma$	417	0.18		0.058	0.36	0.16	0.13
$2^1P_1 \rightarrow 1^3P_0 + \gamma$	513	1.31		0.033	1.5	5.6	5.3
$2^3P_2 \rightarrow 1^1P_1 + \gamma$	404	1.12		0.67	1.3	1.0	0.89
$2^3P_1 \rightarrow 1^1P_1 + \gamma$	347	0.15		0.050	0.045	0.15	0.13

<sup>c</sup>Non relativistic quark model<sup>d</sup>Relativistic quark model

nant contribution to the matrix element, since  $j_0(k_0 r/2)$  becomes very small. M1 transition rates are very sensitive to hyperfine splittings of the levels due to the  $k_0^3$  factor in eqn(11). The resulting M1 radiative transition rates of these states are presented in Table III. The M1 transition rates calculated in our model agree well with the values predicted by other theoretical models.

Using the Van-Royen-Weisskopf relation we have calculated annihilation decay widths like leptonic decay

widths, two -photon and two gluon decay widths with the inclusion of radiative corrections [50]. The resulting leptonic decay widths are listed in Table V. Our predictions for leptonic decay widths have been compared with experiment and other theoretical models and are found to be in good agreement. The Tables VI and VII list the two photon and two gluon decay widths of charmonium states. The two photon and two-gluon decay widths for charmonium states reasonably agree with the available experimental data and with other theoretical models.

TABLE V. Leptonic Decay widths (keV)

State	Present Work $\Gamma_{l+l-}$	Exp $\Gamma_{l+l-}$	[39]	[58]	[69]
$J/\psi$	4.65	$5.55 \pm 0.14$	3.589	4.28	12.13
$\psi(2S)$	2.25	$2.33 \pm 0.07$	1.440	2.25	5.03
$\psi(3S)$	0.98	$0.86 \pm 0.07$	0.975	1.66	3.48
$1^3D_1$	0.35	$0.242 \pm 0.030$	0.096	0.09	0.056
$2^3D_1$	0.68	$0.83 \pm 0.07$	0.112	0.16	0.096

TABLE VI. Two-Photon Decay widths (keV)

State	Present Work $\Gamma$	Exp $\Gamma$	[39]	[70]	[71]
$\eta_c(1S)$	6.89	$7.2 \pm 0.7$	6.812	3.50	7.18
$\eta_c(2S)$	6.68	$7.0 \pm 3.5$	2.625	1.38	1.71
$\eta_c(3S)$	1.23		1.760	0.94	1.21
$\chi_{c0}(1P)$	1.96	$2.36 \pm 0.35$	2.119	1.39	3.28
$\chi_{c0}(2P)$	1.02		1.308	1.11	
$\chi_{c2}(1P)$	0.82	$0.66 \pm 0.07$	0.261	0.44	
$\chi_{c2}(2P)$	0.19		0.168	0.48	

TABLE VII. Two-Gluon Decay widths (MeV)

State	Present Work $\Gamma$	Exp	[39]	[72]	[73]
$\eta_c(1S)$	27.61	$28.6 \pm 2.2$	22.048	15.70	32.209
$\eta_c(2S)$	7.92	$14 \pm 7$	8.496	8.10	
$\eta_c(3S)$	5.67		5.696		
$\chi_{c0}(1P)$	9.67	$10.3 \pm 0.6$	6.114	4.68	10.467
$\chi_{c0}(2P)$	3.67		3.775		
$\chi_{c2}(1P)$	2.15	$1.97 \pm 0.11$	0.633	1.72	1.169
$\chi_{c2}(2P)$	0.59		0.401		
$1^1D_2^e$	0.068		0.014		
$2^1D_2^e$	0.061		0.012		

<sup>e</sup>Without QCD corrections

#### IV. CONCLUSIONS AND OUTLOOK

Basic aim of the present work is to develop a consistent model which could reproduce both the spectra and the decay widths with the same set of parameters and to investigate the effect of instanton potential on masses and the excited states of charmonium. From our analysis, we infer that the present model has the right prediction both for the mass spectrum and decay widths. Since different potentials can reproduce the same spectra the stringent test for any given model is the calculation of the other observables like leptonic, the radiative, two-photon and two-gluon decay widths in addition to mass spectrum. In our earlier work also, we had come to the similar conclusion while investigating light meson spectrum [74] [75]. The differences in the prediction for the decay rates in various theoretical models can be attributed to the differences in mass predictions and wave function effects. From the study of mass spectra and decay properties of  $c\bar{c}$  states in a phenomenological approach we draw up the following conclusions.

(1) Our calculations for the low lying charmonium

states are in good agreement with experimental measurements

- (2) The hyperfine separations are directly related to the spin-spin interaction. The theoretical predictions of our model are remarkably consistent with well established experimental data for the conventional charmonium states. Below the open charm threshold, our theoretical calculations are in agreement with lattice calculations and experimental results.
- (3) The mass splittings between the radial excitations and the ground states also provide an important check on the validity of our potential model.
- (4) The calculations demonstrate that the charmonium masses/mass splittings can be computed with a combination of coulombic, instanton and confinement potentials. Hence, in heavy quark sector, the instanton potential plays the role of OGEP. The instanton effects are quite significant in the central part of the potential.



- (5) They turn out to be rather small in describing the hyperfine mass splittings of the charmonia. This might be due to the spin-dependent part of the potential from the instanton vacuum is an order of magnitude smaller than the central part. The tensor interaction almost does not contribute to the masses.
- (6) Charmonium decays provide a deeper insight on the exact nature of the inter quark forces and decay mechanisms. Using the predicted masses

and the radial wave function at the origin, leptonic decays, two photon and two gluon decays are computed using the Van Royen-Weisskopf relation. The calculated values including the correction factor agree with the experimental values within a few MeV. From our calculations, we conclude that the inclusion of QCD correction factors are of importance for obtaining accurate results for decay rates.

- 
- [1] J. J. Aubert *et al.*, Phys. Rev. Lett. **33**, 1404 (1974).
  - [2] J. E. Augustin *et al.*, Phys. Rev. Lett. **33**, 1406 (1974).
  - [3] K. Olive *et al.*, Chinese Physics C **38**, 090001 (2014).
  - [4] W.-J. Deng, H. Liu, L.-C. Gui, and X.-H. Zhong, Phys. Rev. D **95**, 034026 (2017).
  - [5] C. T. H. Davies, K. Hornbostel, G. P. Lepage, A. J. Lidsey, J. Shigemitsu, and J. Sloan, Phys. Rev. D **52**, 6519 (1995).
  - [6] J. J. Dudek and E. Rrapaj (for the Hadron Spectrum Collaboration), Phys. Rev. D **78**, 094504 (2008).
  - [7] E. Eichten, K. Gottfried, T. Kinoshita, K. Lane, and T.-M. Yan, Physical Review Letters **36**, 500 (1976).
  - [8] E. Eichten, K. Gottfried, T. Kinoshita, K. Lane, and T.-M. Yan, Physical Review D **17**, 3090 (1978).
  - [9] E. Eichten, K. Gottfried, T. Kinoshita, K. Lane, and T.-M. Yan, Physical Review D **21**, 203 (1980).
  - [10] S. Godfrey and N. Isgur, Phys. Rev. D **32**, 189 (1985).
  - [11] D. Stanley and D. Robson, Physical Review D **21**, 3180 (1980).
  - [12] D. Ebert, R. Faustov, and V. Galkin, Physical Review D **67**, 014027 (2003).
  - [13] S. Gershtein, V. Kiselev, A. Likhoded, and A. Tkabladze, Physical Review D **51**, 3613 (1995).
  - [14] L. P. Fulcher, Physical Review D **44**, 2079 (1991).
  - [15] L. P. Fulcher, Physical Review D **60**, 074006 (1999).
  - [16] L. P. Fulcher, Physical Review D **50**, 447 (1994).
  - [17] S. N. Gupta, J. M. Johnson, W. W. Repko, and C. J. Suchyta III, Physical Review D **49**, 1551 (1994).
  - [18] S. N. Gupta and J. M. Johnson, Physical Review D **53**, 312 (1996).
  - [19] C. S. Vanamali and K. B. V. Kumar, Phys. Rev. C **94**, 054002 (2016).
  - [20] R. Bhaduri, L. Cohler, and Y. Nogami, Il Nuovo Cimento A (1965-1970) **65**, 376 (1981).
  - [21] L. Burakovsky and T. Goldman, Nucl. Phys. A **625**, 220 (1997).
  - [22] S. Gershtein *et al.*, Sov. J. Nucl. Phys **48**, 327 (1988).
  - [23] Y.-Q. Chen and Y.-P. Kuang, Phys. Rev. D **46**, 1165 (1992).
  - [24] S. S. Gershtein, V. V. Kiselev, A. K. Likhoded, and A. V. Tkabladze, Phys. Rev. D **51**, 3613 (1995).
  - [25] M. Baldicchi and G. M. Prosperi, Phys. Rev. D **62**, 114024 (2000).
  - [26] W. Kwong and J. L. Rosner, Phys. Rev. D **44**, 212 (1991).
  - [27] K. B. Vijaya Kumar, Bhaghyesh, and A. P. Monteiro, Indian Journal of Physics **87**, 493 (2013).
  - [28] A. P. Monteiro and K. B. Vijaya Kumar, Chinese physics C **34**, 1396 (2010).
  - [29] A. P. Monteiro and K. B. Vijaya Kumar, Communications in theoretical Physics **53**, 325 (2010).
  - [30] A. P. Monteiro and K. B. Vijaya Kumar, Indian Journal of Pure & Applied Physics **48**, 240 (2010).
  - [31] A. P. Monteiro, M. Bhat, and K. B. Vijaya Kumar, International Journal of Modern Physics A **32**, 1750021 (2017).
  - [32] M. Bhat, A. P. Monteiro, and K. B. Vijaya Kumar, International Journal of Modern Physics E **26**, 1750037 (2017).
  - [33] K. B. Vijaya Kumar, A. K. Rath, and S. B. Khadkikar, Pramana **48**, 997 (1997).
  - [34] A. P. Monteiro, K. B. Vijaya Kumar, and Bhaghyesh, Communications in Theoretical Physics **56**, 476 (2011).
  - [35] Bhaghyesh, K. B. Vijaya Kumar, and Y.-L. Ma, International Journal of Modern Physics A **27**, 1250011 (2012).
  - [36] Bhaghyesh, K. B. Vijaya Kumar, and A. P. Monteiro, Chinese Physics C **35**, 997 (2011).
  - [37] Bhaghyesh and K. B. Vijaya Kumar, Chinese Physics C **37**, 023103 (2013).
  - [38] Bhaghyesh and K. B. Vijaya Kumar, Communications in Theoretical Physics **55**, 1044 (2011).
  - [39] Bhaghyesh and K. B. Vijaya Kumar, International Journal of Modern Physics A **27**, 1250127 (2012).
  - [40] U. T. Yakhshiev, H.-C. Kim, M. M. Musakhanov, E. Hiyama, and B. Turimov, Chin. Phys. C **41**, 083102 (2017).
  - [41] W.-J. Deng, H. Liu, L.-C. Gui, and X.-H. Zhong, arXiv preprint arXiv:1608.00287 (2016).
  - [42] W. Kwong and J. L. Rosner, Phys. Rev. D **38**, 279 (1988).
  - [43] V. Novikov, L. Okun, M. Shifman, A. Vainshtein, M. Voloshin, and V. Zakharov, Phys. Rep. **41**, 1 (1978).
  - [44] N. Brambilla *et al.*, CERN Yellow Report, CERN-2005-005 (2004).
  - [45] W. Caswell and G. Lepage, Phys. Lett. B **167**, 437 (1986).
  - [46] G. T. Bodwin, E. Braaten, and G. P. Lepage, Phys. Rev. D **51**, 1125 (1995).
  - [47] N. Brambilla, A. Pineda, J. Soto, and A. Vairo, Phys. Lett. B **470**, 215 (1999).
  - [48] B. A. Thacker and G. P. Lepage, Phys. Rev. D **43**, 196 (1991).
  - [49] W. Kwong, P. B. Mackenzie, R. Rosenfeld, and J. L. Rosner, Phys. Rev. D **37**, 3210 (1988).
  - [50] R. Royen and V. F. Weisskopf, Il Nuovo Cimento A **50**, 617 (1967).

- [51] D. Faiman and A. W. Hendry, Phys. Rev. **180**, 1572 (1969).
- [52] R. P. Feynman, M. Kislinger, and F. Ravndal, Phys. Rev. D **3**, 2706 (1971).
- [53] Bhaghyesh, K. B. Vijaya Kumar, and A. P. Monteiro, Journal of Physics G: Nuclear and Particle Physics **38**, 085001 (2011).
- [54] D. I. Dyakonov and V. Y. Petrov, Nuclear Physics B **245**, 259 (1984).
- [55] E. V. Shuryak, Nuclear Physics B **203**, 93 (1982).
- [56] C. P. et al., Chinese Physics C **40**, 100001 (2016).
- [57] D. Ebert, R. N. Faustov, and V. O. Galkin, The European Physical Journal C-Particles and Fields **71**, 1 (2011).
- [58] S. F. Radford and W. W. Repko, Physical Review D **75**, 074031 (2007).
- [59] L. Cao, Y.-C. Yang, and H. Chen, Few-Body Systems **53**, 327 (2012).
- [60] J. Segovia, D. R. Entem, F. Fernandez, and E. Hernandez, International Journal of Modern Physics E **22**, 1330026 (2015).
- [61] T. Kawanai and S. Sasaki, Phys. Rev. D **92**, 094503 (2015).
- [62] D. Mohler, Proceedings, 5th International Workshop on Charm Physics (Charm 2012): Honolulu, Hawaii, USA, May 14-17, 2012 (2012).
- [63] M. Okamoto *et al.* (CP-PACS Collaboration), Phys. Rev. D **65**, 094508 (2002).
- [64] W. Lucha and F. F. Schöberl, arXiv preprint hep-ph/9601263 (1996).
- [65] J. Zhang, arXiv:1311.370v (2013).
- [66] T. Barnes, S. Godfrey, and E. S. Swanson, Phys. Rev. D **72**, 054026 (2005).
- [67] W.-J. Deng, H. Liu, L.-C. Gui, and X.-H. Zhong, arXiv preprint arXiv:1608.00287 (2016).
- [68] B.-Q. Li and K.-T. Chao, Phys. Rev. D **79**, 094004 (2009).
- [69] T. Barnes, arXiv preprint hep-ph/0406327 (2004).
- [70] C. R. Münz, Nuclear Physics A **609**, 364 (1996).
- [71] O. Lakhina and E. S. Swanson, Physical Review D **74**, 014012 (2006).
- [72] A. Parmar, B. Patel, and P. C. Vinodkumar, Nuclear Physics A **848**, 299 (2010).
- [73] J. T. Lavery, S. F. Radford, and W. W. Repko, arXiv preprint arXiv:0901.3917 (2009).
- [74] K. B. Vijayakumar, B. Hanumaiah, and Pepin, The European Physical Journal A **19**, 247 (2004).
- [75] Bhavyashri, K. B. Vijayakumar, B. Hanumaiah, S. Sarangi, and S.-G. Zhou, Journal of Physics G: Nuclear and Particle Physics **31**, 981 (2005).