SPECIAL VALUES OF ZETA-FUNCTIONS OF REGULAR SCHEMES

STEPHEN LICHTENBAUM

§0 Introduction.

In this paper we will give a conjectural formula (Conjecture 3.1) for the special values $\zeta^*(X,r)$ of the scheme zeta-function of a regular scheme X projective and flat of dimension d (so relative dimension d-1) over Spec $\mathbb Z$ at a rational integer r, in terms of singular, de Rham, and Weil-ètale motivic cohomology, valid up to sign and powers of 2.

We can factor the map from X to Spec \mathbb{Z} uniquely through Spec O_K , where K is a number field and the generic fiber of X over Spec O_K is a smooth connected algebraic variety over K. We will construct complexes made up of variants of these cohomology groups, and the conjectured formula will give the special value as a product of Euler characteristics of these complexes, equipped with suitable integral structures. We will prove that, if $d \leq 2$ this conjecture is compatible with Serre's conjectured functional equation for the zeta-function, and if d > 2 this compatibility is true modulo two previously existing conjectures which are true in dimension ≤ 2 .

We will discuss how this related to previous work on this subject. Beilinson, building on a previous conjecture of Deligne, gave a special values conjecture ([RSS]) for $r \leq 0$ up to a rational number. More specifically, if one writes the scheme zeta function in the usual way as the product $\prod_{j=0}^{2d-2} L_j(X,s)^{(-1)^{j+1}}$, Beilinson gave a formula for the special value $L_j(X,r)$ up to a rational number. Bloch and Kato ([BK]) gave a formula up to sign for this special value when the weight $j-2r \leq -3$. Fontaine and Perrin-Riou ([Fo]) gave such a conjectured formula for all integers r. Flach and Morin ([FM1]) gave a conjectured formula for special values of the zeta-function taking into account real primes. and more conceptual than that of Fontaine and Perrin-Riou. In a very recent preprint [FM2], they also giver a proof of the compatibility of their conjecture with the functional equation, under essentially the same hypotheses as in the present paper. Fontaine asserts that the compatibility of his conjecture would follow from a local conjecture which still remains far from being proven.

In this paper, as in ([FM1]), we only consider the zeta-function and not the associated L-functions. We believe that the L-function conjectures are probably not completely correct, basically because of torsion phenomena in cohomology, which necessitate correction terms in analogous formulas for special values of zeta-functions of varieties over finite fields. We note that the wonderful formula of Bloch, Kato, and T. Saito, which plays an extremely important role in the proof of compatibility is only valid for Euler characteristics, and not for individual cohomology groups, which forces the restriction to zeta-functions. We also remark that throughout this paper we work with the scheme zeta-function of X.

Recently Niranjan Ramachandran and I [(LR]) have shown that if X is an arithmetic surface and r = 1, rhe main conjecture of the paper is equivalent to the conjecture of Birch and Swinnerton-Dyer for the Jacobian of X. A similar result has been proved by Flach and Siebert ([FS]) for the main conjecture of [FM1].

The equivalence of the conjecture in this article with the conjecture of Flach-Morin, and of either or both of these conjectures with the original conjecture of Fontaine and Perrin-Riou remains open, and is a very interesting and important question.

There are two basic approaches to zeta-function conjectures: One (the Tamagawa approach) involves writing the formula as a product of local formulas (one for each prime p), and then using the product formula to show that although the individual factors may depend on choices, (possibly of a differential) the product does not. This approach is used by Tate in his Bourbaki talk [T] on the conjecture of Birch and Swinnerton-Dyer for abelian varieties, by Fontaine and Perrin-Riou([Fo]), and by Flach-Morin. ([FM1]).

The other just involves just working with the infinite primes and for example, choosing a particularly good differential. This approach was used by the author in his conjectures on the Dedekind zeta function ([Li3]), by Silverman in stating the conjecture of Birch and Swinnerton-Dyer for elliptic curves in his book ([Si]) on elliptic curves, and in this paper. The basic idea of this paper is that by making the infinite prime part of the formula of Fontaine and Perrin -Riou more precise, we can dispense with the detailed local analysis.

We should make it clear that, even to state our conjecture, we have to assume the validity of other previous conjectures.

First, we need the conjecture, which is very far from being proved, that the zeta-function of X, which converges for Re(s) > d, can be meromorphically continued to the entire plane, so we can talk about $\zeta^*(X, r)$ for r < d.

Second, we assume that the etale motivic cohomology groups that we will define are finitely generated.

Third, we assume that the Beilinson regulator maps and various Arakelov intersection pairings induce isomorphisms on complex vector spaces.

For the proof of compatibility, we need the theorem that the groups $H_{et}^{2r+1}(X,\mathbb{Z}(r))$ and $H_{et}^{(2(d-r)+1)}(X,\mathbb{Z}(d-r))$ are finite and Pontriagin dual to each other. This has been proved by Flach and Morin [FM1] if $d \leq 2$, and under some restrictions in the general case..

We also need, for the full Bloch-Kato-Saito theorem, resolution of singularities for arithmetic schemes.

Let X be a scheme of finite type over Spec \mathbb{Z} , and Krull dimension d. Let x denote a closed point of X, and let N(x) be the order of the residue field $\kappa(x)$. Recall that x is closed if and only if $\kappa(x)$ is finite. Let G be any meromorphic function on \mathbb{C} , let r be a rational integer, and let a_r be the order of the zero of G(s) at s = r. Let $G^*(r)$ be the limit as s approaches r of $G(s)(s-r)^{-a_r}$. If G is a zeta-function, $G^*(r)$ is referred to as a special value of G.

Definition 0.1. [Se1]: The zeta function $\zeta(X,s)$ of X is defined to be $\prod_x (1-N(x)^{-s})^{-1}$, where x runs over the closed points of X.

The product defining $\zeta(X,s)$ is well known to converge for Re(s) > d and is conjectured to have a meromorphic continuation to the entire plane. We will tacitly assume this conjecture in what follows. It is further conjectured ([Se2]. [Bl3]) that there exists a Γ -factor $\Gamma(X,s)$ and a positive rational number A, the conductor, such that if we let $\phi(X,s) = A^{s/2}\zeta(X,s)\Gamma(X,s)$ then $\phi(X,s)$ satisfies the functional equation $\phi(X,s) = \pm \phi(X,d-s)$.

Now let X be regular, and projective and flat over Spec \mathbb{Z} . The basic idea behind our conjectured formula is to start with Fontaine's "Deligne-Beilinson" conjectures [Fo], which give the special values up to a rational number in terms of determinants of maps of complex vector spaces with given rational structures. These complex vector spaces come from singular and de Rham cohomology, and from Weil-étale motivic cohomology. We replace the rational structures by integral structures, and take determinants with respect to these. The singular cohomology of course has a natural integral structure, and the Weil-étale groups conjecturally do also. We define an integral structure on the de Rham groups by using derived exterior powers. We should note that these derived exterior powers have

an important role to play even in the number ring case (d = 1).

We also introduce the orders of naturally occurring finite cohomology groups into the picture. Finally we replace the period maps in Fontaine's picture by "modified" period maps, where we divide by special values of the Gamma function.

Our conjectural formula expresses the special values of the zeta-function in terms of the product of Euler characteristics of exact sequences of complex vector spaces with integral structures. The complex vector spaces will be derived from singular cohomology, de Rham cohomology, and Weil-ètale motivic cohomology. For the exact formula, see Section 3. The maps between them arise from Beilinson's conjectures, Arakelov height pairings, and periods.

The plan of this paper is as follows:

Section 1: Integral structures and Euler characteristics.

Section 2: The groups and maps involved in the conjecture,

Section 3: Statement of the conjecture.

Section 4: The Euler characteristic of the period map.

Section 5: Serre's functional equation and Γ - function identities.

Section 6: Compatibility of the conjecture with the functional equation.

Section 7. The case of number rings.

As we move along we will explain how our definitions of groups and maps relate to those of Fontaine and Perrin-Riou [FP].

We would like to thank Spencer Bloch, Matthias Flach, Thomas Geisser, Baptiste Morin, Niranjan Ramachandran, and Takeshi Saito for many helpful conversations.

§.1. Integral structures and Euler characteristics.

Let $V_0, V_1 \dots V_n$ be finite-dimensional complex vector spaces, and

$$V_* = 0 \rightarrow V_0 \rightarrow V_1 \rightarrow \cdots \rightarrow V_n \rightarrow 0$$

be an exact sequence.

Let B_i be a lattice spanned by a basis for the vector space V_i . Let ΛV denote the highest exterior power of V and ΛB denote the highest exterior power of B. The alternating tensor product of the $\Lambda V_i's$ is canonically isomorphic to $\mathbb C$ and the alternating tensor product of the $\Lambda B_i's$ is isomorphic to $\mathbb Z$. The natural inclusions of B_i in V_i induce a map from $\mathbb Z$ to $\mathbb C$

and the determinant $det(V_*, B_*)$ in $\mathbb{C}^*/\pm 1$ of the pair (V_*, B_*) is defined to be the image of a generator of \mathbb{Z} in \mathbb{C} .

Definition 1.1. Let V_* be a sequence of finite-dimensional complex vector spaces. An integral structure on V_* is a sequence of pairs (A_*, a_*) where A_* is a lattice in V_* and a_* is a positive rational number.

An example of an integral structure on a finite-dimensional complex vector space V comes from a finitely generated abelian group M with a homomorphism from M to V whose image is a lattice M_0 in V and whose kernel is the torsion subgroup M_{tor} of M. The integral structure is then $(M_0, |M_{tor}|)$.

Definition 1.2. An integral structure (V_*, A_*) is torsion-free if each a_j is equal to 1.

Definition 1.3. Let (A_*, a_*) be an integral structure on the finite exact complex V_* . We define the Euler characteristic of (A_*, a_*) to be det $(V_*, A_*) \prod a_j^{(-1)^{j+1}}$

Definition 1.4. Let (A_*, a_*) be an integral structure on V_* , (B_*, b_*) be an integral structure on W_* , and ϕ_* be a map of complexes from V_* to W_* such that ϕ_j is an isomorphism for all j. Let \det_j be the determinant of ϕ_j with respect to the lattices A_j and B_j , and let $\chi(\phi_j) = \det_j b_j/a_j$. Define the Euler characteristic $\chi(\phi)$ to be $\prod_{j=0}^n (-1)^{j+1} \chi(\phi_j)$.

Definition 1.5. Let $\tilde{A} = (A_*, a_*)$ be an integral structure on V_* . Let V_*^{\vee} be the sequence of dual vector spaces to V_* . We define the dual integral structure \tilde{A}^{\vee} on V_*^{\vee} to be (A_*^{\vee}, a_*^{-1}) , where A_*^{\vee} is the lattice dual to A_* .:

Proposition 1.6. The Euler characteristic of the dual integral structure is equal to either to the Euler characteristic of the original integral structure or to its inverse, depending on whether n is odd or even.

Proof. Well-known

- §2. The groups and maps involved in the conjecture.
- §2.1 Weil-étale motivic cohomology...

As before, Let X be a regular scheme, projective and flat over Spec \mathbb{Z} Let X_0 be the fiber of X over Spec \mathbb{Q} , and let K be the algebraic closure of \mathbb{Q} in the function field of X, Let O_K be the ring of integers in K. We may regard X as a scheme projective and flat over Spec O_K .

We will first define Weil-ètale motivic cohomology groups and then discuss their relation to the groups defined by Fontaine and Perrin-Riou ([Fo] and [Fl]).

Let r be an integer, and j a non-negative integer. We would like to define a Weil-étale site and complexes of sheaves $\mathbb{Z}(r)$ on this site whose cohomology groups $H_W^j(X,\mathbb{Z}(r))$ would be Weil-ètale motivic cohomology, but unfortunately we do not know how to do this. Instead, for $j \leq 2r$ we define $H_W^j(X,\mathbb{Z}(r))$ to be the hypercohomology groups $H_{et}^j(X,\mathbb{Z}(r))$, where $\mathbb{Z}(r)$ denotes Bloch's higher Chow group complex sheafified for the étale topology ([Bl1].[Le]). Sometimes these groups are referred to as étale motivic cohomology. For $j \geq 2r + 1$, we define $H_W^j(X,\mathbb{Z}(r))$ to be $h^j(RHom(R\Gamma_{et}(X,\mathbb{Z}(d-r)),\mathbb{Z}[-2d-1])$, so we have the exact sequence

$$(2.1.1)$$

$$0 \to Ext^{1}(H_{et}^{2d+2-j}(X, \mathbb{Z}(d-r)), \mathbb{Z}) \to H_{W}^{j}(X, \mathbb{Z}(r)) \to Hom(H_{et}^{2d+1-j}(X, \mathbb{Z}(d-r)), \mathbb{Z}) \to 0$$

If we had our hypothetical Weil-étale site, with section functor denoted by Γ_W , this would follow, up to 2-torsion, from a duality theorem which asserted that $R\Gamma_W(X,\mathbb{Z}(d-r))$ was isomorphic to $RHom(R\Gamma_W(X,\mathbb{Z}(r)),\mathbb{Z}[-2d-1])$. The analogue of this theorem, assuming the usual conjectures, is true for Weil-étale cohomology in the geometric case, as shown in [Ge]). We note here that in [FM1], Flach and Morin have constructed such a complex of abelian groups, which satisfies this duality theorem assuming that standard finiteness conjectures hold.

The group $H^{2r}_W(X,\mathbb{Z}(r))$ is by definition $H^{2r}_{et}(X,\mathbb{Z}(r))$, and by standard arguments this agrees with the group $H^{2r}_{Zar}(X,\mathbb{Z}(r))$ of codimension r cycles on X modulo rational equivalence after tensoring with \mathbb{Q} . Hence there is a cycle map ϕ from $H^{2r}_W(X,\mathbb{Z}(r))$ to singular cohomology with rational coefficients. Let $H^{2r}_W(X,\mathbb{Z}(r))^1$ denote Ker ϕ (cycles homologous to zero) and $H^{2r}_W(X,\mathbb{Z}(r))^2$ denote Image ϕ (cycles modulo homological equivalence.).

We have the exact sequence

$$0 \to H^{2(d-r)}_W(X, \mathbb{Z}(d-r))^1 \to H^{2(d-r)}_W(X, \mathbb{Z}(d-r)) \to H^{2(d-r)}_W(X, \mathbb{Z}(d-r))^2 \to 0$$

Conjecture 2.1.1. The groups $H^j_{et}(X,\mathbb{Z}(r))$ are finitely generated for $j \leq 2r+1$, and finite for j = 2r+1.

This implies

Conjecture 2.1.2. The cohomology groups $H_W^j(X,\mathbb{Z}(r))$ are finitely generated for all j.

Assuming the truth of this conjecture, we give the complex vector space $H_W^j(X,\mathbb{Z}(r))_{\mathbb{C}}$ the standard integral structure $H_W^j(X,\mathbb{Z}(r))$.

We also need

Conjecture 2.1.3. The finite groups $H_{et}^{2r+1}(X,\mathbb{Z}(r))$ and $H_{et}^{2(d-r)+1}(X,\mathbb{Z}(d-r))$ are Pontriagin duals.

Flach and Morin show in [FM1, Proposition 3.4] that this follows from Conjecture 2.1.2 for $d \leq 2$ and is under some restrictions in the general case.

§2.2 Singular and de Rham cohomology.

We also will have need of singular cohomology groups. Let $X_{\mathbb{C}} = X \times_{\mathbb{Z}} \mathbb{C}$. Complex conjugation c acts on $H_B^j(X_{\mathbb{C}}, \mathbb{Z})$ via the natural action of conjugation on \mathbb{C} . If r is even (resp. odd), let $\tilde{H}_B^j(X, \mathbb{C}(r))^+$ and $\tilde{H}_B^j(X, \mathbb{Z}(r))^+$ be the set of elements y in $H_B^j(X_{\mathbb{C}}, \mathbb{C})$ and $H_B^j(X_{\mathbb{C}}, \mathbb{Z})$ such that c(y) = y (resp. c(y) = -y). We define $H_B^j(X, \mathbb{C}(r))$ (resp. $H_B^j(X, \mathbb{C}(r))^+$) to be $H_B^j(X, \mathbb{C}(r))$ (resp. $\tilde{H}_B(X_{\mathbb{C}}, \mathbb{C})^+$) and its standard integral structure given by mapping $H_B^j(X_{\mathbb{C}}, \mathbb{Z})$ (resp. $\tilde{H}_B^j(X_{\mathbb{C}}, \mathbb{Z})^+$) to $H_B^j(X_{\mathbb{C}}, \mathbb{C})$ (resp. $\tilde{H}_B(X_{\mathbb{C}}, \mathbb{C})^+$) via the natural map followed by multiplication by $(2\pi i)^r$.

Let $\Omega = \Omega_{X_{\mathbb{C}}/\mathbb{C}}$. The de Rham cohomology group $H^j_{DR}(X_{\mathbb{C}},\mathbb{C})$ has the Hodge decomposition

$$\prod_{i+k=j} H^i(X_{\mathbb{C}}, \Lambda^k \Omega)$$

which gives rise to the Hodge filtration $G_m = \prod_{k \geq m} H^{j-k}(X, \Lambda^k \Omega)$.

 $H^j_{DR}(X_{\mathbb{C}},\mathbb{C}(r))$ is defined to be $H^j_{DR}(X_{\mathbb{C}},\mathbb{C})$ but with the Hodge filtration F given by $F_m = G_{m+r}$. If M is $H^j(X,\mathbb{Z}(r))$ we define $t_M = t_{j,r}$ to be $H^j_{DR}(X_{\mathbb{C}},\mathbb{C}(r))/F_0 =$

 $= \prod_{k < r} H^{j-k}(X_{\mathbb{C}}, \Lambda^k \Omega_{X_{\mathbb{C}}/\mathbb{C}}) = \prod_{\sigma} \prod_{k < r} H^{j-k}(X_{\sigma}, \Lambda^k \Omega_{X_{\sigma}/\mathbb{C}}). \text{ Here } \sigma \text{ runs through all embeddings of the number field } K \text{ into } \mathbb{C}. \text{ and } X_{\sigma} = X \times_{O_K} \mathbb{C} \text{ where the map from } O_K \text{ to } \mathbb{C} \text{ is induced by } \sigma. \text{ The standard integral structure on } t_M \text{ is given by}$

 $\prod_{\sigma} \prod_{k < r} H^{j-k}(X, \lambda^k \Omega_{X/O_K})$, where λ^k denotes the kth derived exterior power. (See the appendix for a discussion of derived exterior powers).

§2.3 Maps between cohomology groups.

Let $M=M_{j,r}$ be the motive $H^j(X,\mathbb{Z}(r))$. Let $M^*=H^{2d-2-j}(X,\mathbb{Z}(d-1-r))$, and $N=M^*(1)=H^{2d-2-j}(X,d-r)$. The classical period map $\alpha_{j,0}$ maps $H_B(M_{j,0})_{\mathbb{C}}=H^j_B(X_{\mathbb{C}},\mathbb{C})$ to $H_{DR}(M)_{\mathbb{C}}=H^j_{DR}(X_{\mathbb{C}},\mathbb{Z}_{\mathbb{C}})$, which induces a map $\alpha_{j,r}=(2\pi i)^r\alpha_{j,0}$) from $H_B(M)^+_{\mathbb{C}}=H^j_B(X,\mathbb{Z}(r))^+_{\mathbb{C}}$ to $(t_M)_{\mathbb{C}}$. We now define a new map γ_M (which we call the enhanced period map) as follows: $H_{DR}(M)$ has a decreasing Hodge filtration $F_{q'}(M)$. Let $H^{q'}=F_{q'}/F_{q'+1}$. Let $h_{q'}$ be the dimension of $H_{q'}$. Then $H_{DR}(M)$ has the direct sum Hodge decomposition $\coprod H^{q'}$. We decompose $\alpha_M=\alpha_{j,r}$ into the direct sum of the maps $\alpha^{q'}(M)$ where $\alpha^{q'}$ is the map α_M followed by the projection onto $H^{q'}$. Let Γ be the usual gamma-function. Recall that the weight w(M) of M is equal to j-2r, Now let $\gamma^{q'}(M)$ be $\Gamma^*(-w(M)+q'))\alpha^{q'}(M)$ and let $\gamma_{j,r}$ be the isomorphism $\tilde{\gamma}_M=\coprod_{q'}\gamma^{q'}(M)$. Since $h_{q'}=h(p,q)$, where p+q=j and q'=q-r, the determinant of $\gamma_{j,r}$ is equal to the determinant of $\alpha_{j,r}$ multiplied by $\prod_{q'}\Gamma^*(-w(M)+q')^{h_{q'}}$ which is equal to $\prod_p\Gamma^*(r-p)^{h(p,q)}$, where p+q=j and the product is over all p between 0 and j.

Consider the following diagram of exact sequences:

$$0 \longrightarrow (H_B(M)^+)_{\mathbb{C}} \stackrel{i}{\longrightarrow} (H_B(M))_{\mathbb{C}} \stackrel{p}{\longrightarrow} (H_B(M)^-)_{\mathbb{C}} \longrightarrow 0$$

$$\downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$0 \longrightarrow (F_0(M))_{\mathbb{C}} \stackrel{j}{\longrightarrow} (H_{DR}(M))_{\mathbb{C}} \stackrel{q}{\longrightarrow} (t_M)_{\mathbb{C}} \longrightarrow 0$$

Let $\gamma_M = q \circ \tilde{\gamma}_M \circ i$. Let $\beta_M = p \circ \tilde{\gamma}_M^{-1} \circ j$. Diagram -chasing immediately shows that $\tilde{\gamma}_M$ induces isomorphisms from Ker γ_M to Ker β_M and from Coker γ_M to Coker β_M .

Proposition 2.3.1. The exact sequence of complex vector spaces

$$2.3.1(M) 0 \to Ker\gamma_M \to (H_B(M)^+)_{\mathbb{C}} \to (t_M)_{\mathbb{C}} \to Coker\gamma_M \to 0$$

is dual to the exact sequence

$$2.3.2(N) 0 \to Ker\beta_N \to (F_0(N))_{\mathbb{C}} \to (H_B^-(N))_{\mathbb{C}} \to Coker\beta_N \to 0$$

.

Proof. $F_0(N)_{\mathbb{C}}$ is the Serre dual of $(t_M)_{\mathbb{C}}$. $H_B(N)^-$ may be identified with $H_B(M^*)^+$, which is the Poincarè dual of $H_B(M)^+$. $H_B^{2d-2}(X,\mathbb{C}(d-1))$ may be canonically identified

with $H^{2d-2}_{DR}(X,\mathbb{C})$. Poincarè duality is compatible with Serre duality, which implies the proposition.

We from now on choose an arbitrary basis for Ker γ_M , the basis for Ker β_M induced by the isomorphism between Ker γ_M and Ker β_M , the basis for Coker γ_N induced by the above duality, and the basis for Coker β_N induced by the isomorphism between Coker γ_N and Coker β_N We will use these integral structures on the various kernels and cokernels,

If A is a finitely generated abelian group, Let A_{tor} denote the torsion subgroup of A and \bar{A} denote A/A_{tor} .

If $\phi: A \to B$ is a homomorphism of finitely generated abelian groups, let ϕ be the induced homomorphism from \bar{A} to \bar{B} and let ϕ_{tor} be the induced homomorphism from A_{tor} to B_{tor} .

Lemma 2.3.2. Let $0 \to A \xrightarrow{f} B \xrightarrow{g} C \to 0$ be an exact sequence of finitely generated abelian groups. There is a natural isomorphism from Ker $\bar{g}/Im\bar{f}$ to Coker g_{tor} , and the determinant of $0 \to \bar{A} \to \bar{B} \to \bar{C} \to 0$ is equal to the Euler characteristic $|B_{tor}|/|A_{tor}||C_{tor}|$ of $0 \to A_{tor} \to B_{tor} \to C_{tor} \to 0$.

Proof. Exercise

Proposition 2.3.3. $\chi(2.3.2(M)) = \chi(\tilde{\gamma}_M)\chi(2.3.1(M))$

Proof. Let Λ denote highest exterior power. Let $A_1^* = H_B(M)^+$, let $A_2^* = H_B(M)$ and let $A_3^* = H_B(M)^-$. Let A_j be a generator of $\Lambda(\bar{A}_j^*)$. Let $B_1^* = F_0(M)$, $B_2^* = H_{DR}(M)$, and $B_3^* = t_M$. Let B_j be a generator of $\Lambda(\bar{B}_j^*)$. Let C_1 be a generator of any lattice L_1 in $\Lambda(Ker(\beta_M)) = \Lambda(Ker(\gamma_M))$ and C_2 be a generator of any lattice L_2 in $\Lambda(Coker(\beta_M)) = \Lambda(Coker(\gamma_M))$.

Let $A = A_2/A_1A_3$ and $B = B_2/B_1B_3$. Let $a_j = |(A_j^*)_{tor}|$ and $b_j = |(B_j^*)_{tor}|$. Let $\chi(A_{tor}) = a_1a_3/a_2$ and $\chi(B_{tor}) = b_1b_3/b_2$. $\chi_{tor}(2.3.1(M)) = b_3/a_1$, and $\chi_{tor}(2.3.2(M)) = a_3/b_1$.

We have
$$\chi_{tor}(2.3.1(M))/\chi_{tor}(2.3.2(M)) = b_1b_3/a_1a_3$$
, and $\chi(2.3.1(M))/\chi(2.3.2(M)) =$

$$= (det(2.3.1(M))/det(2.3.2(M)))(\chi_{tor}(2.3.2(M))/\chi_{tor}(2.3.1(M))) =$$

$$= B_1 B_3 a_1 a_3 / A_1 A_3 b_1 b_3 == (B_2 det(B)/A_2 det(A))(\chi(A_{tor} a_2/\chi(B_{tor} b_2 a_2)))$$

Lemma 2.3.2 implies that this equals B_2a_2/b_2A_2 , which is $\chi(\bar{\gamma}_M)^{-1}$.

Beilinson defines Chern class maps from the algebraic K-theory groups to Deligne cohomology. Let $\gamma_{j,r} = \gamma_M$ where $M = h^j(X, \mathbb{Z}(r))$. In our language, Beilinson's map becomes

a map $c_{j,r}$ (for $j \leq 2r-2$) from $H_W^{j+1}(X,\mathbb{Z}(r))_{\mathbb{C}}$ to Coker $\gamma_{j,r}$. Let N be the motive $M^*(1) = h^{2d-2-j}(X,\mathbb{Z}(d-r))$. Since Ker γ_N may be identified with the dual of Coker $\gamma_{2d-2-j,d-r}$, we also have the dual $b_{j,r}$ (for $j \geq 2d-2r$) of Beilinson's Chern class map which maps Ker $\gamma_{2d-2-j,d-r}$ to $H_W^{2d-j}(X,\mathbb{Z}(d-r))_{\mathbb{C}}$. We have:

Conjecture 2.3.4. (Beilinson) If $j \leq 2r - 3$ the maps $c_{j,r}$ and $b_{j,r}$ are isomorphisms.

Recall that by definition $H^{2r+1}_W(X,\mathbb{Z}(r))_{\mathbb{C}}$ is dual to $H^{2(d-r)}_{et}(X,\mathbb{Z}(d-r))_{\mathbb{C}}$ which is the same as $H^{2(d-r)}_{Zar}(X,\mathbb{Z}(d-r))_{\mathbb{C}}$, i.e. codimension (d-r) cycles on X modulo rational equivalence. Recall that $H^{2(d-r)}_{et}(X,\mathbb{Z}(d-r))^1$ denote cycles homologically equivalent to zero, and $H^{2(d-r)}_{et}(X,\mathbb{Z}(d-r))^2$ denote cycles modulo homological equivalence. Let $H^{2r+1}_W(X,\mathbb{Z}(r))^1_{\mathbb{C}}$ be the dual of $H^{2d-r}_{et}(X,\mathbb{Z}(d-r))^1_{\mathbb{C}}$

Conjecture 2.3.5. (Beilinson) There is an exact sequence

$$0 \to H_W^{2r-1}(X, \mathbb{Z}(r))_{\mathbb{C}} \stackrel{c_{2r-2,r}}{\to} Coker(\gamma_M) \to H_W^{2r+1}(X, \mathbb{Z}(r))_{\mathbb{C}}^2 \to 0$$
with $M = H^{2r-2}(X, \mathbb{Z}(r))$.

This is a slightly different but more natural variant of Beilinson's original conjecture, and it is implicitly used by Fontaine ([Fo]).

Conjecture 2.3.6. There is an exact sequence

$$0 \to H^{2r}_W(X, \mathbb{Z}(r))^2_{\mathbb{C}} \to Ker(\gamma_N) \to H^{2r+2}_W(X, \mathbb{Z}(r))_{\mathbb{C}} \to 0$$

This is the dual of conjecture 2.3.5, with M replaced by $N = M^*(1) = H^{2d-2r}(X, \mathbb{Z}(d-r))$

Conjecture 2.3.7. There is an isomorphism $(H^{2r+1}_W(X,\mathbb{Z}(r))^1)_{\mathbb{C}}$ to $(H^{2r}_W(X,\mathbb{Z}(r))^1)_{\mathbb{C}}$

This is the non-degeneracy of the Arakelov intersection pairing restricted to finite cycles homologous to zero, where it is independent of metrics.

§.3 The statement of the conjecture.

We would like to first explain the relationship between Weil-ètale motivic cohomology groups and the groups which occur in Fontaine's Deligne-Beilinson conjecture. We look at the motive $M = H^j(X, \mathbb{Z}(r))$, Recall that $N = M^*(1)$ is $H^{2d-2-j}(X, \mathbb{Z}(d-r))$.

Fontaine starts with a projective non-singular algebraic variety X_0 over Spec \mathbb{Q} . He chooses a regular model X for X_0 projective and flat over Spec \mathbb{Z} . He conjectures that the following six-term sequence is always exact:

$$0 \to H^0_f(M)_{\mathbb{C}} \to Ker(\gamma_M) \to H^1_c(M)_{\mathbb{C}} \to H^1_f(M)_{\mathbb{C}} \to Coker(\gamma_M) \to H^2_c(M)_{\mathbb{C}} \to 0$$

.

If $j \leq 2r-2$, Fontaine's $H_f^1(M)$ is our $H_W^{j+1}(X,\mathbb{Z}(r))_{\mathbb{Q}}$, (We are using motivic cohomology instead of algebraic K-theory, but these two groups agree after tensoring with \mathbb{Q}). (Actually, Fontaine's group is the image of K(X) in $K(X_0)$, but we conjecture that the natural map is always injective.)

If j = 2r - 1, Fontaine's $H_f^1(M)$ is the group of codimension r cycles on X_0 homologically equivalent to zero, tensored with \mathbb{Q} ,

If
$$j \ge 2r$$
, $H_f^1(M) = 0$

Fontaine's $H_f^0(M)$ is zero unless j=2r, in which case it equals the group of codimension r cycles on X_0 modulo homological equivalence, tensored with \mathbb{Q} .

Fontaine's $H_c^i(M)$ is the \mathbb{Q} -dual of $H_f^{2-i}(N)$ for i=1,2.

For each j and r with $j \leq 2r - 3$ or $j \geq 2r + 1$, we will define a sequence of integral structures A(j,r) and A'(j,r). A(j,r) is given by:

$$(j \le 2r - 3)$$
 $c_{j,r} : H_W^{j+1}(X, \mathbb{Z}(r))_{\mathbb{C}} \to Coker(\gamma_{j,r})$

while A'(j,r) is given by:

$$(j \ge 2r+1)$$
 $b_{i,r}: Ker(\gamma_{i,r}) \to H_W^{j+2}(X,\mathbb{Z}(r))_{\mathbb{C}}$

If 1 < r < d we define an integral structure C(r) given by: C(r) =

$$0 \to H_W^{2r-1}(X, \mathbb{Z}(r))_{\mathbb{C}} \stackrel{c_{2r-2,r}}{\to} Coker(\gamma_{2r-2,r}) \to H_W^{2r+1}(X, \mathbb{Z}(r))_{\mathbb{C}} \stackrel{e_r}{\to} \\ \to H_W^{2r}(X, \mathbb{Z}(r))_{\mathbb{C}} \to Ker(\gamma_{2r,r}) \stackrel{b_{2r,r}}{\to} H_W^{2r+2}(X.\mathbb{Z}(r))_{\mathbb{C}} \to 0$$

Here e_r is induced by the Arakelov intersection pairing. We give these vector spaces the standard integral structures previously defined in §. 2.2.

Conjectures 2.3.4, 2.3.5, 2.3.6 and 2.3.7 imply that these sequences are exact.

We give degrees to the terms of these complexes by requiring that $Ker(\gamma_M)$ has even degree and $Coker(\gamma_M)$ has odd degree. (These sequences are all truncations of modified versions of Fontaine's six-term sequence in [Fo] and this convention makes the degrees agree)

Finally we define exact sequences $B(j,r)_{\mathbb{C}}$ given for all j and r by

$$0 \to (Ker(\gamma_M) \to (H_B^j(X, \mathbb{Z}(r))^+)_{\mathbb{C}} \stackrel{\gamma_M}{\to} t_{j,r} \to Coker(\gamma_M) \to 0.$$

We put $Ker(\gamma_M)$ in degree zero.

The integral structures on the cohomology groups here are induced by the standard integral structures defined in §2.2.

Let
$$\chi_{A,C}(X,r) = \chi(C(r)) \prod_{j=0}^{\min(2d-1,2r-3)} (\chi(A(j,r))^{(-1)^j} \prod_{j=\max(0,2r+1)}^{2d-1} \chi(A'(j,r))^{(-1)^j}$$

Let $\chi_B(X,r) = \prod_{j=0}^{\infty} \chi(B(j,r))^{(-1)^j}$ Let $\chi(X,r) = \chi_{A,C}(X,r)\chi_B(X,r)^{(-1)}$

Conjecture 3.1. Give all groups in the above exact sequences their standard integral structures.

$$\zeta^*(X, r) = \chi(X, r)$$

up to sign and powers of 2. (Note that A(j,r) and B(j,r) are torsion for $j \geq 2d-1$, and zero for j large).

If $j \neq 2r - 1$ each of the terms Ker (γ_M) and Coker (γ_M) occurs exactly twice in the conjecture with degrees of opposite parity, so the conjecture is independent of the choice of integral structure. If j = 2r - 1, Ker (γ_M) and Coker (γ_M) are both zero.

Proposition 3.2. a)If $0 \le j \le 2r - 3$, then $2(d - r) + 1 \le 2d - 2 - j \le 2d - 2$ and the integral structures $A'(2d - 2 - j, d - r)_{tf}$ and $A(j,r)_{tf}$ are dual. Hence det(A(j,r)) = det(A'(2d - 2 - j, d - r))

b) The integral structures $C(r)_{tf}$ and $C(d-r)_{tf}$ are dual. Hence det(C(r)) = det(C(d-r)).

Proposition 3.3. $\chi_{A,C}(X,r) = \chi_{A,C}(X,d-r)$

Proof. $\chi(C(r)) = \det(C(r))/\tan(C(r))$. C(r) is dual to C(d-r), so $\det(C(r)) = \det(C(d-r))$. $\chi(A(j,r)) = \det(A(j,r))/\tan(A(j,r))$. $\chi(A'(j,r)) = \det(A'(j,r))/\tan(A'(j,r))$. $\Lambda(\bar{j},r)$ is dual to $\Lambda'(2d-\bar{2}-j,d-r)$ so $\det(A(j,r)) = \det(A'(2d-2-j,d-r))$ and $\det(A'(j,r)) = \det(A(2d-2-j,d-r))$.

On the other hand

$$torC(r)\prod_{j=0}^{2r-3}tor(A(j,r))^{(-1)^{j}}\prod_{j=2r+1}^{2d-1}tor(A'(j,r)^{(-1)^{j}}=\prod_{j=1}^{2d+1}|H^{j}(X,\mathbb{Z}(r))_{tor}|^{(-1)^{j}}$$

which is equal by duality to

$$\prod_{j=1}^{2d+1} |H^{j}(X, \mathbb{Z}(d-r))_{tor}|^{(-1)^{j}}$$

which is equal to

$$tor(C(d-r)) \prod_{j=0}^{2d-2r-3} tor(A(j,d-r))^{(-1)^{j}} \prod_{j=2d-2r+1}^{2d-1} tor(A'(j,d-r))^{(-1)^{j}}$$

§4 The Euler characteristic of the period map.

Let K be the integral closure of \mathbb{Q} in the function field of X. Then we can view X as a scheme over $S = \operatorname{Spec} O_K$, and $X_{\mathbb{C}}$ is canonically isomorphic to $\prod_{\sigma} X \times_S \operatorname{Spec} \mathbb{C}$, where the product is taken over all embeddings σ of K in \mathbb{C} . Let $\kappa(v)$ be the residue field of the closed point v, and let $X_v = X \times_S \operatorname{Spec} \kappa(v)$

Recall that A is the positive rational number which appears in the conjectured functional equation $A^{s/2}\Gamma(X,s)\zeta(X,s) = \pm A^{(d-s)/2}\Gamma(X,d-s)\zeta(X,d-s)$ for $\zeta(X,s)$. Let $\omega_{X/S}$ be the relative canonical class of X over S.

Definition 4.1.
$$A'_v = (\Delta_X . \Delta_X)_v = (-1)^d c_{dX_v}^X (\Omega_{X_v/O_v}) \in CH_0(X_{\kappa(v)}).$$
 $A' = (\Delta_X . \Delta_X)_S = \prod_v A'_v.$

(This definition is taken from [KS]. We will not use it directly in what follows. What we need is stated in Theorems 4.2 and 4,3)

Here the product is taken over all closed points v of S such that X is not smooth over S at v. Note that the Chern class $c_{d_{X_v}}^X(\Omega)$ is equal to 1 if X is smooth over S at v.

Theorem 4.2. (Bloch-Kato-Saito) If $d \leq 3$, A' = A. If strong resolution of singularities holds for schemes of finite type over Spec \mathbb{Z} , then A' = A in general.

Proof. This is the main result (Theorem 6.2.3) of [KS](the conductor formula of Bloch). Let $O_v = O_K$ localized at v. Let $X_v = X \times_{O_K} O_v$. **Theorem 4.3.** The Euler characteristic of the cone $C_{m,v}$ of the map from

 $R\Gamma(X_v, \lambda^m \Omega_{X_v/O_v})$ to $R\Gamma(X_v, RHom(\lambda^{d-1-m}\Omega_{X_v/O_v}, \omega_{X_v/O_v}))$ induced by Serre duality is equal to $(A')_v^{(-1)^{m+1}}$.

Proof. This is Corollary 4.9 of [Sa].

Theorem 4.4. The Euler characteristic of the classical period map α from de Rham to singular cohomology of $X_{\mathbb{C}}$ with respect to the canonical integral structures is $(A')^{d/2}(2\pi i)^{\chi(X_{\mathbb{C}})(d-1)/2}$.

This theorem will follow from the following propositions.

Proposition 4.5. Let H^j be $H^j_{DR}(X_{\mathbb{C}}, \mathbb{C}) = \prod_{\sigma} H^j_{DR}(X_{\sigma}, \mathbb{C})$ with its canonical integral structure. Let G^j be $H^j_{DR}(X_{\mathbb{C}}, \mathbb{C})$ with the integral structure given by

 $\prod_{\sigma} \prod_{k=0}^{j} Ext_{X_{\sigma}}^{2d-2-j+k}(\lambda^{d-1-k}\Omega_{X/O_{K}}, \omega_{X/O_{K}}). \text{ Then the Euler characteristic } \chi_{I} \text{ of the identity map from } H^{*} \text{ to } G^{*} \text{ with respect to the given integral structures is } (A')^{d}.$

Proof. Let $\Omega = \Omega_{X_{\sigma}/\mathbb{C}}$ and $\Omega_K = \Omega_{X/O_K}$, By Serre duality, $H^{j-k}(X_{\sigma}, \Lambda^k\Omega)$ is canonically isomorphic to $Ext_{X_{\sigma}}^{k-j+1-d}(\Lambda^{d-1-k}\Omega, \omega_{\mathbb{C}})$. So $A_{j,k} = H^{j-k}(X, \lambda^k\Omega_K)$ and $B_{j,k} = Ext_X^{j-k}(\lambda^{d-1-k}\Omega_K, \omega)$ give two different integral structures on $H^{j-k}(X_{\sigma}, \Lambda^k\Omega)$.

Let $d_{j,k}$ be the determinant of the identity map with respect to these two integral structures. (This is independent of the choice of basis for the integral structures, up to sign.) Then Theorem 4.3 asserts that $\prod_j (d_{j,k})^{(-1)^{j-k}} \cdot \chi((A_{j,k})_{tor})^{-1} \chi((B_{j,k})_{tor}) = (A')_{\sigma}$ if k is even and is equal to $(A')_{\sigma}^{-1}$ if k is odd. By Serre duality, $\chi((A_{j,k})_{tor}) = (\chi((B_{j,k})_{tor})^{-1}$.

We conclude easily from this that $\prod_k \prod_j (d_{j,k})^{(-1)^j} (\chi(\lambda^k(\Omega))_{tor}))^{-2} = (A')^d$ and the product on the left is the Euler characteristic of the identity map.

Let Q_{DR} denote the de Rham cup product, and Q_B denote the cup product in singular cohomology. Let $\alpha^j: H^j_{DR}(X_{\mathbb{C}}, \mathbb{C}) \to H^j_B(X_{\mathbb{C}}, \mathbb{C})$ be the classical period map, We know that Q_{DR} and Q_B are compatible, i.e. $Q_B(\alpha^j(a), \alpha^k(b)) = \alpha^{j+k}(Q_{DR}(a, b))$

Proposition 4.6. Let $[v_{i,j}]$ be a basis of $H^j_{DR}(X_{\mathbb{C}}, \mathbb{C})$. Let $[u_{i,j}]$ be a basis of $H^j_B(X_{\mathbb{C}}, \mathbb{C})$ coming from a basis of $H^j(X_{\mathbb{C}}, \mathbb{Z})$ modulo torsion. Let E_j be the matrix of α^j with respect to the bases $[v_{i,j}]$ and $[u_{i,j}]$. Then

$$\left(\prod_{i} (\det E^{j})^{(-1)^{j}}\right)^{2} = \prod_{i} \det(Q_{DR}(v_{i,j}.v_{k,2d-2-j}))^{(-1)^{j}} (2\pi i)^{\chi(X_{\mathbb{C}})(d-1)}$$

Proof. Fix j. Since on the top-dimensional cohomology group $H_{DR}^{2d-2}(X_{\mathbb{C}}, \mathbb{C})$, $\phi(1_{DR}) = (2\pi i)^{d-1}1_B$, we have $\phi(Q_{DR}(v_{i,j}, v_{k,2d-2-j})) = Q_B(\phi(v_{i,j}), \phi(v_{k,2d-2-j}))$ which implies $(2\pi i)^{(d-1)B_j} \det(Q_{DR}(v_{i,j}, v_{k,2d-2-j})) = \det Q_B(\phi^j(v_{i,j}), \phi^{2d-2-j}(v_{k,2d-2-j})) = \det (E^j) \det(E^{2d-2-j}) \det(Q_B(u_{i,j}, u_{k,2d-2-j})) = \pm \det(E^j) \det(E^{2d-2-j})$. Multiplying by $(-1)^j$ and taking products over j, we obtain the proposition.

Proposition 4.7. Let $d_j = \prod_k d_{j,k}$ be the determinant of the identity map with respect to the two given integral structures on $H^j_{DR}(X_{\mathbb{C}}, \mathbb{C})$. Then a) $d_j = \det(Q_{DR}(x_{i,j}, x_{k,2d-2-j}))$ b) $\prod d_j^{(-1)^j} = \prod \det((Q_{DR}(x_{i,j}, x_{k,2d-2-j})^{(-1)^j})$

Proof. By the compatibility of cup product and Serre duality, $det((Q_{DR}x_{i,j}, y_{k,2d-2-j}))$ is equal to 1. Then 4.7a follows from the definition of d_j , and of course implies 4.7b..

Proof of Theorem 4.4:

Recall that by definition and Serre duality $\chi(\alpha) = \prod_j (\det E^j)^{(-1)^j} \chi((H_{DR})_{tor})^{(-2)}$ and $\chi_I = \prod_j d_j^{(-1)^j} \chi((H_{DR})_{tor})^{-1}$. Substituting x for v in Proposition 4.6 and using Proposition 4.5 we obtain

$$(A')_{\sigma}^{d}(\chi(H_{DR}^{*}(X)_{tor})^{2} = ((\prod_{i} det(Q_{DR}([x_{i,j}], [x_{k,2d-2-j}])^{(-1)^{j}}))$$

Proposition 4.6 implies

$$(A')_{\sigma}^{d}(\chi(H_{DR}^{*}(X)_{tor})^{2} = (2\pi i)^{-\chi(X_{\mathbb{C}})(d-1)}(\prod det(\alpha^{j})^{(-1)^{j}})^{-2}$$

which implies

$$(2\pi i)^{\chi(X_{\mathbb{C}})(d-1)/2} (A')_{\sigma}^{d/2} = \det(\alpha^*) \chi((H_{DR}^*(X))_{tor})^{(-1)}$$

By Poincarè duality, $\chi(H_B^*(X_{\mathbb{C}}, \mathbb{Z})_{tor}) = 1$, so

$$\det(\alpha^*)\chi((H_{DR}^*(X))_{tor})^{(-1)}\chi((H_B^*(X_{\mathbb{C}},\mathbb{Z})_{tor}) = (2\pi i)^{\chi(X_{\mathbb{C}})(d-1)/2}(A')_{\sigma}^{d/2}$$

$$\chi(\alpha) = (2\pi i)^{(\chi(X_{\mathbb{C}})(d-1)/2} (A')_{\sigma}^{d/2}$$

Taking the product over σ , we get Theorem 4.4.

Corollary 4.8. The Euler characteristic χ_r of the classical period map α_r from

$$H^*_{DR}(X_{\mathbb{C}}, \mathbb{C}(r)) \ to \ H^*_B(X, \mathbb{C}, \mathbb{C}(r)) \ is \ (A')^{d/2} (2\pi i)^{\chi(X_{\mathbb{C}})((d-1)/2-r)}$$
.

This follows from the definition of twisting by r.

Recall that
$$\Gamma_{r,j} = \prod_{p+q=j} \Gamma^*(r-p)^{h(p,q)}$$
. Let $\Gamma_r = \prod_{r,j} \Gamma_{r,j}^{(-1)^{j+1}}$

Corollary 4.9. The Euler characteristic $\chi(\gamma_r)$ of the enhanced period map γ_r is

$$\Gamma_r(A')^{d/2} (2\pi i)^{\chi(X_{\mathbb{C}})(((d-1)/2-r))}$$

Proof: By the remarks at the beginning of section 2.3, $\chi(\gamma_r)$ is equal to $\chi(\alpha_r)$ multiplied by $\prod_{j=0}^{2d-2} \Gamma_{r,j}^{(-1)^{j+!}} = \Gamma_r$

§5.1 Serre's functional equation and Γ -function identities

Let X_0 be a smooth projective algebraic variety of dimension d-1 over the number field K. Let j be a non-negative integer, and let $L^j(X_0, s)$ be the L-function attached by Serre in [Se2] to the j-th cohomology group of X_0 ,

Let σ be an embedding of K into \mathbb{C} , and let v be the place of K induced by σ . Let K_v be the completion of K at v. Let $X_v = X_0 \times_K \mathbb{C}$, where σ maps F into \mathbb{C} , and let $\Omega = \Omega_{X_v/\mathbb{C}}$. Recall that Hodge theory gives us a decomposition $H^j_{DR}(X_v) = \coprod H^{p,q}_v$, where the sum is taken over pairs (p,q) such that p+q=j and $H^{p,q}_v = H^q(X_v,\Omega^p)$. Let c be the automorphism of X_v induced by complex conjugation acting on \mathbb{C}/K_v . Then if j is even and equal to 2n, c acts as an involution on $H^{n,n}_v$. Let $h_v(p,q)$ be the dimension of $H^{p,q}_v$.

Then
$$H_v^{n,n} = H_v^{(n,+)} \oplus H_v^{(n,-)}$$
, where

$$H_v^{(n,+)} = \{x \in H_v^{n,n}, c(x) = (-1)^n x\}$$

$$H_v^{(n,-)} = \{x \in H_v^{n,n}, c(x) = (-1)^{(n+1)}x\}$$

Let
$$h_v(n, +) = dim H_v^{(n,+)}$$
, and $h_v(n-) = dim H_v^{(n,-)}$.

Let B_v^j be the rank of $H^j(X_v, \mathbb{Z})$ and let $(B_v^j)^+$ be the rank of the subgroup of $H^j(X_v, \mathbb{Z})$ left fixed by c. Let $(B_v^j)^- = B_v^j - (B_v^j)^+$. Note that if j = 2n, $(B_v^j)^+$ is equal to $\Sigma h_v(p,q) + h_v(n,+)$ if n is even and is equal to $\Sigma h_v(p,q) + h_v(n,-)$ if n is odd, where the sum is taken

over all pairs (p, q) where p < q and p + q = j. Let $(B_v^{j,r})^-$ be $(B_v^j)^-$ if r is even, and $(B_v^j)^+$ if r is odd.

Let
$$\Gamma_{\mathbb{C}}(s) = (2\pi)^{-s}\Gamma(s)$$
. Let $\Gamma_{\mathbb{R}}(s) = \pi^{-s/2}\Gamma(s/2)$.

Serre gives the functional equation $\phi^j(s) = \pm \phi(j+1-s)$, where $\phi(s) = L^j(s) A_j^{s/2} \Gamma^j(s)$, A_j is a certain positive integer, and $\Gamma^j(s)$ is described as follows:

 $\Gamma^j(s) = \prod_v \Gamma^j_v(s)$ where $\Gamma^j_v(s) = \prod_{p+q=j} (\Gamma_{\mathbb{C}}(s-inf(p,q))^{h_v(p,q)})$ if v is a complex place of F,

 $\Gamma_v(s) = \Gamma_{\mathbb{R}}(s-n)^{h_v(n,+)}\Gamma_{\mathbb{R}}(s-n+1)^{h_v(n,-)}\prod_{p< j-p}\Gamma_{\mathbb{C}}(s-p)^{h_v(p,j-p)}$, if v is a real place of F.

We observe that it is an easy computation that $\Gamma_v^j(s) = \Gamma_v^{2d-2-j}(s+d-j-1)$, so that with our earlier observation that at least in the smooth case $L^j(s) = L^{2d-2-j}(s+d-j-1)$ we obtain that Serre's functional equation is equivalent to the functional equation $\phi^j(s) = \pm \phi^{2d-2-j}(d-s)$.

Theorem 5.1.1. Let v be a real place of K.

If j is even, $(\Gamma_v^j)^*(r)/(\Gamma_v^{2d-2-j})^*(d-r)$ is equal up to sign and powers of 2 to $\prod_p (\Gamma^*(r-p))^{h_v(p,q)} \pi^{-B_v^j(r-j/2)+(B_v^{j,r})^-}$. (The products run over $0 \le p \le B_v^j$).

If j is odd, $(\Gamma_v^j)^*(r)/(\Gamma_v^{2d-2-j})^*(d-r)$ is equal up to sign and powers of 2 to $\prod_p \Gamma^*(r-p)^{h_v(p,q)} \pi^{-B_v^j(r-(j+1)/2)}$

. Proof. We consider the case when j is even. (The case when j is odd is similar, but simpler). Let j=2n. Fix v and let q=j-p. First look at terms where $p\neq q$. Let p'=d-1-p and q'=d-1-q, so p'+q'=2d-2-j. We have

$$\prod_{p < q} \Gamma_{\mathbb{C}}^*(r-p)^{h_v(p,q)} / \prod_{p' < q'} \Gamma_{\mathbb{C}}^*(d-r-p')^{h_v(p',q')} =$$

,

$$\prod_{p < q} \Gamma_{\mathbb{C}}^*(r-p)^{h_v(p,q)} / \prod_{p > q} \Gamma_{\mathbb{C}}^*(1-r+p)^{h_v(p,q)}$$

since $h_v(p,q) = h_v(p',q')$ by Serre duality.

By definition of $\Gamma_{\mathbb{C}}$, this is equal to

$$\prod_{p < q} \Gamma^*(r-p)^{h_v(p,q)} / \prod_{p > q} \Gamma^*(1-r+p)^{h_v(p,q)}$$

multiplied by

$$(2\pi)^{-(\sum_{p< q}(h_v(p,q)(r-p)-\sum_{p>q}(h_v(p,q)(1-r+p)))}$$

This product is then equal to

$$\pm \prod_{p \neq n} \Gamma^*(r-p)^{h_v(p,q)} (2\pi)^{-(\Sigma_{p,q}(h_v(p,q))((r-p)-(1-r+j-p))}$$

because of the relation $\Gamma^*(r) = \pm \Gamma^*(1-r)^{-1}$ for integral r which follows immediately from the functional equation for the Gamma function. We then obtain:

(5.1.1)
$$\pm \prod_{p \neq n} \Gamma^*(r-p)^{h_v(p,q)} (2\pi)^{-(B_v^j - h_v(n,n))(r-(j+1)/2)}$$

We now look at the terms involving n with v still fixed.

We first observe that the functional equation for the Gamma function implies that $\Gamma^*(a/2)\Gamma^*((2-a)/2)$ equals $\pm \pi$ if a is an odd integer and equals ± 1 if a is an even integer. We compute:

$$\Gamma_{\mathbb{D}}^*((r-n))^{h_v(n,+)}\Gamma_{\mathbb{D}}^*((r-n+1))^{h_v(n,-)}$$

multiplied by

$$\Gamma_{\mathbb{R}}^*((d-r-(d-1-n))^{-h_v(n,+)}\Gamma_{\mathbb{R}}^*(d+1-r-(d-1-n))^{-h_v(n,-)}$$

which we rewrite as

(5.1.2)
$$(\Gamma^*((r-n)/2)(\Gamma^*(1-r+n)/2)^{-1})^{h_v(n,+)}$$

multiplied by

(5.1.3)
$$\Gamma^*((r-n+1)/2)(\Gamma^*((n+2-r)/2)^{-1})^{h_v(n,-)}$$

multiplied by

(5.1.4)
$$\pi^{-(h_v(n,+)(2r-2n-1)/2+h_v(n,-)(2r-2n-1)/2)}$$

First assume that r - n is odd. By the functional equation,

(5.1.5)
$$\Gamma^*((1-r+n)/2)^{-1} = \pm \Gamma^*(r-n+1)/2$$

(5.1.6)
$$\Gamma^*((n+2-r)/2)^{-1} = \pm \pi^{-1}\Gamma^*((r-n)/2)$$

Now recall the duplication formula for the Gamma function;

(5.1.7)
$$\Gamma(2s) = 2^{2s-1}\Gamma(s)\Gamma(s+1/2)/\sqrt{\pi}$$

Then (5.1.2) becomes (using (5.1.5) and (5.1.7))

$$(\pm 2^{n-r+1}\Gamma^*(r-n)\sqrt{\pi})^{h_v(n,+)}$$

and (5.1.3) becomes

$$(\pm 2^{n-r}\Gamma^*(r-n)\sqrt{\pi})^{-h_v(n,-)}$$

while (5.1.4) is

(5.1.8)
$$\pi^{-h_v(n,n)(r-(j+1)/2)}$$

So, up to sign and powers of 2, our product has become

(5.1/9)
$$\Gamma^*(r-n)^{h_v(n,n)} \pi^{(h_v(n,n)(r-(j+1)/2) + h_v(n,+)/2 - h_v(n,-)/2}$$

Multiplying (5.1.1) by (5.1.9) we get

(5.1.10)
$$\prod_{n} \Gamma^*(r-p)^{h_v(p,q)} \pi^{-(B_v^j(r-(j+1/2))) + h_v(n,+)/2 - h_v(n,-)/2)}$$

which equals

(5.1.11)
$$\prod_{p} \Gamma^*(r-p)^{h_v(p,q)} \pi^{-(B_v^j(r-j/2) + (B_v^j)^+}$$

if n is even (so r odd) and equals

(5.1.12)
$$\prod_{p} \Gamma^*(r-p)^{h_v(p,q)} \pi^{-B_v^j(r-j/2) + (B_v^j)^-}$$

if n is odd (so r even). since $(B_j - h_v(n, n)/2) + h_v(n, +)$ is equal to $(B_v^j)^+$ if n is even and $(B_v^j)^-$ if n is odd.

The proof for r-n even is identical, except for switching $h_v(n,+)$ and $h_v(n,-)$.

Theorem 5.1.2. Let v be a complex place of K. Then $(\Gamma_v^j)^*(r)/(\Gamma_v^{2d-2-j})^*(d-r)$ is equal up to sign and powers of 2 to $(\prod_p \Gamma^*(r-p))^{2h(p,q)} \pi^{-B_v^j(2r-(j+1))}$

Proof. Let q = j - p, p' = d - 1 - p and q' = d - 1 - q. Note that $0 \le p, q, p', q', \le d - 1$. We write $(\Gamma^j)_v^*(r)/(\Gamma^{2d-2-j})_v^*(d-r) = 0$

$$\prod_{p} \Gamma_{\mathbb{C}}^*(r - Inf(p, q))^{h_v(p, q)} / \prod_{p'} \Gamma_{\mathbb{C}}(d - r - Inf(p', q'))^{h_v(p', q')}$$

which is equal to

(5.1.13)
$$\prod_{p} \Gamma^*(r - Inf(p, q))^{(h_v(p, q))} / \prod_{p'} \Gamma^*(d - r - Inf(p', q'))^{h_v(p', q')}$$

multiplied by

$$(5.1.14) (2\pi)^{-\Sigma_p h_v(p,q)(r-Inf(p,q)) - \Sigma_{p'} h_v(p',q')(d-r-Inf(p',q'))}$$

Since $h_v(p', q') = h_v(p, q)$, the functional equation for the Gamma function transforms (5.1.13) into (up to sign)

(5.1.15)
$$\prod_{p} \Gamma^{*}(r - Inf(p, q))^{h_{v}(p, q)} \prod_{p} \Gamma^{*}(r - Sup(p, q))^{h_{v}(p, q)}$$

,

which in turn equals

$$(\prod_{p} \Gamma^*(r-p)^{h_v(p,q)})^2$$

Now (5.1.14) easily transforms into $(\pi^{2r-j-1})^{-\Sigma_p h_v(p,q)}$, which becomes $\pi^{-(B_v^j(2r-(j+1)))}$.

§.6 Compatibility of the conjecture with the functional equation.

Starting with Serre's conjectured functional equation for cohomological L-functions described in the previous section, Bloch, Kato, and T. Saito conclude that the following functional equation holds for the zeta-function of X;

Conjecture 6.1. Let
$$\phi(X,s) = \zeta(X,s)A^{s/2}\Gamma(X,s)$$
. Then $\phi(X,s) = \pm \phi(X,d-s)$

Here the constant A = A(X) is obtained by taking the alternating product of the constants A_j which occur in the conjectured functional equation for Serre's L-function L_j and modifying it by terms coming from degenerate fibers. It is still a positive rational number.

The generic fiber X_0 of X is a projective algebraic variety smooth over a number field $K = H^0(X_0, O_{X_0})$. Let $\Gamma(X, s) = \prod_j \prod_\sigma \Gamma(X_{v(\sigma)}^j, s)^{(-1)^j}$, where Γ_v^j was defined in the previous section. If we rewrite our conjecture as $\zeta^*(X, r) = \chi(X, r)$, then what we want to show is that $\zeta^*(X, r)/\zeta^*(X, d-r) = \chi(X, r)/\chi(X, d-r)$, up to sign and powers of 2, where the left-hand side is computed by the functional equation.

Proposition 6.2. Conjecture 6.1 implies

$$\zeta^*(X,r)/\zeta^*(X,d-r) = A^{d/2-r} \prod_j (\prod_{p+q=j} \Gamma^*(r-p)^{h(p,q)})^{(-1)^j} \pi^{-\chi(X_{\mathbb{C}})(r-(d-1)/2) + \chi^-(X_{\mathbb{C}},\mathbb{Z}(r))}$$

.

This is an immediate consequence of Theorems 5.1.1 and 5.1.2, remembering that $B_j = B_{2d-2-j}$.

We now wish to compute $\chi(X,r)/\chi(X,d-r)$ and show that it agrees with the expression in Proposition 6.2. We first recall that $\chi(X,r)=\chi_{A,C}(X,r)\chi_B(X,r)$ and that $\chi_{A,C}(X,r)/\chi_{A,C}(X,d-r)=1$, by Proposition 3.3.

We now have to look at $\chi_B(X,r)/\chi_B(X,d-r)$.

Lemma 6.3. Let $\chi_{i,k}$ be the Euler characteristic of the identity map from the complex vector space $H^i(X_{\mathbb{C}}, \Lambda^k \Omega_{X_{\mathbb{C}}})$ with the integral structure $H^i(X, \lambda^k \Omega_X)$ to the same vector space with the integral structure $\underline{RHom}(H^{d-1-i}(X, \lambda^{d-1-k}\Omega_X), \omega)$. Then $\prod_i \chi_{i,k}^{(-1)^i} = (A')^{(-1)^k}$.

Proof. This follows immediately from Theorem 4.3.

Lemma 6.4. Let $\theta_{j,r}$ be the Euler characteristic of the identity map from the complex vector space $t_M = \coprod_{0 \leq k < r} H^{j-k}(X_{\mathbb{C}}, \Lambda^k \Omega_{X_{\mathbb{C}}})$ with the integral structure $\coprod_{0 \leq k < r} H^{j-k}(X, \lambda^k \Omega_X)$ to the same vector space with the integral structure given by

$$\underline{RHom}(H^{d-1-j+k}(X,\lambda^{d-1-k}\Omega_X),\omega)$$

. Then
$$\prod_{j} \theta_{j,r}^{(-1)^{j}} = (A')^{r}$$
.

Proof. This follows immediately from Lemma 6.3

Lemma 6.5. Let η_j be the Euler characteristic of the identity map from $H_B^j(X_{\mathbb{C}}, \mathbb{C})^+$ (if r is odd) or $H_B^j(X_{\mathbb{C}}, \mathbb{C})^-$ (if r is even) with the integral structure $H_B^j(X, \mathbb{Z}(r-1))^-$ to the same vector space with the integral structure $H_B^j(X, \mathbb{Z}(r))^+$ or $H_B^j(X, \mathbb{Z}(r))^-$. Then $\prod \eta_j^{(-1)^j} = (2\pi i)^{\chi^-(X_{\mathbb{C}}, \mathbb{Z}(r))}$

Proof. This follows immediately from the definitions.

Let $\gamma(j,r)$ denote the integral structure $0 \to Ker\gamma_* \to H_B^j(X,\mathbb{C}(r))^+ \to t(j,r)_{\mathbb{C}} \to Coker\gamma_* \to 0$

Let $\beta(j,r)$ denote the integral structure $0 \to Ker\beta_* \to F_0(j,r)_{\mathbb{C}} \to H_B^j(X,\mathbb{C}(r))^- \to Coker\beta_* \to 0$

Proposition 6.6. a) The integral structure $\beta(j,r)$ is dual to the integral structure $\delta(2d-2-j,d-r)$ given as follows:

$$0 \to Ker\gamma_* \to (H_B^{2d-2-j}(X, \mathbb{Z}(d-1-r))_{\mathbb{C}})^- \to t(2d-2-j, d-r)_{\mathbb{C}} \to Coker\gamma_* \to 0$$

where the singular cohomology group has its standard structure and t(2d-2-j, d-r) has the integral structure dual to the standard one on $F_0(j,r)$.

b)
$$det(\beta(j,r)) = det(\delta(2d-2-j,d-r))$$

Proof. Part a) follows immediately from the compatibility of Serre and Poincarè duality, and then b) follows immediately.

Let
$$\chi(\gamma(r)) = \prod_{j=0}^{2d-2} \chi(\gamma(j,r))^{(-1)^j}$$
, $\chi(\beta(r)) = \prod_{j=0}^{2d-2} \chi(\beta(j,r))^{(-1)^j}$. and $\chi(\epsilon(d-r)) = \prod_{j=0}^{2d-2} \chi(\epsilon(j,d-r))^{(-1)^j}$.

Corollary 6.7.
$$\chi(\beta(r)) = \chi(\delta(d-r))$$

Proof. This follows from the preceding proposition and a consideration of torsion, exactly as in Proposition 3.3.

Note that $\gamma(2d-2-j,d-r)$ is the integral structure $\delta(2d-2-j,d-r)$ except that the singular cohomology group now has the integral structure given by $H_B^{2d-2-j}(X,\mathbb{Z}(d-r)_{\mathbb{C}})^+$ and the tangent space has the standard integral structure rather than the one coming from duality.

Proposition 6.8.
$$\chi(\gamma(d-r)) = (A')^{d-r}(2\pi i)^{-\chi^-(X_{\mathbb{C}},\mathbb{Z}(r))}\chi(\delta(d-r))$$

Proof. This follows immediately from Lemmas 6.4, 6.5 and the previous remark. Note also that $\chi^-(X_{\mathbb{C}}, \mathbb{Z}(d-1-r)) = \chi^-(X_{\mathbb{C}}, \mathbb{Z}(r))$ because Poincarè duality is compatible with complex conjugation.

Proposition 6.9.
$$\chi(\gamma(r))/\chi(\beta(r)) = \Gamma_r(A')^{d/2}(2\pi i)^{\chi(X_{\mathbb{C}})((d-1)/2-r)}$$

Proof. This follows from Proposition 2.3.3 and Corollary 4.9

Proposition 6.10.
$$\chi(\gamma(d-r))/\chi(\gamma(r)) = \Gamma_r(A')^{d/2-r}(2\pi i)^{\chi(X_{\mathbb{C}})((d-1)/2-r)+\chi^{-}(X_{\mathbb{C}},\mathbb{Z}(r))}$$

Proof; This follows immediately from Propositions 6.8, 6.9, and Corollary 6.7

Theorem 6.11.
$$\chi_B(X, d-r)/\chi_B(X, r) = \Gamma_r(A')^{d/2-r} (2\pi i)^{\chi(X_{\mathbb{C}})(r-(d-1)/2)+\chi^-(X_{\mathbb{C}}, \mathbb{Z}(r))}$$

Proof.
$$\chi_B(X, d-r)/\chi_B(X, r) = \chi(\gamma(d-r))/\chi(\gamma(r))$$

Proposition 6.2 and Theorem 6.11 immediately imply the compatibility of our conjecture with the functional equation, if we replace A by A'.

Theorem 6.12. If $d \leq 2$, our conjecture is compatible with the functional equation.

This follows from Theorem 6.11 and the main theorem of [KS].

§7. The case of number rings

Let F be a number field with ring of integers O_F , class number h, number of roots of unity w, and discriminant d_F . Let $X = \text{Spec } O_F$. We will explain how our conjecture for

X and r reduces to standard theorems if r = 0 or r = 1 and well-known conjectures if r < 0 or r > 1.

We begin with r = 0.

We know that $H^{j}_{et}(X, \mathbb{Z}(1)) = 0$ if j < 1, $H^{1}_{et}(X, \mathbb{Z}(1)) = O_{F}^{*}$, $H^{2}_{et}(X, \mathbb{Z}(1)) = Pic(X)$, $H^{3}(X, \mathbb{Z}(1)) = 0$ (up to 2-torsion) and $H^{0}_{et}(X, \mathbb{Z}(0)) = \mathbb{Z}$.

It immediately follows from the definitions that $H_W^0(X, \mathbb{Z}(0)) = \mathbb{Z}$, $H_W^1(X, \mathbb{Z}(0)) = 0$, and $H^3(X, \mathbb{Z}(0)) = \mu_F^{\vee}$, the dual of the roots of unity in F.

We also have the exact sequence $0 \to Pic(X)^{\vee} \to H^2_W(X,\mathbb{Z}(0)) \to Hom(O_F^*,\mathbb{Z}) \to 0$.

We also have $t_{j,0} = 0$ for all j. It follows that A(j,0) is always equal to 0.

We also see that A'(j,0) = 0 unless j = 1, when A(1,0) reduces to $(\mu_F)^{\vee}$ in degree 2, so $\chi(A'(1,0)) = w$.

C(0) becomes $0 \to \mathbb{C} \to \mathbb{C}^{r_1+r_2} \to Hom(O_F^*, \mathbb{C}) \to 0$, with the integral structure on the last term being $H^2_W(X, \mathbb{Z}(2))$ and the second map being the dual of the classical regulator. So $\chi(C(0)) = hR$, and $\chi(X, 0) = hR/w$. As is well-known, $\zeta^*(X, 0) = -hR/w$.

We now consider the case when r = 1.

A(j,1) is easily seen to be zero for j < -1.

The complex part of $\mathbb{C}(1)$ is given by $0 \to O_F^* \otimes \mathbb{C} \to \mathbb{C}^{r_1+r_2} \to \mathbb{C} \to 0$, with the last two terms getting the standard bases and the first term a basis coming from O_F^* , so $det(\mathbb{C}(1))$ is the classical regulator R. The Euler characteristic of $\mathbb{C}(1)_{tor}$ is

$$|H_W^!(X,\mathbb{Z}(1))_{tor}|/|H^2(X,\mathbb{Z}(1))_{tor}| = w/h$$

. It follows that the Euler characteristic of $\mathbb{C}(1)$ is hR/w.

Since
$$t_{j,1} = 0$$
 for $j \ge 0$, and $H_W^j(X, \mathbb{Z}(1)) = 0$ for $j \ge 5$, $A'(j,1) = 0$ for $j \ge 3$.
Finally, $B(j,1) = 0$ if $j \ne 0$, and $B(0,1)$ is given by

$$(7.1) 0 \to \mathbb{C}^{r_2} \to O_F \otimes \mathbb{C} \to \mathbb{C}^{r_1 + r_2} \to 0$$

.

Note that we have the usual map θ mapping $O_F \otimes \mathbb{C}$ to $\mathbb{C}^{r_1+2r_2}$ by sending $x \otimes 1$ to the collection of $\sigma(x)$ as σ runs through the embeddings of F in \mathbb{C} . The map from \mathbb{C}^{r_2} to $O_F \otimes \mathbb{C}$ is given by the natural inclusion of \mathbb{C}^{r_2} in $\mathbb{C}^{r_1+2r_2}$ multiplied by $2\pi i$, followed by the inverse of θ . Since the determinant of θ with respect to the usual bases is \sqrt{d}_F , we see that

the determinant of (7,1) is equal to $(2\pi i)^{r_2}/\sqrt{d_F}$. Hence the Euler characteristic $\chi(X,1)$ is equal to $hR(2\pi i)^{r_2}/w\sqrt{d_F}$ which is equal to the usual formula $hR(2\pi)^{r_2}2^{r_1}/w\sqrt{|d_F|}$ for $\zeta^*(X,1)$ up to a power of 2.

Now let r < 0.

The only non-zero groups $H_W^k(X,\mathbb{Z}(r))$ occur when k=2 or k=3, so A(j,r) is equal to zero for all j in the appropriate range, C(r)=0, and A'(j,r)=0 unless j=0 or j=1.

B(j,r)=0 unless j=0. B(0,r) reduces to the isomorphism $Ker(b_{0,r})\to H^2_W(X,\mathbb{Z}(r))^+$ so we give $Ker(b_{0,r})$ the integral structure induced from $H^2_W(X,\mathbb{Z}(r))^+$, and $\chi(B(0,r))=1$.

 $A'(0,r)_{\mathbb{C}}$ is given by $Ker(b_{0,r}) \to H^2_W(X,\mathbb{Z}(r))_{\mathbb{C}}$. $H^2_W(X,\mathbb{Z}(r))_{\mathbb{C}}$ is dual to $H^1_W(X,\mathbb{Z}(1-r))_{\mathbb{C}}$ which is canonically isomorphic to $K_{1-2r}(O_F)_{\mathbb{C}}$, and $Ker(b_{0,r})$ is canonically dual to $Coker(c_{0,1-r})$, where $c_{0,1-r}$ is the Beilinson regulator map. We conclude that det(A'(0,r)) is dual to the determinant D_{1-r} of the Beilinson regulator map with respect to the natural bases coming from singular cohomology and K-theory.

Then $\chi(X,r)$ is equal to $D_{1-r}|H_W^2(X,\mathbb{Z}(r))_{tor}|/|H_W^3(X,\mathbb{Z}(r))|$, which is equal to $D_{1-r}|H_W^2(X,\mathbb{Z}(1-r))|/|H_W^1(X,\mathbb{Z}(1-r))|$ which in turn is equal to $D_{1-r}|K_{-2r}(O_F)|/|K_{1-2r}(O_F)|$, up to 2-torsion.

This is essentially what was conjectured in [Li3] to be $\zeta^*(X,r)$

Finally, let r > 1. By definition, since j has to be between 2r + 1 and 2d - 1, there is no contribution from A'(j,r)

 $A(j,r)_{\mathbb{C}}$ is the complex $H_W^{j+1}(X,\mathbb{Z}(r))_{\mathbb{C}} \to Coker(\gamma_{j,r})$, which is only non-zero when j=0, in which case the map is the Beilinson regulator $c_{0,r}$, Since j has to be either 0 or 1, the torsion Euler characteristic is $|H^2(X,\mathbb{Z}(r))|/|H^1(X,\mathbb{Z}(r))_{tor}$ which up to 2-torsion is $K_{2r-2}(O_F)|/|K_{2r-1}(O_F)_{tor}|$. So letting R_r be the determinant of the Beilinson regulator map, the Euler characteristic $\chi(A,r) = (|K_{2r-2}(O_F)|/|K_{2r-1}(O_F)_{tor}|)R_r$, up to 2-torsion.

Now $B(j,r)_{\mathbb{C}}$ obviously is zero if $j \neq 0$. Let $s(r) = r_2$ If r is even and $s(r) = r_1 + r_2$ if r is odd. Then by the same arguments as in the case when r = 1 we have $det(B(0,r)_{\mathbb{C}} = (2\pi i)^{rs(r)} \sqrt{(d_F)}$.

The integral structure on $t_{j,r}$ is given by $\coprod_{0 \leq k < r} H^{j-k}(X, \lambda^k(\Omega))$ In the appendix, we compute that $\lambda^k(\Omega)$ is equal to $\Omega[k-1]$. So $t_{j,r} = \coprod_{k < r} H^{j-k}(X, \Omega[k-1])$. Since Ω only has cohomology in dimension 0, we see that $t_{j,r} = 0$ unless j = 1, and the order of $H^0(X, \Omega)$ is d_F so $\chi_{tor}(t_{j,r}) = d_F^{-r}$, and $\chi(B(j,r)) = (2\pi i)^{rs(r)} d_F^{(r-1/2)}$.

Putting everything together, we get that our conjecture says that $\zeta(X,r) = \chi(X,r)$,

where

$$\chi(X,r) = ((K_{2r-2}(O_F))/|K_{2r-1}(O_F)_{tor}|)((2\pi i)^{rs(r)}d_F^{r-1/2}R_r$$

up to 2-torsion, which is compatible with our conjecture for s = 1 - r via the functional equation, which of course here is a well-known theorem.

Appendix: Derived Exterior Powers

Let \mathcal{A} be an abelian category. Let $S\mathcal{A}$ denote the category of simplicial objects of \mathcal{A} and $C\mathcal{A}$ denote the category of homological chain complexes of objects of \mathcal{A} ending in degree zero. There are well-known functors $N: S\mathcal{A} \to C\mathcal{A}$ and $K: C\mathcal{A} \to S\mathcal{A}$ such that NK is the identity and KN is naturally equivalent to the identity functor. N and K also preserve homotopies. Let Λ^k denote k-th exterior power. Let X be a scheme and \mathcal{A} be the category of coherent locally free sheaves on X. If Q, is in $S\mathcal{A}$ with Q_n a locally free sheaf on X for all n, we define $\Lambda^k Q$, to be $\Lambda^k(Q_n)$ in $S\mathcal{A}$.

Proposition A.1.

Let X be a regular scheme projective over Spec \mathbb{Z} . Write X as a closed subscheme of a projective space $P = (P^n)_{\mathbb{Z}}$ such that I is the sheaf of ideals defining X. Then the complex of locally free sheaves $C_{X,P} = I/I^2 \to \Omega_{P/\mathbb{Z}}$ defines an element in the derived category of locally free sheaves on X which is independent of the choice of embedding of X into P.

Proof. If we have two different embeddings of X in P_1 and P_2 , take the Segré embedding of $P_1 \times P_2$ in P_3 , and compare successively the complexes defined by the embeddings into P_1 and P_2 with the product embedding into P_3 . (For details, see [LS]).

Definition A.2. $\lambda^k \Omega_{X/\mathbb{Z}} = N \Lambda^k K C_{X,P}$.

We see easily that this is independent of the choice of embedding.

We begin by recalling the following fact ([H], Exercise 5.16 (d)):

Lemma A.3. Let (X, O_X) be a ringed space, and let $0 \to F' \to F \to F'' \to 0$ be an exact sequence of locally free sheaves of O_X -modules. Then there exists a finite filtration of $\Lambda^r F$:

$$\Lambda^r F = G^0 \supseteq G^1 \supseteq \cdots \supseteq G^r \supseteq G^{r+1} = 0$$

with quotients $G^p/G^{p+1} = \Lambda^p F' \otimes \Lambda^{r-p} F''$.

Theorem A.4. Let (X, O_X) be a scheme such that every coherent sheaf on X has a finite resolution by locally free sheaves. Let $0 \to F' \to F \to F'' \to 0$ be an exact sequence of coherent sheaves on X. Let r be a positive integer, Then there exist objects in the derived category $G^0, G^1, \ldots G^{r+1}$ and maps from G^p to G^{p+1} such that we have $G^0 = \lambda^r(F), G^{r+1} = 0$, and exact triangles $G^p \to G^{p+1} \to \lambda^p(F') \otimes^L \lambda^{r-p}(F'') \to G^p[1]$.

Proof, This is an easy corollary of the definition of derived exterior power and the previous lemma.

Theorem A.5. Let A be a ring and M an A-module. a) $\lambda^0 M$ is canonically homotopic to the complex consisting only of A in degree 0.

- b) $\lambda^1 M$ is canonically homotopic to the complex consisting only of M in degree 0.
- c) If M has projective dimension r then $\lambda^k M$ is represented by a complex of length kr.

Proof: a) and b) are obvious and c) follows immediately from a theorem of Dold and Puppe ([DP].

Theorem A.6. Let $X = Spec\ O_F$, and $\Omega = \Omega_{O_F/\mathbb{Z}}$. Then $\lambda^k(\Omega)$ is isomorphic to $\Omega[k-1]$

Proof. Let D be the inverse different of O_F over \mathbb{Z} . We use induction on k, applying Theorem A.4 to the exact sequence $0 \to O_F \to D \to \Omega \to 0$, and using that $\lambda^k(O_F) = \lambda^k(D) = 0$ for k > 1.

References

- [Bl1] Bloch, S. Algebraic cycles and higher K-theory, Adv, Math 61 (1986). 267-304
- [Bl2] Bloch, S. Algebraic Cycles and the Beilinson Conjectures, Contemp. Math. Vol. 58 Part I, (1986), 65-78
- [Bl3] Bloch, S. De Rham cohomology and conductors of curves. Duke Math. J. 54~(1987) no. 2~295--308
- [BK] Bloch, S.; Kato, K. L-functions and Tamagawa numbers of motives. The Grothen-dieck Festschrift, Vol. I, 333-400, Progr. Math., 86, Birkhauser Boston, Boston, MA, 1990.
- [DP] Dold, A. and Puppe, D. Homologie nicht-aadditiven Funktoren. Anwendungen. Annales de l'Institut Fourier 11 (1961) 201-312
- [Fl] Flach, M. The equivariant Tamagawa number conjecture: a survey. With an appendix by C. Greither. Contemp. Math., 358, Stark's conjectures: recent work and new directions, 79-125, Amer. Math. Soc., Providence, RI, 2004.

- [FM] Flach, M. and Morin, B., Weil-étale cohomology and Zeta values of proper regular arithmetic schemes. Doc, Math 23 (2018) 1425-1560
- [FM2] Flach, M. and Morin, B. Compatibility of Special value conjectures with the functional equation of Zeta functions. arXiv 2005.04829
- [Fo] Fontaine, J-M. Valeurs spéciales des Fonctions-L des motifs, Sem. Bourbaki 34 (1991-1992), Exp. No. 751
- [FP] Fontaine, J.-M. and Perrin-Riou, B. Autour des conjectures de Bloch et Kato: co-homologie galoisienne et valeurs de fonctions L. Motives (Seattle, WA, 1991), 599706, Proc. Sympos. Pure Math., 55, Part 1, Amer. Math. Soc., Providence, RI, 1994.
- [FS] Flach, M. amd Siebel, D., Special values of the zeta-function of an algebraic surface. 2019, arxiv: 1909.07465
- [Ge] Geisser, T. Arithmetic cohomology over finite fields and special values of ζ -functions, Duke Math. Journal 133 (2006), 27-57
 - [H] Harsthorne, R. Algebraic Geometry, Springer-Verlag (1977)
- [KS] Kato, K., Saito, T., On the Conductor formula of Bloch. Publ. Math. Inst. Hautes Etudes Sci. 100 (2004) 5-151
- [Le] Levine, M. Techniques of localization in the theory of algebraic cycles, J. Alg.Geom. 10 (2001), 298-363
- [Li1] Lichtenbaum, S. The Weil-étale topology for number rings. Ann. of Math. (2) 170 (2009), no. 2, 657-683.
- [Li2] Lichtenbaum, S. Euler Characteristics and Special Values of Zeta-Functions, Proceedings of Fields Institute conference on algebraic cycles in honor of Spencer Bloch, Fields Institute communications, volume 56, (2009)
- [Li3] Lichtenbaum, S. Values of Zeta-Functions, Etale Cohomology, and Algebraic K-Theory. in Springer Lecture Notes in Mathematics (342) 1973
- [LR] Lichtenbaum, S. and Ramachandranm N. Values of zeta functions of algebraic surfaces at $s=1.\ 2021.\ arxiv:2101.06530$
- [LS] Lichtenbaum, S, and Schlessinger, M. The cotangent complex of a morphism. Trans. Amer. Math. Soc., 126 No. 1, (1967), 41-70
- [M] Morin, B. Zeta functions of regular arithmetic schemes at s=0. Duke Math. J. 87, 1263-1336. 98/91653'———
 - [R] Rapoport, M. Comparison of the regulators of Beilinson and of Borel, in Beilinson's

conjectures on special values of L-functions, (M. Rapoport, N. Schappacher, P.Schneider, eds.), Perspectives in Math. 4 Acad. Press 1989

RSS Rapoport, M, Schappacher, N., and Schneider, S., Beilinson's conjectures on special values of L-functions, Perspectives in Math. 4 Acad, Press 1989,

[Sa] Saito, T. Parity in Bloch's conductor formula in even dimensions. Journal de Théorie des Nombres de Bordeaux 16(2004) no, 2, 403-421

[Se1] Serre, J.-P., Zeta and L-functions, in Arithmetical Algebraic Geometry, Harper and Row (1965), 82-92

[Se2] Serre, J.-P., Facteurs locaux des fonctions zêta des variétiés algèbriques (définitions et conjectures) Séminaire Delange-Pisot-Poitou, (1969/70), no. 1

[[T] Tate, J., On the conjectures of Birch and Swinnerton-Dyer and a geometric analog Dix exposs sur la cohomologie des schmas, 189214, Adv. Stud. Pure Math., 3, North-Holland, Amsterdam, 1968.