

CANONICAL CORRELATION COEFFICIENTS OF HIGH-DIMENSIONAL GAUSSIAN VECTORS: FINITE RANK CASE

BY ZHIGANG BAO^{*}, JIANG HU[†], GUANGMING PAN[‡] AND WANG ZHOU[§]

*Hong Kong University of Science and Technology
Northeast Normal University
Nanyang Technological University
and National University of Singapore*

Consider a Gaussian vector $\mathbf{z} = (\mathbf{x}', \mathbf{y}')'$, consisting of two sub-vectors \mathbf{x} and \mathbf{y} with dimensions p and q respectively. With n independent observations of \mathbf{z} , we study the correlation between \mathbf{x} and \mathbf{y} , from the perspective of the Canonical Correlation Analysis. We investigate the high-dimensional case: both p and q are proportional to the sample size n . Denote by Σ_{uv} the population cross-covariance matrix of random vectors \mathbf{u} and \mathbf{v} , and denote by S_{uv} the sample counterpart. The canonical correlation coefficients between \mathbf{x} and \mathbf{y} are known as the square roots of the nonzero eigenvalues of the canonical correlation matrix $\Sigma_{xx}^{-1}\Sigma_{xy}\Sigma_{yy}^{-1}\Sigma_{yx}$. In this paper, we focus on the case that Σ_{xy} is of finite rank k , i.e. there are k nonzero canonical correlation coefficients, whose squares are denoted by $r_1 \geq \dots \geq r_k > 0$. We study the sample counterparts of $r_i, i = 1, \dots, k$, i.e. the largest k eigenvalues of the sample canonical correlation matrix $S_{xx}^{-1}S_{xy}S_{yy}^{-1}S_{yx}$, denoted by $\lambda_1 \geq \dots \geq \lambda_k$. We show that there exists a threshold $r_c \in (0, 1)$, such that for each $i \in \{1, \dots, k\}$, when $r_i \leq r_c$, λ_i converges almost surely to the right edge of the limiting spectral distribution of the sample canonical correlation matrix, denoted by d_+ . When $r_i > r_c$, λ_i possesses an almost sure limit in $(d_+, 1]$, from which we can recover r_i 's in turn, thus provide an estimate of the latter in the high-dimensional scenario. We also obtain the limiting distribution of λ_i 's under appropriate normalization. Specifically, λ_i possesses Gaussian type fluctuation if $r_i > r_c$, and follows Tracy-Widom distribution if $r_i < r_c$. Some applications of our results are also discussed.

1. Introduction. In multivariate analysis, the most general and favorable method to investigate the relationship between two random vectors \mathbf{x} and \mathbf{y} , is the Canonical Correlation Analysis (CCA), which was raised in the seminal work of Hotelling [27]. CCA aims at finding two sets of basis vectors, such that the correlations between the projections of the variables \mathbf{x} and \mathbf{y} onto these basis vectors are mutually maximized, namely, seeking vectors $\mathbf{a} = \mathbf{a}_1$ and $\mathbf{b} = \mathbf{b}_1$ to maximize the correlation coefficient

$$\rho \equiv \rho(\mathbf{a}, \mathbf{b}) := \frac{\text{Cov}(\mathbf{a}'\mathbf{x}, \mathbf{b}'\mathbf{y})}{\sqrt{\text{Var}(\mathbf{a}'\mathbf{x})} \cdot \sqrt{\text{Var}(\mathbf{b}'\mathbf{y})}}. \quad (1.1)$$

Conventionally, $\rho_1 := \rho(\mathbf{a}_1, \mathbf{b}_1)$ is called the *first canonical correlation coefficient*. Having obtained the first m canonical correlation coefficients $\rho_i, i = 1, \dots, m$ and the corresponding vector

^{*}Z.G. Bao was supported by a startup fund from HKUST

[†]J. Hu was partially supported by CNSF 11301063

[‡]G.M. Pan was partially supported by the Ministry of Education, Singapore, under grant # ARC 14/11

[§] W. Zhou was partially supported by the Ministry of Education, Singapore, under grant MOE2015-T2-2-039(R-155-000-171-112).

MSC 2010 subject classifications: 62H20, 60B20, 60F99

Keywords and phrases: Canonical correlation analysis, Random Matrices, MANOVA ensemble, High-dimensional data, finite rank perturbation, largest eigenvalues

pairs $(\mathbf{a}_i, \mathbf{b}_i), i = 1, \dots, m$, one can proceed to seek vectors $(\mathbf{a}_{m+1}, \mathbf{b}_{m+1})$ maximizing ρ subject to the constraint that $(\mathbf{a}'_{m+1}\mathbf{x}, \mathbf{b}'_{m+1}\mathbf{y})$ is uncorrelated with $(\mathbf{a}'_i\mathbf{x}, \mathbf{b}'_i\mathbf{y})$ for all $i = 1, \dots, m$. Analogously, we call ρ_i the *ith canonical correlation coefficient* if it is nonzero. Denoting by Σ_{uv} the population cross-covariance matrix of arbitrary two random vectors \mathbf{u} and \mathbf{v} , it is well known that $r_i := \rho_i^2$ is the *ith largest eigenvalue of the (population) canonical correlation matrix*

$$\Sigma_{xx}^{-1}\Sigma_{xy}\Sigma_{yy}^{-1}\Sigma_{yx}.$$

Let $\mathbf{z}_i = (\mathbf{x}'_i, \mathbf{y}'_i)', i = 1, \dots, n$ be n independent observations of the vector $\mathbf{z} := (\mathbf{x}', \mathbf{y}')' \sim N(\boldsymbol{\mu}, \Sigma)$ with mean vector $\boldsymbol{\mu}$ and covariance matrix

$$\Sigma = \begin{pmatrix} \Sigma_{xx} & \Sigma_{xy} \\ \Sigma_{yx} & \Sigma_{yy} \end{pmatrix},$$

We can study the canonical correlation coefficients via their sample counterparts. To be specific, we employ the notation S_{uv} to represent the sample cross-covariance matrix for arbitrary two random vectors \mathbf{u} and \mathbf{v} , where the implicit sample size of $(\mathbf{u}', \mathbf{v}')'$ is assumed to be n , henceforth. Then the square of the *ith sample canonical correlation coefficient* is defined as the *ith largest eigenvalue of the sample canonical correlation matrix* (CCA matrix in short)

$$S_{xx}^{-1}S_{xy}S_{yy}^{-1}S_{yx},$$

denoted by λ_i in the sequel.

Let p and q be the dimensions of the sub-vectors \mathbf{x} and \mathbf{y} , respectively. In the classical low-dimensional setting, i.e., both p and q are fixed but n is large, one can safely use λ_i to estimate r_i , considering the convergence of the sample cross-covariance matrices towards their population counterparts. However, nowadays, due to the increasing demand in the analysis of high-dimensional data springing up in various fields such as genomics, signal processing, microarray, finance and proteomics, putting forward a theory on high-dimensional CCA is much needed. So far, there are only a handful of works devoted to this topic. Fujikoshi in [20] derived the asymptotic distributions of the canonical correlation coefficients when q is fixed while p is proportional to n . Oda et al. in [39] considered the problem of testing for redundancy in high dimensional canonical correlation analysis. Recently, with certain sparsity assumption, the theoretical results and potential applications of high-dimensional sparse CCA have been discussed in [23, 22]. In the null case, i.e., \mathbf{x} and \mathbf{y} are independent, the Tracy-Widom law for the largest canonical correlation coefficients has been studied in [32, 26, 25], when p, q, n are proportional. Recently, in [34] Johnstone and Onatski derived the asymptotics of the likelihood ratio processes of CCA corresponding to the null hypothesis of no spikes and the alternative of a single spike.

In this paper, we will work with the following high-dimensional setting.

ASSUMPTION 1.1 (On the dimensions). *We assume that $p := p(n)$, $q := q(n)$, and*

$$p/n = c_1 \rightarrow y_1 \in (0, 1), \quad q/n = c_2 \rightarrow y_2 \in (0, 1), \quad \text{as } n \rightarrow \infty, \quad \text{s.t. } y_1 + y_2 \in (0, 1).$$

Without loss of generality, we always work with the additional assumption

$$p > q, \quad \text{thus } c_1 > c_2.$$

Observe that here y_1 and y_2 are asymptotic parameters, while c_1 and c_2 are non-asymptotic parameters. Hence, in general, a c_1, c_2 -dependent random variable $X(c_1, c_2)$ can not serve as a limiting target of a random sequence $X_n(c_1, c_2), n \geq 1$. Nevertheless, to ease the presentation, from time to time, we still write $X_n(c_1, c_2) \rightarrow X(c_1, c_2)$, if $X_n(c_1, c_2) - X(c_1, c_2) \rightarrow 0$, where the convergence could be in distribution, in probability or a.s., etc.

Let $\bar{\mathbf{x}}$ and $\bar{\mathbf{y}}$ be the sample means of n samples $\{\mathbf{x}_i\}_{i=1}^n$ and $\{\mathbf{y}_i\}_{i=1}^n$ respectively, and use the notation $\hat{\mathbf{x}}_i := \mathbf{x}_i - \bar{\mathbf{x}}$ and $\hat{\mathbf{y}}_i := \mathbf{y}_i - \bar{\mathbf{y}}$ for $i = 1, \dots, n$. We can then write

$$S_{ab} = \frac{1}{n-1} \sum_{i=1}^n \hat{\mathbf{a}}_i \hat{\mathbf{b}}_i', \quad \mathbf{a}, \mathbf{b} = \mathbf{x} \text{ or } \mathbf{y}$$

It is well known that there exist $n-1$ i.i.d. Gaussian vectors $\tilde{\mathbf{z}}_i = (\tilde{\mathbf{x}}_i', \tilde{\mathbf{y}}_i')' \sim N(\mathbf{0}, \Sigma)$, such that

$$S_{ab} = \frac{1}{n-1} \sum_{i=1}^{n-1} \tilde{\mathbf{a}}_i \tilde{\mathbf{b}}_i', \quad \mathbf{a}, \mathbf{b} = \mathbf{x} \text{ or } \mathbf{y}.$$

For simplicity, we recycle the notation \mathbf{x}_i and \mathbf{y}_i to replace $\tilde{\mathbf{x}}_i$ and $\tilde{\mathbf{y}}_i$, and work with n instead of $n-1$, noticing that such a replacement on sample size is harmless to Assumption 1.1. Hence, we can and do assume that \mathbf{z} is centered in the sequel and denote

$$S_{ab} = \frac{1}{n} \sum_{i=1}^n \mathbf{a}_i \mathbf{b}_i', \quad \mathbf{a}, \mathbf{b} = \mathbf{x} \text{ or } \mathbf{y}.$$

Furthermore, for brevity, we introduce the notation

$$C_{xy} := S_{xx}^{-1} S_{xy} S_{yy}^{-1} S_{yx}, \quad (1.2)$$

for the CCA matrix, and C_{yx} can be analogously defined via switching the roles of x and y in (1.2). Notice that C_{xy} and C_{yx} possess the same non-zero eigenvalues.

By our assumption $p > q$, there are at most q non-zero canonical correlations, either population ones or sample ones. An elementary fact is that $\lambda_i, r_i \in [0, 1]$ for all $i = 1, \dots, q$. Note that $\lambda_i, i = 1, \dots, q$ are also eigenvalues of the $q \times q$ matrix C_{yx} , whose empirical spectral distribution (ESD) will be denoted by

$$F_n(x) := \frac{1}{q} \sum_{i=1}^q \mathbf{1}_{\{\lambda_i \leq x\}}. \quad (1.3)$$

We use the following notations for the data matrices

$$\mathcal{X} := \mathcal{X}_n = (\mathbf{x}_1, \dots, \mathbf{x}_n), \quad \mathcal{Y} := \mathcal{Y}_n = (\mathbf{y}_1, \dots, \mathbf{y}_n).$$

Let $\text{vec}(\mathcal{X}) = (\mathbf{x}_1', \dots, \mathbf{x}_n')'$ be the vectorization of the matrix \mathcal{X} , and define $\text{vec}(\mathcal{Y})$ be the analogously. We see that $\text{vec}(\mathcal{X}) \sim N(\mathbf{0}, I_n \otimes \Sigma_{xx})$ and $\text{vec}(\mathcal{Y}) \sim N(\mathbf{0}, I_n \otimes \Sigma_{yy})$. Our aim, in this work, is to study the asymptotic behavior of a few largest sample canonical correlation coefficients $\sqrt{\lambda_i}$'s, and try to get the information about the population ones $\rho_i = \sqrt{r_i}$ from the sample ones. We will focus on the case of finite rank, i.e., there is some fixed nonnegative integer k , such that

$$r_1 \geq \dots \geq r_k \geq r_{k+1} = \dots = r_q = 0.$$

Specifically, we make the following assumption throughout the work.

ASSUMPTION 1.2 (On the rank of the population matrix). *We assume that $\text{rank}(\Sigma_{xy}) \leq k$ for some fixed positive integer k . Furthermore, setting $r_0 = 1$, we denote by k_0 the nonnegative integer satisfying*

$$1 = r_0 \geq \dots \geq r_{k_0} > r_c \geq r_{k_0+1} \geq \dots r_k \geq r_{k+1} = 0, \quad (1.4)$$

where

$$r_c \equiv r_c(c_1, c_2) := \sqrt{\frac{c_1 c_2}{(1 - c_1)(1 - c_2)}}. \quad (1.5)$$

In Section 1.2 we will state our main results. Before that, we introduce in Section 1.1 some known results in the *null case*, i.e. $k = 0$, which will be the starting point of our discussion.

1.1. *The null case: MANOVA ensemble.* At first, we introduce some known results on the limiting behavior of $\{\lambda_i\}_{i=1}^q$ in the null case, i.e. \mathbf{x} and \mathbf{y} are independent, or else, $r_i = 0$ for all $i = 1, \dots, q$. It is elementary to see that the canonical correlation coefficients are invariant under the block diagonal transformation $(\mathbf{x}_i, \mathbf{y}_i) \rightarrow (\mathbf{A}\mathbf{x}_i, \mathbf{B}\mathbf{y}_i)$, for any $p \times p$ matrix \mathbf{A} and $q \times q$ matrix \mathbf{B} , as long as both of them are nonsingular. Hence, without loss of generality, in this section, we tentatively assume that $\Sigma_{xx} = I_p$ and $\Sigma_{yy} = I_q$. Under our high-dimensional setting, i.e. Assumption 1.1, it is known that λ_i 's do not converge to 0 even in the null case, instead, they typically spread out over an interval contained in $[0, 1]$. Specifically, we have the following theorem on $F_n(x)$ (c.f. (1.3)), which is essentially due to Wachter [45].

THEOREM 1.3. *When \mathbf{x} and \mathbf{y} are independent Gaussian and Assumption 1.1 holds, almost surely, F_n converges weakly to a deterministic probability distribution $F(x)$ with density*

$$f(x) = \frac{1}{2\pi c_2} \frac{\sqrt{(d_+ - x)(x - d_-)}}{x(1 - x)} \mathbf{1}(d_- \leq x \leq d_+), \quad (1.6)$$

where

$$d_{\pm} = \left(\sqrt{c_1(1 - c_2)} \pm \sqrt{c_2(1 - c_1)} \right)^2 \quad (1.7)$$

REMARK 1.4. *In the null case, the convergence of the ESD of the CCA matrix actually holds under a more general distribution assumption, see [48].*

Conventionally, we call $F(x)$ in Theorem 1.3 the limiting spectral distribution (LSD) of C_{yx} . One might note that $F(x)$ is also the LSD of the so-called MANOVA ensemble with appropriately chosen parameters, which is widely studied in the Random Matrix Theory (RMT). Actually, when \mathbf{x} and \mathbf{y} are Gaussian and independent, the CCA matrix C_{yx} is exactly a MANOVA matrix. To see this, we note that $P_x := \mathcal{X}'(\mathcal{X}\mathcal{X}')^{-1}\mathcal{X}$ is a projection matrix independent of Y . Hence, we can write

$$C_{yx} = (\mathcal{Y}(I - P_x)\mathcal{Y}' + \mathcal{Y}P_x\mathcal{Y}')^{-1}\mathcal{Y}P_x\mathcal{Y}'.$$

Using Cochran's theorem, we see that $\mathcal{Y}(I - P_x)\mathcal{Y}'$ and $\mathcal{Y}P_x\mathcal{Y}'$ are independent Wishart,

$$\mathcal{Y}(I - P_x)\mathcal{Y}' \sim \mathcal{W}_q(I_q, n - p), \quad \mathcal{Y}P_x\mathcal{Y}' \sim \mathcal{W}_q(I_q, p).$$

Hereafter, we use the notation $\mathcal{W}_m(\cdot, \cdot)$ to denote the Wishart distribution $\text{Wishart}_m(\cdot, \cdot)$ for short. Consequently, $\lambda_i, i = 1, \dots, q$ are known to possess the following joint density function,

$$p_n(\lambda_1, \dots, \lambda_q) = C_n \prod_{i < j} |\lambda_i - \lambda_j| \prod_{i=1}^q (1 - \lambda_i)^{(n-p-q-1)/2} \lambda_i^{(p-q-1)/2} \mathbf{1}(\lambda_i \in [0, 1]),$$

where C_n is the normalizing constant, see Muirhead [38], page 112, for instance. Or else, one can refer to [32], for more related discussions. In the context of RMT, the point process possessing the above joint density is also called Jacobi ensemble.

Throughout the paper, we will say that an n -dependent event $A \equiv A(n)$ holds with *overwhelming probability*, if for any given positive number ℓ , there exists a constant C_ℓ such that

$$\mathbb{P}(A) \geq 1 - C_\ell n^{-\ell}.$$

Especially, for any fixed integer $K \geq 0$, we have $\cap_{i=1}^{n^K} A_i$ holds with overwhelming probability if A_i holds with overwhelming probability individually with the common C_ℓ 's. The next known result concerns the convergence of the largest eigenvalues.

THEOREM 1.5. *When \mathbf{x} and \mathbf{y} are independent and Assumption 1.1 holds, we have*

$$\lambda_i - d_+ \xrightarrow{a.s.} 0, \quad (1.8)$$

for any fixed positive integer i . Moreover, for any small constant $\varepsilon > 0$,

$$\lambda_1 \leq d_+ + \varepsilon \quad (1.9)$$

holds with overwhelming probability.

REMARK 1.6. *The estimate (1.9), is actually implied by some existing results in the literature directly. For example, one can refer to the small deviation estimate of the largest eigenvalue of the Jacobi ensemble in [36]. Moreover, (1.8) is a direct consequence of (1.9) and Theorem 1.3.*

REMARK 1.7. *As we mentioned in Introduction, on the fluctuation level, λ_1 possesses a Type 1 Tracy-Widom limit after appropriate normalization. Such a result has been established recently in [32, 26, 25].*

1.2. *Finite rank case.* We now turn to the case we are interested in: the finite rank case. To wit, Assumption 1.2 holds. It will be clear that the CCA matrix in such a finite rank case can be viewed as a finite rank perturbation of that in the null case. Consequently, the global behavior, especially the LSD, turns out to coincide with the null case. However, finite rank perturbation may significantly alter the behavior of the extreme eigenvalues, when the perturbation is strong enough. Similar problems have been studied widely for various random matrix models, not trying to be comprehensive, we refer to the spiked sample covariance matrices [31, 6, 41, 4, 5, 19], the deformed Wigner matrices [12, 42, 18, 13, 37], the deformed unitarily invariant matrices [9, 35], and some other deformed models [11, 10, 46]. In this work, for our CCA matrix C_{xy} , we study the limits and the fluctuations of its largest eigenvalues, i.e. squares of the largest sample canonical correlation coefficients, under Assumption 1.2. Our main results are the following three theorems. Let

$$\gamma_i := r_i(1 - c_1 + c_1 r_i^{-1})(1 - c_2 + c_2 r_i^{-1}). \quad (1.10)$$

Recall r_c and d_+ defined in (1.5) and (1.7), respectively. It is easy to check that $\gamma_i \geq d_+$ if $r_i \geq r_c$.

THEOREM 1.8 (Limits). *Under Assumptions 1.1 and 1.2, the squares of the largest canonical correlation coefficients exhibit the following convergence as $n \rightarrow \infty$.*

(i): (Outliers) *For $1 \leq i \leq k_0$, we have*

$$\lambda_i - \gamma_i \xrightarrow{a.s.} 0.$$

(ii): (Sticking eigenvalues) *For each fixed $i \geq k_0 + 1$, we have*

$$\lambda_i - d_+ \xrightarrow{a.s.} 0.$$

The next two results are on fluctuations of λ_i 's. We need the following definition.

DEFINITION 1.9. *For two (possibly) n -dependent numbers $a(n), b(n) \in \mathbb{C}$, we say $a(n)$ is well separated from $b(n)$, if there exists a small positive constant ε such that $|a(n) - b(n)| \geq \varepsilon$ for sufficiently large n .*

The next theorem is on the fluctuations of the outliers.

THEOREM 1.10 (Fluctuations of the outliers). *Suppose that Assumptions 1.1 and 1.2 hold. Let l_0 be the cardinality of the set $\Gamma := \{r_1, \dots, r_{k_0}\}$ (not counting multiplicity), and denote by $r_1 = r_{(1)} > \dots > r_{(l_0)} = r_{k_0}$ the l_0 different values in Γ . Set $n_0 = 0$ and denote by n_l the multiplicity of $r_{(l)}$ in Γ for $1 \leq l \leq l_0$. Let $J_l = [\sum_{i=0}^{l-1} n_i + 1, \sum_{i=0}^l n_i] \cap \mathbb{Z}$ for $1 \leq l \leq l_0$. If $r_{(\ell)}$ is well separated from $r_c, 1, r_{(\ell-1)}$ and $r_{(\ell+1)}$, the n_l -dimensional random vector*

$$\left\{ \frac{\sqrt{n}(\lambda_j - \gamma_j)}{\xi(r_j)}, j \in J_l \right\}$$

converges weakly to the distribution of the ordered eigenvalues of an n_l -dimensional symmetric Gaussian random matrix $G = (g_{ij})$ with independent (up to symmetry) entries $g_{ij} \sim N(0, 1 + \delta_{ij})$. Here

$$\xi^2(r_j) := \frac{(1 - r_j)^2 \left(2(1 - c_1)(1 - c_2)r_j + c_1 + c_2 - 2c_1c_2 \right) \left((1 - c_1)(1 - c_2)r_j^2 - c_1c_2 \right)}{r_j^2}. \quad (1.11)$$

REMARK 1.11. *The assumption that r_ℓ is away from $r_c, 1, r_{(\ell-1)}$ and $r_{(\ell+1)}$ by a constant distance $\varepsilon > 0$ (well separated) is not necessary. It is possible to reduce ε to some n -dependent distance $n^{-\alpha}$ for sufficiently small α . But we do not pursue this direction here.*

Our last theorem is on the fluctuation of the sticking eigenvalues.

THEOREM 1.12 (Fluctuations of the sticking eigenvalues). *Suppose that Assumptions 1.1 and 1.2 hold. In addition, we assume that r_c is well separated from r_{k_0} and r_{k_0+1} . There exists a CCA matrix in the null case with the same parameters p, q, n , whose nonzero eigenvalues are denoted by $\lambda_1 > \lambda_2 > \dots > \lambda_q$, such that for any fixed positive integer m and any small constant $\varepsilon > 0$, we have*

$$\max_{1 \leq i \leq m} |\lambda_{k_0+i} - \lambda_i| \leq n^{-1+\varepsilon} \quad (1.12)$$

in probability. This implies

$$\frac{n^{\frac{2}{3}}(\lambda_{k_0+1} - d_+)}{\xi_{tw}} \implies F_1, \quad (1.13)$$

where F_1 is the Type 1 Tracy-Widom distribution and

$$\xi_{tw}^3 = \frac{d_+^2(1 - d_+)^2}{\sqrt{c_1c_2(1 - c_1)(1 - c_2)}}.$$

REMARK 1.13. *From (1.12), it is easy to conclude (1.13), by using the Tracy-Widom limit of λ_1 derived in [32, 26, 25]. Observe that in these references, the Tracy-Widom law is stated for the logit transform of λ_1 , i.e., $\log(\lambda_1/(1 - \lambda_1))$. Using a Taylor expansion, it is elementary to check that $n^{\frac{2}{3}}(\lambda_1 - d_+)/\xi_{tw} \implies F_1$ from the Tracy-Widom law for the logit transform of λ_1 (see Theorem 1 and Section 2.1.1 in [32] for more details).*

REMARK 1.14. *Similarly to Remark 1.11, the assumption that r_c is away from r_{k_0} and r_{k_0+1} by a constant distance can be weakened. But we do not pursue this direction here.*

To illustrate the result in Theorems 1.8, 1.10 and 1.12, we did some numerical simulations. The different limiting behavior of λ_i in (i) and (ii) of Theorem 1.8 can be observed in Figure 1. The fluctuation for λ_i in Theorem 1.10 and Theorem 1.12 can be seen from Figures 2 and 1.




FIG 1. We chose a Gaussian vector $\mathbf{z} = (\mathbf{x}', \mathbf{y}')'$ with $p = 500$ and $q = 1000$. The sample size is $n = 5000$. Hence, $c_1 = 0.1$, $c_2 = 0.2$. Then $r_c \approx 0.17$ and $d_+ = 0.5$. We chose $k = 4$ and $(r_1, r_2, r_3, r_4) = (0.5, 0.4, 0.3, 0.16)$. Then $\gamma_1 \approx 0.66$, $\gamma_2 \approx 0.6$, $\gamma_3 \approx 0.54$ and $\gamma_4 \approx 0.50$. The above simulation result is based on 1000 replications. The abscises of the vertical segments represent d_+ and $\{\gamma_1, \gamma_2, \gamma_3\}$. The four histograms represent the distributions of the largest four eigenvalues of the CCA matrix respectively.

1.3. Organization and notations. Our paper is organized as follows. In Section 2, we display some applications of our results. We introduce in Section 3 some necessary preliminaries. In Section 4, we will reformulate the sample canonical correlation matrix in the finite rank case as a perturbation of that in the null case, thereby obtaining a determinant equation for the largest eigenvalues. By solving the limiting counterpart of the determinant equation, we can get the limits of the largest eigenvalues in Section 5, i.e. prove Theorem 1.8. In Section 6 we derive the fluctuations of the outliers (Theorem 1.10), and in Section 7, we study the fluctuations of the sticking eigenvalues (Theorem 1.12). Sections 8 and 9 are then devoted to proving some technical lemmas used in previous sections.

Throughout the paper, the notation C represents some generic constants whose values which may vary from line to line. The notation $\mathbf{0}_{k \times \ell}$ is used to denote the k by ℓ null matrix, which will be abbreviated to $\mathbf{0}_k$ if $k = \ell$, and from time to time we also use the abbreviation $\mathbf{0}$ if the dimension $k \times \ell$ is clear from the context. For any matrix \mathbf{A} , its (i, j) th entry will be written as \mathbf{A}_{ij} . When \mathbf{A} is square, we denote by $\text{Spec}(\mathbf{A})$ its spectrum. For a function $f : \mathbb{C} \rightarrow \mathbb{C}$ and an Hermitian matrix \mathbf{A} with spectral decomposition $\mathbf{U}_\mathbf{A} \boldsymbol{\Lambda}_\mathbf{A} \mathbf{U}_\mathbf{A}^*$, we define $f(\mathbf{A})$ as usual, in the sense of functional calculus, to wit, $f(\mathbf{A}) = \mathbf{U}_\mathbf{A} f(\boldsymbol{\Lambda}_\mathbf{A}) \mathbf{U}_\mathbf{A}^*$, where $f(\boldsymbol{\Lambda}_\mathbf{A})$ is the diagonal matrix obtained via mapping the eigenvalues of \mathbf{A} to their images under f . We will conventionally use the notations $\|\mathbf{A}\|$ and $\|\mathbf{A}\|_{\text{HS}}$ to represent the operator norm and Hilbert-Schmidt norm of a matrix \mathbf{A} , respectively, while $\|\mathbf{b}\|$ stands for the Euclidean norm of a vector \mathbf{b} . Throughout this paper, we use $o_p(1)$ to denote a scalar negligible (in probability) or a fixed-dimensional random matrix with negligible (in probability) entries. And the notations $o_{\text{a.s.}}(1)$, $O_p(1)$ and $O_{\text{a.s.}}(1)$ are used in a similar way.

For brevity, we further introduce the notations

$$\varpi := c_1 c_2, \quad \vartheta := (1 - c_1)(1 - c_2). \quad (1.14)$$

2. Applications. In this section, we discuss three applications of our results Theorems 1.8, 1.10 and 1.12 in hypothesis testing, estimation of the number and the values of the canonical correlation coefficients (CCC for short). At the end, we present an experiment on a real rain forest data.




FIG 2. We chose a Gaussian vector $\mathbf{z} = (\mathbf{x}', \mathbf{y}')'$ with $p = 500$ and $q = 1000$. The sample size is $n = 5000$. Hence, $c_1 = 0.1$, $c_2 = 0.2$. Then $r_c \approx 0.17$ and $d_+ = 0.5$. We chose $k = 4$ and $(r_1, r_2, r_3, r_4) = (0.5, 0.4, 0.4, 0.16)$. Then $\gamma_1 \approx 0.66$ and $\gamma_2 = \gamma_3 \approx 0.6$. This result is based on 1000 replications. The upper panel shows that the frequency histogram of λ_1 compared to its Gaussian limits (dashed lines). The lower panels show that the contour plots of empirical joint density function of λ_2 and λ_3 (left plot, after centralisation and scaling) and contour plots of their limits (right plot). The empirical joint density function is displayed by using the two-dimensional kernel density estimates.

2.1. *Application 1: Power of testing for the independence of two high dimensional normal vectors.* Testing for the independence of random vectors is a very classical problem. For two normal random vectors \mathbf{x} and \mathbf{y} with dimensions p and q respectively, we consider the test

$$H_0 : \Sigma_{xy} = \mathbf{0}_{p \times q} \quad \text{v.s.} \quad H_1 : \text{not } H_0.$$

Currently, for high dimensional cases, there are three kinds of widely discussed test procedures: (i) Corrected likelihood ratio tests (see [29, 30]); (ii) Trace tests (see [29, 49, 28, 47, 8]); (iii) Largest eigenvalue tests (see [32, 33, 26]). It has been shown by numerical results in [8, 26] that if $\Sigma_{\mathbf{xy}}$ is sparse, the corrected likelihood ratio tests and trace tests fail and the largest eigenvalue tests works well. In the following, we propose a statistic based on the CCC of \mathbf{x} and \mathbf{y} , and show that it is powerful against finite rank case. It is well known that testing for the independence of two high dimensional normal vectors equals to testing their first canonical correlation coefficient being zero or not, i.e.,

$$H_0 : r_1 = 0 \quad \text{v.s.} \quad H_1 : r_1 > 0.$$

Therefore, a natural test statistic is the first eigenvalue λ_1 of the sample canonical correlation matrix C_{xy} . Then according to Section 2.1.1.in [32] and Theorem 2.1 in [26], under the null hypothesis and Assumption 1.1 we have

$$\frac{n^{2/3}(\lambda_1 - d_+)}{\xi_{tw}} \Rightarrow F_1. \quad (2.1)$$

Therefore, we reject H_0 if

$$\lambda_1 > n^{-2/3} q_\alpha \xi_{tw} + d_+, \quad (2.2)$$

where q_α is the $1 - \alpha$ quantile of Tracy-Widom distribution F_1 . If the sample is under the Assumption 1.2 and $r_1 > r_c$, then this test will be able to detect the alternative hypothesis with a power tending to one as the dimension tends to infinity.

THEOREM 2.1 (Power function). *Suppose that the assumptions in Theorems 1.10 and 1.12 hold with $r_1 > r_c$. Then as $n \rightarrow \infty$, the power function of the test procedure (2.2)*

$$\text{Power} = \mathbb{P} \left(\frac{\sqrt{n}(\lambda_1 - \gamma_1)}{\xi(r_1)} > \frac{n^{-1/6} q_\alpha \xi_{tw}}{\xi(r_1)} + \frac{\sqrt{n}(d_+ - \gamma_1)}{\xi(r_1)} \right) \rightarrow 1.$$

PROOF. Under the conditions in Theorems 1.10 and 1.12, for any α , we have $\frac{n^{-1/6} q_\alpha \xi_{tw}}{\xi(r_1)} \rightarrow 0$. From the conclusion of Theorem 1.8 and $r_1 > r_c$, we have $\lim_{n \rightarrow \infty} (d_+ - \gamma_1) < 0$, which implies $\lim_{n \rightarrow \infty} \frac{\sqrt{n}(d_+ - \gamma_1)}{\xi(r_1)} \rightarrow -\infty$. In addition, by applying Theorem 1.10, $\frac{\sqrt{n}(\lambda_1 - \gamma_1)}{\xi(r_1)}$ converges weakly to the distribution of the largest eigenvalue of a symmetric Gaussian random matrix. Therefore, combining the above results, we then complete the proof of Theorem 2.1. \square

2.2. *Application 2: Estimating the number of the population CCC.* As a fundamental problems in CCA, the estimation of the number and the values of the population CCC are widely investigated. In this subsection, we first apply our results to determine the number of the outliers of high dimensional population CCC (not counting multiplicity). Actually, due to the threshold in our results, we cannot detect the population CCC which are smaller than $\sqrt{r_c}$. To the best of our knowledge, there is no effective method in general to successfully detect the population spikes below the threshold when the dimension is large, even for the simpler spike PCA problem. The reason can be find in the recent work [3] for instance.

We propose our estimator of the number of the outliers of population CCC k_0 by the number of eigenvalues of the sample canonical correlation matrix which are larger than d_+ :

$$\hat{k}_0 := \max\{i : \lambda_i \geq d_+ + \epsilon_n\}, \quad (2.3)$$

where ϵ_n is a sequence of positive number only depending on n and satisfying $\epsilon_n \sqrt{n} \rightarrow 0$ and $\epsilon_n n^{2/3} \rightarrow \infty$. Then the estimator \hat{k}_0 is weakly consistent according to the following theorem.

THEOREM 2.2 (Weak consistency of \hat{k}_0). *Suppose that the assumptions in Theorems 1.10 and 1.12 hold. As $n \rightarrow \infty$, the estimator \hat{k}_0 in (2.3) is weakly consistent for the number of the outliers of population CCC k_0 , that is*

$$\mathbb{P}(\hat{k}_0 = k_0) \rightarrow 1. \quad (2.4)$$

PROOF. According to the definition of \hat{k}_0 in (2.3), we have that

$$\begin{aligned} \mathbb{P}(\hat{k}_0 = k_0) &= \mathbb{P}(\lambda_{k_0} \geq d_+ + \epsilon_n, \lambda_{k_0+1} < d_+ + \epsilon_n) \\ &\geq 1 - \mathbb{P}(\lambda_{k_0} < d_+ + \epsilon_n) - \mathbb{P}(\lambda_{k_0+1} \geq d_+ + \epsilon_n). \end{aligned} \quad (2.5)$$

By Theorem 1.10 and $\epsilon_n \sqrt{n} \rightarrow 0$, we obtain that

$$\mathbb{P}(\lambda_{k_0} < d_+ + \epsilon_n) = \mathbb{P}(\sqrt{n}(\lambda_{k_0} - \gamma_{k_0}) < \sqrt{n}(d_+ - \gamma_{k_0} + \epsilon_n)) \rightarrow 0. \quad (2.6)$$

Analogously, from Theorem 1.12 and $\epsilon_n n^{2/3} \rightarrow \infty$, we also have

$$\mathbb{P}(\lambda_{k_0+1} \geq d_+ + \epsilon_n) = \mathbb{P}(n^{2/3}(\lambda_{k_0+1} - d_+) \geq n^{2/3}\epsilon_n) \rightarrow 0,$$

which together with (2.5) and (2.6) implies (2.4). Therefore, we complete the proof. \square

REMARK 2.3. *Although in theory, any sequence ϵ_n which satisfies the conditions $\epsilon_n \sqrt{n} \rightarrow 0$ and $\epsilon_n n^{2/3} \rightarrow \infty$ is applicable, we have to choose one in practice. Thus, it is worth to notice that the smaller ϵ_n one chooses, the easier one overestimates k_0 and vice versa. In the simulation, based on the idea in the paper [40, 46], we choose ϵ_n to be $\log \log(n)/n^{2/3}$, which makes our estimator conservative.*

In the following, we report a short simulation result to illustrate the performance of our estimator. For comparison, we also show the result of other three estimators in [21] by using the following model selection criteria: AIC, BIC and C_p respectively, which are

$$\hat{k}_A = \arg \min_{j \in \{0,1,\dots,q\}} AIC_j, \quad \hat{k}_B = \arg \min_{j \in \{0,1,\dots,q\}} BIC_j, \quad \hat{k}_C = \arg \min_{j \in \{0,1,\dots,q\}} CP_j.$$

Here for $j \in \{1, \dots, q\}$,

$$\begin{aligned} AIC_j &= -n \log \left[\prod_{i=j+1}^q (1 - \lambda_i) \right] - 2(p-j)(q-j), \\ BIC_j &= -n \log \left[\prod_{i=j+1}^q (1 - \lambda_i) \right] - \log(n)(p-j)(q-j), \\ CP_j &= n \sum_{i=j+1}^q \frac{\lambda_i}{1 - \lambda_i} - 2(p-j)(q-j) \end{aligned}$$

and $AIC_0 = BIC_0 = CP_0 = 0$.

According to the results in Table 1 and Table 2, we can find that the performance of our estimator is much better than AIC, BIC and C_p estimators when the dimensions are big. If the dimensions are small and the sample size is big enough, then besides our estimator, the AIC and C_p estimators perform well, but BIC estimator performs badly. That coincides with the conclusion in [21], which showed if q is fixed and $p/n \rightarrow c_1 > 0$ but small, AIC and C_p estimators are weakly consistent but BIC estimator is not.

	$p=10, r_c = 0.0071$				$p=60, r_c = 0.0445$				$p=110, r_c = 0.0854$			
	\hat{k}_0	\hat{k}_A	\hat{k}_B	\hat{k}_C	\hat{k}_0	\hat{k}_A	\hat{k}_B	\hat{k}_C	\hat{k}_0	\hat{k}_A	\hat{k}_B	\hat{k}_C
≤ 2	0	0	0	0	0	0	375	0	0	0	1000	0
3	0	0	0	0	90	0	625	0	95	0	0	0
4	1000	937	1000	935	1000	846	0	463	905	488	0	0
5	0	63	0	65	0	151	0	465	0	458	0	26
≥ 6	0	0	0	0	0	3	0	72	0	54	0	974
	$p=160, r_c = 0.1305$				$p=210, r_c = 0.1809$				$p=260, r_c = 0.2379$			
	\hat{k}_0	\hat{k}_A	\hat{k}_B	\hat{k}_C	\hat{k}_0	\hat{k}_A	\hat{k}_B	\hat{k}_C	\hat{k}_0	\hat{k}_A	\hat{k}_B	\hat{k}_C
≤ 2	0	0	1000	0	0	0	1000	0	61	0	1000	0
3	714	0	0	0	27	0	0	0	939	0	0	0
4	286	62	0	0	793	0	0	0	0	0	0	0
5	0	349	0	0	0	1	0	0	0	0	0	0
≥ 6	0	589	0	1000	0	999	0	1000	0	1000	0	1000

TABLE 1

We fix $(r_1, r_2, r_3, r_4) = (0.8, 0.6, 0.4, 0.2)$, $r_5 = \dots = r_q = 0$, $p/q = 2$, $n = 1000$, and let $\{p, q\}$ vary. The results are obtained based on 1000 replications with centered normal distributions. The tuning parameter ϵ_n is chosen to be $\log \log(n)/n^{2/3}$. We present the counting numbers of AIC, BIC, C_p estimators proposed in [21] and our estimator which are equal to the values in the first column. The correct numbers for the estimators are marked **bold**. Notice that as r_c changes along with $\{p, q\}$, the true value of k_0 are different.

2.3. Application 3: Estimating the values of the population CCC. As a measure to detect the correlation level between two random vectors, the population CCC are also important. Traditionally, one uses the sample CCC to estimate the population ones directly, since under

	$p=10, q=5, n=100$				$p=60, q=30, n=600$				$p=110, q=55, n=1100$			
	\hat{k}_0	\hat{k}_A	\hat{k}_B	\hat{k}_C	\hat{k}_0	\hat{k}_A	\hat{k}_B	\hat{k}_C	\hat{k}_0	\hat{k}_A	\hat{k}_B	\hat{k}_C
≤ 2	44	0	88	0	0	0	1000	0	0	0	1000	0
3	877	65	732	39	186	2	0	0	22	0	0	0
4	79	871	180	876	814	744	0	118	978	559	0	2
5	0	64	0	85	0	238	0	525	0	400	0	51
≥ 6	0	0	0	0	0	16	0	357	0	41	0	947
	$p=160, q=80, n=1600$				$p=210, q=105, n=2100$				$p=260, q=130, n=2600$			
	\hat{k}_0	\hat{k}_A	\hat{k}_B	\hat{k}_C	\hat{k}_0	\hat{k}_A	\hat{k}_B	\hat{k}_C	\hat{k}_0	\hat{k}_A	\hat{k}_B	\hat{k}_C
≤ 2	0	0	1000	0	0	0	1000	0	0	0	1000	0
3	6	0	0	0	3	0	0	0	0	0	0	0
4	994	426	0	0	1000	289	0	0	1000	188	0	0
5	0	482	0	0	0	547	0	0	0	546	0	0
≥ 6	0	92	0	1000	0	164	0	1000	0	266	0	1000

TABLE 2

We fix $(r_1, r_2, r_3, r_4) = (0.8, 0.6, 0.4, 0.2)$, $r_5 = \dots r_q = 0$, $c_1 = 0.1$, $c_2 = 0.05$, $r_c = 0.077$, $k_0 = 4$, and let $\{p, q, n\}$ change. The results are obtained based on 1000 replications with centered normal distributions. The tuning parameter ϵ_n is chosen to be $\log \log(n)/n^{2/3}$. We present the counting numbers of AIC, BIC, C_p estimators proposed in [21] and our estimator which are equal to the values in the first column. The correct numbers are marked **bold**.

the traditional assumption that p, q are fixed and $n \rightarrow \infty$, the sample CCC tend to the population ones almost surely (see Chapter 12 in [1]). However, it has been noticed that sample CCC suffer from inflation, which occurs when the number of observations is not sufficient (see [43, 15, 50]). But the good news is that now we can easily explain the reason of the inflation properties of the sample CCC by our Theorem 1.8. In this subsection, we will give an approach to estimate the population CCC which are bigger than $\sqrt{r_c}$.

Solving the equation (1.10) via replacing γ_i by λ_i , we have two solutions

$$\hat{r}_i := \frac{2c_1c_2 - c_1 - c_2 + \lambda_i \pm \sqrt{(\lambda_i - d_-)(\lambda_i - d_+)}}{2(c_1c_2 - c_1 - c_2 + 1)}. \quad (2.7)$$

Notice that the product of two solutions equals to $c_1c_2/[(1 - c_1)(1 - c_2)] = r_c^2$. Therefore, according to the fact that $\lambda_i > d_+$ and $\hat{r}_i \rightarrow r_i > r_c$ when $n \rightarrow \infty$, we should choose a plus sign in (2.7). Hence, the estimator of r_i is chosen to be

$$\hat{r}_i := \phi(\lambda_i) = \frac{2c_1c_2 - c_1 - c_2 + \lambda_i + \sqrt{(\lambda_i - d_-)(\lambda_i - d_+)}}{2(c_1c_2 - c_1 - c_2 + 1)}, \quad (2.8)$$

Then according to Theorem 1.8, we have the following theorem.

THEOREM 2.4. *Under the same conditions of Theorem 1.8, for any $1 \leq i \leq k_0$, we have almost surely*

$$\hat{r}_i \rightarrow r_i.$$

Next, we display a short simulation result in Table 3 and Table 4, which are the sample means and standard deviations (s.d.) of the estimators \hat{r}_i . In the simulation, we first estimate k_0 by (2.3) with the tuning parameter $\epsilon_n = \log \log(n)/n^{2/3}$, and then for $i \leq \hat{k}_0$, we obtain the estimators \hat{r}_i by (2.8). According to the results we find that our estimator is excellent, especially when the population CCC are not equal to each other. If the multiplicity of some CCC is bigger

	\hat{r}_1		\hat{r}_2		\hat{r}_3		\hat{r}_4	
	mean	s.d.	mean	s.d.	mean	s.d.	mean	s.d.
$p=10, q=5, n=100$	0.798	0.036	0.588	0.066	0.362	0.072	0.241	0.033
$p=60, q=30, n=600$	0.800	0.016	0.598	0.027	0.392	0.035	0.190	0.029
$p=110, q=55, n=1100$	0.799	0.012	0.599	0.020	0.397	0.025	0.189	0.026
$p=160, q=80, n=1600$	0.800	0.010	0.598	0.017	0.397	0.020	0.191	0.023
$p=210, q=105, n=2100$	0.800	0.008	0.600	0.014	0.397	0.019	0.194	0.020
$p=260, q=130, n=2600$	0.800	0.008	0.599	0.013	0.399	0.017	0.194	0.018

TABLE 3

We fix $(r_1, r_2, r_3, r_4) = (0.8, 0.6, 0.4, 0.2)$, $r_5 = \cdots r_q = 0$, $c_1 = 0.1$, $c_2 = 0.05$, $r_c = 0.077$, $k_0 = 4$, and let $\{p, q, n\}$ vary. The results are obtained based on 1000 replications with centered normal distributions. The tuning parameter ϵ_n is chosen to be $\log \log(n)/n^{2/3}$.

	\hat{r}_1		\hat{r}_2		\hat{r}_3		\hat{r}_4	
	mean	s.d.	mean	s.d.	mean	s.d.	mean	s.d.
$p=10, q=5, n=100$	0.800	0.037	0.594	0.061	0.409	0.066	0.288	0.051
$p=60, q=30, n=600$	0.799	0.015	0.599	0.027	0.419	0.029	0.356	0.031
$p=110, q=55, n=1100$	0.799	0.011	0.599	0.020	0.417	0.021	0.370	0.021
$p=160, q=80, n=1600$	0.800	0.010	0.600	0.015	0.415	0.018	0.376	0.018
$p=210, q=105, n=2100$	0.800	0.008	0.600	0.014	0.413	0.015	0.381	0.016
$p=260, q=130, n=2600$	0.800	0.007	0.600	0.014	0.412	0.014	0.383	0.015

TABLE 4

We fix $(r_1, r_2, r_3, r_4) = (0.8, 0.6, 0.4, 0.4)$, $r_5 = \cdots r_q = 0$, $c_1 = 0.1$, $c_2 = 0.05$, $r_c = 0.077$, $k_0 = 4$, and let $\{p, q, n\}$ vary. The results are obtained based on 1000 replications with centered normal distributions. The tuning parameter ϵ_n is chosen to be $\log \log(n)/n^{2/3}$.

than one, as the results shown in Table 4, there should be some \hat{r}_i 's close to each other. In this case, although $r_i = r_{i+1}$ for certain multiple population CCC, the estimator \hat{r}_i may differ from \hat{r}_{i+1} by a certain amount since our λ_i 's are ordered. Suppose now we can determine $r_i = r_{i+1}$ from the information in the CCA matrix, we may then use the average of \hat{r}_i and \hat{r}_{i+1} to get more precise estimate of both r_i and r_{i+1} . Hence, determining if there are multiple r_i 's would be important for the estimate of r_i as well, although sometimes directly using \hat{r}_i is sufficient. In the following, we propose a statistic to test the hypothesis:

$$H_0 : r_{j_0-1} > r_{j_0} = r_{j_0+1} = \cdots = r_{j_0+j_1-1} > r_{j_0+j_1} \quad \text{v.s.} \quad H_1 : \text{not } H_0, \quad (2.9)$$

where $j_0 \geq 1$, $j_0 + j_1 - 1 \leq k_0$ and $r_0 = 1$. If this test is not rejected, we then estimate $r_{j_0} = r_{j_0+1} = \cdots = r_{j_0+j_1-1}$ by $j_1^{-1} \sum_{i=j_0}^{j_0+j_1-1} \hat{r}_i$. Under the null hypothesis in (2.9), using Theorem 1.10, we know that the j_1 -dimensional random vector

$$\left\{ \frac{\sqrt{n}(\lambda_j - \gamma_j)}{\xi(r_{j_0})}, j \in \{j_0, \dots, j_1 - 1\} \right\}$$

converges weakly to the distribution of the ordered eigenvalues of j_1 -dimensional symmetric Gaussian random matrix G . Thus we can naturally use the testing statistics

$$T_n := \frac{\sqrt{n}(\lambda_{j_0} - \lambda_{j_0+j_1-1})}{\xi(\hat{r}_{j_0})},$$

where $\hat{r}_{j_0} = j_1^{-1} \sum_{i=j_0}^{j_0+j_1-1} \hat{r}_i$, to test the hypothesis (2.9). We have the following theorem.

THEOREM 2.5. *Suppose that the assumptions in Theorem 1.10 hold. Under the null hypothesis of (2.9), we have for any $x \in \mathbb{R}$,*

$$|\mathbb{P}(T_n \leq x) - \mathbb{P}(\lambda_1^G - \lambda_{j_1}^G \leq x)| \rightarrow 0,$$

where λ_1^G and $\lambda_{j_1}^G$ are the largest and smallest eigenvalues of j_1 -dimensional symmetric Gaussian random matrix G (defined in Theorem 1.10) respectively. In addition, if the alternative hypothesis H_1 is: there exists some $j_2 \in [0, j_1 - 2] \cap \mathbb{Z}$ such that $r_{j_0+j_2}$ and $r_{j_0+j_2+1}$ are well separated (c.f. Definition 1.9), we have the power function

$$\text{Power} = \mathbb{P}(T_n > q_\alpha(j_1)) \rightarrow 1,$$

where $q_\alpha(j_1)$ is the $1 - \alpha$ quantile of the distribution $\lambda_1^G - \lambda_{j_1}^G$.

This theorem can be easily obtained by using Skorohod strong representation theorem and Theorem 1.10, thus we omit the proof here. For convenient application, under the null hypothesis the 95% empirical quantile of the distribution $\lambda_1^G - \lambda_{j_1}^G$ is shown in Table 5 with some different j_1 by 10^8 bootstrap replicates.

	$j_1 = 2$	$j_1 = 3$	$j_1 = 4$	$j_1 = 5$	$j_1 = 6$	$j_1 = 7$	$j_1 = 8$	$j_1 = 9$	$j_1 = 10$	\cdots
$q_{5\%}(j_1)$	3.462	4.593	5.459	6.191	6.838	7.424	7.964	8.468	8.942	\cdots

TABLE 5

The 5% empirical quantile of the distribution $\lambda_1^G - \lambda_{j_1}^G$ with 4 digits.

Finally, we give a procedure of estimating the population CCC. The algorithm is shown in Table 6.

Step 1: $\{\lambda_1 \geq \dots \geq \lambda_q\} = \text{Eigenvalue}(C_{yx})$;
 Step 2: Test whether $r_1 = 0$.
 If not reject, then continue;
 Step 3: $\hat{k}_0 := \max\{i : \lambda_i \geq d_+ + \epsilon_n\}$;
 Step 4: $\hat{r}_i = \phi(\lambda_i)$, $i = 1, \dots, \hat{k}_0$;
 Step 5: If there exists some $\{j_0, \dots, j_0 + j_1 - 1\}$,
 such that $\frac{\sqrt{n}(\lambda_{j_0} - \lambda_{j_0+j_1-1})}{\xi(\hat{r}_{j_0})} \leq q5\%(j_1)$,
 then reset $\hat{r}_{j_0} = \dots = \hat{r}_{j_0+j_1-1} = j_1^{-1} \sum_{i=j_0}^{j_0+j_1-1} \hat{r}_i$.
 Step 6: $\hat{\rho}_i = \sqrt{\hat{r}_i}$.

TABLE 6

Algorithm for estimating the population CCC.

2.4. The limestone grassland community data. To illustrate the application of canonical correlation, we apply our result to a limestone grassland community data which can be obtained from Table A-2 of [24]. This data records eight species (\mathbf{x} , $p = 8$) and six soil variables (\mathbf{y} , $q = 6$) from forty-five ($n = 44$) 10m \times 10m stands in a limestone grassland community in Anglesey, North Wales. This experiment concerns relationships between the abundances of several plant species and associated soil characteristics. More explanations can be found in Chapter 7 of [24]. Notice that although the dimensions p, q and the sample size n are not so “large”, from the simulations (some results are not reported in this manuscript), we can already observe the high-dimensional effect when the dimensions are bigger than 5 and sample sizes are bigger than 20, i.e., the classical estimates are already not reliable. The eigenvalues of C_{yx} are as follows:

$$\lambda_1 = 0.829, \quad \lambda_2 = 0.520, \quad \lambda_3 = 0.359, \quad \lambda_4 = 0.107, \quad \lambda_5 = 0.094, \quad \lambda_6 = 0.038.$$

In addition, we have $d_+ = 0.533$ and $\xi_{tw}^3 = 0.468$. Then we use (2.1) to test the hypothesis $r_1 = 0$, and obtain the p-value to be 3.71×10^{-5} . Thus we have strong evidence to reject the null hypothesis, which suggests the existence of a linear relationship between \mathbf{x} and \mathbf{y} . Since there is only one eigenvalue of C_{yx} which is bigger than d_+ , we can only determine the existence of the first CCC $\hat{\rho}_1 = \sqrt{\hat{r}_1} = 0.864$, which is estimated by (2.8). For the rest CCC, we have no enough evidence to ensure their existence.

3. Preliminaries. In this section, we introduce some basic notions and known technical results escorting our proofs and calculations in the subsequent sections.

For any given probability distribution $\sigma(\lambda)$, its Stieltjes transform is known as

$$s_\sigma(z) := \int \frac{1}{\lambda - z} d\sigma(\lambda), \quad z \in \mathbb{C}^+ := \{\omega \in \mathbb{C} : \Im \omega > 0\}.$$

From the definition, we can immediately get the fact that $\Im s_\sigma(z) > 0$ for $z \in \mathbb{C}^+$. Actually, the definition of $s_\sigma(z)$ can be extended to the domain $\mathbb{C} \setminus \text{supp}(\sigma)$, by setting $s_\sigma(\bar{z}) = \overline{s_\sigma(z)}$, where $\text{supp}(\sigma)$ represents the support of $\sigma(\lambda)$. Then $s_\sigma(z)$ is holomorphic on $\mathbb{C} \setminus \text{supp}(\sigma)$.

In the sequel, we will also need the Stieltjes transform of the MANOVA matrices $A(A+B)^{-1}$ and $\mathcal{A}(\mathcal{A}+\mathcal{B})^{-1}$, where

$$A \sim \mathcal{W}_p(I_p, q), \quad B \sim \mathcal{W}_p(I_p, n - q), \quad \mathcal{A} \sim \mathcal{W}_q(I_q, p), \quad \mathcal{B} \sim \mathcal{W}_q(I_q, n - p)$$

From [2], we know that the Stieltjes transform of $A(A+B)^{-1}$ converges (a.s.) to

$$\check{s}(z) := \frac{z - c_1 - c_2 - \sqrt{(z - d_-)(z - d_+)}}{2c_1 z(z - 1)} - \frac{1}{z}, \quad (3.1)$$

and the Stieltjes transform of $\mathcal{A}(\mathcal{A} + \mathcal{B})^{-1}$ converges to

$$\tilde{s}(z) := \frac{z - c_1 - c_2 - \sqrt{(z - d_-)(z - d_+)}}{2c_2 z(z - 1)} - \frac{1}{z}. \quad (3.2)$$

Note that $\tilde{s}(z)$ is the Stieltjes transform of the distribution given in (1.6), according to the discussions in Section 1.1.

Throughout the paper, we will often use the well known large deviation result of the extreme eigenvalues of Wishart matrices. Assume that $S \sim \mathcal{W}_{n_1}(I_{n_1}, n_2)$ for some $n_1 := n_1(n)$ and $n_2 := n_2(n)$ satisfying $n_1/n \rightarrow a_1 \in (0, 1)$, $n_2/n \rightarrow a_2 \in (0, 1)$ as n tends to infinity, and $a := a_1/a_2 \in (0, 1)$. Denoting $\lambda_1(n_2^{-1}S)$ and $\lambda_{n_1}(n_2^{-1}S)$ the largest and the smallest eigenvalues of $n_2^{-1}S$ respectively, it is well known that for any given positive number $\varepsilon > 0$,

$$(1 + \sqrt{a})^2 + \varepsilon \geq \lambda_1(n_2^{-1}S) \geq \lambda_{n_1}(n_2^{-1}S) \geq (1 - \sqrt{a})^2 - \varepsilon \quad (3.3)$$

holds with overwhelming probability (see Theorem 2.13 of [14])

4. Determinant equation. In this section, we derive a determinant equation for the eigenvalues of C_{xy} which are not in the spectrum of the null case. We will see that the canonical correlation matrix in the finite rank case can be viewed as a finite rank perturbation of that in the null case.

As mentioned above, the canonical correlation coefficients are invariant under the block diagonal transformation $(\mathbf{x}_i, \mathbf{y}_i) \rightarrow (\mathbf{A}\mathbf{x}_i, \mathbf{B}\mathbf{y}_i)$, for any $p \times p$ matrix \mathbf{A} and $q \times q$ matrix \mathbf{B} , as long as both of them are nonsingular. Hence, to study λ_i , we can start with the following setting

$$\begin{pmatrix} \mathcal{X} \\ \mathcal{Y} \end{pmatrix} = \Sigma^{1/2} V, \quad \Sigma = \begin{pmatrix} I_p & T \\ T' & I_q \end{pmatrix},$$

where V is a $(p + q) \times n$ matrix with i.i.d. $N(0, 1)$ entries, and

$$T = \text{diag}(\sqrt{r_1}, \dots, \sqrt{r_k}) \oplus \mathbf{0}_{(p-k) \times (q-k)},$$

e.g. see Muirhead [38], page 530, formula (7). Apparently, there exists a Gaussian matrix \mathcal{W} independent of \mathcal{Y} such that

$$\begin{pmatrix} \mathcal{X} \\ \mathcal{Y} \end{pmatrix} = \begin{pmatrix} \mathcal{W} + T\mathcal{Y} \\ \mathcal{Y} \end{pmatrix}. \quad (4.1)$$

Here

$$\text{vec}(\mathcal{W}) \sim N(\mathbf{0}, I_n \otimes (I_p - TT')).$$

Hence, \mathcal{W} is a $p \times n$ matrix with independent entries. More specifically, $w_{ij} \sim N(0, 1 - r_i)$. According to the above definitions, we have

$$\Sigma_{xx} = I_p, \quad \Sigma_{yy} = I_q, \quad \Sigma_{xy} = T, \quad \Sigma_{yx} = T'.$$

For brevity, from now on, we use symbols W, X, Y to denote the scaled data matrices, i.e.,

$$W := \frac{1}{\sqrt{n}} \mathcal{W}, \quad X := \frac{1}{\sqrt{n}} \mathcal{X}, \quad Y := \frac{1}{\sqrt{n}} \mathcal{Y}. \quad (4.2)$$

Correspondingly, we introduce the notations

$$S_{ww} = WW', \quad S_{wy} = WY', \quad S_{yw} = YW', \quad S_{yy} = YY'. \quad (4.3)$$

In light of (4.1), we have

$$\begin{aligned} S_{xx} &= S_{ww} + TS_{yw} + S_{wy}T' + TS_{yy}T', \\ S_{xy} &= S_{wy} + TS_{yy}, \quad S_{yx} = S_{yw} + S_{yy}T'. \end{aligned} \quad (4.4)$$

Hence, due to the assumption that $\text{rank}(T) = k$, C_{xy} can be regarded as a finite rank perturbation of C_{wy} . Note that, with probability 1, the eigenvalues of C_{xy} are the solutions for λ of the characteristic equation

$$\det(D) := \det(S_{xy}S_{yy}^{-1}S_{yx} - \lambda S_{xx}) = 0. \quad (4.5)$$

In light of (4.4), it is equivalent to

$$\det(S_{wy}S_{yy}^{-1}S_{yw} - \lambda S_{ww} + (1 - \lambda)(TS_{yw} + S_{wy}T' + TS_{yy}T')) = 0.$$

Denote by

$$P_y = Y'(YY')^{-1}Y. \quad (4.6)$$

Then both P_y and $I_n - P_y$ are projections with $\text{rank}(P_y) = q$ and $\text{rank}(I_n - P_y) = n - q$ almost surely. Further, we decompose the matrix W and T as follows,

$$W = \begin{pmatrix} W_1 \\ W_2 \end{pmatrix}, \quad T = \begin{pmatrix} T_1 \\ \mathbf{0} \end{pmatrix}, \quad (4.7)$$

where W_1 (resp. T_1) is a $k \times n$ (resp. $k \times q$) matrix composed by the first k rows of W (resp. T). Correspondingly, we can introduce the notations like $S_{yw_2} = YW_2'$, $S_{w_2w_2} = W_2W_2'$, analogously to (4.3). Applying the decompositions in (4.7), we can rewrite (4.5) in the following form

$$\det(D) = \det \begin{pmatrix} D_{11} & D_{12} \\ D_{21} & D_{22} \end{pmatrix} = 0 \quad (4.8)$$

where

$$\begin{aligned} D_{11} &\equiv D_{11}(\lambda) := W_1P_yW_1' - \lambda W_1W_1' + (1 - \lambda)(T_1YW_1' + W_1Y'T_1' + T_1YY'T_1'), \\ D_{12} &\equiv D_{12}(\lambda) := W_1P_yW_2' - \lambda W_1W_2' + (1 - \lambda)T_1YW_2', \\ D_{21} &\equiv D_{21}(\lambda) := W_2P_yW_1' - \lambda W_2W_1' + (1 - \lambda)W_2Y'T_1' \end{aligned} \quad (4.9)$$

and

$$D_{22} \equiv D_{22}(\lambda) := W_2P_yW_2' - \lambda W_2W_2' = S_{w_2w_2}(C_{w_2y} - \lambda), \quad (4.10)$$

where C_{w_2y} is defined analogously to (1.2). In case D_{22} is invertible, we have the equation $\det(D) = \det(D_{22})\det(D_{11} - D_{12}D_{22}^{-1}D_{21}) = 0$, which implies

$$\det(D_{11}(\lambda) - D_{12}(\lambda)D_{22}^{-1}(\lambda)D_{21}(\lambda)) = 0. \quad (4.11)$$

Apparently, the sufficient condition of $D_{22}(\lambda)$ being invertible is that λ is not an eigenvalue of C_{w_2y} . Notice that since k is fixed and W_2 is independent of Y , we can apply the results in Theorems 1.3 and 1.5 to the matrix C_{w_2y} as well. Hence, if we want to investigate the eigenvalues which are not in $\text{Spec}(C_{w_2y})$, it suffices to solve (4.11) and find the properties of the limits of its solutions.

For brevity, we introduce the notation

$$M_n(z) := D_{11}(z) - D_{12}(z)D_{22}^{-1}(z)D_{21}(z) \quad (4.12)$$

which is a well defined $k \times k$ matrix-valued function for all $z \in \mathbb{C} \setminus \text{Spec}(C_{w_2y})$. using (4.9), (4.10) and the fact $YP_y = Y$, we can write

$$M_n(z) = \mathcal{Z}_1 \mathcal{A}(z) \mathcal{Z}_1', \quad (4.13)$$

where

$$\mathcal{Z}_1 = W_1 + T_1 Y, \quad \mathcal{A}(z) = (P_y - z) - (P_y - z) W_2' (W_2 (P_y - z) W_2')^{-1} W_2 (P_y - z). \quad (4.14)$$

Hereafter, we often use z to represent zI in short when the dimension of the identity matrix is clear.

5. Limits. In this section, we provide the proof of Theorem 1.8 based on several lemmas, whose proofs will be postponed. Our discussion consists of two parts, aimed at (i) and (ii) in Theorem 1.8, respectively: (1) For the outliers, we locate them by deriving the limits of the solutions to the equation (4.11); (2) For the eigenvalues sticking to d_+ , we simply use Cauchy's interlacing property to get the conclusion.

- *The outliers*

To locate the outliers, we start with the equation (4.11), i.e., $\det M_n(z) = 0$. Intuitively, if $M_n(z)$ is *close* to some deterministic matrix-valued function $M(z)$ in certain sense, it is reasonable to expect that the solutions of (4.11) are close to those of the equation $\det[M(z)] = 0$. Such an implication can be explicitly formulated in the *location lemma* given later, see Lemma 5.2. To state the result, we introduce more notations. Set for any positive constant δ the domain

$$\mathcal{D} \equiv \mathcal{D}(\delta) := \{z \in \mathbb{C} : d_+ + \delta < \Re z \leq 2, |\Im z| \leq 1\}. \quad (5.1)$$

Define the functions $s(z), m_i(z) : \mathcal{D} \rightarrow \mathbb{C}$ as follows:

$$s(z) := \frac{z - c_1 - c_2 - \sqrt{(z - d_-)(z - d_+)}}{2(z - 1)}. \quad (5.2)$$

$$m_i(z) := (c_2 - (1 + c_1)z - (1 - z)s(z))(1 - r_i) + (1 - z)\left(1 - \frac{1 - z}{c_2}s(z)\right)r_i, \quad (5.3)$$

where the square root is specified as the one with positive imaginary part in case $z \in \mathbb{C}^+$, and $s(\bar{z}) = \overline{s(z)}$. It is elementary to see that $s(z)$ and $m_i(z)$'s are all holomorphic on $\mathbb{C} \setminus [d_-, d_+]$ and thus also on \mathcal{D} . In addition, from the definition (5.2), $s(z)$ satisfies the following equation

$$(z - 1)s^2(z) + (c_1 + c_2 - z)s(z) - c_1 c_2 = 0. \quad (5.4)$$

Recall $\check{s}(z)$ and $\tilde{s}(z)$ from (3.1) and (3.2). We see that

$$\check{s}(z) = \frac{1}{c_1 z} s(z) - \frac{1}{z}, \quad \tilde{s}(z) = \frac{1}{c_2 z} s(z) - \frac{1}{z}. \quad (5.5)$$

Further, we define the diagonal matrix

$$M(z) := \text{diag}(m_1(z), \dots, m_k(z)).$$

Recall δ in the definition of \mathcal{D} in (5.1). We introduce the following event

$$\Xi_1 \equiv \Xi_1(n, \delta) := \left\{ \|C_{w_2y}\| \leq d_+ + \frac{\delta}{2} \right\}, \quad (5.6)$$

Note that, Theorem 1.5 tells us that Ξ_1 holds with overwhelming probability. Hence, we have $\|(C_{w_2y} - z)^{-1}\| = O(1)$ for all $z \in \mathcal{D}$ with overwhelming probability. This also implies that, in Ξ_1 , $M_n(z)$ is holomorphic on \mathcal{D} almost surely.

Our main technical task for the limits part is the following lemma.

LEMMA 5.1. *For any given $\delta > 0$, and any sufficiently small $\varepsilon > 0$,*

$$\sup_{z \in \mathcal{D}} \sup_{i,j=1,\dots,k} |(M_n)_{ij}(z) - M_{ij}(z)| \leq n^{-\varepsilon}$$

holds almost surely.

The proof of Lemma 5.1 will be postponed to Section 8. The following location lemma is a direct consequence of Lemma 5.1. At first, we remark here, it will be clear that the solutions of the equation $\det(M(z)) = 0$ can only be real.

LEMMA 5.2 (The location lemma). *For any given and sufficiently small $\delta > 0$, let $z_1 > \dots > z_\kappa$ be the solutions in $(d_+ + \delta, 1)$ of the equation $\det(M(z)) = 0$, with multiplicities n_1, \dots, n_κ respectively. Then for any fixed $\eta > 0$ and each $i \in \{1, \dots, \kappa\}$, almost surely, there exists $z_{n,i,1} > \dots > z_{n,i,\kappa_i}$ with multiplicities $m_{i,1}, \dots, m_{i,\kappa_i}$ respectively, satisfying $\sum_{j=1}^{\kappa_i} m_{i,j} = n_i$, such that*

$$\sup_{j=1,\dots,\kappa_i} |z_{n,i,j} - z_i| \leq \eta, \quad (5.7)$$

and $\cup_{i=1}^s \{z_{n,i,j}, j = 1, \dots, \kappa_i\}$ is the collection of all solutions (multiplicities counted) of the equation $\det M_n(z) = 0$ in $(d_+ + \delta, 1)$.

PROOF OF LEMMA 5.2 WITH LEMMA 5.1 GRANTED. At first, as mentioned above, in Ξ_1 , $M_n(z)$ is holomorphic on \mathcal{D} . Moreover, according to (3.3), both Y and W are bounded by some constant in operator norm with overwhelming probability. In addition, we know that Ξ_1 holds with overwhelming probability. Hence, according to the definition of $M_n(z)$ in (4.12), one sees that $\|M_n(z)\|$ is bounded uniformly on \mathcal{D} with overwhelming probability. Therefore, the entries of $M_n(z)$ are all bounded in magnitude with overwhelming probability as well. In addition, it is clear that $M(z)$ is holomorphic and its entries are also bounded in magnitude on \mathcal{D} . Hence, Lemma 5.1 implies that almost surely,

$$\sup_{z \in \mathcal{D}} |\det(M_n(z)) - \det(M(z))| \leq n^{-\varepsilon},$$

taking into account the fact that the determinant is a multivariate polynomial of the matrix entries. It is obvious that $\det[M_n(z)]$ only has real roots since the equation (4.5) does. Then by Rouché's theorem, we can immediately get (5.7). \square

Now, with the aid of Lemma 5.2, we prove (i) of Theorem 1.8.

PROOF OF (I) OF THEOREM 1.8. According to Lemma 5.2, it suffices to solve $\det(M(z)) = 0$ in $(d_+ + \delta, 1]$ to get z_i , for sufficiently small $\delta > 0$. By the definition of $M(z)$, we shall solve the equation

$$m_i(z) = 0. \quad (5.8)$$

It will be clear that there is a unique simple solution for the above equation. We denote it by γ_i in the sequel.

First, if $r_i = 1$, we get from the definition (5.3) that

$$m_i(\gamma_i) = (1 - \gamma_i) \left(1 - \frac{1 - \gamma_i}{c_2} s(\gamma_i) \right) = 0.$$

It is elementary to check that the only solution is $\gamma_i = 1$. If $r_i < 1$, we introduce the shorthand notation

$$t_i = \frac{r_i}{1 - r_i}. \quad (5.9)$$

According to the definition in (5.3), we see that (5.8) is equivalent to

$$\begin{aligned} & \sqrt{\gamma_i^2 + 2(2c_1c_2 - c_1 - c_2)\gamma_i + (c_1 - c_2)^2} (c_2 - t_i\gamma_i + t_i) \\ &= t_i\gamma_i^2 + (t_i(c_2 - c_1 - 1) + c_2 - 2c_1c_2)\gamma_i + (c_1 - c_2)t_i + c_1c_2 - c_2^2. \end{aligned} \quad (5.10)$$

Notice that if $\gamma_i \in (d_+, 1]$, we have

$$\gamma_i^2 + 2(2c_1c_2 - c_1 - c_2)\gamma_i + (c_1 - c_2)^2 = (\gamma_i - d_-)(\gamma_i - d_+) > 0$$

and $c_2 - t_i\gamma_i + t_i > 0$. Thus under the restriction that

$$t_i\gamma_i^2 + (t_i(c_2 - c_1 - 1) + c_2 - 2c_1c_2)\gamma_i + (c_1 - c_2)t_i + c_1c_2 - c_2^2 \geq 0, \quad (5.11)$$

solving (5.10), we get

$$\gamma_i = \frac{(1 + t_i^{-1}c_1)(1 + t_i^{-1}c_2)}{1 + t_i^{-1}} = r_i(1 - c_1 + c_1r_i^{-1})(1 - c_2 + c_2r_i^{-1}). \quad (5.12)$$

Applying (5.9) and (5.12), it is elementary to check that (5.11) is equivalent to

$$r_i \geq r_c = \sqrt{\frac{c_1c_2}{(1 - c_1)(1 - c_2)}}. \quad (5.13)$$

Note that

$$\begin{aligned} \gamma_i &= c_1(1 - c_2) + c_2(1 - c_1) + r_i(1 - c_1)(1 - c_2) + r_i^{-1}c_1c_2 \\ &\geq c_1(1 - c_2) + c_2(1 - c_1) + 2\sqrt{c_1c_2(1 - c_1)(1 - c_2)} = d_+ \end{aligned} \quad (5.14)$$

Moreover, equality holds in the second step of (5.14) only if $r_i = r_c$. Hence, (5.8) has solution in (d_+, ∞) only if $r_i > r_c$, with the solution γ_i given by (5.12). Now, what remains is to check $\gamma_i < 1$. It again follows from (5.13) by elementary calculation. Then by (5.7) in Lemma 5.1, we get that

$$\lambda_i - \gamma_i \xrightarrow{\text{a.s.}} 0, \quad \text{for } i = 1, \dots, k_0.$$

Hence, we conclude the proof of (i) of Theorem 1.8. \square

Now, we proceed to the proof of (ii) .

- *The Sticking eigenvalues*

PROOF OF (II) OF THEOREM 1.8. At first, according to the proof of (i) of Theorem 1.8, we see that with overwhelming probability, there are exactly k_0 largest eigenvalues (multiplicities counted) of C_{xy} in the interval $(d_+ + \delta, 1]$, for any sufficiently small $\delta > 0$. Hence, we see that $\limsup_{n \rightarrow \infty} \lambda_i \leq d_+ + \delta$ for all $i \geq k_0 + 1$ almost surely. Moreover, by the Cauchy's interlacing property and (1.8), we always have $\liminf_{n \rightarrow \infty} \lambda_i \geq d_+ - \delta$ almost surely for any fixed i . Since δ can be arbitrarily small, we get the conclusion that $\lambda_i - d_+ \xrightarrow{\text{a.s.}} 0$ for any fixed $i \geq k_0 + 1$. Hence, we complete the proof. \square

6. Fluctuations of the outliers. For the fluctuations of the outliers, we first derive a CLT for the entries of M_n , which together with (4.11) will imply a CLT for the outliers. Recall from Lemma 5.2 the notations n_1, \dots, n_κ as the multiplicities of the solutions of $\det[M(z)] = 0$. Set

$$J_l = \left\{ \sum_{i=1}^{l-1} n_i + 1, \dots, \sum_{i=1}^l n_i \right\}.$$

We derive the central limit theorem for the n_l -packed sample eigenvalues $\{\sqrt{n}(\lambda_j - \gamma_j), j \in J_l\}$. Note that γ_j 's are all the same for $j \in J_l$, and they are l -th largest solution of equation $\det[M(z)] = 0$ with multiplicity n_l . Actually, from Lemma 5.1 and Lemma 5.2, for $j \in J_l$, it would not be difficult to conclude that for any sufficiently small $\varepsilon > 0$,

$$\sup_{1 \leq \alpha, \beta \leq k} |(M_n)_{\alpha\beta}(\lambda_j) - M_{\alpha\beta}(\gamma_j)| \leq n^{-\varepsilon} \quad (6.1)$$

almost surely. However, for Theorem 1.10, we need to take a step further to consider the fluctuation of the difference in (6.1). We split the task into three steps, and the results are collected in Lemmas 6.1-6.3. The first lemma is to expand $M_n(\lambda_j)$ around $M_n(\gamma_j)$. Recall the notations ϑ, ϖ defined in (1.14).

LEMMA 6.1. *Suppose that the assumptions in Theorem 1.10 hold. Especially, $j \in \{1, \dots, k_0\}$ such that r_j is well separated from r_c and 1. As $n \rightarrow \infty$, we have*

$$\sqrt{n}[M_n(\lambda_j) - M_n(\gamma_j)] = -\sqrt{n}(\lambda_j - \gamma_j)(\Delta + o_p(1)),$$

where Δ is a diagonal matrix defined as

$$\Delta = \frac{(1 - c_1)\vartheta r_j^2 + c_1^2 c_2}{\vartheta r_j^2 - \varpi} (I_k - T_1 T_1') + \frac{(1 - c_1)\vartheta r_j^3 + c_1(3c_1 c_2 - c_1 - 2c_2 + 1)r_j - 2c_1^2 c_2}{r_j(\vartheta r_j^2 - \varpi)} T_1 T_1',$$

and $o_p(1)$ represents a generic $k \times k$ matrix with negligible (in probability) entries.

Recall $\mathcal{A}(z)$ defined in (4.14). We further introduce the following matrices

$$\begin{aligned} \mathcal{M}_1 &\equiv \mathcal{M}_1(z) := W_1 \mathcal{A}(z) W_1', \\ \mathcal{M}_2 &\equiv \mathcal{M}_2(z) := T_1 Y \mathcal{A}(z) W_1', \\ \mathcal{M}_3 &\equiv \mathcal{M}_3(z) := T_1 Y \mathcal{A}(z) Y' T_1'. \end{aligned} \quad (6.2)$$

According to the definitions in (4.12), (4.9) and (4.10), it is easy to check that

$$M_n(z) = \mathcal{M}_1 + \mathcal{M}_2 + \mathcal{M}_2' + \mathcal{M}_3. \quad (6.3)$$

We denote the SVD of Y by

$$Y = U_y \Lambda_y V_y. \quad (6.4)$$

It is well known that U_y and V_y are Haar distributed, and U_y, Λ_y, V_y are mutually independent (see (3.9) of [16] for instance). Then we see that

$$P_y = V_y' \Lambda_y' (\Lambda_y \Lambda_y')^{-1} \Lambda_y V_y. \quad (6.5)$$

is independent of U_y . With the above notations, we introduce

$$\mathcal{B} \equiv \mathcal{B}(z) := \Lambda_y V_y \mathcal{A}(z), \quad \mathcal{D} \equiv \mathcal{D}(z) := \Lambda_y V_y \mathcal{A}(z) V_y' \Lambda_y'. \quad (6.6)$$

Therefore, we can write

$$\mathcal{M}_1 = W_1 \mathcal{A} W_1', \quad \mathcal{M}_2 = T_1 U_y \mathcal{B} W_1', \quad \mathcal{M}_3 = T_1 U_y \mathcal{D} U_y' T_1'. \quad (6.7)$$

Set

$$\mathcal{M}(\gamma_j) := \frac{1}{n} \text{tr} \mathcal{A}(\gamma_j) \cdot (I_k - T_1 T_1') + \frac{1}{q} \text{tr} \mathcal{D}(\gamma_j) \cdot T_1 T_1'$$

The next lemma depicts the fluctuation of $M_n(\gamma_j)$ around $\mathcal{M}(\gamma_j)$.

LEMMA 6.2. *Suppose that the assumptions in Theorem 1.10 hold. For any $j \in \{1, \dots, k_0\}$ such that r_j is well separated from r_c and 1, we have that*

$$\sqrt{n} \left(M_n(\gamma_j) - \mathcal{M}(\gamma_j) \right)$$

converges weakly to a $k \times k$ symmetric Gaussian matrix \mathcal{R} with independent (up to symmetry) mean zero entries and the variance of $\mathcal{R}_{\alpha\beta}$ is

$$\begin{aligned} & \frac{1 + \delta_{\alpha\beta}}{r_j^2(\vartheta r_j^2 - \varpi)} \left(((1 - c_1)r_j + c_1)^2 (\vartheta r_j - \varpi)^2 ((1 - c_2)r_j^2 + c_2)(1 - r_\alpha)(1 - r_\beta) \right. \\ & \quad + (1 - c_2)((1 - c_1)r_j + c_1)^2 (1 - r_j)^2 (\vartheta r_j - \varpi)^2 (r_\alpha(1 - r_\beta) + r_\beta(1 - r_\alpha)) \\ & \quad \left. + ((1 - c_1)r_j + c_1)^2 (\vartheta r_j - \varpi)^2 ((1 - c_2)r_j^2 + 2c_1(1 - c_2)r_j + c_1(1 - 2c_2))r_\alpha r_\beta \right). \end{aligned} \quad (6.8)$$

Finally, we claim that the difference between $\mathcal{M}(\gamma_j)$ and $M(\gamma_j)$ is of order $o_p(\frac{1}{\sqrt{n}})$.

LEMMA 6.3. *Suppose that the assumptions in Theorem 1.10 hold. For any $j \in \{1, \dots, k_0\}$ such that r_j is well separated from r_c and 1, we have that*

$$\sqrt{n}(\mathcal{M}(\gamma_j) - M(\gamma_j)) = o_p(1),$$

where $o_p(1)$ represents a generic $k \times k$ matrix with negligible (in probability) entries.

The proofs of Lemmas 6.1-6.3 will be postponed to Section 9. With the aid of these lemmas, we can now prove Theorem 1.10.

PROOF OF THEOREM 1.10. Note that $\det M_n(\lambda_j) = 0$, and $M(\gamma_j)$ is a diagonal matrix with diagonal elements $m_i(\gamma_j)$, $i = 1, \dots, k$ (c.f. (5.3)). In addition, for any $l = 1, \dots, l_0$, $m_i(\gamma_j)$ is zero if $i \in J_l$ and nonzero otherwise. Therefore by Lemmas 6.1-6.3, the definition of T_1 and Skorohod strong representation theorem, we conclude that there exists a random matrix \mathcal{R} defined in Lemma 6.2, such that for any $j, \alpha, \beta \in J_l$,

$$\sqrt{n} \left((M_n)_{\alpha\beta}(\lambda_j) - M_{\alpha\beta}(\gamma_j) \right) = -\delta_{\alpha\beta} \Delta_{jj} \sqrt{n}(\lambda_j - \gamma_j)(1 + o_p(1)) + \mathcal{R}_{\alpha\beta} + o_p(1). \quad (6.9)$$

Let $\mathcal{N}^{(l)}$ be a $k \times k$ diagonal matrix with $\mathcal{N}_{jj}^{(l)} = n^{1/4}$ if $j \in J_l$ and $\mathcal{N}_{jj}^{(l)} = 1$ otherwise. Apparently we have $\det[\mathcal{N}^{(l)}] = n^{n_l/4} \neq 0$. Combining Theorem 1.8 with Lemmas 6.1-6.3, we conclude that

$$\mathcal{N}^{(l)} M_n(\lambda_j) \mathcal{N}^{(l)} = M(\gamma_j) + \left(\sqrt{n} (M_n)_{\alpha\beta}(\lambda_j) \mathbf{1}(j, \alpha, \beta \in J_l) \right)_{k \times k} + o_p(1).$$

Notice that $M(\gamma_j)$ is a diagonal matrix with J_l diagonal elements being zeros and

$$\left(\sqrt{n} (M_n)_{\alpha\beta}(\lambda_j) \mathbf{1}(j, \alpha, \beta \in J_l) \right)_{k \times k}$$

is a null matrix except the $J_l \times J_l$ block. According to $\det(\mathcal{N}^{(l)} M_n(\lambda_j) \mathcal{N}^{(l)}) = 0$ and (6.9), we obtain that

$$\det \left(- \lim_{n \rightarrow \infty} \left(\sqrt{n}(\lambda_j - \gamma_j)(I_{n_l} + o_p(1)) + \mathcal{R}(J_l \times J_l) / \Delta(j, j) \right) \right) = 0, \quad (6.10)$$

where $\mathcal{R}(J_l \times J_l)$ represents the $J_l \times J_l$ block of \mathcal{R} . From (6.10), we see that $\lim_{n \rightarrow \infty} \sqrt{n}(\lambda_j - \gamma_j)$ is the eigenvalue of $\mathcal{R}(J_l \times J_l) / \Delta_{jj}$. Since the eigenvalues of C_{yx} are simple almost surely, we see that the n_l random variables $\{\sqrt{n}(\lambda_j - \gamma_j), j \in J_l\}$ converge in probability to the set of eigenvalues of the matrix $\mathcal{R}(J_l \times J_l) / \Delta_{jj}$. The variance in (1.11) can be obtained from the definition of Δ in Lemma 6.1, and also (6.8).

Therefore, we complete the proof of Theorem 1.10. \square

7. Fluctuations of the sticking eigenvalues. In this section, we prove Theorem 1.12. When $k = 1$, the main idea is to compare the eigenvalues of C_{xy} with those of C_{w_2y} directly. Then we use the Tracy-Widom law for the largest eigenvalues of C_{w_2y} to conclude the fluctuation of the sticking eigenvalues of C_{xy} . For more general k , we use the argument for the case $k = 1$ recursively. Our proof strategy is inspired by the work [10].

For the comparison, throughout this section, we denote by $\tilde{\lambda}_1 \geq \tilde{\lambda}_2 \geq \dots \geq \tilde{\lambda}_q$ the ordered eigenvalues of C_{yw_2} , while use $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_q$ to denote the ordered eigenvalues of C_{yx} as before.

Our aim is to show the following proposition.

PROPOSITION 7.1. *Suppose that the assumptions in Theorem 1.12 holds. Then for any small positive constant ε and any fixed integer $i \geq 1$, the following holds in probability*

$$|\lambda_{k_0+i} - \tilde{\lambda}_i| \leq n^{-1+\varepsilon}. \quad (7.1)$$

With the aid of Proposition 7.1, we can prove Theorem 1.12 below.

PROOF OF THEOREM 1.12. Let

$$\hat{X} := W + T\hat{Y}, \quad (7.2)$$

where \hat{Y} is a i.i.d. copy of Y and \hat{Y} is independent of Y . Denote by $C_{y\hat{x}}$ via replacing X by \hat{X} in C_{yx} . Then $C_{y\hat{x}}$ is a CCA matrix in the null case. We first compare the eigenvalues of $C_{y\hat{x}}$, denoted by $\hat{\lambda}_i$'s, with the eigenvalues of C_{yw_2} , i.e., $\tilde{\lambda}_i$. Applying Proposition 7.1 to the case $r_1 = \dots = r_k = 0$, we can get $\max_{1 \leq i \leq m} |\hat{\lambda}_i - \tilde{\lambda}_i| \leq n^{-1+\varepsilon}$ in probability. This together with (7.1) implies (1.12). This together with Remark 1.13 completes the proof of Theorem 1.12. \square

To prove Proposition 7.1. We will need the following lemmas.

LEMMA 7.2. *For any $i = 0, \dots, q$, and $j = 1, \dots, k$, the diagonal entry $(M_n)_{jj}(z)$ is a decreasing function in $z \in (\tilde{\lambda}_{i+1}, \tilde{\lambda}_i)$, where $\tilde{\lambda}_0 = \infty$ and $\tilde{\lambda}_{q+1} = -\infty$.*

PROOF. Recall (4.13) and (4.14). It is elementary to check that

$$\frac{\partial \mathcal{A}(z)}{\partial z} = -\mathcal{Q}(z)(\mathcal{Q}(z))^T,$$

where

$$\mathcal{Q}(z) := I + (P_y - z)W_2'(W_2(P_y - z)W_2')^{-1}W_2$$

Note that, as long as $z \notin \text{Spec}(C_{w_2y})$, $\frac{\partial \mathcal{A}(z)}{\partial z}$ is a well-defined negative semi-definite matrix. Hence,

$$\frac{\partial}{\partial z}(M_n)_{jj}(z) = \left(\mathcal{Z}_1 \frac{\partial \mathcal{A}(z)}{\partial z} \mathcal{Z}_1' \right)_{jj} \leq 0$$

in the interval $(\tilde{\lambda}_{i+1}, \tilde{\lambda}_i)$ for any $i = 0, \dots, q$, which means that $(M_n)_{jj}(z)$ is decreasing. \square

Let $\mathbf{m} \geq 1$ be any fixed integer. For any small constants $\varepsilon, \delta > 0$, we introduce the following random domain

$$\Omega \equiv \Omega(\mathbf{m}, \varepsilon, \delta) := (\tilde{\lambda}_1 + n^{-1+\varepsilon}, d_+ + \delta) \cup \left\{ \cup_{i=1}^{\mathbf{m}} (\tilde{\lambda}_{i+1} + n^{-1+\varepsilon}, \tilde{\lambda}_i - n^{-1+\varepsilon}) \right\} \quad (7.3)$$

We also need the following lemma.

LEMMA 7.3. *With the above notations, we have the following:*

(i): *If $i \neq j$, for any small $\varepsilon, \delta > 0$, there exists some positive constant $c > 0$, such that*

$$\sup_{z \in \Omega} |(M_n)_{ij}(z)| \leq n^{-c}$$

holds with overwhelming probability.

(ii): *For any small $\varepsilon > 0$ and any small $\varepsilon' > 0$, there exists a sufficiently small δ , such that*

$$\sup_{z \in \Omega} |(M_n)_{ii}(z) - m_i(d_+)| \leq \varepsilon'$$

holds with overwhelming probability.

The proof of Lemma 7.3 is postponed to Appendix B.

In the sequel, we first show the proof of Proposition 7.1 in the case of $k = 1$. At the end of this section, we will extend the discussion to general k case. Note that when $k = 1$, We have $k_0 = 1$ or $k_0 = 0$, which means $r_1 > r_c$ or $r_1 < r_c$, respectively. We restate Proposition 7.1 for $k = 1$ in the following proposition.

PROPOSITION 7.4. *Suppose that $k = 1$. We have the following*

(i): *If $r_1 > r_c$, then for any fixed integer $m \geq 1$, and any $\varepsilon > 0$, we have*

$$|\lambda_{m+1} - \tilde{\lambda}_m| \leq n^{-1+\varepsilon}$$

in probability.

(ii): *If $r_1 < r_c$, then for any fixed integer $m \geq 1$, and any $\varepsilon > 0$, we have*

$$|\lambda_m - \tilde{\lambda}_m| \leq n^{-1+\varepsilon}$$

in probability.

With Lemma 7.2 and Lemma 7.3, we prove Proposition 7.4.

PROOF OF PROPOSITION 7.4. Since $k = 1$, we have $\det M_n = M_n(z) = M_{11}(z)$. Applying Lemma 7.3 with $k = 1$, we see that for any small ε' , there is a sufficiently small $\delta > 0$ such that

$$\sup_{z \in \Omega} |\det M_n(z) - m_1(d_+)| \leq \varepsilon' \quad (7.4)$$

with overwhelming probability. Since we assume that r_1 is well separated from r_c , it is easy to check from the definition (5.3) that $|m_1(d_+)| \geq c$ for some positive constant c . Hence, choosing

ε' in (7.4) sufficiently small, we see that $|\det M_n(z)| \neq 0$ uniformly on Ω , with overwhelming probability. That means, with overwhelming probability, there is no eigenvalue of C_{xy} in Ω .

Let $\mathbf{w}_i, i = 1, \dots, p$ be the rows of W . Note that, in case of $k = 1$, we have $\mathbf{w}_1 = W_1$. Moreover, P_w is the projection onto the subspace spanned by $\{\mathbf{w}_1 + T_1 Y, \mathbf{w}_2, \dots, \mathbf{w}_p\}$, and P_{w_2} is the projection onto the subspace spanned by $\{\mathbf{w}_2, \dots, \mathbf{w}_p\}$. Then, by Cauchy interlacing, we know that $\text{Spec}(P_w P_y P_w)$ and $\text{Spec}(P_{w_2} P_y P_{w_2})$ are interlacing. This implies that

$$\lambda_1 \geq \tilde{\lambda}_1 \geq \lambda_2 \geq \dots \geq \tilde{\lambda}_{q-1} \geq \lambda_q \geq \tilde{\lambda}_q, \quad (7.5)$$

since the nonzero eigenvalues of C_{wy} (resp. $C_{w_2 y}$) are the same as those of $P_w P_y P_w$ (resp. $P_{w_2} P_y P_{w_2}$).

In case (i): $r_1 > r_c$, from Theorem 1.8 we know that $\lambda_1 \rightarrow \gamma_1 > d_+ + \delta$ almost surely, for any sufficiently small $\delta > 0$. Since λ_i 's are solutions to $\det M_n(z) = 0$, and $m_1(d_+) \neq 0$, we see from (7.4) and (7.5) that

$$\lambda_i \in [\tilde{\lambda}_{i+1}, \tilde{\lambda}_{i+1} + n^{-1+\varepsilon}] \cup [\tilde{\lambda}_i - n^{-1+\varepsilon}, \tilde{\lambda}_i], \quad i = 2, \dots, \mathbf{m}.$$

in probability.

In case (ii): $r_1 < r_c$, from Theorem 1.8 we know that $\lambda_1 \rightarrow d_+$ in probability. Hence, we have

$$\lambda_1 \in [\tilde{\lambda}_1, \tilde{\lambda}_1 + n^{-1+\varepsilon}], \quad \lambda_i \in [\tilde{\lambda}_{i+1}, \tilde{\lambda}_{i+1} + n^{-1+\varepsilon}] \cup [\tilde{\lambda}_i - n^{-1+\varepsilon}, \tilde{\lambda}_i], \quad i = 2, \dots, \mathbf{m}.$$

Therefore, to prove Proposition 7.4, it suffices to check that for all $i = 2, \dots, \mathbf{m}$, λ_i is in $[\tilde{\lambda}_{i+1}, \tilde{\lambda}_{i+1} + n^{-1+\varepsilon}]$ or $[\tilde{\lambda}_i - n^{-1+\varepsilon}, \tilde{\lambda}_i]$. Note that $\det M_n(z) = (M_n)_{11}(z)$ is decreasing in the interval $(\tilde{\lambda}_{i+1}, \tilde{\lambda}_i)$, according to Lemma 7.2. Furthermore, since $s(d_+) = \frac{d_+ - c_1 - c_2}{2(d_+ - 1)}$, it is elementary to check

$$m_j(d_+) > 0, \quad \text{if } r_j > r_c, \quad m_j(d_+) < 0, \quad \text{if } r_j < r_c.$$

Therefore, if $r_1 > r_c$, according to (7.4), we have $\det M_n(z) > 0$ on $(\tilde{\lambda}_{i+1} + n^{-1+\varepsilon}, \tilde{\lambda}_i - n^{-1+\varepsilon})$. By the monotonicity of $\det M_n(z)$, we see that $\det M_n(z) = 0$ can only be achieved if $z \in [\tilde{\lambda}_i - n^{-1+\varepsilon}, \tilde{\lambda}_i]$. Hence, in case of $r_1 \geq r_c$, we have $\lambda_i \in [\tilde{\lambda}_i - n^{-1+\varepsilon}, \tilde{\lambda}_i]$ for all $i = 2, \dots, \mathbf{m}$. In contrast, when $r_1 < r_c$, we have $\lambda_i \in [\tilde{\lambda}_{i+1}, \tilde{\lambda}_{i+1} + n^{-1+\varepsilon}]$ for all $i = 2, \dots, \mathbf{m}$.

This concludes the proof of Proposition 7.4. □

Now, we prove Proposition 7.1.

PROOF OF PROPOSITION 7.1. For general k , we need more notations. For $1 \leq a \leq k$, let

$$W_{1a} := \begin{pmatrix} \mathbf{w}_1 \\ \vdots \\ \mathbf{w}_a \end{pmatrix}, \quad T_{1a} := \begin{pmatrix} t_1 & & \\ & \ddots & \\ & & t_a \end{pmatrix}$$

be the $a \times n$ matrices composed of the first a rows of W and T , respectively, and we write

$$W = \begin{pmatrix} W_{1a} \\ W_{2a} \end{pmatrix}, \quad T = \begin{pmatrix} T_{1a} \\ T_{2a} \end{pmatrix}.$$

We take the above as the definitions of W_{2a} and T_{2a} . Further, we denote

$$X_a := W_{2a} + T_{2a} Y.$$

Denote the eigenvalues of C_{x_ay} by

$$\lambda_{1,a} \geq \lambda_{2,a} \geq \cdots \quad (7.6)$$

Especially, with the above notations, we have $\tilde{\lambda}_i = \lambda_{i,k}$ and $\lambda_i = \lambda_{i,0}$. Our aim is compare $\lambda_{i,a-1}$'s with $\lambda_{i,a}$'s. First of all, by Cauchy interlacing, we have

$$\lambda_{i-1,a} \geq \lambda_{i,a-1} \geq \lambda_{i,a}. \quad (7.7)$$

Further, if $\lambda_{i,a-1} \notin \text{Spec}(C_{x_ay})$, then $\lambda_{i,a-1}$ is a solution to the equation

$$\mathbf{z}_{a-1} \mathcal{P}_a \mathbf{z}'_{a-1} = 0, \quad (7.8)$$

where

$$\mathbf{z}_{a-1} := \mathbf{w}_{a-1} + t_{a-1} \mathbf{y}_{a-1},$$

and

$$\mathcal{P}_a(z) := (P_y - z) - (P_y - z)X'_a(X_a(P_y - z)X'_a)^{-1}X_a(P_y - z).$$

We start with the case $a = k - 1$ and compare $\lambda_{i,a-1}$'s with $\lambda_{i,a}$'s. This is equivalent to discuss the case $k = 2$. Hence, we can set $k = 2$ and $a = 1$. We recall the decomposition of D in (4.8). By Schur complement, we have

$$D^{-1} = \begin{pmatrix} D_{11} & D_{12} \\ D_{21} & D_{22} \end{pmatrix}^{-1} = \begin{pmatrix} (M_n)^{-1} & * \\ * & * \end{pmatrix}, \quad (7.9)$$

where $*$'s represent other blocks, whose explicit expressions are irrelevant to us. Observe that M_n (c.f. (4.12)) is 2×2 by the assumption $k = 2$. On the other hand, using a different decomposition, we can also get by Schur complement that

$$D^{-1} = \begin{pmatrix} \mathbf{z}_1(P_y - z)\mathbf{z}'_1 & \mathbf{z}_1(P_y - z)X_2 \\ X'_2(P_y - z)\mathbf{z}'_1 & X_2(P_y - z)X'_2 \end{pmatrix}^{-1} = \begin{pmatrix} (\mathbf{z}_1 \mathcal{P}_2 \mathbf{z}'_1)^{-1} & * \\ * & * \end{pmatrix}. \quad (7.10)$$

From (7.9) and (7.10), we obtain

$$\mathbf{z}_1 \mathcal{P}_2 \mathbf{z}'_1 = \frac{1}{(M_n^{-1})_{11}}. \quad (7.11)$$

Then, we first consider the eigenvalues of C_{x_1y} , which is a perturbation of C_{w_2y} . Recall the notations introduced in (7.6). Using Proposition 7.4, we see that the following statements hold in probability:

(i): If $r_2 > r_c$, we have for any fixed integer $m \geq 0$

$$|\lambda_{m+1,1} - \tilde{\lambda}_m| \leq n^{-1+\varepsilon}. \quad (7.12)$$

(ii): If $r_2 < r_c$, we have for any fixed integer $m \geq 0$,

$$|\lambda_{m,1} - \tilde{\lambda}_m| \leq n^{-1+\varepsilon}. \quad (7.13)$$

According to the assumption $r_1 \geq r_2$, it suffices to consider the following three cases:

$$(i)': r_1 \geq r_2 > r_c, \quad (ii)': r_1 > r_c > r_2, \quad (iii)': r_c > r_1 > r_2.$$

First, we consider case (i)'. Applying Theorem 1.8, we see that

$$\lambda_{1,0} \rightarrow \gamma_1, \quad \lambda_{2,0} \rightarrow \gamma_2, \quad \lambda_{m,0} \rightarrow d_+$$

almost surely. Recall Cauchy interlacing formula (7.7). We see that $\lambda_{m,1} \geq \lambda_{m+1,0} \geq \lambda_{m+1,1}$. We first show

$$\lambda_{m+1,0} \notin (\tilde{\lambda}_m + n^{-1+\varepsilon}, \tilde{\lambda}_{m-1} - n^{-1+\varepsilon}) \quad (7.14)$$

in probability. Then according to (7.12), we have

$$\lambda_{m+1,0} \notin (\lambda_{m+1,1} + 2n^{-1+\varepsilon}, \lambda_{m,1} - 2n^{-1+\varepsilon}) \quad (7.15)$$

in probability.

To see (7.14), we use (7.11), which together with Lemma 7.3 implies

$$\sup_{z \in \Omega} |\mathbf{z}_1 \mathcal{P}_2(z) \mathbf{z}'_1 - m_1(d_+)| \leq \varepsilon'.$$

Hence, (7.14) and (7.15) follow. Further, similarly to Lemma 7.2, we can show that $\mathbf{z}_1 \mathcal{P}_2(z) \mathbf{z}'_1$ is decreasing in $(\lambda_{m,1}, \lambda_{m+1,1})$. Then, by the sign of $m_1(d_+)$ and the fact that $\lambda_{i,0}$ is the solution to the equation $\mathbf{z}_1 \mathcal{P}_2(z) \mathbf{z}'_1 = 0$, we see that

$$|\lambda_{m+1,0} - \lambda_{m,1}| \leq 2n^{-1+\varepsilon} \quad (7.16)$$

in probability. This together with (7.12) further implies that

$$|\lambda_{m+1,0} - \tilde{\lambda}_{m-1}| \leq 3n^{-1+\varepsilon}. \quad (7.17)$$

Then we get the conclusion for case (i)', by slightly modifying ε . For case (ii)', it suffices to go through the above discussion, with $(\tilde{\lambda}_{m-1}, \tilde{\lambda}_m)$ replaced by $(\tilde{\lambda}_m, \tilde{\lambda}_{m+1})$ in (7.14) and (7.18). For case (iii)', we first need replace $(\tilde{\lambda}_{m-1}, \tilde{\lambda}_m)$ by $(\tilde{\lambda}_m, \tilde{\lambda}_{m+1})$ in (7.14) and (7.18). Then, instead of (7.16), we have

$$|\lambda_{m+1,0} - \lambda_{m+1,1}| \leq 2n^{-1+\varepsilon}$$

since the sign of $m_1(d_+)$ changed. This together with (7.13) further implies that

$$|\lambda_{m+1,0} - \tilde{\lambda}_{m+1}| \leq 3n^{-1+\varepsilon}. \quad (7.18)$$

The above discussion on the case $k = 2$ simply gives the result for $\lambda_{i,k-1}$ for general k . Then, we apply the above argument again to compare $\lambda_{i,k-3}$'s with $\lambda_{i,k-2}$'s. This is equivalent to consider the eigenvalues of C_{xy} with $k = 3$. Using the argument recursively, we can finally get the result for λ_i 's for general k . This concludes the proof. \square

8. Proof of Lemma 5.1. To ease the presentation, we introduce the following notations

$$\begin{aligned} \Phi &\equiv \Phi(z) := D_{22}^{-1}(z) = (W_2(P_y - z)W'_2)^{-1}, \\ \Upsilon &\equiv \Upsilon(z) := (Y(P_{w_2} - z)Y')^{-1}, \end{aligned} \quad (8.1)$$

Observe that $\text{rank}(P_{w_2}) = p - k$ and $\text{rank}(P_y) = q$ almost surely. For any $m \times \ell$ rectangle matrix A , using SVD, it is easy to check the identity

$$A(A'A - zI_\ell)^{-1}A' = AA'(AA' - zI_m)^{-1} = I_m + z(AA' - zI_m)^{-1}. \quad (8.2)$$

Moreover, we can write

$$\Phi(z) = S_{w_2 w_2}^{-\frac{1}{2}} (\tilde{C}_{w_2 y} - z)^{-1} S_{w_2 w_2}^{-\frac{1}{2}}, \quad (8.3)$$

where we introduced the symmetric matrix

$$\tilde{C}_{w_2 y} := S_{w_2 w_2}^{-\frac{1}{2}} S_{w_2 y} S_{yy}^{-1} S_{y w_2} S_{w_2 w_2}^{-\frac{1}{2}}. \quad (8.4)$$

Using (8.3) and (8.2), it is not difficult to check

$$S_{yy}^{-1/2} S_{y w_2} \Phi S_{w_2 y} S_{yy}^{-1/2} = I_q + z S_{yy}^{1/2} \Upsilon S_{yy}^{1/2}, \quad (8.5)$$

from which we immediately get

$$S_{y w_2} \Phi S_{w_2 y} = S_{yy} + z S_{yy} \Upsilon S_{yy}. \quad (8.6)$$

Further, we introduce the following notations for Wishart matrices

$$\begin{aligned} E &:= W_2 P_y W_2', & H &:= W_2 (I - P_y) W_2', \\ \mathcal{E} &:= Y P_{w_2} Y', & \mathcal{H} &:= Y (I - P_{w_2}) Y'. \end{aligned} \quad (8.7)$$

By Cochran's Theorem, we know that E and H are independent, so are \mathcal{E} and \mathcal{H} . In addition, in light of (4.2), we have

$$\begin{aligned} nE &\sim \mathcal{W}_{p-k}(I_{p-k}, q), & nH &\sim \mathcal{W}_{p-k}(I_{p-k}, n-q), \\ n\mathcal{E} &\sim \mathcal{W}_q(I_q, p-k), & n\mathcal{H} &\sim \mathcal{W}_q(I_q, n-p+k). \end{aligned} \quad (8.8)$$

Observe that

$$\begin{aligned} S_{w_2 w_2} &= E + H, & S_{yy} &= \mathcal{E} + \mathcal{H}, \\ C_{w_2 y} &= (E + H)^{-1} E, & C_{y w_2} &= (\mathcal{E} + \mathcal{H})^{-1} \mathcal{E}, \end{aligned} \quad (8.9)$$

where $C_{w_2 y}$ and $C_{y w_2}$ are defined analogously to (1.2). From (8.9), we can also write

$$\Phi(z) = ((1-z)E - zH)^{-1} \quad \Upsilon(z) = ((1-z)\mathcal{E} - z\mathcal{H})^{-1}. \quad (8.10)$$

Therefore, we have

$$\begin{aligned} S_{yy} \Upsilon S_{yy} &= (\mathcal{E} + \mathcal{H}) ((1-z)\mathcal{E} - z\mathcal{H})^{-1} (\mathcal{E} + \mathcal{H}) \\ &= (1-z)^{-1} \mathcal{E} - z^{-1} \mathcal{H} + (z(1-z))^{-1} ((1-z)\mathcal{H}^{-1} - z\mathcal{E}^{-1})^{-1} \end{aligned} \quad (8.11)$$

Following from (8.6) and (8.11) we obtain

$$(1-z) S_{y w_2} \Phi S_{w_2 y} = \mathcal{E} + ((1-z)\mathcal{H}^{-1} - z\mathcal{E}^{-1})^{-1}. \quad (8.12)$$

In the sequel, we use the notation

$$\Psi \equiv \Psi_n(z) := ((1-z)\mathcal{H}^{-1} - z\mathcal{E}^{-1})^{-1}. \quad (8.13)$$

Then, we can write

$$(1-z) S_{y w_2} \Phi S_{w_2 y} = \mathcal{E} + \Psi. \quad (8.14)$$

Recall the SVD in (6.4). Note that $W_1, W_2, U_y, \Lambda_y, V_y$ are mutually independent. According to the definitions of $\mathcal{A}, \mathcal{B}, \mathcal{D}$ in (4.14) and (6.6) the, definition of Φ in (8.1), and (6.5), we see that (W_1, U_y) is independent of $\mathcal{A}, \mathcal{B}, \mathcal{D}$. Hence, we can condition on W_2, Λ_y, V_y and thus condition on $\mathcal{A}, \mathcal{B}, \mathcal{D}$ and use the randomness of W_1, U_y in the sequel.

Our first step is to use a $k \times q$ Gaussian matrix to approximate $T_1 U_y$. For simplicity, we denote

$$\mathcal{T} := \text{diag}(\sqrt{r_1}, \dots, \sqrt{r_k}).$$

Apparently, $\mathcal{T}^2 = T_1 T_1'$. More specifically, we claim the following lemma.

LEMMA 8.1. *With the above notations, there exists a $k \times q$ Gaussian matrix G_1 with i.i.d. $N(0, \frac{1}{q})$ entries, such that for any sufficiently large $K > 0$,*

$$\begin{aligned}\mathcal{M}_2 &= \mathcal{T}G_1\mathcal{B}W_1' + O\left(\frac{(\log n)^K}{n}\right), \\ \mathcal{M}_3 &= \mathcal{T}G_1\mathcal{D}G_1'\mathcal{T} - \frac{1}{q}\text{tr } \mathcal{D} \cdot \mathcal{T}(G_1G_1' - I)\mathcal{T} + O\left(\frac{(\log n)^K}{n}\right),\end{aligned}\quad (8.15)$$

hold uniformly in $z \in \mathcal{D}$, with overwhelming probability. Here $O(\frac{(\log n)^K}{n})$ represents a generic $k \times k$ random matrix with all entries of order $O(\frac{(\log n)^K}{n})$.

The proof of Lemma 8.1 will be postponed to Appendix. In the sequel, we condition on $\mathcal{A}, \mathcal{B}, \mathcal{D}$, and use the randomness of W_1 and G_1 only. First, we introduce the following well-known large deviation inequalities for Gaussian which will be frequently used in the sequel.

LEMMA 8.2. *Assume that $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ are two i.i.d. $N(0, n^{-1}I_n)$ vectors. Let $A \in \mathbb{C}^{n \times n}$ and $\mathbf{b} \in \mathbb{R}^n$ be independent of \mathbf{x} and \mathbf{y} . The following three statements hold with overwhelming probability for any sufficiently large constant $K > 0$:*

$$\begin{aligned}(i) : |\mathbf{x}'A\mathbf{x} - \frac{1}{n}\text{tr } A| &\leq \frac{(\log n)^K}{n}\|A\|_{HS}, \quad (ii) : |\mathbf{x}'A\mathbf{y}| \leq \frac{(\log n)^K}{n}\|A\|_{HS}, \\ (iii) : |\mathbf{x}'\mathbf{b}| &\leq \frac{(\log n)^K}{\sqrt{n}}\|\mathbf{b}\|_2.\end{aligned}$$

With the aid of Lemma 8.2, we can derive the following.

LEMMA 8.3. *For any given $\delta > 0$ in the definition of \mathcal{D} (c.f. (5.1)), and any sufficiently large constant $K > 0$, the following inequalities hold uniformly on \mathcal{D} , with overwhelming probability,*

$$\begin{aligned}\sup_{i,j} \left| (\mathcal{M}_1)_{ij} - \delta_{ij}(1 - r_i) \left(c_2 - (1 + c_1)z - (1 - z)\frac{1}{n}\text{tr } (E\Phi) \right) \right| &\leq \frac{(\log n)^K}{\sqrt{n}}, \\ \sup_{i,j} \left| (\mathcal{M}_2)_{ij} \right| &\leq \frac{(\log n)^K}{\sqrt{n}}, \quad \sup_{i,j} \left| (\mathcal{M}_3)_{ij} - \delta_{ij}r_i(1 - z)\frac{1}{q}\text{tr } (\mathcal{H} - \Psi) \right| \leq \frac{(\log n)^K}{\sqrt{n}}.\end{aligned}\quad (8.16)$$

For simplicity, we denote by $\mathbf{w}_i = (w_{i1}, \dots, w_{in})$ and $\mathbf{b}_i := (b_{i1}, \dots, b_{iq})$ the i -th rows of W_1 and G_1 , respectively.

PROOF OF LEMMA 8.3. In light of the definitions in (8.7) and (4.14), and also (8.10), it is elementary to check that

$$\begin{aligned}\frac{1}{n}\text{tr } \mathcal{A} &= \frac{q}{n} - z - \frac{1}{n}\text{tr } (\Phi((1 - z)^2E + z^2H)) \\ &= \frac{q}{n} - z - z\frac{p}{n} - (1 - z)\frac{1}{n}\text{tr } (E\Phi).\end{aligned}\quad (8.17)$$

In addition, using the fact that $\|W_2\| \leq C$ with overwhelming probability (c.f. (3.3)), it is not difficult to see that for all $z \in \mathcal{D}$,

$$\|\mathcal{A}\|_{HS}^2 \leq Cn(+ \|\Phi\|^2). \quad (8.18)$$

with overwhelming probability. In addition, we have

$$\|\Phi\| \leq \|(C_{w_2y} - z)^{-1}\| \|S_{w_2w_2}^{-1}\| = O(1) \quad (8.19)$$

with overwhelming probability, where we used the fact that the event Ξ_1 holds with overwhelming probability, and also (3.3). Hence, $\|\mathcal{A}\|_{\text{HS}} \leq C\sqrt{n}$ holds with overwhelming probability on \mathcal{D} .

Then, conditioning on the matrix \mathcal{A} and use the randomness of \mathbf{w}_i 's only, the first statement in (8.16) follows from Lemma 8.2 (i), (ii) immediately.

For the second estimate in (8.16), we condition on \mathcal{B} , and use the randomness of \mathbf{w}_i 's and \mathbf{b}_i 's. Then the conclusion follows from Lemma 8.2 (ii) and the fact that

$$\|\mathcal{B}\|_{\text{HS}} \leq C\sqrt{n} \quad (8.20)$$

holds uniformly on \mathcal{D} , with overwhelming probability. To see (8.20), we recall the definition of \mathcal{B} from (6.6). Similarly to (8.18),

$$\|\mathcal{B}\|_{\text{HS}}^2 \leq Cn(1 + \|\Phi\|^2).$$

Using (8.19) again, we see that (8.20) holds.

For the third estimate, we use the randomness of \mathbf{b}_i 's and condition on the \mathcal{D} . Then the conclusion again follows from Lemma 8.2 (i), (ii), the fact

$$\frac{1}{q} \text{tr } \mathcal{D} = \frac{1-z}{q} \text{tr} \left(S_{yy} - (1-z)S_{yw_2}\Phi S_{w_2y} \right) = \frac{1-z}{q} \text{tr} (\mathcal{H} - \Psi) \quad (8.21)$$

and the fact that

$$\|\mathcal{D}\|_{\text{HS}} \leq C\sqrt{n} \quad (8.22)$$

holds with overwhelming probability. The proof of (8.22) is similar to that of (8.20). We thus omit the details.

This completes the proof of Lemma 8.3. \square

Recall $s(z)$ in (5.2). For brevity, we further introduce the following function

$$\varrho(z) := \frac{1-z}{c_2} s(z) - c_1. \quad (8.23)$$

Hence, what remains is to estimate the normalized traces of $E\Phi$, \mathcal{H} and Ψ . We collect the results in the following lemma, whose proof will be postponed to Appendix A.

LEMMA 8.4. *For any given $z \in \mathcal{D}$, we have the following estimates:*

$$\begin{aligned} \frac{1}{q} \text{tr } \mathcal{H} &= \frac{n-p}{n} + o_p\left(\frac{1}{\sqrt{n}}\right), & \frac{1}{n} \text{tr } E\Phi(z) &= s(z) + o_p\left(\frac{1}{\sqrt{n}}\right), \\ \frac{1}{q} \text{tr } \Psi(z) &= \varrho(z) + o_p\left(\frac{1}{\sqrt{n}}\right). \end{aligned} \quad (8.24)$$

Further, for any sufficiently small constant $\varepsilon > 0$, we have the uniform estimate

$$\sup_{z \in \mathcal{D}} \left| \frac{1}{n} \text{tr } E\Phi(z) - s(z) \right| = O_{a.s.}(n^{-\varepsilon}), \quad \sup_{z \in \mathcal{D}} \left| \frac{1}{q} \text{tr } \Psi(z) - \varrho(z) \right| = O_{a.s.}(n^{-\varepsilon}).$$

With the aid of Lemmas 8.3 and 8.4, we can now prove Lemma 5.1.

PROOF OF LEMMA 5.1. According to the decomposition in (6.3), we see from Lemmas 8.3 and 8.4 that

$$\begin{aligned} (M_n)_{ij}(z) &= (\mathcal{M}_1)_{ij}(z) + (\mathcal{M}_2)_{ij}(z) + (\mathcal{M}_2)_{ji}(z) + (\mathcal{M}_3)_{ij}(z) \\ &= \delta_{ij} \left((c_2 - (1+c_1)z - (1-z)s(z))(1-r_i) + (1-z)(1-c_1-\varrho(z))r_i \right) + O_{a.s.}(n^{-\varepsilon}) \\ &= \delta_{ij} m_i(z) + O_{a.s.}(n^{-\varepsilon}) \end{aligned}$$

uniformly in $z \in \mathcal{D}$ and i, j . Here in the last step we used the definitions in (5.3) and (8.23). This completes the proof of Lemma 5.1. \square

9. Proofs of Lemma 6.1-6.3. In this section we prove Lemmas 6.1-6.3. First, from (1.7), (1.10) and (1.14), it is easy to check

$$\sqrt{(\gamma_j - d_-)(\gamma_j - d_+)} = \frac{\vartheta r_j^2 - \varpi}{r_j}.$$

This further implies

$$s(\gamma_j) = -\frac{\varpi}{\vartheta r_j - \varpi}, \quad s'(\gamma_j) = \frac{\varpi \vartheta r_j^2}{(\vartheta r_j - \varpi)^2 (\vartheta r_j^2 - \varpi)}. \quad (9.1)$$

Moreover, we need the following lemma whose proof is postponed to Appendix A.

LEMMA 9.1. *Suppose that the assumptions in Theorem 1.10 hold. For any $j \in \{1, \dots, k_0\}$ such that r_j is well separated from r_c and 1, we have the following estimates*

$$\begin{aligned} \frac{1}{n} \text{tr } H\Phi(\gamma_j) &= \frac{-c_1 + (1 - \gamma_j)s(\gamma_j)}{\gamma_j} + o_p(1), \\ \frac{1}{n} \text{tr } (H\Phi(\gamma_j)H\Phi(\gamma_j)) &= \frac{\gamma_j(1 - \gamma_j)^2 s'(\gamma_j) + (\gamma_j^2 - 1)s(\gamma_j) + c_1}{\gamma_j^2} + o_p(1), \\ \frac{1}{n} \text{tr } (E\Phi(\gamma_j)H\Phi(\gamma_j)) &= (1 - \gamma_j)s'(\gamma_j) - s(\gamma_j) + o_p(1), \\ \frac{1}{n} \text{tr } (E\Phi(\gamma_j)E\Phi(\gamma_j)) &= \gamma_j s'(\gamma_j) + s(\gamma_j) + o_p(1). \end{aligned}$$

Then, recall the definitions of \mathcal{E} and \mathcal{H} in (8.7), Ψ in (8.13). We have the following lemma whose proof is also postponed to Appendix A.

LEMMA 9.2. *Suppose that the assumptions in Theorem 1.10 hold. For any $j \in \{1, \dots, k_0\}$ such that r_j is well separated from r_c and 1, we have the following results:*

$$\begin{aligned} \sup_{\alpha, \beta} \left| \Psi_{\alpha\beta}(\gamma_j) - \delta_{\alpha\beta} \varrho(\gamma_j) \right| &= o_p(1), \\ \sup_{\alpha, \beta} \left| \left(\Psi(\gamma_j)(\mathcal{H}^{-1} + \mathcal{E}^{-1})\Psi(\gamma_j) \right)_{\alpha\beta} - \delta_{\alpha\beta} \varrho'(\gamma_j) \right| &= o_p(1). \end{aligned}$$

9.1. Proof of Lemma 6.1. Recall the decomposition of M_n in (6.3). In the sequel, we consider $\mathcal{M}_1, \mathcal{M}_2$ and \mathcal{M}_3 separately.

Recall \mathcal{M}_1 in (6.2), $\mathcal{A}(z)$ in (4.14) and $\Phi(z)$ in (8.10). Using the identity

$$\Phi(z_1) - \Phi(z_2) = (z_1 - z_2)\Phi(z_1)W_2W_2'\Phi(z_2), \quad (9.2)$$

it is not difficult to write

$$\sqrt{n}(\mathcal{M}_1(\lambda_j) - \mathcal{M}_1(\gamma_j)) = \sqrt{n}(\lambda_j - \gamma_j) \sum_{a=0}^4 W_1 \mathcal{A}_a W_1'. \quad (9.3)$$

where

$$\begin{aligned} \mathcal{A}_0 &:= -I_n, \quad \mathcal{A}_1 := -P_y W_2' \Phi(\gamma_j) W_2 W_2' \Phi(\lambda_j) W_2 P_y, \\ \mathcal{A}_2 &= \mathcal{A}_3' := P_y W_2' \Phi(\lambda_j) W_2 + \gamma_j P_y W_2' \Phi(\gamma_j) W_2 W_2' \Phi(\lambda_j) W_2, \\ \mathcal{A}_4 &:= -(\lambda_j + \gamma_j) W_2' \Phi(\lambda_j) W_2 - \gamma_j^2 W_2' \Phi(\gamma_j) W_2 W_2' \Phi(\lambda_j) W_2. \end{aligned} \quad (9.4)$$

Since $W_1 W_1' \rightarrow (I_k - T_1 T_1')$ almost surely, we have

$$W_1 \mathcal{A}_0 W_1' = -(I_k - T_1 T_1') + o_p(1).$$

The equation (9.2) together with Theorem 1.8, (8.19) and the fact that the event Ξ_1 in (5.6) holds with overwhelming probability implies that

$$\|\Phi(\lambda_j)\| = O_p(1), \quad \|\Phi(\lambda_j) - \Phi(\gamma_j)\| = o_p(1). \quad (9.5)$$

By (9.5), Lemma 8.2 and Lemma 9.1, we have

$$\begin{aligned} W_1 \mathcal{A}_1 W_1' &= -\frac{1}{n} \text{tr} (P_y W_2' \Phi(\gamma_j) W_2 W_2' \Phi(\gamma_j) W_2 P_y) \cdot (I_k - T_1 T_1') + o_p(1) \\ &= -s'(\gamma_j)(I_k - T_1 T_1') + o_p(1). \end{aligned}$$

Here we used the definition of $E = W_2 P_y W_2'$, and the cyclicity of trace.

Analogously, we obtain for $a = 2, 3$

$$W_1 \mathcal{A}_a W_1' = (s(\gamma_j) + \gamma_j s'(\gamma_j))(I_k - T_1 T_1') + o_p(1).$$

and

$$W_1 \mathcal{A}_4 W_1' = (c_1 - s(\gamma_j) - \gamma_j s'(\gamma_j))(I_k - T_1 T_1') + o_p(1).$$

Combining the above estimate yields

$$\begin{aligned} &\sqrt{n}(\mathcal{M}_1(\lambda_j) - \mathcal{M}_1(\gamma_j)) \\ &= \sqrt{n}(\lambda_j - \gamma_j) \left((c_1 - 1 + s(\gamma_j) + (\gamma_j - 1)s'(\gamma_j))(I_k - T_1 T_1') + o_p(1) \right). \end{aligned} \quad (9.6)$$

Analogously, for \mathcal{M}_2 , using Lemma 8.2, we will get

$$\sqrt{n}(\mathcal{M}_2(\lambda_j) - \mathcal{M}_2(\gamma_j)) = \sqrt{n}(\lambda_j - \gamma_j) o_p(1), \quad (9.7)$$

At the end, we consider \mathcal{M}_3 (c.f. (6.2)). By (8.14), we can write

$$\mathcal{M}_3 \equiv \mathcal{M}_3(z) := (1 - z)T_1(\mathcal{H} - \Psi)T_1'.$$

Notice that by definition in (8.13), we have

$$\Psi = \mathcal{H}\Upsilon\mathcal{E}. \quad (9.8)$$

And we also have the expansion

$$\Psi(z_1) - \Psi(z_2) = (z_1 - z_2)\Psi(z_2)(\mathcal{H}^{-1} + \mathcal{E}^{-1})\Psi(z_1) = (z_1 - z_2)\mathcal{H}\Upsilon(z_2)Y Y' \Upsilon(z_1)\mathcal{E}. \quad (9.9)$$

Furthermore, analogously to (8.19), we have

$$\|\Upsilon\| \leq \|(C_{yw_2} - z)^{-1}\| \|S_{yy}^{-1}\| = O(1) \quad (9.10)$$

with overwhelming probability, for $z \in \mathcal{D}$. Hence, from (9.8)-(9.10), we also have the analogues of (8.19) and (9.5)

$$\|\Psi(\gamma_j)\| = O_p(1), \quad \|\Psi(\lambda_j)\| = O_p(1), \quad \|\Psi(\lambda_j) - \Psi(\gamma_j)\| = o_p(1).$$

Then approximating $\Phi(\lambda_j)$ by $\Phi(\gamma_j)$, and using Lemma 9.2, we obtain

$$\sqrt{n}(\mathcal{M}_3(\lambda_j) - \mathcal{M}_3(\gamma_j))$$

$$\begin{aligned}
&= -\sqrt{n}(\lambda_j - \gamma_j)T_1(\mathcal{H} - \Psi(\lambda_j) + (1 - \gamma_j)\Psi(\lambda_j)(\mathcal{H}^{-1} + \mathcal{E}^{-1})\Psi(\gamma_i))T_1' \\
&= -\sqrt{n}(\lambda_j - \gamma_j)\left(\left(1 - c_1 - \varrho(\gamma_j) + (1 - \gamma_j)\varrho'(\gamma_j)\right)T_1T_1' + o_p(1)\right) \\
&= -\sqrt{n}(\lambda_j - \gamma_j)\left(\left(1 - 2\frac{(1 - \gamma_j)s(\gamma_j)}{c_2} + \frac{(1 - \gamma_j)^2s'(\gamma_j)}{c_2}\right)T_1T_1' + o_p(1)\right), \tag{9.11}
\end{aligned}$$

where in the second step we used the fact $\mathcal{H}_{ij} = \delta_{ij}(1 - c_1) + o_p(1)$, and in the last step we used the definition (8.23).

Therefore, combining (9.1) with the equations (9.6), (9.7) and (9.11), we can conclude the proof of Lemma 6.1 by elementary calculation.

9.2. Proof of Lemma 6.2. In this subsection, we prove Lemma 6.2. Recall the matrices defined in (6.6). Let

$$\mathring{\mathcal{D}} := \mathcal{D} - \frac{1}{q}\text{tr } \mathcal{D} \cdot I.$$

In according to (6.3), (6.7) and Lemma 8.1, we can write

$$M_n - \frac{1}{q}\text{tr } \mathcal{D} \cdot \mathcal{T}^2 = \begin{pmatrix} W_1 & \mathcal{T}G_1 \end{pmatrix} \begin{pmatrix} \mathcal{A} & \mathcal{B}' \\ \mathcal{B} & \mathring{\mathcal{D}} \end{pmatrix} \begin{pmatrix} W_1' \\ (\mathcal{T}G_1)' \end{pmatrix} + o_p\left(\frac{1}{\sqrt{n}}\right). \tag{9.12}$$

In the sequel, we condition on $\mathcal{A}, \mathcal{B}, \mathcal{D}$, and use the randomness of W_1 and G_1 only. Recall the notation $\mathbf{w}_i = (w_{i1}, \dots, w_{in})$ as the i -th rows of W_1 and further denote by $\mathbf{d}_i := (d_{i1}, \dots, d_{iq})$ the i -th row of $\mathcal{T}G_1$.

According to (9.12), we see that

$$\begin{aligned}
&\sqrt{n}\left((M_n)_{ij}(\gamma_j) - \delta_{ij}\left((1 - r_i)\frac{1}{n}\text{tr } \mathcal{A} + r_i\frac{1}{q}\text{tr } \mathcal{D}\right)\right) \\
&= \sqrt{n}\left(\mathbf{w}_i\mathcal{A}\mathbf{w}_j' - \delta_{ij}(1 - r_i)\frac{1}{n}\text{tr } \mathcal{A}\right) + \sqrt{n}\left(\mathbf{d}_i\mathcal{B}\mathbf{w}_j' + \mathbf{w}_i\mathcal{B}'\mathbf{d}_j'\right) \\
&\quad + \sqrt{n}\mathbf{d}_i\mathring{\mathcal{D}}\mathbf{d}_j' + o_p(1) =: Q_{ij} + o_p(1). \tag{9.13}
\end{aligned}$$

Let \mathbb{E}_0 be the conditional expectation w.r.t. $\mathcal{A}, \mathcal{B}, \mathcal{D}$. Set

$$f_{\mathbf{c}}(t) := \exp\left(it \sum_{i \leq j} c_{ij}Q_{ij}\right), \quad \phi_{\mathbf{c}}(t) := \mathbb{E}_0 f_{\mathbf{c}}(t) \tag{9.14}$$

for any collection of fixed numbers $\mathbf{c} := (c_{ij})_{i \leq j}$. It suffices to show that for some deterministic $s \equiv s(\mathbf{c})$,

$$\phi_{\mathbf{c}}'(t) = -s^2 t \phi_{\mathbf{c}}(t) + o_p(1). \tag{9.15}$$

From (9.15) one can solve $\phi_{\mathbf{c}}(t) = \exp(-\frac{s^2 t^2}{2} + o_p(1))$, which further implies

$$\mathbb{E} f_{\mathbf{c}}(t) = \mathbb{E} \phi_{\mathbf{c}}(t) \rightarrow \exp(-\frac{s^2 t^2}{2}).$$

This is known as Tikhomirov's method for CLT [44].

To establish (9.15), we start with the definitions in (9.13) and (9.14). Then we have

$$\phi_{\mathbf{c}}'(t) = i\sqrt{n} \sum_{i \leq j} c_{ij} \sum_{\ell=1}^q \left(\mathbb{E}_0 \left(w_{i\ell} (\mathcal{A}\mathbf{w}_j')_{\ell} f_{\mathbf{c}}(t) \right) - \delta_{ij}(1 - r_i) \frac{1}{n} \text{tr } \mathcal{A} \phi_{\mathbf{c}}(t) \right)$$

$$\begin{aligned}
& + i\sqrt{n} \sum_{i \leq j} c_{ij} \sum_{\ell=1}^q \mathbb{E}_0 \left(\left(w_{j\ell}(\mathbf{d}_i \mathcal{B})_\ell + w_{i\ell}(\mathcal{B}' \mathbf{d}'_j)_\ell \right) f_{\mathbf{c}}(t) \right) \\
& + i\sqrt{n} \sum_{i \leq j} c_{ij} \sum_{\ell=1}^q \mathbb{E}_0 \left(d_{i\ell}(\mathring{\mathcal{D}} \mathbf{d}'_j)_\ell f_{\mathbf{c}}(t) \right).
\end{aligned} \tag{9.16}$$

Using integration by parts formula for Gaussian

$$\mathbb{E} g h(g) = \sigma^2 \mathbb{E} h'(g), \quad g \sim N(0, \sigma^2), \tag{9.17}$$

we see that

$$\begin{aligned}
\mathbb{E}_0 \left(w_{i\ell}(\mathcal{A} \mathbf{w}'_j)_\ell f_{\mathbf{c}}(t) \right) &= \frac{1}{n} \delta_{ij} (1 - r_i) \mathcal{A}_{\ell\ell} \phi_{\mathbf{c}}(t) + \frac{it}{n} (1 - r_i) \sum_{\alpha \leq \beta} c_{\alpha\beta} \mathbb{E}_0 \left(\frac{\partial Q_{\alpha\beta}}{\partial w_{i\ell}} (\mathcal{A} \mathbf{w}'_j)_\ell f_{\mathbf{c}}(t) \right) \\
\mathbb{E}_0 \left(\left(w_{j\ell}(\mathbf{d}_i \mathcal{B})_\ell + w_{i\ell}(\mathcal{B}' \mathbf{d}'_j)_\ell \right) f_{\mathbf{c}}(t) \right) \\
&= \frac{it}{n} \sum_{\alpha \leq \beta} c_{\alpha\beta} \mathbb{E}_0 \left(\left((1 - r_j) \frac{\partial Q_{\alpha\beta}}{\partial w_{j\ell}} (\mathbf{d}_i \mathcal{B})_\ell + (1 - r_i) \frac{\partial Q_{\alpha\beta}}{\partial w_{i\ell}} (\mathcal{B}' \mathbf{d}'_j)_\ell \right) f_{\mathbf{c}}(t) \right). \\
\mathbb{E}_0 \left(d_{i\ell}(\mathring{\mathcal{D}} \mathbf{d}'_j)_\ell f_{\mathbf{c}}(t) \right) &= \frac{1}{q} \delta_{ij} r_i \mathring{\mathcal{D}}_{\ell\ell} \phi_{\mathbf{c}}(t) + \frac{it}{q} \sum_{\alpha \leq \beta} c_{\alpha\beta} r_i \mathbb{E}_0 \left(\frac{\partial Q_{\alpha\beta}}{\partial d_{i\ell}} (\mathring{\mathcal{D}} \mathbf{d}'_j)_\ell f_{\mathbf{c}}(t) \right).
\end{aligned} \tag{9.18}$$

Plugging (9.18) into (9.16) yields

$$\phi'_{\mathbf{c}}(t) = -t \mathbb{E}_0 \left(Z f_{\mathbf{c}}(t) \right), \tag{9.19}$$

where

$$\begin{aligned}
Z &:= \frac{1}{\sqrt{n}} \sum_{i \leq j} c_{ij} \sum_{\alpha \leq \beta} c_{\alpha\beta} \sum_{\ell} \left((1 - r_i) \frac{\partial Q_{\alpha\beta}}{\partial w_{i\ell}} (\mathcal{A} \mathbf{w}'_j)_\ell \right. \\
&\quad \left. + ((1 - r_j) \frac{\partial Q_{\alpha\beta}}{\partial w_{j\ell}} (\mathbf{d}_i \mathcal{B})_\ell + (1 - r_i) \frac{\partial Q_{\alpha\beta}}{\partial w_{i\ell}} (\mathcal{B}' \mathbf{d}'_j)_\ell) + \frac{n}{q} r_i \frac{\partial Q_{\alpha\beta}}{\partial d_{i\ell}} (\mathring{\mathcal{D}} \mathbf{d}'_j)_\ell \right).
\end{aligned} \tag{9.20}$$

Now, note that

$$\begin{aligned}
\frac{\partial Q_{\alpha\beta}}{\partial w_{i\ell}} &= \delta_{\alpha i} \sqrt{n} \left((\mathcal{A} \mathbf{w}'_\beta)_\ell + (\mathcal{B}' \mathbf{d}'_\beta)_\ell \right) + \delta_{\beta i} \sqrt{n} \left((\mathbf{w}_\alpha \mathcal{A})_\ell + (\mathbf{d}_\alpha \mathcal{B})_\ell \right), \\
\frac{\partial Q_{\alpha\beta}}{\partial d_{i\ell}} &= \delta_{\alpha i} \sqrt{n} \left((\mathcal{B} \mathbf{w}'_\beta)_\ell + (\mathring{\mathcal{D}} \mathbf{d}'_\beta)_\ell \right) + \delta_{\beta i} \sqrt{n} \left((\mathbf{w}_\alpha \mathcal{B})_\ell + (\mathbf{d}_\alpha \mathring{\mathcal{D}})_\ell \right).
\end{aligned} \tag{9.21}$$

Substituting (9.21) into (9.20), and using Lemma 8.2, together with the bounds $\|\mathcal{A}\|, \|\mathcal{B}\|, \|\mathring{\mathcal{D}}\| \leq C$ with overwhelming probability on \mathcal{D} , it is not difficult to get

$$\begin{aligned}
Z &= \sum_{i \leq j} c_{ij}^2 (1 + \delta_{ij}) \left((1 - r_i)(1 - r_j) \frac{1}{n} \text{tr } \mathcal{A}^2 + (r_i(1 - r_j) + r_j(1 - r_i)) \frac{1}{q} \text{tr } \mathcal{B} \mathcal{B}' \right. \\
&\quad \left. + \frac{n}{q^2} r_i r_j \text{tr } \mathring{\mathcal{D}}^2 \right) + o_p(1) =: s_n^2 + o_p(1).
\end{aligned} \tag{9.22}$$

Plugging (9.22) into (9.19), we get

$$\phi'_{\mathbf{c}}(t) = -t s_n^2 \phi_{\mathbf{c}}(t) + o_p(1). \tag{9.23}$$

This implies that Q_{ij} 's (c.f. (9.13)) are asymptotically independent Gaussians (up to symmetry), and the limiting variance of Q_{ij} is given by the limit of

$$(1 - r_i)(1 - r_j) \frac{1}{n} \text{tr } \mathcal{A}^2 + (r_i(1 - r_j) + r_j(1 - r_i)) \frac{1}{q} \text{tr } \mathcal{B} \mathcal{B}' + \frac{n}{q^2} r_i r_j \text{tr } \mathring{\mathcal{D}}^2.$$

Hence, in the sequel, it suffices to estimate the normalized traces of \mathcal{A}^2 , $\mathcal{B} \mathcal{B}'$, and $\mathring{\mathcal{D}}^2$. The result is formulated in the following lemma, whose proof will be also postponed to Appendix A.

LEMMA 9.3. *Suppose that the assumptions in Theorem 1.10 hold. For any $j \in \{1, \dots, k_0\}$ such that r_j is well separated from r_c and 1, we have*

$$\begin{aligned} \frac{1}{n} \text{tr } \mathcal{A}^2 &= -(1 - \gamma_j)^2 s(\gamma_j) + \gamma_j (1 - \gamma_j)^2 s'(\gamma_j) + \gamma_j^2 (1 - c_1) + (1 - 2\gamma_j) c_2 + o_p(1), \\ \frac{1}{q} \text{tr } \mathcal{B} \mathcal{B}' &= (1 - \gamma_j)^2 \left(1 - \frac{1}{c_2} s(\gamma_j) + \frac{\gamma_j (1 - \gamma_j)}{c_2} s'(\gamma_j) \right) + o_p(1), \\ \frac{1}{q} \text{tr } \mathring{\mathcal{D}}^2 &= (1 - \gamma_j)^2 \left(-c_1 + c_2 + \left(\frac{(\gamma_j - 1)^2}{c_2} + \frac{(2c_2 - 1)}{c_2} \gamma_j + \frac{c_1 - c_2}{c_2} \right) s(\gamma_j) \right. \\ &\quad \left. - \frac{(1 + (c_2 - 1)\gamma_j)(1 - \gamma_j)}{c_2^2} s^2(\gamma_j) + \frac{\gamma_j (1 - \gamma_j)^2}{c_2} s'(\gamma_j) \right) + o_p(1). \end{aligned} \quad (9.24)$$

With the above lemma and (9.23), we conclude the proof of Lemma 6.2.

9.3. *Proof of Lemma 6.3.* It is equivalent to show the following lemma.

LEMMA 9.4. *Suppose that the assumptions in Theorem 1.10 hold. For any $j \in \{1, \dots, k_0\}$ such that r_j is well separated from r_c and 1, we have*

$$\begin{aligned} \frac{1}{n} \text{tr } \mathcal{A}(\gamma_j) &= c_2 - (1 + c_1) \gamma_j - (1 - \gamma_j) s(\gamma_j) + o_p\left(\frac{1}{\sqrt{n}}\right), \\ \frac{1}{q} \text{tr } \mathcal{D}(\gamma_j) &= (1 - \gamma_j)(1 - c_1 - \varrho(\gamma_j)) + o_p\left(\frac{1}{\sqrt{n}}\right). \end{aligned} \quad (9.25)$$

PROOF. The results follows from Lemma 8.4, (8.17) and (8.21) immediately. \square

With Lemma 9.4, we can conclude the proof of Lemma 6.3 easily from the definitions in (5.3) and (8.23).

APPENDIX A

In this appendix, we prove Lemma 8.1, Lemma 8.4, Lemma 9.1, Lemma 9.2, and Lemma 9.3.

PROOF OF LEMMA 8.1. Let \mathbf{u}_i be the i -th row of U_y . We only state the details for the proof of the case $k = 2$. For general fixed k , the proof is just a straightforward extension. It is well known that a Haar orthogonal matrix can be obtained from a Gaussian matrix with i.i.d. entries via Gram-Schmidt orthogonalization. Especially, there exists a pair of i.i.d. Gaussian vectors $\mathbf{g}_1, \mathbf{g}_2 \in \mathbb{R}^q$ with i.i.d. $N(0, \frac{1}{q})$ entries, such that

$$(\mathbf{u}_1, \mathbf{u}_2) = \left(\frac{\mathbf{g}_1}{\|\mathbf{g}_1\|}, \frac{\mathbf{g}_2 - \langle \mathbf{g}_2, \mathbf{g}_1 \rangle \frac{\mathbf{g}_1}{\|\mathbf{g}_1\|^2}}{\|\mathbf{g}_2 - \langle \mathbf{g}_2, \mathbf{g}_1 \rangle \frac{\mathbf{g}_1}{\|\mathbf{g}_1\|^2}\|} \right). \quad (\text{A.1})$$

Using Lemma 8.2, it is elementary to see that $\|\mathbf{g}_i\|^2 - 1 = O(\frac{(\log n)^K}{n})$ and

$$\|\mathbf{g}_i\| = 1 - \frac{1}{2}(\|\mathbf{g}_i\|^2 - 1) + O(\frac{(\log n)^K}{n}), \quad \mathbf{g}_1 A \mathbf{g}_2' = O(\frac{(\log n)^K}{\sqrt{n}}) \quad (\text{A.2})$$

hold with overwhelming probability, for any $q \times q$ matrix A independent of \mathbf{g}_i 's with $\|A\|_{\text{HS}} \leq C\sqrt{n}$. Hence, for any $\mathbf{d} \in \mathbb{R}^q$ independent of \mathbf{u}_i 's, with $\|\mathbf{d}\| \leq C$, we have

$$\mathbf{u}_i \mathbf{d}' = \left(1 - \frac{1}{2}(\|\mathbf{g}_i\|^2 - 1) + O(\frac{(\log n)^K}{n})\right) \mathbf{g}_i \mathbf{d}' = \mathbf{g}_i \mathbf{d}' + O(\frac{(\log n)^K}{n}), \quad (\text{A.3})$$

with overwhelming probability. This proves the first approximation in Lemma 8.1 if we regard the columns of $\mathcal{B}W_1'$ as \mathbf{d}' above.

For the second approximation in Lemma 8.1, we again use (A.1) and (A.2) to see that

$$\begin{aligned} \mathbf{u}_i \mathcal{D} \mathbf{u}_i' &= \left(1 - \frac{1}{2}(\|\mathbf{g}_i\|^2 - 1) + O(\frac{(\log n)^K}{n})\right)^2 \mathbf{g}_i \mathcal{D} \mathbf{g}_i', \\ \mathbf{u}_1 \mathcal{D} \mathbf{u}_2 &= (1 + O(\frac{(\log n)^K}{\sqrt{n}})) \mathbf{g}_1 \mathcal{D} \mathbf{g}_2' - (1 + O(\frac{(\log n)^K}{\sqrt{n}})) \mathbf{g}_1 \mathbf{g}_2' \mathbf{g}_1 D \mathbf{g}_1'. \end{aligned} \quad (\text{A.4})$$

Hence, with overwhelming probability, we have

$$\begin{aligned} \mathbf{u}_i \mathcal{D} \mathbf{u}_i' &= \mathbf{g}_i \mathcal{D} \mathbf{g}_i' - (\|\mathbf{g}_i\|^2 - 1) \frac{1}{q} \text{tr } \mathcal{D} + O(\frac{(\log n)^K}{n}), \\ \mathbf{u}_1 \mathcal{D} \mathbf{u}_2 &= \mathbf{g}_1 \mathcal{D} \mathbf{g}_2' - \mathbf{g}_1 \mathbf{g}_2' \frac{1}{q} \text{tr } \mathcal{D} + O(\frac{(\log n)^K}{n}). \end{aligned}$$

This proves the second approximation in Lemma 8.1 in case of $k = 2$.

For more general fixed k , it suffices to write $(\mathbf{u}_1, \dots, \mathbf{u}_k)$ in terms of $(\mathbf{g}_1, \dots, \mathbf{g}_k)$, using Gram-Schmidt orthogonalization. The proof is analogous. We omit the details. Hence, we completed the proof of Lemma 8.1. \square

PROOF OF LEMMA 8.4. Recall (8.8). It is elementary to check the first estimate of (8.24). For the second estimate, we use the fact

$$\Phi E = \left((E + H)^{-1} E - z \right)^{-1} (E + H)^{-1} E = (C_{w_2 y} - z)^{-1} C_{w_2 y}, \quad (\text{A.5})$$

from which we have

$$\frac{1}{n} \text{tr } E \Phi = \frac{1}{n} \text{tr } (C_{w_2 y} - z)^{-1} C_{w_2 y} = c_1 + z \frac{1}{n} \text{tr } (C_{w_2 y} - z)^{-1} + O(\frac{1}{n}). \quad (\text{A.6})$$

Recall $\check{s}(z)$ defined in (5.5). Now we claim that for any small constant $\varepsilon > 0$,

$$\check{s}_n(z) := \frac{1}{p} \text{tr} (C_{w_2 y} - z)^{-1} = \check{s}(z) + O_p(n^{-1+\varepsilon}). \quad (\text{A.7})$$

for any $z \in \mathcal{D}$. This together with (5.5) and (A.6) implies that

$$\frac{1}{n} \text{tr} E\Phi = s(z) + O_p(n^{-1+\varepsilon}) \quad (\text{A.8})$$

for any $z \in \mathcal{D}$.

Hence, what remains is to show (A.7). To this end, we first focus on a sub domain of \mathcal{D} :

$$\mathcal{D}_+ \equiv \mathcal{D}_+(\delta) := \left\{ z \in \mathbb{C} : d_+ + \delta < \Re z \leq 2, \frac{1}{n} \leq |\Im z| \leq 1 \right\}$$

Note that for $z \in \mathcal{D}_+$, the imaginary part of z is away from 0 by at least $\frac{1}{n}$. Then, we have the trivial bound

$$\sup_{z \in \mathcal{D}_+} \|(C_{w_2 y} - z)^{-1}\| \leq n, \quad (\text{A.9})$$

deterministically. Further, since the event Ξ_1 (c.f. (5.6)) holds with overwhelming probability, it is easy to see that

$$\sup_{z \in \mathcal{D}} \|(C_{w_2 y} - z)^{-1}\| \leq C, \quad (\text{A.10})$$

with overwhelming probability.

The deterministic bound and the high probability event (A.10) together implies that

$$\sup_{z \in \mathcal{D}_+} \mathbb{E} \|(C_{w_2 y} - z)^{-1}\|^\ell \leq C_\ell \quad (\text{A.11})$$

for some positive constant C_ℓ , for any given $\ell \in \mathbb{N}$. Further, we introduce a smooth cutoff function

$$\mathcal{Q} \equiv \mathcal{Q}(E, H) = \chi\left(\frac{1}{p} \text{tr} (E + H)^{-1}\right) = \chi\left(\frac{1}{p} \text{tr} S_{w_2 w_2}^{-1}\right), \quad (\text{A.12})$$

where $\chi(x) = 1$ if $0 \leq x \leq n^\varepsilon$, while $\chi(x) = 0$ if $x \geq n^\varepsilon + 1$, and $|\chi'(x)| \leq C$ for all $x \in [n^\varepsilon, n^\varepsilon + 1]$. From (3.3), we see that $\frac{1}{p} \text{tr} S_{w_2 w_2}^{-1} = O(1)$ with overwhelming probability. Hence, $\mathcal{Q} = 1$ with overwhelming probability. In turn, we see that $\mathcal{Q} = 1$ implies deterministically that $\|S_{w_2 w_2}^{-1}\| = O(n^{1+\varepsilon})$.

Now, to prove (A.7) for a given $z \in \mathcal{D}_+$, it suffices to show the following two

$$\text{Var}(\check{s}_n(z)\mathcal{Q}) = O\left(\frac{1}{n^2}\right), \quad \mathbb{E}(\check{s}_n(z)\mathcal{Q}) = \check{s}(z) + o\left(\frac{1}{\sqrt{n}}\right) \quad (\text{A.13})$$

since $\mathcal{Q} = 1$ with overwhelming probability. We first show the variance bound above. We write

$$E = \mathfrak{X}\mathfrak{X}^*, \quad H = \mathfrak{Y}\mathfrak{Y}^*, \quad \mathfrak{X} = (x_{ij})_{p-k,q}, \quad \mathfrak{Y} = (y_{ij})_{p-k,n-q},$$

and all x_{ij} 's and y_{ij} 's are i.i.d. $N(0, \frac{1}{n})$. We then regard $\check{s}_n(z)\mathcal{Q} = \frac{1}{p} \text{tr} E\Phi\mathcal{Q}$ as a function of the Gaussian variables x_{ij} 's and y_{ij} 's. Using Poincaré inequality for Gaussian, we have

$$\text{Var}(\check{s}_n(z)) \leq \frac{1}{n} \mathbb{E} \left(\sum_{i,j} \left| \frac{\partial(\frac{1}{p} \text{tr} E\Phi(z))\mathcal{Q}}{\partial x_{ij}} \right|^2 + \sum_{\alpha,\beta} \left| \frac{\partial(\frac{1}{p} \text{tr} E\Phi(z))\mathcal{Q}}{\partial y_{\alpha\beta}} \right|^2 \right). \quad (\text{A.14})$$

Observe that

$$\begin{aligned} \frac{\partial((\frac{1}{p}\text{tr } E\Phi)\mathcal{Q})}{\partial x_{ij}} &= \frac{1}{p} \left((\mathfrak{X}\Phi)_{ji} + (\Phi\mathfrak{X})_{ij} + z(\mathfrak{X}\Phi E\Phi)_{ji} + z(\Phi E\Phi\mathfrak{X})_{ij} \right) \mathcal{Q} \\ &\quad - \frac{1}{p^2} \text{tr } E\Phi \chi' \left(\frac{1}{p} \text{tr } S_{w_2 w_2}^{-1} \right) \left((\mathfrak{X}^* S_{w_2 w_2}^{-2})_{ji} + (S_{w_2 w_2}^{-2} \mathfrak{X})_{ij} \right) \end{aligned}$$

Then it is not difficult to see that

$$\begin{aligned} \frac{1}{n} \mathbb{E} \left(\sum_{i,j} \left| \frac{\partial \frac{1}{p} \text{tr } E\Phi(z)}{\partial x_{ij}} \right|^2 \right) &\leq \frac{C}{n^2} \mathbb{E} \left(\left(\sum_{\alpha=1}^3 \|E\|^\alpha \|\Phi\|^{\alpha+1} \right) \mathcal{Q}^2 \right) \\ &\quad + \frac{C}{n^2} \mathbb{E} \left(\left(\|E\|^2 \|S_{w_2 w_2}^{-1}\|^4 \right) \left(\left(\frac{1}{p} \text{tr } E\Phi \right) \chi' \left(\frac{1}{p} \text{tr } S_{w_2 w_2}^{-1} \right) \right)^2 \right) \end{aligned} \quad (\text{A.15})$$

for some positive constant C . Note that $\mathcal{Q} \neq 0$ only when $\frac{1}{p} \text{tr } S_{w_2 w_2}^{-1} \leq n^\varepsilon + 1$, and similarly $\chi' \left(\frac{1}{p} \text{tr } S_{w_2 w_2}^{-1} \right) \neq 0$ only if $\frac{1}{p} \text{tr } S_{w_2 w_2}^{-1} \in [n^\varepsilon, n^\varepsilon + 1]$. Hence, if the random variables in two expectations are nonzero, we have the bound $\|S_{w_2 w_2}^{-1}\| = O(n^{1+\varepsilon})$. In addition, we also know from (3.3) that $\|S_{w_2 w_2}^{-1}\| = O(1)$ holds with overwhelming probability. These facts imply that

$$\mathbb{E} \|S_{w_2 w_2}^{-1}\|^a \mathcal{Q}^b = O(1), \quad (\text{A.16})$$

for any fixed $a, b > 0$. The same bound holds if we replace \mathcal{Q} by $\chi' \left(\frac{1}{p} \text{tr } S_{w_2 w_2}^{-1} \right)$. Furthermore, it is well known that

$$\mathbb{E} \|E\|^\ell = O(1) \quad (\text{A.17})$$

for any given $\ell > 0$. Then combining the first inequality in (8.19), with (A.11), (A.16) and (A.17), we can obtain from (A.15) that

$$\frac{1}{n} \mathbb{E} \left(\sum_{i,j} \left| \frac{(\partial \frac{1}{p} \text{tr } E\Phi(z)) \mathcal{Q}}{\partial x_{ij}} \right|^2 \right) \leq \frac{C}{n^2} \mathbb{E} \left(\left(\sum_{\alpha=1}^3 \|E\|^\alpha \|\Phi\|^{\alpha+1} \right) \mathcal{Q}^2 \right) = O\left(\frac{1}{n^2}\right)$$

uniformly in $z \in \mathcal{D}_+$. For the second term in (A.14), we can get the same bound analogously. This completes the proof of the variance estimate in (A.13).

The estimate of mean in (A.13) was obtained in [2] for all

$$z \in \{w : \Re w \in [d_- - c, d_+ + c], \Im w \geq c\}$$

for any small positive constant c . The proof in [2] take the advantages of the lower bound $\Im z \geq c$, which guarantees an order 1 upper bound of $\|(C_{w_2 y} - z)^{-1}\|$. Here for $z \in \mathcal{D}_+$, we have (A.11) instead. This high order moment bound is sufficient for us to extend the discussion in [2] to the region \mathcal{D}_+ easily. Hence we omit the proof of (A.13), and refer to [2] for the interested readers.

Furthermore, using Chebyshev's inequality, (A.13) and the fact that $\mathcal{Q} = 1$ with overwhelming probability, we see that for any $z \in \mathcal{D}_+$

$$\mathbb{P}(|\check{s}_n(z) - \check{s}(z)| \geq n^{-\varepsilon}) = O(n^{-2+2\varepsilon}).$$

Now, we define the lattice $\mathcal{S}_+ = \mathcal{D} \cap (n^{-\varepsilon} \mathbb{Z} \times n^{-\varepsilon} i\mathbb{Z})$, where $\mathbb{Z} = \{\pm 1, \pm 2, \dots\}$. Then apparently $\sharp(\mathcal{S}_+) = O(n^{2\varepsilon})$. This implies that

$$\mathbb{P} \left(\bigcup_{z \in \mathcal{S}_+} \{|\check{s}_n(z) - \check{s}(z)| \geq n^{-\varepsilon}\} \right) = O(n^{-2+4\varepsilon}).$$

In addition, for any $z \in \mathcal{D}$, one can find a $z' \in \mathcal{S}_+$ such that $|z - z'| = O(n^{-\varepsilon})$. Then by continuity of $\check{s}_n(z)$ and $\check{s}(z)$ on \mathcal{D} , i.e., $|\check{s}'_n(z)|, |\check{s}'(z)| \leq \eta^{-2}$, we can easily get

$$\mathbb{P}\left(\bigcup_{z \in \mathcal{D}} \{|\check{s}_n(z) - \check{s}(z)| \geq Cn^{-\varepsilon}\}\right) = O(n^{-2+4\varepsilon}).$$

By choosing ε sufficiently small, we can then show that the second estimate in (8.24) holds uniformly in \mathcal{D} if we replace $o_p(\frac{1}{\sqrt{n}})$ by $O_{a.s.}(n^{-\varepsilon})$.

Next, we show the third estimate in (8.24). Recall (8.8). We write

$$\mathcal{H} = \mathcal{X}\mathcal{X}^*, \quad \mathcal{E} = \mathcal{Y}\mathcal{Y}^*, \quad \mathcal{X} = (\tilde{x}_{ij})_{q, n-p+k}, \quad \mathcal{Y} = (\tilde{y}_{ij})_{q, p-k} \quad (\text{A.18})$$

and all \tilde{x}_{ij} 's and \tilde{y}_{ij} 's are i.i.d. $N(0, \frac{1}{n})$. Analogously to the proof of the second estimate in (8.24), we can prove the result for any fixed $z \in \mathcal{D}_+$ at first, then extend the proof to whole \mathcal{D} by continuity. For a fixed $z \in \mathcal{D}_+$, we again split the proof into two parts

$$\text{Var}\left(\frac{1}{q} \text{tr} \Psi(z) \tilde{\mathcal{Q}}\right) = O\left(\frac{1}{n^2}\right), \quad (\text{A.19})$$

and

$$\mathbb{E} \frac{1}{q} \text{tr} \Psi(z) \tilde{\mathcal{Q}} = \varrho(z) + o\left(\frac{1}{\sqrt{n}}\right), \quad (\text{A.20})$$

where

$$\tilde{\mathcal{Q}}(\mathcal{E}, \mathcal{H}) = \chi\left(\frac{1}{q} \text{tr} (\mathcal{E} + \mathcal{H})^{-1}\right) = \chi\left(\frac{1}{q} \text{tr} S_{yy}^{-1}\right). \quad (\text{A.21})$$

Similarly, here we used the fact that $\tilde{\mathcal{Q}} = 1$ with overwhelming probability. The proof of (A.19) can be obtained by Poincaré inequality again. We omit the details. In the sequel, we show how to derive (A.20) by integration by parts formula of Gaussian measures. To ease the presentation, we get rid of the factor $\tilde{\mathcal{Q}}$ in the discussion, and do the reasoning as if $\|S_{yy}^{-1}\| = O(1)$ deterministically. The rigorous justification of the following derivation requires one to put $\tilde{\mathcal{Q}}$ back into each step.

Using the integration by parts formula for Gaussian variables (c.f. (9.17)), we arrive at

$$\begin{aligned} \frac{1}{q} \mathbb{E} \text{tr} \Psi(z) &= \frac{1}{q} \sum_{i=1}^q \sum_{j=1}^{n-p+k} \mathbb{E} \tilde{x}_{ij} (\mathcal{X}^* \Upsilon \mathcal{E})_{ji} = \frac{1}{qn} \sum_{i=1}^q \sum_{j=1}^{n-p+k} \mathbb{E} \frac{\partial (\mathcal{X}^* \Upsilon \mathcal{E})_{ji}}{\partial x_{ij}} \\ &= \frac{1}{qn} \mathbb{E} \sum_{i=1}^q \sum_{j=1}^{n-p+k} \left((\Upsilon \mathcal{E})_{ii} + z (\mathcal{X}^* \Upsilon \mathcal{X})_{jj} (\Upsilon \mathcal{E})_{ii} + z (\mathcal{X}^* \Upsilon)_{ji} (\mathcal{X}^* \Upsilon \mathcal{E})_{ji} \right) \\ &= \frac{n-p+k}{qn} \mathbb{E} \text{tr} (\mathcal{E} \Upsilon) + \frac{z}{qn} \mathbb{E} \text{tr} (\mathcal{H} \Upsilon) \text{tr} (\mathcal{E} \Upsilon) + \frac{z}{qn} \mathbb{E} \text{tr} (\mathcal{H} \Upsilon \mathcal{E} \Upsilon). \end{aligned} \quad (\text{A.22})$$

Note that for any $z \in \mathcal{D}_+$, one has

$$\frac{z}{qn} \mathbb{E} \text{tr} (\mathcal{H} \Upsilon \mathcal{E} \Upsilon) = O\left(\frac{1}{n}\right). \quad (\text{A.23})$$

Similarly to the variance bound in (A.13), it is not difficult to show

$$\text{Var}\left(\frac{1}{p} \text{tr} (\mathcal{H} \Upsilon)\right) = O\left(\frac{1}{n^2}\right), \quad \text{Var}\left(\frac{1}{p} \text{tr} (\mathcal{E} \Upsilon)\right) = O\left(\frac{1}{n^2}\right). \quad (\text{A.24})$$

To be precise, in order to obtain (A.23) and (A.24), the tracial quantities shall be multiplied by \tilde{Q} . But as we mentioned before, we omit it from the presentation. Combining (A.22)- (A.24), we have

$$\begin{aligned} \frac{1}{q} \mathbb{E} \text{tr} \Psi(z) &= \frac{n-p+k}{qn} \mathbb{E} \text{tr} (\mathcal{E} \Upsilon) + \frac{z}{qn} \mathbb{E} \text{tr} (\mathcal{H} \Upsilon) \mathbb{E} \text{tr} (\mathcal{E} \Upsilon) + O\left(\frac{1}{n}\right) \\ &= \frac{n-p+k}{qn} \mathbb{E} \text{tr} (\mathcal{E} \Upsilon) + \frac{1}{qn} \left((1-z) \mathbb{E} \text{tr} (\mathcal{E} \Upsilon) - q \right) \mathbb{E} \text{tr} (\mathcal{E} \Upsilon) + O\left(\frac{1}{n}\right) \\ &= \left(\frac{n-p+k}{qn} - \frac{1}{n} \right) \mathbb{E} \text{tr} (\mathcal{E} \Upsilon) + \frac{1-z}{qn} (\mathbb{E} \text{tr} (\mathcal{E} \Upsilon))^2 + O\left(\frac{1}{n}\right) \end{aligned}$$

where we used the fact

$$z \mathbb{E} \text{tr} (\mathcal{H} \Upsilon) = (1-z) \mathbb{E} \text{tr} (\mathcal{E} \Upsilon) - q.$$

Analogously to (A.13), we can show

$$\frac{1}{q} \mathbb{E} \text{tr} (\mathcal{E} \Upsilon) = \frac{z}{q} \mathbb{E} \text{tr} (C_{yw_2} - zI_q)^{-1} + 1 = z\tilde{s}(z) + 1 + o\left(\frac{1}{\sqrt{n}}\right),$$

where $\tilde{s}(z)$ is defined in (3.2). Then by (5.5), we have

$$\frac{1}{q} \mathbb{E} \text{tr} (\mathcal{E} \Upsilon) = \frac{1}{c_2} s(z) + o\left(\frac{1}{\sqrt{n}}\right). \quad (\text{A.25})$$

Combining the above results we can conclude that for any given $z \in \mathcal{D}_+$

$$\frac{1}{q} \mathbb{E} \text{tr} \Psi(z) = \frac{(1-c_1-c_2)s(z) + (1-z)s^2(z)}{c_2} + o\left(\frac{1}{\sqrt{n}}\right) = \varrho(z) + o\left(\frac{1}{\sqrt{n}}\right).$$

where we used (5.4). The extension of the result to the whole \mathcal{D} and the proof for the uniform bound are the same as the second estimate in (8.24). We thus omit the details.

This completes the proof of Lemma 8.4. □

PROOF OF LEMMA 9.1. Using the fact $((1-z)E - zH)\Phi(z) = I$ and the second estimate of Lemma 8.4, we see that

$$\frac{1}{n} \text{tr} (H\Phi(\gamma_j)) = \frac{1-\gamma_j}{\gamma_j n} \text{tr} E\Phi - \frac{p-k}{\gamma_j n} = \frac{-c_1 + (1-\gamma_j)s(\gamma_j)}{\gamma_j} + o_p\left(\frac{1}{\sqrt{n}}\right). \quad (\text{A.26})$$

Now, observe that

$$\begin{aligned} \frac{\partial}{\partial z} \left(\frac{1}{n} \text{tr} (H\Phi(z)) \right) \Big|_{z=\gamma_j} &= \frac{1}{n} \text{tr} (H\Phi(\gamma_j) W_2 W_2' \Phi(\gamma_j)) \\ &= -\frac{1}{\gamma_j n} \text{tr} H\Phi(\gamma_j) + \frac{1}{\gamma_j n} \text{tr} H\Phi(\gamma_j) E\Phi(\gamma_j) \\ &= -\frac{1}{1-\gamma_j} \frac{1}{n} \text{tr} H\Phi(\gamma_j) + \frac{1}{\gamma_j n} \text{tr} H\Phi(\gamma_j) H\Phi(\gamma_j). \end{aligned} \quad (\text{A.27})$$

Hence, to prove the second and third estimate in Lemma 9.1, it suffices to consider the LHS of (A.27). Although the limit and differentiation cannot be exchanged in general, here we can indeed obtain from (A.26) that

$$\frac{\partial}{\partial z} \left(\frac{1}{n} \text{tr} (H\Phi(z)) \right) \Big|_{z=\gamma_j} = \frac{d}{dz} \frac{1-z}{z} s(z) \Big|_{z=\gamma_j} + O_p(n^{-\frac{1}{4}}). \quad (\text{A.28})$$

To see this, it suffices to consider the non-limiting ratio

$$\frac{\frac{1}{n} \operatorname{tr} (H(\Phi(\gamma_j + n^{-\frac{1}{4}})) - \Phi(\gamma_j))}{n^{-\frac{1}{4}}} = \frac{1}{n} \operatorname{tr} (H\Phi(\gamma_j)W_2W_2')\Phi(\gamma_j) + O_p(n^{-\frac{1}{4}}), \quad (\text{A.29})$$

which follows by using the expansion (9.2) twice. Applying the estimate in (A.26) to the LHS of (A.29) leads to (A.28).

Therefore, combining (A.26), (A.27) and (A.28) yields the second and the third estimates in Lemma 9.1. To prove the last estimate in Lemma 9.1, we can consider the derivative of $\frac{1}{n} \operatorname{tr} E\Phi$ instead of $\frac{1}{n} \operatorname{tr} H\Phi$ and go through the above procedure. We omit the details.

This concludes the proof of Lemma 9.1. \square

PROOF OF LEMMA 9.2. From the identity (8.14), we have

$$\Psi = (1 - z)S_{yw_2}\Phi S_{w_2y} - \mathcal{E}.$$

Analogously to the proof of the third estimate in (8.16), we can use the spectral decomposition (6.4). Then we write

$$S_{yw_2}\Phi S_{w_2y} = U_y\Lambda_y V_y W_2' \Phi W_2 V_y' \Lambda_y' U_y'.$$

Again, similarly to the Gaussian approximation used in (8.1), we can approximate the α -th row and β -th column of U_y by Gaussian vectors. Then Lemma 8.2 leads us to the estimates

$$\Psi_{\alpha\beta}(\gamma_j) = \delta_{\alpha\beta} \frac{1}{q} \operatorname{tr} \Psi(\gamma_j) + O_p\left(\frac{(\log n)^K}{\sqrt{n}}\right) = \varrho(\gamma_j) + O_p\left(\frac{(\log n)^K}{\sqrt{n}}\right). \quad (\text{A.30})$$

where the last step follows from Lemma 8.4.

Next, for the second estimate in Lemma 9.2, we first notice that

$$\begin{aligned} \Psi(\gamma_j)(\mathcal{H}^{-1} + \mathcal{E}^{-1})\Psi(\gamma_j) &= \frac{\partial}{\partial z} \Psi(z) \Big|_{z=\gamma_j} = \frac{\partial}{\partial z} \left((1 - z)S_{yw_2}\Phi S_{w_2y} \right) \Big|_{z=\gamma_j} \\ &= -S_{yw_2}\Phi S_{w_2y} + (1 - z)S_{yw_2}\Phi S_{w_2w_2}\Phi S_{w_2y}. \end{aligned}$$

Again, applying the SVD of Y , we can derive

$$\left(\Psi(\gamma_j)(\mathcal{H}^{-1} + \mathcal{E}^{-1})\Psi(\gamma_j) \right)_{\alpha\beta} = \delta_{\alpha\beta} \frac{1}{q} \frac{\partial}{\partial z} \operatorname{tr} \Psi(z) \Big|_{z=\gamma_j} + O_p(n^{-\frac{1}{2}+\varepsilon})$$

Then similarly to (A.28), we have

$$\left(\Psi(\gamma_j)(\mathcal{H}^{-1} + \mathcal{E}^{-1})\Psi(\gamma_j) \right)_{\alpha\beta} = \delta_{\alpha\beta} \varrho'(\gamma_j) + o_p(1).$$

This concludes the proof of Lemma 9.2. \square

PROOF OF LEMMA 9.3. First, we estimate $\frac{1}{n} \operatorname{tr} \mathcal{A}^2$. From the definition in (4.14), we have

$$\begin{aligned} \frac{1}{n} \operatorname{tr} (\mathcal{A}(\gamma_j))^2 &= \frac{1}{n} \operatorname{tr} (P_y - \gamma_j)^2 + \frac{1}{n} \operatorname{tr} ((P_y - \gamma_j)^2 W_2' \Phi(\gamma_j) W_2)^2 \\ &\quad - \frac{2}{n} \operatorname{tr} ((P_y - \gamma_j)^3 W_2' \Phi(\gamma_j) W_2). \end{aligned} \quad (\text{A.31})$$

It is easy to check

$$\frac{1}{n} \operatorname{tr} (P_y - \gamma_j)^2 = (1 - 2\gamma_j)c_2 + \gamma_j^2 + o_p(1).$$

For the second term of (A.31), using Lemma 9.1 we obtain

$$\begin{aligned} \frac{1}{n} \text{tr} ((P_y - \gamma_j)^2 W_2' \Phi(\gamma_j) W_2)^2 &= \frac{1}{n} \text{tr} ((1 - \gamma_j)^2 \Phi(\gamma_j) E + \gamma_j^2 \Phi(\gamma_j) H)^2 \\ &= (1 - \gamma_j)(1 - 3\gamma_j)s(\gamma_j) + \gamma_j(1 - \gamma_j)^2 s'(\gamma_j) + \gamma_j^2 c_1 + o_p(1). \end{aligned}$$

Similarly, we have

$$\begin{aligned} \frac{1}{n} \text{tr} ((P_y - \gamma_j I_n)^3 W_2' \Phi(\gamma_j) W_2) &= \frac{1}{n} \text{tr} ((1 - \gamma_j)^3 \Phi(\gamma_j) E - \gamma_j^3 \Phi(\gamma_j) H) \\ &= (1 - \gamma_j)(1 - 2\gamma_j)s(\gamma_j) + \gamma_j^2 c_1 + o_p(1). \end{aligned}$$

Substituting the above results to (A.31), we get the first estimate in (9.24).

Next, we investigate $\frac{1}{q} \text{tr} \mathcal{B} \mathcal{B}'$. From the definition in (6.6), it is elementary to check

$$\text{tr} \mathcal{B} \mathcal{B}' = (1 - \gamma_j)^2 \text{tr} (S_{yy} - S_{yw_2} \Phi S_{w_2 y} + (\gamma_j - \gamma_j^2) S_{yw_2} \Phi S_{w_2 w_2} \Phi S_{w_2 y}). \quad (\text{A.32})$$

Recall (8.14). Then, with overwhelming probability, we have

$$\frac{1}{q} \text{tr} S_{yw_2} \Phi(\gamma_j) S_{w_2 y} = \frac{1}{1 - \gamma_j} \frac{1}{q} \text{tr} (\mathcal{E} + \Psi(\gamma_j)) = \frac{c_1 + \varrho(\gamma_j)}{1 - \gamma_j} + o_p(1), \quad (\text{A.33})$$

by (8.8) and Lemma 8.4.

Observe that

$$S_{yw_2} \Phi(z) S_{w_2 w_2} \Phi(z) S_{w_2 y} = \frac{\partial}{\partial z} S_{yw_2} \Phi(z) S_{w_2 y} = \frac{\partial}{\partial z} z S_{yy} \Upsilon(z) S_{yy}, \quad (\text{A.34})$$

where in the last step we used (8.6). Using (8.11) and (8.13), we see that

$$\begin{aligned} \frac{\partial}{\partial z} z S_{yy} \Upsilon(z) S_{yy} &= \frac{\partial}{\partial z} \left(\frac{z}{1 - z} \mathcal{E} - \mathcal{H} + \frac{1}{1 - z} \Psi(z) \right) \\ &= \frac{1}{(1 - z)^2} \mathcal{E} + \frac{1}{(1 - z)^2} \Psi(z) + \frac{1}{1 - z} \Psi(z) (\mathcal{E}^{-1} + \mathcal{H}^{-1}) \Psi(z). \end{aligned} \quad (\text{A.35})$$

Furthermore, from Lemma 9.2 we have

$$\begin{aligned} \frac{1}{q} \text{tr} \mathcal{E} &= c_1 + o_p(1), \quad \frac{1}{q} \text{tr} \Psi(\gamma_j) = \varrho(\gamma_j) + o_p(1), \\ \frac{1}{q} \text{tr} \Psi(\gamma_j) (\mathcal{E}^{-1} + \mathcal{H}^{-1}) \Psi(\gamma_j) &= \varrho'(\gamma_j) + o_p(1). \end{aligned} \quad (\text{A.36})$$

Combining (A.34), (A.35) and (A.36) yields

$$\frac{1}{q} \text{tr} S_{yw_2} \Phi(\gamma_j) S_{w_2 w_2} \Phi(\gamma_j) S_{w_2 y} = \frac{c_1 + \varrho(\gamma_j)}{(1 - \gamma_j)^2} + \frac{\varrho'(\gamma_j)}{1 - \gamma_j} + o_p(1). \quad (\text{A.37})$$

Therefore, we get the second estimate in (9.24) by substituting (A.33) and (A.37) into (A.32).

At the end, we prove the last estimate in Lemma 9.3. Note that

$$\begin{aligned} \frac{1}{q} \text{tr} \mathcal{D}_n^2 &= (1 - \gamma_j^2) \frac{1}{q} \text{tr} (S_{yy} - (1 - \gamma_j) S_{yw_2} \Phi S_{w_2 y})^2 \\ &= (1 - \gamma_j)^2 \frac{1}{q} \text{tr} (\mathcal{H} - \Psi)^2 = (1 - \gamma_j)^2 \left(\frac{1}{q} \text{tr} \mathcal{H}^2 - \frac{2}{q} \text{tr} \mathcal{H} \Psi + \frac{1}{q} \text{tr} \Psi^2 \right). \end{aligned} \quad (\text{A.38})$$

First, it is not difficult to check

$$\frac{1}{q} \text{tr } \mathcal{H}^2 \xrightarrow{\text{P}} \left(1 + \frac{c_2}{1 - c_1}\right) (1 - c_1)^2 = (1 - c_1 + c_2)(1 - c_1). \quad (\text{A.39})$$

We further claim that

$$\begin{aligned} \frac{1}{q} \text{tr } \mathcal{H} \Psi &= \frac{1 - c_1 - c_1 c_2}{c_2} (1 - \gamma_j) s(\gamma_j) + \frac{1}{c_2} (1 - \gamma_j)^2 s^2(\gamma_j) - c_1 (1 - c_1) + o_{\text{P}}(1), \\ \frac{1}{q} \text{tr } \Psi^2 &= \left(\frac{\gamma_j(\gamma_j - 1)}{c_2} - 2c_1(1 - \gamma_j) + \frac{1 - c_1 - c_2}{c_2} (1 - 2\gamma_j) \right) s(\gamma_j) \\ &\quad + \frac{(1 - \gamma_j)(2 - 3\gamma_j)}{c_2} s^2(\gamma_j) + \frac{\gamma_j(1 - \gamma_j)^2}{c_2} s'(\gamma_j) - (1 - c_1 - c_2)c_1 + o_{\text{P}}(1). \end{aligned} \quad (\text{A.40})$$

The proof of (A.40) will be postponed to the end. Consequently, we have

$$\begin{aligned} \frac{1}{q} \text{tr } \mathcal{D}_n^2 &= (1 - \gamma_j)^2 \left((1 - c_1 + c_2) + \left(\frac{\gamma_j(\gamma_j - 1)}{c_2} - 2(1 - \gamma_j) - \frac{1 - c_1 - c_2}{c_2} \right) s(\gamma_j) \right. \\ &\quad \left. - \frac{\gamma_j(1 - \gamma_j)}{c_2} s^2(\gamma_j) + \frac{\gamma_j(1 - \gamma_j)^2}{c_2} s'(\gamma_j) \right) + o_{\text{P}}(1). \end{aligned}$$

In addition, we see that

$$\frac{1}{q} \text{tr } \mathcal{D}_n = (1 - \gamma_j) \frac{1}{q} \text{tr } (\mathcal{H} - \Psi) = (1 - \gamma_j) \left(1 - \frac{1 - \gamma_j}{c_2} s(\gamma_j) \right) + o_{\text{P}}(1).$$

Then, by definition, we get the last estimate in (9.3).

Hence, what remains is to prove (A.40). Analogously to the proof of Lemma 8.4, by continuity, we can work with

$$\tilde{\gamma}_j = \gamma_j + i \frac{1}{n} \in \mathcal{D}_+$$

instead of γ_j . Recall $\tilde{\mathcal{Q}}$ defined in (A.21). Similarly to (A.13), also (A.19) and (A.20), we can split the tasks into two steps: first, we show that

$$\text{Var} \left(\left(\frac{1}{q} \text{tr } \mathcal{H} \Psi(\tilde{\gamma}_j) \right) \tilde{\mathcal{Q}} \right) = O\left(\frac{1}{n^2}\right), \quad \text{Var} \left(\left(\frac{1}{q} \text{tr } \Psi^2(\tilde{\gamma}_j) \right) \tilde{\mathcal{Q}} \right) = O\left(\frac{1}{n^2}\right) \quad (\text{A.41})$$

and then we estimate $\mathbb{E} \left(\left(\frac{1}{q} \text{tr } \mathcal{H} \Psi(\tilde{\gamma}_j) \right) \tilde{\mathcal{Q}} \right)$ and $\mathbb{E} \left(\left(\frac{1}{q} \text{tr } \Psi^2(\tilde{\gamma}_j) \right) \tilde{\mathcal{Q}} \right)$. The proof of (A.41) is analogous to the proof of (A.13). Hence, we omit it. In the sequel, we briefly state the estimates of $\mathbb{E} \left(\left(\frac{1}{q} \text{tr } \mathcal{H} \Psi(\tilde{\gamma}_j) \right) \tilde{\mathcal{Q}} \right)$ and $\mathbb{E} \left(\left(\frac{1}{q} \text{tr } \Psi^2(\tilde{\gamma}_j) \right) \tilde{\mathcal{Q}} \right)$, using integration by parts formula for Gaussian measure. Again, for brevity, we get rid of the factor $\tilde{\mathcal{Q}}$ in the discussion, and do the reasoning as if $\|S_{yy}^{-1}\| = O(1)$ deterministically. A rigorous justification of the following derivation requires one to put $\tilde{\mathcal{Q}}$ back into each step.

To ease the presentation, we omit $\tilde{\gamma}_j$ from the notations in the following discussions. First, from (9.8) we have

$$\frac{1}{q} \text{tr } \mathcal{H} \Psi = \frac{1}{q} \text{tr } \mathcal{H}^2 \Upsilon \mathcal{E}, \quad \frac{1}{q} \text{tr } \Psi^2 = \frac{1}{q} \text{tr } (\mathcal{H} \Upsilon \mathcal{E})^2.$$

We use integration by parts (c.f. (9.17)) again. Recall (A.18). We have

$$\frac{1}{q} \mathbb{E} \text{tr } \mathcal{H}^2 \Upsilon \mathcal{E} = \frac{1}{q} \sum_{\alpha, \beta} \mathbb{E} x_{\alpha\beta} (\mathcal{X}' \mathcal{H} \Upsilon \mathcal{E})_{\beta\alpha}$$

$$\begin{aligned}
&= \frac{1}{qn} \sum_{\alpha, \beta} \mathbb{E} \left((\mathcal{H}\Upsilon\mathcal{E})_{\alpha\alpha} + (\mathcal{X}')_{\beta\alpha} (\mathcal{X}'\Upsilon\mathcal{E})_{\beta\alpha} + (\mathcal{X}'\mathcal{X})_{\beta\beta} (\Upsilon\mathcal{E})_{\alpha\alpha} \right. \\
&\quad \left. + \gamma_j ((\mathcal{X}'\mathcal{H}\Upsilon)_{\beta\alpha} (\mathcal{X}'\Upsilon\mathcal{E})_{\beta\alpha} + (\mathcal{X}'\mathcal{H}\Upsilon\mathcal{X})_{\beta\beta} (\Upsilon\mathcal{E})_{\alpha\alpha}) \right) \\
&= \frac{n-p+k}{n} \frac{1}{q} \mathbb{E} \text{tr} (\mathcal{H}\Upsilon\mathcal{E}) + \frac{1}{qn} \mathbb{E} \text{tr} \mathcal{H} \text{tr} \mathcal{E}\Upsilon + \gamma_j \frac{1}{qn} \mathbb{E} \text{tr} \mathcal{H}^2 \Upsilon \text{tr} \mathcal{E}\Upsilon + O\left(\frac{1}{n}\right) \\
&= \frac{n-p+k}{n} \frac{1}{q} \mathbb{E} \text{tr} (\mathcal{H}\Upsilon\mathcal{E}) + \frac{1}{qn} \mathbb{E} \text{tr} \mathcal{H} \mathbb{E} \text{tr} \mathcal{E}\Upsilon + \gamma_j \frac{1}{qn} \mathbb{E} \text{tr} \mathcal{H}^2 \Upsilon \mathbb{E} \text{tr} \mathcal{E}\Upsilon + O\left(\frac{1}{n}\right),
\end{aligned}$$

where the last step follows from the variance bounds $O(\frac{1}{n^2})$ for all the normalized tracial quantities above. The proofs of these variance bounds are again similar to the proof of the first bound in (A.13).

Now, note that three terms on the RHS above can be computed easily. More specifically, we first recall (A.25). Then, we also note that

$$\frac{1}{q} \mathbb{E} \text{tr} (\mathcal{H}\Upsilon\mathcal{E}) = \frac{1}{q} \mathbb{E} \text{tr} \Psi(\gamma_j) = \varrho(\gamma_j) + o(1) \quad (\text{A.42})$$

and

$$\begin{aligned}
\frac{1}{n} \mathbb{E} \text{tr} \mathcal{H}^2 \Upsilon(\gamma_j) &= \frac{1}{\gamma_j n} \mathbb{E} \text{tr} \mathcal{H} \Upsilon(\gamma_j) (\gamma_j \mathcal{H} - (1 - \gamma_j) \mathcal{E}) + \frac{1 - \gamma_j}{\gamma_j n} \mathbb{E} \text{tr} \mathcal{H} \Upsilon(\gamma_j) \mathcal{E} \\
&= -\frac{1}{\gamma_j n} \mathbb{E} \text{tr} \mathcal{H} + \frac{1 - \gamma_j}{\gamma_j n} \mathbb{E} \text{tr} \Psi(\gamma_j) = \frac{c_2(c_1 - 1)}{\gamma_j} + \frac{1 - \gamma_j}{\gamma_j} c_2 \varrho(\gamma_j) + o(1).
\end{aligned}$$

Hence, we have

$$\begin{aligned}
\frac{1}{q} \mathbb{E} \text{tr} \mathcal{H}^2 \Upsilon \mathcal{E} &= ((1 - c_1) + (1 - \gamma_j) s(\gamma_j)) \varrho(\gamma_j) + O\left(\frac{1}{n}\right) \\
&= \frac{1 - c_1 - c_1 c_2}{c_2} (1 - \gamma_j) s(\gamma_j) + \frac{1}{c_2} (1 - \gamma_j)^2 s^2(\gamma_j) - c_1 (1 - c_1) + o(1).
\end{aligned}$$

Analogously, after a tedious calculation using integration by parts, we will get

$$\begin{aligned}
\frac{1}{q} \mathbb{E} \text{tr} \Psi^2 &= \frac{1}{q} \sum_{\alpha, \beta} \mathbb{E} x_{\alpha\beta} \left(\mathcal{X}' \Upsilon \mathcal{E} \mathcal{H} \Upsilon \mathcal{E} \right)_{\beta\alpha} \\
&= \frac{n-p+k}{n} \frac{1}{q} \mathbb{E} \text{tr} \mathcal{H} \Upsilon \mathcal{E} \Upsilon \mathcal{E} + \frac{1}{qn} \mathbb{E} \text{tr} \mathcal{H} \Upsilon \mathcal{E} \mathbb{E} \text{tr} \mathcal{E} \Upsilon + z \frac{1}{qn} \mathbb{E} \text{tr} \mathcal{H} \Upsilon \mathbb{E} \text{tr} \mathcal{H} \Upsilon \mathcal{E} \Upsilon \mathcal{E} \\
&\quad + z \frac{1}{qn} \mathbb{E} \text{tr} \mathcal{H} \Upsilon \mathcal{H} \Upsilon \mathcal{E} \mathbb{E} \text{tr} \mathcal{E} \Upsilon + O\left(\frac{1}{n}\right). \quad (\text{A.43})
\end{aligned}$$

The tracial quantities on the RHS can now be computed. For example, the terms like $\mathcal{H} \Upsilon \mathcal{E} \Upsilon \mathcal{E}$ can be obtained via taking derivative of $\mathcal{H} \Upsilon \mathcal{E}$. More specifically, we have

$$\begin{aligned}
\frac{\partial}{\partial z} \frac{1}{q} \mathbb{E} \text{tr} \mathcal{H} \Upsilon(z) \mathcal{E} \Big|_{z=\gamma_j} &= \frac{1}{q} \mathbb{E} \text{tr} \mathcal{H} \Upsilon(\gamma_j) (\mathcal{E} + \mathcal{H}) \Upsilon(\gamma_j) \mathcal{E} \\
&= \frac{1}{1 - \gamma_j} \frac{1}{q} \mathbb{E} \text{tr} \mathcal{H} \Upsilon(\gamma_j) \mathcal{E} + \frac{1}{1 - \gamma_j} \frac{1}{q} \mathbb{E} \text{tr} \mathcal{H} \Upsilon(\gamma_j) \mathcal{H} \Upsilon(\gamma_j) \mathcal{E} \\
&= -\frac{1}{\gamma_j} \frac{1}{q} \mathbb{E} \text{tr} \mathcal{H} \Upsilon(\gamma_j) \mathcal{E} + \frac{1}{\gamma_j} \frac{1}{q} \mathbb{E} \text{tr} \mathcal{H} \Upsilon(\gamma_j) \mathcal{E} \Upsilon(\gamma_j) \mathcal{E}.
\end{aligned}$$

From the above and (A.42), we get

$$\frac{1}{q} \mathbb{E} \text{tr} \mathcal{H} \Upsilon(\gamma_j) \mathcal{H} \Upsilon(\gamma_j) \mathcal{E} = (1 - \gamma_j) \varrho'(\gamma_j) - \varrho(\gamma_j) + o(1)$$

$$\begin{aligned}
&= -\frac{2(1-\gamma_j)}{c_2}s(\gamma_j) + \frac{(1-\gamma_j)^2}{c_2}s'(\gamma_j) + c_1 + o(1), \\
\frac{1}{q}\mathbb{E}\text{tr } \mathcal{H}\Upsilon(\gamma_j)\mathcal{E}\Upsilon(\gamma_j)\mathcal{E} &= \gamma_j\varrho'(\gamma_j) + \varrho(\gamma_j) + o(1) \\
&= \frac{1-2\gamma_j}{c_2}s(\gamma_j) + \frac{\gamma_j(1-\gamma_j)}{c_2}s'(\gamma_j) - c_1 + o(1).
\end{aligned}$$

Further, we recall (A.25), from which we also have

$$\frac{1}{q}\mathbb{E}\text{tr } (\mathcal{H}\Upsilon) = -\frac{1}{\gamma_j} + \frac{1-\gamma_j}{c_2\gamma_j}s(\gamma_j) + o(1).$$

Plugging all these estimates into (A.43), we get

$$\begin{aligned}
\frac{1}{q}\mathbb{E}\text{tr } \Psi^2(\gamma_j) &= \frac{2(1-\gamma_j)(1-2\gamma_j)}{c_2}s^2(\gamma_j) + \left(\frac{1-c_1-c_2}{c_2}\gamma_j(1-\gamma_j) + 2\frac{\gamma_j(1-\gamma_j)^2}{c_2}s(\gamma_j)\right)s'(\gamma_j) \\
&\quad + \left(-2c_1(1-\gamma_j) + \frac{1-c_1-c_2}{c_2}(1-2\gamma_j)\right)s(\gamma_j) + o(1).
\end{aligned} \tag{A.44}$$

Taking the derivative for (5.4), one can derive

$$2(\gamma_j - 1)s(\gamma_j)s'(\gamma_j) = -s^2(\gamma_j) + s(\gamma_j) - (c_1 + c_2 - \gamma_j)s'(\gamma_j) \tag{A.45}$$

Using (A.45) to (A.44), we obtain the second estimate in (A.40).

This completes the proof of Lemma 9.3. □

APPENDIX B

In this section, we prove Lemma 7.3. First, from Lemma 5.1, $(M_n)_{ij}(z)$ can be approximated by $\delta_{ij}m_i(z)$ up to an error $O(n^{-\varepsilon})$ if $z \in \mathcal{D}$. We now claim that this approximation also holds in the following domain \tilde{D} : we set for any $\tau > 0, \varepsilon_0$ the domain

$$\tilde{D} \equiv \tilde{D}(\delta, \varepsilon_0) := \left\{ z = x + i\eta : d_+ - \delta \leq x \leq 1 - \delta, n^{-1+\varepsilon_0} \leq \eta \leq 1 \right\}.$$

LEMMA B.1. *For any small positive constants δ and ε_0 , there exists some positive constant $c > 0$, such that*

$$\sup_{z \in \tilde{D}} \sup_{i,j=1,\dots,k} |(M_n)_{ij}(z) - \delta_{ij}m_i(z)| \leq n^{-c}$$

in probability.

For simplicity, in the sequel, we denote

$$\eta_0 = n^{-1+\varepsilon_0}.$$

To prove Lemma B.1, we need a local law for the Stieltjes transform of MANOVA matrix. Recall $\check{s}(z)$ from (3.1). We also recall the Stieltjes transform $\check{s}_n(z)$ from (A.7).

LEMMA B.2 (Local law of MONOVA at the right edge). *For any small $\delta > 0$ and $\varepsilon_0 > 0$, we have*

$$\left| \check{s}_n(z) - \check{s}(z) \right| \leq \frac{N^{\varepsilon'}}{N\eta} \tag{B.1}$$

uniformly on \tilde{D} with overwhelming probability, for any small constant $\varepsilon' > 0$.

PROOF. Observe that

$$\begin{aligned} ((E + H)^{-1}E - z)^{-1} &= ((1 - z)H^{-1}E - zI)^{-1}(I + H^{-1}E) \\ &= \frac{1}{(1 - z)^2} \left(H^{-1}E - \frac{z}{1 - z}I \right)^{-1} + \frac{1}{1 - z}I. \end{aligned}$$

Hence, by (8.9) and (A.7) we have

$$\check{s}_n(z) = \frac{1}{(1 - z)^2} \hat{s}_n\left(\frac{z}{1 - z}\right) + \frac{1}{1 - z} + O\left(\frac{1}{n}\right), \quad (\text{B.2})$$

where

$$\hat{s}_n(\omega) := \frac{1}{p} \text{tr} \left(H^{-1}E - \omega \right)^{-1}$$

is the Stieltjes transform of the F matrix $H^{-1}E$. Observe that $\Im(\frac{z}{1-z}) \sim \Im z$ when $\Re z \in [d_+ - \delta, 1 - \delta]$. A local law for $\hat{s}_n(\omega)$ at the right edge of $H^{-1}E$ with $\Im \omega \geq \eta_0$ has been proved in [26] (see Theorem 7.1 therein). Then by (B.2), Lemma B.2 follows immediately. \square

Denoting by $\tilde{\gamma}_i \equiv \tilde{\gamma}_{n,i}$ the i -th largest n -quantile of $f(x)$ in (1.6), i.e.,

$$\int_{d_-}^{\tilde{\gamma}_i} f(x) dx = \frac{n - i + 1}{n}. \quad (\text{B.3})$$

COROLLARY B.3 (Rigidity of eigenvalues near the right edge). *For any small δ in Lemma B.2, there exists some $c \equiv c(\delta) \in (0, 1)$ such that*

$$|\tilde{\lambda}_i - \tilde{\gamma}_i| \leq n^{\varepsilon'} n^{-\frac{2}{3}} i^{-\frac{1}{3}}, \quad \forall i = 1, \dots, \lfloor cn \rfloor. \quad (\text{B.4})$$

holds with overwhelming probability, for any small constant $\varepsilon' > 0$

PROOF. Denoting by $\hat{\lambda}_1 \geq \hat{\lambda}_2 \geq \dots \geq \hat{\lambda}_{p-k}$ the ordered eigenvalues of the F matrix $H^{-1}E$. Observe that $C_{w_2y} = (E + H)^{-1}E = H^{-1}E(H^{-1}E + I)^{-1}$. Hence, we have

$$\tilde{\lambda}_i = \frac{\hat{\lambda}_i}{1 + \hat{\lambda}_i}.$$

Further we set $\hat{\gamma}_i$ to be the solution to

$$\tilde{\gamma}_i = \frac{\hat{\gamma}_i}{1 + \hat{\gamma}_i}.$$

In [26], it is proved that $|\hat{\lambda}_1 - \hat{\gamma}_1| \leq n^{-\frac{2}{3} + \varepsilon'}$ with overwhelming probability. This implies directly that

$$|\tilde{\lambda}_1 - \tilde{\gamma}_1| \leq n^{-\frac{2}{3} + \varepsilon'}. \quad (\text{B.5})$$

with overwhelming probability. The rigidity for general i then follows from (B.5), Lemma B.2, and the square root decay of the density function in (1.6). The proof is standard, we omit the details. For interested readers, we refer to the proof of Theorem 2.2 in [17]. \square

With the aid of Lemma B.2, we are now at the stage to prove Lemma B.1.

PROOF OF LEMMA B.1. Here we only need to argue that the proof of Lemma 5.1 still works well for $z \in \tilde{D}$. It suffices to go through the proofs of Lemmas 8.1, 8.3 and 8.4 again, with the aid of Lemma B.2 and Corollary B.3.

First, analogously to Lemma 8.1, we will show that

$$\mathcal{M}_2 = \mathcal{T}G_1 \mathcal{B}W'_1 + O(n^{-\frac{1}{2}+2\varepsilon_0}), \quad \mathcal{M}_3 = \mathcal{T}G_1 \mathcal{D}G'_1 T'_1 + O(n^{-\frac{1}{2}+2\varepsilon_0}), \quad (\text{B.6})$$

hold with overwhelming probability on \tilde{D} . Here the second term on the RHS of the second equation in (8.15) has been absorbed into the error term $O(n^{-\frac{1}{2}+2\varepsilon_0})$. Similarly to the proof of Lemma 8.1, we only show the details for the case of $k = 2$. Recall (A.3) and (A.4). Denote by \mathbf{w}_ℓ the ℓ -th column of W_1 . It suffices to show that with overwhelming probability,

$$\mathbf{g}_1 \mathcal{D}(z) \mathbf{g}'_1 = O(n^{\varepsilon_0}), \quad \mathbf{g}_1 \mathcal{D}(z) \mathbf{g}'_2 = O(n^{\varepsilon_0}), \quad \mathbf{g}'_i \mathcal{B}(z) \mathbf{w}_j = O(n^{\varepsilon_0}) \quad (\text{B.7})$$

hold for $z \in \tilde{D}$. We only show the proof for the first two estimates above. The last one can be checked analogously. We use Lemma 8.2 again. For the first two estimates in (B.7), it suffices to show that

$$\frac{1}{n} \text{tr } \mathcal{D}(z) = O(n^{\varepsilon_0}), \quad \frac{1}{n^2} \|\mathcal{D}(z)\|_{HS}^2 = \frac{1}{n^2} \text{tr } \mathcal{D}(z) (\mathcal{D}(z))^* = O(n^{-\frac{\varepsilon_0}{2}}) \quad (\text{B.8})$$

hold with overwhelming probability on \tilde{D} . We start with the first estimate in (B.8). From the first equation of (8.21) and (8.3), we have

$$\begin{aligned} \left| \frac{1}{n} \text{tr } \mathcal{D}(z) \right| &\leq |1 - z| \left| \frac{1}{n} \text{tr } S_{yy} \right| + |1 - z|^2 \left| \frac{1}{n} \text{tr } S_{yw_2} \Phi(z) S_{w_2 y} \right| \\ &\leq C \left(1 + \frac{1}{n} \text{tr } |\tilde{C}_{w_2 y} - z| \right) \end{aligned}$$

with overwhelming probability, where we also used (3.3). Observe that $\tilde{C}_{w_2 y}$ has the same eigenvalues as $C_{w_2 y}$. Hence, we have on \tilde{D} ,

$$\frac{1}{n} \text{tr } |\tilde{C}_{w_2 y} - z| = \frac{1}{n} \sum_{i=1}^{p-k} \frac{1}{|\tilde{\lambda}_i - z|} = O((\log n)^K) \quad (\text{B.9})$$

with overwhelming probability, for some sufficiently large $K > 0$. The proof of (B.9) from Corollary B.3 is standard. We refer to the proof of (3.43) in [7] for a similar argument.

For the second estimate, we have

$$\frac{1}{n^2} \text{tr } \mathcal{D}(z) (\mathcal{D}(z))^* \leq |1 - z| \left| \frac{1}{n^2} \text{tr } S_{yy} (\mathcal{D}(z))^* \right| + |1 - z|^2 \left| \frac{1}{n^2} \text{tr } S_{yw_2} \Phi(z) S_{w_2 y} (\mathcal{D}(z))^* \right|$$

By Von Neumann's trace inequality, we can easily get

$$\left| \frac{1}{n^2} \text{tr } S_{yy} (\mathcal{D}(z))^* \right| \leq \frac{1}{n^2} \text{tr } S_{yy} \|(\mathcal{D}(z))^*\|_{\text{op}} \leq \frac{C}{n\eta_0} = Cn^{-\varepsilon_0}$$

with overwhelming probability. Analogously, using (3.3), (8.3) and (B.9), we have

$$\left| \frac{1}{n^2} \text{tr } S_{yw_2} \Phi(z) S_{w_2 y} (\mathcal{D}(z))^* \right| \leq \frac{C}{n^2 \eta_0} \text{tr } |\tilde{C}_{w_2 y} - z| \leq n^{-\frac{\varepsilon_0}{2}}$$

with overwhelming probability.

Next, we shall check that an analogue of Lemma 8.3 holds as well on \tilde{D} . Specifically, we shall show, that on \tilde{D} , the following estimates hold with overwhelming probability.

$$\sup_{i,j} \left| (\mathcal{M}_1)_{ij} - \delta_{ij} \left(c_2 - (1 + c_1)z - (1 - z) \frac{1}{n} \text{tr } (E\Phi) \right) \right| \leq n^{-\frac{\varepsilon_0}{8}},$$

$$\sup_{i,j} |(\mathcal{M}_2)_{ij}| \leq n^{-\frac{\varepsilon_0}{8}}, \quad \sup_{i,j} |(\mathcal{M}_3)_{ij} - \delta_{ij} t_i^2 (1-z) \frac{1}{q} \text{tr}(\mathcal{H} - \Psi)| \leq n^{-\frac{\varepsilon_0}{8}}. \quad (\text{B.10})$$

Similarly to the proof of Lemma 8.3, we start with (6.7) and (B.6) and use the large deviation inequality in Lemma 8.2. It suffices to show that

$$\frac{1}{n^2} \|\mathcal{A}\|_{HS}^2 \leq n^{-\frac{\varepsilon_0}{2}}, \quad \frac{1}{n^2} \|\mathcal{B}\|_{HS}^2 \leq n^{-\frac{\varepsilon_0}{2}}, \quad \frac{1}{n^2} \|\mathcal{D}\|_{HS}^2 \leq n^{-\frac{\varepsilon_0}{2}} \quad (\text{B.11})$$

holds on \tilde{D} with overwhelming probability. Note that the last estimate of (B.11) has already been proved in (B.8). The first two can be proved analogously. We thus omit the details.

At the end, we show that an analogue of Lemma 8.4 holds on the domain \tilde{D} as well. Apparently, the first estimate in (8.24) still holds. In the sequel, we will show that

$$\begin{aligned} \sup_{z \in \tilde{D}} \left| \frac{1}{n} \text{tr} E\Phi(z) - s(z) \right| &\leq n^{-\frac{\varepsilon_0}{2}}, \\ \sup_{z \in \tilde{D}} \left| \frac{1}{q} \text{tr} \Psi(z) - \varrho(z) \right| &\leq n^{-\frac{\varepsilon_0}{2}}. \end{aligned} \quad (\text{B.12})$$

with overwhelming probability. Note that the first estimate in (B.12) follows from (A.6) and Lemma B.2 by choosing $\varepsilon' = \frac{\varepsilon_0}{2}$ therein.

For the second estimate in (B.12), we first define

$$\mathfrak{P} \equiv \mathfrak{P}(z) := \frac{1}{n} \text{tr} S_{yy} \Upsilon(z) S_{yy} - \frac{1}{n} \text{tr} \Upsilon(z) S_{yy}. \quad (\text{B.13})$$

Our aim is to show that

$$\sup_{z \in \tilde{D}} |\mathfrak{P}(z)| \leq C n^{-\frac{1}{2}} \quad (\text{B.14})$$

with overwhelming probability.

We postpone the proof of (B.14) to the end, and show how to prove the second estimate in (B.12) with the aid of (B.14). First, from (8.6), (8.9) and (8.14), we can write

$$\Psi(z) = z(1-z) S_{yy} \Upsilon S_{yy} + (1-z) \mathcal{H} - z \mathcal{E}.$$

Hence, from (B.13) and (B.14), we have

$$\frac{1}{q} \text{tr} \Psi(z) = z(1-z) \frac{1}{q} \text{tr} \Upsilon(z) S_{yy} + (1-z) \frac{1}{q} \text{tr} \mathcal{H} - z \frac{1}{q} \text{tr} \mathcal{E} + O(n^{-\frac{1}{2}}) \quad (\text{B.15})$$

on \tilde{D} , with overwhelming probability. It is elementary to see that

$$\frac{1}{q} \text{tr} \mathcal{H} = 1 - c_1 + O(n^{-\frac{1}{2}+\varepsilon}), \quad \frac{1}{q} \text{tr} \mathcal{E} = c_1 + O(n^{-\frac{1}{2}+\varepsilon}) \quad (\text{B.16})$$

with overwhelming probability. Furthermore, according to (8.9) and (8.10), we can write

$$\frac{1}{q} \text{tr} \Upsilon(z) S_{yy} = \frac{1}{q} \text{tr} (C_{yw_2} - z)^{-1} =: \tilde{s}_n(z). \quad (\text{B.17})$$

Since C_{w_2y} and C_{yw_2} share the same nonzero eigenvalues, in light of (A.7), we have

$$\frac{c_1}{c_2} \tilde{s}_n(z) = \tilde{s}_n(z) + \frac{c_2 - c_1}{c_2} \frac{1}{z}. \quad (\text{B.18})$$

From (5.5), we also have

$$\frac{c_1}{c_2} \tilde{s}(z) = \tilde{s}(z) + \frac{c_2 - c_1}{c_2} \frac{1}{z}. \quad (\text{B.19})$$

Hence, combining (B.1), (B.18) and (B.19) yields

$$\sup_{z \in \tilde{\mathcal{D}}} \left| \tilde{s}_n(z) - \tilde{s}(z) \right| \leq \frac{N^{\varepsilon'}}{N\eta_0}. \quad (\text{B.20})$$

Plugging (B.17), (B.20) and (B.16) into (B.15), we can get (B.12) easily by choosing $\varepsilon' = \frac{\varepsilon_0}{2}$.

Hence, what remains is to prove (B.14). It suffices to show the one point estimate for any given $z \in \tilde{\mathcal{D}}$. The uniform bound follows from the continuity and the definition of overwhelming probability. More specifically, similarly to the proof of Lemma 8.4, we introduce the lattice $\tilde{\mathcal{S}} := \tilde{\mathcal{D}} \cap (n^{-3}\mathbb{Z} \times n^{-3}\mathbb{Z})$. According to the definition of with overwhelming probability, we see that if we can prove $|\mathfrak{P}(z)| \leq n^{-\frac{1}{2}}$ for any $z \in \tilde{\mathcal{S}}$ with overwhelming probability individually, we also have the uniform bound $\sup_{z \in \tilde{\mathcal{S}}} |\mathfrak{P}(z)| \leq n^{-\frac{1}{2}}$ with overwhelming probability. Moreover, for any $z' \in \tilde{\mathcal{D}}$, there is one $z \in \tilde{\mathcal{S}}$ such that $|z - z'| \leq n^{-3}$. Further, we have the following bound uniformly in all $z \in \tilde{\mathcal{D}}$

$$|\mathfrak{P}'(z)| \leq C \|\Upsilon'(z)\| = O\left(\frac{1}{\eta_0^2}\right) = O(n^2), \quad (\text{B.21})$$

conditioning on the event $\|S_{yy}\|, \|S_{yy}^{-1}\| \leq C$, which itself holds with overwhelming probability. By the continuity in (B.21), we can further obtain (B.14) from (B.21).

Therefore, in the sequel, we only show the proof of $|\mathfrak{P}(z)| \leq n^{-\frac{1}{2}}$ for one $z \in \tilde{\mathcal{D}}$. For brevity, we omit z from the notation. We estimate the high order moments of \mathfrak{P} . To this end, we set

$$\mathfrak{H}(m_1, m_2) = \mathfrak{P}^{m_1} \bar{\mathfrak{P}}^{m_2}.$$

With this notation, we can write

$$\mathbb{E}\mathfrak{H}(m, m) = \mathbb{E} \frac{1}{n} \text{tr} S_{yy} \Upsilon S_{yy} \mathfrak{H}(m-1, m) - \mathbb{E} \frac{1}{n} \text{tr} \Upsilon S_{yy} \mathfrak{H}(m-1, m). \quad (\text{B.22})$$

We apply the Gaussian integral by parts (c.f. (9.17)) to the first term above. Then we get

$$\begin{aligned} \mathbb{E} \left(\frac{1}{n} \text{tr} S_{yy} \Upsilon S_{yy} \mathfrak{H}(m-1, m) \right) &= \frac{1}{n} \sum_{i,j} \mathbb{E} \left(y_{ij} (Y' \Upsilon S_{yy})_{ji} \mathfrak{H}(m-1, m) \right) \\ &= \mathbb{E}(\mathfrak{c}_1 \mathfrak{H}(m-1, m)) + \mathbb{E}(\mathfrak{c}_2 \mathfrak{H}(m-2, m)) + \mathbb{E}(\mathfrak{c}_3 \mathfrak{H}(m-1, m-1)) \end{aligned} \quad (\text{B.23})$$

where

$$\begin{aligned} \mathfrak{c}_1 &= \frac{1}{n^2} \sum_{i,j} \frac{\partial (Y' \Upsilon S_{yy})_{ji}}{\partial y_{ij}}, & \mathfrak{c}_2 &= \frac{1}{n^2} \sum_{i,j} (Y' \Upsilon S_{yy})_{ji} \frac{\mathfrak{P}}{\partial y_{ij}}, \\ \mathfrak{c}_3 &= \frac{1}{n^2} \sum_{i,j} (Y' \Upsilon S_{yy})_{ji} \frac{\bar{\mathfrak{P}}}{\partial y_{ij}}. \end{aligned}$$

The estimates of \mathfrak{c}_2 and \mathfrak{c}_3 all nearly the same. We thus focus on the former. By elementary calculation, it is not difficult to derive

$$\mathfrak{c}_1 = \frac{1}{n^2} \sum_{i,j} \left((\Upsilon S_{yy})_{ii} - (Y' \Upsilon)_{ji} ((P_{w_2} - z) Y' \Upsilon S_{yy})_{ji} \right)$$

$$\begin{aligned}
& - (Y' \Upsilon Y (P_{w_2} - z))_{jj} (\Upsilon S_{yy})_{ii} + (Y' \Upsilon)_{ji} y_{ij} + (Y' \Upsilon Y)_{jj} \\
& = \frac{1}{n} \text{tr} \Upsilon S_{yy} - \frac{1}{n^2} \text{tr} \Upsilon Y (P_{w_2} - z) Y' \Upsilon S_{yy} \\
& \quad - \frac{1}{n^2} \text{tr} \Upsilon Y (P_{w_2} - z) Y' \text{tr} \Upsilon S_{yy} + \frac{1}{n^2} \text{tr} \Upsilon S_{yy} + \frac{q}{n^2} \text{tr} \Upsilon S_{yy} \\
& = \frac{1}{n} \text{tr} \Upsilon S_{yy}.
\end{aligned} \tag{B.24}$$

where in the last step we used the definition of Υ in (8.1). Similarly, we can derive

$$\mathfrak{c}_2 = \frac{2}{n^3} \text{tr} S_{yy}^2 \Upsilon S_{yy} \Upsilon = \frac{2}{n^3} \text{tr} S_{yy} (C_{yw_2} - z)^{-2} \tag{B.25}$$

Further, substituting (B.23) and (B.24) into (B.22), we have

$$\mathbb{E} \mathfrak{H}(m, m) = \mathbb{E}(\mathfrak{c}_2 \mathfrak{H}(m-2, m)) + \mathbb{E}(\mathfrak{c}_3 \mathfrak{H}(m-1, m-1))$$

Now, by Young's inequality, we have

$$\mathbb{E} \mathfrak{H}(m, m) \leq \frac{(\log n)^m}{m} (\mathbb{E} |\mathfrak{c}_2|^m + \mathbb{E} |\mathfrak{c}_3|^m) + \frac{2(m-1)}{m} \frac{1}{(\log n)^{\frac{m}{m-1}}} \mathbb{E} \mathfrak{H}(m, m).$$

This implies that

$$\mathbb{E} \mathfrak{H}(m, m) \leq C \frac{(\log n)^m}{m} (\mathbb{E} |\mathfrak{c}_2|^m + \mathbb{E} |\mathfrak{c}_3|^m). \tag{B.26}$$

Using (B.25), we see that

$$\mathbb{E} |\mathfrak{c}_2|^m \leq (\mathbb{E} \|S_{yy}\|^{2m})^{\frac{1}{2}} (\mathbb{E} (\frac{2}{n^3} \text{tr} |C_{yw_2} - z|^{-2})^{2m})^{\frac{1}{2}}. \tag{B.27}$$

Applying (B.20) and the fact $\tilde{s}(z) = O(1)$, we observe that

$$\frac{1}{n^3} \text{tr} |C_{yw_2} - z|^{-2} = \frac{1}{n^2 \eta} \frac{1}{n} \Im \text{tr} (C_{yw_2} - z)^{-1} = \frac{c_2}{n^2 \eta} \Im \tilde{s}_n(z) = O(n^{-1-\varepsilon_0}) \tag{B.28}$$

on \tilde{D} with overwhelming probability. Moreover, we also have the deterministic bound

$$\frac{1}{n^3} \text{tr} |C_{yw_2} - z|^{-2} = O(\frac{1}{n^2 \eta_0^2}) = O(n^{-2\varepsilon_0}).$$

Hence, we have

$$(\mathbb{E} (\frac{1}{n^3} \text{tr} |C_{yw_2} - z|^{-2})^{2m})^{\frac{1}{2}} \leq n^{-m(1+\frac{\varepsilon_0}{2})} \tag{B.29}$$

Using (3.3), we also have

$$(\mathbb{E} \|S_{yy}\|^{2m})^{\frac{1}{2}} \leq K^m \tag{B.30}$$

for some positive constant K . Plugging (B.29) and (B.30) into (B.27), we obtain

$$\mathbb{E} |\mathfrak{c}_2|^m \leq n^{-m(1+\frac{\varepsilon_0}{4})}.$$

The same bound holds for \mathfrak{c}_3 . Hence, we obtain from (B.26) that

$$\mathbb{E} |\mathfrak{P}|^{2m} = \mathbb{E} \mathfrak{H}(m, m) \leq n^{-m}.$$

Since m can be arbitrary fixed positive integer, we get $|\mathfrak{P}(z)| \leq n^{-\frac{1}{2}}$ with overwhelming probability for any $z \in \tilde{D}$.

This completes the proof of Lemma B.1. \square

With the aid of Lemma B.1, Lemma B.2, and Corollary B.3, we prove Lemma 7.3.

PROOF OF LEMMA 7.3. For any $z \in \Omega$, we set $z_0 = z + i\eta_0$, where $\eta_0 = n^{-1+\varepsilon_0}$. Our strategy is to compare $M_n(z)$ with $M_n(z_0)$, and use Lemma B.1 to conclude Lemma 7.3. We first show that

$$\sup_{z \in \Omega} |(M_n)_{ij}(z_0) - (M_n)_{ij}(z)| \leq n^{-c} \quad (\text{B.31})$$

with overwhelming probability. To see this, we first observe from (4.13) that

$$M_n(z_0) - M_n(z) = \mathcal{Z}_1(\mathcal{A}(z_0) - \mathcal{A}(z))\mathcal{Z}_1.$$

For $i = 0, \dots, 4$, let $\mathcal{A}_i(z_0, z)$ be the matrices obtained via replacing (λ_j, γ_j) by (z_0, z) in \mathcal{A}_i defined in (9.4). Similarly to (9.3), we have

$$M_n(z_0) - M_n(z) = i\eta_0 \sum_{a=0}^4 \mathcal{Z}_1 \mathcal{A}_a(z_0, z) \mathcal{Z}_1.$$

Since $\mathcal{A}_0(z_0, z) = -I_n$, apparently we have

$$\eta_0 (\mathcal{Z}_1 \mathcal{A}_0(z_0, z) \mathcal{Z}_1)_{ij} = O(n^{-1+\varepsilon_0})$$

with overwhelming probability, by simply using the definition of \mathcal{Z}_1 in (4.14) and Lemma 8.2. Hence, it suffices to show the estimate the entries of $\eta_0 \mathcal{Z}_1 \mathcal{A}_a(z_0, z) \mathcal{Z}_1$ for $a = 1, \dots, 4$. In the sequel, we only present the details for $a = 1$. The others can be handled similarly. From the definitions of \mathcal{Z}_1 (c.f. (4.14)) and $\mathcal{A}_1(z_0, z)$, it suffices to estimate the entries of

$$\eta_0 W_1 \mathcal{A}_1(z_0, z) W_1', \quad \eta_0 T_1 Y \mathcal{A}_1(z_0, z) W_1', \quad \eta_0 T_1 Y \mathcal{A}_1(z_0, z) Y' T_1'. \quad (\text{B.32})$$

For the first term, observe that W_1 is independent of $\mathcal{A}_1(z_0, z)$, and also the event Ω . Hence, by Lemma 8.2, we have

$$\eta_0 (W_1 \mathcal{A}_1(z_0, z) W_1')_{ij} = \frac{\delta_{ij}(1 - r_i)}{n^{2-\varepsilon_0}} \text{tr} \mathcal{A}_1(z_0, z) + O\left(\frac{(\log n)^K}{n^{2-\varepsilon_0}} \|\mathcal{A}_1(z_0, z)\|_{\text{HS}}\right)$$

uniformly on Ω , with overwhelming probability. Note that by the definition of the domain Ω in (7.3), we see that $\sup_{z \in \Omega} \|C_{w_2 y} - z\| \leq n^{1-\varepsilon}$. This together with (3.3), (8.3) and (B.9) implies that

$$\begin{aligned} \frac{1}{n^{2-\varepsilon_0}} |\text{tr} \mathcal{A}_1(z_0, z)| &= \frac{1}{n^{2-\varepsilon_0}} |\text{tr} P_y W_2' \Phi(z) W_2 W_2' \Phi(z_0) W_2 P_y| \\ &\leq \frac{1}{n^{1-\varepsilon_0+\varepsilon}} \text{tr} |\tilde{C}_{w_2 y} - z_0|^{-1} = O\left(\frac{(\log n)^K}{n^{\varepsilon-\varepsilon_0}}\right) \end{aligned}$$

holds with overwhelming probability, where in the second step we bounded all the matrices except $(\tilde{C}_{w_2 y} - z_0)^{-1}$ by their operator norms. Similarly, we have

$$\begin{aligned} \frac{(\log n)^{2K}}{n^{4-2\varepsilon_0}} \|\mathcal{A}_1(z_0, z)\|_{\text{HS}}^2 &\leq C \frac{(\log n)^{2K}}{n^{4-2\varepsilon_0}} \|\Phi(z)\|^2 \text{tr} \Phi(z_0) \Phi(\bar{z}_0) \\ &\leq C \frac{(\log n)^{2K}}{n^{2-2\varepsilon_0+2\varepsilon}} \text{tr} |\tilde{C}_{w_2 y} - z_0|^{-2} = O\left(\frac{(\log n)^K}{n^{2\varepsilon-3\varepsilon_0}}\right) \end{aligned}$$

with overwhelming probability, where in the last step we used (B.28). Choosing $\varepsilon > 3\varepsilon_0$ (say), we obtain

$$\eta_0 (W_1 \mathcal{A}_1(z_0, z) W_1')_{ij} = O(n^{-\varepsilon_0}) \quad (\text{B.33})$$

with overwhelming probability.

For the entries of the last two matrices in (B.32), we first do the approximation by choosing $\varepsilon > 3\varepsilon_0$ (say)

$$\begin{aligned}\eta_0 T_1 Y \mathcal{A}_1(z_0, z) W_1' &= \eta_0 \mathcal{T} G_1 \Lambda_y V_y \mathcal{A}_1(z_0, z) W_1' + O(n^{-\varepsilon_0}), \\ \eta_0 T_1 Y \mathcal{A}_1(z_0, z) Y' T_1' &= \eta_0 \mathcal{T} G_1 \mathcal{A}_1(z_0, z) G_1' \mathcal{T} + O(n^{-\varepsilon_0}).\end{aligned}\tag{B.34}$$

The proof of the above estimates is similar to (B.6). We omit the details. The estimates of the entries of the first term on the RHS of two equations in (B.34) can be derived in the same way as (B.33), using Lemma 8.2. We thus omit the details. Furthermore, the estimates of the entries of $\eta_0 \mathcal{Z}_1 \mathcal{A}_a(z_0, z) \mathcal{Z}_1$ for $a = 2, 3, 4$ are similar.

Hence, we completed the proof of (B.31). Observe that for any sufficiently small $\delta > 0$, $\Omega \subset [d_+ - \delta, d_+ + \delta] \subset [d_+ - \delta, 1 - \delta]$ almost surely, in light of (1.8). Hence, (B.31) together with Lemma B.1 implies that

$$\sup_{z \in \Omega} \sup_{i,j} |(M_n)_{ij} - \delta_{ij} m_i(z)| \leq n^{-c}$$

in probability, by using the continuity of the function $m_i(z)$. Further, when δ is sufficiently small, we also have

$$\sup_{z \in \Omega} \sup_i |m_i(z) - m_i(\delta_+)| \leq \varepsilon'$$

in probability, by the continuity of $m_i(z)$. This concludes the proof of Lemma 7.3. \square

REFERENCES

- [1] T. Anderson. (2003). An introduction to multivariate statistical analysis. Third Edition. Wiley New York.
- [2] Z. D. Bai, J. Hu, G.M. Pan, W. Zhou. (2015). Convergence of the empirical spectral distribution function of Beta matrices. *Bernoulli*, **21**(3), 1538-1574.
- [3] Z. Bai, Y. Fujikoshi, and K. P. Choi. High-dimensional consistency of AIC and BIC for estimating the number of significant components in principal component analysis. *The Annals of Statistics*, Accepted.
- [4] Z. D. Bai, J. Yao. (2008). Central limit theorems for eigenvalues in a spiked population model. *Ann. Inst. Henri Poincaré, Probab. Stat. Gauthier-Villars*, **44**(3), 447.
- [5] J. Baik, G. Ben Arous, S. Péché. (2005). Phase transition of the largest eigenvalue for nonnull complex sample covariance matrices. *Ann. Probab.*, 1643-1697.
- [6] J. Baik, J. W. Silverstein. (2006). Eigenvalues of large sample covariance matrices of spiked population models. *J. Multivariate Anal.*, **97**(6), 1382-1408.
- [7] Z.G. Bao, G.M. Pan, W. Zhou. (2013). Local density of the spectrum on the edge for sample covariance matrices with general population. Preprint. Available at <http://www.ntu.edu.sg/home/gmpan/publications.html>.
- [8] Z.G. Bao, J. Hu, G. Pan, and W. Zhou. (2017). Test of independence for high-dimensional random vectors based on freeness in block correlation matrices. *Electron. J. Statist.*, **11**(1):1527-1548.
- [9] S. T. Belinschi, H. Bercovici, M. Capitaine, and M. Février. (2012) Outliers in the spectrum of large deformed unitarily invariant models. Preprint.
- [10] F. Benaych-Georges, A. Guionnet, M. Maida. (2011). Fluctuations of the extreme eigenvalues of finite rank deformations of random matrices. *Electron. J. Probab.*, **16**(60), 1621-1662.
- [11] F. Benaych-Georges, N. R. Rao. (2011). The eigenvalues and eigenvectors of finite, low rank perturbations of large random matrices. *Adv. Math.*, **227**(1), 494-521.
- [12] M. Capitaine, C. Donati-Martin, D. Féral. (2009). The largest eigenvalues of finite rank deformation of large Wigner matrices: convergence and nonuniversality of the fluctuations. *Ann. Probab.*, 1-47.
- [13] M. Capitaine, C. Donati-Martin, D. Féral. (2009). Central limit theorems for eigenvalues of deformations of Wigner matrices. *Ann. Inst. Henri Poincaré, Probab. Stat.*, **48**(1), 107-133.
- [14] K. R. Davidson, S. J. Szarek. (2001). Local operator theory, random matrices and Banach spaces. *Handbook of the geometry of Banach spaces*, **1** (317-366): 131.
- [15] P. Dutilleul, B. Pelletier, and G. Alpargu. (2008). Modified tests for assessing the multiple correlation between one spatial process and several others. *Journal of Statistical Planning and Inference*, **138**(5):1402-1415.

- [16] A. Edelman, N. R. Rao. (2005). Random matrix theory. *Acta Numerica*, **14**, 233-297.
- [17] L. Erdős, H.-T. Yau, J. Yin. (2012). Rigidity of eigenvalues of generalized Wigner matrices. *Adv. Math.*, **229**(3): 1435-1515.
- [18] D. Féral, S. Péché. (2007). The largest eigenvalue of rank one deformation of large Wigner matrices. *Comm. Math. Phys.*, **272**(1), 185-228.
- [19] D. Féral, S. Péché. (2009). The largest eigenvalues of sample covariance matrices for a spiked population: diagonal case. *J. Math. Phys.*, **50** (7), 073302.
- [20] Y. Fujikoshi. (2016). High-Dimensional Asymptotic Distributions of Characteristic Roots in Multivariate Linear Models and Canonical Correlation Analysis. Technical report.
- [21] Y. Fujikoshi and T. Sakurai. (2016). High-dimensional consistency of rank estimation criteria in multivariate linear model. *Journal of Multivariate Analysis*, 149:199–212.
- [22] C. Gao, Z. Ma, Z. Ren, HH Zhou. (2015). Minimax estimation in sparse canonical correlation analysis. *The Annals of Statistics*. **43**(5), 2168-97.
- [23] C. Gao, Z. Ma, HH Zhou. (2014). Sparse CCA: Adaptive estimation and computational barriers. *arXiv preprint arXiv:1409.8565*.
- [24] R. Gittins. (1985). Canonical Analysis, volume 12 of *Biomathematics*. Springer Berlin Heidelberg, Berlin, Heidelberg.
- [25] X. Han, G. Pan, Q. Yang. (2016). A unified matrix model including both CCA and F matrices in multivariate analysis: the largest eigenvalue and its applications. *arXiv:1606.04417*.
- [26] X. Han, G. Pan, and B. Zhang. (2016). The Tracy-Widom law for the largest eigenvalue of F type matrices. *Annals of Statistics*, **44** (4): 1564–1592,
- [27] H. Hotelling. (1936). Relations between two sets of variates. *Biometrika*, 321-377.
- [28] M. Hyodo, N. Shutoh, T. Nishiyama, and T. Pavlenko. (2015). Testing block-diagonal covariance structure for high-dimensional data. *Statistica Neerlandica*, 69(4):460–482.
- [29] D. Jiang, Z. Bai, and S. Zheng. (2013). Testing the independence of sets of large-dimensional variables. *Science China Mathematics*, 56(1):135–147.
- [30] T. Jiang and F. Yang. (2013). Central limit theorems for classical likelihood ratio tests for high-dimensional normal distributions. *The Annals of Statistics*, 41(4):2029–2074.
- [31] I. M. Johnstone. (2001). On the distribution of the largest eigenvalue in principal components analysis. *Annals of statistics*, 295-327.
- [32] I. M. Johnstone. (2008). Multivariate analysis and Jacobi ensembles: Largest eigenvalue, Tracy-Widom limits and rates of convergence. *Ann. Statist.*, 36(6): 2638.
- [33] I. M. Johnstone. (2009). Approximate null distribution of the largest root in multivariate analysis. *The Annals of Applied Statistics*, 3(4):1616–1633.
- [34] I.M. Johnstone, A. Onatski. (2015). Testing in high-dimensional spiked models. *arXiv:1509.07269*.
- [35] V. Kargin. (2015). Subordination for the sum of two random matrices. *Ann. Probab.* **43** (4): 2119-2150.
- [36] B. Katz-Moses. (2012). Small Deviations for the beta-Jacobi Ensemble. PhD thesis.
- [37] A. Knowles, J. Yin. (2013). The isotropic semicircle law and deformation of Wigner matrices. *Comm. Pure Appl. Math.*, **66**(11),1663-1749.
- [38] R. J. Muirhead. (1982). *Aspects of Multivariate Statistical Theory*. Wiley, New York.
- [39] R. Oda, H. Yanagihara and Y. Fujikoshi (2016). Asymptotic non-null distributions of test statistics for redundancy in the high-dimensional canonical correlation analysis. Technical report.
- [40] D. Passemier and J. Yao. (2012). On determining the number of spikes in a high-dimensional spiked population model. *Random Matrices: Theory and Applications*, 01(01):1150002.
- [41] D. Paul. (2007). Asymptotics of sample eigenstructure for a large dimensional spiked covariance model. *Statist. Sinica*, **17** (4), 1617-1642.
- [42] S. Péché. (2006). The largest eigenvalue of small rank perturbations of Hermitian random matrices. *Probab. theory and related fields*, **134**(1), 127-173.
- [43] A. C. Rencher and F. C. Pun. (1980). Inflation of R^2 in Best Subset Regression. *Technometrics*, 22(1):49
- [44] A.N. Tikhomirov.(1981). On the convergence rate in the central limit theorem for weakly dependent random variables. *Theory Probab. Appl.*, **25**(4): 790-809.
- [45] K. W. Wachter. (1980). The limiting empirical measure of multiple discriminant ratios. *Ann. Statist.*, 937-957.
- [46] Q. Wang and J. Yao. (2017). Extreme eigenvalues of large-dimensional spiked Fisher matrices with application. *The Annals of Statistics*, 45(1):415–460.
- [47] Y. Yamada, M. Hyodo, and T. Nishiyama. (2017). Testing block-diagonal covariance structure for high-dimensional data under non-normality. *Journal of Multivariate Analysis*, 155(4):305–316.
- [48] Y. R. Yang, G. M. Pan. (2012) The convergence of the empirical distribution of canonical correlation coefficients. *Electron. J. Probab.*, **17**, 64.

- [49] Y. Yang and G. Pan. (2015). Independence test for high dimensional data based on regularized canonical correlation coefficients. *The Annals of Statistics*, 43(2):467–500.
- [50] S. Zheng, D. Jiang, Z. Bai, and X. He. (2014) Inference on multiple correlation coefficients with moderately high dimensional data. *Biometrika*, 101(3):748–754.

DEPARTMENT OF MATHEMATICS,
HONG KONG UNIVERSITY OF SCIENCE AND TECHNOLOGY,
HONG KONG
E-MAIL: mazgbao@ust.hk

KLASMOE AND SCHOOL OF MATHEMATICS & STATISTICS
NORTHEAST NORMAL UNIVERSITY
P. R. CHINA 130024
E-MAIL: huj156@nenu.edu.cn

DIVISION OF MATHEMATICAL SCIENCES,
NANYANG TECHNOLOGICAL UNIVERSITY,
SINGAPORE 637371
E-MAIL: gmpan@ntu.edu.sg

DEPARTMENT OF STATISTICS AND APPLIED PROBABILITY,
NATIONAL UNIVERSITY OF SINGAPORE,
SINGAPORE 117546
E-MAIL: stazw@nus.edu.sg
URL: <http://www.sta.nus.edu.sg/stazw/>