

# Functional Regression on Manifold with Contamination

BY ZHENHUA LIN

*Department of Statistics and Applied Probability, National University of Singapore,  
117546, Singapore  
stalz@nus.edu.sg*

AND FANG YAO

*Department of Probability and Statistics, School of Mathematical Sciences, Center for  
Statistical Science, Peking University, Beijing, 100871, China  
fyao@math.pku.edu.cn*

## SUMMARY

We propose a new method for functional nonparametric regression with a predictor that resides on a finite-dimensional manifold but is only observable in an infinite-dimensional space. Contamination of the predictor due to discrete/noisy measurements is also accounted for. By using functional local linear manifold smoothing, the proposed estimator enjoys a polynomial rate of convergence that adapts to the intrinsic manifold dimension and the contamination level. This is in contrast to the logarithmic convergence rate in the literature of functional nonparametric regression. We also observe a phase transition phenomenon regarding the interplay of the manifold dimension and the contamination level. We demonstrate that the proposed method has favorable numerical performance relative to commonly used methods via simulated and real data examples.

*Some key words:* Contaminated functional data; Functional nonparametric regression; Intrinsic dimension; Local linear manifold smoothing; Phase transition.

## 1. INTRODUCTION

Regression with a functional predictor is of central importance in the field of functional data analysis, the field that has been advanced by Ramsay & Silverman (1997, 2002) and many others. Early development of functional regression focuses on functional linear models (Cardot et al., 1999; Yao et al., 2005b; Yuan & Cai, 2010). Extensions of linear models include generalized linear regression (Cardot & Sarda, 2005; Müller & Stadtmüller, 2005), additive models (Müller & Yao, 2008), quadratic models (Yao & Müller, 2010), among others. These works prescribe specific forms of the regression model, and are regarded as functional parametric regression models (Ferraty & Vieu, 2006) that entail efficient estimation procedures and hence are well studied in the literature.

In contrast, functional nonparametric regression that does not impose structural constraints on the regression function has received less attention. The first landmark development of nonparametric functional data analysis is the monograph of Ferraty & Vieu (2006). Recent advances in this direction include the Nadaraya–Watson estimator (Ferraty et al., 2012) and the  $k$ -nearest-neighbor estimator (Kudraszow & Vieu, 2013). The development of functional nonparametric regression is hindered by a theoretical barrier that is formulated in Mas (2012) and is linked to the small ball probability problem (Delaigle & Hall, 2010). Essentially, in a rather general setting, the minimax rate of nonparametric regression on a generic functional space is slower than any polynomial of the sample size, which differs markedly from the polynomial minimax rates for many functional parametric regression procedures (e.g. Hall & Keilegom, 2007; Yuan & Cai, 2010, for functional linear regression). These endeavors on functional nonparametric regression do not exploit the intrinsic structure that is common in practice. For instance, Chen & Müller (2012) suggested that functional data often possess a low-dimensional manifold structure which can be utilized for more efficient representation. By contrast, we exploit the nonlinear low-dimensional structure for functional nonparametric regression.

Our method, which we call functional regression on manifold, assumes the model

$$Y = g(X) + \varepsilon, \tag{1}$$

where  $Y$  is a scalar response,  $X$  is a functional predictor sampled from an unknown manifold  $\mathcal{M}$ ,  $\varepsilon$  is the error term independent of  $X$ , and  $g$  is some unknown functional to be estimated. In reality, the functional predictor  $X$  is rarely fully observed. To accommodate

this common practice, we assume that  $X$  is recorded at a grid of points with noise. The model (1) features a manifold structure  $\mathcal{M}$  that underlies the functional predictor  $X$  and is assumed to be a finite-dimensional but potentially nonlinear submanifold of the function space  $\mathcal{L}^2(D)$ , the space of square integrable functions defined on a compact domain  $D \subset \mathbb{R}$ . For a background on both finite-dimensional and infinite-dimensional manifolds, we refer readers to Lang (1995) and Lang (1999).

Data analysis with a manifold structure has been extensively studied in the statistical literature. For example, techniques have been invented to learn an unknown manifold based on a point cloud, such as locally linear embedding (Roweis & Saul, 2000; Wu & Wu, 2018), isomap (Tenenbaum et al., 2000), t-SNE (van der Maaten & Hinton, 2008), among many others. Supervised learning on an unknown manifold has also been investigated, such as estimation of functions defined on a manifold (Aswani et al., 2011; Cheng & Wu, 2013; Sober et al., 2019) and estimation of the gradient of such functions (Mukherjee et al., 2010). In addition, data analysis on a known manifold has been studied, such as fundamentals related to the Fréchet mean (Bhattacharya & Patrangenaru, 2003, 2005; Bhattacharya & Lin, 2017), manifold-valued function estimation (Yuan et al., 2012; Lin et al., 2016; Cornea et al., 2017; Lin et al., 2019), manifold-valued principal component analysis (Huckemann et al., 2010; Panaretos et al., 2014), classification on manifolds (Yao & Zhang, 2019+), and nonparametric manifold-valued inference (Patrangenaru & Ellingson, 2015).

However, the literature specifically relating functional data to manifolds is scarce. Zhou & Pan (2014) investigated functional principal component analysis on an irregular domain. Chen & Müller (2012) and Lila & Aston (2016) considered the representation and principal component analysis of functional data sampled from a manifold. Manifold-valued random functions were studied by Su et al. (2014), Dai & Müller (2018) and Lin & Yao (2019). To the best of our knowledge, we are the first to consider a manifold structure in functional regression where a global representation of the low-dimensional functional predictor  $X$  can be inefficient. For illustration, Example 1 in Supplementary Material exhibits a random process taking values in a one-dimensional submanifold of  $\mathcal{L}^2([0, 1])$  while having an infinite number of components in its Karhunen–Loève expansion.

When estimating the regression functional  $g$  in (1), we explicitly account for the hidden manifold structure by estimating the tangent spaces of the manifold. Specifically, we

first recover the observed functional predictors from their discrete/noisy measurements, and then adopt the local linear manifold smoothing (Cheng & Wu, 2013). While our approach and the one of Cheng & Wu (2013) share the same intrinsic manifold setup, they fundamentally differ in the ambient aspect, which raises challenging issues unique to functional data. First, functional data naturally live in an infinite-dimensional ambient space, while the Euclidean data considered by Cheng & Wu (2013) have a finite ambient dimension. Second, the effect of noise/sampling in the observed functional data needs to be explicitly treated, since functional data are discretely and noisily recorded in practice, which then introduces contamination of the functional predictor. This contamination issue is not encountered in Cheng & Wu (2013), or is only considered for linear regression of multivariate data (Aswani et al., 2011; Loh & Wainwright, 2012). Moreover, the contamination has an intrinsic dimension that grows with the sample size and thus is coupled with the ambiently infinite dimensionality.

The main contributions of this article are as follows. First, by exploiting structural information of the predictor, our proposal entails an effective estimation procedure that adapts to the unknown manifold structure and the contamination level while maintains the flexibility of functional nonparametric regression. Second, by careful theoretical analysis, we confirm that the regression functional  $g$  can be estimated at a polynomial convergence rate of the sample size, especially when only the contaminated functional predictors are available. This provides a new angle to functional nonparametric regression that is subject to a logarithmic rate (Mas, 2012). Third, the contamination on predictors is explicitly treated and is shown to be an integrated part of the convergence rate, which has not been well studied even in classical functional linear regression (Hall & Keilegom, 2007). Finally, we discover that, the polynomial convergence rate exhibits a phase transition phenomenon, depending on the interplay between the manifold dimension and the contamination level. This type of phase transition has not yet been discovered in functional regression, and shares at least the same importance of those concerning estimation of mean/covariance functions (e.g. Cai & Yuan, 2011; Zhang & Wang, 2016). In addition, during our theoretical development, we obtain some results that are generally useful with their own merit, such as the consistency of the estimated intrinsic dimension and tangent spaces of the manifold in the presence of contamination.

## 2. ESTIMATION OF FUNCTIONAL REGRESSION ON MANIFOLD

## 2.1. Step I: Recovery of Functional Data

We assume that each predictor  $X_i$  is observed at  $m_i$  design points  $T_{i1}, \dots, T_{im_i} \in D$ . Denote the observed value at  $T_{ij}$  by  $X_{ij}^* = X_i(T_{ij}) + \zeta_{ij}$ , where  $\zeta_{ij}$  is random noise with mean zero and is independent of all  $X_i$  and  $T_{ij}$ . The collection  $\mathbb{X}_i = \{(T_{i1}, X_{i1}^*), \dots, (T_{im_i}, X_{im_i}^*)\}$  represents all measurements for the realization  $X_i$ , and  $\{\mathbb{X}_1, \dots, \mathbb{X}_n\}$  constitutes the observed data for the predictor. We shall clarify that, although each trajectory  $X_i$  as a whole function resides on the manifold  $\mathcal{M}$ , the  $m_i$ -dimensional vector  $\mathbb{V}_i = (X_i(T_{i1}), \dots, X_i(T_{im_i}))$  does not. Consequently, the manifold assumption in Cheng & Wu (2013) is violated for  $\mathbb{V}_i$ .

When  $\inf_i m_i$  is sufficiently large or grows with the sample size, a scenario commonly referred to as the dense design, we may recover each function  $X_i$  based on the observed data  $\mathbb{X}_i$  by individual smoothing estimation. Popular smoothing techniques include the local linear smoother (Fan, 1993) and spline smoothing (Ramsay & Silverman, 2005), among others. By applying one of these methods, we obtain an estimate  $\hat{X}_i$  of  $X_i$ , referred to as the contaminated version of  $X_i$  that is used in the subsequent steps to estimate  $g$ . To be specific, we consider the local linear estimate of  $X_i(t)$  given by  $\hat{b}_1$  with

$$(\hat{b}_1, \hat{b}_2) = \arg \min_{(b_1, b_2) \in \mathbb{R}^2} \frac{1}{m_i} \sum_{j=1}^{m_i} \left\{ X_{ij}^* - b_1 - b_2(T_{ij} - t) \right\}^2 K\left(\frac{T_{ij} - t}{h_i}\right), \quad (2)$$

where  $K$  is a compactly supported symmetric density function and  $h_i$  is the bandwidth. Calculation shows that  $\hat{b}_1 = (R_0 S_2 - R_1 S_1) / (S_0 S_2 - S_1^2)$ , where for  $r = 0, 1$  and 2,

$$S_r(t) = \frac{1}{m_i h_i} \sum_{j=1}^{m_i} K\left(\frac{T_{ij} - t}{h_i}\right) \left(\frac{T_{ij} - t}{h_i}\right)^r, R_r(t) = \frac{1}{m_i h_i} \sum_{j=1}^{m_i} K\left(\frac{T_{ij} - t}{h_i}\right) \left(\frac{T_{ij} - t}{h_i}\right)^r X_{ij}^*.$$

The estimate  $\hat{b}_1$  does not have a finite mean squared error, as its denominator is zero with a positive probability for a finite sample. To overcome this issue, we adopt the technique of ridging (Fan, 1993; Seifert & Gasser, 1996; Hall & Marron, 1997) to estimate  $X_i(t)$  by the following ridged local linear estimate

$$\hat{X}_i(t) = \frac{R_0 S_2 - R_1 S_1}{S_0 S_2 - S_1^2 + \delta 1_{\{|S_0 S_2 - S_1^2| < \delta\}}}, \quad (3)$$

where  $\delta > 0$  is a sufficiently small constant that depends on  $m_i$ , e.g.,  $\delta = m_i^{-2}$ .

When  $\sup_i m_i$  is relatively small or bounded by a constant, a scenario commonly referred to as the sparse design, to recover  $X_i$ , the procedure proposed by Yao et al. (2005a) can be adopted to recover individual functions. We refer readers to Supplementary Material for the details of such procedure.

## 2.2. Step II: Estimation of the Manifold Dimension and Tangent Space

To characterize the manifold structure, we shall first estimate the intrinsic dimension  $d$  of the manifold  $\mathcal{M}$ . We adopt the maximum likelihood estimator proposed by Levina & Bickel (2004), substituting the unobservable  $X_i$  with the contaminated version  $\hat{X}_i$ . For a given  $x \in \mathcal{M}$ , define  $\hat{G}_i(x) = \|x - \hat{X}_i\|_{\mathcal{L}^2}$  and let  $\hat{G}_{(k)}(x)$  be the  $k$ th order statistic of  $\hat{G}_1(x), \dots, \hat{G}_n(x)$ . Then the intrinsic dimension  $d$  is estimated by

$$\hat{d} = \frac{1}{k_2 - k_1 + 1} \sum_{k=k_1}^{k_2} \hat{d}_k, \quad (4)$$

with

$$\hat{d}_k = \frac{1}{n} \sum_{i=1}^n \hat{d}_k(\hat{X}_i), \quad \hat{d}_k(x) = \left\{ \frac{1}{k-1} \sum_{j=1}^{k-1} \log \frac{\hat{G}_{(k)}(x) + \Delta}{\hat{G}_{(j)}(x) + \Delta} \right\}^{-1}, \quad (5)$$

where  $\Delta$  is a positive constant depending on  $n$ , and  $k_1, k_2$  are tuning parameters. This  $\Delta$  regularizes  $\hat{d}_k(x)$  in order to overcome the additional variability introduced by the contamination on the predictor. We conveniently set  $\Delta = 1/\log \bar{m}$  with  $\bar{m} = n^{-1} \sum_{i=1}^n m_i$ , while refer readers to Levina & Bickel (2004) for the choice of  $k_1$  and  $k_2$ . When the observed data are sparsely sampled, the distance  $\hat{G}_i(x)$  can be better estimated by the procedure of Peng & Müller (2008).

Now we proceed to estimate the tangent space at the given point  $x$  as follows.

- A neighborhood of  $x$  is determined by a tuning parameter  $h_{pca} > 0$ , denoted by  $\hat{\mathcal{N}}_{\mathcal{L}^2}(h_{pca}, x) = \{\hat{X}_i : \|x - \hat{X}_i\|_{\mathcal{L}^2} < h_{pca}, i = 1, \dots, n\}$ .
- Compute the local empirical covariance function

$$\hat{\mathcal{C}}_x(s, t) = \frac{1}{|\hat{\mathcal{N}}_{\mathcal{L}^2}(h_{pca}, x)|} \sum_{\hat{X} \in \hat{\mathcal{N}}_{\mathcal{L}^2}(h_{pca}, x)} \{\hat{X}(s) - \hat{\mu}_x(s)\} \{\hat{X}(t) - \hat{\mu}_x(t)\} \quad (6)$$

and obtain the eigenfunctions  $\hat{\varphi}_1, \hat{\varphi}_2, \dots, \hat{\varphi}_{\hat{d}}$  corresponding to the first  $\hat{d}$  leading eigenvalues, where  $\hat{\mu}_x = |\hat{\mathcal{N}}_{\mathcal{L}^2}(h_{pca}, x)|^{-1} \sum_{\hat{X} \in \hat{\mathcal{N}}_{\mathcal{L}^2}(h_{pca}, x)} \hat{X}$  is the local mean function and  $|\hat{\mathcal{N}}_{\mathcal{L}^2}(h_{pca}, x)|$  denotes the number of observations in  $\hat{\mathcal{N}}_{\mathcal{L}^2}(h_{pca}, x)$ .

- Estimate the tangent space at  $x$  by  $\widehat{T_x\mathcal{M}} = \text{span}\{\hat{\varphi}_1, \hat{\varphi}_2, \dots, \hat{\varphi}_{\hat{d}}\}$ , the linear space spanned by the first  $\hat{d}$  estimated eigenfunctions.

### 2.3. Step III: Local Linear Regression on the Tangent Space

Finally, we utilize the local manifold structure by projecting all  $\hat{X}_i$  onto the estimated tangent space  $\widehat{T_x\mathcal{M}}$  and obtain the local coordinate  $\hat{\xi}_i = (\langle \hat{X}_i, \hat{\varphi}_1 \rangle, \dots, \langle \hat{X}_i, \hat{\varphi}_{\hat{d}} \rangle)^T$  for  $\hat{X}_i$ . Then, the estimate of  $g(x)$  is given by

$$\hat{g}(x) = e_1^T (\hat{Q}^T \hat{W} \hat{Q})^{-1} \hat{Q}^T \hat{W} \mathcal{Y}, \quad \hat{Q} = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ \hat{\xi}_1 & \hat{\xi}_2 & \cdots & \hat{\xi}_n \end{pmatrix}^T, \quad (7)$$

$\hat{W} = \text{diag}(K_{h_{reg}}(\|x - \hat{X}_1\|_{\mathcal{L}^2}), K_{h_{reg}}(\|x - \hat{X}_2\|_{\mathcal{L}^2}), \dots, K_{h_{reg}}(\|x - \hat{X}_n\|_{\mathcal{L}^2}))$  with  $K_h(t) = K(t/h)/h^{\hat{d}}$  and the bandwidth  $h_{reg}$ ,  $\mathcal{Y} = (Y_1, \dots, Y_n)^T$ , and  $e_1^T = (1, 0, \dots, 0)$  is an  $n \times 1$  vector. Here, the matrix  $\hat{Q}$  incorporates the estimated geometric structure that is encoded by the local eigenbasis  $\hat{\varphi}_1, \dots, \hat{\varphi}_{\hat{d}}$ . We emphasize that, in the above estimation procedure which is illustrated by the diagram in the left panel of Figure 1, all steps are based on the contaminated sample  $\{\hat{X}_1, \dots, \hat{X}_n\}$ , rather than the unavailable functions  $X_1, \dots, X_n$ . When the predictor  $x$  is also only measured at  $m_x$  discrete points  $t_1, \dots, t_{m_x}$ , we impute it by the procedures in Section 2.1, and replace  $x$  in (5)–(7) with the imputed curve  $\tilde{x}$  to obtain an estimate of  $g(\tilde{x})$ .

### 2.4. Tuning Parameter Selection

There are several tuning parameters to be determined in our estimation procedure. For the parameters  $k_1$  and  $k_2$  in (4) to estimate the intrinsic dimension,  $k_1 = 10$  and  $k_2 = 20$  are suggested by Levina & Bickel (2004). However, we found that  $k_1 = 20$  and  $k_2 = 30$  work better generally in our setting, perhaps partially due to the contamination that requires a relatively larger local neighborhood to offset.

For the individual smoothing presented in Section 2.1, we adopt the following leave-one-out cross-validation to select the bandwidth  $h_i$  (Fan & Gijbels, 1996; Lee & Solo, 1999). Let  $\hat{X}_{i,h,-j}(x)$  be the leave-one-out estimate of  $X_i(t)$ , i.e., the estimate computed according to (3) using all of  $(T_{i1}, X_{i1}^*), \dots, (T_{im_i}, X_{im_i}^*)$  but  $(T_{ij}, X_{ij}^*)$ . We then select  $h_i$  from a pool of candidates to minimize the cross-validation error  $\text{cv}(h) = \sum_{j=1}^{m_i} \{X_{ij}^* - \hat{X}_{i,h,-j}(T_{ij})\}^2$ .




Fig. 1: Illustration of functional regression on a manifold (left panel) and illustration of the asymptotic result of (9) for  $d = 2$  (right panel).

For the bandwidths  $h_{pca}$  in (6) and  $h_{reg}$  in (7), we choose the pair  $(h_{pca}, h_{reg})$  from a pool  $\mathbb{H}$  of candidate pairs to minimize the following leave-one-out cross-validation error  $CV(h_{pca}, h_{reg}) = \sum_{i=1}^n \{Y_i - \hat{g}_{h_{pca}, h_{reg}, -i}(\hat{X}_i)\}^2$ , where  $\hat{g}_{h_{pca}, h_{reg}, -i}$  denotes the leave-one-out estimate of  $g$  with parameters  $h_{pca}, h_{reg}$  without using the pair  $(\hat{X}_i, Y_i)$ . The pool  $\mathbb{H}$  shall be constructed in the way that every  $\hat{\mathcal{N}}_{\mathcal{L}^2}(h_{pca}, \hat{X}_i)$  contains at least  $\hat{d} + 1$  samples for every pair  $(h_{pca}, h_{reg})$  in  $\mathbb{H}$  to ensure sufficient data for local estimation.

### 3. THEORETICAL PROPERTIES

We focus on the scenario that  $\inf_i m_i$  increases with the sample size  $n$ , while leave the one that  $\sup_i m_i \leq m_0 < \infty$  for future research due to elevated challenges. Without loss of generality, assume  $m_i \asymp m$  where  $a_n \asymp b_n$  denotes  $0 < \liminf a_n/b_n < \limsup a_n/b_n < \infty$ . We further assume that  $\zeta_{ij}$ , and similarly,  $T_{ij}$  and  $X_i$ , are independently and identically distributed, while emphasize that the development below can be modified to accommodate fixed designs, weak dependence and/or heterogeneous distributions. This generality will require considerably heavier technicalities without adding further insight, and is not pursued here.

The discrepancy between  $\hat{X}_i$  and  $X_i$ , quantified by  $\|\hat{X}_i - X_i\|_{\mathcal{L}^2}$ , is termed the contamination of  $X_i$ . The decay of this contamination is intimately linked to the consistency of our estimates of the intrinsic dimension, the tangent space, and eventually the regression functional  $g(x)$ . Moreover, the convergence rate of  $\hat{g}(x)$  is found to exhibit a phase transition phenomenon depending on the interplay between the intrinsic dimension and the decay of contamination. To set the stage, we start with a property of contamina-



tion in recovery of functional data by the individual smoothing approach in Section 2.1. Specifically, we study the  $p$ th moment of contamination when  $\hat{X}_i$  is the ridged local linear estimate in (3). Our result below for an arbitrary  $p$ th moment is not present in the literature (e.g., Fan, 1993, for  $p = 2$  only).

Let  $\Sigma(\nu, L)$  denote the Hölder class with an exponent  $\nu$  and an Hölder constant  $L$ , which represents the set of  $\ell = \lfloor \nu \rfloor$  times differentiable functions  $F$  whose derivative  $F^{(\ell)}$  satisfies  $|F^{(\ell)}(t) - F^{(\ell)}(s)| \leq L|t - s|^{\nu - \ell}$  for  $s, t \in D$ , where  $\lfloor \nu \rfloor$  denotes the largest integer strictly smaller than  $\nu$ . We require the following mild assumptions, and assume  $h_i \asymp h_0$  without loss of generality.

- (A1)  $K$  is a differentiable kernel with a bounded derivative,  $\int_{-1}^1 K(u)du = 1$ ,  $\int_{-1}^1 uK(u)du = 0$ , and  $\int_{-1}^1 |u|^p K(u)du < \infty$  for all  $p > 0$ .
- (A2) The sampling density  $f_T$  is bounded away from zero and infinity, i.e., for some constants  $C_{T,1}, C_{T,2} \in (0, \infty)$ ,  $C_{T,1} = \inf_{t \in D} f_T(t) \leq \sup_{t \in D} f_T(t) = C_{T,2}$ .
- (A3)  $X \in \Sigma(\nu, L_X)$ , where  $L_X > 0$  is a random quantity and the constant  $\nu \in (0, 2]$  quantifies the smoothness of the process.
- (A4) For all  $r \geq 1$ ,  $E \sup_t |X(t)|^r < \infty$ ,  $E(L_X)^r < \infty$  and  $E|\zeta|^r < \infty$ .

The condition  $E \sup_t |X(t)|^r < \infty$  holds rather generally (Li & Hsing, 2010; Zhang & Wang, 2016), compared to a stronger assumption on  $X$  given in (A.1) of Hall et al. (2006). The following proposition is an immediate consequence of Lemma S.1 in the Supplementary Material, and hence its proof is omitted.

**PROPOSITION 1.** *For any  $p \geq 1$ , assume  $E|\zeta|^p < \infty$ . Under the assumptions (A1)–(A3), for the estimate  $\hat{X}$  in (3) with  $h_0 \asymp m^{-\frac{1}{2\nu+1}}$  and  $\delta = m^{-2}$ , we have*

$$\{E(\|\hat{X} - X\|_{\mathcal{L}^2}^p \mid X)\}^{1/p} = O(m^{-\frac{\nu}{2\nu+1}}) \left\{ \sup_t |X(t)| + L_X \right\}. \quad (8)$$

Furthermore, if the assumption (A4) also holds, then  $(E\|\hat{X} - X\|_{\mathcal{L}^2}^p)^{1/p} = O(m^{-\frac{\nu}{2\nu+1}})$ .

When  $X$  is deterministic as in nonparametric regression, the rate in (8) for  $p = 2$  coincides with that in Tsybakov (2008). In addition, the  $p$ th order of the contamination  $\|\hat{X}_i - X_i\|_{\mathcal{L}^2}$  decays at a polynomial rate that depends on  $\nu$ , but not the order  $p$ .

To analyze the asymptotic property of  $\hat{g}(x)$ , we make the following assumptions.

- (B1) The probability density  $f$  of  $X$  on  $\mathcal{M}$  satisfies  $C_{f,1} = \inf_{x \in \mathcal{M}} f(x) \leq \sup_{x \in \mathcal{M}} f(x) = C_{f,2}$  for some constants  $0 < C_{f,1} \leq C_{f,2} < \infty$ .
- (B2) The regression functional  $g$  has a bounded second derivative.

For (B1), since the functional predictor resides on a low-dimensional manifold, the existence of a density can be safely assumed. We also make the following assumption on the imputed trajectories in Section 2.1.

- (B3)  $\hat{X}_1, \dots, \hat{X}_n$  are independently and identically distributed. For some  $\beta \in (0, \infty)$  and all  $p \geq 1$ ,  $\{E(\|\hat{X} - X\|_{\mathcal{L}^2}^p | X)\}^{1/p} \leq C_p m^{-\beta} \eta(X)$  for some constant  $C_p$  depending only on  $p$  and some nonnegative function  $\eta(X)$  depending only on  $X$  such that  $E\{\eta(X)\}^p < \infty$ .

Under the assumptions (A1)–(A4), by Proposition 1, the imputed functions  $\hat{X}_1, \dots, \hat{X}_n$  by individual smoothing via local linear estimation (3) satisfies (B3) with  $\beta = \nu/(2\nu + 1)$ . Therefore, (B3) could be replaced with the more concrete assumptions (A1)–(A4). It can be relaxed to accommodate heterogeneous data distributions and weakly dependent functional data by modifying our proofs. Also, it is possible to accommodate imputed functions that are attained by borrowing information across individuals (e.g., Yao et al., 2005a), which is beyond our scope here and can be a topic of future research.

The contamination of the predictor  $X$  renders the true neighborhood  $\mathcal{N}_{\mathcal{L}^2}(h_{pca}, x) = \{X_i : \|X_i - x\|_{\mathcal{L}^2} < h_{pca}\}$  inaccessible. However, we can show that the contaminated one  $\hat{\mathcal{N}}_{\mathcal{L}^2}(h_{pca}, x)$  is a good estimate; see Section S.2 and Lemma 8 in Supplementary Material for details. Consequently, the local manifold structure can be consistently estimated in the sense of the following theorem.

**THEOREM 1.** *Suppose that the assumptions (B1) and (B3) hold.*

- (a)  $\hat{d}$  is a consistent estimator of  $d$  when  $\min\{k_1, k_2\} \rightarrow \infty$  and  $\max\{k_1, k_2\}/m \rightarrow 0$ .
- (b) If  $h_{pca} \rightarrow 0$  and  $h_{pca} \gtrsim \max\{m^{-\beta+\epsilon}, n^{-1/(d+2)}\}$  for an arbitrarily small but fixed constant  $\epsilon > 0$ , then the eigenbasis  $\{\hat{\varphi}_k\}_{k=1}^d$  derived from  $\hat{\mathcal{C}}_x$  in (6) is close to an orthonormal basis  $\{\phi_k\}_{k=1}^d$  of  $T_x\mathcal{M}$ , in the sense that, for each  $x \in \mathcal{M}$ ,

$$\hat{\varphi}_k = \phi_k + O_P(h_{pca}^{3/2})u_k + O_P(h_{pca})u_k^\perp \quad (k = 1, \dots, d), \quad (9)$$

where  $u_k \in T_x\mathcal{M}$ ,  $u_k^\perp \perp T_x\mathcal{M}$ , and  $\|u_k\|_{\mathcal{L}^2} = \|u_k^\perp\|_{\mathcal{L}^2} = 1$ .

In light of Theorem 1(a), we shall from now on present the subsequent results by conditioning on the event  $\hat{d} = d$ . For part (b) that is illustrated in the right panel of Figure 1, the condition  $h_{pca} \gtrsim m^{-\beta+\epsilon}$  suggests that  $h_{pca}$  shall be larger than the contamination by an arbitrarily small polynomial order of  $m$ . This is required to ensure that the discrepancy between the estimated local neighborhood  $\hat{N}_{\mathcal{L}^2}(h_{pca}, x)$  and the uncontaminated neighborhood  $\mathcal{N}_{\mathcal{L}^2}(h_{pca}, x) = \{X_i : \|x - X_i\|_{\mathcal{L}^2} < h_{pca}, i = 1, \dots, n\}$  is asymptotically negligible, suggested by Lemma 8 in Supplementary Material. The curvature at  $x$  is a constant that is absorbed into the  $O_P$  terms, and thus does not influence the asymptotic rate. However, practically it is often more difficult to estimate the tangent structure at a point with larger curvature.

We are ready to state the results on the estimated regression functional. Recall that  $\hat{g}(x)$  in (7) is obtained by applying the local linear smoother to the coordinates of contaminated predictors within the estimated tangent space at  $x$ . It is well known that the local linear estimator does not suffer from boundary effects, i.e., the first order behavior of the estimator on the boundary is the same as in the interior (Fan, 1992). However, the contamination of the predictor has different impact, and we shall address the interior and boundary cases separately. Denote  $\mathcal{X} = \{(X_1, \hat{X}_1), \dots, (X_n, \hat{X}_n)\}$  and  $\mathcal{M}_h = \{x \in \mathcal{M} : \inf_{y \in \partial\mathcal{M}} \mathfrak{d}(x, y) \leq h\}$ , where  $\partial\mathcal{M}$  denotes the boundary of  $\mathcal{M}$  and  $\mathfrak{d}(\cdot, \cdot)$  denotes the distance function on  $\mathcal{M}$ . For points sufficiently far away from the boundary of  $\mathcal{M}$ , we have the following result about the convergence rate of the estimator  $\hat{g}(x)$ .

**THEOREM 2.** *Assume that (A1) and (B1)–(B3) hold. Let  $x \in \mathcal{M} \setminus \mathcal{M}_{h_{reg}}$  and  $h_{pca}$  satisfies the conditions of Theorem 1(b). For an arbitrarily small but fixed constant  $\epsilon > 0$ , suppose that  $h_{reg} \rightarrow 0$ ,  $h_{reg} > h_{pca}$ , and  $\min\{nh_{reg}, m^\beta h_{reg}^{5/3+\epsilon}\} \rightarrow \infty$ . Then*

$$E \left[ \{\hat{g}(x) - g(x)\}^2 \mid \mathcal{X} \right] = O_P \left( h^4 + \frac{1}{m^{2\beta} h_{reg}^{2+2\epsilon}} + \frac{1}{nh^d} \right). \quad (10)$$

*In addition, if  $h_{pca} \asymp \max\{m^{-\beta}, n^{-1/(d+2)}\}$ , and  $h_{reg} \asymp n^{-1/(d+4)}$  when  $m \gtrsim n^{(3+\epsilon)/\{\beta(d+4)\}}$  and  $h_{reg} \asymp m^{-\beta/(3+\epsilon)}$  otherwise, then*

$$E \left[ \{\hat{g}(x) - g(x)\}^2 \mid \mathcal{X} \right] = O_P \left( n^{-\frac{4}{d+4}} + m^{-\frac{4\beta}{3+\epsilon}} \right). \quad (11)$$

We emphasize the following observations from this theorem. First, according to our analysis in Supplementary Material, the first two terms on the right hand side of (10)

correspond to the bias while the last term stems from the variability of the estimator. This suggests that, under the conditions of the theorem, the contamination has impact on the asymptotic bias but not the variance. Second, the convergence rate of  $\hat{g}(x)$  is a polynomial of the sample size  $n$  and the sampling rate  $m$ . This is in contrast with traditional functional nonparametric regression methods that do not exploit the intrinsic structure and thus cannot reach a polynomial rate of convergence.

Third, the rate in (11) consists of two terms, one related to the intrinsic dimension  $d$  and the sample size  $n$ , and the other related to  $m$  and  $\beta$  that together characterize the contamination of the predictor. As  $\epsilon > 0$  is arbitrary, the transition of these two terms occurs at the rate  $m_0 \asymp n^{3/\{\beta(d+4)\}}$ . When the sampling rate falls below  $m_0$ , the contamination term dominates the convergence rate in (11). Otherwise, the intrinsic dimension and sample size determine the rate. This phase transition, although sharing the similar spirit of Cai & Yuan (2011) and Zhang & Wang (2016), has a different interpretation, as follows. When the contamination level is low, the manifold structure can be estimated reliably and utilized for regression. In contrast, when the contamination is in a high level, for example, where  $m$  or  $\beta$  is small, the manifold structure is buried by noise and cannot be well exploited. Finally, it is observed that the phase transition threshold  $m_0$  increases with the intrinsic dimension  $d$  that indicates the complexity of a manifold. This interesting finding suggests that, although a complex manifold makes the estimation more challenging, for example, leading to a slower rate, such manifold is more resistant to contamination.

In our setup, the actual observed predictor is  $\mathbb{X}_i = \{(T_{i1}, X_{i1}^*), \dots, (T_{im}, X_{im}^*)\}$  and is an  $m_i$ -dimensional random vector. Moreover, the distribution of this random vector is fully supported on  $\mathbb{R}^{m_i}$  due to the presence of the noise  $\zeta_{ij}$ , and thus the support of the distribution of the recovered trajectory  $\hat{X}_i$  is also  $m_i$ -dimensional. Smoothness of functional data could help tighten the distribution of  $\hat{X}_i$ , but does not reduce its dimension. As  $m_i$  goes to infinity, it might then raise a serious concern of the curse of dimensionality. In this sense, the polynomial rate and phase transition phenomenon in Theorem 2 are remarkable: when  $\inf_i m_i$  surpasses certain threshold, by exploiting the low-dimensional manifold structure, the growing dimension of the contamination can be defeated with the aid of smoothness.

The following theorem characterizes the behavior of  $\hat{g}$  on the boundary of  $\mathcal{M}$ .

THEOREM 3. Assume that (A1) and (B1)–(B3) hold. Let  $x \in \mathcal{M}_{h_{reg}}$  and  $h_{pca}$  satisfies the conditions of Theorem 1(b). For an arbitrarily small but fixed constant  $\epsilon > 0$ , suppose that  $h_{reg} \rightarrow 0$ ,  $h_{reg} > h_{pca}$ , and  $\min\{nh_{reg}, m^\beta h_{reg}^{3+\epsilon}\} \rightarrow \infty$ . Then

$$E \left[ \{\hat{g}(x) - g(x)\}^2 \mid \mathcal{X} \right] = O_P \left( h^4 + \frac{1}{m^{2\beta} h_{reg}^{4+2\epsilon}} + \frac{1}{nh^d} \right).$$

In addition, if  $h_{pca} \asymp \max\{m^{-\beta}, n^{-1/(d+2)}\}$ , and  $h_{reg} \asymp n^{-1/(d+4)}$  when  $m \gtrsim n^{(4+\epsilon)/\{\beta(d+4)\}}$  and  $h_{reg} \asymp m^{-\beta/(4+\epsilon)}$  otherwise, then

$$E \left[ \{\hat{g}(x) - g(x)\}^2 \mid \mathcal{X} \right] = O_P \left( n^{-\frac{4}{d+4}} + m^{-\frac{4\beta}{4+\epsilon}} \right). \quad (12)$$

By comparing the above with Theorem 2, we see that the effect of the intrinsic dimension on convergence is the same, regardless where  $\hat{g}$  is evaluated on the manifold. However, the effect of contamination behaves differently, due to the fact that the second order behavior of the local linear estimator depends on the location and needs to be considered when there is contamination of  $X$ . Moreover, we see that the phase transition occurs at  $\mathbf{m}_1 = n^{4/\{\beta(d+4)\}} \gg \mathbf{m}_0$ , and when the contamination dominates, the convergence is slightly slower for boundary points than for interior points. This is the price we pay for the boundary effect when the predictor is contaminated, which is in contrast with the classical result on the local linear estimator (Fan, 1993).

#### 4. SIMULATION STUDY

To demonstrate the performance of our framework, we conduct simulation studies for three different manifolds, namely, the three-dimensional rotation group  $SO(3)$ , the Klein bottle and the mixture of two Gaussian densities.

- $SO(3)$  manifold: we set  $X_i(t) = \sum_{k=1}^9 z_{ik} b_k(t)$ , where  $b_{2\ell-1}(t) = \cos\{(2\ell-1)\pi t/10\}/5^{1/2}$  and  $b_{2\ell}(t) = \sin\{(2\ell-1)\pi t/10\}/5^{1/2}$ . To generate the random variables  $z_{ik}$ , for a vector  $r = (r_1, r_2, r_3)$  and a variable  $\theta \in \mathbb{R}$ , we define

$$R(r, \theta) = (1 - \cos \theta) r r^T + \begin{pmatrix} \cos \theta & -r_3 \sin \theta & r_2 \sin \theta \\ r_3 \sin \theta & \cos \theta & -r_1 \sin \theta \\ -r_2 \sin \theta & r_1 \sin \theta & \cos \theta \end{pmatrix}.$$

Denoting  $e_2 = (0, 1, 0)^T$  and  $e_3 = (0, 0, 1)^T$ , we set  $(z_{i1}, \dots, z_{i9})^T = \text{vec}(Z_i)$  with Euler angle parameterization  $Z_i = R(e_3, u_i)R(e_2, v_i)R(e_3, w_i)$ , where  $(u_i, v_i)$  are uniformly sampled from the two-dimensional sphere  $S^2 = [0, 2\pi) \times [0, \pi]$ , and  $w_i$  are uniformly sampled from the unit circle  $S^1 = [0, 2\pi)$ .

- Klein bottle: we set  $X_i(t) = \sum_{k=1}^4 z_{ik} b_k(t)$  with  $b_k(t)$  as in the  $SO(3)$  setting. We set  $z_{i1} = (2 \cos v_i + 1) \cos u_i$ ,  $z_{i2} = (2 \cos v_i + 1) \sin u_i$ ,  $z_{i3} = 2 \sin v_i \cos(u_i/2)$  and  $z_{i4} = 2 \sin v_i \sin(u_i/2)$ , where  $u_i$  and  $v_i$  independently sampled from the uniform distribution on  $(0, 2\pi)$ . Here  $(u, v) \mapsto (z_1, z_2, z_3, z_4)$  is a parameterization of the Klein bottle with an intrinsic dimension  $d = 2$ .
- Gaussian mixture: we set  $X_i$  to  $X_i(t) = \exp\{-(t - u_i)^2/2\}/(2\pi)^{1/2} + \exp\{-(t - v_i)^2/2\}/(2\pi)^{1/2}$  with  $(v_1, v_2)^T$  uniformly sampled from a circle with diameter 0.5, similar to that used in Chen & Müller (2012).

The functional predictor  $X_i$  is observed at  $m_i$  points  $T_{i1}, \dots, T_{im_i}$  in the interval  $[0, 1]$  with heteroscedastic measurement errors  $\zeta_{ij} \sim N(0, \sigma_{ij}^2)$ , where  $\sigma_{ij}$  is determined by the signal-to-noise ratio  $\text{snr}_X = \text{Var}\{X(T_{ij})|T_{ij}\}/\sigma_{ij}^2 = 4$ . The response is generated by  $Y_i = 4 \sin(4Z_i) \cos(Z_i^2) + 2\Gamma(1 + Z_i/2) + \varepsilon_i$  with  $Z_i = \int_0^1 X_i^2(t)tdt$  and  $\Gamma(\alpha) = \int_0^\infty s^{\alpha-1}e^{-s}ds$ . The noise  $\varepsilon_i$  added to the response  $Y$  is a centered Gaussian variable with variance  $\sigma_\varepsilon^2$  that is determined by the signal-to-noise ratio  $\text{snr}_Y = \text{var}(Y)/\sigma_\varepsilon^2 = 2$ . To see the impact of the manifold structure on regression, we normalize the functional predictor in all settings to the unit scale, i.e., multiplying  $X$  by the constant  $c = 1/(E\|X\|^2)^{1/2}$  so that the resultant  $X$  satisfies  $E\|X\|^2 = 1$ . Such scaling does not change the geometric structure of manifolds but the size. In order to account for at least 95% of variance of data, we find empirically that more than 10 principal components are needed in all settings, i.e., the dimensions of the contaminated data are considerably larger than their intrinsic dimensions.

For evaluation, we generate independent test data of size 5000, and compute the root mean square error using the test data. In the test data, each predictor is also discretely measured and contaminated by noise in the same way of the training sample. We compare our method with nonparametric estimators based on functional Nadaraya–Watson smoothing, functional conditional expectation, functional mode, functional conditional median and multi-method that averages estimates from the methods of functional conditional expectation, functional mode and functional conditional median (Ferraty & Vieu,

2006). Functional linear regression is also included to illustrate the impact of nonlinear relationship. The tuning parameters in these methods, such as the number of principal components for functional linear regression and the bandwidth for the nonparametric methods, are selected by 10-fold cross-validation.

We consider the scenario of dense functional data here, while refer readers to Supplementary Material for simulation studies for sparsely observed data. Specifically, we set  $m_i = m = 100$  and  $T_{ij} = t_j$ , where  $t_1, \dots, t_m$  are equally spaced over  $[0, 1]$ . Three sample sizes are considered, namely,  $n = 250, 500, 1000$ . We repeat each study 100 times independently, and the results are presented in Table 1. First, we observe that the proposed method enjoys favorable numerical performance in all simulation settings. Second, as the sample size grows, the reduction in root mean square error is more prominent for the proposed method than for the others. For example, the relative reduction from  $n = 250$  ( $n = 500$ , respectively) to  $n = 500$  ( $n = 1000$ , respectively) is 25.5% (22.7%, respectively) for our method, but 8.49% (2.75%, respectively) for the functional Nadaraya–Watson estimator. This may provide some numerical evidence that the proposed estimator has a faster convergence rate. Furthermore, it also provides evidence for the polynomial rate stated in Theorem 2 and 3. Based on these theorems the relative reduction is expected to be  $1 - (n_1/n_2)^{2/(d+4)}$  when the sample size increases from  $n_1$  to  $n_2$ , as the data is sufficiently dense and thus the convergence rate is dominated by the intrinsic dimension. For the setting of Klein bottle, it is about 20.6%, and the empirical relative reduction is 22.7% from  $n_1 = 500$  to  $n_2 = 1000$ . Similar observations can be made for other settings. In contrast, the existing kernel methods perform no better than a logarithmic rate, providing numerical evidence for the theory of Mas (2012). Third, as the intrinsic dimension goes up, the relative reduction in root mean square error for our estimator decreases, suggesting that the intrinsic dimension plays an important role in the convergence rate. Finally, different manifolds result in different constants hidden in the  $O_P$  terms in Theorem 2 and 3. For example, those in the  $SO(3)$  setting seem relatively smaller than their counterparts in the setting of Klein bottle according to Table 1.

## 5. REAL DATA EXAMPLES

We apply our method to analyze three real datasets. For the purpose of evaluation, we train our method on 75% of each dataset and reserve the other 25% as test data. The

Table 1: Results of simulation studies for densely observed data

	$SO(3)$ Manifold			Klein Bottle			Gaussian Mixture		
	$n = 250$	$n = 500$	$n = 1000$	$n = 250$	$n = 500$	$n = 1000$	$n = 250$	$n = 500$	$n = 1000$
FLR	22.1 (0.34)	21.8 (0.23)	21.6 (0.20)	61.3 (0.62)	61.2 (0.39)	6.09 (0.35)	29.6 (1.43)	29.0 (1.26)	28.8 (0.99)
FNW	16.2 (0.58)	15.7 (0.43)	15.5 (0.32)	31.8 (4.05)	29.1 (1.79)	28.3 (0.68)	18.7 (1.46)	17.5 (0.83)	17.0 (0.65)
FCE	15.3 (0.66)	14.1 (0.52)	13.2 (0.30)	29.7 (1.46)	27.1 (1.04)	26.1 (0.81)	21.1 (1.32)	20.4 (0.93)	19.8 (0.64)
FMO	25.4 (1.16)	23.0 (0.94)	22.0 (0.85)	46.2 (3.07)	41.2 (2.20)	38.3 (1.87)	35.9 (2.80)	33.6 (2.05)	32.2 (1.61)
FCM	20.2 (0.60)	18.6 (0.52)	17.2 (0.35)	39.1 (2.67)	33.9 (1.61)	30.9 (1.02)	27.3 (1.71)	25.1 (1.05)	23.2 (0.83)
MUL	18.2 (0.59)	16.6 (0.48)	15.4 (0.31)	34.0 (2.13)	30.0 (1.24)	27.7 (0.92)	24.6 (1.49)	23.1 (1.07)	21.8 (0.81)
FREM	10.1 (0.72)	8.16 (0.56)	6.38 (0.25)	16.5 (1.39)	12.3 (1.11)	9.51 (0.74)	10.5 (1.32)	8.08 (0.86)	6.12 (0.75)

FLR, functional linear regression; FNW, functional Nadaraya–Watson smoothing; FCE, functional conditional expectation; FMO, functional mode, FCM, functional conditional median; MUL, multi-method; FREM, the proposed functional regression on manifold; MSP, meat spectrometric data; DTI, diffusion tensor imaging data; SBP, systolic blood pressure data. The numbers outside of parentheses are the Monte Carlo average of root mean square error based on 100 independent simulation replicates, and the numbers in parentheses are the corresponding standard error.

root mean square error is computed on the held-out test data. We repeat this 100 times based on random partitions of the datasets, and summarize the results in Table 2.

The first application is to predict the fat content of a piece of meat based on a spectrometric curve for the meat using the Tecator dataset with 215 meat samples (Ferraty & Vieu, 2006). For each sample, the spectrometric curve for a piece of finely chopped pure meat was measured at 100 different wavelengths from 850 to 1050nm. Along with the spectrometric curves, the fat content for each piece of meat was recorded. Comparing to the analytic chemistry required for measuring the fat content, obtaining a spectrometric curve is less time and cost consuming. As in Ferraty & Vieu (2006), we predict the fat content based on the first derivative curves approximated by the difference quotient between measurements at adjacent wavelengths, shown in the left panel of Figure 2. It is seen that there are some striking patterns around the middle wavelengths. The proposed method is able to capture these patterns by a low-dimensional manifold structure. For example, functional linear regression uses 15.7 principal components on average with a standard error 1.07, while the intrinsic dimension estimated by our method is 5.05 with a standard error 0.62. Thus, our method predicts the fat content more accurately than the others by a significant margin according to Table 2.

The second example studies the relationship between cognitive function and brain microstructure in the corpus callosum of patients with multiple sclerosis, a common demyelinating disease caused by inflammation in the brain. Demyelination refers to the damage to myelin that protects axons and helps nerve signal to travel faster. It occurs in the white matter of the brain and can potentially lead to loss of mobility or even






Fig. 2: first derivatives of meat spectrometric curves (left panel), fractional anisotropy profiles (middle panel), and systolic blood pressure over time (right panel).

cognitive impairment (Jongen et al., 2012). Diffusion tensor imaging, a technique that can produce high-resolution images of white matter tissues by tracing water diffusion within the tissues, is an important method to examine potential myelin damage in the brain. For example, from such images, some properties of white matter, such as fractional anisotropy of water diffusion, can be derived. It has been shown that fractional anisotropy is related to multiple sclerosis (Ibrahim et al., 2011).

To predict cognitive performance based on fractional anisotropy profiles, we utilize the data collected at Johns Hopkins University and the Kennedy-Krieger Institute. The data contains  $n = 340$  profiles from multiple sclerosis patients and paced auditory serial addition test scores that quantify cognitive function (Gronwall, 1977), where each profile was recorded at a grid of 93 points. In the middle panel of Figure 2, we show all fractional anisotropy profiles, and observe that the data is considerably more complex than the spectrometric data. The average of estimated intrinsic dimensions is 5.82 with a standard error 0.098. By contrast, the average number of principal components for functional linear regression is 11.98 with a standard error 5.22. According to Table 2, our method enjoys the most accurate prediction, while all other functional nonparametric methods deteriorate substantially.

Our third example concerns systolic blood pressure of healthy men using an anonymous data from the Baltimore longitudinal study of aging. In the study, 1590 healthy male volunteers were scheduled to visit the Gerontology Research Center bi-annually. Systolic blood pressure and current age were recorded during each visit. The design of the data is sparse and irregular, as many visits were missed by participants or not on the schedule; see Pearson et al. (1997) for more details. Our study aims to predict the average systolic blood pressure in late middle age, between 55 and 60, based on the blood pressure

Table 2: Results for real data analysis

	FLR	FNW	FCE	FMO	FCM	MUL	FREM
MSP	2.56 (0.43)	2.42 (0.33)	1.97 (0.35)	2.66 (0.46)	2.82 (0.45)	2.31 (0.35)	1.06 (0.34)
DTI	1.14 (0.09)	1.28 (0.12)	1.36 (0.13)	1.78 (0.16)	1.25 (0.14)	1.33 (0.13)	0.96 (0.09)
SBP	1.34 (0.18)	1.57 (0.17)	1.64 (0.16)	2.33 (0.26)	1.68 (0.19)	1.76 (0.17)	1.15 (0.11)

FLR, functional linear regression; FNW, functional Nadaraya–Watson smoothing; FCE, functional conditional expectation; FMO, functional mode, FCM, functional conditional median; MUL, multi-method; MSP, meat spectrometric data; FREM, the proposed functional regression on manifold; DTI, diffusion tensor imaging data; SBP, systolic blood pressure data. The numbers outside of parentheses are the Monte Carlo average of root mean square error based on 100 independent simulation replicates, and the numbers in parentheses are the corresponding standard error. The results for the diffusion tensor imaging data and systolic blood pressure data are scaled by 0.1 for visualization.

trajectory between age 45 and 55. By excluding subjects with at most one visit between age 45 and 55 and no visit between 55 and 60, we obtain a subset of the data with  $n = 323$  subjects and on average 4.2 visits per subject, shown in the right panel of Figure 2. The average of estimated intrinsic dimensions is 2.4 with a standard error 0.069, while the average number of principal components for functional linear regression is 4 with a standard error 2.01. Based on Table 2, our method outperforms others significantly.

#### ACKNOWLEDGEMENT

Fang Yao’s research is partially supported by National Natural Science Foundation of China (Key Grant 11931001 and General Grant 11871080), and the Key Laboratory of Mathematical Economics and Quantitative Finance (Peking University), Ministry of Education.

#### SUPPLEMENTARY MATERIAL

Additional details and simulation studies for sparse functional data, the proofs of main theorems, auxiliary results, and technical lemmas with proofs are collected in an online Supplementary Material for space economy.

#### REFERENCES

- ASWANI, A., BICKEL, P. & TOMLIN, C. (2011). Regression on manifolds: Estimation of the exterior derivative. *The Annals of Statistics* **39**, 48–81.
- BHATTACHARYA, R. & LIN, L. (2017). Omnibus CLTs for Fréchet means and nonparametric inference on non-Euclidean spaces. *Proceedings of the American Mathematical Society* **145**, 413–428.

- BHATTACHARYA, R. & PATRANGENARU, V. (2003). Large sample theory of intrinsic and extrinsic sample means on manifolds. I. *The Annals of Statistics* **31**, 1–29.
- BHATTACHARYA, R. & PATRANGENARU, V. (2005). Large sample theory of intrinsic and extrinsic sample means on manifolds. II. *The Annals of Statistics* **33**, 1225–1259.
- CAI, T. & YUAN, M. (2011). Optimal estimation of the mean function based on discretely sampled functional data: Phase transition. *The Annals of Statistics* **39**, 2330–2355.
- CARDOT, H., FERRATY, F. & SARDA, P. (1999). Functional linear model. *Statistics & Probability Letters* **45**, 11–22.
- CARDOT, H. & SARDA, P. (2005). Estimation in generalized linear models for functional data via penalized likelihood. *Journal of Multivariate Analysis* **92**, 24–41.
- CHEN, D. & MÜLLER, H. (2012). Nonlinear manifold representations for functional data. *The Annals of Statistics* **40**, 1–29.
- CHENG, M. & WU, H. (2013). Local linear regression on manifolds and its geometric interpretation. *Journal of the American Statistical Association* **108**, 1421–1434.
- COIFMAN, R., LAFON, S., LEE, A. B., MAGGIONI, M., NADLER, B., WARNER, F. & ZUCKER, S. W. (2005). Geometric diffusions as a tool for harmonic analysis and structure definition of data: Diffusion maps. *PNAS* **102**, 7426–7431.
- CORNEA, E., ZHU, H., KIM, P. & IBRAHIM, J. G. (2017). Regression models on Riemannian symmetric spaces. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **79**, 463–482.
- DAI, X. & MÜLLER, H.-G. (2018). Principal component analysis for functional data on Riemannian manifolds and spheres. *Annals of Statistics* **46**, 3334–3361.
- DELAIGLE, A. & HALL, P. (2010). Defining probability density for a distribution of random functions. *The Annals of Statistics* **38**, 1171–1193.
- FAN, J. (1992). Design-adaptive nonparametric regression. *Journal of the American Statistical Association* **87**, 998–1004.
- FAN, J. (1993). Local linear regression smoothers and their minimax efficiencies. *The Annals of Statistics* **21**, 196–216.
- FAN, J. & GIJBELS, I. (1996). *Local Polynomial Model ling and Its Applications*. London: Chapman and Hall.
- FERRATY, F., KEILEGOM, I. V. & VIEU, P. (2012). Regression when both response and predictor are functions. *Journal of Multivariate Analysis* **109**, 10–28.
- FERRATY, F. & VIEU, P. (2006). *Nonparametric Functional Data Analysis: Theory and Practice*. New York: Springer-Verlag.
- GRONWALL, D. M. A. (1977). Paced auditory serial-addition task: A measure of recovery from concussion. *Perceptual and Motor Skills* **44**, 367–373.
- HALL, P. & KEILEGOM, I. V. (2007). Two sample tests in functional data analysis starting from discrete data. *Statistica Sinica* **17**, 1511–1531.
- HALL, P. & MARRON, J. S. (1997). On the shrinkage of local linear curve estimators. *Statistics and Computing* **516**, 11–17.
- HALL, P., MÜLLER, H.-G. & WANG, J.-L. (2006). Properties of principal component methods for functional and longitudinal data analysis. *The Annals of Statistics* **34**, 1493–1517.

- HUCKEMANN, S., HOTZ, T. & MUNK, A. (2010). Intrinsic shape analysis: Geodesic PCA for Riemannian manifolds modulo isometric Lie group actions. *Statistica Sinica* **20**, 1–58.
- IBRAHIM, I., TINTERA, J., SKOCH, A., F., J., P., H., MARTINKOVA, P., ZVARA, K. & RASOVA, K. (2011). Fractional anisotropy and mean diffusivity in the corpus callosum of patients with multiple sclerosis: the effect of physiotherapy. *Neuroradiology* **53**, 917–926.
- JONGEN, P., TER HORST, A. & BRANDS, A. (2012). Cognitive impairment in multiple sclerosis. *Minerva Medica* **103**, 73–96.
- KUDRASZOW, N. L. & VIEU, P. (2013). Uniform consistency of kNN regressors for functional variables. *Statistics & Probability Letters* **83**, 1863–1870.
- LANG, S. (1995). *Differential and Riemannian Manifolds*. New York: Springer.
- LANG, S. (1999). *Fundamentals of Differential Geometry*. New York: Springer.
- LEE, T. C. & SOLO, V. (1999). Bandwidth selection for local linear regression: A simulation study. *Computational Statistics* **14**, 515–532.
- LEVINA, E. & BICKEL, P. (2004). Maximum likelihood estimation of intrinsic dimension. *Advances in Neural Information* **17**, 777–784.
- LI, Y. & HSING, T. (2010). Uniform convergence rates for nonparametric regression and principal component analysis in functional/longitudinal data. *The Annals of Statistics* **38**, 3321–3351.
- LILA, E. & ASTON, J. (2016). Smooth principal component analysis over two-dimensional manifolds with an application to neuroimaging. *The Annals of Applied Statistics* **10**, 1854–1879.
- LIN, L., MU, N., CHEUNG, P. & DUNSON, D. (2019). Extrinsic Gaussian processes for regression and classification on manifolds. *Bayesian Analysis* **14**, 887–906.
- LIN, L., THOMAS, B. S., ZHU, H. & DUNSON, D. B. (2016). Extrinsic local regression on manifold-valued data. *Journal of the American Statistical Association* **112**, 1261–1273.
- LIN, Z. & YAO, F. (2019). Intrinsic Riemannian functional data analysis. *The Annals of Statistics* **47**, 3533–3577.
- LOH, P.-L. & WAINWRIGHT, M. J. (2012). High-dimensional regression with noisy and missing data: Provable guarantees with non-convexity. *The Annals of Statistics* **40**, 1637–1664.
- MAS, A. (2012). Lower bound in regression for functional data by representation of small ball probabilities. *Electronic Journal of Statistics* **6**, 1745–1778.
- MUKHERJEE, S., WU, Q. & ZHOU, D.-X. (2010). Learning gradients on manifolds. *Bernoulli* **16**, 181–207.
- MÜLLER, H. G. & STADTMÜLLER, U. (2005). Generalized functional linear models. *The Annals of Statistics* **33**, 774–805.
- MÜLLER, H. G. & YAO, F. (2008). Functional additive models. *Journal of the American Statistical Association* **103**, 1534–1544.
- PANARETOS, V. M., PHAM, T. & YAO, Z. (2014). Principal flows. *Journal of the American Statistical Association* **109**, 424–436.
- PATRANGENARU, V. & ELLINGSON, L. (2015). *Nonparametric Statistics on Manifolds and Their Applications to Object Data Analysis*. CRC Press.
- PEARSON, J., MORRELL, C., BRANT, L., LANDIS, P. & FLEG, J. (1997). Age-associated changes in blood pressure in a longitudinal study of healthy men and women. *Journal of Gerontology: Medical Sciences* **52**, 177–183.

- PENG, J. & MÜLLER, H.-G. (2008). Distance-based clustering of sparsely observed stochastic processes, with applications to online auctions. *The Annals of Applied Statistics* **2**, 1056–1077.
- RAMSAY, J. O. & SILVERMAN, B. (2002). *Applied Functional Data Analysis: Methods and Case Studies*. New York: Springer.
- RAMSAY, J. O. & SILVERMAN, B. W. (1997). *Functional Data Analysis*. New York: Springer-Verlag.
- RAMSAY, J. O. & SILVERMAN, B. W. (2005). *Functional Data Analysis*. Springer Series in Statistics. New York: Springer, 2nd ed.
- ROWEIS, S. T. & SAUL, L. K. (2000). Nonlinear dimensionality reduction by locally linear embedding. *Science* **290**, 2323–2326.
- SEIFERT, B. & GASSER, T. (1996). Finite-sample variance of local polynomials: analysis and solutions. *Journal of the American Statistical Association* **91**, 267–275.
- SOBER, B., AIZENBUD, Y. & LEVIN, D. (2019). Approximation of functions over manifolds: A moving least-squares approach. *arxiv*.
- SU, J., KURTEK, S., KLASSEN, E. & SRIVASTAVA, A. (2014). Statistical analysis of trajectories on Riemannian manifolds: Bird migration, hurricane tracking, and video surveillance. *The Annals of Applied Statistics* **8**, 530–552.
- TENENBAUM, J. B., SILVA, V. D. & LANGFORD, J. C. (2000). A global geometric framework for nonlinear dimensionality reduction. *Science* **290**, 2319–2323.
- TSYBAKOV, A. B. (2008). *Introduction to Nonparametric Estimation*. New York: Springer.
- VAN DER MAATEN, L. & HINTON, G. (2008). Visualizing data using t-SNE. *Journal of Machine Learning Research* **9**, 2579–2605.
- WU, H.-T. & WU, N. (2018). Think globally, fit locally under the manifold setup: Asymptotic analysis of locally linear embedding. *The Annals of Statistics* **46**, 3805–3837.
- YAO, F. & MÜLLER, H. G. (2010). Functional quadratic regression. *Biometrika* **97**, 49–64.
- YAO, F., MÜLLER, H.-G. & WANG, J.-L. (2005a). Functional data analysis for sparse longitudinal data. *Journal of the American Statistical Association* **100**, 577–590.
- YAO, F., MÜLLER, H. G. & WANG, J.-L. (2005b). Functional linear regression analysis for longitudinal data. *The Annals of Statistics* **33**, 2873–2903.
- YAO, Z. & ZHANG, Z. (2019+). Principal boundary on riemannian manifolds. *Journal of the American Statistical Association*, to appear.
- YUAN, M. & CAI, T. T. (2010). A reproducing kernel Hilbert space approach to functional linear regression. *The Annals of Statistics* **38**, 3412–3444.
- YUAN, Y., ZHU, H., LIN, W. & MARRON, J. S. (2012). Local polynomial regression for symmetric positive definite matrices. *Journal of Royal Statistical Society: Series B (Statistical Methodology)* **74**, 697–719.
- ZHANG, X. & WANG, J.-L. (2016). From sparse to dense functional data and beyond. *The Annals of Statistics* **44**, 2281–2321.
- ZHOU, L. & PAN, H. (2014). Principal component analysis of two-dimensional functional data. *Journal of Computational and Graphical Statistics* **23**, 779–801.

# Supplementary Material for “Functional Regression on Manifold with Contamination”

BY ZHENHUA LIN

*Department of Statistics and Applied Probability, National University of Singapore,*

*117546, Singapore*

stalz@nus.edu.sg

AND FANG YAO

*Department of Probability and Statistics, School of Mathematical Sciences, Center for*

*Statistical Science, Peking University, Beijing, 100871, China*

fyao@math.pku.edu.cn

## SUMMARY

Section S.1 contains details and simulation studies for sparsely observed functional data, Section S.2 contains auxiliary examples and results, Section S.3 provides proofs to the main theorems, and Section S.4 contains technical lemmas.

### S.1. DETAILS AND SIMULATION STUDIES FOR SPARSE DESIGN

When the functional data are sparsely observed, we adopt the procedure proposed by Yao et al. (2005) to recover individual functions, as follows. First, the local linear smoother (Fan, 1993) is adopted to produce an estimate  $\hat{\mu}$  of the global mean function and an estimate  $\hat{C}$  of the global covariance function of  $X$  by pooling all observed data; see Zhang & Wang (2016) for more details. Then estimates  $\hat{\psi}_k$  and  $\hat{\lambda}_k$  of the eigenfunctions and eigenvalues, respectively, are obtained by solving the eigen-equations  $\int_D \hat{C}(s, t) \hat{\psi}_k(s) ds = \hat{\lambda}_k \hat{\psi}_k(t)$ . The global principal component scores  $\xi_{ij}$  are estimated by  $\hat{\xi}_{ik} = \hat{\lambda}_k \hat{\phi}_{ik}^T \hat{\Sigma}_i^{-1} (a_i - b_i)$ , where  $\hat{\phi}_{ik}^T = (\hat{\psi}_k(T_{i1}), \dots, \hat{\psi}_k(T_{im_i}))$ ,  $a_i = (X_{i1}^*, \dots, X_{im_i}^*)^T$ ,  $b_i = (\hat{\mu}(T_{i1}), \dots, \hat{\mu}(T_{im_i}))^T$ , and  $\hat{\Sigma}_i$  is an  $m_i \times m_i$  matrix whose element in the  $j$ th row and  $l$ th column is  $\hat{C}(T_{ij}, T_{il}) + \hat{\sigma}_\zeta^2 1_{j=l}$  with  $\hat{\sigma}_\zeta^2$  being the estimate of the variance of the

Table S.1: Results of simulation studies for sparsely observed data

	$SO(3)$ Manifold			Klein Bottle			Gaussian Mixture		
	$n = 250$	$n = 500$	$n = 1000$	$n = 250$	$n = 500$	$n = 1000$	$n = 250$	$n = 500$	$n = 1000$
FLR	24.0 (0.27)	23.8 (0.27)	23.6 (0.25)	61.4 (0.51)	61.3 (0.44)	61.1 (0.35)	43.2 (1.64)	42.0 (1.65)	41.1 (1.63)
FNW	23.9 (0.24)	23.9 (0.26)	23.8 (0.24)	60.9 (0.59)	60.2 (0.62)	59.7 (0.62)	49.7 (3.65)	47.2 (1.05)	46.2 (0.76)
FCE	25.0 (0.42)	24.9 (0.33)	24.8 (0.31)	62.5 (0.89)	62.2 (0.75)	61.9 (0.56)	49.5 (1.12)	49.2 (0.83)	48.8 (0.78)
FMO	33.5 (1.61)	32.8 (1.19)	32.0 (1.45)	83.9 (3.70)	82.7 (3.06)	81.5 (2.64)	63.8 (3.06)	62.4 (2.16)	60.2 (2.30)
FCM	25.5 (0.60)	25.1 (0.42)	24.4 (0.30)	65.3 (1.56)	63.4 (1.15)	61.5 (0.74)	50.5 (1.52)	49.4 (1.05)	47.7 (0.76)
MUL	26.3 (0.59)	26.0 (0.45)	25.5 (0.39)	66.3 (1.32)	65.4 (1.12)	64.2 (0.79)	51.6 (1.35)	50.9 (0.99)	49.5 (0.89)
FREM	24.9 (0.96)	24.4 (0.87)	23.9 (0.68)	56.1 (2.62)	52.0 (1.19)	50.1 (0.64)	37.1 (2.11)	34.7 (1.98)	32.7 (1.46)

FLR, functional linear regression; FNW, functional Nadaraya–Watson smoothing; FCE, functional conditional expectation; FMO, functional mode, FCM, functional conditional median; MUL, multi-method; FREM, the proposed functional regression on manifold; MSP, meat spectrometric data; DTI, diffusion tensor imaging data; SBP, systolic blood pressure data. The numbers outside of parentheses are the Monte Carlo average of root mean square error based on 100 independent simulation replicates, and the numbers in parentheses are the corresponding standard error.

noise  $\zeta$ . Finally,  $X_i(t)$  is estimated by  $\hat{X}_i(t) = \hat{\mu}(t) + \sum_{j=1}^{\mathcal{K}} \hat{\xi}_{ij} \hat{\psi}_j(t)$ , where  $\mathcal{K}$  is a tuning parameter whose selection is discussed in Yao et al. (2005).

To illustrate the numerical performance of the proposed method for sparsely and irregularly observed data, we adopt the same setting from Section 4 for dense data, except that now  $m_i \sim 1 + \text{Poisson}(3)$  and  $T_{ij} \sim \text{uniform}(0, 1)$ . In this new setting, the average number of observations per curve is 4 and the observed time points are irregularly scattered. From the results presented in Table S.1, we observe that, the proposed method is comparable to other methods for the  $SO(3)$  manifold while exhibits a clear advantage for the other two manifolds. In addition, as the data are rather sparse, the contamination is expected to dominate the convergence rate in (11) and (12). Thus, we observe that the root mean square error decreases slowly with the sample size, in contrast with the fast rate observed in the case of dense data.

Since the contamination is of a high level in this setting, the structure of the  $SO(3)$  manifold might be buried by the contamination and thus could not be exploited. This might explain why the proposed method shares a similar performance with the functional linear regression or is even slightly outperformed by the latter. Also the performance of sophisticated regression methods like nonparametric regression methods is generally more sensitive to the noise level of the predictor, especially when the predictor resides in a space of higher dimension. This might explain why in the setting of the  $SO(3)$  manifold, almost all nonparametric regression methods listed in Table S.1 perform no better than the functional linear regression which is perhaps the simplest parametric method in functional regression.

## S.2. AUXILIARY EXAMPLE AND RESULTS

*Example 1.* Let  $S^1 = \{v_\omega = (\cos \omega, \sin \omega) : \omega \in [0, 2\pi)\}$  denote the unit circle regarded as a one-dimensional Riemannian manifold. Let  $D = [0, 1]$  and denote  $\phi_1, \phi_2, \dots$  a complete orthonormal basis of  $\mathcal{L}^2(D)$ . Define map  $X(v_\omega) = \sqrt{C} \sum_k k^{-c} \{\cos(k\omega)\phi_{2k-1} + \sin(k\omega)\phi_{2k}\}$  with  $c > 3/2$  and  $C = 1/\sum_k k^{-2c+2} \in (0, \infty)$ . According to Proposition S.1,  $X$  is an isometric embedding of  $S^1$  into  $\mathcal{L}^2(D)$ . Then  $\mathcal{M} = X(S^1)$  is a submanifold of  $\mathcal{L}^2(D)$ . Moreover, *no finite-dimensional* linear subspace of  $\mathcal{L}^2(D)$  fully encompasses  $\mathcal{M}$ . A consequence of this observation is that, a random process taking samples from such  $\mathcal{M}$  might have an infinite number of eigenfunctions, even though  $\mathcal{M}$  is merely one-dimensional, as we shall exhibit in the following. Let us treat  $S^1$  as a probability space endowed with the uniform probability measure, and define random variables  $\xi_{2k-1}(v_\omega) = \sqrt{C}k^{-c} \cos(k\omega)$  and  $\xi_{2k}(v_\omega) = \sqrt{C}k^{-c} \sin(k\omega)$ . Then  $X = \sum_k \xi_k \phi_k$  can be regarded as a random process with samples from  $\mathcal{M}$ . It is easy to check that  $E(\xi_k \xi_j) = 0$  if  $k \neq j$ ,  $E(\xi_k) = 0$ , and  $E\xi_{2k-1}^2 = E\xi_{2k}^2 = C\pi k^{-2c}$ , which implies that  $E(\|X\|_{\mathcal{L}^2}^2) < \infty$ . One can see that the eigenfunctions of the covariance operator of  $X$  are exactly  $\phi_k$ . Therefore,  $X = \sum_k \xi_k \phi_k$  is the Karhunen-Loève expansion of the random process  $X$ , which clearly includes an infinite number of principal components, while  $X$  is intrinsically sampled from the one-dimensional manifold  $\mathcal{M}$ .

**PROPOSITION 1.** *The embedding  $X$  defined in Example 1 is an isometric embedding. Moreover, there is no finite-dimensional linear subspace of  $\mathcal{L}^2(D)$  that fully contains the image  $X(S^1)$ .*

*Proof.* Let  $V = \{(\cos \omega, \sin \omega) : \omega \in (a, b)\}$  be a local neighborhood of  $v$ , and let  $\psi(v) = \omega \in (a, b)$  for  $v = (v_1, v_2) = (\cos \omega, \sin \omega) \in V$ . Then  $\psi$  is a chart of  $S^1$ . Let  $U$  be open in  $\mathcal{L}^2$  such that  $X(v) \in U$ . Since  $\mathcal{L}^2$  is a linear space, the identity map  $I$  serves as a chart. Let  $X_{U,V} : \psi(V) \rightarrow \mathcal{L}^2$  denote the map  $X \circ \psi^{-1}$ . Let  $\vartheta = \sqrt{C} \sum_k k^{-c+1} \{-\sin(k\omega)\phi_{2k-1} +$



$\cos(k\omega)\phi_{2k}\}$ . It defines a linear map from  $\mathbb{R}$  to  $\mathcal{L}^2$ , denoted by  $\Theta(t) = t\vartheta \in \mathcal{L}^2$ . Then,

$$\begin{aligned} A(t) &\equiv t^{-2} \|X_{U,V}(\omega + t) - X_{U,V}(\omega) - \Theta(t)\|^2 \\ &= C \sum_{k=1}^{\infty} \left\{ \frac{k^{-c} \cos(k\omega + kt) - k^{-c} \cos(k\omega) + tk^{-c+1} \sin(k\omega)}{t} \right\}^2 + \\ &\quad C \sum_{k=1}^{\infty} \left\{ \frac{k^{-c} \sin(k\omega + kt) - k^{-c} \sin(k\omega) - tk^{-c+1} \cos(k\omega)}{t} \right\}^2 \\ &\equiv CB_1^2(t) + CB_2^2(t). \end{aligned}$$

By Lipschitz property of the function

$$B_{1,k}(t) \equiv k^{-c} \cos(k\omega + kt) - k^{-c} \cos(k\omega) + tk^{-c+1} \sin(k\omega),$$

we conclude that  $|B_{1,k}(t)| \leq t \sup_t |B'_{1,k}(t)| \leq 2k^{-c+1}t$ . This implies that  $\sup_t B_1^2(t) \leq \sum_k 4k^{-2c+2} < \infty$ . By similar reasoning,  $\sup_t B_2^2(t) < \infty$  and hence  $\sup_t A(t) < \infty$ . We now apply the dominated convergence theorem to conclude that

$$\lim_{t \rightarrow 0} A(t) = C \lim_{t \rightarrow 0} \{B_1^2(t) + B_2^2(t)\} = C \sum_{k=1}^{\infty} \lim_{t \rightarrow 0} \left\{ \frac{B_{1,k}(t)}{t} \right\}^2 + C \sum_{k=1}^{\infty} \lim_{t \rightarrow 0} \left\{ \frac{B_{2,k}(t)}{t} \right\}^2 = 0.$$

By recalling that the tangent space  $T_{v_\omega} S^1$  at  $v_\omega$  is  $\mathbb{R}$  and the tangent space  $T_{X(v_\omega)} \mathcal{L}^2(D)$  at  $X(v_\omega)$  is  $\mathcal{L}^2(D)$ , the above shows that the differential map  $X_{*,v_\omega} : T_{v_\omega} S^1 \rightarrow T_{X(v_\omega)} \mathcal{L}^2(D)$  at  $v_\omega$  is given by the linear map  $\Theta$ , i.e.,

$$X_{*,v_\omega}(t) = \Theta(t) = t \sum_k \sqrt{C} k^{-c+1} \{-\sin(k\omega)\phi_{2k-1} + \cos(k\omega)\phi_{2k}\},$$

and the embedded tangent space at  $v_\omega$  is  $\text{span}\{-\sum_k k^{-c+1} \sin(k\omega)\phi_{2k-1} + \sum_k k^{-c+1} \cos(k\omega)\phi_{2k}\}$ . As this differential map is injective at all  $v \in S^1$ ,  $X$  is indeed an immersion. Since  $S^1$  is compact,  $X$  is also an embedding, and the image  $X(S^1)$  is a submanifold of  $\mathcal{L}^2(D)$ .

To show that  $X$  is isometric, note that the tangent space of  $S^1$  at  $v$  is the real line  $\mathbb{R}$ , equipped with the usual inner product  $\langle s, t \rangle = st$  for  $s, t \in \mathbb{R}$ . Let  $\langle\langle f_1, f_2 \rangle\rangle = \int_D f_1(t)f_2(t)dt$  for  $f_1, f_2 \in \mathcal{L}^2(D)$  denote the canonical inner product of  $\mathcal{L}^2(D)$ . Recall-

ing the definition  $C = 1/\sum_{k=1}^{\infty} k^{-2c+2}$  in the example, we deduce that

$$\begin{aligned}\langle\langle X_{*,v_\omega}(s), X_{*,v_\omega}(t) \rangle\rangle &= \langle\langle \Theta(s), \Theta(t) \rangle\rangle = st \langle\langle \vartheta, \vartheta \rangle\rangle \\ &= Cst \sum_{k=1}^{\infty} \{k^{-2c+2} \sin^2(k\omega) + k^{-2c+2} \cos^2(k\omega)\} \\ &= Cst \sum_{k=1}^{\infty} k^{-2c+2} = st = \langle s, t \rangle,\end{aligned}$$

which shows that  $X$  is isometric.

Finally, to show that there is no finite-dimensional linear subspace of  $\mathcal{L}^2(D)$  that fully contains  $X(S^1)$ , we take the strategy of “proof by contradiction” to assume that  $H$  is a finite-dimensional linear subspace of  $\mathcal{L}^2(D)$  such that  $X(S^1) \subset H$ . Since  $H$  is finite-dimensional, there exists  $0 \neq \varphi \in \mathcal{L}^2(D)$  such that  $\varphi \perp H$  and hence  $\varphi \perp X(S^1)$ , or more specifically,  $\langle\langle \varphi, x \rangle\rangle = 0$  for each  $x \in X(S^1)$ . As  $\phi_1, \phi_2, \dots$  form a complete orthonormal basis of  $\mathcal{L}^2(D)$ , we can find real numbers  $a_1, a_2, \dots$  such that  $\varphi = \sum_k (a_{2k-1} \phi_{2k-1} + a_{2k} \phi_{2k})$ . Then,  $\langle\langle \varphi, x \rangle\rangle = 0$  for each  $x \in X(S^1)$  is equivalent to  $\sum_k k^{-c} \{a_{2k-1} \cos(k\omega) + a_{2k} \sin(k\omega)\} = 0$  for all  $\omega$ . Since  $\cos(k\omega)$  and  $\sin(k\omega)$ , as functions of  $\omega$ , are orthogonal, it implies that  $a_{2k-1} = 0$  and  $a_{2k} = 0$  for all  $k$ , which indicates that  $\varphi = 0$ . However, by assumption,  $\varphi \neq 0$ , and we draw a contradiction.  $\square$

The contamination of the predictor  $X$  poses substantial challenge on the estimation of the manifold structure. For instance, the quality of the tangent space at  $x$ , denoted by  $T_x \mathcal{M}$ , crucially depends on a bona fide neighborhood around  $x$ , while the contaminated neighborhood  $\hat{\mathcal{N}}_{\mathcal{L}^2}(h_{pca}, x)$  and the inaccessible true neighborhood  $\mathcal{N}_{\mathcal{L}^2}(h_{pca}, x) = \{X_i : \|X_i - x\|_{\mathcal{L}^2} < h_{pca}\}$  might contain different observations. Fortunately we can show that they are not far apart in the sense of Proposition 2. In practice, we suggest to choose  $\max(h_{reg}, h_{pca}) < \min\{2/\tau, \text{inj}(\mathcal{M})\}/4$ , where  $\tau$  is the condition number of  $\mathcal{M}$  and  $\text{inj}(\mathcal{M})$  is the injectivity radius of  $\mathcal{M}$  (Cheng & Wu, 2013), so that  $\hat{\mathcal{N}}_{\mathcal{L}^2}(h_{pca}, x)$  provides a good approximation of the true neighborhood of  $x$  within the manifold.

**PROPOSITION 2.** For  $0 < \varrho < \beta$ , define  $h_- = h_{pca} - m^{-(\beta+\varrho)/2}$  and  $h_+ = h_{pca} + m^{-(\beta+\varrho)/2}$ . Let  $Z_i = 1_{\{\hat{X}_i \in \mathbb{B}_{h_{pca}}^{\mathcal{L}^2}(x)\}}$ ,  $V_{i0} = 1_{\{X_i \in \mathbb{B}_{h_-}^{\mathcal{L}^2}(x)\}}$  and  $V_{i1} = 1_{\{X_i \in \mathbb{B}_{h_+}^{\mathcal{L}^2}(x)\}}$ . Under the assumption (B3) and  $\log m \gtrsim \log n$ ,  $\text{pr}(\forall i : V_{i0} \leq Z_i \leq V_{i1}) \rightarrow 1$  as  $n \rightarrow \infty$ .

Hence one can always obtain lower and upper bounds for quantities involving  $Z_i$  in terms of  $V_{i0}$  and  $V_{i1}$ , i.e., with large probability, it is equivalent to substitute  $Z_i$  with  $V_{i0}$  and  $V_{i1}$  in our analysis.

*Proof.* We first bound the following event

$$\begin{aligned} \text{pr}(\forall i : Z_i \leq V_{i1}) &= \prod_{i=1}^n \{1 - \text{pr}(Z_i > V_{i1})\} = \{1 - \text{pr}(Z > V_1)\}^n \\ &= \{1 - \text{pr}(Z = 1, V_1 = 0)\}^n \geq \left\{1 - \text{pr}(\|\hat{X} - X\| \geq m^{-(\beta+\varrho)/2})\right\}^n \\ &\geq \left(1 - c_1^p m^{p(\beta+\varrho)/2} m^{-p\beta}\right)^n \geq \left(1 - c_1^p m^{-2}\right)^n \rightarrow 1, \end{aligned}$$

where  $c_1 > 0$  is some constant, and  $p > 0$  is a constant that is sufficiently large. Similarly, we can deduce that

$$\text{pr}(\forall i : V_{i0} \leq Z_i) \rightarrow 1,$$

and the conclusion  $\text{pr}(\forall i : V_{0i} \leq Z_i \leq V_{1i}) \rightarrow 1$  follows.  $\square$

We now address the case that the predictor  $x$  is not fully observed. It is reasonable to assume that  $x$  comes from the same source of the data, in the sense that its smoothed version  $\tilde{x}$  has the same contamination level as those  $\hat{X}_1, \dots, \hat{X}_n$ , as per (B3). To be specific, assume that

$$(B4) \quad \text{the estimate } \tilde{x} \text{ is independent of } X_1, \dots, X_n \text{ and } \mathbb{X}_1, \dots, \mathbb{X}_n. \text{ Also } \{E\|\tilde{x} - x\|_{\mathcal{L}^2}^p\}^{1/p} \leq C'_p m^{-\beta} \text{ for all } p \geq 1, \text{ where } C'_p \text{ is a constant depending on } p \text{ only.}$$

Note that the independent condition in (B4) is satisfied if  $t_1, \dots, t_{m_x}$  are independent of  $X_1, \dots, X_n$  and  $\mathbb{X}_1, \dots, \mathbb{X}_n$ . The second part of (B4) is met if assumptions similar to (A1)–(A4) hold also for  $x$  and  $t_1, \dots, t_{m_x}$ , according to Proposition 1.

**THEOREM 4.** *With the conditions (A1), (B1)–(B3), and the additional assumption (B4), the equation (11) holds when  $x \in \mathcal{M} \setminus \mathcal{M}_{h_{reg}}$ , and the equation (12) holds when  $x \in \mathcal{M}_{h_{reg}}$ , both with  $\hat{g}(x)$  replaced by  $\hat{g}(\tilde{x})$ .*

*Proof.* We first observe that

$$E \left[ \{\hat{g}(\tilde{x}) - g(x)\}^2 \mid \mathcal{X} \right] \leq 2E \left[ \{\hat{g}(\tilde{x}) - \hat{g}(x)\}^2 \mid \mathcal{X} \right] + 2E \left[ \{\hat{g}(x) - g(x)\}^2 \mid \mathcal{X} \right].$$

To derive the order for the first term, we shall point out that, with Lemma S.7, S.8 and S.9, by following almost the same lines of argument, the conclusions of Theorem 1 hold for  $\tilde{x}$ . This means, by working on  $\tilde{x}$  instead of  $x$ , we still have a consistent estimate of the intrinsic dimension and a good estimate of the tangent space at  $x$ . Given this, it is not difficult but somewhat tedious to verify that the argument in the proof of Theorem 2 and 3 still holds for  $\tilde{x}$ , with care for the discrepancy  $\|\tilde{x} - x\|_{\mathcal{L}^2}$  instead of the discrepancy  $\|\hat{X}_i - X_i\|_{\mathcal{L}^2}$ . This argument also shows that the order of the first term is the same as the second one (this is expected since  $\tilde{x}$  has the same contamination level of those  $\hat{X}_i$ ), and hence the conclusion of the theorem follows.  $\square$

### S.3. PROOFS OF MAIN THEOREMS

To reduce notational burden,  $\mathcal{L}^2(D)$  is simplified by  $\mathcal{L}^2$ , and we shall use  $\|\cdot\|$  to denote the norm  $\|\cdot\|_{\mathcal{L}^2}$  when no confusion arises.

*Proof of Theorem 1.* (a) Without loss of generality, assume  $x = 0$ . Let  $\tilde{G}_j = \hat{G}_j + \Delta$  and  $\tilde{G}_{(1)}, \tilde{G}_{(2)}, \dots, \tilde{G}_{(k)}$  be the associated order statistics of  $\tilde{G}_1, \tilde{G}_2, \dots, \tilde{G}_k$ . Also note that the estimator in Levina & Bickel (2004) is still consistent if  $G$  is replaced with  $\check{G} \equiv G + \Delta$ . Then,

$$\begin{aligned} & \left| \frac{1}{k-1} \sum_{j=1}^{k-1} \log \frac{\hat{G}_{(k)} + \Delta}{\hat{G}_{(j)} + \Delta} - \frac{1}{k-1} \sum_{j=1}^{k-1} \log \frac{\check{G}_{(k)}}{\check{G}_{(j)}} \right| \\ & \leq \left| \log \tilde{G}_{(k)} - \log \check{G}_{(k)} \right| + \left| \frac{1}{k-1} \sum_{j=1}^k \left( \log \tilde{G}_{(j)} - \log \check{G}_{(j)} \right) \right| \equiv I_1 + I_2. \end{aligned} \quad (1)$$

For  $I_1$ , let  $q$  and  $p$  be the indices such that  $\tilde{G}_{(k)} = \hat{G}_q$  and  $\check{G}_p = \check{G}_{(k)}$ , respectively. For the case  $q = p$ , we have  $|\tilde{G}_{(k)} - \check{G}_{(k)}| = |\tilde{G}_p - \check{G}_p| \leq \left| \|\hat{X}_p\| - \|X_p\| \right| + \Delta \leq \|\hat{X}_p - X_p\| + \Delta$  by reverse triangle inequality. When  $q \neq p$ , it is seen that  $\tilde{G}_p < \tilde{G}_{(k)} = \tilde{G}_q$  and  $G_q < G_p = G_{(k)}$ . If  $\tilde{G}_{(k)} > \check{G}_{(k)}$ , then  $|\tilde{G}_{(k)} - \check{G}_{(k)}| \leq |\tilde{G}_{(k)} - \check{G}_q| = |\tilde{G}_q - G_q| \leq \max_{1 \leq j \leq k} \{\|\hat{X}_j - X_j\|\}$ . Otherwise,  $|\tilde{G}_{(k)} - \check{G}_{(k)}| \leq |\tilde{G}_p - \check{G}_p| \leq \max_{1 \leq j \leq k} \{\|\hat{X}_j - X_j\|\}$ . Now,  $\text{pr}(\forall 1 \leq j \leq k : \|\hat{X}_j - X_j\| > \epsilon) \leq \sum_{j=1}^k \text{pr}(\|\hat{X}_j - X_j\| > \epsilon) \leq kE\|\hat{X}_j - X_j\|^r \epsilon^{-r} = O(km^{-r\beta}) = o(1)$  for a sufficiently large constant  $r$ . Therefore,  $|\tilde{G}_{(k)} - \check{G}_{(k)}|$  converges to zero in probability, or  $\tilde{G}_{(k)}$  converges to  $\check{G}_{(k)}$  in probability. By Slutsky's lemma,  $\log \tilde{G}_{(k)}$  converges to  $\log \check{G}_{(k)}$  in probability and hence  $I_1 = o_P(1)$ .

For  $I_2$ , we first observe that

$$I_2 = \left| \frac{1}{k-1} \sum_{j=1}^k \left( \log \tilde{G}_{(j)} - \log \check{G}_{(j)} \right) \right| = \left| \frac{1}{k-1} \sum_{j=1}^k \left( \log \tilde{G}_j - \log \check{G}_j \right) \right|.$$

By Markov's inequality, for any fixed  $\epsilon > 0$ ,

$$\text{pr}(I_2 > \epsilon) \leq \frac{EI_2}{\epsilon} \leq \frac{kE|\log \tilde{G} - \log \check{G}|}{(k-1)\epsilon} = o(1),$$

where the last equality is obtained by Lemma S.7. We then deduce that  $I_2 = o_P(1)$ .

Together with  $I_1 = o_P(1)$  and (1), this implies that

$$\left| \frac{1}{k-1} \sum_{j=1}^{k-1} \log \frac{\hat{G}_{(k)} + \Delta}{\hat{G}_{(j)} + \Delta} - \frac{1}{k-1} \sum_{j=1}^{k-1} \log \frac{\check{G}_{(k)}}{\check{G}_{(j)}} \right| \rightarrow 0 \quad \text{in probability.}$$

Now we apply the argument in Levina & Bickel (2004) to conclude that  $\hat{d}$  is a consistent estimator.

(b) Let  $h = h_{pca}$ , and  $\{\tilde{\phi}_k\}_{k=1}^d$  be an orthonormal basis system for  $T_x \mathcal{M}$  and  $\{\psi_k\}_{k=1}^\infty$  be an orthonormal basis of  $\mathcal{L}^2$ . Without loss of generality, assume that  $\mathcal{M}$  is properly rotated and translated so that  $\psi_k = \tilde{\phi}_k$  for  $k = 1, 2, \dots, d$ , and  $x = 0 \in \mathcal{L}^2$ . The sample covariance operator based on observations in  $\hat{\mathcal{N}}_{\mathcal{L}^2}(h, x)$  is denoted by  $\hat{\mathcal{C}}_x$  as in (6). It is seen that  $\hat{\mathcal{C}}_x = n^{-1} \sum_{i=1}^n (\hat{X}_i - \hat{\mu}_x)(\hat{X}_i - \hat{\mu}_x)Z_i$ , where  $Z_i = 1_{\{\hat{X}_i \in \mathbb{B}_h^{\mathcal{L}^2}(x)\}}$  and  $\hat{\mu}_x = n^{-1} \sum_{i=1}^n \hat{X}_i Z_i$ . Let  $\mathcal{H}_1 = \text{span}\{\psi_k : k = 1, 2, \dots, d\}$  and  $\mathcal{H}_2$  be the complementary subspace of  $\mathcal{H}_1$  in  $\mathcal{L}^2$ , so that  $\mathcal{L}^2 = \mathcal{H}_1 \oplus \mathcal{H}_2$ . Let  $\mathcal{P}_j : \mathcal{L}^2 \rightarrow \mathcal{H}_j$  be projection operators, and we define operator  $\mathcal{A} = \mathcal{P}_1 \hat{\mathcal{C}}_x \mathcal{P}_1$ ,  $\mathcal{B} = \mathcal{P}_2 \hat{\mathcal{C}}_x \mathcal{P}_2$ ,  $\mathcal{D}_{12} = \mathcal{P}_1 \hat{\mathcal{C}}_x \mathcal{P}_2$  and  $\mathcal{D}_{21} = \mathcal{P}_2 \hat{\mathcal{C}}_x \mathcal{P}_1$ . Then  $\hat{\mathcal{C}}_x = \mathcal{A} + \mathcal{B} + \mathcal{D}_{12} + \mathcal{D}_{21}$ . Note that  $\mathcal{D}_{12} + \mathcal{D}_{21}$  is self-adjoint. Therefore, if  $y = \sum_{k=1}^\infty a_k \psi_k \in \mathcal{L}^2$ ,

$$\begin{aligned} \|\mathcal{D}_{12} + \mathcal{D}_{21}\|_{op} &= \sup_{\|y\|=1} \langle (\mathcal{D}_{12} + \mathcal{D}_{21})y, y \rangle = \sup_{\|y\|=1} \left( \langle \mathcal{P}_1 \hat{\mathcal{C}}_x \mathcal{P}_2 y, y \rangle + \langle \mathcal{P}_2 \hat{\mathcal{C}}_x \mathcal{P}_1 y, y \rangle \right) \\ &= 2 \sup_{\|y\|=1} \left( \sum_{k=d+1}^\infty \sum_{j=1}^d a_j a_k \langle \hat{\mathcal{C}}_x \psi_j, \psi_k \rangle \right) \leq 2 \sup_{\|y\|=1} \left( \sum_{k=d+1}^\infty \sum_{j=1}^d |a_j a_k| \cdot \left| \langle \hat{\mathcal{C}}_x \psi_j, \psi_k \rangle \right| \right) \\ &\leq 2 \sup_{j \leq d} \sup_{k \geq d+1} \left| \langle \hat{\mathcal{C}}_x \psi_j, \psi_k \rangle \right| \sup_{\|y\|=1} \left\{ \sum_{k=d+1}^\infty \sum_{j=1}^d (a_j^2 + a_k^2) \right\} \leq 2 \sup_{j \leq d} \sup_{k \geq d+1} \left| \langle \hat{\mathcal{C}}_x \psi_j, \psi_k \rangle \right|. \end{aligned}$$

From Lemma S.9,  $\|\mathcal{D}_{12} + \mathcal{D}_{21}\|_{op} = O_P(h^{d+3} + n^{-1/2}h^{d/2+3} + m^{-\beta}h^{d+1})$ . Similarly, we have  $\|\mathcal{B}\|_{op} = O_P(h^{d+4} + n^{-1/2}h^{d/2+4} + m^{-\beta}h^{d+1})$ , and  $\mathcal{A} = \pi_{d-1}f(x)d^{-1}h^{d+2}I_d +$

$O_P\left(n^{-1/2}h^{d/2+2} + m^{-\beta}h^{d+1}\right)$ , where  $\pi_{d-1}$  is the volume of the  $d-1$  dimensional unit sphere, and  $I_d$  is the identity operator on  $\mathcal{H}_1$ .

Let  $a_n = n^{-1/2}h^{-d/2}$  and  $b_n = m^{-\beta}h^{-1}$ . Then we have

$$\hat{\mathcal{C}}_x = \pi_{d-1}f(x)d^{-1}h^{d+2}\{I_d + O_P(a_n + b_n)\tilde{\mathcal{A}} + O_P(h^2 + b_n)\tilde{\mathcal{B}} + O_P(h + b_n)(\tilde{\mathcal{D}}_{12} + \tilde{\mathcal{D}}_{21})\}$$

where  $\tilde{\mathcal{A}}$ ,  $\tilde{\mathcal{B}}$ ,  $\tilde{\mathcal{D}}_{12}$  and  $\tilde{\mathcal{D}}_{21}$  are operators with norm equal to one, and  $\tilde{\mathcal{D}}_{12}$  is the adjoint of  $\tilde{\mathcal{D}}_{21}$ . With the choice of  $\varrho$ , we have  $\hat{\mathcal{C}}_x = \pi_{d-1}f(x)d^{-1}h^{d+2}\{I_d + O_P(\sqrt{h})\tilde{\mathcal{A}} + O_P(h)\tilde{\mathcal{B}} + O_P(h)(\tilde{\mathcal{D}}_{12} + \tilde{\mathcal{D}}_{21})\}$ . The same perturbation argument done in Singer & Wu (2012) leads to the desired result.  $\square$

*Proof of Theorem 2.* To reduce notions, let  $h = h_{reg}$  and fix  $x \in \mathcal{M} \setminus \mathcal{M}_{h_{reg}}$ . Let  $\{\hat{\varphi}_k\}_{k=1}^d$  be the orthonormal set determined by local FPCA and  $\{\phi_k\}_{k=1}^d$  the associated orthonormal basis of  $T_x\mathcal{M}$ . Let  $\{\psi_k\}_{k=1}^\infty$  be an orthonormal basis of  $\mathcal{L}^2$ . Without loss of generality, assume  $\mathcal{M}$  is properly rotated and translated so that  $x = 0 \in \mathcal{L}^2$  and  $\psi_k = \phi_k$  for  $k = 1, 2, \dots, d$ . Let  $\mathbf{g} = (g(X_1), g(X_2), \dots, g(X_n))^T$ . Then we have

$$E\{\hat{g}(x) \mid \mathcal{X}\} = e_1^T(\hat{Q}^T \hat{W} \hat{Q})^{-1} \hat{Q}^T \hat{W} \mathbf{g}.$$

Take  $z = \exp_x(t\theta)$ , where  $t = O(h)$ ,  $\theta \in T_x\mathcal{M}$ ,  $\|\theta\|_{\mathcal{L}^2} = 1$ , and  $\exp_x$  denotes the exponential map of  $\mathcal{M}$  at  $x$ . By Theorem 1, we have  $\langle \theta, \hat{\varphi}_k \rangle = \langle \theta, \psi_k \rangle + O_P(h_{pca}^{3/2})$  and  $\langle \Pi_x(\theta, \theta), \hat{\varphi}_k \rangle = O_P(h_{pca})$ . By Lemma A.2.2. of Cheng & Wu (2013), we have

$$t\theta = y - t^2\Pi_x(\theta, \theta)/2 + O(t^3). \quad (2)$$

Therefore, for  $k = 1, 2, \dots, d$ ,  $\langle t\theta, \psi_k \rangle = \langle t\theta, \hat{\varphi}_k - O_P(h_{pca}^{3/2})u_k \rangle = \langle z, \hat{\varphi}_k \rangle - t^2\langle \Pi_x(\theta, \theta), \hat{\varphi}_k \rangle/2 + O_P(hh_{pca}^{3/2} + h^2h_{pca}) = \langle z, \hat{\varphi}_k \rangle + O_P(hh_{pca}^{3/2} + h^2h_{pca})$ . Since  $\theta \in T_x\mathcal{M}$ , we have  $\theta = \sum_{k=1}^d \langle \theta, \psi_k \rangle \phi_k$ . Let  $\mathbf{z} = (\langle z, \hat{\varphi}_1 \rangle, \langle z, \hat{\varphi}_2 \rangle, \dots, \langle z, \hat{\varphi}_d \rangle)^T$ . By (2), it is easy to see that

$$\begin{aligned} g(z) - g(x) &= t\theta \nabla g(x) + \text{Hess } g(x)(\theta, \theta)t^2/2 + O_P(t^3) \\ &= \sum_{k=1}^d \langle t\theta, \psi_k \rangle \nabla_{\phi_k} g(x) + \frac{1}{2} \sum_{j,k=1}^d \langle t\theta, \psi_j \rangle \langle t\theta, \psi_k \rangle \text{Hess } g(x)(\phi_j, \phi_k) + O_P(h^3) \\ &= \mathbf{z}^T \nabla g(x) + \frac{1}{2} \mathbf{z}^T \text{Hess } g(x) \mathbf{z} + O_P(h^{5/2}). \end{aligned}$$

Due to the smoothness of  $g$ , the compactness of  $\mathcal{M}$  and the compact support of  $K$ , we have  $\mathbf{g} = Q[g(x) \nabla g(x)]^T + H/2 + O_P(h^{5/2})$ , where  $H = [\xi_1^T \text{Hess } g(x) \xi_1, \xi_2^T \text{Hess } g(x) \xi_2, \dots, \xi_n^T \text{Hess } g(x) \xi_n]^T$ ,  $\xi_i = (\xi_{i1}, \dots, \xi_{id})^T$ ,  $\xi_{ij} = \langle X_i, \psi_j \rangle$ , and

$$Q = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ \xi_1 & \xi_2 & \cdots & \xi_n \end{pmatrix}^T.$$

Then the conditional bias is

$$\begin{aligned} E\{\hat{g}(x) - g(x) \mid \mathcal{X}\} &= e_1^T (\hat{Q}^T \hat{W} \hat{Q})^{-1} \hat{Q}^T \hat{W} \mathbf{g} - g(x) \\ &= e_1^T \left( \frac{1}{n} \hat{Q}^T \hat{W} \hat{Q} \right)^{-1} \frac{1}{n} \hat{Q}^T \hat{W} (Q - \hat{Q}) \begin{bmatrix} g(x) \\ \nabla g(x) \end{bmatrix} \end{aligned} \quad (3)$$

$$+ e_1^T \left( \frac{1}{n} \hat{Q}^T \hat{W} \hat{Q} \right)^{-1} \frac{1}{n} \hat{Q}^T \hat{W} \left\{ \frac{1}{2} H + O_P(h^{5/2}) \right\}. \quad (4)$$

Now we analyze the term in (3). Let  $Z = 1_{\hat{X} \in \mathbb{B}_h^{\mathcal{L}^2}(x)}$ . By Lemma S.8,  $EZ \asymp h^d$ . Then, by Hölder's inequality, for any fixed  $\epsilon > 0$ , we choose a constant  $q > 1$  and a sufficiently large  $p > 0$  so that  $1/q + 1/p = 1$  and  $EZ \|\hat{X} - X\| = (EZ)^{1/q} (E\|\hat{X} - X\|^p)^{1/p} = O(h^{d-\epsilon d} m^{-\beta})$ . Therefore,

$$\frac{1}{n} \hat{Q}^T \hat{W} (Q - \hat{Q}) \begin{bmatrix} g(x) \\ \nabla g(x) \end{bmatrix} = \begin{bmatrix} \frac{1}{n} \sum_{i=1}^n K_h(\|\hat{X}_i - x\|) (\xi_i - \hat{\xi}_i)^T \nabla g(x) \\ \frac{1}{n} \sum_{i=1}^n K_h(\|\hat{X}_i - x\|) (\xi_i - \hat{\xi}_i)^T \nabla g(x) \hat{\xi}_i \end{bmatrix} = O_P(h^{-1-\epsilon d} m^{-\beta}), \quad (5)$$

since

$$\begin{aligned} \left| \frac{1}{n} \sum_{i=1}^n K_h(\|\hat{X}_i - x\|) (\xi_i - \hat{\xi}_i)^T \nabla g(x) \right| &\leq \frac{1}{n} \sum_{i=1}^n K_h(\|\hat{X}_i - x\|) \|\xi_i - \hat{\xi}_i\|_{\mathbb{R}^d} \|\nabla g(x)\| \\ &\leq \left\{ \sup_v |K(v)| \right\} \|\nabla g(x)\| \left( \frac{1}{n} \sum_{i=1}^n Z_i \|\xi_i - \hat{\xi}_i\|_{\mathbb{R}^d} \right) = O_P(h^{-1-\epsilon d} m^{-\beta}), \end{aligned}$$

and similarly,  $n^{-1} \sum_{i=1}^n K_h(\|\hat{X}_i - x\|) (\xi_i - \hat{\xi}_i)^T \nabla g(x) \hat{\xi}_i = O_P(h^{-1-\epsilon d} m^{-\beta}) \mathbb{1}_{d \times 1}$ .

For  $\hat{Q}^T \hat{W} \hat{Q}$ , a direct calculation shows that

$$\frac{1}{n} \hat{Q}^T \hat{W} \hat{Q} = \begin{bmatrix} n^{-1} \sum_{i=1}^n K_h(\|\hat{X}_i - x\|) & n^{-1} \sum_{i=1}^n K_h(\|\hat{X}_i - x\|) \hat{\xi}_i^T \\ n^{-1} \sum_{i=1}^n K_h(\|\hat{X}_i - x\|) \hat{\xi}_i & n^{-1} \sum_{i=1}^n \hat{\xi}_i^T K_h(\|\hat{X}_i - x\|) \hat{\xi}_i \end{bmatrix}.$$

It is easy to check that  $n^{-1} \sum_{i=1}^n K_h(\|\hat{X}_i - x\|) = n^{-1} \sum_{i=1}^n K_h(\|X_i - x\|) + O_P(h^{-1-\epsilon d} m^{-\beta})$ , and note that the choice of  $h$  ensures that  $h^{1+\epsilon d} \gg m^{-\beta}$ . Similar calcu-

lation shows that  $\frac{1}{n} \sum_{i=1}^n K_h(\|\hat{X}_i - x\|) \hat{\xi}_i^T = \frac{1}{n} \sum_{i=1}^n K_h(\|X_i - x\|) \xi_i^T + O_P(h^{-1-\epsilon d} m^{-\beta})$  and also

$$\frac{1}{n} \sum_{i=1}^n \hat{\xi}_i^T K_h(\|\hat{X}_i - x\|) \hat{\xi}_i = \frac{1}{n} \sum_{i=1}^n \xi_i^T K_h(\|X_i - x\|) \xi_i + O_P(h^{-1-\epsilon d} m^{-\beta}).$$

Therefore,

$$\frac{1}{n} \hat{Q}^T \hat{W} \hat{Q} = \frac{1}{n} Q^T W Q + O_P(h^{-1-\epsilon d} m^{-\beta}) \mathbb{1}_{d \times 1} \mathbb{1}_{d \times 1}^T, \quad (6)$$

with

$$\frac{1}{n} Q^T W Q = \begin{bmatrix} n^{-1} \sum_{i=1}^n K_h(\|X_i - x\|) & n^{-1} \sum_{i=1}^n K_h(\|X_i - x\|) \xi_i^T \\ n^{-1} \sum_{i=1}^n K_h(\|X_i - x\|) \xi_i & n^{-1} \sum_{i=1}^n \xi_i K_h(\|X_i - x\|) \xi_i^T \end{bmatrix}.$$

By Lemma S.4, S.5 and S.6, we have

$$\begin{aligned} & \frac{1}{n} Q^T W Q \\ &= \begin{bmatrix} f(x) & h^2 u_{1,2} d^{-1} \nabla f(x)^T \\ h^2 u_{1,2} d^{-1} \nabla f(x) & h^2 u_{1,2} d^{-1} f(x) I_d \end{bmatrix} + \begin{bmatrix} O(h^2) + O_P(n^{-\frac{1}{2}} h^{-\frac{d}{2}}) & O(h^3) + O_P(n^{-\frac{1}{2}} h^{-\frac{d}{2}+1}) \\ O(h^3) + O_P(n^{-\frac{1}{2}} h^{-\frac{d}{2}+1}) & O(h^{7/2}) + O_P(n^{-\frac{1}{2}} h^{-\frac{d}{2}+2}) \end{bmatrix}, \end{aligned}$$

where  $u_{q,k}$  are constants defined in Section S.2. Therefore, combined with (6), it yields

$$\begin{aligned} \frac{1}{n} \hat{Q}^T \hat{W} \hat{Q} &= \begin{bmatrix} f(x) & h^2 u_{1,2} d^{-1} \nabla f(x)^T \\ h^2 u_{1,2} d^{-1} \nabla f(x) & h^2 u_{1,2} d^{-1} f(x) I_d \end{bmatrix} \\ &+ \begin{bmatrix} O(h^2) + O_P(n^{-\frac{1}{2}} h^{-\frac{d}{2}}) + O_P(h^{-1-\epsilon d} m^{-\beta}) & O(h^3) + O_P(n^{-\frac{1}{2}} h^{-\frac{d}{2}+1}) + O_P(h^{-1-\epsilon d} m^{-\beta}) \\ O(h^3) + O_P(n^{-\frac{1}{2}} h^{-\frac{d}{2}+1}) + O_P(h^{-1-\epsilon d} m^{-\beta}) & O(h^{7/2}) + O_P(n^{-\frac{1}{2}} h^{-\frac{d}{2}+2}) + O_P(h^{-1-\epsilon d} m^{-\beta}) \end{bmatrix}. \end{aligned}$$

Then, by the binomial inverse theorem and matrix blockwise inversion, we have

$$\begin{aligned} \left( \frac{1}{n} \hat{Q}^T \hat{W} \hat{Q} \right)^{-1} &= \begin{bmatrix} \frac{1}{f(x)} & -\frac{\nabla^T f(x)}{f(x)^2} \\ -\frac{\nabla f(x)}{f(x)^2} & h^{-2} \frac{d}{u_{1,2} f(x)} I_d \end{bmatrix} \\ &+ \begin{bmatrix} O_P(h^2 + h^{-1-\epsilon d} m^{-\beta} + n^{-\frac{1}{2}} h^{-\frac{d}{2}}) & O_P(h + h^{-3-\epsilon d} m^{-\beta} + n^{-\frac{1}{2}} h^{-\frac{d}{2}-1}) \\ O_P(h + h^{-3-\epsilon d} m^{-\beta} + n^{-\frac{1}{2}} h^{-\frac{d}{2}-1}) & O_P(h^{-\frac{1}{2}} + h^{-5-\epsilon d} m^{-\beta} + n^{-\frac{1}{2}} h^{-\frac{d}{2}-2}) \end{bmatrix}. \end{aligned} \quad (7)$$

Together with (5), it implies that

$$e_1^T (\hat{Q}^T \hat{W} \hat{Q})^{-1} \hat{Q}^T \hat{W} \left\{ (Q - \hat{Q}) \begin{bmatrix} g(x) \\ \nabla g(x) \end{bmatrix} \right\} = O_P(h^{-1-\epsilon d} m^{-\beta}). \quad (8)$$



Now we analyze (4) with a focus on the term  $\hat{Q}^T \hat{W} H$ . A calculation similar to those in Lemma S.5 and S.6 shows that  $\frac{1}{n} \sum_{i=1}^n K_h(\|X_i - x\|) \xi_i^T \text{Hess } g(x) \xi_i = h^2 u_{1,2} d^{-1} f(x) \Delta g(x) + O_P(h^{7/2} + n^{-1/2} h^{-d/2+2})$  and  $\frac{1}{n} \sum_{i=1}^n K_h(\|X_i - x\|) \xi_i^T \text{Hess } g(x) \xi_i \xi_i = O_P(h^4 + n^{-\frac{1}{2}} h^{-\frac{d}{2}+3})$ . Therefore,

$$\frac{1}{n} Q^T W H = \begin{bmatrix} h^2 u_{1,2} d^{-1} f(x) \Delta g(x) + O_P(h^{7/2} + n^{-1/2} h^{-d/2+2}) \\ h^4 + n^{-\frac{1}{2}} h^{-\frac{d}{2}+3} \end{bmatrix}$$

and hence

$$\frac{1}{n} \hat{Q}^T \hat{W} H = \begin{bmatrix} h^2 u_{1,2} d^{-1} f(x) \Delta g(x) + O_P(h^{7/2} + n^{-1/2} h^{-d/2+2} + h^{-1-\epsilon d} m^{-\beta}) \\ O_P(h^4 + n^{-\frac{1}{2}} h^{-\frac{d}{2}+3} + h^{-1-\epsilon d} m^{-\beta}) \end{bmatrix}.$$

The condition on  $h$  implies that  $n^{-\frac{1}{2}} h^{-\frac{d}{2}} \ll 1$ . With (7), we conclude that

$$e_1^T (\hat{Q}^T \hat{W} \hat{Q})^{-1} \hat{Q}^T \hat{W} \left\{ \frac{1}{2} H + O_P(h^3) \right\} = \frac{1}{2d} h^2 u_{1,2} \Delta g(x) + O_P(h^3 + n^{-\frac{1}{2}} h^{-\frac{d}{2}+2} + h^{-1-\epsilon d} m^{-\beta}).$$

Combining this equation with (4) and (8), we immediately see that the conditional bias is

$$E\{\hat{g}(x) - g(x) \mid \mathcal{X}\} = \frac{1}{2d} h^2 u_{1,2} \Delta g(x) + O_P(h^3 + n^{-\frac{1}{2}} h^{-\frac{d}{2}+2} + h^{-1-\epsilon d} m^{-\beta}). \quad (9)$$

Now we analyze the conditional variance. Simple calculation shows that

$$\text{var}\{\hat{g}(x) \mid \mathcal{X}\} = n^{-1} \sigma_\varepsilon^2 e_1^T (n^{-1} \hat{Q}^T \hat{W} \hat{Q})^{-1} (n^{-1} \hat{Q}^T \hat{W} \hat{W} \hat{Q}) (n^{-1} \hat{Q}^T \hat{W} \hat{Q})^{-1} e_1^T \quad (10)$$

and

$$\frac{1}{n} \hat{Q}^T \hat{W} \hat{W} \hat{Q} = \frac{1}{n} Q^T W W Q + O_P(m^{-\beta} h^{-d-1-\epsilon d}) \mathbb{1}_{(d+1) \times (d+1)}. \quad (11)$$

In addition,

$$\frac{1}{n} Q^T W W Q = \begin{bmatrix} \frac{1}{n} \sum_{i=1}^n K_h^2(\|X_i - x\|) & \frac{1}{n} \sum_{i=1}^n K_h^2(\|X_i - x\|) \xi_i^T \\ \frac{1}{n} \sum_{i=1}^n K_h^2(\|X_i - x\|) \xi_i & \frac{1}{n} \sum_{i=1}^n K_h^2(\|X_i - x\|) \xi_i \xi_i^T \end{bmatrix}.$$

With Lemma S.4, S.5 and S.6, we can show that

$$\begin{aligned} & \frac{h^d}{n} Q^T W W Q \\ &= \begin{bmatrix} u_{2,0} \sigma^2 f(x) & h^2 d^{-1} u_{2,2} \sigma^2 \nabla f(x) \\ h^2 d^{-1} u_{2,2} \sigma^2 \nabla^T f(x) & h^2 d^{-1} u_{2,2} \sigma^2 f(x) I_d \end{bmatrix} + \begin{bmatrix} O(h^2) + O_P(n^{-\frac{1}{2}} h^{-\frac{d}{2}}) & O_P(h^3 + n^{-\frac{1}{2}} h^{-\frac{d}{2}+1}) \\ O_P(h^3 + n^{-\frac{1}{2}} h^{-\frac{d}{2}+1}) & O_P(h^{\frac{7}{2}} + n^{-\frac{1}{2}} h^{-\frac{d}{2}+2}) \end{bmatrix}. \end{aligned}$$

Combined with (7), the above equation implies that

$$n^{-1}\sigma_\varepsilon^2 e_1^T (n^{-1}\hat{Q}^T \hat{W} \hat{Q})^{-1} (n^{-1}Q^T W W Q) (n^{-1}\hat{Q}^T \hat{W} \hat{Q})^{-1} e_1^T = O_P(n^{-1}h^{-d}). \quad (12)$$

Also,

$$\begin{aligned} n^{-1}\sigma_\varepsilon^2 e_1^T (n^{-1}\hat{Q}^T \hat{W} \hat{Q})^{-1} \mathbb{1}_{(d+1) \times (d+1)} (n^{-1}\hat{Q}^T \hat{W} \hat{Q})^{-1} e_1^T O_P(h^{-d-1-\epsilon d} m^{-\beta}) \\ = O_P\left(m^{-\beta} n^{-1} h^{-d-1-\epsilon d} (1 + h^{-2-\epsilon d} m^{-\beta} + h^{-4-2\epsilon d} m^{-2\beta})\right). \end{aligned} \quad (13)$$

Combining the above result with (10), (11), (12), (13) and the condition on  $h$ , gives the following conditional variance

$$\text{var}\{\hat{g}(x) \mid \mathcal{X}\} = \frac{1}{nh^d} \frac{u_{2,0}\sigma_\varepsilon^2}{f(x)} + O_P\left(n^{-1}h^{-d}(h + n^{-\frac{1}{2}}h^{-\frac{d}{2}})\right). \quad (14)$$

Finally, the rate for  $E[\{\hat{g}(x) - g(x)\}^2 \mid \mathcal{X}]$  is derived from (9) and (14).  $\square$

*Proof of Theorem 3.* The proof is similar to the proof for Theorem 2. Below we shall only discuss those that are different. Let  $h = h_{reg}$  to reduce notational burden. We first have

$$\begin{aligned} \frac{1}{n} Q^T W Q &= f(x) v \kappa_1 v \\ &+ \begin{bmatrix} O(h) + O_P(n^{-\frac{1}{2}}h^{-\frac{d}{2}}) & O(h^2) + O_P(n^{-\frac{1}{2}}h^{-\frac{d}{2}+1}) \\ O(h^2) + O_P(n^{-\frac{1}{2}}h^{-\frac{d}{2}+1}) & O(h^3) + O_P(n^{-\frac{1}{2}}h^{-\frac{d}{2}+2}) \end{bmatrix} \end{aligned}$$

where

$$\kappa_q = \begin{bmatrix} \kappa_{11,q} & \kappa_{12,q} \\ \kappa_{12,q}^T & \kappa_{22,q} \end{bmatrix}, \kappa_{22,q} = (\kappa_{22,q,j,k})_{j,k=1}^d, v = \begin{bmatrix} 1 & 0 \\ 0 & hI_d \end{bmatrix}.$$

Then,

$$\begin{aligned} \frac{1}{n} \hat{Q}^T \hat{W} \hat{Q} &= f(x) v \kappa_1 v \\ &+ \begin{bmatrix} O(h) + O_P(n^{-\frac{1}{2}}h^{-\frac{d}{2}}) + O_P(h^{-1-\epsilon d} m^{-\beta}) & O(h^2) + O_P(n^{-\frac{1}{2}}h^{-\frac{d}{2}+1}) + O_P(h^{-1-\epsilon d} m^{-\beta}) \\ O(h^2) + O_P(n^{-\frac{1}{2}}h^{-\frac{d}{2}+1}) + O_P(h^{-1-\epsilon d} m^{-\beta}) & O(h^3) + O_P(n^{-\frac{1}{2}}h^{-\frac{d}{2}+2}) + O_P(h^{-1-\epsilon d} m^{-\beta}) \end{bmatrix} \end{aligned}$$

and also

$$\begin{aligned} \left(\frac{1}{n}\hat{Q}^T\hat{W}\hat{Q}\right)^{-1} &= v^{-1}\kappa_1^{-1}v^{-1}/f(x) \\ &+ \begin{bmatrix} O_P(h + h^{-2-\epsilon d}m^{-\beta} + n^{-\frac{1}{2}}h^{-\frac{d}{2}}) & O_P(1 + h^{-4-\epsilon d}m^{-\beta} + n^{-\frac{1}{2}}h^{-\frac{d}{2}-1}) \\ O_P(1 + h^{-4-\epsilon d}m^{-\beta} + n^{-\frac{1}{2}}h^{-\frac{d}{2}-1}) & O_P(h^{-1} + h^{-5-\epsilon d}m^{-\beta} + n^{-\frac{1}{2}}h^{-\frac{d}{2}-2}) \end{bmatrix}. \end{aligned}$$

This implies that

$$\begin{aligned} e_1^T(\hat{Q}^T\hat{W}\hat{Q})^{-1}\hat{Q}^T\hat{W}\left\{(Q - \hat{Q})\begin{bmatrix} g(x) \\ \nabla g(x) \end{bmatrix}\right\} &= e_1^T\begin{bmatrix} 1 & h^{-1} \\ h^{-1} & h^{-2} \end{bmatrix}O_P(h^{-1-\epsilon d}m^{-\beta})\mathbb{1}_{d \times 1} \\ &= O_P(h^{-1-\epsilon d}m^{-\beta}). \end{aligned}$$

By Lemma (4), (5) and (6), one can show that

$$\frac{1}{n}Q^TW H = \begin{bmatrix} h^2 f(x) \int_{\frac{1}{h}\mathbb{D}_x} K(\|\theta\|_{\mathbb{R}^d}) \text{Hess } g(x) \theta d\theta + O_P(h^3 + n^{-1/2}h^{-d/2+2}) \\ h^3 f(x) \int_{\frac{1}{h}\mathbb{D}_x} K(\|\theta\|_{\mathbb{R}^d}) \theta^T \text{Hess } g(x) \theta \theta d\theta + O_P(h^{7/2} + n^{-\frac{1}{2}}h^{-\frac{d}{2}+3}) \end{bmatrix},$$

and hence

$$\frac{1}{n}\hat{Q}^T\hat{W}H = \begin{bmatrix} h^2 f(x) \int_{\frac{1}{h}\mathbb{D}_x} K(\|\theta\|_{\mathbb{R}^d}) \text{Hess } g(x) \theta d\theta + O_P(h^3 + n^{-1/2}h^{-d/2+2} + h^{-1-\epsilon d}m^{-\beta}) \\ h^3 f(x) \int_{\frac{1}{h}\mathbb{D}_x} K(\|\theta\|_{\mathbb{R}^d}) \theta^T \text{Hess } g(x) \theta \theta d\theta + O_P(h^{7/2} + n^{-\frac{1}{2}}h^{-\frac{d}{2}+3} + h^{-1-\epsilon d}m^{-\beta}) \end{bmatrix}.$$

With the condition on  $h$ , from the above we deduce that

$$\begin{aligned} e_1^T(\hat{Q}^T\hat{W}\hat{Q})^{-1}\hat{Q}^T\hat{W}\left\{\frac{1}{2}H(x) + O_P(h^3)\right\} &= h^2 \frac{\text{tr}(\text{Hess } g(x)\kappa_{22,1})}{2(\kappa_{11,1} - \kappa_{12,1}^T \kappa_{22,1}^{-1} \kappa_{12,1})} \\ &+ O_P(h^{5/2} + n^{-\frac{1}{2}}h^{-\frac{d}{2}+2} + h^{-2-\epsilon d}m^{-\beta}). \end{aligned}$$

Then, the conditional bias is

$$E\{\hat{g}(x) - g(x) \mid \mathcal{X}\} = h^2 \frac{\text{tr}(\text{Hess } g(x)\kappa_{22,1})}{2(\kappa_{11,1} - \kappa_{12,1}^T \kappa_{22,1}^{-1} \kappa_{12,1})} + O_P(h^{5/2} + n^{-\frac{1}{2}}h^{-\frac{d}{2}+2} + h^{-2-\epsilon d}m^{-\beta}).$$

For the conditional variance, by Lemma 4, 5 and 6, we have

$$\frac{1}{n}Q^TWWQ = f(x)v\kappa_2vh^{-d} + h^{-d} \begin{bmatrix} O(h) + O_P(n^{-\frac{1}{2}}h^{-\frac{d}{2}}) & O_P(h^2 + n^{-\frac{1}{2}}h^{-\frac{d}{2}+1}) \\ O_P(h^2 + n^{-\frac{1}{2}}h^{-\frac{d}{2}+1}) & O_P(h^{\frac{5}{2}} + n^{-\frac{1}{2}}h^{-\frac{d}{2}+2}) \end{bmatrix}.$$

Thus,

$$\begin{aligned} & n^{-1} \sigma_\varepsilon^2 e_1^T (n^{-1} \hat{Q}^T \hat{W} \hat{Q})^{-1} (n^{-1} Q^T W W Q) (n^{-1} \hat{Q}^T \hat{W} \hat{Q})^{-1} e_1^T \\ &= \frac{1}{n h^d} \frac{\sigma_\varepsilon^2}{f(x)} e_1^T \kappa_{22,1}^{-1} \kappa_{22,2} \kappa_{22,1}^{-1} e_1 + O_P \left( n^{-1} h^{-d} (h^{\frac{1}{2}} + m^{-\beta} h^{-4-\epsilon d} + n^{-\frac{1}{2}} h^{-\frac{d}{2}}) \right). \end{aligned} \quad (15)$$

Also, similar to (13), we have

$$\begin{aligned} & 1 n^{-1} \sigma_\varepsilon^2 e_1^T (n^{-1} \hat{Q}^T \hat{W} \hat{Q})^{-1} \mathbb{1}_{(d+1) \times (d+1)} (n^{-1} \hat{Q}^T \hat{W} \hat{Q})^{-1} e_1^T O_P(h^{-d-1-\epsilon d} n^{-\beta}) \\ &= n^{-1} O_P(h^{-2} + m^{-\beta} h^{-5-\epsilon d} + n^{-\frac{1}{2}} h^{-\frac{d}{2}-2}) O_P(h^{-d-1-\epsilon d} m^{-\beta}). \\ &= O_P \left( n^{-1} h^{-d} h^{-1-\epsilon d} m^{-\beta} (h^{-2} + m^{-\beta} h^{-5-\epsilon d} + n^{-\frac{1}{2}} h^{-\frac{d}{2}-2}) \right). \end{aligned} \quad (16)$$

With the condition on  $h$ , combining (10), (11), (15) and (16), we conclude that the conditional variance is

$$\text{var}\{\hat{g}(x) \mid \mathcal{X}\} = \frac{1}{n h^d} \frac{\sigma_\varepsilon^2}{f(x)} e_1^T \kappa_{22,1}^{-1} \kappa_{22,2} \kappa_{22,1}^{-1} e_1 + O_P(n^{-1} h^{-d}).$$

#### S.4. TECHNICAL LEMMAS

Here we collect some technical lemmas that will be used in the proofs of the main theorems. The lemma below is used to establish Proposition 1. Its proof depends on Lemma 2 and 3.

**LEMMA 1.** *Suppose  $h_0 \rightarrow 0$ , and  $m h_0 \rightarrow \infty$ . For any  $p \geq 1$ , assume  $E|\zeta|^p < \infty$ . Under the assumptions (A1)–(A3), for the estimate  $\hat{X}$  in (3) with a proper choice of  $\delta$ ,*

$$\{E(\|\hat{X} - X\|_{\mathcal{L}^2}^p \mid X)\}^{1/p} = O \left( m^{-1/2} h_0^{-1/2} \right) \left\{ \sup_t |X(t)| + L_X \right\} + O(h_0^\nu) L_X, \quad (17)$$

where  $O(\cdot)$  does not depend on  $X$ .

*Proof.* In order to reduce notations, let  $h = h_0$ . Denoting  $\Delta = \delta \mathbb{1}_{|S_0 S_2 - S_1^2| < \delta}$  with  $\delta = m^{-2}$ , according to (3), we have

$$\hat{X}(t) - X(t) = \frac{S_2(R_0 - S_0 X)}{S_0 S_2 - S_1^2 + \Delta} - \frac{S_1(R_1 - S_1 X)}{S_0 S_2 - S_1^2 + \Delta} - \frac{\Delta X}{S_0 S_2 - S_1^2 + \Delta} \equiv I_1 + I_2 + I_3.$$

Therefore,

$$\|\hat{X} - X\|^p \leq c_p (\|I_1\|^p + \|I_2\|^p + \|I_3\|^p) \quad (18)$$

for some constant  $c_p$  depending on  $p$  only.

For  $I_1$ , we have

$$\begin{aligned} \|I_1\|^p &= \left[ \int_D \left\{ \frac{S_2(R_0 - S_0X)}{S_0S_2 - S_1^2 + \Delta} \right\}^2 dt \right]^{p/2} \leq \left[ \int_D \{S_2(R_0 - S_0X)\}^4 dt \int_D \left( \frac{1}{S_0S_2 - S_1^2 + \Delta} \right)^4 dt \right]^{p/4} \\ &\leq \left\{ \int_D S_2^8 dt \int_D (R_0 - S_0X)^8 dt \right\}^{p/8} \left\{ \int_D \left( \frac{1}{S_0S_2 - S_1^2 + \Delta} \right)^4 dt \right\}^{p/4}. \end{aligned}$$

This also shows that, for  $p \geq 2$ ,

$$\begin{aligned} E(\|I_1\|^p | X) &\leq \left( E \left\{ \left[ \int_D S_2^8 dt \int_D (R_0 - S_0X)^8 dt \right]^{p/4} | X \right\} \right)^{1/2} \left( E \left\{ \left[ \int_D \left( \frac{1}{S_0S_2 - S_1^2 + \Delta} \right)^4 dt \right]^{p/2} | X \right\} \right)^{1/2} \\ &\leq \left( E \left[ \int_D S_2^{4p} dt \right] E \left[ \int_D (R_0 - S_0X)^{4p} dt | X \right] \right)^{1/4} \left( E \left[ \int_D \left\{ \frac{1}{S_0S_2 - S_1^2 + \Delta} \right\}^{2p} dt \right] \right)^{1/2} \\ &= \left\{ \int_D E(S_2^{4p}) dt \right\}^{1/4} \left[ \int_D E\{(R_0 - S_0X)^{4p} | X\} dt \right]^{1/4} \left[ \int_D E \left\{ \left( \frac{1}{S_0S_2 - S_1^2 + \Delta} \right)^{2p} \right\} dt \right]^{1/2}. \end{aligned} \tag{19}$$

Let  $E_{0,X} = E(R_0 - S_0X | X)$  and  $\ell$  be the largest integer strictly less than  $\nu$ . By Taylor expansion,

$$\begin{aligned} E_{0,X} &= E \left[ \frac{1}{mh} \sum_{j=1}^m K \left( \frac{T_j - t}{h} \right) \{X(T_j) + \zeta_j - X(t)\} | X \right] \\ &= \frac{1}{mh} \sum_{j=1}^m E \left[ K \left( \frac{T_j - t}{h} \right) \left\{ \frac{X^{(\ell)}(t + \tau_j(T_j - t)) - X^{(\ell)}(t)}{\ell!} (T_j - t)^\ell \right\} | X \right], \end{aligned}$$

where  $\tau_j \in [0, 1]$ . Hence, with  $u_\nu$  denoting  $\int_{-1}^1 K(s) |s|^\nu ds$ , we have

$$\begin{aligned} |E_{0,X}| &\leq \frac{1}{mh} \sum_{j=1}^m E \left\{ K \left( \frac{T_j - t}{h} \right) \left| \frac{X^{(\ell)}(t + \tau_j(T_j - t)) - X^{(\ell)}(t)}{\ell!} (T_j - t)^\ell \right| | X \right\}, \\ &\leq \frac{1}{h\ell!} L_X E \left\{ K \left( \frac{T - t}{h} \right) |T - t|^\nu \right\} \leq \frac{L_X C_{T,2}}{\ell!} h^\nu u_\nu. \end{aligned} \tag{20}$$

Let  $\sigma_{r,X} = E \left( [h^{-1}K \{(T_j - t)/h\} \{X(T_j) + \zeta_j - X(t)\} - E_{0,X}]^r \mid X \right)$ . Then, for  $r \geq 2$ ,

$$\begin{aligned} \sigma_{r,X} &= E \left( \left[ \frac{1}{h} K \left( \frac{T_j - t}{h} \right) \{X(T_j) + \zeta_j - X(t)\} - E_{0,X} \right]^r \mid X \right) \\ &\leq 3^r E \left[ \left\{ \frac{1}{h} K \left( \frac{T_j - t}{h} \right) |X(T_j) - X(t)| \right\}^r \mid X \right] + 3^r E \left[ \left\{ \frac{1}{h} K \left( \frac{T_j - t}{h} \right) |\zeta_j| \right\}^r \mid X \right] + 3^r |E_{0,X}|^r \\ &\leq 2 \cdot 3^r \left\{ \sup_t |X(t)|^r \right\} E \left[ \left\{ \frac{1}{h} K \left( \frac{T_j - t}{h} \right) \right\}^r \right] + 3^r E \left[ \left\{ \frac{1}{h} K \left( \frac{T_j - t}{h} \right) \right\}^r |\zeta_j|^r \right] + 3^r |E_{0,X}|^r \\ &\leq 2 \cdot 3^r \left\{ \sup_t |X(t)|^r \right\} h^{1-r} C_{T,2}^r + 3^r h^{1-r} C_{T,2}^r E|\zeta|^r + 3^r L_X^r C_{T,2}^r h^{r\nu} u_\nu^r. \end{aligned} \quad (21)$$

With (20) and (21), by Lemma 2, conditioning on  $X$ , we have  $E \{(R_0 - S_0 X - E_{0,X})^{4p} \mid X\} \leq c_1(p) C_{T,2}^{4p} \{\sup_t |X(t)|^{4p} + L_X^{4p} u_\nu^{4p}\} m^{-2p} h^{-2p}$ , where  $c_j(p)$  denote a constant depending on  $p$  only for any  $j$ . This implies that

$$\begin{aligned} E \{(R_0 - S_0 X)^{4p} \mid X\} &\leq 2^{4p} E \{(R_0 - S_0 X - E_{0,X})^{4p} \mid X\} + 2^{4p} E \{(E_{0,X})^{4p} \mid X\} \\ &\leq 2^{4p} c_1(p) C_{T,2}^{4p} \{\sup_t |X(t)|^{4p} + L_X^{4p} u_\nu^{4p}\} m^{-2p} h^{-2p} + (2C_{T,2} u_\nu)^{4p} h^{4p\nu} L_X^{4p}. \end{aligned} \quad (22)$$

By a similar argument, we can show that  $E(S_2 - ES_2)^{4p} \leq c_2(p) m^{-2p} h^{-2p}$ . Also, it is easy to check that  $C_{T,1} u_2 \leq ES_2 \leq C_{T,2} u_2$  with  $u_q$  denoting  $\int_{-1}^1 K(u) |u|^q du$  and hence

$$E(S_2^{4p}) \leq 2^{4p} E(S_2 - ES_2)^{4p} + 2^{4p} |ES_2|^{4p} \leq C_{T,2}^{4p} u_2^{4p} + c_2(p) m^{-2p} h^{-2p} = O(1). \quad (23)$$

The same argument leads to  $E\{S_0 S_2 - S_1^2 - E(S_0 S_2 - S_1^2)\}^{2p} \leq c_3(p) m^{-p} h^{-p}$ . Note that  $\inf_t E(S_0 S_2 - S_1^2) > 0$  so that  $\{E(S_0 S_2 - S_1^2)\}^{-1} = O(1)$ . By Lemma 3, this also implies that

$$\int_D E \left\{ \left( \frac{1}{S_0 S_2 - S_1^2 + \Delta} \right)^{2p} \right\} dt = O(1) + O(m^{-p} h^{-p} + m^{-4p}) = O(1). \quad (24)$$

Putting (19), (22), (23) and (24) together, we conclude that

$$E(\|I_1\|^p \mid X) = c_4(p) \left[ \left\{ \sup_t |X(t)|^p + L_X^p \right\} m^{-p/2} h^{-p/2} + h^{p\nu} L_X^p \right]. \quad (25)$$

The same rate for  $I_2$  can be obtained in a similar fashion. For  $I_3$ , we have

$$\|I_3\|^p = \left\{ \int_D \left( \frac{\Delta X}{S_0 S_2 - S_1^2 + \Delta} \right)^2 dt \right\}^{p/2} \leq \Delta^p \sup_t |X(t)|^p \left\{ \int_D \left( \frac{1}{S_0 S_2 - S_1^2 + \Delta} \right)^2 dt \right\}^{p/2}.$$

Therefore, with (24),

$$E(\|I_3\|^p \mid X) \leq \Delta^p \sup_t |X(t)|^p \int_D E \left\{ \left( \frac{1}{S_0 S_2 - S_1^2 + \Delta} \right)^{2p} \right\} dt \leq O(m^{-2p}) \sup_t |X(t)|^p. \quad (26)$$

Now, by (25) and (26), observing that  $m^{-2p}$  is asymptotically dominated by  $m^{-p/2}h^{-p/2}$ , with (18) we conclude that

$$E(\|\hat{X} - X\|^p \mid X) = O(m^{-p/2}h^{-p/2}) \{ \sup_t |X(t)|^p + L_X^p \} + O(h^{p\nu}) L_X^p. \quad (27)$$

Then the statement (17) follows from  $p \geq 1$ .  $\square$

**LEMMA 2.** *Suppose  $V_1, V_2, \dots, V_N$  are i.i.d. random variables with mean zero. Assume  $EV^r \leq a^r h^{1-r} + b^r h^{\kappa r}$  for some constants  $a, b, \kappa \geq 0$  and  $r \geq 2$ . Suppose  $h \geq N^{-1}$ . Then, for any integer  $p \geq 1$ , we have*

$$E \left\{ \left( \frac{1}{N} \sum_{j=1}^N V_j \right)^{2p} \right\} \leq (2p)^{p-1} (2p)! p 2^p (a^{2p} + b^{2p}) N^{-p} h^{-p}.$$

*Proof.* By the multinomial theorem,

$$\left( \sum_{j=1}^N V_j \right)^{2p} = \sum_{k_1 + \dots + k_N = 2p} \binom{2p}{k_1, k_2, \dots, k_N} \prod_{j=1}^N V_j^{k_j}.$$

For each  $\prod_{j=1}^N V_j^{k_j}$ , if  $k_j = 1$  for some  $j$ , then it is zero. Let  $S(k_1, \dots, k_N)$  denote the number of non-zero  $k_j$ . Then, there are in total  $\binom{N}{1}$  items such that  $S(k_1, \dots, k_N) = 1$ , at most  $\binom{N}{2}(2p)$  items such that  $S(k_1, \dots, k_N) = 2$ , etc. In fact, there are at most  $\binom{N}{q}(2p)^{q-1}$  items such that  $S(k_1, \dots, k_N) = q$ , for every  $q \leq p$ . Note that by the assumption,  $EV^r \leq a^r h^{1-r} + b^r h^{\nu \kappa r} \leq (a^r + b^r) h^{1-r}$  for  $r \geq 2$ . Therefore,

$$\begin{aligned} E \left( \sum_{j=1}^N V_j \right)^{2p} &\leq \sum_{q=1}^p \binom{N}{q} (2p)^{q-1} \binom{2p}{k_1, k_2, \dots, k_N} 2^q (a^{2p} h^{q-2p} + b^{2p} h^{q-2p}) \\ &\leq (2p)^{p-1} (2p)! 2^p (a^{2p} + b^{2p}) \sum_{q=1}^p \frac{N!}{(N-q)! q!} h^{q-2p} \\ &\leq (2p)^{p-1} (2p)! p 2^p (a^{2p} + b^{2p}) \sum_{q=1}^p N^p h^{q-2p} \\ &\leq (2p)^{p-1} (2p)! p 2^p (a^{2p} + b^{2p}) N^p h^{-p}. \end{aligned}$$

Multiplying both sides by  $N^{-2p}$  yields the conclusion of the lemma.  $\square$

LEMMA 3. Suppose  $D \subset \mathbb{R}^p$  is compact set and  $S_N(t)$  for  $t \in D$  is a sequence of random processes defined on  $D$ . For  $b_N > 0$ , define  $\delta_N(t) = 2b_N 1_{\{|S(t)| \leq b_N\}}$ . Suppose for some constant  $c_0$ ,  $0 < c_0 \leq ES_N(t) < \infty$  for all  $t$  and sufficiently large  $N$ . Also, assume  $\lim_{N \rightarrow \infty} \inf_{t \in D} ES_N(t) > 0$ . For a sequence of  $a_N \rightarrow 0$  and any  $p > 0$ , if  $b_N^{-r} a_N^p \leq 1$  for some  $p$  and  $r > 0$ , and

(1) if  $E(\sup_{t \in D} |S_N(t) - ES(t)|^p) = O(a_N^p)$ , then we have

$$E \sup_{t \in D} \left| \frac{1}{S_N(t) + \delta_N(t)} - \frac{1}{ES_N(t)} \right|^r = O(a_N^r + b_N^r);$$

(2) if  $\int_D E(|S_N(t) - ES_N(t)|^p) dt = O(a_N^p)$ , then we have

$$E \int \left| \frac{ES_N(t)}{S_N(t) + \delta_N(t)} - 1 \right|^r dt = O(a_N^r + b_N^r).$$

*Proof.* From now on, we shall suppress  $N$  when there is no confusion raised. Let  $\tilde{S}(t) = S(t) + \delta(t)$  and  $v(t) = E\tilde{S}(t) - E\delta(t) = ES(t)$ . For a fixed  $t$  that is suppressed below,

$$\begin{aligned} \left| \frac{v}{\tilde{S}} - 1 \right|^r &\leq \left( \frac{|\tilde{S} - v|^r}{|\tilde{S}|^r} \right) 1_{\{|\tilde{S} - v| \leq v/2\}} + \left( \frac{|\tilde{S} - v|^r}{|\tilde{S}|^r} \right) 1_{\{|\tilde{S} - v| > v/2\}} \\ &\leq (v/2)^{-r} |\tilde{S} - v|^r + b_N^{-r} |\tilde{S} - v|^r 1_{\{|\tilde{S} - v| > v/2\}} \\ &\leq c_r (v/2)^{-r} |S - ES|^r + c_r (v/2)^{-r} \delta^r + b_N^{-r} |\tilde{S} - v|^r 1_{\{|\tilde{S} - v| > v/2\}} \\ &\equiv I_1 + I_2 + I_3, \end{aligned}$$

where  $c_r > 0$  is a constant independent of  $t$ .

(1) By assumption,  $E \sup_t I_1 = O(a_N^r)$ . Since  $|\delta| \leq 2b_N$ , we have  $E \sup_t I_2 = O(b_N^r)$ .

Also,

$$E \sup_t I_3 \leq c_p b_N^{-r} \sup_t (v/2)^{-p} E(\sup_t |S - v|^{p+r} + \sup_t |\delta|^{p+r}) = O(b_N^{-r} a_N^{p+r}) = O(a_N^r + b_N^r)$$

for a sufficiently large  $p$ , and a constant  $c_p > 0$ .

(2) By the assumption,  $\int I_1 dt = O(a_N^r)$ . Since  $|\delta| \leq 2b_N$ , we have  $\int I_2 dt = O(b_N^r)$ . Also,

$$\int I_3 dt \leq b_N^{-r} \int \{v(t)/2\}^{-p} E|\tilde{S}(t) - v(t)|^{p+r} dt = O(b_N^{-r} a_N^{p+r}) = O(a_N^r + b_N^r).$$

Therefore, the conclusion of part (2) follows.  $\square$



In order to state the following lemmas, we shall first establish some notations and convention. Let  $u_{q,k} = \int_{\mathbb{B}_1^{\mathbb{R}}(0)} K^q(\|u\|_{\mathbb{R}^d}) \|u\|_{\mathbb{R}^d}^k du$ . We also identify  $T_x \mathcal{M}$  with  $\mathbb{R}^d$ . Let  $\mathbb{D}_{x,h}$  denote the set  $\{\theta \in \mathbb{R}^d : \exp_x(\theta) \in \mathbb{B}_h^{\mathcal{M}}(x)\}$ , where  $\exp_x$  denotes the exponential map at  $x$ . Let  $\kappa_{11,q} = \int_{h^{-1}\mathbb{D}_x} K^q(\|u\|_{\mathbb{R}^d}) du$ ,  $\kappa_{12,q,j} = \int_{h^{-1}\mathbb{D}_{x,h}} K^q(\|u\|_{\mathbb{R}^d}) \theta_j du$  and  $\kappa_{22,q,j,k} = \int_{h^{-1}\mathbb{D}_x} K^q(h^{-1}\|\theta\|_{\mathbb{R}^d}) \theta_j \theta_k d\theta$ , where  $\theta_j$  denotes the  $j$ th component of  $\theta$ . Let  $\pi_{d-1}$  denote the volume of the unit sphere  $S^{d-1}$ . The following three lemmas are based on Lemma A.2.5 of Cheng & Wu (2013) and hence their proofs are omitted.

LEMMA 4. *Suppose  $K$  is a kernel function compactly supported in  $[-1, 1]$  and continuously differentiable in  $[0, 1]$ . Let  $h \geq h_{pca}$ .*

1. *If  $x \in \mathcal{M} \setminus \mathcal{M}_h$ , then*

$$n^{-1} \sum_{i=1}^n h^{-d} K^q \left( \frac{\|X_i - x\|_{\mathcal{L}^2}}{h} \right) = u_{q,0} f(x) + O(h^2) + O_P(n^{-\frac{1}{2}} h^{-\frac{d}{2}})$$

2. *If  $x \in \mathcal{M}_h$ , then*

$$n^{-1} \sum_{i=1}^n h^{-d} K^q \left( \frac{\|X_i - x\|_{\mathcal{L}^2}}{h} \right) = f(x) \kappa_{11,q} + O(h) + O_P(n^{-\frac{1}{2}} h^{-\frac{d}{2}}).$$

LEMMA 5. *Suppose  $K$  is a kernel function compactly supported in  $[-1, 1]$  and continuously differentiable in  $[0, 1]$ . Let  $h \geq h_{pca}$  and  $\hat{\varphi}_k$  be the estimate in Theorem 1. Then,*

(1) *if  $x \in \mathcal{M} \setminus \mathcal{M}_h$ ,*

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n h^{-d} K^q \left( \frac{\|X_i - x\|_{\mathcal{L}^2}}{h} \right) \langle X_i - x, \hat{\varphi}_k \rangle \\ &= h^2 u_{q,1} d^{-1} \nabla_{\phi_k} f(x) + O_P(h^3 + n^{-\frac{1}{2}} h^{-\frac{d}{2}+1} + h^2 h_{pca}^{3/2} + h^3 h_{pca}). \end{aligned}$$

(2) *if  $x \in \mathcal{M}_h$ ,*

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n h^{-d} K^q \left( \frac{\|X_i - x\|_{\mathcal{L}^2}}{h} \right) \langle X_i - x, \hat{\varphi}_k \rangle \\ &= h \kappa_{12,q,k}(x) + O_P(h^2 + n^{-\frac{1}{2}} h^{-\frac{d}{2}+1} + h h_{pca}^{3/2} + h^2 h_{pca}). \end{aligned}$$

LEMMA 6. *Suppose  $K$  is a kernel function compactly supported in  $[-1, 1]$  and continuously differentiable in  $[0, 1]$ . Let  $h \geq h_{pca}$ .*

(1) If  $x \in \mathcal{M} \setminus \mathcal{M}_h$ , then

$$\begin{aligned} & n^{-1} \sum_{i=1}^n h^{-d} K^q \left( \frac{\|X_i - x\|_{\mathcal{L}^2}}{h} \right) \langle X_i - x, \hat{\varphi}_j \rangle \langle X_i - x, \hat{\varphi}_k \rangle \\ &= \begin{cases} h^2 u_{q,2} d^{-1} f(x) + O_P(h^{\frac{7}{2}} + n^{-\frac{1}{2}} h^{-\frac{d}{2}+2}) & \text{if } 1 \leq j = k \leq d \\ O_P(h^{\frac{7}{2}} + n^{-\frac{1}{2}} h^{-\frac{d}{2}+2}) & \text{otherwise.} \end{cases} \end{aligned}$$

(2) If  $x \in \mathcal{M}_h$ , then

$$\begin{aligned} & n^{-1} \sum_{i=1}^n h^{-d} K^q \left( \frac{\|X_i - x\|_{\mathcal{L}^2}}{h} \right) \langle X_i - x, \hat{\varphi}_j \rangle \langle X_i - x, \hat{\varphi}_k \rangle \\ &= h^2 f(x) \kappa_{22,q,j,k}(x) + O_P(h^3 + n^{-\frac{1}{2}} h^{-\frac{d}{2}+2} + h^2 h_{pca}^{3/2}). \end{aligned}$$

In order to prove Theorem 1, we establish the following auxiliary lemmas.

LEMMA 7. Let  $\tilde{G} = \hat{G} + \Delta$  and  $\check{G} = G + \Delta$  with  $\Delta = 1/\log m$ . Then  $E|\log \tilde{G} - \log \check{G}| = o(1)$ . This result also holds for  $\hat{C}_{\tilde{x}}$ , if  $\tilde{x}$  is independent of  $\hat{X}_1, \dots, \hat{X}_n$ , and that  $\{E\|\tilde{x} - x\|^p\}^{1/p} = O(m^{-\beta})$  for all  $p \geq 1$ .

*Proof.* By Jensen's inequality and the concavity of  $\log(\cdot)$ ,

$$\begin{aligned} E(\log \check{G} - \log \tilde{G}) &\leq \log E \frac{\check{G}}{\tilde{G}} = \log E \frac{\|X - x\| + \Delta}{\|\hat{X} - x\| + \Delta} \leq \log E \frac{\|\hat{X} - x\| + \|\hat{X} - X\| + \Delta}{\|\hat{X} - x\| + \Delta} \\ &\leq \log(1 + \Delta^{-1} E\|\hat{X} - X\|) \equiv a_n \end{aligned}$$

with  $a_n \geq 0$  and  $a_n \rightarrow 0$ . For the other direction, we first observe that

$$\begin{aligned} E \left( \frac{\tilde{G}}{\check{G}} \right)^{1/4} &= E \left\{ \frac{1}{\check{G}^{1/4}} E(\tilde{G}^{1/4} | X) \right\} \leq E \left[ \frac{1}{\check{G}^{1/4}} E\{\|\hat{X} - X\|^{1/4} + (\|X - x\| + \Delta)^{1/4} | X\} \right] \\ &\leq E \left[ \frac{C_1^{1/4} m^{-\beta/4} \{\eta(X)\}^{1/4}}{\check{G}^{1/4}} + 1 \right] \leq 1 + \Delta^{-1/4} E \left[ C_1^{1/4} m^{-\beta/4} \{\eta(X)\}^{1/4} \right] \\ &= 1 + O(m^{-\beta/4} (\log m)^{1/4}) \end{aligned}$$

where  $C_1 > 0$  is some constant. This implies that

$$\frac{1}{4} E(\log \tilde{G} - \log \check{G}) = E \log \left( \frac{\tilde{G}}{\check{G}} \right)^{1/4} \leq \log E \left( \frac{\tilde{G}}{\check{G}} \right)^{1/4} \equiv \frac{b_n}{4}$$

with  $b_n \geq 0$  and  $b_n \rightarrow 0$ , or equivalently,

$$E(\log \check{G} - \log \tilde{G}) \geq -b_n.$$

Therefore  $E|\log \tilde{G} - \log \tilde{G}| \leq a_n + b_n = o(1)$ . Following almost the same lines, we can deduce the same result for  $\tilde{G}(\tilde{x})$ , i.e., the quantity  $\tilde{G}(x)$  when  $x$  is replaced with  $\tilde{x}$ .  $\square$

LEMMA 8. *Let  $\tilde{x}$  be an estimate of  $x$  such that  $\tilde{x}$  is independent of  $X$  and  $\hat{X}$ , and that  $\{E\|\tilde{x} - x\|^p\}^{1/p} = O(m^{-\beta})$  for all  $p \geq 1$ . Suppose  $0 < a < \beta$ ,  $h \gtrsim m^{-a}$  and  $h_+ = h + m^{-(\beta+a)/2}$ . Let  $\tilde{Z} = 1_{\{\hat{X} \in \mathbb{B}_h^{\mathcal{L}^2}(\tilde{x})\}}$  and  $V = 1_{\{X \in \mathbb{B}_{h_+}^{\mathcal{L}^2}(x)\}}$ . If  $F$  is a positive functional of  $X$  and  $\hat{X}$  such that  $E\{F(X, \hat{X})V\} = O(h^b)$  for some  $b \geq 0$ , and  $E\{F(X, \hat{X})\}^q < \infty$  for some  $q > 1$ , then we have  $E\{F(X, \hat{X})|\tilde{Z} - V|\} = O(h^b)$  and  $E\{F(X, \hat{X})\tilde{Z}\} = O(h^b)$ . Also,  $E\{F(X, \hat{X})|Z - \tilde{Z}\} = o(h^b)$  with  $Z$  denoting  $1_{\{\hat{X} \in \mathbb{B}_h^{\mathcal{L}^2}(x)\}}$ .*

*Proof.* Let  $\kappa = m^{-(\beta+a)/2}$  and  $\tilde{V} = 1_{\{X \in \mathbb{B}_{h_+}^{\mathcal{L}^2}(\tilde{x})\}}$ . Choose  $r > 1$  such that  $r^{-1} + q^{-1} = 1$ . To reduce notational burden, we simply use  $F$  to denote  $F(X, \hat{X})$ .

We shall first establish that  $E(F|V - \tilde{V}|) = O(h^b)$ . To this end, we observe that

$$E(F|V - \tilde{V}|) \leq E(FV1_{\tilde{V}=0}) + E(F\tilde{V}1_{V=0}).$$

For the first term, for any fixed  $s \geq 2rab/(\beta - a)$ , we have

$$\begin{aligned} E(FV1_{\tilde{V}=0}) &= E\{F1_{\|X-x\| \leq h, \|X-\tilde{x}\| \geq h}\} \\ &\leq E\{F1_{\|\tilde{x}-x\| \geq \kappa, \|X-x\| \leq h} + F1_{h-\kappa \leq \|X-x\| \leq h}\} \\ &\leq E\{F1_{\|\tilde{x}-x\| \geq \kappa}\} + E\{FV\} \\ &\leq \{EF^q\}^{1/q} \text{pr}(\|\tilde{x} - x\| \geq \kappa)^{1/r} + O(h^b) \\ &\leq \{EF^q\}^{1/q} \left(m^{s(\beta+a)/2} E\|\tilde{x} - x\|^s\right)^{1/r} + O(h^b) \\ &= O(m^{s(\beta+a)/(2r)-s\beta/r} + h^b) = O(h^b). \end{aligned}$$

Similar result can be derived for the second term. Thus, we prove that  $E(F|V - \tilde{V}|) = O(h^b)$ .

Define  $h_- = h - \kappa$ ,  $\tilde{U} = 1_{\{X \in \mathbb{B}_{h_-}^{\mathcal{C}^2}(\tilde{x})\}}$ . Note that  $\tilde{U} \leq \tilde{V}$ . Then, by Hölder inequality, we have

$$\begin{aligned}
E(F|\tilde{Z} - \tilde{V}|) &= E(F1_{\tilde{Z}=1}1_{\tilde{V}=0}) + E(F1_{\tilde{Z}=0}1_{\tilde{U}=1}) + E(F1_{\tilde{Z}=0}1_{\tilde{V}=1}1_{\tilde{U}=0}) \\
&\leq E(F1_{\tilde{Z}=1}1_{\tilde{V}=0}) + E(F1_{\tilde{Z}=0}1_{\tilde{U}=1}) + E(F\tilde{V}1_{\tilde{U}=0}) \\
&\leq (EF^q)^{1/q} \left\{ (E1_{\tilde{Z}=1}1_{\tilde{V}=0})^{1/r} + (E1_{\tilde{Z}=0}1_{\tilde{U}=1})^{1/r} \right\} + O(h^b) \\
&\leq 2(EF^q)^{1/q} \{ \text{pr}(\|X - \hat{X}\| \geq m^{-(\beta+a)/2}) \}^{1/r} + O(h^b) \\
&\leq 2(EF^q)^{1/q} \left( m^{s(\beta+a)/2} E\|X - \hat{X}\|^s \right)^{1/r} + O(h^b) \\
&= O\left(m^{s(\beta+a)/(2r)-s\beta/r}\right) + O(h^b) = O(h^b).
\end{aligned}$$

Then  $E(F|\tilde{Z} - V|) \leq E(F|\tilde{Z} - \tilde{V}|) + E(F|\tilde{V} - V|) = O(h^b)$ . Since  $|E(F\tilde{Z}) - E(FV)| \leq E(F|\tilde{Z} - V|)$ , the result  $E(F\tilde{Z}) = O(h^b)$  follows.  $\square$

LEMMA 9. Suppose  $\{\psi_k\}_{k=1}^\infty$  is an orthonormal basis of  $\mathcal{H}$  and  $x \in \mathcal{M}$  is fixed. Assume that  $\psi_1, \dots, \psi_d$  span the tangent space  $T_x\mathcal{M}$ . Let  $\pi_{d-1}$  be the volume of the  $d-1$  dimensional unit sphere  $S^{d-1}$  and  $\hat{\mathcal{C}}_x$  the sample covariance operator based on  $\hat{\mathcal{N}}_{\mathcal{L}^2}(h, x)$  for some  $h \gtrsim m^{-a}$  with  $0 < a < \beta$ . Then,

$$\begin{aligned}
\sup_{j \leq d} \sup_{k \geq d+1} \left| \langle \hat{\mathcal{C}}_x \psi_j, \psi_k \rangle \right| &= O_P \left( h^{d+4} + n^{-1/2} h^{d/2+3} + m^{-\beta} h^{d+1} \right), \\
\sup_{j, k \geq d+1} \left| \langle \hat{\mathcal{C}}_x \psi_j, \psi_k \rangle \right| &= O_P \left( h^{d+4} + n^{-1/2} h^{d/2+4} + m^{-\beta} h^{d+1} \right), \\
\sup_{1 \leq j \neq k \leq d} \left| \langle \hat{\mathcal{C}}_x \psi_j, \psi_k \rangle \right| &= O_P \left( h^{d+3} + n^{-1/2} h^{d/2+3} + m^{-\beta} h^{d+1} \right),
\end{aligned}$$

$$\text{for } 1 \leq k \leq d: \quad \langle \hat{\mathcal{C}}_x \psi_k, \psi_k \rangle = \pi_{d-1} f(x) d^{-1} h^{d+2} + O_P \left( n^{-1/2} h^{d/2+2} + m^{-\beta} h^{d+1} \right).$$

The above results hold also for  $\hat{\mathcal{C}}_{\tilde{x}}$ , if  $\tilde{x}$  is independent of  $\hat{X}_1, \dots, \hat{X}_n$ , and that  $\{E\|\tilde{x} - x\|^p\}^{1/p} = O(m^{-\beta})$  for all  $p \geq 1$ .

*Proof.* Denote  $Z_i = 1_{\{\hat{X}_i \in \mathbb{B}_h^{\mathcal{L}^2}(x)\}}$ . Then  $\hat{\mathcal{C}}_x$  can be written as  $\hat{\mathcal{C}}_x = n^{-1} \sum_{i=1}^n (\hat{X}_i - \hat{\mu}_x) \otimes (\hat{X}_i - \hat{\mu}_x) Z_i$ , where  $\hat{\mu}_x = \sum_{i=1}^n \hat{X}_i Z_i$ . For any  $y, z$  such that  $\|y\|_{\mathcal{L}^2} = \|z\|_{\mathcal{L}^2} = 1$ , we have

$$\begin{aligned}
\langle \hat{\mathcal{C}}_x y, z \rangle &= \langle n^{-1} \sum_{i=1}^n Z_i (\hat{X}_i - \hat{\mu}_x) \otimes (\hat{X}_i - \hat{\mu}_x) y, z \rangle = n^{-1} \sum_{i=1}^n \langle \hat{X}_i - \hat{\mu}_x, y \rangle \langle \hat{X}_i - \hat{\mu}_x, z \rangle Z_i \\
&= n^{-1} \sum_{i=1}^n \langle X_i - \mu_x, y \rangle \langle X_i - \mu_x, z \rangle Z_i + n^{-1} \sum_{i=1}^n \langle (\hat{X}_i - X_i) - (\hat{\mu}_x - \mu_x), y \rangle \langle \hat{X}_i - \hat{\mu}_x, z \rangle Z_i \\
&\quad + n^{-1} \sum_{i=1}^n \langle \hat{X}_i - \hat{\mu}_x, y \rangle \langle (\hat{X}_i - X_i) - (\hat{\mu}_x - \mu_x), z \rangle Z_i \\
&\quad + n^{-1} \sum_{i=1}^n \langle (\hat{X}_i - X_i) - (\hat{\mu}_x - \mu_x), y \rangle \langle (\hat{X}_i - X_i) - (\hat{\mu}_x - \mu_x), z \rangle Z_i \\
&\equiv I_1 + I_2 + I_3 + I_4,
\end{aligned}$$

where  $\mu_x = \sum_{i=1}^n X_i Z_i$ . Before we proceed to analyze  $I_1, I_2, I_3$  and  $I_4$ , we prepare some calculations.

First, it can be checked that

$$\|\hat{\mu}_x - \mu_x\| = \left\| \frac{1}{n} \sum_{i=1}^n \{(X_i - \mu_x) + (\hat{X}_i - X_i)\} Z_i \right\| \leq \frac{1}{n} \sum_{i=1}^n \|\hat{X}_i - X_i\| Z_i.$$

This implies that

$$\sum_{i=1}^n \|\hat{\mu}_x - \mu_x\| Z_i \leq \sum_{i=1}^n \|\hat{\mu}_x - \mu_x\| \leq \sum_{i=1}^n \|\hat{X}_i - X_i\| Z_i \quad (28)$$

and

$$\sum_{i=1}^n \|\hat{\mu}_x - \mu_x\|^2 Z_i \leq \sum_{i=1}^n \|\hat{\mu}_x - \mu_x\|^2 \leq 2 \sum_{i=1}^n \|\hat{X}_i - X_i\|^2 Z_i. \quad (29)$$

Now we analyze  $I_2$ . It is seen that

$$\begin{aligned}
E \sup_{y,z} |I_2| &\leq E \sup_{y,z} \left| n^{-1} \sum_{i=1}^n \langle (\hat{X}_i - X_i) - (\hat{\mu}_x - \mu_x), y \rangle \langle \hat{X}_i - \hat{\mu}_x, z \rangle Z_i \right| \\
&\leq E \sup_{y,z} n^{-1} \sum_{i=1}^n \|(\hat{X}_i - X_i) - (\hat{\mu}_x - \mu_x)\| \|\hat{X}_i - \hat{\mu}_x\| Z_i \\
&\leq E \sup_{y,z} n^{-1} \sum_{i=1}^n \|(\hat{X}_i - X_i) - (\hat{\mu}_x - \mu_x)\| h Z_i \\
&\leq n^{-1} h E \sum_{i=1}^n \left( \|\hat{X}_i - X_i\| + \|\hat{\mu}_x - \mu_x\| \right) Z_i \\
&\leq 2hE \left( \|\hat{X}_i - X_i\| Z_i \right), \quad (30)
\end{aligned}$$

where the second inequality follows from Cauchy-Schwarz inequality and  $\|y\|_{\mathcal{L}^2} = \|z\|_{\mathcal{L}^2} = 1$ , the third one is from the fact that  $\hat{\mu}_x, \hat{X}_i \in \mathbb{B}_h^{\mathcal{L}^2}(x)$  for  $Z_i = 1$ , the fourth one follows from triangle inequality, and the fifth is based on (28). Now, let  $h_1 = h + m^{-(\beta+a)/2}$  and  $V_i = 1_{\{X_i \in \mathbb{B}_{h_1}^{\mathcal{L}^2}(x)\}}$ . Based on the assumption (B3),  $E(\|\hat{X}_i - X_i\|V_i) = O(m^{-\beta}h_1^{d+1})$ . Then, by Lemma 8,  $E(\|\hat{X}_i - X_i\|Z_i) = O(m^{-\beta}h_1^{d+1}) = O(m^{-\beta}h^{d+1})$ , since  $h_1 \asymp h$ . With (30), this shows that  $E \sup_{y,z} |I_2| = O(m^{-\beta}h^{d+1})$ . Similarly,  $E \sup_{y,z} |I_3| = O(m^{-\beta}h^{d+1})$ .

For  $I_4$ , we have

$$\begin{aligned}
E \sup_{y,z} |I_4| &= E \sup_{y,z} n^{-1} \left| \sum_{i=1}^n \langle (\hat{X}_i - X_i) - (\hat{\mu}_x - \mu_x), y \rangle \langle (\hat{X}_i - X_i) - (\hat{\mu}_x - \mu_x), z \rangle Z_i \right| \\
&\leq E \sup_{y,z} n^{-1} \sum_{i=1}^n \|(\hat{X}_i - X_i) - (\hat{\mu}_x - \mu_x)\|^2 Z_i \\
&\leq 2En^{-1} \sum_{i=1}^n (\|\hat{X}_i - X_i\|^2 + \|\hat{\mu}_x - \mu_x\|^2) Z_i \\
&\leq 4En^{-1} \sum_{i=1}^n \|\hat{X}_i - X_i\|^2 Z_i = 4E(\|\hat{X}_i - X_i\|^2 Z_i),
\end{aligned}$$

where the first inequality comes from Cauchy-Schwarz inequality and  $\|y\|_{\mathcal{L}^2} = \|z\|_{\mathcal{L}^2} = 1$ , the second one follows from triangle inequality, and the third is based on (29). By Lemma 8, we can show that  $E(\|\hat{X}_i - X_i\|^2 Z_i) = O(m^{-2\beta}h^d)$ .

The order of  $I_1$  is given in Lemma 10. Therefore, the results of the lemma follow. It is easy but somewhat tedious to check that the above derivation also holds for  $\tilde{x}$ .  $\square$

LEMMA 10. For  $0 < a < \beta$ , let  $h \gtrsim m^{-a}$  and  $h_1 = h + n^{-(\beta+a)/2}$ . Denote  $Z_i = 1_{\{\hat{X}_i \in \mathbb{B}_h^{\mathcal{L}^2}(x)\}}$  and  $\Xi_{j,k} = n^{-1} \sum_{i=1}^n \langle X_i - \mu_x, \psi_j \rangle \langle X_i - \mu_x, \psi_k \rangle Z_i$ , where  $\{\psi_k\}$  are defined in Lemma 9. Then, we have

$$E \sup_{j \leq d} \sup_{k \geq d+1} |\Xi_{j,k}| = O(h^{d+3}), \quad (31)$$

$$E \sup_{j,k \geq d+1} |\Xi_{j,k}| = O(h^{d+4}), \quad (32)$$

$$\text{for } 1 \leq j \neq k \leq d: \quad E \Xi_{j,k} = O(h^{d+3}), \quad (33)$$

$$\text{for } 1 \leq k \leq d: \quad E \Xi_{k,k} = O(h^{d+2}). \quad (34)$$

Also,

$$\begin{aligned}
\text{var} \left( \sup_{j \leq d} \sup_{k \geq d+1} |\Xi_{j,k}| \right) &= O(n^{-1}h^{d+6}), \\
\text{var} \left( \sup_{j,k \geq d+1} |\Xi_{j,k}| \right) &= O(n^{-1}h^{d+8}), \\
\text{for } 1 \leq j \neq k \leq d : \quad \text{var}(|\Xi_{j,k}|) &= O(n^{-1}h^{d+6}), \\
\text{for } 1 \leq k \leq d : \quad \text{var}(\Xi_{k,k}) &= O(n^{-1}h^{d+4}).
\end{aligned} \tag{35}$$

*Proof.* Let  $V_i = 1_{\{X_i \in \mathbb{B}_{h_1}^{\mathcal{L}^2}(x)\}}$ . Since  $h_1 \asymp h$ , we shall use them exchangeably when no confusion arises. Let  $\mathcal{P}_1$  be projection into  $\{\psi_1, \dots, \psi_d\}$  and  $\mathcal{P}_2$  be projection into  $\{\psi_{d+1}, \dots\}$ . First, for  $1 \leq j \leq d$ ,

$$\begin{aligned}
E \sup_{k \geq d+1} |\Xi_{j,k}| &\leq E \sup_{k \geq d+1} \left| n^{-1} \sum_{i=1}^n \langle X_i - \mu_x, \psi_j \rangle \langle X_i - \mu_x, \psi_k \rangle Z_i \right| \\
&\leq E \sup_{k \geq d+1} \left| n^{-1} \sum_{i=1}^n \langle X_i, \psi_j \rangle \langle X_i, \psi_k \rangle Z_i \right| \\
&\quad + E \sup_{k \geq d+1} \left| n^{-1} \sum_{i=1}^n \langle \mu_x, \psi_j \rangle \langle \mu_x, \psi_k \rangle Z_i \right| \\
&\equiv I_1 + I_2.
\end{aligned} \tag{36}$$

It is seen that  $I_1$  is the dominant term, which we evaluate below (utilizing the fact that  $\Pi_x(\theta, \theta) \perp T_x \mathcal{M}$ ):

$$\begin{aligned}
I_1 &= E \sup_{k \geq d+1} \left| n^{-1} \sum_{i=1}^n \langle X_i, \psi_j \rangle \langle X_i, \psi_k \rangle Z_i \right| \leq \frac{1}{n} E \sum_{i=1}^n \sup_{k \geq d+1} |\langle X_i, \psi_j \rangle \langle X_i, \psi_k \rangle Z_i| \\
&= E \sup_{k \geq d+1} |\langle X_i, \psi_j \rangle \langle X_i, \psi_k \rangle Z_i| = E \sup_{k \geq d+1} |\langle X_i, \psi_j \rangle \langle \mathcal{P}_2 X_i, \psi_k \rangle Z_i|. \\
&\leq E (|\langle X_i, \psi_j \rangle| \|\mathcal{P}_2 X_i\| |Z_i|).
\end{aligned} \tag{37}$$

Since by Lemma A.2.4 of Cheng & Wu (2013),

$$\begin{aligned}
&E |\langle X_i, \psi_j \rangle| \|\mathcal{P}_2 X_i\| |V_i| \\
&\leq \int_{S^{d-1}} \int_0^{h_1} \langle t\theta, \psi_j \rangle \|t^2 \Pi_x(\theta, \theta)\|_{\mathcal{L}^2} f(\exp(t\theta)) t^{d-1} dt d\theta + O(h^{d+5}) \\
&= O(h^{d+3}),
\end{aligned}$$

we can apply Lemma 8 to conclude  $E(\|\langle X_i, \psi_j \rangle\| \|\mathcal{P}_2 X_i\| Z_i) = O(h^{d+3})$ , and hence with (37), we assert that  $I_1 = O(h^{d+3})$ . This proves (31). The result (32) is obtained in a similar way.

For (33), by the same argument that leads to (36), we can show that for  $1 \leq j \neq k \leq d$ ,  $E\Xi_{j,k}$  is dominated by

$$En^{-1} \sum_{i=1}^n \langle X_i, \psi_j \rangle \langle X_i, \psi_k \rangle Z_i = E \langle X_i, \psi_j \rangle \langle X_i, \psi_k \rangle Z_i.$$

Now, because

$$\begin{aligned} & E \langle X_i, \psi_j \rangle \langle X_i, \psi_k \rangle V_i \\ &= \int_{S^{d-1}} \int_0^{h_1} \langle t\theta, \psi_j \rangle \langle t\theta, \psi_k \rangle [f(x) + t\nabla_\theta f(x)] t^{d-1} dt d\theta + O(h^{d+4}) \\ &= O(h^{d+3}), \end{aligned}$$

where the second equality is based on the fact that the second fundamental form is self-adjoint, by Lemma 8, (33) follows. The result (34) is derived in a similar fashion.

Let  $\chi_{i,k} = n^{-1} \langle X_i, \psi_j \rangle \langle X_i, \psi_k \rangle Z_i$ . Then  $\Xi_{j,k} = \sum_{i=1}^n \chi_{i,k}$ . Then by Theorem 11.1 of Boucheron et al. (2016), we have

$$\text{var} \left( \sup_{k \geq d+1} \Xi_{j,k} \right) = \text{var} \left( \sup_{k \geq d+1} \sum_{i=1}^n \chi_{i,k} \right) \leq \sum_{i=1}^n E \sup_{k \geq d+1} \chi_{i,k}^2 = n E \sup_{k \geq d+1} \chi_{i,k}^2. \quad (38)$$

The term  $E \sup_{k \geq d+1} \chi_{i,k}^2$  can be computed as follows:

$$\begin{aligned} E \sup_{k \geq d+1} \chi_{i,k}^2 &= E \sup_{k \geq d+1} \left[ n^{-1} \langle X_i, \psi_j \rangle \langle \mathcal{P}_2 X_i, \psi_k \rangle Z_i \right]^2 \\ &\leq n^{-2} E \sup_{k \geq d+1} \|X_i\|^2 \|\mathcal{P}_2 X_i\|^2 \|\psi_j\|^2 \|\psi_k\|^2 Z_i = n^{-2} E \|X_i\|^2 \|\mathcal{P}_2 X_i\|^2 Z_i. \end{aligned}$$

Since

$$E \|X_i\|^2 \|\mathcal{P}_2 X_i\|^2 V_i = \int_{S^{d-1}} \int_0^{h_1} \|t\theta\|^2 \|t^2 \Pi_x(\theta, \theta)\|^2 f(\exp(t\theta)) t^{d-1} dt d\theta + O(h^{d+8}) = O(h^{d+6}),$$

we apply Lemma 8 to conclude  $E \|X_i\|^2 \|\mathcal{P}_2 X_i\|^2 Z_i = O(h^{d+6})$ . Therefore,  $E \sup_{k \geq d+1} \chi_{i,k}^2 = O(n^{-2} h^{d+6})$ . With (38), we show that

$$\text{var} \left( \sup_{k \geq d+1} \Xi_{j,k} \right) = O(n^{-1} h^{d+6}).$$

Other results are derived in the same way.  $\square$



## REFERENCES

- BOUCHERON, S., LUGOSI, G. & MASSART, P. (2016). *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford, United Kingdom: Oxford University Press.
- CHENG, M. & WU, H. (2013). Local linear regression on manifolds and its geometric interpretation. *Journal of the American Statistical Association* **108**, 1421–1434.
- FAN, J. (1993). Local linear regression smoothers and their minimax efficiencies. *The Annals of Statistics* **21**, 196–216.
- LEVINA, E. & BICKEL, P. (2004). Maximum likelihood estimation of intrinsic dimension. *Advances in Neural Information* **17**, 777–784.
- SINGER, A. & WU, H.-T. (2012). Vector diffusion maps and the connection laplacian. *Communications on Pure and Applied Mathematics* **65**, 1067–1144.
- YAO, F., MÜLLER, H.-G. & WANG, J.-L. (2005). Functional data analysis for sparse longitudinal data. *Journal of the American Statistical Association* **100**, 577–590.
- ZHANG, X. & WANG, J.-L. (2016). From sparse to dense functional data and beyond. *The Annals of Statistics* **44**, 2281–2321.

