

# Fast multi-output relevance vector regression

Youngmin Ha\*

Adam Smith Business School, University of Glasgow, UK

(First draft: April 14, 2017)

This paper aims to decrease the time complexity of multi-output relevance vector regression from  $O(VM^3)$  to  $O(V^3 + M^3)$ , where  $V$  is the number of output dimensions,  $M$  is the number of basis functions, and  $V < M$ . The experimental results demonstrate that the proposed method is more competitive than the existing method, with regard to computation time. MATLAB codes are available at <http://www.mathworks.com/matlabcentral/fileexchange/49131>.

*Keywords:* Relevance vector regression; Relevance vector machine; Sparse Bayesian learning

## 1. Introduction

When it comes to multi-input nonparametric nonlinear regression or classification, the following three methods can be considered: support vector machine (SVM), relevance vector machine (RVM), and Gaussian process (GP) regression or classification.

SVM, invented by [Cortes and Vapnik \(1995\)](#), is a popular machine learning tool. It uses kernel trick (RVM and GP also use the kernel trick) and makes classification and regression computationally efficient for multidimensional inputs. However, its disadvantage is that a user needs to choose a proper value of the error/margin trade-off parameter ‘ $C$ ’ (the proper value can be found by k-fold cross-validation).

RVM, invented by [Tipping \(2001\)](#),<sup>1</sup> avoids estimating the error/margin trade-off parameter ‘ $C$ ’ of SVM (and in regression the additional insensitivity parameter ‘ $\varepsilon$ ’ ([Vapnik 2000](#)), or ‘ $\nu$ ’ ([Schölkopf et al. 2000](#))), which wastes computation time. Moreover, RVM allows probabilistic predictions (i.e. both mean and variance of a Gaussian distribution) although SVM predicts only mean values (the error bar estimation of SVM is possible with additional computation ([Gao et al. 2002](#), [Chu et al. 2004](#))).

GP regression or classification, invented by [Gibbs \(1997\)](#), also does not need estimating ‘ $C$ ’ (and the additional parameter of regression ‘ $\varepsilon$ ’ or ‘ $\nu$ ’). Furthermore, the predictive variance of GP regression or classification changes over an input vector  $\mathbf{x}_*$ : the predictive variance is smaller at the denser region of training samples, while the predictive variance of RVM is almost constant over  $\mathbf{x}_*$ .

Support vector regression (SVR), relevance vector regression (RVR), and GP regression are for multi-input but single-output regression, and they have been extended as multi-input and multi-output regression to model correlated outputs: multi-output SVR ([Pérez-Cruz et al. 2002](#), [Vazquez and Walter 2003](#), [Tuia et al. 2011](#)), multi-output RVR ([Thayananthan 2005](#), [Thayananthan et al. 2008](#)), and multi-output GP regression ([Boyle and Frean 2004](#), [Bonilla et al. 2007](#), [Alvarez and Lawrence 2009](#)).

The multi-output relevance vector regression (MRVR) algorithm by [Thayananthan \(2005](#), Chap-

\*Email: [y.ha.1@research.gla.ac.uk](mailto:y.ha.1@research.gla.ac.uk)

<sup>1</sup> Single-output relevance vector regression is easily explained in <http://static1.squarespace.com/static/58851af9ebbd1a30e98fb283/t/58902f4a6b8f5ba2ed9d3bfe/1485844299331/RVM+Explained.pdf>.

ter 6), [Thayananthan \*et al.\* \(2008\)](#) uses the Bayes' theorem and the kernel trick to perform regression, but it has the limitation of low computational efficiency. Therefore, a new faster algorithm is proposed in this paper: it uses the matrix normal distribution to model correlated outputs, while the existing algorithm uses the multivariate normal distribution.

The contributions of this paper are:

- in Section 4, to propose a new algorithm with less time complexity than the existing MRVR algorithm by [Thayananthan \(2005, Chapter 6\)](#), [Thayananthan \*et al.\* \(2008\)](#);
- in Section 5, to present Monte Carlo simulation results to compare between the existing and the proposed MRVR algorithm in terms of accuracy and computation time.

The rest of this paper is organised as follows: Section 2 lists related work. Section 3 and Section 4 describe the existing and proposed algorithms of MRVR, respectively. Section 5 shows the experimental results by using MATLAB, and Section 6 gives the conclusion.

## 2. Related work

The computation time of the single-output SVM has decreased: [Guo and Zhang \(2007\)](#) reduced the SVR training time by reducing the number of training examples. [Catanzaro \*et al.\* \(2008\)](#) accelerated the SVM computation by using both faster sequential algorithms and parallel computation on a graphics processing unit (GPU). [Chang and Lin \(2011\)](#) made fast and efficient C++ software of SVM.

The computation time of the single-output RVM has decreased: [Tipping and Faul \(2003\)](#) proposed a new marginal likelihood maximisation algorithm with efficient addition/deletion of candidate basis functions. [Ben-Shimon and Shmilovici \(2006\)](#) partitioned training samples into small chunks to avoid the inverse of a large matrix (the matrix inversion is the most computationally expensive part of RVM). [Yang \*et al.\* \(2010\)](#) accelerated RVM computation by parallelising the matrix inversion operation on a GPU.

The computation time of the single-output GP regression has decreased. [Seeger \*et al.\* \(2003\)](#) reduced GP regression training time by approximating the likelihood of training data. [Shen \*et al.\* \(2006\)](#) reduced both the training and prediction time of GP regression by using an approximation method, based on a tree-type multiresolution data structure. [Srinivasan \*et al.\* \(2010\)](#) accelerated linear algebra operations of GP regression on a GPU, and [Gramacy \*et al.\* \(2014\)](#) made a GPU accelerated version of GP regression approximation.

## 3. Existing method

The following subsections describe the existing method of MRVR ([Thayananthan 2005, Chapter 6](#)), ([Thayananthan \*et al.\* 2008](#)).

### 3.1. Model specification

$V$ -dimensional multi-output regression upon an input  $\mathbf{x} \in \mathbb{R}^{U \times 1}$  ( $U$ -dimensional column vector), a weight  $\mathbf{W} \in \mathbb{R}^{M \times V}$  ( $M$ -by- $V$  matrix), and an output function  $\mathbf{y}(\mathbf{x}; \mathbf{W}) \in \mathbb{R}^{1 \times V}$  ( $V$ -dimensional row vector) (upright bold lower case letters denote vectors, and upright bold capital letters denote matrices) is expressed as

$$\mathbf{y}(\mathbf{x}; \mathbf{W}) = (\mathbf{W}^\top \boldsymbol{\phi}(\mathbf{x}))^\top, \quad (3.1)$$

where  $\mathbf{W}^\top$  is the transpose of the matrix  $\mathbf{W}$  (the objective of this paper is to estimate proper values of  $\mathbf{W}$ ), and  $\boldsymbol{\phi}(\mathbf{x}) = [\phi_1(\mathbf{x}) \ \phi_2(\mathbf{x}) \ \dots \ \phi_M(\mathbf{x})]^\top \in \mathbb{R}^{M \times 1}$  is the transformed input by nonlinear and fixed basis functions. In other words, the output  $\mathbf{y}(\mathbf{x}; \mathbf{W})$  is a linearly weighted sum of the transformed input  $\boldsymbol{\phi}(\mathbf{x})$ .

Given a data set of input-target pairs  $\{\mathbf{x}_i \in \mathbb{R}^{U \times 1}, \mathbf{t}_i \in \mathbb{R}^{1 \times V}\}_{i=1}^N$ , where  $N$  is the number of training samples, it is assumed that the targets  $\mathbf{t}_i$  are samples from the model  $\mathbf{y}(\mathbf{x}_i; \mathbf{W})$  with additive noise:

$$\mathbf{t}_i = \mathbf{y}(\mathbf{x}_i; \mathbf{W}) + \boldsymbol{\epsilon}_i, \quad (3.2)$$

where  $\mathbf{W} \in \mathbb{R}^{(N+1) \times V}$  is the weight,  $\boldsymbol{\epsilon}_i \in \mathbb{R}^{1 \times V}$  are independent samples from a Gaussian noise process with mean zero and a covariance matrix  $\boldsymbol{\Omega} \in \mathbb{R}^{V \times V}$ , and  $\boldsymbol{\Omega}$  is decomposed as the diagonal matrix of the variances  $\mathbf{D} \in \mathbb{R}^{V \times V}$  and the correlation matrix  $\mathbf{R} \in \mathbb{R}^{V \times V}$ :

$$\boldsymbol{\Omega} = \mathbf{D}^{\frac{1}{2}} \mathbf{R} \mathbf{D}^{\frac{1}{2}}, \quad (3.3)$$

where  $\mathbf{D} = \text{diag}(\sigma_1^2, \sigma_2^2, \dots, \sigma_V^2)$ , and  $\sigma_j^2$  is the variance of the  $j$ -th dimension's noise.

Because of the ignorance of  $\mathbf{R}$  by [Thayananathan \(2005\)](#) and the assumption of independent Gaussian noise, the likelihood of the data set can be given by the product of the Gaussian distributions:

$$p(\mathbf{T}|\mathbf{W}, \boldsymbol{\sigma}) = \prod_{j=1}^V (2\pi\sigma_j^2)^{-\frac{N}{2}} \exp\left(-\frac{1}{2\sigma_j^2} \|\boldsymbol{\tau}_j - \boldsymbol{\Phi} \mathbf{w}_j\|^2\right), \quad (3.4)$$

where  $\mathbf{T} = \begin{bmatrix} \mathbf{t}_1 \\ \mathbf{t}_2 \\ \vdots \\ \mathbf{t}_N \end{bmatrix} \in \mathbb{R}^{N \times V}$ ,  $\boldsymbol{\sigma} = [\sigma_1 \ \sigma_2 \ \dots \ \sigma_V] \in \mathbb{R}_{\geq 0}^{1 \times V}$ ,  $\boldsymbol{\tau}_j \in \mathbb{R}^{N \times 1}$  is the  $j$ -th column vector of  $\mathbf{T}$ ,

$\mathbf{w}_j \in \mathbb{R}^{(N+1) \times 1}$  is the  $j$ -th column vector of  $\mathbf{W}$ ,  $\boldsymbol{\Phi} = [\boldsymbol{\phi}(\mathbf{x}_1) \ \boldsymbol{\phi}(\mathbf{x}_2) \ \dots \ \boldsymbol{\phi}(\mathbf{x}_N)]^\top \in \mathbb{R}^{N \times (N+1)}$  is a design matrix,  $\boldsymbol{\phi}(\mathbf{x}) = [1 \ K(\mathbf{x}, \mathbf{x}_1) \ \dots \ K(\mathbf{x}, \mathbf{x}_N)]^\top \in \mathbb{R}^{(N+1) \times 1}$ , and  $K(\mathbf{x}, \mathbf{x}')$  is a kernel function. For clarity, the implicit conditioning on the input  $\mathbf{x}_i, \forall i$  is omitted in Eq. (3.4) and the subsequent expressions.

An assumption to avoid over-fitting in estimation of  $\mathbf{W}$  is

$$p(\mathbf{W}|\boldsymbol{\alpha}) = \prod_{j=1}^V \prod_{i=0}^N \mathcal{N}(0, \alpha_i^{-1}). \quad (3.5)$$

This means the prior distribution of  $\mathbf{w}_j$  is zero-mean Gaussian with inverse variances  $\boldsymbol{\alpha} = [\alpha_0 \ \alpha_1 \ \dots \ \alpha_N]^\top \in \mathbb{R}_{>0}^{(N+1) \times 1}$ , which are  $N+1$  hyperparameters ([Tipping 2001](#)), and  $\mathbf{w}_j$  and  $\mathbf{w}_{j'}$  ( $j \neq j'$ ) have the same distribution as  $\prod_{i=0}^N \mathcal{N}(0, \alpha_i^{-1})$ .

### 3.2. Inference

By both the Bayes' theorem and the property of  $p(\mathbf{T}|\mathbf{W}, \boldsymbol{\alpha}, \boldsymbol{\sigma}) = p(\mathbf{T}|\mathbf{W}, \boldsymbol{\sigma})$ ,<sup>1</sup> the posterior probability distribution function over  $\mathbf{W}$  is decomposed as

$$p(\mathbf{W}|\mathbf{T}, \boldsymbol{\alpha}, \boldsymbol{\sigma}) = \frac{p(\mathbf{T}|\mathbf{W}, \boldsymbol{\sigma}) p(\mathbf{W}|\boldsymbol{\alpha}, \boldsymbol{\sigma})}{p(\mathbf{T}|\boldsymbol{\alpha}, \boldsymbol{\sigma})}, \quad (3.6)$$

<sup>1</sup> In the case that the weight  $\mathbf{W}$  is given, its inverse variances  $\boldsymbol{\alpha}$  are redundant in the calculation of the conditional probability of the target  $\mathbf{T}$ .

and it is given by the product of multivariate Gaussian distributions:

$$p(\mathbf{W}|\mathbf{T}, \boldsymbol{\alpha}, \boldsymbol{\sigma}) = \prod_{j=1}^V (2\pi)^{-\frac{N+1}{2}} |\boldsymbol{\Sigma}_j|^{-\frac{1}{2}} \exp\left(-\frac{1}{2} (\mathbf{w}_j - \boldsymbol{\mu}_j)^\top \boldsymbol{\Sigma}_j^{-1} (\mathbf{w}_j - \boldsymbol{\mu}_j)\right), \quad (3.7)$$

where the  $j$ -th posterior covariance and mean are, respectively:

$$\boldsymbol{\Sigma}_j = \left(\sigma_j^{-2} \boldsymbol{\Phi}^\top \boldsymbol{\Phi} + \mathbf{A}\right)^{-1}, \quad (3.8)$$

$$\boldsymbol{\mu}_j = \sigma_j^{-2} \boldsymbol{\Sigma}_j \boldsymbol{\Phi}^\top \boldsymbol{\tau}_j, \quad (3.9)$$

where  $\mathbf{A} = \text{diag}(\alpha_0, \alpha_1, \dots, \alpha_N) \in \mathbb{R}^{(N+1) \times (N+1)}$ .

In the case of uniform hyperpriors  $\boldsymbol{\alpha}$  and  $\boldsymbol{\sigma}$ , maximising a posteriori  $p(\boldsymbol{\alpha}, \boldsymbol{\sigma}|\mathbf{T}) \propto p(\mathbf{T}|\boldsymbol{\alpha}, \boldsymbol{\sigma}) p(\boldsymbol{\alpha}) p(\boldsymbol{\sigma})$  is equivalent to maximising the marginal likelihood  $p(\mathbf{T}|\boldsymbol{\alpha}, \boldsymbol{\sigma})$ , which is given by

$$p(\mathbf{T}|\boldsymbol{\alpha}, \boldsymbol{\sigma}) = \prod_{j=1}^V (2\pi)^{-\frac{N}{2}} |\sigma_j^2 \mathbf{I} + \boldsymbol{\Phi} \mathbf{A}^{-1} \boldsymbol{\Phi}^\top|^{-\frac{1}{2}} \exp\left(-\frac{1}{2} \boldsymbol{\tau}_j^\top (\sigma_j^2 \mathbf{I} + \boldsymbol{\Phi} \mathbf{A}^{-1} \boldsymbol{\Phi}^\top)^{-1} \boldsymbol{\tau}_j\right). \quad (3.10)$$

### 3.3. Marginal likelihood maximisation

The same method of accelerating the univariate relevance vector machine (Tipping and Faul 2003) is used to accelerate the existing algorithm.

The log of Eq. (3.10) is an objective function:

$$\mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\sigma}) = -\frac{1}{2} \sum_{j=1}^V \left( N \log(2\pi) + \log |\mathbf{C}_j| + \boldsymbol{\tau}_j^\top \mathbf{C}_j^{-1} \boldsymbol{\tau}_j \right), \quad (3.11)$$

where  $\mathbf{C}_j = \sigma_j^2 \mathbf{I} + \boldsymbol{\Phi} \mathbf{A}^{-1} \boldsymbol{\Phi}^\top \in \mathbb{R}^{N \times N}$ , and by considering the dependence of  $\mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\sigma})$  on a single hyperparameter  $\alpha_i, i \in \{0, 1, \dots, N\}$ ,  $\mathbf{C}_j$  is decomposed as the following two parts:

$$\begin{aligned} \mathbf{C}_j &= \sigma_j^2 \mathbf{I} + \sum_{m \neq i} \alpha_m^{-1} \boldsymbol{\phi}_m \boldsymbol{\phi}_m^\top + \alpha_i^{-1} \boldsymbol{\phi}_i \boldsymbol{\phi}_i^\top \\ &= \mathbf{C}_{-i,j} + \alpha_i^{-1} \boldsymbol{\phi}_i \boldsymbol{\phi}_i^\top, \end{aligned} \quad (3.12)$$

where  $\mathbf{C}_{-i,j} \in \mathbb{R}^{N \times N}$  is  $\mathbf{C}_j$  with the contribution of a basis vector  $\boldsymbol{\phi}_i \in \mathbb{R}^{N \times 1}$  removed, and

$$\boldsymbol{\phi}_i = \begin{cases} [1 \ 1 \ \dots \ 1]^\top, & \text{if } i = 0 \\ [K(\mathbf{x}_i, \mathbf{x}_1) \ K(\mathbf{x}_i, \mathbf{x}_2) \ \dots \ K(\mathbf{x}_i, \mathbf{x}_N)]^\top, & \text{otherwise} \end{cases}. \quad (3.13)$$

The determinant and inverse matrix of  $\mathbf{C}_j$  are, respectively:

$$|\mathbf{C}_j| = |\mathbf{C}_{-i,j}| \left(1 + \alpha_i^{-1} \boldsymbol{\phi}_i^\top \mathbf{C}_{-i,j}^{-1} \boldsymbol{\phi}_i\right), \quad (3.14)$$

by Sylvester's determinant theorem, and

$$\mathbf{C}_j^{-1} = \mathbf{C}_{-i,j}^{-1} - \frac{\mathbf{C}_{-i,j}^{-1} \boldsymbol{\Phi}_i \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i,j}^{-1}}{\alpha_i + \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i,j}^{-1} \boldsymbol{\Phi}_i}, \quad (3.15)$$

by Woodbury matrix identity. From these,  $\mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\sigma})$  in Eq. (3.11) can be decomposed into  $\mathcal{L}(\boldsymbol{\alpha}_{-i}, \boldsymbol{\sigma})$ , the marginal likelihood with  $\boldsymbol{\Phi}_i$  excluded, and  $\ell(\alpha_i, \boldsymbol{\sigma})$ , the isolated marginal likelihood of  $\boldsymbol{\Phi}_i$ :

$$\begin{aligned} \mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\sigma}) &= -\frac{1}{2} \sum_{j=1}^V \left( N \log(2\pi) + \log |\mathbf{C}_{-i,j}| + \boldsymbol{\tau}_j^\top \mathbf{C}_{-i,j}^{-1} \boldsymbol{\tau}_j \right) \\ &\quad - \frac{1}{2} \sum_{j=1}^V \left( -\log \alpha_i + \log \left( \alpha_i + \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i,j}^{-1} \boldsymbol{\Phi}_i \right) - \frac{\left( \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i,j}^{-1} \boldsymbol{\tau}_j \right)^2}{\alpha_i + \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i,j}^{-1} \boldsymbol{\Phi}_i} \right) \\ &= \mathcal{L}(\boldsymbol{\alpha}_{-i}, \boldsymbol{\sigma}) + \frac{1}{2} \sum_{j=1}^V \left( \log \alpha_i - \log (\alpha_i + s_{i,j}) + \frac{q_{i,j}^2}{\alpha_i + s_{i,j}} \right) \\ &= \mathcal{L}(\boldsymbol{\alpha}_{-i}, \boldsymbol{\sigma}) + \ell(\alpha_i, \boldsymbol{\sigma}), \end{aligned} \quad (3.16)$$

where  $s_{i,j}$  and  $q_{i,j}$  are defined as, respectively:

$$s_{i,j} \stackrel{\text{def}}{=} \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i,j}^{-1} \boldsymbol{\Phi}_i, \quad (3.17a)$$

$$q_{i,j} \stackrel{\text{def}}{=} \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i,j}^{-1} \boldsymbol{\tau}_j. \quad (3.17b)$$

To avoid the matrix inversion of  $\mathbf{C}_{-i}$  in Eq. (3.17), which requires the time complexity of  $O(N^3)$ ,  $s'_{i,j}$  and  $q'_{i,j}$  are computed as, respectively (by the Woodbury matrix identity):<sup>1</sup>

$$\begin{aligned} s'_{i,j} &= \boldsymbol{\Phi}_i^\top \mathbf{C}_j^{-1} \boldsymbol{\Phi}_i \\ &= \sigma_j^{-2} \boldsymbol{\Phi}_i^\top \boldsymbol{\Phi}_i - \sigma_j^{-4} \boldsymbol{\Phi}_i^\top \boldsymbol{\Phi} \boldsymbol{\Sigma}_j \boldsymbol{\Phi}^\top \boldsymbol{\Phi}_i, \end{aligned} \quad (3.18a)$$

$$\begin{aligned} q'_{i,j} &= \boldsymbol{\Phi}_i^\top \mathbf{C}_j^{-1} \boldsymbol{\tau}_j \\ &= \sigma_j^{-2} \boldsymbol{\Phi}_i^\top \boldsymbol{\tau}_j - \sigma_j^{-4} \boldsymbol{\Phi}_i^\top \boldsymbol{\Phi} \boldsymbol{\Sigma}_j \boldsymbol{\Phi}^\top \boldsymbol{\tau}_j, \end{aligned} \quad (3.18b)$$

and then  $s_{i,j}$  and  $q_{i,j}$  in Eq. (3.17) are computed as, respectively:<sup>2</sup>

$$s_{i,j} = \frac{\alpha_i s'_{i,j}}{\alpha_i - s'_{i,j}}, \quad (3.19a)$$

$$q_{i,j} = \frac{\alpha_i q'_{i,j}}{\alpha_i - s'_{i,j}}. \quad (3.19b)$$

<sup>1</sup>  $s'_{i,j} = \sigma_j^{-2} \boldsymbol{\Phi}_i^\top \boldsymbol{\Phi}_i$  and  $q'_{i,j} = \sigma_j^{-2} \boldsymbol{\Phi}_i^\top \boldsymbol{\tau}_j$  when  $\alpha_i = \infty, \forall i$ .

<sup>2</sup>  $s_{i,j} = s'_{i,j}$  and  $q_{i,j} = q'_{i,j}$  when  $\alpha_i = \infty$ .

$\mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\sigma})$  has a unique maximum with respect to  $\alpha_i$  when the following equation is satisfied:

$$\frac{\partial \ell(\alpha_i, \boldsymbol{\sigma})}{\partial \alpha_i} = \frac{1}{2} \sum_{j=1}^V \left( \frac{1}{\alpha_i} - \frac{1}{\alpha_i + s_{i,j}} - \frac{q_{i,j}^2}{(\alpha_i + s_{i,j})^2} \right) = 0, \quad (3.20)$$

which is a  $(2V - 1)$ -th order polynomial equation of  $\alpha_i$ . This implies that:

- If  $\boldsymbol{\phi}_i$  is “in the model” (i.e.  $\alpha_i < \infty$ ) and  $\alpha_i$  in Eq. (3.20) has at least one positive real root ( $\alpha_i > 0$  as  $\alpha_i$  is inverse variance); then,  $\alpha_i$  is re-estimated,
- If  $\boldsymbol{\phi}_i$  is “in the model” (i.e.  $\alpha_i < \infty$ ) yet  $\alpha_i$  in Eq. (3.20) does not have any positive real root; then,  $\boldsymbol{\phi}_i$  may be deleted (i.e.  $\alpha_i$  is set to be  $\infty$ ),
- If  $\boldsymbol{\phi}_i$  is “out of the model” (i.e.  $\alpha_i = \infty$ ) yet  $\alpha_i$  in Eq. (3.20) has at least one positive real root; then,  $\boldsymbol{\phi}_i$  may be added (i.e.  $\alpha_i$  is set to be a finite value).

In addition,  $\frac{\partial \mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\sigma})}{\partial \sigma_j^2} = 0$  leads to that  $\mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\sigma})$  has a unique maximum with respect to  $\sigma_j^2$  when:

$$\sigma_j^2 = \frac{\|\boldsymbol{\tau}_j - \boldsymbol{\Phi} \boldsymbol{\mu}_j\|^2}{N - \sum_{i=1}^{N+1} \gamma_{i,j}}, \quad (3.21)$$

where  $\gamma_{i,j} \stackrel{\text{def}}{=} 1 - \alpha_{(i-1)} \Sigma_{j,ii}$ , and  $\Sigma_{j,ii}$  is the  $i$ -th diagonal element of  $\boldsymbol{\Sigma}_j \in \mathbb{R}^{(N+1) \times (N+1)}$ .

### 3.4. Expectation-maximisation (EM) algorithm

Algorithm 1, an EM algorithm to maximise the marginal likelihood, starts without any basis vector (i.e.  $M = 0$ ) and selects the basis vector  $\boldsymbol{\phi}_i$  which gives the maximum change of the marginal likelihood  $\mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\sigma})$  of Eq. (3.11) at every iteration.

For efficient computation of the EM algorithm, quantities  $\boldsymbol{\Phi} \in \mathbb{R}^{N \times M}$ ,  $\boldsymbol{\Sigma}_j \in \mathbb{R}^{M \times M}$ , and  $\boldsymbol{\mu}_j \in \mathbb{R}^{M \times 1}$  contain only  $M$  ( $M \leq N + 1$  is always satisfied) basis functions that are currently included in the model (i.e.  $\boldsymbol{\phi}_i$  satisfying  $\alpha_i < \infty$ ), and the diagonal matrix  $\mathbf{A}$  consists of  $M$  hyperparameters of  $\alpha_i$  that are currently included in the model (i.e.  $\alpha_i$  satisfying  $\alpha_i < \infty$ ). Additionally, Eq. (3.21) is rewritten as

$$\sigma_j^2 = \frac{\|\boldsymbol{\tau}_j - \boldsymbol{\Phi} \boldsymbol{\mu}_j\|^2}{N - \sum_{i=1}^M \gamma'_{i,j}}, \quad (3.22)$$

where  $\gamma'_{i,j} \stackrel{\text{def}}{=} 1 - \alpha'_i \Sigma_{j,ii}$ ,  $\alpha'_i$  is the  $i$ -th non-infinity value of  $\boldsymbol{\alpha}$ , and  $\Sigma_{j,ii}$  is the  $i$ -th diagonal element of  $\boldsymbol{\Sigma}_j \in \mathbb{R}^{M \times M}$ .

From Eq. (3.11), the change in the marginal likelihood can be written as

$$\begin{aligned} 2\Delta \mathcal{L} &= 2(\mathcal{L}(\tilde{\boldsymbol{\alpha}}, \boldsymbol{\sigma}) - \mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\sigma})) \\ &= \sum_{j=1}^V \left( \log \frac{|\mathbf{C}_j|}{|\tilde{\mathbf{C}}_j|} + \boldsymbol{\tau}_j^T (\mathbf{C}_j^{-1} - \tilde{\mathbf{C}}_j^{-1}) \boldsymbol{\tau}_j \right), \end{aligned} \quad (3.23)$$

where updated quantities are denoted by a tilde (e.g.,  $\tilde{\boldsymbol{\alpha}}$  and  $\tilde{\mathbf{C}}_j$ ). Eq. (3.23) differs according to whether  $\alpha_i$  is re-estimated, added, or deleted:

---

**Algorithm 1:** Existing EM algorithm of MRVR.

---

**Input:**  $\mathbf{T} \in \mathbb{R}^{N \times V}$ , and  $\boldsymbol{\phi}_i \in \mathbb{R}^{N \times 1}$ ,  $\forall i = \{0, 1, \dots, N\}$ , where  $N$  is the number of training samples, and  $V$  is the number of output dimensions

**Output:**  $\boldsymbol{\Sigma}_j \in \mathbb{R}^{M \times M}$ ,  $\boldsymbol{\mu}_j \in \mathbb{R}^{M \times 1}$ , and  $\sigma_j$ ,  $\forall j = \{1, 2, \dots, V\}$ , where  $M$  is the number of basis functions in the model

// Initialisation

```

1  $\alpha_i \leftarrow \infty, \forall i = \{0, 1, \dots, N\}$ 
2 for  $j \leftarrow 1$  to  $V$  do
3    $\bar{t}_j \leftarrow \frac{1}{N} \sum_{i=1}^N t_{i,j}$ 
4    $\sigma_j^2 \leftarrow \frac{0.1}{N-1} \sum_{i=1}^N (t_{i,j} - \bar{t}_j)^2$ 
5 end
6 convergence  $\leftarrow$  false,  $n \leftarrow 1$ ,  $M \leftarrow 0$ , where  $n$  is the iteration number, and  $M$  is the number of basis functions.
7 while convergence = false do
8   // maximisation step
9   for  $i \leftarrow 0$  to  $N$  do
10    for  $j \leftarrow 1$  to  $V$  do
11      Update  $s'_{i,j}$  and  $q'_{i,j}$  using Eq. (3.18), and Update  $s_{i,j}$  and  $q_{i,j}$  using Eq. (3.19).
12    end
13    switch the number of positive real roots of Eq. (3.20) do
14      case 0 do
15         $\tilde{\alpha}_i \leftarrow \infty$ 
16      case 1 do
17         $\tilde{\alpha}_i \leftarrow$  the positive real root of Eq. (3.20)
18      otherwise do
19         $\tilde{\alpha}_i \leftarrow$  one of the positive real roots of Eq. (3.20), which maximises  $2\Delta\mathcal{L}_i$  of either i) Eq. (3.24) if  $\alpha_i < \infty$  or ii) Eq. (3.25) if  $\alpha_i = \infty$ 
20      end
21    end
22    if  $\tilde{\alpha}_i < \infty$  then
23      if  $\alpha_i < \infty$  then  $z_i \leftarrow$  're-estimation'
24      | Update  $2\Delta\mathcal{L}_i$  using Eq. (3.24).
25      else  $z_i \leftarrow$  'addition'
26      | Update  $2\Delta\mathcal{L}_i$  using Eq. (3.25).
27      end
28    else if  $\alpha_i < \infty$  then  $z_i \leftarrow$  'deletion'
29    | Update  $2\Delta\mathcal{L}_i$  using Eq. (3.26).
30    else
31       $2\Delta\mathcal{L}_i \leftarrow -\infty$ 
32    end
33  end
34   $i \leftarrow \arg \max_i 2\Delta\mathcal{L}_i$  // Select  $i$  which gives the greatest increase of the marginal likelihood
35  if  $n \neq 1$  then
36    | Update  $\sigma_j, \forall j$  using Eq. (3.22).
37  end
38  switch  $z_i$  do
39    case 're-estimation' do
40       $\Delta \log \alpha \leftarrow \log \frac{\alpha_i}{\tilde{\alpha}_i}$ 
41       $\alpha_i \leftarrow \tilde{\alpha}_i$ 
42      // Check convergence (convergence criteria are the same as those in (Tipping and Faul 2003))
43      if  $|\Delta \log \alpha| < 0.1$  then
44        convergence  $\leftarrow$  true
45        for  $i \leftarrow 0$  to  $N$  do
46          if  $\alpha_i = \infty$  then // if  $\phi_i$  is "out of the model"
47            if  $\tilde{\alpha}_i < \infty$  then // if  $\phi_i$  may be added
48              convergence  $\leftarrow$  false
49              break for loop
50            end
51          end
52        end
53      end
54    case 'addition' do
55       $\alpha_i \leftarrow \tilde{\alpha}_i$ ,  $M \leftarrow M + 1$ 
56    case 'deletion' do
57       $\alpha_i \leftarrow \infty$ ,  $M \leftarrow M - 1$ 
58    end
59  end
60  // Expectation step
61  Sequentially update i)  $\boldsymbol{\Phi} \in \mathbb{R}^{N \times M}$ ,  $\mathbf{A} \in \mathbb{R}^{M \times M}$ , ii)  $\boldsymbol{\Sigma}_j \in \mathbb{R}^{M \times M}, \forall j$ , and iii)  $\boldsymbol{\mu}_j \in \mathbb{R}^M, \forall j$  using Eq. (3.8) and Eq. (3.9), where  $\boldsymbol{\Phi}$ ,  $\boldsymbol{\Sigma}_j$ , and  $\boldsymbol{\mu}_j$  contain only  $M$  basis functions that are currently included in the model, and the diagonal matrix  $\mathbf{A}$  consists of  $M$  hyperparameters of  $\alpha_i$  that are currently included in the model.
62   $n \leftarrow n + 1$ 
63 end

```

---

**Re-estimation.** as  $\mathbf{C}_j = \mathbf{C}_{-i,j} + \alpha_i^{-1} \boldsymbol{\phi}_i \boldsymbol{\phi}_i^\top$  and  $\tilde{\mathbf{C}} = \mathbf{C}_{-i,j} + \tilde{\alpha}_i^{-1} \boldsymbol{\phi}_i \boldsymbol{\phi}_i^\top$ ,

$$2\Delta\mathcal{L}_i = \sum_{j=1}^V \left( \frac{q_{i,j}^2}{s'_{i,j} + (\tilde{\alpha}_i^{-1} - \alpha_i^{-1})^{-1}} - \log(1 + s'_{i,j}(\tilde{\alpha}_i^{-1} - \alpha_i^{-1})) \right), \quad (3.24)$$

where  $\tilde{\alpha}_i$  is re-estimated  $\alpha_i$ ,

**Addition.** as  $\mathbf{C}_j = \mathbf{C}_{-i,j}$  and  $\tilde{\mathbf{C}}_j = \mathbf{C}_{-i,j} + \tilde{\alpha}_i^{-1} \boldsymbol{\phi}_i \boldsymbol{\phi}_i^\top$ ,

$$2\Delta\mathcal{L}_i = \sum_{j=1}^V \left( \frac{q_{i,j}^2}{\tilde{\alpha}_i + s_{i,j}} + \log \frac{\tilde{\alpha}_i}{\tilde{\alpha}_i + s_{i,j}} \right), \quad (3.25)$$

**Deletion.** as  $\mathbf{C}_j = \mathbf{C}_{-i,j} + \alpha_i^{-1} \boldsymbol{\phi}_i \boldsymbol{\phi}_i^\top$  and  $\tilde{\mathbf{C}}_j = \mathbf{C}_{-i,j}$ ,

$$2\Delta\mathcal{L}_i = \sum_{j=1}^V \left( \frac{q_{i,j}^2}{s'_{i,j} - \alpha_i} - \log \left( 1 - \frac{s'_{i,j}}{\alpha_i} \right) \right). \quad (3.26)$$

### 3.5. Making predictions

We can predict both the mean of  $j$ -th output dimension  $y_{*,j}$  and its variance  $\sigma_{*,j}^2$  from a new input vector  $\mathbf{x}_*$  based on both i) Eq. (3.2), the model specification, and ii) Eq. (3.7), the posterior distribution over the weights, conditioned on the most probable (MP) hyperparameters:  $\boldsymbol{\alpha}_{\text{MP}} \in \mathbb{R}_{>0}^{M \times 1}$  and  $\boldsymbol{\sigma}_{\text{MP}} \in \mathbb{R}_{\geq 0}^{1 \times V}$ , obtained from Algorithm 1. Predictive distribution of  $t_{*,j}$  is normally distributed as

$$p(t_{*,j} | \mathbf{T}, \boldsymbol{\alpha}_{\text{MP}}, \boldsymbol{\sigma}_{\text{MP}}) = \mathcal{N}(t_{*,j} | y_{*,j}, \sigma_{*,j}^2), \quad (3.27)$$

with

$$y_{*,j} = \boldsymbol{\phi}(\mathbf{x}_*)^\top \boldsymbol{\mu}_j, \quad (3.28)$$

and

$$\sigma_{*,j}^2 = \sigma_{\text{MP},j}^2 + \boldsymbol{\phi}(\mathbf{x}_*)^\top \boldsymbol{\Sigma}_j \boldsymbol{\phi}(\mathbf{x}_*), \quad (3.29)$$

where  $\boldsymbol{\phi}(\mathbf{x}_*) \in \mathbb{R}^{M \times 1}$  comes from only  $M$  basis functions that are included in the model after the EM algorithm, and subscript  $j$  refers to the  $j$ -th output dimension. The predictive variance  $\sigma_{*,j}^2$  comprises the sum of two variance components: the estimated noise on the training data  $\sigma_{\text{MP},j}^2$  and that due to the uncertainty in the prediction of the weights  $\boldsymbol{\phi}(\mathbf{x}_*)^\top \boldsymbol{\Sigma}_j \boldsymbol{\phi}(\mathbf{x}_*)$ .

### 3.6. Algorithm complexity

Matrix inversion of  $\boldsymbol{\Sigma}_j \in \mathbb{R}^{M \times M}$  in Eq. (3.8) for all  $j \in \{1, 2, \dots, V\}$  determines i) the time complexity of the existing algorithm as  $O(VM^3)$  and ii) the memory complexity as  $O(VM^2)$ , where  $V$  is the number of output dimensions, and  $M$  is the number of basis functions.<sup>1</sup>

<sup>1</sup> The matrix multiplication to calculate  $s'_{i,j}$  and  $q'_{i,j}$  in Eq. (3.18) for all  $i \in \{1, 2, \dots, N\}$ ,  $j \in \{1, 2, \dots, V\}$  at the 11-th line of Algorithm 1 has the same time complexity because the matrix multiplication  $\boldsymbol{\Phi} \boldsymbol{\Sigma}_j \boldsymbol{\Phi}^\top$  is pre-calculated. In other words, the time complexity of the matrix multiplication is  $O(VM^3)$ , not  $O(NVM^3)$ , because  $\boldsymbol{\Phi} \boldsymbol{\Sigma}_j \boldsymbol{\Phi}^\top$  is independent of  $i$ .



## 4. Proposed method

### 4.1. Model specification

Given a data set of input-target pairs  $\{\mathbf{x}_i \in \mathbb{R}^{U \times 1}, \mathbf{t}_i \in \mathbb{R}^{1 \times V}\}_{i=1}^N$ , where  $N$  is the number of training samples, it is assumed that the targets  $\mathbf{t}_i$  are samples from the model  $\mathbf{y}(\mathbf{x}_i; \mathbf{W})$  with additive noise:

$$\mathbf{t}_i = \mathbf{y}(\mathbf{x}_i; \mathbf{W}) + \boldsymbol{\epsilon}_i, \quad (4.1)$$

where  $\mathbf{W} \in \mathbb{R}^{(N+1) \times V}$  is the weight and  $\boldsymbol{\epsilon}_i \in \mathbb{R}^{1 \times V}$  are independent samples from a Gaussian noise process with mean zero and a covariance matrix  $\boldsymbol{\Omega} \in \mathbb{R}^{V \times V}$ .

Eq. (4.1) can be rewritten, using matrix algebra, as

$$\mathbf{T} = \boldsymbol{\Phi} \mathbf{W} + \mathbf{E}, \quad (4.2)$$

where  $\mathbf{T} = \begin{bmatrix} \mathbf{t}_1 \\ \mathbf{t}_2 \\ \vdots \\ \mathbf{t}_N \end{bmatrix} \in \mathbb{R}^{N \times V}$  is the target,  $\mathbf{E} = \begin{bmatrix} \boldsymbol{\epsilon}_1 \\ \boldsymbol{\epsilon}_2 \\ \vdots \\ \boldsymbol{\epsilon}_N \end{bmatrix} \in \mathbb{R}^{N \times V}$  is

the noise,  $\boldsymbol{\Phi} = [\boldsymbol{\phi}(\mathbf{x}_1) \ \boldsymbol{\phi}(\mathbf{x}_2) \ \dots \ \boldsymbol{\phi}(\mathbf{x}_N)]^\top \in \mathbb{R}^{N \times (N+1)}$  is a design matrix,  $\boldsymbol{\phi}(\mathbf{x}) = [1 \ K(\mathbf{x}, \mathbf{x}_1) \ K(\mathbf{x}, \mathbf{x}_2) \ \dots \ K(\mathbf{x}, \mathbf{x}_N)]^\top \in \mathbb{R}^{(N+1) \times 1}$ , and  $K(\mathbf{x}, \mathbf{x}')$  is a kernel function.

Because of the assumption of independent Gaussian noise, the likelihood of the data set can be given by the matrix Gaussian distribution:

$$p(\mathbf{T}|\mathbf{W}, \boldsymbol{\Omega}) = (2\pi)^{-\frac{VN}{2}} |\boldsymbol{\Omega}|^{-\frac{N}{2}} \exp\left(-\frac{1}{2} \text{tr}(\boldsymbol{\Omega}^{-1} (\mathbf{T} - \boldsymbol{\Phi} \mathbf{W})^\top (\mathbf{T} - \boldsymbol{\Phi} \mathbf{W}))\right), \quad (4.3)$$

where  $\boldsymbol{\Omega} = \frac{\mathbb{E}[\mathbf{E}^\top \mathbf{E}]}{N}$ , and  $\text{tr}$  denotes trace.<sup>1</sup> As I assumed,  $\mathbf{I} = \frac{\mathbb{E}[\mathbf{E} \mathbf{E}^\top]}{\text{tr}(\boldsymbol{\Omega})}$ , which means the noise is independent among rows with the same variance, where  $\mathbf{I}$  is an  $N \times N$  identity matrix. For clarity, the implicit conditioning on the input  $\mathbf{x}_i, \forall i$  is omitted in Eq. (4.3) and the subsequent expressions.

An assumption to avoid over-fitting in the estimation of  $\mathbf{W}$  is

$$p(\mathbf{W}|\boldsymbol{\alpha}, \boldsymbol{\Omega}) = (2\pi)^{-\frac{V(N+1)}{2}} |\boldsymbol{\Omega}|^{-\frac{N+1}{2}} |\mathbf{A}|^{\frac{V}{2}} \exp\left(-\frac{1}{2} \text{tr}(\boldsymbol{\Omega}^{-1} \mathbf{W}^\top \mathbf{A} \mathbf{W})\right), \quad (4.4)$$

where  $\mathbf{A}^{-1} = \text{diag}(\alpha_0^{-1}, \alpha_1^{-1}, \dots, \alpha_N^{-1}) = \frac{\mathbb{E}[\mathbf{W} \mathbf{W}^\top]}{\text{tr}(\boldsymbol{\Omega})}$ . This means the prior distribution of  $\mathbf{W}$  is zero-mean Gaussian with among-row inverse variances  $\boldsymbol{\alpha} = [\alpha_0 \ \alpha_1 \ \dots \ \alpha_N]^\top \in \mathbb{R}_{>0}^{(N+1) \times 1}$ , which are  $N+1$  hyperparameters (Tipping 2001). Eq. (4.4) implies another assumption:  $\boldsymbol{\Omega} = \frac{\mathbb{E}[\mathbf{W}^\top \mathbf{W}]}{\text{tr}(\mathbf{A}^{-1})}$ . Actually, this is unreasonable because the weight  $\mathbf{W}$  has no relationship with the noise  $\mathbf{E}$  (i.e.  $\mathbf{I} = \frac{\mathbb{E}[\mathbf{W}^\top \mathbf{W}]}{\text{tr}(\mathbf{A}^{-1})}$ , which means that the weights of different output dimensions are not correlated, is a reasonable assumption), but it is essential for creating a computationally efficient algorithm.

<sup>1</sup> If  $\boldsymbol{\Omega} = \text{diag}(\sigma_1^2, \sigma_2^2, \dots, \sigma_V^2)$ , then Eq. (4.3) will be Eq. (3.4).

## 4.2. Inference

By both the Bayes' theorem and the property of  $p(\mathbf{T}|\mathbf{W}, \boldsymbol{\alpha}, \boldsymbol{\Omega}) = p(\mathbf{T}|\mathbf{W}, \boldsymbol{\Omega})$ ,<sup>1</sup> the posterior probability distribution function over  $\mathbf{W}$  is decomposed as

$$p(\mathbf{W}|\mathbf{T}, \boldsymbol{\alpha}, \boldsymbol{\Omega}) = \frac{p(\mathbf{T}|\mathbf{W}, \boldsymbol{\Omega}) p(\mathbf{W}|\boldsymbol{\alpha}, \boldsymbol{\Omega})}{p(\mathbf{T}|\boldsymbol{\alpha}, \boldsymbol{\Omega})}, \quad (4.5)$$

and it is given by the matrix Gaussian distribution:<sup>2</sup>

$$p(\mathbf{W}|\mathbf{T}, \boldsymbol{\alpha}, \boldsymbol{\Omega}) = (2\pi)^{-\frac{V(N+1)}{2}} |\boldsymbol{\Omega}|^{-\frac{N+1}{2}} |\boldsymbol{\Sigma}|^{-\frac{V}{2}} \exp\left(-\frac{1}{2} \text{tr}(\boldsymbol{\Omega}^{-1} (\mathbf{W} - \mathbf{M})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{W} - \mathbf{M}))\right), \quad (4.6)$$

where the posterior covariance and mean are, respectively:

$$\boldsymbol{\Sigma} = (\boldsymbol{\Phi}^\top \boldsymbol{\Phi} + \mathbf{A})^{-1}, \quad (4.7)$$

$$\mathbf{M} = \boldsymbol{\Sigma} \boldsymbol{\Phi}^\top \mathbf{T}. \quad (4.8)$$

In the case of uniform hyperpriors  $\boldsymbol{\alpha}$  and  $\boldsymbol{\Omega}$ , maximising a posteriori  $p(\boldsymbol{\alpha}, \boldsymbol{\Omega}|\mathbf{T}) \propto p(\mathbf{T}|\boldsymbol{\alpha}, \boldsymbol{\Omega}) p(\boldsymbol{\alpha}) p(\boldsymbol{\Omega})$  is equivalent to maximising the marginal likelihood  $p(\mathbf{T}|\boldsymbol{\alpha}, \boldsymbol{\Omega})$ , which is given by:

$$p(\mathbf{T}|\boldsymbol{\alpha}, \boldsymbol{\Omega}) = (2\pi)^{-\frac{VN}{2}} |\boldsymbol{\Omega}|^{-\frac{N}{2}} |\mathbf{I} + \boldsymbol{\Phi} \mathbf{A}^{-1} \boldsymbol{\Phi}^\top|^{-\frac{V}{2}} \exp\left(-\frac{1}{2} \text{tr}\left(\boldsymbol{\Omega}^{-1} \mathbf{T}^\top (\mathbf{I} + \boldsymbol{\Phi} \mathbf{A}^{-1} \boldsymbol{\Phi}^\top)^{-1} \mathbf{T}\right)\right). \quad (4.9)$$

## 4.3. Marginal likelihood maximisation

The same method of accelerating the univariate relevance vector machine (Tipping and Faul 2003) is used to accelerate the proposed algorithm.

The log of Eq. (4.9) is an objective function:

$$\mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\Omega}) = -\frac{1}{2} (VN \log(2\pi) + N \log |\boldsymbol{\Omega}| + V \log |\mathbf{C}| + \text{tr}(\boldsymbol{\Omega}^{-1} \mathbf{T}^\top \mathbf{C}^{-1} \mathbf{T})), \quad (4.10)$$

where  $\mathbf{C} = \mathbf{I} + \boldsymbol{\Phi} \mathbf{A}^{-1} \boldsymbol{\Phi}^\top \in \mathbb{R}^{N \times N}$ , and by considering the dependence of  $\mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\Omega})$  on a single hyperparameter  $\alpha_i, i \in \{0, 1, \dots, N\}$ ,  $\mathbf{C}$  is decomposed as the following two parts:

$$\begin{aligned} \mathbf{C} &= \mathbf{I} + \sum_{m \neq i} \alpha_m^{-1} \boldsymbol{\phi}_m \boldsymbol{\phi}_m^\top + \alpha_i^{-1} \boldsymbol{\phi}_i \boldsymbol{\phi}_i^\top \\ &= \mathbf{C}_{-i} + \alpha_i^{-1} \boldsymbol{\phi}_i \boldsymbol{\phi}_i^\top, \end{aligned} \quad (4.11)$$

where  $\mathbf{C}_{-i} \in \mathbb{R}^{N \times N}$  is  $\mathbf{C}$  with the contribution of a basis vector  $\boldsymbol{\phi}_i \in \mathbb{R}^{N \times 1}$  removed, and

$$\boldsymbol{\phi}_i = \begin{cases} [1 \ 1 \ \dots \ 1]^\top, & \text{if } i = 0 \\ [K(\mathbf{x}_i, \mathbf{x}_1) \ K(\mathbf{x}_i, \mathbf{x}_2) \ \dots \ K(\mathbf{x}_i, \mathbf{x}_N)]^\top, & \text{otherwise} \end{cases}. \quad (4.12)$$

<sup>1</sup> In the case that the weight  $\mathbf{W}$  is given, its inverse variances  $\boldsymbol{\alpha}$  are redundant in the calculation of the conditional probability of the target  $\mathbf{T}$ .

<sup>2</sup> The proof is in Appendix.

The determinant and inverse matrix of  $\mathbf{C}$  are, respectively:

$$|\mathbf{C}| = |\mathbf{C}_{-i}| (1 + \alpha_i^{-1} \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i}^{-1} \boldsymbol{\Phi}_i), \quad (4.13)$$

by Sylvester's determinant theorem, and

$$\mathbf{C}^{-1} = \mathbf{C}_{-i}^{-1} - \frac{\mathbf{C}_{-i}^{-1} \boldsymbol{\Phi}_i \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i}^{-1}}{\alpha_i + \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i}^{-1} \boldsymbol{\Phi}_i}, \quad (4.14)$$

by Woodbury matrix identity. From these,  $\mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\Omega})$  in Eq. (4.10) can be decomposed into  $\mathcal{L}(\boldsymbol{\alpha}_{-i}, \boldsymbol{\Omega})$ , the marginal likelihood with  $\boldsymbol{\Phi}_i$  excluded, and  $\ell(\alpha_i, \boldsymbol{\Omega})$ , the isolated marginal likelihood of  $\boldsymbol{\Phi}_i$ :

$$\begin{aligned} \mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\Omega}) &= -\frac{1}{2} (VN \log(2\pi) + N \log |\boldsymbol{\Omega}| + V \log |\mathbf{C}_{-i}| + \text{tr}(\boldsymbol{\Omega}^{-1} \mathbf{T}^\top \mathbf{C}_{-i}^{-1} \mathbf{T})) \\ &\quad - \frac{1}{2} \left( -V \log \alpha_i + V \log (\alpha_i + \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i}^{-1} \boldsymbol{\Phi}_i) - \text{tr} \left( \boldsymbol{\Omega}^{-1} \mathbf{T}^\top \frac{\mathbf{C}_{-i}^{-1} \boldsymbol{\Phi}_i \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i}^{-1}}{\alpha_i + \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i}^{-1} \boldsymbol{\Phi}_i} \mathbf{T} \right) \right) \\ &= \mathcal{L}(\boldsymbol{\alpha}_{-i}, \boldsymbol{\Omega}) + \frac{1}{2} \left( V \log \alpha_i - V \log (\alpha_i + s_i) + \frac{\text{tr}(\boldsymbol{\Omega}^{-1} \mathbf{q}_i^\top \mathbf{q}_i)}{\alpha_i + s_i} \right) \\ &= \mathcal{L}(\boldsymbol{\alpha}_{-i}, \boldsymbol{\Omega}) + \ell(\alpha_i, \boldsymbol{\Omega}), \end{aligned} \quad (4.15)$$

where  $s_i$  and  $\mathbf{q}_i \in \mathbb{R}^{1 \times V}$  are defined as, respectively:

$$s_i \stackrel{\text{def}}{=} \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i}^{-1} \boldsymbol{\Phi}_i, \quad (4.16a)$$

$$\mathbf{q}_i \stackrel{\text{def}}{=} \boldsymbol{\Phi}_i^\top \mathbf{C}_{-i}^{-1} \mathbf{T}. \quad (4.16b)$$

To avoid the matrix inversion of  $\mathbf{C}_{-i}$  in Eq. (4.16), which requires the time complexity of  $O(N^3)$ ,  $s'_i$  and  $\mathbf{q}'_i \in \mathbb{R}^{1 \times V}$  are computed as (by the Woodbury matrix identity):<sup>1</sup>

$$\begin{aligned} s'_i &= \boldsymbol{\Phi}_i^\top \mathbf{C}^{-1} \boldsymbol{\Phi}_i \\ &= \boldsymbol{\Phi}_i^\top \boldsymbol{\Phi}_i - \boldsymbol{\Phi}_i^\top \boldsymbol{\Phi} \boldsymbol{\Sigma} \boldsymbol{\Phi}^\top \boldsymbol{\Phi}_i, \end{aligned} \quad (4.17a)$$

$$\begin{aligned} \mathbf{q}'_i &= \boldsymbol{\Phi}_i^\top \mathbf{C}^{-1} \mathbf{T} \\ &= \boldsymbol{\Phi}_i^\top \mathbf{T} - \boldsymbol{\Phi}_i^\top \boldsymbol{\Phi} \boldsymbol{\Sigma} \boldsymbol{\Phi}^\top \mathbf{T}, \end{aligned} \quad (4.17b)$$

and then  $s_i$  and  $\mathbf{q}_i$  in Eq. (4.16) are computed as:<sup>2</sup>

$$s_i = \frac{\alpha_i s'_i}{\alpha_i - s'_i}, \quad (4.18a)$$

$$\mathbf{q}_i = \frac{\alpha_i \mathbf{q}'_i}{\alpha_i - s'_i}. \quad (4.18b)$$

<sup>1</sup>  $s'_i = \boldsymbol{\Phi}_i^\top \boldsymbol{\Phi}_i$  and  $\mathbf{q}'_i = \boldsymbol{\Phi}_i^\top \mathbf{T}$  when  $\alpha_i = \infty, \forall i$ .

<sup>2</sup>  $s_i = s'_i$  and  $\mathbf{q}_i = \mathbf{q}'_i$  when  $\alpha_i = \infty$ .

$\frac{\partial \ell(\alpha_i, \mathbf{\Omega})}{\partial \alpha_i} = 0$  leads to that  $\mathcal{L}(\mathbf{\alpha}, \mathbf{\Omega})$  has a unique maximum with respect to  $\alpha_i$  when:

$$\alpha_i = \begin{cases} \frac{s_i^2}{\frac{\text{tr}(\mathbf{\Omega}^{-1} \mathbf{q}_i^\top \mathbf{q}_i)}{V} - s_i}, & \text{if } \frac{\text{tr}(\mathbf{\Omega}^{-1} \mathbf{q}_i^\top \mathbf{q}_i)}{V} > s_i \\ \infty, & \text{if } \frac{\text{tr}(\mathbf{\Omega}^{-1} \mathbf{q}_i^\top \mathbf{q}_i)}{V} \leq s_i \end{cases}, \quad (4.19)$$

which implies that:

- If  $\mathbf{\Phi}_i$  is “in the model” (i.e.  $\alpha_i < \infty$ ) and  $\frac{\text{tr}(\mathbf{\Omega}^{-1} \mathbf{q}_i^\top \mathbf{q}_i)}{V} > s_i$ ; then,  $\alpha_i$  is re-estimated,
- If  $\mathbf{\Phi}_i$  is “in the model” (i.e.  $\alpha_i < \infty$ ) yet  $\frac{\text{tr}(\mathbf{\Omega}^{-1} \mathbf{q}_i^\top \mathbf{q}_i)}{V} \leq s_i$ ; then,  $\mathbf{\Phi}_i$  may be deleted (i.e.  $\alpha_i$  is set to be  $\infty$ ),
- If  $\mathbf{\Phi}_i$  is “out of the model” (i.e.  $\alpha_i = \infty$ ) yet  $\frac{\text{tr}(\mathbf{\Omega}^{-1} \mathbf{q}_i^\top \mathbf{q}_i)}{V} > s_i$ ; then,  $\mathbf{\Phi}_i$  may be added (i.e.  $\alpha_i$  is set to be a finite value).

In addition,  $\frac{\partial \mathcal{L}(\mathbf{\alpha}, \mathbf{\Omega})}{\partial \mathbf{\Omega}} = \mathbf{0}$ , where  $\mathbf{0}$  is a  $V \times V$  zero matrix, leads to that  $\mathcal{L}(\mathbf{\alpha}, \mathbf{\Omega})$  has a unique maximum with respect to  $\mathbf{\Omega}$  when:

$$\mathbf{\Omega} = \frac{\mathbf{T}^\top (\mathbf{T} - \mathbf{\Phi} \mathbf{M})}{N}. \quad (4.20)$$

#### 4.4. Expectation-maximisation (EM) algorithm

Algorithm 2, an EM algorithm to maximise the marginal likelihood, starts without any basis vector (i.e.  $M = 0$ ) and selects the basis vector  $\mathbf{\Phi}_i$  which gives the maximum change of the marginal likelihood  $\mathcal{L}(\mathbf{\alpha}, \mathbf{\Omega})$  of Eq. (4.10) at every iteration.

For efficient computation of the EM algorithm, quantities  $\mathbf{\Phi} \in \mathbb{R}^{N \times M}$  and  $\mathbf{\Sigma} \in \mathbb{R}^{M \times M}$  contain only  $M$  ( $M \leq N + 1$  is always satisfied) basis functions that are currently included in the model (i.e.  $\mathbf{\Phi}_i$  which satisfies  $\alpha_i < \infty$ ), and the diagonal matrix  $\mathbf{A}$  consists of  $M$  hyperparameters of  $\alpha_i$  that are currently included in the model (i.e.  $\alpha_i$  which satisfies  $\alpha_i < \infty$ ).

From Eq. (4.10), the change in the marginal likelihood can be written as

$$\begin{aligned} 2\Delta \mathcal{L} &= 2(\mathcal{L}(\tilde{\mathbf{\alpha}}, \mathbf{\Omega}) - \mathcal{L}(\mathbf{\alpha}, \mathbf{\Omega})) \\ &= V \log \frac{|\mathbf{C}|}{|\tilde{\mathbf{C}}|} + \text{tr} \left( \mathbf{\Omega}^{-1} \mathbf{T}^\top (\mathbf{C}^{-1} - \tilde{\mathbf{C}}^{-1}) \mathbf{T} \right), \end{aligned} \quad (4.21)$$

where updated quantities are denoted by a tilde (e.g.,  $\tilde{\mathbf{\alpha}}$  and  $\tilde{\mathbf{C}}$ ). Eq. (4.21) differs according to whether  $\alpha_i$  is re-estimated, added, or deleted:

**Re-estimation.** as  $\mathbf{C} = \mathbf{C}_{-i} + \alpha_i^{-1} \mathbf{\Phi}_i \mathbf{\Phi}_i^\top$  and  $\tilde{\mathbf{C}} = \mathbf{C}_{-i} + \tilde{\alpha}_i^{-1} \mathbf{\Phi}_i \mathbf{\Phi}_i^\top$ ,

$$2\Delta \mathcal{L}_i = \frac{\text{tr}(\mathbf{\Omega}^{-1} \mathbf{q}_i'^\top \mathbf{q}_i')}{s_i' + (\tilde{\alpha}_i^{-1} - \alpha_i^{-1})^{-1}} - V \log(1 + s_i' (\tilde{\alpha}_i^{-1} - \alpha_i^{-1})), \quad (4.22)$$

where  $\tilde{\alpha}_i$  is re-estimated  $\alpha_i$ ,

---

**Algorithm 2:** Proposed EM algorithm of MRVR.

---

**Input:**  $\mathbf{T} \in \mathbb{R}^{N \times V}$  and  $\boldsymbol{\phi}_i \in \mathbb{R}^{N \times 1}, \forall i = \{0, 1, \dots, N\}$ , where  $N$  is the number of training samples, and  $V$  is the number of output dimensions

**Output:**  $\boldsymbol{\Sigma} \in \mathbb{R}^{M \times M}$ ,  $\mathbf{M} \in \mathbb{R}^{M \times V}$ , and  $\boldsymbol{\Omega} \in \mathbb{R}^{V \times V}$ , where  $M$  is the number of basis functions in the model

**// Initilisation**

- 1  $\alpha_i \leftarrow \infty, \forall i = \{0, 1, \dots, N\}$
- 2  $\bar{\mathbf{t}} \leftarrow \frac{1}{N} \sum_{i=1}^N \mathbf{t}_i$ , where  $\bar{\mathbf{t}} \in \mathbb{R}^{1 \times V}$ , and  $\mathbf{t}_i \in \mathbb{R}^{1 \times V}$  is the  $i$ -th row vector of  $\mathbf{T}$ .
- 3  $\boldsymbol{\Omega} \leftarrow \frac{0.1}{N-1} \sum_{i=1}^N (\mathbf{t}_i - \bar{\mathbf{t}})^\top (\mathbf{t}_i - \bar{\mathbf{t}})$ , where  $\boldsymbol{\Omega} \in \mathbb{R}^{V \times V}$  is a covariance matrix.
- 4 convergence  $\leftarrow$  false
- 5  $n \leftarrow 1$ , where  $n$  is the iteration number
- 6  $M \leftarrow 0$ , where  $M$  is the number of basis functions.
- 7 **while** convergence = false **do**
  - // maximisation step**
  - 8 **for**  $i \leftarrow 0$  **to**  $N$  **do**
    - 9 Update  $s'_i$  and  $\mathbf{q}'_i$  using Eq. (4.17).
    - 10 Update  $s_i$  and  $\mathbf{q}_i$  using Eq. (4.18).
    - 11  $\theta_i \leftarrow \frac{\text{tr}(\boldsymbol{\Omega}^{-1} \mathbf{q}_i^\top \mathbf{q}_i)}{V} - s_i$
    - 12 **if**  $\theta_i > 0$  **then**
      - 13 **if**  $\alpha_i < \infty$  **then**  $z_i \leftarrow$  're-estimation'
      - 14  $\tilde{\alpha}_i \leftarrow \frac{s_i^2}{\theta_i}$
      - 15 Update  $2\Delta\mathcal{L}_i$  using Eq. (4.22).
      - 16 **else**  $z_i \leftarrow$  'addition'
      - 17 Update  $2\Delta\mathcal{L}_i$  using Eq. (4.23).
      - 18 **end**
      - 19 **else if**  $\alpha_i < \infty$  **then**  $z_i \leftarrow$  'deletion'
      - 20 Update  $2\Delta\mathcal{L}_i$  using Eq. (4.24).
      - 21 **else**
      - 22  $2\Delta\mathcal{L}_i \leftarrow -\infty$
      - 23 **end**
  - 24 **end**
  - 25  $i \leftarrow \arg \max_i 2\Delta\mathcal{L}_i$  // Select  $i$  which gives the greatest increase of the marginal likelihood
  - 26 **switch**  $z_i$  **do**
    - 27 **case** 're-estimation' **do**
      - 28  $\Delta \log \alpha \leftarrow \log \frac{\alpha_i}{\tilde{\alpha}_i}$
      - 29  $\alpha_i \leftarrow \tilde{\alpha}_i$
      - 30 **// Check convergence** (convergence criteria are the same as those in (Tipping and Faul 2003))
      - 31 **if**  $|\Delta \log \alpha| < 0.1$  **then**
        - 32 convergence  $\leftarrow$  true
        - 33 **for**  $i \leftarrow 0$  **to**  $N$  **do**
          - 34 **if**  $\alpha_i = \infty$  **then** // if  $\phi_i$  is "out of the model"
          - 35 **if**  $\theta_i > 0$  **then** // if  $\phi_i$  may be added
          - 36 convergence  $\leftarrow$  false
          - 37 break for loop
          - 38 **end**
        - 39 **end**
      - 40 **end**
    - 41 **case** 'addition' **do**
      - 42  $\alpha_i \leftarrow \frac{s_i^2}{\theta_i}$
      - 43  $M \leftarrow M + 1$
    - 44 **case** 'deletion' **do**
      - 45  $\alpha_i \leftarrow \infty$
      - 46  $M \leftarrow M - 1$
  - 47 **end**
  - 48 **if**  $n \neq 1$  **then**
    - 49 Update  $\boldsymbol{\Omega}$  using Eq. (4.20).
  - 50 **end**
  - // Expectation step**
  - 51 Sequentially update i)  $\boldsymbol{\Phi} \in \mathbb{R}^{N \times M}$ ,  $\mathbf{A} \in \mathbb{R}^{M \times M}$ , ii)  $\boldsymbol{\Sigma} \in \mathbb{R}^{M \times M}$ , and iii)  $\mathbf{M} \in \mathbb{R}^{M \times V}$  using Eq. (4.7) and Eq. (4.8), where  $\boldsymbol{\Phi}$ ,  $\boldsymbol{\Sigma}$ , and  $\mathbf{M}$  contain only  $M$  basis functions that are currently included in the model, and the diagonal matrix  $\mathbf{A}$  consists of  $M$  hyperparameters of  $\alpha_i$  that are currently included in the model.
  - 52  $n \leftarrow n + 1$
- 53 **end**

---

**Addition.** as  $\mathbf{C} = \mathbf{C}_{-i}$  and  $\tilde{\mathbf{C}} = \mathbf{C}_{-i} + \tilde{\alpha}_i^{-1} \boldsymbol{\phi}_i \boldsymbol{\phi}_i^\top$ , where  $\tilde{\alpha}_i = \frac{s_i^2}{\frac{\text{tr}(\boldsymbol{\Omega}^{-1} \mathbf{q}_i^\top \mathbf{q}_i)}{V} - s_i}$ ,

$$2\Delta\mathcal{L}_i = \frac{\text{tr}(\boldsymbol{\Omega}^{-1} \mathbf{q}_i'^\top \mathbf{q}_i') - V s_i'}{s_i'} + V \log \frac{V s_i'}{\text{tr}(\boldsymbol{\Omega}^{-1} \mathbf{q}_i'^\top \mathbf{q}_i')}, \quad (4.23)$$

**Deletion.** as  $\mathbf{C} = \mathbf{C}_{-i} + \alpha_i^{-1} \boldsymbol{\phi}_i \boldsymbol{\phi}_i^\top$  and  $\tilde{\mathbf{C}} = \mathbf{C}_{-i}$ ,

$$2\Delta\mathcal{L}_i = \frac{\text{tr}(\boldsymbol{\Omega}^{-1} \mathbf{q}_i'^\top \mathbf{q}_i')}{s_i' - \alpha_i} - V \log \left( 1 - \frac{s_i'}{\alpha_i} \right). \quad (4.24)$$

#### 4.5. Making predictions

We can predict both a mean vector  $\mathbf{y}_* \in \mathbb{R}^{1 \times V}$  and a covariance matrix  $\boldsymbol{\Omega}_* \in \mathbb{R}^{V \times V}$  from a new input vector  $\mathbf{x}_* \in \mathbb{R}^{U \times 1}$  based on both i) Eq. (4.2), the model specification, and ii) Eq. (4.6), the posterior distribution over the weights, conditioned on the most probable (MP) hyperparameters:  $\boldsymbol{\alpha}_{\text{MP}} \in \mathbb{R}_{>0}^{M \times 1}$  and  $\boldsymbol{\Omega}_{\text{MP}} \in \mathbb{R}^{V \times V}$ , obtained from Algorithm 2. Predictive distribution of  $\mathbf{t}_*$  is jointly normally distributed as

$$p(\mathbf{t}_* | \mathbf{T}, \boldsymbol{\alpha}_{\text{MP}}, \boldsymbol{\Omega}_{\text{MP}}) = \mathcal{N}(\mathbf{t}_* | \mathbf{y}_*, \boldsymbol{\Omega}_*), \quad (4.25)$$

with

$$\mathbf{y}_* = \boldsymbol{\Phi}(\mathbf{x}_*)^\top \mathbf{M}, \quad (4.26)$$

and

$$\boldsymbol{\Omega}_* = \boldsymbol{\Omega}_{\text{MP}} (1 + \boldsymbol{\Phi}(\mathbf{x}_*)^\top \boldsymbol{\Sigma} \boldsymbol{\Phi}(\mathbf{x}_*)),^1 \quad (4.27)$$

where  $\boldsymbol{\Phi}(\mathbf{x}) \in \mathbb{R}^{M \times 1}$  comes from only  $M$  basis functions that are included in the model after the EM algorithm. The predictive covariance matrix  $\boldsymbol{\Omega}_*$  comprises the two components: the estimated noise on the training data  $\boldsymbol{\Omega}_{\text{MP}}$  and that due to the uncertainty in the prediction of the weights  $\boldsymbol{\Omega}_{\text{MP}} \boldsymbol{\Phi}(\mathbf{x}_*)^\top \boldsymbol{\Sigma} \boldsymbol{\Phi}(\mathbf{x}_*)$ , where  $\boldsymbol{\Phi}(\mathbf{x}_*)^\top \boldsymbol{\Sigma} \boldsymbol{\Phi}(\mathbf{x}_*) \in \mathbb{R}_{\geq 0}$  by the fact that a covariance matrix is always positive semidefinite. They share  $\boldsymbol{\Omega}_{\text{MP}}$  by the assumption of  $\boldsymbol{\Omega} = \frac{\mathbb{E}[\mathbf{E}^\top \mathbf{E}]}{N} = \frac{\mathbb{E}[\mathbf{W}^\top \mathbf{W}]}{\text{tr}(\mathbf{A}^{-1})}$  in Section 4.1.

#### 4.6. Algorithm complexity

Matrix inversion of  $\boldsymbol{\Omega} \in \mathbb{R}^{V \times V}$  in Eq. (4.19) and that of the  $M \times M$  matrix in Eq. (4.7) determine i) the time complexity of the proposed algorithm as  $O(V^3 + M^3)$  and ii) the memory complexity as  $O(V^2 + M^2)$ , where  $V$  is the number of output dimensions, and  $M$  is the number of basis functions.<sup>2</sup>

<sup>1</sup> Eq. (4.27) is obtained by the property that the covariance between two elements  $W_{i,j}$  and  $W_{i',j'}$  is the covariance between the rows  $i$  and  $i'$ , i.e.  $\boldsymbol{\Sigma}$ , multiplied by the covariance between the columns  $j$  and  $j'$ , i.e.  $\boldsymbol{\Omega}_{\text{MP}}$  (Arnold 1981, p. 311).

<sup>2</sup> The matrix multiplication to calculate  $s_i'$  and  $\mathbf{q}_i'$  in Eq. (4.17) for all  $i \in \{1, 2, \dots, N\}$  at the 9-th line of Algorithm 2 does not influence the time complexity because the matrix multiplication  $\boldsymbol{\Phi} \boldsymbol{\Sigma} \boldsymbol{\Phi}^\top$  is pre-calculated. In other words, the time complexity of the matrix multiplication is  $O(M^3)$ , not  $O(NM^3)$ , because  $\boldsymbol{\Phi} \boldsymbol{\Sigma} \boldsymbol{\Phi}^\top$  is independent of  $i$ .


(a) Existing method ( $\sigma_{*,j}^2$  is equal to Eq. (3.29))(b) Proposed method ( $\sigma_{*,j}^2$  is equal to the  $j$ -th diagonal element of  $\mathbf{\Omega}_*$  in Eq. (4.27))

Figure 1.: An example of MRVR (when  $U = 1, V = 2, N = 200$ , and the Gaussian kernel  $K(\mathbf{x}, \mathbf{x}') = \exp\left(-\frac{\|\mathbf{x} - \mathbf{x}'\|^2}{2\lambda^2}\right)$  with a free parameter  $\lambda = 1.6$  is used)

## 5. Experimental results

### 5.1. An example of MRVR

Fig. 1 shows an example of the MRVR results obtained using the two methods when the true function of each output dimension is the sinc function and the linear function, respectively. Fig. 1a and Fig. 1b show slightly different results although the same training samples are used.

### 5.2. Comparisons of the performance

The two methods are compared in terms of i) running time (computation time in INTEL® Core™ i5-3470 CPU and MATLAB® R2013b), ii) the estimation accuracy of the noise covariance matrix, iii) root-mean-square error (RMSE) between true functions and predicted mean values, and iv) the number of relevance vectors (RVs), where RVs are those training vectors associated with the remaining non-zero weights (i.e. the number of basis functions  $M$  is equal to the number of RVs).




Figure 2.: True functions of MC simulations

To measure the estimation accuracy of the noise covariance matrix  $\mathbf{\Omega}$ , entropy loss  $L_1(\mathbf{\Omega}, \hat{\mathbf{\Omega}})$  and quadratic loss  $L_2(\mathbf{\Omega}, \hat{\mathbf{\Omega}})$  are used (each of these is 0 when  $\hat{\mathbf{\Omega}} = \mathbf{\Omega}$  and is positive when  $\hat{\mathbf{\Omega}} \neq \mathbf{\Omega}$ ) (Anderson 1984, pp. 273–274):

$$L_1(\mathbf{\Omega}, \hat{\mathbf{\Omega}}) = \text{tr}(\hat{\mathbf{\Omega}}\mathbf{\Omega}^{-1}) - \log |\hat{\mathbf{\Omega}}\mathbf{\Omega}^{-1}| - V, \quad (5.1)$$

$$L_2(\mathbf{\Omega}, \hat{\mathbf{\Omega}}) = \text{tr} \left( \left( \hat{\mathbf{\Omega}}\mathbf{\Omega}^{-1} - \mathbf{I} \right)^2 \right), \quad (5.2)$$

where the estimated  $V \times V$  covariance matrix of the noise  $\hat{\mathbf{\Omega}}$  is  $\mathbf{\Omega}_{\text{MP}}$  in the case of the proposed method. In the case of the existing method,  $\hat{\mathbf{\Omega}}$  can be obtained using both i) the estimated standard deviation of the noise  $\hat{\mathbf{D}} = \text{diag}(\sigma_{\text{MP},1}, \sigma_{\text{MP},2}, \dots, \sigma_{\text{MP},V})$  in Section 3.5 and ii) the estimated correlation matrix of the noise  $\hat{\mathbf{R}}$ :

$$\hat{\mathbf{\Omega}} = \hat{\mathbf{D}}\hat{\mathbf{R}}\hat{\mathbf{D}}, \quad (5.3)$$

where  $\hat{\mathbf{R}} = \tilde{\mathbf{D}}^{-1}\tilde{\mathbf{\Omega}}\tilde{\mathbf{D}}^{-1}$ ,  $\tilde{\mathbf{\Omega}} = \frac{(\mathbf{T} - \mathbf{\Phi}\tilde{\mathbf{M}})^\top(\mathbf{T} - \mathbf{\Phi}\tilde{\mathbf{M}})}{N-1}$ ,  $\tilde{\mathbf{M}} = [\boldsymbol{\mu}_1 \ \boldsymbol{\mu}_2 \ \dots \ \boldsymbol{\mu}_V] \in \mathbb{R}^{M \times V}$ , and  $\tilde{\mathbf{D}} = \sqrt{\text{diag}(\tilde{\mathbf{\Omega}})}$ .

Monte Carlo (MC) simulations with random covariance matrices of the noise were conducted for the performance comparisons. The noise from the random covariance matrix is added to the true functions in Fig. 2 (each output has a sinc function with a translation in the x-axis), and the two methods of MRVR with the Gaussian kernel were performed using the same training samples for a fair comparison.

Unpaired two-sample  $t$ -tests may be used to compare the two methods to determine whether the performance difference is fundamental or whether it is due to random fluctuations (Simon 2013, pp. 631–635), but the normality assumption of the performance measures (i.e. running time, entropy loss, quadratic loss, RMSE, and the number of RVs) of the two methods must be checked. The Jarque–Bera tests  $JB = \frac{n}{6} (S^2 + \frac{1}{4}(K-3)^2)$  with the number of observations  $n = 101$  and



Table 1.: The number of rejections of the null hypothesis of the Jarque–Bera test

	Running time	Entropy loss	Quadratic loss	RMSE	The number of RVs
Existing method	30	30	30	4	16
Proposed method	30	28	29	3	14

Table 2.: The difference in median values of running time (seconds)

	$N = 50$	$N = 100$	$N = 150$	$N = 200$	$N = 250$	$N = 300$
$V = 1$	<b>0.04</b> ( $3.6 \times 10^{-10}$ )	<b>0.19</b> ( $2.1 \times 10^{-9}$ )	<b>0.40</b> ( $4.9 \times 10^{-9}$ )	<b>1.02</b> ( $3.2 \times 10^{-9}$ )	<b>3.11</b> ( $1.8 \times 10^{-20}$ )	<b>5.95</b> ( $7.1 \times 10^{-18}$ )
$V = 2$	<b>0.20</b> ( $4.8 \times 10^{-25}$ )	<b>0.76</b> ( $1.5 \times 10^{-22}$ )	<b>2.44</b> ( $9.0 \times 10^{-26}$ )	<b>5.95</b> ( $2.1 \times 10^{-25}$ )	<b>17.20</b> ( $1.3 \times 10^{-30}$ )	<b>26.17</b> ( $1.0 \times 10^{-31}$ )
$V = 3$	<b>0.33</b> ( $3.2 \times 10^{-26}$ )	<b>1.70</b> ( $3.8 \times 10^{-32}$ )	<b>5.27</b> ( $5.4 \times 10^{-29}$ )	<b>10.37</b> ( $1.1 \times 10^{-31}$ )	<b>37.74</b> ( $4.7 \times 10^{-34}$ )	<b>64.99</b> ( $2.2 \times 10^{-34}$ )
$V = 4$	<b>0.74</b> ( $2.2 \times 10^{-33}$ )	<b>3.60</b> ( $1.3 \times 10^{-32}$ )	<b>9.69</b> ( $2.1 \times 10^{-33}$ )	<b>24.51</b> ( $2.4 \times 10^{-33}$ )	<b>57.14</b> ( $2.1 \times 10^{-34}$ )	<b>112.18</b> ( $1.2 \times 10^{-34}$ )
$V = 5$	<b>1.05</b> ( $1.1 \times 10^{-33}$ )	<b>5.25</b> ( $5.4 \times 10^{-34}$ )	<b>13.01</b> ( $6.3 \times 10^{-34}$ )	<b>34.09</b> ( $2.2 \times 10^{-34}$ )	<b>92.96</b> ( $1.2 \times 10^{-34}$ )	<b>176.24</b> ( $1.2 \times 10^{-34}$ )

Table 3.: The difference in median values of entropy loss


	$N = 50$	$N = 100$	$N = 150$	$N = 200$	$N = 250$	$N = 300$
$V = 1$	$3.4 \times 10^{-4}$ (0.9904)	$8.0 \times 10^{-5}$ (0.9176)	$2.2 \times 10^{-5}$ (0.9962)	$1.7 \times 10^{-5}$ (0.9520)	$3.7 \times 10^{-5}$ (0.8454)	$5.7 \times 10^{-5}$ (0.9808)
$V = 2$	$3.9 \times 10^{-3}$ (0.4942)	$3.3 \times 10^{-3}$ (0.5964)	$3.4 \times 10^{-3}$ (0.1132)	$6.8 \times 10^{-4}$ (0.3915)	$1.1 \times 10^{-3}$ (0.7434)	$4.0 \times 10^{-4}$ (0.6234)
$V = 3$	$3.2 \times 10^{-2}$ (0.0883)	$8.6 \times 10^{-3}$ (0.2469)	<b><math>6.8 \times 10^{-3}</math></b> (0.0192)	<b><math>8.0 \times 10^{-3}</math></b> (0.0069)	<b><math>5.1 \times 10^{-3}</math></b> (0.0119)	<b><math>1.4 \times 10^{-3}</math></b> (0.0412)
$V = 4$	$3.9 \times 10^{-2}$ (0.1137)	<b><math>3.0 \times 10^{-2}</math></b> (0.0030)	<b><math>1.5 \times 10^{-2}</math></b> (0.0010)	<b><math>1.1 \times 10^{-2}</math></b> (0.0042)	<b><math>9.3 \times 10^{-3}</math></b> (0.0063)	<b><math>4.4 \times 10^{-3}</math></b> (0.0058)
$V = 5$	<b><math>1.1 \times 10^{-1}</math></b> (0.0231)	<b><math>3.0 \times 10^{-2}</math></b> (0.0015)	<b><math>2.9 \times 10^{-2}</math></b> (0.0000)	<b><math>2.5 \times 10^{-2}</math></b> (0.0008)	<b><math>1.4 \times 10^{-2}</math></b> (0.0005)	<b><math>1.2 \times 10^{-2}</math></b> (0.0018)

a 5% significance level for 30 cases ( $V = \{1, 2, 3, 4, 5\}$  and  $N = \{50, 100, 150, 200, 250, 300\}$ ) were conducted, in which the null hypothesis was that the data of the performance measures came from a normal distribution. Table 1 shows the number of rejections of the null hypothesis. Consequently, the  $t$ -test can yield misleading results in the case that the null hypothesis is rejected.


Instead of the  $t$ -test, two-sided Wilcoxon rank sum tests, whose null hypothesis is that two populations have equal median values, were used for the comparisons as they have greater efficiency than the  $t$ -test on non-normal distributions and are nearly as efficient as the  $t$ -test on normal distributions (Montgomery 2013, Chapter 10).

Fig. 3 shows the median values of the performance measures of both methods with various  $V$  and  $N$  values. Entropy loss, quadratic loss, and RMSE decrease as  $N$  increases: the greater the number of training samples, the more accurate the estimation. In contrast, the number of RVs, the number of iterations of each EM algorithm (the same tolerance value of 0.1 for checking the convergence of each EM algorithm was used as in 43th line of Algorithm 1 and 30th line of Algorithm 2), and the running time (only for learning without prediction) of each EM algorithm increase as  $N$  increases: the greater the number of training samples, the greater the computational burden.


Tables 2–6 show i) the difference in the median values of the performance measures, where each median value is obtained from 101 MC simulations, and then the median value of the proposed method is subtracted from that of the existing method (i.e. positive difference values mean that the proposed method is better than the existing method, while negative difference values mean the




(a) The number of iterations




(b) Running time




(c) Entropy loss



(d) Quadratic loss



(e) RMSE



(f) The number of RVs

Figure 2: Median values of MC simulations (the number of simulations is 101)

Table 4.: The difference in median values of quadratic loss

	$N = 50$	$N = 100$	$N = 150$	$N = 200$	$N = 250$	$N = 300$
$V = 1$	$2.8 \times 10^{-3}$ (0.9981)	$-5.7 \times 10^{-4}$ (0.9233)	$4.9 \times 10^{-5}$ (0.9981)	$2.3 \times 10^{-4}$ (0.9405)	$4.4 \times 10^{-5}$ (0.8605)	$2.0 \times 10^{-4}$ (0.9770)
$V = 2$	$5.5 \times 10^{-3}$ (0.5080)	$3.5 \times 10^{-3}$ (0.4526)	$7.0 \times 10^{-3}$ (0.0915)	$2.2 \times 10^{-3}$ (0.3616)	$1.6 \times 10^{-3}$ (0.6824)	$-1.9 \times 10^{-4}$ (0.5782)
$V = 3$	<b><math>8.5 \times 10^{-2}</math></b> (0.0180)	$2.5 \times 10^{-2}$ (0.1154)	<b><math>1.4 \times 10^{-2}</math></b> (0.0109)	<b><math>1.4 \times 10^{-2}</math></b> (0.0029)	<b><math>1.3 \times 10^{-2}</math></b> (0.0051)	<b><math>2.7 \times 10^{-3}</math></b> (0.0263)
$V = 4$	<b><math>1.7 \times 10^{-1}</math></b> (0.0382)	<b><math>8.6 \times 10^{-2}</math></b> (0.0005)	<b><math>5.4 \times 10^{-2}</math></b> (0.0001)	<b><math>2.4 \times 10^{-2}</math></b> (0.0007)	<b><math>2.9 \times 10^{-2}</math></b> (0.0019)	<b><math>1.5 \times 10^{-2}</math></b> (0.0013)
$V = 5$	<b><math>7.3 \times 10^{-1}</math></b> (0.0058)	<b><math>1.8 \times 10^{-1}</math></b> (0.0000)	<b><math>8.5 \times 10^{-2}</math></b> (0.0000)	<b><math>5.5 \times 10^{-2}</math></b> (0.0001)	<b><math>4.8 \times 10^{-2}</math></b> (0.0000)	<b><math>3.7 \times 10^{-2}</math></b> (0.0004)

Table 5.: The difference in median values of RMSE

	$N = 50$	$N = 100$	$N = 150$	$N = 200$	$N = 250$	$N = 300$
$V = 1$	-0.0038 (0.956)	-0.0000 (0.985)	0.0000 (0.967)	-0.0001 (1.000)	-0.0007 (0.965)	0.0001 (0.987)
$V = 2$	-0.0035 (0.898)	-0.0076 (0.544)	-0.0051 (0.562)	-0.0033 (0.758)	-0.0064 (0.646)	-0.0020 (0.977)
$V = 3$	-0.0177 (0.182)	-0.0049 (0.299)	-0.0012 (0.408)	-0.0030 (0.661)	-0.0013 (0.546)	-0.0037 (0.565)
$V = 4$	<b>-0.0121</b> (0.041)	<b>-0.0142</b> (0.034)	<b>-0.0101</b> (0.017)	-0.0037 (0.092)	-0.0038 (0.115)	-0.0029 (0.293)
$V = 5$	<b>-0.0150</b> (0.008)	<b>-0.0026</b> (0.013)	<b>-0.0149</b> (0.001)	<b>-0.0110</b> (0.015)	<b>-0.0068</b> (0.025)	-0.0062 (0.072)

Table 6.: The difference in median values of the number of RVs

	$N = 50$	$N = 100$	$N = 150$	$N = 200$	$N = 250$	$N = 300$
$V = 1$	0 ( $8.8 \times 10^{-1}$ )	1 ( $9.0 \times 10^{-1}$ )	0 ( $8.4 \times 10^{-1}$ )	0 ( $9.6 \times 10^{-1}$ )	0 ( $8.9 \times 10^{-1}$ )	0 ( $9.9 \times 10^{-1}$ )
$V = 2$	-1 ( $5.2 \times 10^{-2}$ )	<b>-1</b> ( $4.8 \times 10^{-4}$ )	<b>-1</b> ( $3.2 \times 10^{-3}$ )	<b>-2</b> ( $5.1 \times 10^{-4}$ )	<b>-2</b> ( $1.5 \times 10^{-3}$ )	<b>-3</b> ( $2.2 \times 10^{-3}$ )
$V = 3$	<b>-3</b> ( $3.6 \times 10^{-7}$ )	<b>-2</b> ( $3.9 \times 10^{-5}$ )	<b>-2</b> ( $1.5 \times 10^{-5}$ )	<b>-3</b> ( $3.4 \times 10^{-7}$ )	<b>-3</b> ( $3.2 \times 10^{-6}$ )	<b>-2</b> ( $2.6 \times 10^{-6}$ )
$V = 4$	<b>-3</b> ( $2.2 \times 10^{-8}$ )	<b>-3</b> ( $4.8 \times 10^{-6}$ )	<b>-2</b> ( $8.6 \times 10^{-5}$ )	<b>-5</b> ( $2.4 \times 10^{-9}$ )	<b>-2</b> ( $2.2 \times 10^{-7}$ )	<b>-2</b> ( $1.8 \times 10^{-4}$ )
$V = 5$	<b>-4</b> ( $3.9 \times 10^{-15}$ )	<b>-3</b> ( $3.9 \times 10^{-10}$ )	<b>-3</b> ( $3.9 \times 10^{-8}$ )	<b>-4</b> ( $9.0 \times 10^{-10}$ )	<b>-3</b> ( $6.9 \times 10^{-7}$ )	<b>-6</b> ( $3.2 \times 10^{-7}$ )

opposite) and ii) the  $p$ -values of the Wilcoxon rank sum tests, which appear inside the brackets. The  $p$ -value is interpreted as the probability that a difference in the median values would be obtained given that the population medians of two methods are equivalent, i.e. the  $p$ -value is not equal to the probability that the population medians are equivalent (Simon 2013, p. 635). Note that statistically significant difference values are marked in bold ( $p$ -value  $< 0.05$ ).

The proposed method is faster than the existing method as shown in Table 2 (all differences are statistically significant). In particular, the time difference is amplified as  $V$  or  $N$  increases. This is because the time complexity of the proposed method  $O(V^3 + M^3)$  is less than that of the existing method  $O(VM^3)$  ( $O(V^3 + M^3) < O(VM^3)$  is satisfied since  $V < M$  is satisfied in most applications). Note that even when the number of input dimensions  $U$  changes, the size of the design matrix  $\Phi$  does not change. Hence,  $U$  does not influence the time complexity of both the methods.

Furthermore, the proposed method achieves higher accuracy in estimating the covariance matrix of the noise  $\mathbf{\Omega}$  than the existing method as shown in Table 3 and Table 4. This is because the proposed method considers the correlation matrix of the noise as Eq. (4.3), but the existing method does not as Eq. (4.3).

However, the proposed method is worse than the existing one in terms of i) the accuracy in predicting the mean values as shown in Table 5 (in particular, the RMSE increases in the region of high  $V$  and low  $N$ ) and ii) the number of RVs as shown in Table 6. This is because the proposed method has the assumption of the weight  $\mathbf{\Omega} = \frac{\mathbb{E}[\mathbf{W}^T \mathbf{W}]}{\text{tr}(\mathbf{A}^{-1})}$ , which behaves as the constraint of the weight. Consequently, the mean values tend to deviate from the true functions, and the number of RVs increases.

The MATLAB codes of the experiment have been uploaded on <http://www.mathworks.com/matlabcentral/fileexchange/49131> to avoid any potential ambiguity of both the methods.

## 6. Conclusion

A new algorithm of MRVR has been proposed. It is more efficient in computing the weight  $\mathbf{W}$  and more accurate in estimating the covariance matrix of the noise  $\mathbf{\Omega}$  than the existing algorithm. Its computational efficiency and accuracy can be attributed to the different model specifications: the existing method expresses the likelihood of the training data as the product of the Gaussian distributions in Eq. (3.4), but the proposed one expresses it as the matrix Gaussian distribution in Eq. (4.3).

However, the proposed method has drawbacks of lower accuracy in estimating the mean of the weight  $\mathbf{M}$  in Eq. (4.8) and higher number of RVs than the existing method. These disadvantages are caused by the assumption  $\mathbf{\Omega} = \frac{\mathbb{E}[\mathbf{W}^T \mathbf{W}]}{\text{tr}(\mathbf{A}^{-1})}$ , which means the weight  $\mathbf{W}$  is related to the noise  $\mathbf{E}$  in Eq. (4.2), but it was indispensable to make MRVR faster.

## References

- Alvarez, M. and Lawrence, N.D., Sparse convolved Gaussian processes for multi-output regression. In *Proceedings of the Advances in Neural Information Processing Systems*, pp. 57--64, 2009.
- Anderson, T.W., *An introduction to multivariate statistical analysis* (2 edn), 1984, Wiley.
- Arnold, S.F., *The theory of linear models and multivariate analysis*, 1981, Wiley.
- Ben-Shimon, D. and Shmilovici, A., Accelerating the relevance vector machine via data partitioning. *Foundations of Computing and Decision Sciences*, 2006, **31**, 27--41.
- Bonilla, E.V., Chai, K.M.A. and Williams, C.K.I., Multi-task Gaussian process prediction. In *Proceedings of the Advances in Neural Information Processing Systems*, pp. 153--160, 2007.
- Boyle, P. and Frean, M.R., Dependent Gaussian Processes.. In *Proceedings of the Advances in Neural Information Processing Systems*, pp. 217--224, 2004.
- Catanzaro, B., Sundaram, N. and Keutzer, K., Fast support vector machine training and classification on graphics processors. In *Proceedings of the Proceedings of the 25th international conference on Machine learning*, pp. 104--111, 2008.
- Chang, C.C. and Lin, C.J., LIBSVM: A library for support vector machines. *ACM Transactions on Intelligent Systems and Technology*, 2011, **2**, 27:1--27.
- Chu, W., Keerthi, S.S. and Ong, C.J., Bayesian support vector regression using a unified loss function. *IEEE Transactions on Neural Networks*, 2004, **15**, 29--44.
- Cortes, C. and Vapnik, V., Support-vector networks. *Machine learning*, 1995, **20**, 273--297.
- Gao, J.B., Gunn, S.R., Harris, C.J. and Brown, M., A probabilistic framework for SVM regression and error bar estimation. *Machine Learning*, 2002, **46**, 71--89.
- Gibbs, M., Bayesian Gaussian processes for classification and regression. PhD thesis, University of Cambridge, 1997.
- Gramacy, R.B., Niemi, J. and Weiss, R.M., Massively parallel approximate Gaussian process regression. *SIAM/ASA Journal on Uncertainty Quantification*, 2014, **2**, 564--584.
- Guo, G. and Zhang, J.S., Reducing examples to accelerate support vector regression. *Pattern Recognition Letters*, 2007, **28**, 2173--2183.

- Montgomery, D.C., *Applied Statistics and Probability for Engineers* (6 edn), 2013, Wiley.
- Pérez-Cruz, F., Camps-Valls, G., Soria-Olivas, E., Pérez-Ruixo, J.J., Figueiras-Vidal, A.R. and Artés-Rodríguez, A., Multi-dimensional function approximation and regression estimation. In *Proceedings of the International Conference on Artificial Neural Networks*, pp. 757–762, 2002.
- Schölkopf, B., Smola, A.J., Williamson, R.C. and Bartlett, P.L., New support vector algorithms. *Neural computation*, 2000, **12**, 1207–1245.
- Seeger, M., Williams, C. and Lawrence, N., Fast forward selection to speed up sparse Gaussian process regression. In *Proceedings of the Workshop on Artificial Intelligence and Statistics 9*, 2003.
- Shen, Y., Ng, A. and Seeger, M., Fast Gaussian process regression using kd-trees. In *Proceedings of the 19th Annual Conference on Neural Information Processing Systems*, 2006.
- Simon, D., *Evolutionary optimization algorithms*, 2013, Wiley.
- Srinivasan, B.V., Qi, H. and Duraiswami, R., GPUML: Graphical processors for speeding up kernel machines. In *Proceedings of the Workshop on High Performance Analytics - Algorithms, Implementations, and Applications*, 2010.
- Thayananthan, A., Template-based pose estimation and tracking of 3D hand motion. PhD thesis, University of Cambridge, 2005.
- Thayananthan, A., Navaratnam, R., Stenger, B., Torr, P.H. and Cipolla, R., Pose estimation and tracking using multivariate regression. *Pattern Recognition Letters*, 2008, **29**, 1302–1310.
- Tipping, M.E., Sparse Bayesian learning and the relevance vector machine. *The journal of machine learning research*, 2001, **1**, 211–244.
- Tipping, M.E. and Faul, A.C., Fast marginal likelihood maximisation for sparse Bayesian models. In *Proceedings of the Proceedings of the ninth international workshop on artificial intelligence and statistics*, Vol. 1, 2003.
- Tuia, D., Verrelst, J., Alonso, L., Pérez-Cruz, F. and Camps-Valls, G., Multioutput support vector regression for remote sensing biophysical parameter estimation. *IEEE Geoscience and Remote Sensing Letters*, 2011, **8**, 804–808.
- Vapnik, V., *The nature of statistical learning theory*, 2000, Springer.
- Vazquez, E. and Walter, E., Multi-output support vector regression. In *Proceedings of the 13th IFAC Symposium on System Identification*, pp. 1820–1825, 2003.
- Yang, D., Liang, G., Jenkins, D.D., Peterson, G.D. and Li, H., High performance relevance vector machine on GPUs. In *Proceedings of the Symposium on Application Accelerators in High-Performance Computing*, 2010.

### Appendix: proof of Eq. (4.6) and Eq. (4.9)

$$\begin{aligned}
& p(\mathbf{W}|\mathbf{T}, \boldsymbol{\alpha}, \boldsymbol{\Omega}) p(\mathbf{T}|\boldsymbol{\alpha}, \boldsymbol{\Omega}) \\
&= p(\mathbf{T}|\mathbf{W}, \boldsymbol{\Omega}) p(\mathbf{W}|\boldsymbol{\alpha}, \boldsymbol{\Omega}) \\
&= \rho \exp\left(-\frac{1}{2} \text{tr}(\boldsymbol{\Omega}^{-1}(\mathbf{T} - \boldsymbol{\Phi}\mathbf{W})^\top (\mathbf{T} - \boldsymbol{\Phi}\mathbf{W}))\right) \exp\left(-\frac{1}{2} \text{tr}(\boldsymbol{\Omega}^{-1}\mathbf{W}^\top \mathbf{A}\mathbf{W})\right), \\
&\quad \text{where } \rho = (2\pi)^{-\frac{VN}{2}} |\boldsymbol{\Omega}|^{-\frac{N}{2}} (2\pi)^{-\frac{V(N+1)}{2}} |\boldsymbol{\Omega}|^{-\frac{N+1}{2}} |\mathbf{A}|^{\frac{V}{2}} \\
&= \rho \exp\left(-\frac{1}{2} \text{tr}(\boldsymbol{\Omega}^{-1}((\mathbf{T}^\top - \mathbf{W}^\top \boldsymbol{\Phi}^\top)(\mathbf{T} - \boldsymbol{\Phi}\mathbf{W}) + \mathbf{W}^\top \mathbf{A}\mathbf{W}))\right) \\
&= \rho \exp\left(-\frac{1}{2} \text{tr}(\boldsymbol{\Omega}^{-1}(\mathbf{T}^\top \mathbf{T} - \mathbf{T}^\top \boldsymbol{\Phi}\mathbf{W} - \mathbf{W}^\top \boldsymbol{\Phi}^\top \mathbf{T} + \mathbf{W}^\top \boldsymbol{\Sigma}^{-1} \mathbf{W}))\right), \text{ where } \boldsymbol{\Sigma} = (\boldsymbol{\Phi}^\top \boldsymbol{\Phi} + \mathbf{A})^{-1} \\
&= \rho \exp\left(-\frac{1}{2} \text{tr}(\boldsymbol{\Omega}^{-1}(\mathbf{T}^\top \mathbf{T} - \mathbf{T}^\top \boldsymbol{\Phi}\mathbf{W} + \mathbf{W}^\top \boldsymbol{\Sigma}^{-1}(\mathbf{W} - \mathbf{M})))\right), \text{ where } \mathbf{M} = \boldsymbol{\Sigma} \boldsymbol{\Phi}^\top \mathbf{T} \\
&= \rho \exp\left(-\frac{1}{2} \text{tr}(\boldsymbol{\Omega}^{-1}(\mathbf{T}^\top \mathbf{T} - \mathbf{T}^\top \boldsymbol{\Phi}\mathbf{W} + \mathbf{W}^\top \boldsymbol{\Sigma}^{-1}(\mathbf{W} - \mathbf{M}) - \mathbf{M}^\top \boldsymbol{\Sigma}^{-1}(\mathbf{W} - \mathbf{M}) + \mathbf{M}^\top \boldsymbol{\Sigma}^{-1}(\mathbf{W} - \mathbf{M})))\right) \\
&= \rho \exp\left(-\frac{1}{2} \text{tr}(\boldsymbol{\Omega}^{-1}((\mathbf{W} - \mathbf{M})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{W} - \mathbf{M}) + \mathbf{M}^\top \boldsymbol{\Sigma}^{-1}(\mathbf{W} - \mathbf{M}) + \mathbf{T}^\top \mathbf{T} - \mathbf{T}^\top \boldsymbol{\Phi}\mathbf{W}))\right) \\
&= \rho \exp\left(-\frac{1}{2} \text{tr}(\boldsymbol{\Omega}^{-1}((\mathbf{W} - \mathbf{M})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{W} - \mathbf{M}) + \mathbf{T}^\top \boldsymbol{\Phi}(\mathbf{W} - \mathbf{M}) + \mathbf{T}^\top \mathbf{T} - \mathbf{T}^\top \boldsymbol{\Phi}\mathbf{W}))\right), \\
&\quad \text{since } \boldsymbol{\Sigma} = \boldsymbol{\Sigma}^\top \\
&= \rho \exp\left(-\frac{1}{2} \text{tr}(\boldsymbol{\Omega}^{-1}((\mathbf{W} - \mathbf{M})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{W} - \mathbf{M}) + \mathbf{T}^\top (\mathbf{T} - \boldsymbol{\Phi}\mathbf{M})))\right) \\
&= \rho \exp\left(-\frac{1}{2} \text{tr}\left(\boldsymbol{\Omega}^{-1}\left((\mathbf{W} - \mathbf{M})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{W} - \mathbf{M}) + \mathbf{T}^\top \left(\mathbf{I} - \boldsymbol{\Phi}(\boldsymbol{\Phi}^\top \boldsymbol{\Phi} + \mathbf{A})^{-1} \boldsymbol{\Phi}^\top\right) \mathbf{T}\right)\right)\right) \\
&= \rho \exp\left(-\frac{1}{2} \text{tr}\left(\boldsymbol{\Omega}^{-1}\left((\mathbf{W} - \mathbf{M})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{W} - \mathbf{M}) + \mathbf{T}^\top (\mathbf{I} + \boldsymbol{\Phi}\mathbf{A}^{-1} \boldsymbol{\Phi}^\top)^{-1} \mathbf{T}\right)\right)\right), \\
&\quad \text{by the Woodbury matrix identity} \\
&= (2\pi)^{-\frac{V(N+1)}{2}} |\boldsymbol{\Omega}|^{-\frac{N+1}{2}} |\boldsymbol{\Sigma}|^{-\frac{V}{2}} \exp\left(-\frac{1}{2} \text{tr}(\boldsymbol{\Omega}^{-1}(\mathbf{W} - \mathbf{M})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{W} - \mathbf{M}))\right) \\
&\quad (2\pi)^{-\frac{VN}{2}} |\boldsymbol{\Omega}|^{-\frac{N}{2}} |\mathbf{I} + \boldsymbol{\Phi}\mathbf{A}^{-1} \boldsymbol{\Phi}^\top|^{-\frac{V}{2}} \exp\left(-\frac{1}{2} \text{tr}\left(\boldsymbol{\Omega}^{-1} \mathbf{T}^\top (\mathbf{I} + \boldsymbol{\Phi}\mathbf{A}^{-1} \boldsymbol{\Phi}^\top)^{-1} \mathbf{T}\right)\right)
\end{aligned}$$