

# Counting Process Based Dimension Reduction Methods for Censored Outcomes\*

Qiang Sun<sup>†</sup> Ruoqing Zhu<sup>‡</sup> Tao Wang<sup>§</sup> Donglin Zeng<sup>¶</sup>

March 1, 2022

## Abstract

We propose a class of dimension reduction methods for right censored survival data using a counting process representation of the failure process. Semiparametric estimating equations are constructed to estimate the dimension reduction subspace for the failure time model. The proposed method addresses two fundamental limitations of existing approaches. First, using the counting process formulation, it does not require any estimation of the censoring distribution to compensate the bias in estimating the dimension reduction subspace. Second, the nonparametric part in the estimating equations is adaptive to the structural dimension, hence the approach circumvents the curse of dimensionality. Asymptotic normality is established for the obtained estimators. We

---

\*The first two authors contribute equally.

<sup>†</sup>Department of Statistical Sciences, University of Toronto, Toronto, ON M5S 3G3; e-mail: [qsun@utstat.toronto.edu](mailto:qsun@utstat.toronto.edu).

<sup>‡</sup>The corresponding author. Department of Statistics, University of Illinois at Urbana-Champaign, Champaign, IL 61820; email: [rqzhu@illinois.edu](mailto:rqzhu@illinois.edu).

<sup>§</sup>Department of Bioinformatics and Biostatistics, Shanghai Jiao Tong University, Shanghai, China; email: [neowangtao@sjtu.edu.cn](mailto:neowangtao@sjtu.edu.cn).

<sup>¶</sup>Department of Biostatistics, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599; email: [dzeng@email.unc.edu](mailto:dzeng@email.unc.edu).

further propose a computationally efficient approach that simplifies the estimation equation formulations and requires only a singular value decomposition to estimate the dimension reduction subspace. Numerical studies suggest that our new approaches exhibit significantly improved performance for estimating the true dimension reduction subspace. We further conduct a real data analysis on a skin cutaneous melanoma dataset from The Cancer Genome Atlas. The proposed method is implemented in the R package “orthoDr”.

**keywords** Sufficient Dimension Reduction; Survival Analysis; Estimating Equations; Semi-parametric Inference; Sliced Inverse Regression.

## 1 Introduction

Dimension reduction is an important problem in regression analysis. It aims to extract a low-dimensional subspace from a  $p$ -dimensional covariates  $X = (X_1, \dots, X_p)^T$ , to predict an outcome of interest  $T$ . The dimension reduction literature often assumes the multiple-index model

$$T = h(B^T X, \epsilon), \quad (1.1)$$

where  $\epsilon$  is a random error independent of  $X$ ,  $B \in \mathbb{R}^{p \times d}$  is a coefficient matrix with  $d < p$ , and  $h(\cdot)$  is a completely unknown link function. This model is equivalent to assuming  $T \perp X \mid B^T X$  (Li, 1991). Since any  $d$  linearly independent vectors in the linear space spanned by the columns of  $B$  also satisfy model (1.1) for some  $h$ , we define  $\mathcal{S}(B)$  to be this linear subspace. Furthermore, we call the intersection of all such subspaces satisfying  $T \perp X \mid B^T X$  the central subspace, denoted by  $\mathcal{S}_{T|X}$ , whose dimension is called the structural dimension. According to Cook (2009),  $\mathcal{S}_{T|X}$  is uniquely defined under mild conditions. The goal of sufficient dimension reduction in (1.1) is to determine the structural dimension and the central subspace using empirical data.

There is an extensive literature on estimating the central subspace for completely observed data, including the seminal paper from [Li \(1991\)](#) and subsequent works such as [Cook & Weisberg \(1991\)](#), [Zhu et al. \(2006\)](#), [Li & Wang \(2007\)](#), [Xia \(2007\)](#), and [Ma & Zhu \(2012\)](#). When  $T$  is subject to right censoring, which frequently occurs in survival analysis, model (1.1) includes many well-known survival models as special cases, for instance, the Cox proportional hazard model ([Cox, 1972](#)), the accelerated failure-time model ([Lin et al., 1998](#)), and the linear transformation models ([Zeng & Lin, 2007](#)).

There has been limited work on estimating the dimension reduction subspace using censored observations. [Li et al. \(1999\)](#) propose a modified sliced inverse regression method which uses the estimate of the conditional survival function to account for censored cases. [Xia et al. \(2010\)](#) propose to estimate the conditional hazard function nonparametrically and utilize its gradient and local linear regression to construct the dimension reduction directions. In [Li et al. \(1999\)](#), a  $p$  dimensional kernel estimation is used for compensating the bias caused by censoring, while in [Xia et al. \(2010\)](#), the estimation procedure also requires a  $p$  dimensional kernel hazard function to provide reliable initial value, and then gradually reduce the dimension to  $d$ . Hence these methods inevitably suffer from the curse of dimensionality. When  $p$  is not small, some alternative approaches such as [Lu & Li \(2011\)](#) adopt an inverse probability weighting scheme, which implicitly requires the correct specification of the censoring mechanism.

In this paper, we propose a counting process-based dimension-reduction framework that leads to four different approaches. The proposed methods enjoy unique advantages and address several limitations of the existing literature. First, our approach is built upon a counting process representation of the underlying survival model. This allows a construction of doubly robust estimating equations, and the resulting estimator is more stable than existing approaches such as [Xia et al. \(2010\)](#). This formulation can avoid the linearity assumption ([Li, 1991](#)) and the estimation of any censoring distribution, which are necessary

components of [Li et al. \(1999\)](#) and [Lu & Li \(2011\)](#). Second, the proposed framework is adaptive to the structural dimension in the sense that the involved nonparametric estimations only depend on the dimension of  $\mathcal{S}(B)$ , which is usually small, thus circumvents the curse of dimensionality. To this end, the proposed method shares similar advantage as [Xia et al. \(2010\)](#). However, computationally, we utilize an optimization approach on the Stiefel manifold ([Wen & Yin, 2013](#)) to solve the estimating equations, which is numerically stable and fast. Last, under some restrictive assumptions, our method reduces to a computationally efficient approach that can directly estimate the dimension reduction subspace without nonparametric smoothing.

## 2 Proposed methods

### 2.1 Semiparametric estimating equations for the central subspace

Throughout the paper, we denote the failure time by  $T$  and the censoring time by  $C$ . Let  $Y = \min(T, C)$  and  $\delta = I(T \leq C)$  be the observed event time and the censoring indicator, respectively. We assume that  $C$  is independent of  $T$  conditional on  $X$ . Let  $N(u) = I(Y \leq u, \delta = 1)$  and  $Y(u) = I(Y \geq u)$  denote the observed counting process and the at-risk process, respectively. Let  $\lambda(u|X)$  be the conditional hazard for  $T$  given  $X$ . Due to [Xia et al. \(2010\)](#), model (1.1) is equivalent to  $\lambda(u|X) = \lambda(u|B^T X)$ . We further let  $dM(u, X) = dM(u, B^T X) = dN(u) - \lambda(u|B^T X)Y(u)du$  be the martingale increment process indexed by  $u$ . This paper centers on constructing estimation equations that are built upon the counting process representation of the survival model. To derive the estimating equations, we follow [Bickel et al. \(1993\)](#) and [Tsiatis \(2007\)](#) to obtain the ortho-complement

of the nuisance tangent space at  $B$  as

$$\mathcal{E}^\perp = \left\{ \int \{ \alpha(u, X) - \alpha^*(u, B^\top X) \} dM(u, X) : \right. \\ \left. \alpha(u, X) \text{ is a measurable function of } X \text{ and } u \right\}, \quad (2.1)$$

where

$$\alpha^*(u, B^\top X) = E\{ \alpha(u, X) | \mathcal{F}_u, B^\top X \},$$

and  $\mathcal{F}_u$  the filtration. The derivation can be found in the Supplementary Material. To estimate  $B$ , we consider the unbiased estimating equations

$$E \left[ \int \{ \alpha(u, X) - \alpha^*(u, B^\top X) \} \{ dN(u) - \lambda(u|B^\top X) Y(u) du \} \right] = 0. \quad (2.2)$$

The sample versions based on  $n$  independent and identical copies  $\{Y_i, \delta_i, X_i\}_{i=1}^n$  are given by

$$n^{-1} \sum_{i=1}^n \left[ \int \{ \alpha(u, X_i) - \alpha^*(u, B^\top X_i) \} \{ dN_i(u) - \lambda(u|B^\top X_i) Y_i(u) du \} \right] = 0, \quad (2.3)$$

where the conditional hazard function will be estimated using the data. For some particular choices of  $\alpha(u, X)$ , this can be implemented utilizing the generalized method of moments ([Hansen, 1982](#)):

$$B = \arg \min_{B \in \Theta} \{ \psi_n(B)^\top \psi_n(B) \}, \quad (2.4)$$

where  $\psi_n(B)$  is the left hand side of (2.3). Several quantities in  $\psi_n(B)$  need to be estimated nonparametrically. For example, the conditional hazard function  $\lambda(u|B^\top X_i)$  at any time point  $u$  can be estimated by

$$\hat{\lambda}(u|B^\top X = z) = \frac{\sum_{i=1}^n K_b(Y_i - u) \delta_i K_h(B^\top X_i - z)}{\sum_{j=1}^n I(Y_j \geq u) K_h(B^\top X_j - z)}, \quad (2.5)$$

for some bandwidths  $b, h$ , and kernel function  $K_h(\cdot) = h^{-1}K(\cdot/h)$ . Details of these nonparametric components will be deferred to Section 3. It is worth noting that this nonparametric

component requires is only a  $d$  dimensional kernel, hence adaptive to the underlying structure.

It is then crucial to choose specific forms of  $\alpha(u, X)$ . Different choices may result in simplifications of the above formulation and/or gain additional theoretical and computational advantages. In the next two sections, we present four different choices, which fall into two categories: the forward and inverse regression schemes. The main differences between the two schemes are whether the counting process  $N(u)$  is used in the definition of  $\alpha(u, X)$ . The forward regression scheme is essentially the estimating equations approach in the normal regression, while the inverse regression scheme utilizes  $N(u)$  to mimic the sliced inverse regression (Li, 1991) conceptually.

## 2.2 Forward regression

In the forward regression scheme, we choose  $\alpha(u, X)$  such that it does not depend on the observed failure process  $N(u)$ . We first notice that, as long as  $\alpha(u, X)$  depends at most on the at-risk process  $Y(u)$ , we can simplify the estimating equations in (2.2) into

$$E\left(\int\left[\alpha(u, X) - E\{\alpha(u, X)|Y(u) = 1, B^T X\}\right]dN(u)\right) = 0. \quad (2.6)$$

We now give one example of  $\alpha(u, X)$  in the following when the structural dimension  $d = 1$ . This requires only a 1-dimensional nonparametric estimation.

**Example 2.1.** With  $\alpha(u, X) = X$ , the population version of the  $p$ -dimensional estimating equations is given by:

$$E\left(\int\left[X - E\{X|Y(u) = 1, B^T X\}\right]dN(u)\right) = 0. \quad (2.7)$$

This formulation reduces to the set of efficient estimating equations for the Cox proportional hazard model when the exponential link is known to be the underlying truth. It can also be used for the transformation models proposed by Zeng & Lin (2007). For some simple

extensions, we could let  $\alpha(u, X) = E\{XY(u)\}X^T$  to obtain  $p$ -by- $p$  dimensional estimating equations, which is suitable for the case of  $d > 1$ . To implement the forward regression method given by (2.7), noticing that  $dN_i(u)$  takes a jump at  $Y_i$  only if  $\delta_i = 1$ , we can estimate  $\psi_n(B)$  in (2.4) using

$$\hat{\psi}_n(B) = \frac{1}{n} \sum_{i=1}^n \{X_i - \hat{E}(X|Y \geq Y_i, B^T X_i)\} \delta_i, \quad (2.8)$$

where  $\hat{E}\{X|Y \geq u, B^T X = z\}$  takes the following form for any given  $u$  and  $z$ ,

$$\frac{\sum_{i=1}^n X_i I(Y_i \geq u) K_h(B^T X_i - z)}{\sum_{i=1}^n I(Y_i \geq u) K_h(B^T X_i - z)}, \quad (2.9)$$

for some choices of bandwidth  $h$  and kernel function  $K_h(\cdot)$ . Again, these details are deferred to Section 3.

### 2.3 Inverse regression

In this section, we focus on the inverse regression scheme. An important property that motivates the development is

$$\{dN(u) | Y(u) = 1, B^T X\} \sim \text{Bernoulli}\{\lambda(u|B^T X)du\}, \quad (2.10)$$

where  $dN(t) = N(t + dt) - N(t)$ . Hence, we can consider the sliced conditional mean of  $X$  given the outcome of  $dN(t)$  among the risk set, i.e.,  $Y(t) = 1$ . This leads to the construction of a local mean difference that is essentially the sliced mean difference for the binary outcome  $dN(u)$  (Cook & Lee, 1999):

$$\varphi(u) = E\{X | dN(u)=1, Y(u)=1\} - E\{X | dN(u)=0, Y(u)=1\}. \quad (2.11)$$

It should be noted that the outcome  $dN(u)$  conditioning on the event  $Y(u) = 1$  depends only on the failure model  $\lambda(u|B^T X)$  (Xia et al., 2010). Hence, by varying the argument  $u$ , the inverse regression curve  $\varphi(u)$  is contained within the central subspace  $\mathcal{S}_{T|X}$ . With this

$\varphi(u)$  established, we consider the function

$$\alpha(u, X) = X\varphi^T(u). \quad (2.12)$$

Then,

$$\alpha(u, X) - \alpha^*(u, B^T X) = [X - E\{X|Y(u) = 1, B^T X\}]\varphi^T(u). \quad (2.13)$$

This particular choice can be implemented by estimating  $E\{X|Y(u) = 1, B^T X\}$  using equation (2.9) and estimating  $\varphi(u)$  using the sliced average based on (2.11):

$$\hat{\varphi}(u) = \frac{\sum_{i=1}^n X_i I(u \leq Y_i < u + h, \delta_i = 1)}{\sum_{i=1}^n I(u \leq Y_i < u + h, \delta_i = 1)} - \frac{\sum_{i=1}^n X_i I(Y_i \geq u)}{\sum_{i=1}^n I(Y_i \geq u)}. \quad (2.14)$$

Based on this choice of  $\alpha$ , we propose two methods that utilize the estimating equations (2.2), and a computationally efficient method that further simplifies the formula to a singular value decomposition problem.

**Example 2.2.** Replacing  $\alpha(u, X) - \alpha^*(u, B^T X)$  in (2.2) by (2.13) leads to the semiparametric inverse regression approach in its population version:

$$E\left(\int [X - E\{X|Y(u) = 1, B^T X\}]\varphi^T(u)dM(u)\right). \quad (2.15)$$

This consists of  $p \times p$  estimating functions, and is able to handle  $d > 1$ . However, the nonparametric estimation part is only  $d$  dimensional as reflected by  $B^T X$ . Furthermore, this formulation enjoys the double robustness property which is illustrated in the Supplementary Material. Similar phenomenon has been observed by [Ma & Zhu \(2012\)](#) in the regression setting but without censoring. This suggests that if one of the terms  $E\{X|Y(u) = 1, B^T X\}$  and  $M(u)$  is estimated incorrectly, we can still obtain consistent estimations of the dimension reduction subspace. In our numerical experiment, we do observe numerical advantage of this approach over its simplified version, which is given in Example 2.3.



To implement this method, a vectorized  $\psi_n(B)$  is given by

$$\widehat{\psi}_n(B) = \text{vec} \left[ \frac{1}{n} \sum_{i=1}^n \sum_{\substack{j=1 \\ \delta_j=1}}^n \left\{ X_i - \widehat{E}(X|Y \geq Y_j, B^T X_i) \right\} \widehat{\varphi}^T(Y_j) \left\{ \delta_i I(j=i) - \widehat{\lambda}(Y_j|B^T X_i) \right\} \right], \quad (2.16)$$

where  $\widehat{E}\{X|Y \geq u, B^T X = z\}$  and  $\widehat{\varphi}^T(u)$  are given in (2.9) and (2.14), respectively, and the conditional hazard function can be estimated by (2.5). We finally apply the generalized method of moments (2.4) to estimate  $B$ .

**Example 2.3.** Similar to the forward regression example, our choice of  $\alpha(u, X)$  in (2.12) depends on at most the at-risk process  $Y(u)$ . Hence, the estimating functions in (2.15) can be simplified to the following counting process inverse regression approach:

$$E \left( \int [X - E\{X|Y(u) = 1, B^T X\}] \varphi^T(u) dN(u) \right). \quad (2.17)$$

Replacing  $dM(u)$  with  $dN(u)$  greatly reduces the computational burden. This can be seen from (2.16), where a conditional hazard function  $\widehat{\lambda}(Y_j|B^T X_i)$  needs to be evaluated at each observed failure time point  $j$  for all observations  $i$ . Of course, by doing this simplification, we lose the double robustness property. The implementation of this approach is a simplified version of (2.16) with:

$$\widehat{\psi}_n(B) = \text{vec} \left[ \frac{1}{n} \sum_{i=1}^n \left\{ X_i - \widehat{E}(X|Y \geq Y_i, B^T X_i) \right\} \delta_i \widehat{\varphi}^T(Y_i) \right], \quad (2.18)$$

where the estimations of nonparametric components are provided previously.

**Example 2.4.** With some additional assumptions,  $B$  can be estimated without any nonparametric smoothing. We make the following definitions:

**Definition 2.5.** For any  $\alpha \in \mathbb{R}^p$  and any  $u > 0$ , the linearity condition (Li, 1991) is satisfied further conditioning on the event  $\{Y(u) = 1\}$ , i.e.,

$$E\{\alpha^T X|Y(u) = 1, B^T X = z\} = c_0(u) + c^T(u)z, \quad (2.19)$$

where  $c_0(u)$  and  $c(u)$  are constants that possibly depend on  $u$ . Furthermore, the time-invariant covariance condition requires

$$\text{Cov}\{X|Y(u) = 1\} = c_1(u)\Sigma, \quad (2.20)$$

where  $c_1(u)$  is some constant that depends on  $u$ .

Noticing that after centering  $X$  at time point  $u$ , if the above two conditions are satisfied, we have

$$\begin{aligned} E(X|Y(u) = 1, B^T X) - E(X|Y(u) = 1) \\ = P\{X - E(X|Y(u) = 1)\}, \end{aligned}$$

where  $P = \Sigma B(B^T \Sigma B)^{-1} B^T$  and the constant term  $c_1(u)$  vanishes. Realizing that by the time-invariant covariance condition,  $P$  remains the same across all time points, plugging in the above equation into (2.17) leads to

$$Q E\left(\int [X - E\{X|Y(u) = 1\}] \varphi^T(u) dN(u)\right) = 0,$$

where  $Q = I - P$ . This is equivalent to deriving the left-singular space of the covariance matrix

$$E\left(\int [X - E\{X|Y(u) = 1\}] \varphi^T(u) dN(u)\right). \quad (2.21)$$

The computation of this approach is extremely simple. Realizing that  $dN(u)$  takes value 1 at at most one time point on the entire time domain, which corresponds to the failure subjects, the covariance form can be estimated by a sum of  $n$  terms. Then we perform singular value decomposition on this sample covariance matrix and obtain its leading left singular vectors, hence no optimization is required. Details are provided in Algorithm (1).

**Remark 2.6.** The two conditions imposed in this example are somehow restrictive and do not always hold. For example, since  $Y(u)$  is a process that depends on both the failure

and censoring distribution, as long as the censoring distribution depends on structures beyond  $B^T X$ , the conditions could be violated. However, many recent works of literature argue that the sliced inverse regression seems to still have satisfactory performances even when the linearity condition do not hold (Li & Dong, 2009; Dong & Li, 2010). Hence, this does not prevent the method from serving as a good explorative tool. The method is also practically very useful since it is served as the initial value when solving our other optimization approaches to speed up the computation.

### 3 Implementation and Algorithm

The implementation of the computationally efficient method given in (2.21) is straightforward since only sliced averaging and eigen-decomposition are required. Algorithm (1) summarizes the estimation procedure.

---

**Algorithm 1** Algorithm for the computationally efficient approach.

---

Input:  $\{(X_i, \delta_i, Y_i), 1 \leq i \leq n\}$ ,  $h > 0$ ,  $k > 0$ .

Step 1: For each  $Y_i$  such that  $\delta_i = 1$ , calculate  $\hat{\varphi}(Y_i)$  using Equation (2.14) and calculate

$$\hat{E}(X|Y > Y_i) \text{ using } \hat{E}(X|Y > u) = \{\sum_{i=1}^n I(Y_i > u)\}^{-1} \{\sum_{i=1}^n X_i I(Y_i > u)\}.$$

Step 2: Calculate  $\hat{M} = n^{-1} \sum_{\delta_i=1} \{X_i - \hat{E}(X|Y_i)\} \hat{\varphi}^T(Y_i)$ .

Step 3: Perform the singular value decomposition:  $\hat{M} = \hat{U} \hat{D} \hat{V}^T$ .

Output:  $\hat{B}$  as the first  $k$  columns of  $\hat{U}$ .

---

It requires numerical optimization to solve the estimating equations of the forward regression approach given in (2.7) and the two inverse regression approaches, given in (2.15) and (2.17), respectively. For all three approaches, we use the corresponding choice of the moment conditions and solve for the minimizer of  $\hat{\psi}_n(B)^T \hat{\psi}_n(B)$ , where  $\hat{\psi}_n(B)$  is specified in (2.8), (2.16) and (2.18) respectively. Existing methods use general-purpose optimization

tools such as the Newton–Raphson to solve for the minimizer, however, dimension reduction methods create an additional difficulty due to the identifiability issue, i.e.,  $B$  is not uniquely defined and the rank may not be preserved if we solve it freely within the space of  $\mathbb{R}^{p \times d}$ . To tackle this, [Ma & Zhu \(2012\)](#) propose to set the upper block (or a selected set of  $d$  rows) of  $B$  as a diagonal matrix and solve for the rest of parameters. However, this requires the pre-knowledge of the location of the important variables. Instead, we propose an orthogonality constrained optimization approach to solve our semiparametric dimension reduction model within the Stiefel manifold ([Edelman et al., 1998](#)):

$$\begin{aligned} & \text{minimize} \quad \widehat{\psi}_n(B)^\top \widehat{\psi}_n(B), \\ & \text{subject to} \quad B^\top B = I_{d \times d}. \end{aligned} \tag{3.1}$$

The advantage of this optimization approach is that we exactly preserves the rank  $d$  of the column space defined  $B$  while not pre-specify the restrictions on any of it entries. The main machinery of this algorithm is the optimization approached proposed by [Wen & Yin \(2013\)](#). The method is a first-order descent algorithm that preserves the update of the parameters within the manifold. In particular, let the gradient matrix be defined as

$$G = \frac{\partial \widehat{\psi}_n(B)^\top \widehat{\psi}_n(B)}{\partial B}. \tag{3.2}$$

Then, utilizing the Cayley transformation, we can update  $B$  to

$$B(\tau_0) = \left( I + \frac{\tau_0}{2} A \right)^{-1} \left( I - \frac{\tau_0}{2} A \right) B, \tag{3.3}$$

where  $A = GB^\top - BG^\top$  is a skew-symmetric matrix, and  $\tau_0$  is a step size. In practice,  $\tau_0$  can be chosen using inexact line search by incorporating the Wolfe conditions ([Nocedal & Wright, 2006](#)). It can be easily verified that if  $B^\top B = I$ , then  $B(\tau_0)^\top B(\tau_0) = I$  for any  $\tau_0 > 0$ , which preserves the constraint exactly. This approach is in-line with traditional dimensional reduction methods which recover the column space of  $B$  rather than treating each entry as a fixed parameter. Moreover, if an upper block diagonal version is desired, we can

easily convert the obtained solution through linear transformations. However, in this case, we can select the largest entries in the estimated  $\widehat{B}$  as the location of the diagonal matrix, instead of pre-specifying the locations. The full algorithm is presented in Algorithm 2. The iteration is stopped when a pre-specified optimization precision  $\varepsilon_0$  is reached. For estimating the nonparametric components (2.5) and (2.9), we exploit the Gaussian kernel and choose the optimal bandwidth  $h = (4/(d+2))^{1/(d+4)} n^{-1/(d+4)} \widehat{\sigma}$  (Silverman, 1986), where  $\widehat{\sigma}$  is the estimated standard deviation. For estimating the conditional hazard function, equation 2.5 bears much computational burden because it requires  $\mathcal{O}(n^2)$  flops to calculate the hazard at any given  $u$  and  $z$ . An alternative approach that greatly reduces the computational cost can be considered using the definition in Dabrowska et al. (1989), given by the following:

$$\widehat{\lambda}(u|B^T X = z) = \frac{\sum_{i=1}^n I(Y_i = u) I(\delta_i = 1) K_h(B^T X_i - z)}{\sum_{j=1}^n I(Y_j \geq u) K_h(B^T X_j - z)} \quad (3.4)$$

Since the indicator  $I(Y_i = u)$  can only take 1 if  $u$  is among the observed survival times. Hence, the numerator essentially requires only a single flop. Based on our experience, the numerical performance of the two versions are very similar. Hence, the above definition is implemented and used in the simulation study. Lastly, the implementation is available through the R package “orthoDr” (Zhao et al., 2017) through the Rcpp (Eddelbuettel & François, 2011) interface.

## 4 Asymptotic Normality

We focus on the semiparametric inverse regression approach, in which  $\widehat{B}$  obtained by solving

$$\frac{1}{n} \text{vec} \left[ \sum_{i=1}^n \int_0^\tau \left\{ \alpha(u, X_i) - \widehat{\alpha}^*(u, \widehat{B}^T X_i) \right\} d\widehat{M}(u, \widehat{B}^T X_i) \right] = 0,$$

To address the identifiability issue of  $B$ , we restrict our attention to the matrices in the form of  $B = (B_u^T, B_\ell^T)^T$ , where the upper sub-matrix  $B_u = I_d \in \mathbb{R}^{d \times d}$  is the  $d \times d$  identity matrix. In this manner, we can view  $B_\ell$  as the unique parameterizations of the subspace

---

**Algorithm 2** The orthogonality constrained optimization algorithm.

---

Input:  $\varepsilon_0, \{(X_i, \delta_i, Y_i), 1 \leq i \leq n\}$ .

Initialize: Obtain  $B^{(0)}$  from the computationally efficient approach in Algorithm 1.

For  $k = 1$  to  $k = \text{max.iter}$ :

Numerically approximate the gradient matrix  $G$  at  $B^{(k)}$ .

Compute the skew-symmetric matrix  $A = GB^T - BG^T$ .

Perform line search for  $\tau_0$  on the path  $B(\tau_0) = (I + \frac{\tau_0}{2}A)^{-1}(I - \frac{\tau_0}{2}A)B$ .

Update  $B^{(k+1)} = B(\tau_0)$ .

Stop if  $\|B^{(k+1)} - B^{(k)}\|_2 \leq \varepsilon_0$ .

Output:  $\hat{B} = B^{(k+1)}$ .

---

$\mathcal{S}(B)$ . We then write  $\beta_\ell = \text{vecl}(B) = \text{vec}(B_\ell)$ , the vector concatenating all free parameters in  $B$ . We need the the following regularity assumptions.

**Assumption 4.1.** There exists a  $\tau$ , such that  $0 < \tau < \infty$  and  $\text{pr}(Y > \tau|X) > 0$ .

**Assumption 4.2.** Let  $f_{B^T X}(z)$  be the density function of  $B^T X$  evaluated at  $z = B^T x$ ,  $f(t, z)$  be the density of  $T$  given  $B^T X = z$ ,  $S(t, x) = \text{pr}(T \geq t|X = x)$  and  $S_c(t, x) = \text{pr}(C \geq t|X = x)$ . Assume that  $f(t, z)$ ,  $f_{B^T X}(z)$ ,  $S(t, z)$  and  $E(S_c(t, X)|z)$  are bounded, and have bounded first and second derivatives with respect to  $t$  and  $z$ , and  $S(t, z)$  is bounded away from zero.

**Assumption 4.3.** The univariate kernel function  $K(x)$  is symmetric with  $\int x^2 K(x) dx < \infty$ . The  $d$ -dimensional kernel function is a product of  $d$  univariate kernel functions, that is  $K(u) = \prod K(u_j)$  for  $u = (u_1, \dots, u_d)^T$ .

Assumptions 4.1 and 4.2 are standard in survival analysis. Assumptions 4.3 is commonly used in kernel estimations. Base on these two assumptions, we can provide the rate for our conditional hazard function estimation and its derivatives. Its easy to see that the

Silverman formula implemented in our numerical approach automatically leads to consistent estimations.

**Lemma 4.4.** Under Assumption 4.3 and 4.2, and assume that the bandwidths satisfy  $h, b \rightarrow 0$ ,  $n b h^{d+2} \rightarrow \infty$ , we have, uniformly for all  $t$  and  $z$ ,

$$\begin{aligned}\widehat{\lambda}(t|z) &= \lambda(t, z) + O_p\left((n b h^d)^{-1/2} + h^2 + b^2\right), \text{ and} \\ \frac{\partial}{\partial z} \widehat{\lambda}(t|z) &= \frac{\partial}{\partial z} \lambda(t, z) + O_p\left((n b h^{d+2})^{-1/2} + h^2 + b^2\right).\end{aligned}$$

Before presenting our main theorem, we also need the convergence result of the  $\alpha^*$  functions. However, we do not want the theoretical result being limited to the choice given in equation (2.2). Instead, we provide general results for any valid choice of the  $\alpha^*$  function, as long as the following condition is satisfied.

**Assumption 4.5.** We assume that for some  $\kappa < 1/2$ , the convergence rate for the following conditional nonparametric estimation holds uniformly over all  $u$  and  $z$ ,

$$\begin{aligned}\text{vec}\left\{\widehat{\alpha}^*(u, z) - \alpha^*(u, z)\right\} &= O_p\left(n^{-1/2+\kappa}\right), \\ \frac{\partial}{\partial z} \text{vec}\left\{\widehat{\alpha}^*(u, z) - \alpha^*(u, z)\right\} &= O_p\left(n^{-1/2+\kappa}\right),\end{aligned}$$

Note that, for most valid choices such as a kernel estimation of the conditional density, when the number of dimension  $d$  is fixed, the rate provide in Lemma 4.4 is essentially valid for  $\widehat{\alpha}^*(u, z)$ , while for conditional expectation estimations, the classical rate of  $O_p((n h^d)^{-1/2} + h^2)$  can be obtained. Hence, with proper choice of the bandwidth, the rate of Assumption 4.5 can always be achieved. We now present the main theorem.

**Theorem 4.6** (Asymptotic Normality). Under Assumptions 4.1-4.5, and the choice of bandwidths specified in Lemma 1, the estimator  $\text{vecl}(\widehat{B})$  is asymptotically normal, that is

$$\sqrt{n} \left( \widehat{\beta}_\ell - \beta_\ell \right) \xrightarrow{d} \mathcal{N}(0, \Sigma),$$

where  $\Sigma = (G^T G)^{-1} G \Sigma_A G^T (G^T G)^{-1}$ ,

$$\Sigma_A = \text{cov}(A(\tau)) = \text{cov} \left( \int_0^\tau \text{vec} \{ \alpha(u, X) - \alpha^*(u, B^T X) \} dM(u, B^T X) \right),$$

$$\text{and } G = E \left( \frac{\partial}{\partial \beta_\ell} \int_0^\tau \text{vec} [ \{ \alpha(u, X_i) - \alpha^*(u, B^T X) \} ] dM(u, B^T X) \right).$$

## 5 Numerical Examples

### 5.1 Simulation Studies

In this section, we examine the finite sample performance of our proposed methods via extensive numerical experiments. Specifically, we carry out the estimation of dimension reduction subspace using the forward regression approach (2.7), the semiparametric inverse regression approach (2.15), the counting process inverse regression approach (2.17) and the computational efficient approach (2.21). All of our methods are implemented through the “orthoDr” package in R. Four alternative approaches are considered: a naive approach that performs sliced inverse regression on the failure observations, carried out using the “dr” package (Weisberg, 2002); the double slicing approach (Li et al., 1999) using R package “censorSIR” provided by Wu et al. (2008); the minimal average variance estimation based on hazard functions in Xia et al. (2010). This implementation is provided by the original author through MATLAB; and the inverse probability of censoring weighted approach based on Lu & Li (2011). We carry out this approach ourselves by using a Cox proportional hazard model to estimate the censoring weights and obtain the reduced space by utilizing the “dr” package with subject weights.

We consider four different settings: Setting 1 is a classical Cox proportional hazard model; Setting 2 is constructed with structural dimension  $d=2$  and directions in the hazard function are changing over time; Setting 3 also has structural dimension equal to two, with the two directions interacting with each other. Setting 4 also has two interacting structural



dimensions, while the failure and the censoring variables have overlap. For each setting, we consider  $p = 6, 12$  and  $18$ . Each experiment is repeated 200 times with sample size  $n = 400$ .

Setting 1: The true survival time  $T$  and the censoring time  $C$  are generated from exponential distributions with rate  $\exp(\beta^\top X)$  and  $\exp(X_4 + X_5 - 1)$  respectively, where  $\beta = (1, 0.5, 0, \dots, 0)^\top$  and  $X_j$  is the  $j$ -th element of  $X$ , for  $1 \leq j \leq p$ . The covariate  $X$  follows from multivariate normal distribution with mean 0 and covariance  $\Sigma = (0.5^{|i-j|})_{ij}$ . The overall censoring rate is around 35.3%.

Setting 2: We generate  $T_1$  and  $T_2$  from exponential distributions with rate  $\exp(\beta_1^\top X)$  and  $\exp(\beta_2^\top X)$  respectively, where  $\beta_1 = (1, 0, 1, 0, \dots, 0)^\top$  and  $\beta_2 = (0, 1, 0, 1, 0, \dots, 0)^\top$ . The true survival time  $T = T_1 I(T_1 < 0.4) + (T_2 + 0.4) I(T_1 \geq 0.4)$ . The censoring time  $C$  is generated from exponential distributions with rate  $\exp(X_5 - X_6 - 2)$ . The covariate  $X$  follows the same distribution as in Setting 1. The overall censoring rate is around 35.1%.

Setting 3: The true survival time  $T$  is generated from Weibull distribution with shape parameter 5 and scale parameter  $\exp(4\beta_2^\top X(\beta_1^\top X - 1))$ , where  $\beta_1 = (1, 0, 1, 0, \dots, 0)^\top$  and  $\beta_2 = (0, 1, 0, 1, 0, \dots, 0)^\top$ . The censoring time  $C$  is generated uniformly from 0 to  $3 \exp(X_5 - X_6 + 0.5)$ . We further draw  $X$  such that  $X_j$ 's follow standard uniform distribution  $U(0, 1)$  independently. The overall censoring rate is around 33.8%.

Setting 4: The true survival time  $T$  is generated from a Cox proportional hazard model with  $\log(T) = -2.5 + \beta_1^\top X + 0.5\beta_1^\top X\beta_2^\top X + 0.25 \log(-\log(1 - u))$  and  $\log(C) = -0.5 + \beta_3^\top X + \log(-\log(1 - u))$ , where  $u$ 's are i.i.d. uniform distributed,  $\beta_1 = (1, 1, 0, \dots, 0)^\top$ ,  $\beta_2 = (0, 0, 1, -1, 0, \dots, 0)^\top$ , and  $\beta_3 = (0, 1, 0, 1, 1, 1, 0, \dots, 0)^\top$ . The covariate  $X$  follows the same distribution as setting 1, except  $\Sigma = (0.25^{|i-j|})$ . The overall censoring rate is around 26.2%.

We first investigate the statistical performance using three different measures: the Frobenius norm distance between the projection matrix  $P$  and its estimator  $\hat{P}$ , where  $P = B(B^\top B)^{-1}B^\top$ ; the trace correlation  $\text{tr}(P\hat{P})/d$ , where  $d$  is the structural dimension;

and the canonical correlation between  $B^T X$  and  $\widehat{B}^T X$ . The results are summarized in Table 1.

Overall, the two inverse regression methods achieve the best performance, followed by the computationally efficient approach. It is worth to point out that the computationally efficient approach, while no non-parametric approximation is required, outperforms existing methods in almost all settings. Among all competing methods, double slicing performs the best in general, while Xia et al. (2010) and Lu & Li (2011) outperforms double slicing in Setting 3 and Setting 4, respectively. In terms of the three error measurements, we found that the Frobenius norm distance is the most informative measurement, while the Trace and Canonical correlations are less sensitive to the performances.

Among the two inverse regression methods, the semiparametric version is slightly better in Settings 3 and 4. The main advantage of the semiparametric version compared with the counting process version is the double robustness, which ensures consistency even when the conditional expectations are not estimated correctly. However, this theoretical advantage does not translate into strong numerical improvements in Settings 1 and 2 especially when  $p$  is large. This is possibly due to the variations in the hazard function estimation, which introduces less numerical stability. In setting 1, forward regression approach achieves the best performance. As discussed in Example 1, this method mimics the efficient estimating equations used in the Cox proportional hazard model and is thus the most efficient method in this setting. In setting 2, The computationally efficient approach performs similarly to the two inverse regression approaches and even outperforms them under large  $p$ . This shows some potential of this approach in higher dimension settings when nonparametric estimations may not be preferred.

One major challenge of solving estimating equations is the computational complexity, especially with nonparametric components. Our proposed method adds additional difficulties with the orthogonality constraints, i.e.,  $B^T B = I_{d \times d}$ . However, with our proposed

orthogonality constrained optimization algorithm, in combination with the Rcpp interface, our implementation can solve the proposed method very efficiently. In addition, parallel computing through OpenMP is utilized to numerically approximate the gradient for each entry of  $B$ . For example, in Setting 2 with  $p = 6$ , the mean computational time of the inverse regression counting process approach is 1.62 seconds, while the time for the semi-parametric version is 8.01 seconds. Table 3 in the Supplementary Material summarizes the computational cost across all settings. All simulations are done on an Intel Xeon E5-2680v4 processor with 5 parallel threads.

We further investigate the variance of the proposed methods. Due to the complicated form of the variance formula, we instead use the bootstrap to obtain an estimation of the standard deviation of the proposed estimators. Using an upper-block-diagonal version of the parameter of interest, we estimate the standard deviation of the parameters based on 100 bootstrap samples and also report the 95% confidence interval. The results show that in setting one, the bootstrap estimator of all the proposed methods approximates the standard deviation closely. In the rest settings, the approximation of the computational efficient approach and counting process inverse regression approach still archives good performance, while semiparametric inverse regression approach slightly over-estimate the standard deviations, leads to a slight over-coverage (around 98%). However, the proposed methods still archive smaller empirical standard deviation on nearly all parameters across all settings. Details are provided in the Supplementary Material.

## 5.2 Skin Cutaneous Melanoma Data Analysis

We apply the proposed method to The Cancer Genome Atlas (TCGA, <http://cancergenome.nih.gov/>) skin cutaneous melanoma dataset. TCGA provides the public with one of the most comprehensive profiling data on more than thirty cancer types. We acquire gene expression and clinical data on a total of 469 patients (156 observed failures) and their mRNA


expression data on 20,531 genes. To produce biologically meaningful results, we preselect 20 genes in this analysis, which are the top 20 genes highly associated with cutaneous melanoma based on meta-analyses of over 145 existing literature (Chatzinasiou et al., 2011). A list of these genes can be found at <http://bioserver-3.bioacademy.gr/Bioserver/MelGene/>. We further include age at diagnosis as a clinical control variable. All covariates are pre-processed to have unit variance and zero mean.

Selecting the number of structural dimensions can be a challenging task, especially with right censored survival model (Xia et al., 2010). To this end, we adopt the validated information criterion developed by Ma & Zhang (2015), which is particularly suited for our generalized method of moments framework. The validated information criterion is constructed by penalizing the quadratic form of the objective function. Interestingly when we apply this method to all of our proposed estimating equation approaches,  $d = 1$  always yields the best fit. Hence we present the results for all method under  $d = 1$ . As a demonstration of the fitted model, we project the design matrix on the estimated direction of the semiparametric inverse regression approach and plot the survival outcome against the projection (Figure 1). A nonparametric estimation of the conditional survival function based on this projection is also produced. From these two figures, we can see a clear trend that subjects with larger values of the projection have lower survival rate. As for comparisons, we perform the competing methods with one structural dimension, and the results can be found in the Supplementary Material (Figure 6). From this simple visualization, it seems that the double slicing method obtains a similar direction with monotone effects on the risk of failure, while the other directions obtained by other methods are non-monotone.

We also observe both similarities and differences among different methods for the identified genes. A pairwise Frobenius norm distance is provided in the Supplementary Material (Table 8). This suggests that the proposed methods have fairly small distances while existing methods mostly do not agree with each other. In addition, double slicing has the

smallest distance with the proposed methods. The proposed methods share some consistent trend of the loadings on the most influential variables. For example, all the proposed methods identify Age as the most important variable with loadings over 0.5. Corresponding this with the survival curve plot in Figure 1, it suggests that higher survival rate is observed for smaller derived direction. This further indicates that patients with younger age tend to have higher survival, which is biologically intuitive. This finding is also consistent with the double slicing method, which identifies Age with loading 0.47. However, other methods do not assign large loading to Age. Here, we present all results with a positive loading of age, and the directions are multiplied by  $-1$  otherwise. Another important variable that all methods agree in terms of signs is MTAP. This gene has been previously reported to have a negative correlation with the progression of melanocytic tumors [Behrmann et al. \(2003\)](#), which justifies the large negative value estimated by the proposed methods. However, the magnitudes in alternative methods are small. Other common genes identified by the proposed methods are MYH7B and CASP8. We mine the literature and found that [Li et al. \(2008\)](#) genotyped putatively functional polymorphisms of CASP8 and found a significant association with lower risk of cutaneous melanoma. The result therein supports the large negative loading of CASP8 gene in our fitted model. For differences across methods, TYR is identified by alternative methods except the double slicing, with extremely large loadings (up to  $-0.78$ ). The enzyme encoded by this gene controls the production of melanin, hence it has been shown to be strongly associated with melanoma ([Gudbjartsson et al., 2008](#)). Although the estimated directions are dominated by this gene, we did not observe a monotone effect of the directions (See Figure 6 in the Supplementary Material).

Figure 1: Fitted direction and survival function of the semiparametric inverse regression



The left figure is the projected direction versus the observed failure (blue dot) and censoring (orange +) times. The right figure is a nonparametric estimation of the survival function based on the projected direction.

## 6 Discussion

In this paper, we proposed a counting process based dimension reduction framework for censored outcomes. A family of generalized method of moments based approaches has been constructed for estimating the dimension reduction subspace. The main advantage of the proposed method is that it requires only a  $d$ -dimensional (instead of  $p$ ) nonparametric kernel estimation while no censoring distribution is modeled. The reduced dimension of the nonparametric estimation circumvents the difficulties of many existing methods and improves the efficiency when the total dimension  $p$  is too large for kernel methods. Our simulation study suggests that the proposed method outperforms existing methods in a variety of settings. To efficiently solve the proposed estimating equations, we further introduce an orthogonality constrained optimization method that solves the parameters within a Stiefel manifold. With implementations in the R package “orthoDr” through C++, the counting process version of the estimators can be solved within a few seconds. However, the martingale version requires significant more calculations due to the local estimation the hazard function, hence requiring a few minutes to solve. We believe that there is still potential

room to improve the computational performance. In addition, our computational efficient approach requires only a singular value decomposition and has satisfactory performances. However, it does not enjoy the same theoretical guarantee without restraint conditions on the covariates. Further relaxation of these conditions is of great interest.

Our framework can be possibly extended to more general settings. First, by imposing penalization on the estimating equations, it is possible to extend the proposed method to moderately high dimensional data. Sparse estimation of the  $B$  parameter may help both interpretations and improves the prediction accuracy of subsequent nonparametric models. The second direction is to search for an alternative construction of the  $\alpha$  functions. Throughout our developments, we used the  $\phi(u)$  function which is motivated by the inverse regression of a Bernoulli distribution. It would be interesting to investigate the possibilities of a “SAVE” type of  $\alpha$  function that may detect more complicated model structure. We can also consider using  $\alpha(u, X) = B^T X \varphi^T(u)$ , which is another valid choice. Lastly, it is also interesting to extend this framework to a time-varying coefficient setting, where we may let the dimension reduction space  $\mathcal{S}$  to change over time  $t$ .

## 7 Supplementary Material

Supplementary material available at Biometrika online includes a derivation of the orthocomplement of the nuisance tangent space; the proof of the double robustness property for the semiparametric inverse regression approach; the proof of Lemma 4.4 and Theorem 4.6; and additional simulation and data analysis results.

## 8 Acknowledgement

The first two authors contribute equally. This work is partially supported by the NIH and the NSERC.

# Supplementary Material

## A Tangent Space Derivation

Before we give the tangent space, we first derive the nuisance tangent spaces  $\mathcal{E}_1, \mathcal{E}_2$  and  $\mathcal{E}_3$  in the following proposition. The proof follows the similar argument for proving the nuisance tangent spaces of cox regression model (Tsiatis, 2007) and is thus omitted for simplicity.

**Proposition A.1.** The nuisance tangent spaces  $\mathcal{E}_1, \mathcal{E}_2$  and  $\mathcal{E}_3$  have the following forms

$$\begin{aligned}\mathcal{E}_1 &= \left\{ \int \alpha(u, B^T X) dM(u, B^T X) : \alpha(u, B^T X) \text{ is measurable} \right\}, \\ \mathcal{E}_2 &= \left\{ \int \alpha(u, X) dM_C(u, X) : \alpha(u, X) \text{ is measurable} \right\}, \text{ and} \\ \mathcal{E}_3 &= \left\{ \alpha(X) : E\{\alpha(X)\} = 0 \right\},\end{aligned}$$

Next, we give a proof of tangent space  $\mathcal{E}^\perp$ , which is defined as

$$\mathcal{E}^\perp = \left\{ \int \{\alpha(u, X) - \alpha^*(u, B^T X)\} dM(u, X) : \alpha(u, X) \text{ is measurable} \right\}.$$

where

$$\alpha^*(u, B^T X) = E\{\alpha(u, X) | \mathcal{F}_u, B^T X\}.$$

*Proof of Proposition A.1.* For a fully nonparametric model, the nuisance tangent space is the whole Hilbert space  $\mathcal{H}$  with each element having mean zero. Therefore, if we put no restriction on the hazard function  $\lambda(t|X)$  and write the associated nuisance tangent space as  $\mathcal{E}_1^*$ , we obtain

$$\begin{aligned}\mathcal{H} &= \mathcal{E}_1^* \oplus \mathcal{E}_2 \oplus \mathcal{E}_3, \text{ where} \\ \mathcal{E}_1^* &= \left\{ \int \alpha(u, X) dM(u, X) : \alpha(u, X) \text{ is measurable} \right\}.\end{aligned}$$



The orthogonal completion of  $\mathcal{E}$  satisfies that  $\mathcal{E}^\perp \subset \mathcal{E}_1^*$  and  $\mathcal{E}^\perp \perp \mathcal{E}_1$ . In order to identify  $\mathcal{E}^\perp$ , it suffices to take an arbitrary element in  $\mathcal{E}_1^*$  and find its residual after projecting it onto  $\mathcal{E}_1$ . To find the projection, we must derive  $\alpha^*(u, B^T X) \in \mathcal{E}_1$  such that

$$E\left(\int (\alpha(u, X) - \alpha^*(u, B^T X))^T dM(u, X) \int a(u, B^T X) dM(u, X)\right) = 0.$$

The covariance of martingale stochastic integrals above can be computed by finding the expectation of the predictable covariance process ([Fleming & Harrington, 2011](#)):

$$\begin{aligned} & E\left(\int \{\alpha(u, X) - \alpha^*(u, B^T X)\}^T dM(u, X) \int a(u, B^T X) dM(u, X)\right) \\ &= E\left(\int \left[E\{\alpha(u, X) | \mathcal{F}_u, B^T X\} - \alpha^*(u, B^T X)\right]^T a(u, B^T X) \lambda(u | B^T X) Y(u) du\right) = 0, \end{aligned}$$

where  $a(u, B^T X)$  is arbitrary and  $\mathcal{F}_u$  is the filtration. Thus we must have

$$\alpha^*(u, B^T X) = E\{\alpha(u, X) | \mathcal{F}_u, B^T X\}.$$

This completes the proof. □

## B Proof of the Double Robustness Property for the semi-parametric inverse regression approach

Recall that, for the semiparametric inverse regression approach, we solve the sample version of the following estimating functions

$$E\left[\int \left\{E(X|Y(u)) - E(X|Y(u), B^T X)\right\} \varphi^T(u) dM(u)\right] = 0.$$

For simplicity, we will use the random function to denote

$$F(X, u) = \left\{E(X|Y(u)) - E(X|Y(u), B^T X)\right\} \varphi^T(u).$$

Case 1: Suppose  $M(u)$  is misspecified as  $M^*(u)$ . Then we have that

$$\begin{aligned}
E\{F(X, u)dM^*(u)\} &= E[E\{F(X, u)dM^*(u)|Y(u), X\}] \\
&= E[F(X, u)E\{dM^*(u)|Y(u), X\}] \\
&= E[E\{F(X, u)|Y(u), B^T X\}E\{dM^*(u)|Y(u), B^T X\}] \\
&= 0,
\end{aligned}$$

where the last equation is due to the fact that  $E\{F(X, u)|Y(u), B^T X\} = 0$ . Hence we have

$$E\left[\int F(X, u)dM^*(u)\right] = 0.$$

Case 2: Suppose the function  $F(X, u)$  is misspecified to  $F^*(X, u)$ . With a similar argument, we can show that  $E\{\int F^*(X, u)dM(u)\} = 0$  due to the fact that  $E\{dM(u)|Y(u), B^T X\} = 0$ . This completes the proof.

## C Proofs of asymptotic results

### C.1 Proof of Lemma 4.4

*Proof of Lemma 4.4.* Without loss of generality, we prove the results for  $d = 1$ . We start with the convergence rate of  $\hat{\lambda}(t|B^T X = B^T x)$ . Let  $f_{B^T X}(z)$  be the true density function of  $B^T X$  evaluated at  $z = B^T x$ , and

$$A_i = \frac{1}{n} \sum_{j=1}^n I(Y_j \geq u) K_h(B^T X_j - z) - f_{B^T X}(z) E\{I(Y \geq u)|z\}.$$

Suppose we have proved that  $A_i = O_p((nh)^{-1/2} + h^2)$  uniformly. Let  $z_i = B^T x_i$ . We have

$$\begin{aligned}
\hat{\Lambda}(t|z) &= \sum_{i=1}^n \frac{K_b(Y_i - t) \delta_i K_h(B^T X_i - z)}{\sum_{j=1}^n I(Y_j \geq t) K_h(B^T X_j - z)} \\
&= \underbrace{\frac{1}{n} \sum_{i=1}^n \frac{K_b(Y_i - t) \delta_i K_h(B^T X_i - z)}{f_{B^T X}(z) E\{I(Y \geq t)|z\}}}_{\text{I}} (1 + O(A_i)).
\end{aligned}$$

We bound the term (I) first. Let  $S(t, x) = \text{pr}(T \geq t | X = x)$  and  $S_c(t, x) = \text{pr}(C \geq t | X = x)$ .

Then the expectation of (I) can be written as

$$\begin{aligned}
& E \left[ \frac{K_b(Y_i - t) \delta_i K_h(B^T X_i - z)}{f_{B^T X}(z) S(t, z) E\{S_c(t, X_i) | z\}} \right] \\
&= \iint \frac{K_b(y_i - t) K_h(z_i - z) E\{S_c(y_i, X_i | B^T x_i)\}}{f_{B^T X}(B^T x) S(t, B^T x) E\{S_c(t, X_i) | B^T x\}} f(y_i, B^T x_i) f_{B^T X}(B^T x_i) dy_i dz_i \\
&= \iint \frac{K(v) K(u) E\{S_c(t + bv, X_i | z + hu)\}}{f_{B^T X}(z) S(t, z) E\{S_c(t, X_i) | z\}} f(t + bv, z + hu) f_{B^T X}(z + hu) dv du \\
&= \iint K(v) K(u) \lambda(t, z) dv du \\
&\quad + \frac{h^2 \partial^2}{2 \partial z^2} \iint \frac{K(v) K(u) E\{S_c(t, X_i | z^*)\}}{f_{B^T X}(z) S(t, z) E\{S_c(t, X_i) | z\}} f(t, z^*) f_{B^T X}(z^*) u^2 dv du \\
&\quad + \frac{b^2 \partial^2}{2 \partial t^2} \iint \frac{K(v) K(u) E\{S_c(t^*, X_i | z)\}}{f_{B^T X}(z) S(t, z) E\{S_c(t, X_i) | z\}} f(t^*, z) f_{B^T X}(z) v^2 dv du \\
&= \lambda(t, z) + \frac{h^2}{2} \sup \left\{ \frac{\frac{\partial^2}{\partial z^2} (f(t, z^*) f_{B^T X}(z^*) E\{S_c(y_i, X_i | z^*)\})}{f_{B^T X}(z) S(t, z) E\{S_c(t, X_i) | z\}} \right\} \int K(u) u^2 du \\
&\quad + \frac{b^2}{2} \sup \left\{ \frac{\frac{\partial^2}{\partial b^2} (f(t^*, z) E\{S_c(t^*, X_i | z)\})}{S(t, z) E\{S_c(t, X_i) | z\}} \right\} \int K(v) v^2 dv \\
&= \lambda(t, z) + O(h^2 + b^2),
\end{aligned}$$

where we set  $z_i = z + hu$  and  $y_i = t + bv$  in the second equality,  $z^*$  and  $t^*$  are some convex combinations of  $z$  and  $z + hu$ ,  $t$  and  $t + bv$  respectively. The last equality is a consequence of the assumed assumptions.

Next, we bound the variance of  $\hat{\lambda}(t, z)$ :

$$\begin{aligned}
\text{var} \left\{ \hat{\lambda}(t, z) \right\} &= \text{var} \left\{ \frac{1}{n} \sum_{i=1}^n \frac{K_b(Y_i - t) \delta_i K_h(B^T X_i - z)}{f_{B^T X}(z) E\{I(Y \geq t) | z\}} (1 + O(A_i)) \right\} \\
&\leq 2 \text{var} \left\{ \frac{1}{n} \sum_{i=1}^n \frac{K_b(Y_i - t) \delta_i K_h(B^T X_i - z)}{f_{B^T X}(z) E\{I(Y \geq t) | z\}} \right\} \\
&\quad + 2 \text{var} \left\{ \frac{1}{n} \sum_{i=1}^n \frac{K_b(Y_i - t) \delta_i K_h(B^T X_i - z)}{f_{B^T X}(z) E\{I(Y \geq t) | z\}} O(A_i) \right\}.
\end{aligned}$$

We upper bound the first quantity first.

$$\begin{aligned}
& 2 \operatorname{var} \left\{ \frac{1}{n} \sum_{i=1}^n \frac{K_b(Y_i - t) \delta_i K_h(B^\top X_i - z)}{f_{B^\top X}(z) E\{I(Y \geq t)|z\}} \right\} \\
&= \frac{2}{n} \left( E \left\{ \frac{K_b(Y_i - t) \delta_i K_h(B^\top X_i - z)}{f_{B^\top X}(z) E\{I(Y \geq t)|z\}} \right\}^2 - \lambda^2(t, z) \right) + O(h^2/n + b^2/n) \\
&= \frac{2}{n} E \left\{ \frac{K_b(Y_i - t) \delta_i K_h(B^\top X_i - z)}{f_{B^\top X}(z) E\{I(Y \geq t)|z\}} \right\}^2 + O(1/n) \\
&= \frac{1}{nhb} \iint \frac{K^2(v) K^2(u) E\{S_c(t + bv, X_i|z + hu)\}}{f_{B^\top X}^2(z) S^2(t, z) E^2\{S_c(t, X_i)|z\}} f(t + bv, z + hu) f_{B^\top X}(z + hu) dv du + O(1/n) \\
&= \frac{1}{nhb} \iint \frac{K^2(v) K^2(u)}{f_{B^\top X}(z) S^2(t, z) E\{S_c(t, X_i)|z\}} f(t, z) dv du \\
&\quad + \frac{h \partial^2}{nb \partial z^2} \iint \frac{K^2(v) K^2(u) E\{S_c(t, X_i|z^*)\}}{f_{B^\top X}^2(z) S^2(t, z) E^2\{S_c(t, X_i)|z\}} f(t, z^*) f_{B^\top X}(z^*) u^2 dv du \\
&\quad + \frac{b \partial^2}{nh \partial t^2} \iint \frac{K^2(v) K^2(u) E\{S_c(t^*, X_i|z)\}}{f_{B^\top X}^2(z) S^2(t, z) E^2\{S_c(t, X_i)|z\}} f(t^*, z) f_{B^\top X}(z) u^2 dv du + O(1/n) \\
&= O(1/(nhb)),
\end{aligned}$$

uniformly over all  $t$  and  $z$ . Following the similar argument, we can show that

$$\begin{aligned}
& 2 \operatorname{var} \left\{ \frac{1}{n} \sum_{i=1}^n \frac{K_b(Y_i - t) \delta_i K_h(B^\top X_i - z)}{f_{B^\top X}(z) E\{I(Y \geq t)|z\}} O(A_i) \right\} \\
&= O \left( \frac{1}{nhb} O^2(A_i) \right) = O((nhb)^{-2}b + (nhb)^{-1}h^4).
\end{aligned}$$

In the last equality, we use the fact that  $A_i^2 = O_p(1/(nh) + h^4)$  uniformly, which can be proved by following a simplified argument in [Mack & Silverman \(1982\)](#). Combining the above results together, we obtain

$$|\widehat{\lambda}(t, z) - \lambda(t, z)| = O_p((nhb)^{-1/2} + h^2 + b^2),$$

uniformly for all  $t$  and  $z$ . Using a similar argument, after a some long algebra, we can prove the second and the third statements. We do not repeat the details here.

□

## C.2 Proof of Theorem 4.6

Recall that  $\beta_\ell = \text{vecl}(B)$  and let  $\beta = \text{vec}(B)$ . Let  $\tilde{\beta}_\ell = \text{vecl}(\tilde{B})$  and  $\Omega = \{\tilde{B} : \|\tilde{\beta}_\ell - \beta_\ell\|_2 \leq Cn^{-1/2}\}$ , or equivalently  $\Omega = \{\tilde{\beta} : \|\tilde{\beta}_\ell - \beta_\ell\|_2 \leq Cn^{-1/2}\}$  for some  $C > 0$ . We are ready to prove the main theorem of this paper.

*Proof of Theorem 4.6.* Without loss of generality, we prove the theorem under the generic case in which  $\alpha^*(u, B^\top X)$  is arbitrary. For simplicity, let

$$S_n(\hat{B}) = \frac{1}{n} \text{vec} \left[ \sum_{i=1}^n \int_0^\tau \left\{ \alpha(u, X_i) - \hat{\alpha}^*(u, \hat{B}^\top X_i) \right\} d\widehat{M}(u, \hat{B}^\top X_i) \right].$$

We sometimes write  $M(u, B^\top X_i)$  as  $M_i(u)$  and  $S_n(B)$  as  $S_n(\beta_\ell)$ . Following Jureckova (1971) and Tsiatis (1990), it suffices to show that  $S_n(\beta_\ell)$  is uniformly asymptotically linear in a small neighborhood of the true parameter  $\beta_\ell$ . That is, we need to show, there exists some linear operator,  $G_n$ , such that

$$\sup_{\|\hat{\beta}_\ell - \beta_\ell\|_2 \leq Cn^{-1/2}} n^{1/2} \|S_n(\hat{\beta}_\ell) - S_n^*(\beta_\ell) - G_n(\hat{\beta}_\ell - \beta_\ell)\|_2 = o_p(1), \text{ where}$$

$$S_n^*(\beta_\ell) = S_n^*(B) = \frac{1}{n} \text{vec} \left[ \sum_{i=1}^n \int_0^\tau \left\{ \alpha(u, X_i) - \alpha^*(u, B^\top X_i) \right\} dM(u, B^\top X_i) \right].$$

We then expand  $S_n(\hat{\beta}_\ell)$  as follows,

$$\begin{aligned} 0 = S_n(\hat{\beta}_\ell) &= \frac{1}{n} \text{vec} \left[ \sum_{i=1}^n \int_0^\tau \left\{ \alpha(u, X_i) - \hat{\alpha}^*(u, \hat{B}^\top X_i) \right\} d\widehat{M}(u, \hat{B}^\top X_i) \right] \\ &= \underbrace{\frac{1}{n} \text{vec} \left[ \sum_{i=1}^n \int_0^\tau \left\{ \alpha(u, X_i) - \hat{\alpha}^*(u, B^\top X_i) \right\} d\widehat{M}(u, B^\top X_i) \right]}_{\text{I}} \\ &\quad + \underbrace{\frac{1}{n} \sum_{i=1}^n \frac{\partial}{\partial z} \text{vec} \left[ \left\{ \alpha(u, X_i) - \hat{\alpha}^*(u, B^\top X_i) \right\} d\widehat{M}(u, B^\top X_i) \right] (\hat{B} - B)^\top X_i}_{\text{II}} + o_p \left( n^{-\frac{1}{2}} \right), \end{aligned}$$

where  $z$  is a vectorized argument of  $B^\top X$ . It remains to analyze terms (I) and (II) respec-

tively. For term (I), we obtain that

$$\begin{aligned}
\text{I} &= \frac{1}{n} \text{vec} \left[ \sum_{i=1}^n \int_0^\tau \left\{ \alpha(u, X_i) - \alpha^*(u, B^\top X_i) \right\} dM(u, B^\top X_i) \right] \\
&\quad + \underbrace{\frac{1}{n} \text{vec} \left[ \sum_{i=1}^n \int_0^\tau \left\{ \alpha(u, X_i) - \alpha^*(u, B^\top X_i) \right\} d \left( \widehat{M}(u, B^\top X_i) - M(u, B^\top X_i) \right) \right]}_{\text{I}_1} \\
&\quad + \underbrace{\frac{1}{n} \text{vec} \left[ \sum_{i=1}^n \int_0^\tau \left\{ \alpha^*(u, B^\top X_i) - \widehat{\alpha}^*(u, B^\top X_i) \right\} d \left\{ \widehat{M}(u, B^\top X_i) - M(u, B^\top X_i) \right\} \right]}_{\text{I}_2} \\
&\quad + \underbrace{\frac{1}{n} \text{vec} \left[ \sum_{i=1}^n \int_0^\tau \left\{ \alpha^*(u, B^\top X_i) - \widehat{\alpha}^*(u, B^\top X_i) \right\} dM(u, B^\top X_i) \right]}_{\text{I}_3}.
\end{aligned}$$

We bound terms  $\text{I}_1$ ,  $\text{I}_2$  and  $\text{I}_3$ , respectively. For  $\text{I}_1$ , we can applied Lemma 4.4. Note that for certain choice of the bandwidth  $b$  and  $h$ , we can always find some  $\kappa < 1/2$  such that  $\widehat{\lambda}(t|z) = \lambda(t, z) + O_p(n^{-1/2+\kappa})$ . Hence, for simplification, we use  $\kappa$  throughout the rest of this proof.

$$\begin{aligned}
\text{I}_1 &= \frac{1}{n} \text{vec} \left[ \sum_{i=1}^n \int_0^\tau \left\{ \alpha(u, X_i) - \alpha^*(u, B^\top X_i) \right\} \left\{ \lambda(u, B^\top X_i) - \widehat{\lambda}(u, B^\top X_i) \right\} Y_i(u) du \right] \\
&= O_p(n^{-1+\kappa}) O_p \left( n^{-\frac{1}{2}} \text{vec} \left[ \sum_{i=1}^n \int_0^\tau \left\{ \alpha(u, X_i) - \alpha^*(u, B^\top X_i) \right\} \lambda(u, B^\top X_i) Y_i(u) du \right] \right) \\
&= O_p(n^{-1+\kappa}) O_p(1) = O_p(n^{-1+\kappa}),
\end{aligned}$$

where the last inequality is due to central limit theorem. Following the some algebra and using Assumption 4.5, we can show that  $\text{I}_2 = O_p(n^{-1+\kappa})$  and  $\text{I}_3 = O_p(n^{-1+\kappa})$ . Combining the bounds for the above three terms above, we obtain that

$$\begin{aligned}
\text{I} &= \frac{1}{n} \text{vec} \left[ \sum_{i=1}^n \int_0^\tau \left\{ \alpha(u, X_i) - \alpha^*(u, B^\top X_i) \right\} dM(u, B^\top X_i) \right] + O_p(n^{-1+\kappa}) \\
&= \frac{1}{n} \text{vec} \left[ \sum_{i=1}^n \int_0^\tau \left\{ \alpha(u, X_i) - \alpha^*(u, B^\top X_i) \right\} dM(u, B^\top X_i) \right] + o_p(n^{-1/2}),
\end{aligned}$$

where we use the fact that  $\kappa < 1/2$  with proper choice of  $b$  and  $h$ . For term II, in a similar

argument, we shall obtain that

$$\begin{aligned}\text{II} &= \frac{1}{n} \sum_{i=1}^n \frac{\partial}{\partial z} \left( \int_0^\tau \text{vec}[\{\alpha(u, X_i) - \alpha^*(u, B^\top X_i)\}] dM(u, B^\top X_i) \right) (\widehat{B} - B)^\top X_i + o_p(n^{-1/2}) \\ &= \frac{1}{n} \sum_{i=1}^n \frac{\partial}{\partial \beta_\ell} \left( \int_0^\tau \text{vec}[\{\alpha(u, X_i) - \alpha^*(u, B^\top X_i)\}] dM(u, B^\top X_i) \right) (\widehat{\beta}_\ell - \beta_\ell) + o_p(n^{-1/2}),\end{aligned}$$

where we rewrite the first term in the right hand side as the derivative with respect to  $\beta_\ell$ .

For simplicity, define

$$\begin{aligned}A_i(\tau) &= \int_0^\tau \text{vec} \{ \alpha(u, X_i) - \alpha^*(u, B^\top X_i) \} dM(u, B^\top X_i), \\ \text{and } G_n &= \frac{1}{n} \sum_{i=1}^n \frac{\partial}{\partial \beta_\ell} \left( \int_0^\tau \text{vec}[\{\alpha(u, X_i) - \alpha^*(u, B^\top X_i)\}] dM(u, B^\top X_i) \right).\end{aligned}$$

Combing the asymptotically linear expansion with terms I and II,  $S_n(\widehat{B})$  can be written as

$$\widehat{S}_n(\widehat{B}) = \frac{1}{n} \sum_{i=1}^n A_i(\tau) + G_n(\beta_\ell - \widehat{\beta}_\ell) + o_p(n^{-1/2}) = 0.$$

Therefore,  $\widehat{\beta}_\ell - \beta_\ell$  can be written as

$$\sqrt{n}(\widehat{\beta}_\ell - \beta_\ell) = (G_n^\top G_n)^{-1} G_n^\top \left\{ \frac{1}{\sqrt{n}} \sum_{i=1}^n A_i(\tau) \right\} + o_p(1),$$

where we implicitly assume that  $G_n^\top G_n$  is invertible, which requires that at the number of estimating equations is larger than the number of parameters. In other words, this assume that the estimating equations are rich enough to recover the  $d(p-d)$ -dimensional vector of parameters  $\beta_\ell$ . This finishes the proof.  $\square$

## D Computational Time

This computational time for all proposed method is presented in Table 3. All simulations are done on an Intel Xeon E5-2680v4 processor with 5 parallel threads.

## E Additional Simulation Results

This section includes parameter estimations and standard deviation results in the simulation. For each estimated parameter matrix  $\widehat{B}$ , we linearly transform the columns of this

matrix such that the block sub-matrix is an identity matrix. For example, in settings 2 and 3, we require the first two rows to be a block diagonal matrix, i.e., the transformed parameter matrix is obtained through  $\widehat{B}\widehat{B}_{[1:2, 1:2]}^{-1}$ , where  $B_{[1:2, 1:2]}$  is the upper-block (first two rows) of  $\widehat{B}$ . In setting 1, a one dimensional case, only the first row is used, hence is essentially  $\widehat{B}\widehat{B}_{[1, 1]}^{-1}$ . In setting 4, the first and the third rows are used. Parameter estimations of other rows are reported in the following tables. To estimate the standard deviation, we use 100 bootstrapped samples of the training data to estimate the parameters, then translate each parameter estimate into the form with a diagonal matrix at the specified rows. Then for each parameter in the rest of matrix, since the parameters asymptotically follow a joint Gaussian distribution, we use 1.4826 times the median absolute deviation of the 100 bootstrap replicates as an estimation of the standard error, following the fact that this is a consistent estimator of the standard deviation in the normal distribution. The reason that we use the median absolute deviation instead of a regular standard deviation estimator is to ensure robustness such that extreme values (possibly due to convergence issue caused by duplicated samples) in the bootstrap estimations do not dominate the result.

## F Additional results of TCGA data analysis


This section contains additional results of the TCGA skin cutaneous melanoma data analysis. Table 8 is the pairwise distance measure of the first direction estimated by all methods. This shows that the proposed methods mostly agree with each other. Figure 6 presents the fitted models of competing approaches.

## References

BEHRMANN, I., WALLNER, S., KOMYOD, W., HEINRICH, P. C., SCHUIERER, M., BUETNER, R. & BOSSERHOFF, A.-K. (2003). Characterization of methylthioadenosin phos-




Figure 2: Boxplot of parameter estimates (Setting 1,  $p = 6$ )




each paramter, the boxes are (from left to right, with proposed methods colored): DS (Li et al., 1999); IPCW-SIR (Lu & Li, 2011); hMave (Xia et al., 2010); The forward regression; The computational efficient approach; The counting process inverse regression; The semiparametric inverse regression.

Figure 3: Boxplot of parameter estimates (Setting 2,  $p = 6$ )




For each paramter, the boxes are (from left to right, with proposed methods colored): DS (Li et al., 1999); IPCW-SIR (Lu & Li, 2011); hMave (Xia et al., 2010); The forward regression; The computational efficient approach; The counting process inverse regression; The semiparametric inverse regression. Proposed methods are colored.

Figure 4: Boxplot of parameter estimates (Setting 3,  $p = 6$ )




For each paramter, the boxes are (from left to right, with proposed methods colored): DS (Li et al., 1999); IPCW-SIR (Lu & Li, 2011); hMave (Xia et al., 2010); The forward regression; The computational efficient approach; The counting process inverse regression; The semiparametric inverse regression. Proposed methods are colored.

Figure 5: Boxplot of parameter estimates (Setting 4,  $p = 6$ )



The true parameter is For each paramter, the boxes are (from left to right, with proposed methods colored): DS (Li et al., 1999); IPCW-SIR (Lu & Li, 2011); hMave (Xia et al., 2010); The forward regression; The computational efficient approach; The counting process inverse regression; The semiparametric inverse regression. Proposed methods are colored.

Figure 6: Estimated directions and fitted survival functions of alternative methods



First row: Li et al. (1999); Second row: Lu & Li (2011); Third row: Xia et al. (2010). For each row, the left panel is the projected direction versus the observed failure (blue dot) and censoring (orange +) times. The right panel is a nonparametric estimation of the survival function based on the projected direction.

- phorylase (mtap) expression in malignant melanoma. *The American journal of pathology* **163**, 683–690.
- BICKEL, P. J., KLAASSEN, C. A., RITOV, Y. & WELLNER, J. A. (1993). *Efficient and Adaptive Estimation for Semiparametric Models*. Johns Hopkins University Press Baltimore.
- CHATZINASIIOU, F., LILL, C. M., KYPREOU, K., STEFANAKI, I., NICOLAOU, V., SPYROU, G., EVANGELOU, E., ROEHR, J. T., KODELA, E., KATSAMBAS, A. et al. (2011). Comprehensive field synopsis and systematic meta-analyses of genetic association studies in cutaneous melanoma. *Journal of the National Cancer Institute* **103**, 1227–1235.
- COOK, R. D. (2009). *Regression Graphics: Ideas for Studying Regressions Through Graphics*. John Wiley & Sons.
- COOK, R. D. & LEE, H. (1999). Dimension reduction in binary response regression. *Journal of the American Statistical Association* **94**, 1187–1200.
- COOK, R. D. & WEISBERG, S. (1991). Discussion of sliced inverse regression for dimension reduction. *Journal of the American Statistical Association* **86**, 328–332.
- COX, D. R. (1972). Regression models and life-tables. *Journal of the Royal Statistical Society: Series B* **34**, 187–220.
- DABROWSKA, D. M. et al. (1989). Uniform consistency of the kernel conditional kaplan-meier estimate. *The Annals of Statistics* **17**, 1157–1167.
- DONG, Y. & LI, B. (2010). Dimension reduction for non-elliptically distributed predictors: second-order methods. *Biometrika* **97**, 279–294.
- EDDELBUEITTEL, D. & FRANÇOIS, R. (2011). Rcpp: Seamless R and C++ integration. *Journal of Statistical Software* **40**, 1–18.

- EDELMAN, A., ARIAS, T. A. & SMITH, S. T. (1998). The geometry of algorithms with orthogonality constraints. *SIAM journal on Matrix Analysis and Applications* **20**, 303–353.
- FLEMING, T. R. & HARRINGTON, D. P. (2011). *Counting Processes and Survival Analysis*. John Wiley & Sons.
- GUDBJARTSSON, D. F., SULEM, P., STACEY, S. N., GOLDSTEIN, A. M., RAFNAR, T., SIGURGEIRSSON, B., BENEDIKTSDOTTIR, K. R., THORISDOTTIR, K., RAGNARSSON, R., SVEINSDOTTIR, S. G. et al. (2008). Asip and tyr pigmentation variants associate with cutaneous melanoma and basal cell carcinoma. *Nature genetics* **40**, 886–891.
- HANSEN, L. P. (1982). Large sample properties of generalized method of moments estimators. *Econometrica* **50**, 1029–1054.
- JURECKOVA, J. (1971). Nonparametric estimate of regression coefficients. *The Annals of Mathematical Statistics* **42**, 1328–1338.
- LI, B. & DONG, Y. (2009). Dimension reduction for nonelliptically distributed predictors. *The Annals of Statistics* **37**, 1272–1298.
- LI, B. & WANG, S. (2007). On directional regression for dimension reduction. *Journal of the American Statistical Association* **102**, 997–1008.
- LI, C., ZHAO, H., HU, Z., LIU, Z., WANG, L.-E., GERSHENWALD, J. E., PRIETO, V. G., LEE, J. E., DUVIC, M., GRIMM, E. A. et al. (2008). Genetic variants and haplotypes of the caspase-8 and caspase-10 genes contribute to susceptibility to cutaneous melanoma. *Human Mutation* **29**, 1443–1451.
- LI, K.-C. (1991). Sliced inverse regression for dimension reduction. *Journal of the American Statistical Association* **86**, 316–327.

- LI, K.-C., WANG, J.-L. & CHEN, C.-H. (1999). Dimension reduction for censored regression data. *The Annals of Statistics* **27**, 1–23.
- LIN, D., WEI, L. & YING, Z. (1998). Accelerated failure time models for counting processes. *Biometrika* **85**, 605–618.
- LU, W. & LI, L. (2011). Sufficient dimension reduction for censored regressions. *Biometrics* **67**, 513–523.
- MA, Y. & ZHANG, X. (2015). A validated information criterion to determine the structural dimension in dimension reduction models. *Biometrika* **102**, 409–420.
- MA, Y. & ZHU, L. (2012). A semiparametric approach to dimension reduction. *Journal of the American Statistical Association* **107**, 168–179.
- MACK, Y. P. & SILVERMAN, B. W. (1982). Weak and strong uniform consistency of kernel regression estimates. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete* **61**, 405–415.
- NOCEDAL, J. & WRIGHT, S. J. (2006). *Numerical Optimization*. Springer.
- SILVERMAN, B. W. (1986). *Density Estimation for Statistics and Data Analysis*. CRC press.
- TSIATIS, A. (2007). *Semiparametric Theory and Missing Data*. Springer Science & Business Media.
- TSIATIS, A. A. (1990). Estimating regression parameters using linear rank tests for censored data. *The Annals of Statistics* **18**, 354–372.
- WEISBERG, S. (2002). Dimension reduction regression in r. *Journal of Statistical Software* **7**, 1–22.

- WEN, Z. & YIN, W. (2013). A feasible method for optimization with orthogonality constraints. *Mathematical Programming* **142**, 397–434.
- WU, T., SUN, W., YUAN, S., CHEN, C.-H. & LI, K.-C. (2008). A method for analyzing censored survival phenotype with gene expression data. *BMC bioinformatics* **9**, 417.
- XIA, Y. (2007). A constructive approach to the estimation of dimension reduction directions. *The Annals of Statistics* **35**, 2654–2690.
- XIA, Y., ZHANG, D. & XU, J. (2010). Dimension reduction and semiparametric estimation of survival models. *Journal of the American Statistical Association* **105**, 278–290.
- ZENG, D. & LIN, D. (2007). Maximum likelihood estimation in semiparametric regression models with censored data. *Journal of the Royal Statistical Society: Series B* **69**, 507–564.
- ZHAO, R., ZHANG, J. & ZHU, R. (2017). *orthoDr: An Orthogonality Constrained Optimization Approach for Semi-Parametric Dimension Reduction Problems*. R package version 0.3.0.
- ZHU, L., MIAO, B. & PENG, H. (2006). On sliced inverse regression with high-dimensional covariates. *Journal of the American Statistical Association* **101**, 630–643.

Table 1: Simulation results: Mean ( $\times 10^2$ ) and standard deviations ( $\times 10^2$ , in parenthesis) of Frobenius norm distance (Frob), trace correlation (Tr) and canonical correlation (CCor).

Setting 1 (d=1)				$p = 6$			$p = 12$			$p = 18$		
	Frob	Tr	CCor	Frob	Tr	CCor	Frob	Tr	CCor	Frob	Tr	CCor
Naive	54 (12)	85 (6)	94 (3)	66 (12)	78 (8)	92 (3)	73 (11)	73 (8)	91 (3)			
DS	33 (10)	94 (4)	98 (2)	46 (11)	89 (5)	97 (2)	53 (10)	85 (5)	96 (2)			
IPCW-SIR	64 (13)	78 (9)	91 (4)	75 (11)	71 (9)	89 (4)	80 (11)	68 (9)	89 (4)			
hMave	68 (12)	76 (8)	86 (5)	73 (11)	73 (8)	86 (5)	79 (10)	68 (8)	84 (5)			
Forward	21 (6)	98 (1)	99 (0)	33 (8)	94 (3)	99 (1)	39 (7)	92 (3)	98 (1)			
CP-SIR	26 (9)	96 (3)	99 (1)	40 (10)	91 (4)	98 (1)	49 (9)	88 (4)	97 (1)			
IR-CP	23 (7)	97 (2)	99 (0)	35 (8)	94 (3)	98 (1)	41 (7)	91 (3)	98 (1)			
IR-Semi	23 (8)	97 (2)	99 (0)	37 (8)	93 (3)	98 (1)	44 (8)	90 (4)	98 (1)			
Setting 2 (d=2)				$p = 6$			$p = 12$			$p = 18$		
	Frob	Tr	CCor	Frob	Tr	CCor	Frob	Tr	CCor	Frob	Tr	CCor
Naive	67 (19)	88 (6)	96 (3)	87 (18)	80 (8)	91 (6)	106 (17)	71 (9)	87 (6)			
DS	44 (13)	95 (3)	98 (1)	68 (15)	88 (5)	94 (3)	84 (11)	82 (5)	92 (3)			
IPCW-SIR	83 (19)	82 (8)	94 (3)	98 (17)	75 (9)	90 (6)	114 (16)	67 (9)	86 (8)			
hMave	114 (31)	65 (16)	74 (16)	139 (19)	51 (12)	64 (12)	151 (14)	43 (10)	59 (10)			
Forward	102 (1)	49 (1)	100 (0)	105 (2)	48 (1)	99 (0)	107 (2)	46 (1)	99 (0)			
CP-SIR	37 (11)	96 (2)	98 (1)	61 (12)	90 (4)	96 (2)	78 (10)	85 (4)	93 (2)			
IR-CP	49 (19)	93 (6)	96 (3)	73 (20)	86 (8)	92 (4)	90 (17)	79 (8)	89 (5)			
IR-Semi	39 (14)	96 (3)	98 (2)	65 (16)	89 (6)	94 (3)	83 (15)	82 (6)	91 (3)			
Setting 3 (d=2)				$p = 6$			$p = 12$			$p = 18$		
	Frob	Tr	CCor	Frob	Tr	CCor	Frob	Tr	CCor	Frob	Tr	CCor
Naive	72 (23)	86 (9)	96 (5)	99 (22)	74 (11)	88 (11)	116 (18)	66 (10)	82 (13)			
DS	40 (14)	95 (3)	99 (1)	60 (13)	91 (4)	97 (3)	73 (15)	86 (6)	95 (5)			
IPCW-SIR	113 (26)	66 (13)	81 (14)	129 (15)	58 (9)	74 (12)	133 (11)	55 (7)	74 (12)			
hMave	40 (18)	95 (6)	99 (3)	66 (27)	87 (12)	94 (9)	89 (29)	78 (14)	89 (12)			
Forward	100 (0)	50 (0)	100 (0)	100 (0)	50 (0)	100 (0)	101 (0)	50 (0)	100 (0)			
CP-SIR	34 (11)	97 (2)	99 (1)	55 (11)	92 (3)	97 (2)	67 (11)	88 (4)	96 (3)			
IR-CP	30 (14)	97 (3)	99 (1)	46 (14)	94 (4)	99 (1)	58 (15)	91 (5)	97 (4)			
IR-Semi	19 (8)	99 (1)	100 (0)	29 (8)	98 (1)	100 (0)	40 (11)	96 (2)	99 (1)			
Setting 4 (d=2)				$p = 6$			$p = 12$			$p = 18$		
	Frob	Tr	CCor	Frob	Tr	CCor	Frob	Tr	CCor	Frob	Tr	CCor
Naive	33 (9)	97 (2)	99 (1)	52 (10)	93 (3)	97 (1)	66 (10)	89 (4)	95 (2)			
DS	49 (12)	94 (3)	95 (3)	62 (11)	90 (4)	94 (3)	71 (11)	87 (4)	93 (3)			
IPCW-SIR	35 (9)	97 (2)	98 (1)	52 (10)	93 (3)	97 (1)	64 (10)	89 (3)	95 (2)			
hMave	142 (3)	50 (2)	59 (4)	145 (5)	47 (4)	57 (4)	149 (7)	45 (6)	55 (6)			
Forward	101 (0)	50 (0)	100 (0)	102 (1)	49 (0)	99 (0)	102 (1)	49 (0)	99 (0)			
CP-SIR	36 (7)	97 (1)	98 (1)	51 (8)	93 (2)	97 (1)	63 (8)	90 (3)	95 (1)			
IR-CP	22 (9)	99 (2)	99 (1)	42 (15)	95 (4)	98 (2)	57 (17)	91 (5)	96 (3)			
IR-Semi	13 (5)	99 (0)	100 (0)	24 (7)	98 (1)	99 (0)	34 (10)	97 (2)	99 (1)			

DS: [Li et al. \(1999\)](#); IPCW-SIR: [Lu & Li \(2011\)](#); hMave: [Xia et al. \(2010\)](#); Forward: forward regression; CP-SIR: the computational efficient approach; IR-CP: the counting process inverse regression approach; IR-Semi: the semiparametric inverse regression approach.



Table 2: SKCM data analysis results: the loading vectors ( $\times 10^2$ ) of the first structural dimension.

	Naive	DS	IPCW-SIR	hMave	Forward	CP-SIR	IR-CP	IR-Semi
Age	16	47	10	0	60	59	53	54
TYRP1	-16	-5	-9	24	18	11	39	30
OCA2	18	17	14	-6	21	19	22	5
TYR	-60	-9	-65	-78	-19	-27	-19	9
SLC45A2	11	24	23	14	30	28	16	17
CDKN2A	6	-28	-2	-12	-9	-7	-2	-11
MX2	2	-2	-2	-12	-19	-13	-30	-27
MTAP	-15	-8	-10	-14	-31	-36	-35	-30
MITF	56	-9	43	5	-13	-12	2	-27
VDR	5	-18	-9	10	-10	-6	-4	2
CCND1	-20	35	-21	-5	16	17	18	16
MYH7B	10	-27	5	-4	-29	-32	-30	-48
ATM	-16	-22	2	28	-4	7	0	6
PLA2G6	-22	-16	-21	7	4	-5	-11	-3
CASP8	15	-39	21	-13	-26	-24	-18	-14
AFG3L1	12	26	18	-15	17	10	-6	-9
CDK10	3	8	2	25	-7	-1	9	8
PARP1	-9	3	-22	17	14	18	8	18
CLPTM1L	-8	-5	17	2	-6	-6	-2	-6
ERCC5	-14	25	-17	-7	12	13	22	3
FTO	-3	-3	-8	14	15	17	7	5

DS: [Li et al. \(1999\)](#); IPCW-SIR: [Lu & Li \(2011\)](#); hMave: [Xia et al. \(2010\)](#); Forward: forward regression; CP-SIR: the computational efficient approach; IR-CP: the counting process inverse regression approach; IR-Semi: the semiparametric inverse regression approach.

Table 3: Mean computational time (in seconds)

Dimension	Setting 1			Setting 2		
	$p = 6$	$p = 12$	$p = 18$	$p = 6$	$p = 12$	$p = 18$
Forward	< 1	1	1	< 1	1	1
CP-SIR	< 1	< 1	< 1	< 1	< 1	< 1
IR-CP	< 1	1	1	2	6	11
IR-Semi	1	4	13	8	56	162

Dimension	Setting 3			Setting 4		
	$p = 6$	$p = 12$	$p = 18$	$p = 6$	$p = 12$	$p = 18$
Forward	< 1	< 1	1	< 1	< 1	1
CP-SIR	< 1	< 1	< 1	< 1	< 1	< 1
IR-CP	2	5	8	1	3	7
IR-Semi	12	47	146	6	27	96

Forward: forward regression; CP-SIR: the computational efficient approach; IR-CP: the counting process inverse regression approach; IR-Semi: the semiparametric inverse regression approach. All simulations are done on an Intel Xeon E5-2680v4 processor with 5 parallel threads.

Table 4: Mean ( $\times 10^4$ ) and standard deviation ( $\times 10^4$ ) of parameter estimations (Setting 1,  $p = 6$ ).

		$\beta_2 = 0.5$	$\beta_3 = 0$	$\beta_4 = 0$	$\beta_5 = 0$	$\beta_6 = 0$
Naive	mean	480	-3	267	263	-13
	sd	158	127	137	140	120
DS	mean	477	2	53	55	-6
	sd	127	103	138	146	93
IPCW-SIR	mean	482	-3	343	352	-17
	sd	185	136	166	173	146
hMave	mean	482	2	-380	-394	0
	sd	178	145	155	161	123
Forward	mean	536	36	-3	-1	3
	sd	98	80	81	81	75
	$\widehat{\text{sd}}$	99	80	82	82	74
	coverage	956	920	940	948	952
CP-SIR	mean	511	0	-37	-29	2
	sd	129	104	102	94	90
	$\widehat{\text{sd}}$	134	105	103	105	94
	coverage	960	964	956	960	972
IR-CP	mean	546	45	-3	-4	1
	sd	108	86	88	93	79
	$\widehat{\text{sd}}$	108	88	91	89	82
	coverage	934	924	932	952	968
IR-Semi	mean	520	13	-5	-4	1
	sd	115	91	95	99	80
	$\widehat{\text{sd}}$	117	94	96	98	85
	coverage	932	960	948	944	956

DS: [Li et al. \(1999\)](#); IPCW-SIR: [Lu & Li \(2011\)](#); hMave: [Xia et al. \(2010\)](#);  
Forward: forward regression; CP-SIR: the computational efficient approach;  
IR-CP: the counting process inverse regression; IR-Semi: the semiparametric inverse regression. For the proposed methods, “ $\widehat{\text{sd}}$ ” is the bootstrap estimation of the standard deviation, and “coverage” is the coverage rate of the 95% confidence interval.

Table 5: Mean ( $\times 10^4$ ) and standard deviation ( $\times 10^4$ ) of parameter estimations (Setting 2,  $p = 6$ ).

		$\beta_{13} = 1$	$\beta_{14} = 0$	$\beta_{15} = 0$	$\beta_{16} = 0$	$\beta_{23} = 0$	$\beta_{24} = 1$	$\beta_{25} = 0$	$\beta_{26} = 0$
Naive	mean	1029	-3	134	-138	37	1101	377	-351
	sd	206	209	135	126	412	1080	317	302
DS	mean	1027	-31	12	-8	-2	996	38	-21
	sd	196	195	130	121	265	289	217	223
IPCW-SIR	mean	1033	-14	275	-272	89	1347	551	-504
	sd	230	301	185	159	785	3514	1123	660
hMave	mean	1294	413	1742	-2036	-634	1245	-2692	2902
	sd	12873	11019	35093	36345	10599	8001	31410	32855
CP-SIR	mean	989	3	-17	11	9	979	-8	23
	sd	194	175	134	112	231	220	169	148
	$\widehat{\text{sd}}$	187	194	140	127	248	257	182	161
	coverage	926	959	950	988	983	975	979	983
IR-CP	mean	980	-125	31	-38	45	1127	7	106
	sd	266	347	193	168	466	556	335	262
	$\widehat{\text{sd}}$	270	321	208	188	438	507	338	293
	coverage	967	1000	983	1000	992	992	979	992
IR-Semi	mean	960	-37	-21	-10	79	1054	64	46
	sd	189	176	123	117	224	269	194	154
	$\widehat{\text{sd}}$	196	215	148	133	289	317	228	201
	coverage	963	983	979	983	992	988	975	983

DS: [Li et al. \(1999\)](#); IPCW-SIR: [Lu & Li \(2011\)](#); hMave: [Xia et al. \(2010\)](#); Forward: forward regression; CP-SIR: the computational efficient approach; IR-CP: the counting process inverse regression; IR-Semi: the semiparametric inverse regression. For the proposed methods, “ $\widehat{\text{sd}}$ ” is the bootstrap estimation of the standard deviation, and “coverage” is the coverage rate of the 95% confidence interval.

Table 6: Mean ( $\times 10^4$ ) and standard deviation ( $\times 10^4$ ) of parameter estimations (Setting 3,  $p = 6$ ).

		$\beta_{13} = 1$	$\beta_{14} = 0$	$\beta_{15} = 0$	$\beta_{16} = 0$	$\beta_{23} = 0$	$\beta_{24} = 1$	$\beta_{25} = 0$	$\beta_{26} = 0$
Naive	mean	1017	-26	59	-60	97	818	293	-271
	sd	184	924	275	242	1107	6038	1787	1601
DS	mean	1000	8	1	3	28	1035	1	26
	sd	53	52	44	42	287	344	261	236
IPCW-SIR	mean	1133	156	53	-460	542	1552	374	-1592
	sd	1080	2164	1951	2036	4062	7741	6845	7293
hMave	mean	984	-5	-46	33	-81	969	-139	38
	sd	310	136	63	175	1890	848	325	1057
CP-SIR	mean	998	8	-2	0	13	1043	35	-12
	sd	56	53	42	43	278	313	189	226
	$\widehat{\text{sd}}$	60	61	45	45	295	313	211	208
	coverage	960	972	964	952	964	960	972	932
IR-CP	mean	1003	5	-1	2	21	1012	-6	14
	sd	47	42	40	40	340	189	206	195
	$\widehat{\text{sd}}$	53	55	47	47	358	237	216	210
	coverage	996	988	976	984	968	984	984	972
IR-Semi	mean	1002	4	-2	1	8	1006	-6	7
	sd	31	31	27	27	179	106	98	105
	$\widehat{\text{sd}}$	37	39	33	33	214	149	124	125
	coverage	984	992	988	984	984	988	996	976

DS: [Li et al. \(1999\)](#); IPCW-SIR: [Lu & Li \(2011\)](#); hMave: [Xia et al. \(2010\)](#); Forward: forward regression; CP-SIR: the computational efficient approach; IR-CP: the counting process inverse regression; IR-Semi: the semiparametric inverse regression. For the proposed methods, “ $\widehat{\text{sd}}$ ” is the bootstrap estimation of the standard deviation, and “coverage” is the coverage rate of the 95% confidence interval.

Table 7: Mean ( $\times 10^4$ ) and standard deviation ( $\times 10^4$ ) of parameter estimations (Setting 4,  $p = 6$ ).

		$\beta_{12} = 1$	$\beta_{14} = 0$	$\beta_{15} = 0$	$\beta_{16} = 0$	$\beta_{22} = 0$	$\beta_{24} = -1$	$\beta_{25} = 0$	$\beta_{26} = 0$
Naive	mean	1083	93	83	79	-17	-1043	-2	-11
	sd	132	101	71	82	232	199	141	159
DS	mean	922	-75	-79	-82	-250	-1293	-253	-268
	sd	116	131	92	91	226	263	167	199
IPCW-SIR	mean	1102	107	98	96	-21	-1062	-32	-24
	sd	135	106	82	89	242	225	154	160
hMave	mean	149	-837	-864	-860	951	-122	950	928
	sd	239	194	310	265	419	329	539	443
CP-SIR	mean	873	-127	-122	-125	34	-989	19	13
	sd	102	91	63	70	199	164	126	140
	$\widehat{\text{sd}}$	99	92	71	70	192	178	137	135
	coverage	672	692	620	560	932	948	956	940
IR-CP	mean	975	53	18	16	8	-1007	-2	-3
	sd	62	57	40	42	168	144	104	118
	$\widehat{\text{sd}}$	71	65	49	48	212	175	138	139
	coverage	944	876	952	960	992	984	992	972
IR-Semi	mean	990	19	9	5	10	-1007	0	-6
	sd	50	45	33	34	91	82	64	68
	$\widehat{\text{sd}}$	54	49	37	37	120	106	81	82
	coverage	952	952	964	948	992	972	976	972

DS: [Li et al. \(1999\)](#); IPCW-SIR: [Lu & Li \(2011\)](#); hMave: [Xia et al. \(2010\)](#); Forward: forward regression; CP-SIR: the computational efficient approach; IR-CP: the counting process inverse regression; IR-Semi: the semiparametric inverse regression. For the proposed methods, “ $\widehat{\text{sd}}$ ” is the bootstrap estimation of the standard deviation, and “coverage” is the coverage rate of the 95% confidence interval.

Table 8: Pairwise distance measures ( $\times 10^2$ ) among all methods

	Naive	DS	IPCW-SIR	hMave	Forward	CP-SIR	IR-CP
DS	141						
IPCW-SIR	63	141					
hMave	131	141	125				
Forward	141	87	141	135			
CP-SIR	140	92	140	129	31		
IR-CP	140	108	140	130	67	66	
IR-Semi	137	115	138	138	80	81	76