

Maximizing free energy gain

Artemy Kolchinsky¹, Iman Marvian², Can Gokler³, Zi-Wen Liu⁴, Peter Shor⁵,
Oles Shtanko⁴, Kevin Thompson³, David Wolpert^{1,6,7}, Seth Lloyd^{1,2,4,8,*}

1. Santa Fe Institute, 2. MIT Research Laboratory of Electronics
3. Harvard Engineering and Applied Sciences, 4. MIT Physics, 5. MIT Mathematics
6. MIT Aero Astro, 7. Arizona State University, 8. MIT Mechanical Engineering

* to whom correspondence should be addressed: slloyd@mit.edu

Free energy is energy that is available to do work. Maximizing the free energy gain and the gain in work that can be extracted from a system is important for a wide variety of physical and technological processes, from energy harvesting processes such as photosynthesis to energy storage systems such as fuels and batteries. This paper extends recent results from non-equilibrium thermodynamics [1-20] and quantum resource theory [21-24] to derive closed-form solutions for the maximum possible gain in free energy and extractable work that can be obtained by varying the initial states of classical and quantum stochastic processes. Simple formulae allow the comparison the free energy increase for the optimal procedure with that for a sub-optimal procedure. The problem of finding the optimal free-energy harvesting procedure is shown to be convex and solvable via gradient descent.

Consider a green plant: when the sun rises in the morning, the plant begins to perform photosynthesis and to harvest energy. When the sun sets in the evening the plant stops photosynthesizing. Free energy is energy that is available to do work: it is energy that is unencumbered by entropy, and so is ‘free.’ A plant that has managed to harvest more energy during the course of the day, and to increase its free energy by a larger amount

come evening, will in general have an evolutionary advantage over a plant that attains a smaller increase in free energy. Given a physical system that obeys a particular dynamics of interaction with its surroundings, described by a classical or quantum-mechanical stochastic process, the goal of this paper is to identify the initial probability distribution over the states of the system that leads to the maximum increase in the system's free energy during that process, and to compare the optimal increase in free energy with the increase when the process begins in a sub-optimal state. We include possible constraints on energy, volume, particle number, etc. We show that the problem of finding the optimal initial state for increasing the available free energy and extractable work is convex and can be solved, e.g., by gradient descent: a species in which successive generations get better and better at harvesting free energy is headed for the global optimum.

The last few decades have seen a revolution in non-equilibrium statistical mechanics [1-20], with the realization that many thermodynamic processes are governed by exact and unexpected relations such as the Jarzynski equality [1] and the Crooks fluctuation theorem [2]. A recent example of such a result is the the Kolchinsky-Wolpert theorem [11], which governs the amount of work dissipated in a stochastic process. It provides a simple formula that allows the comparison between the minimum amount of work that could be dissipated, and the amount actually dissipated, as a function of the initial distribution of states for the process. This paper extends the proof of the Kolchinsky-Wolpert theorem to the problem of free energy increase, and to the closely related problem of maximizing the amount of work that can be extracted from a physical system. We compare the free energy and extractable work available to a system at the beginning of a classical or quantum stochastic process, with the amount available at the end of the process. We derive formulae for the maximum increase in free energy increase and extractable work that can be obtained by varying the system's initial probability distribution/quantum state. Our results hold hold when temperature, pressure, etc., vary over the process, and support constraints on energy, volume, etc. In contrast with the resource theory of quantum thermodynamics [21-23], which typically looks at how much work can be extracted from states, we focus on how much work can be extracted from stochastic processes. As noted above, maximum free energy increase/extractable work is important, for example, for photosynthetic organisms who must harvest energy from the sun at different temperatures

and different environmental conditions throughout the day.

Free energy

Free energy measures the amount of energy available in a system to do work when it can exchange energy with an environment at temperature τ [12-24]. Variants of free energy are obtained by considering the cases where the system can exchange other globally conserved quantities with the environment: for example the system could exchange volume (e.g., via a piston) with an environment at pressure p , exchange charge with an environment at voltage V , or exchange molecules with an environment at chemical potential μ , etc. The conserved quantities exchanged with the environment are sometimes called collective variables, and their corresponding potentials (temperature, pressure, etc.), are called generalized conjugate forces [18]. For simplicity of exposition, we first consider just the case of energy and temperature.

The non-equilibrium free energy [13-15] of a system described by a probability distribution $p(x)$, relative to an environment at temperature τ , is

$$F(p) = \langle E \rangle - \tau S = \sum_x p(x) E(x) - \tau \left(- \sum_x p(x) \ln p(x) \right), \quad (1)$$

where $\langle E \rangle$ is the average energy of the system, S is its entropy, and τ is the temperature. We set Boltzmann's constant $k_B = 1$. The non-equilibrium free energy is related to the maximum amount of work W^{ex} that can be extracted from the system by turning on, controlling, and then turning off an interaction with an environment at temperature τ : $W^{ex} = F(p) - F_{th}(p_{th})$, where $F_{th}(p_{th}) = \langle E_{th} \rangle - \tau S_{th}$ is the free energy of the thermal state of the system at temperature τ , characterized by probabilities that obey a Boltzmann-Gibbs distribution, $p_{th}(x) \propto e^{-E(x)/\tau}$. A useful expression for the extractable work is

$$W^{ex} = F(p) - F_{th}(p_{th}) = \tau D(p||p_{th}), \quad (2)$$

where $D(p||p_{th}) = - \sum_x p(x) \ln (p_{th}(x)/p(x))$ is the Kullback-Leibler divergence, also known as the relative entropy. This amount of work can be extracted, for example, by performing a ‘quench’ – a rapid change in the initial energies $E(x) \rightarrow \hat{E}(x) = -\tau \log(p_x)$ – followed by a slow isothermal deformation of the energies $\hat{E}(x)$ back to their final values $E(x)$ [13-15]. As the example of photosynthesis shows, extractable work is a useful and important quantity: it represents a system's energetic ‘cash on hand.’

Free energy increase

Now look at how free energy and extractable work increase over the course of a stochastic process. We assume that the dynamics of the stochastic process itself is fixed: our goal is to maximize free energy increase over all possible initial probability distributions for the system. We begin by deriving a simple criterion for when it is possible to increase free energy at all.

Consider a stochastic process defined by conditional probabilities $p(x_1|x_0)$ for output states x_1 given input state x_0 . In the supplementary material we consider quantum-mechanical processes as well: all our results hold equally for classical and for quantum stochastic processes. The system has the initial probability distribution $p_0(x_0)$, and final distribution $p_1(x_1) = \sum_{x_0} p(x_1|x_0)p_0(x_0)$. We want to compare the free energy/extractable work available at the beginning of the process, with the same quantity after the process has taken place. Note that we are not looking at the amount of work actually extracted in the course the process: instead we consider the amount of work available if we were to extract it at the start of the process, and compare it with the amount available if we were to extract it at the end of the process. We allow the final energy function $E_1(x)$ and temperature τ_1 at the end of the process to be different from the initial energy function $E_0(x)$ and temperature τ_0 : free energies and extractable work are defined relative to E_0 , τ_0 , p_0 at the beginning of the process, and E_1 , τ_1 , p_1 at the end of the process.

First, consider processes that take an initial thermal distribution to a final thermal distribution, possibly at a different temperature (generalized Gibbs-preserving processes): $p_0^{th} \rightarrow p_1^{th}$. The change in extractable work over the process is equal to

$$\Delta W^{ex} = \tau_1 D(p_1 \| p_1^{th}) - \tau_0 D(p_0 \| p_0^{th}). \quad (3)$$

When $\tau_1 \leq \tau_0$, we immediately have $\Delta W^{ex} \leq 0$, because the relative entropy is non-increasing under stochastic processes (the data processing inequality for the K-L divergence). That is, if the process is Gibbs preserving, and the final temperature is less than or equal to the initial temperature, then the amount of extractable work can't increase, no matter how one prepares the system initially.

By contrast, suppose that the process is not Gibbs preserving: $p_0^{th} \rightarrow p_1 \neq p_1^{th}$. Now

if we prepare the system in a thermal state, the extractable work increases:

$$W_1^{ex} - W_0^{ex} = \tau_1 D(p_1 \| p_1^{th}) - \tau_0 D(p_0^{th} \| p_0^{th}) = \tau_1 D(p_1 \| p_{th}) > 0. \quad (4)$$

Note that equation (4) holds for any value of τ_1 , independent of whether it is less than or greater than τ_0 . Combining equations (3-4) implies

Theorem 1:

(1a) A stochastic process allows an increase in extractable work if it is not Gibbs-preserving: i.e., it fails to map an initial thermal state to a final thermal state.

(1b) A Gibbs-preserving stochastic process allows an increase in extractable work only if the final temperature is higher than the initial temperature.

That is, *any* process that does not preserve thermal equilibrium allows the harvesting of free energy. As long as the process doesn't map thermal states to thermal states, there is free energy to be gained. Conversely, processes that preserve thermal equilibrium can't be used to harvest free energy if the temperature does not increase over the course of the process.

Maximizing free energy increase

We now derive a formula for the maximum possible gain in free energy/extractable work. Fix the stochastic process $p(x_1|x_0)$ and look for the initial distribution $q_0(x_0)$ that maximizes the free energy increase $\Delta F(q) = F_1(q_1) - F_0(q_0)$, where the energy function $E_1(x_1)$ at time t_1 need not be the same as the energy function $E_0(x_0)$ at time t_0 . Define a Lagrangian

$$\mathcal{L} = F(q_1) - F(q_0) - \lambda \left(\sum_{x_0} q_0(x_0) - 1 \right), \quad (5)$$

where λ is a Lagrange multiplier to enforce the normalization constraint. The Lagrange equation $\partial \mathcal{L} / \partial q_0(x_0) = 0$ then implies

$$\sum_{x_1} p(x_1|x_0) (E_1(x_1) + \tau \ln q_1(x_1)) - (E_0(x_0) + \tau \ln q_0(x_0)) - \lambda = 0. \quad (6)$$

Here, we require that q_0 has full support, i.e., it lies in the interior of the set of possible initial distributions. Extending an argument from [1], however, we can show that optimal

solutions do indeed have full support under a wide range of circumstances: a sufficient condition for the initial distribution that maximizes the free energy gain to have full support is for the conditional probabilities $p(x_1|x_0)$ to be non-zero. If a solution to equation (6) has full support then it gives the maximum free energy gain and the maximum gain in extractable work, as will now be seen.

A solution that maximizes free energy gain also maximizes the gain in extractable work because these two quantities differ by a constant, the difference in equilibrium free energy between the beginning and end of the process. So maximizing the free energy gain also maximizes the gain in extractable work. Compare the maximum free energy gain with the free energy gain for a different initial distribution $r_0(x_0)$. Multiplying equation (6) by $r_0(x_0) - q_0(x_0)$, and summing over x_0 yields

$$\sum_{x_0} (r_0(x_0) - q_0(x_0)) \frac{\partial \mathcal{L}}{\partial q_0(x_0)} = 0. \quad (7)$$

Writing equation (7) out explicitly yields

$$\Delta F(q) = \Delta E(r) + \tau \left(- \sum_{x_0} r_0(x_0) \ln q_0(x_0) + \sum_{x_1} r_1(x_1) \ln q_1(x_1) \right). \quad (8)$$

Equation (8) allows us to compare the free energy increase when the initial distribution is some other, sub-optimal $r_0(x_0)$ to the maximum free energy increase. Subtracting $\Delta F(r)$ from equation (8) immediately implies

Theorem 2:

$$\Delta F(q) - \Delta F(r) = \tau (D(r_0||q_0) - D(r_1||q_1)). \quad (9)$$

Since stochastic processes do not increase the relative entropy (data processing), the right-hand side is non-negative, confirming that any solution q_0 to the Lagrange equations (6) does indeed give the maximum free energy increase, as long as that optimal solution has full support. Below, we will see that the right-hand side of equation (9) is convex as a function of r_0 , so that there is in fact either a unique maximum, or a single, connected manifold of maxima.

Note that while the left-hand side of equation (9) is about the change of free energy during a stochastic process, the right-hand side of this equality consists purely of information-theoretic quantities, multiplied by the temperature. The change in the relative

entropy can be understood as the loss of distinguishability during the stochastic process: it does not explicitly depend on the system Hamiltonian. Indeed, the formula for extractable work, equation (2), is a simple example of a relation between information-theoretic and physical quantities. Such relations have been found very useful in the resource theory of thermodynamics [20-23]. In fact, equation (2) can be thought as a special case of theorem 2, in which the stochastic process is a thermalizing process, i.e. a process that maps any input state to the thermal state. Theorem 2 gives a relation between information-theoretic and physical quantities that holds for *any* stochastic process.

Theorem 2 gives a simple formula for the lost increase in free energy that comes from preparing the system in the ‘wrong’ initial distribution: the lost free energy is given by comparing the relative entropy between the ‘wrong’ and ‘right’ (i.e., optimal) distributions at the beginning of the process with the the same relative entropy at the end of the process. This relative entropy is multiplied by the temperature τ . It is important to note that the stochastic process itself need not take place by interaction with an environment at temperature τ : rather τ is the temperature of the environment with which the system is going to ‘cash in’ in its free energy by interacting with that environment at some time in the future.

Consider a system such as a photosynthetic organism that undergoes the same stochastic process many times – day after day. Each time it has the opportunity to vary its initial state r to try to improve the amount of free energy increase. In the supplementary material, we prove

Theorem 3: $D(r_0||q_0) - D(r_1||q_1)$ is convex as a function of r_0 .

That is, the problem of trying to maximize free energy increase is a problem in convex optimization, and so is in general tractable: there are no local optima to the problem of maximizing free energy increase, and the maximum can be found by gradient descent. A plant that improves its initial conditions for energy harvesting during the day is by definition, going in the right direction.

Optimal increase of generalized free energies, together with constraints

Theorem 2 can be readily extended to include generalized free energies, including constraints on inputs and outputs [16-23]. Look at the generalized non-equilibrium free

energy (sometimes called a thermodynamic potential)

$$G = \sum_j \beta_j \sum_x r(x) b_j(x) + \tau \sum_x r(x) \ln r(x) \equiv +\vec{\beta} \cdot \langle \vec{B} \rangle - \tau S. \quad (10)$$

Here S is the entropy, τ is the temperature, the $b_j(x)$ are the possible values of the collective variables B_j , and β_j are the corresponding generalized conjugate forces. For example, take B_0 to be the energy, with $\beta_0 = 1$; then B_1 could be volume, with β_1 equal to the pressure, B_2 could be charge, with β_2 equal to voltage, B_3 could be particle number, with β_3 equal to chemical potential, etc.

The method of Lagrange multipliers can now be used to find the initial distribution q_0 that maximizes the generalized free energy increase $\Delta G = G_1 - G_0$, and so also the extractable work $\Delta G - \Delta G_{th}$, where ΔG_{th} is the change in the equilibrium free energy from beginning to end of the process, subject to constraints on the expectation values for observables C_k^0 and C_k^1 at the beginning and the end of the process. By allowing the temperature and potentials to vary from the beginning to end of the the process, we allow the comparison of the case where one ‘cashes in’ one’s free energy at the beginning of the process, at temperature τ_0 and relative to collective variables and potentials B_j^0, β_j^0 , with the case where one cashes in one’s free energy at the end of the process, at temperature τ_1 and relative to collective variables and potentials B_j^1, β_j^1 . As shown in the supplementary material, comparing the generalized free energy increase for another initial distribution r_0 with the maximum free energy increase then yields

Theorem 4:

$$\Delta G(q_0) - \Delta G(r_0) = \tau_0 D(r_0 \| q_0) - \tau_1 D(r_1 \| q_1). \quad (11)$$

$\Delta G(q_0) - \Delta G(r_0)$ is also the difference between the maximum extractable work, and the extractable work obtained by preparing the system in the sub-optimal initial distribution r_0 . As above, the extractable work differs from the non-equilibrium free energy by a term equal to the thermal free energy G_{th} : but the ΔG_{th} terms are the same for q_0 and r_0 and so cancel out in equation (11).

Theorem 4 provides an expression for the maximum free energy increase in a broad range of situations, e.g., when the initial and final temperatures are different, or when constraints are applied to the initial and/or final states.

Note that when the temperature at the end of the process is less than or equal to the temperature at the beginning, $\tau_1 \leq \tau_0$, then the right hand side of equation (11) is always non-negative by the data processing inequality. In this case, any solution to the Lagrange equations is guaranteed to give the maximum free energy increase. The proof of theorem 3 shows that the right hand side of equation is convex and so guarantees that the set of solutions forms a connected manifold. For example, theorem 4 can be applied to the situation when photosynthesis begins during the hot day and ends in the cool evening. The formula for free energy, $F = E - \tau S$, shows that one can always harvest more work by interacting with an environment at lower temperature: the cool evening is a better time to cash in one's free energy than the hot day. If $\tau_1 > \tau_0$, any solution to the Lagrange equations still gives the maximum free energy increase as long as the maximum does not occur on the boundary of the solution space.

We show in the supplementary material that theorems 1-4 hold for quantum mechanical systems evolving according to completely positive maps.

Discussion: This paper discussed the creation of free energy by stochastic and quantum mechanical processes. We showed that a wide variety of stochastic processes support increase in free energy and in extractable work if and only if they fail to map thermal states to thermal states. We then derived formulae for the maximum increase in free energy and extractable work allowed by a physical process, and for the difference in free energy increase between the optimal initial state preparation and sub-optimal preparation. This difference is determined by the difference in Kullback-Leibler divergence between the initial and final probability distributions. For macroscopic systems, the deficit in free energy harvested by a sub-optimal system preparation may itself be a macroscopic quantity. For a living system that requires free energy to survive and to reproduce, there is considerable evolutionary pressure to increase the amount of free energy harvested. The convexity of the increase in Kullback-Liebler divergence means that finding the optimal initial distribution for harvesting free energy is a problem in convex optimization: there are no local optima, and the problem of finding the global optimum can be solved using, e.g., gradient descent. For example, a species where each generation does slightly better at harvesting free energy will eventually approach the global maximum rate.

Acknowledgements: This work was supported by NSF under an INSPIRE program. S.L. was supported by ARO and AFOSR. AK and DHW would like to thank the Santa Fe Institute for helping to support this research. This paper was made possible through the support of Grant No. TWCF0079/AB47 from the Templeton World Charity Foundation, Grant No. FQXi-RH13-1349 from the FQXi foundation, and Grant No. CHE-1648973 from the U.S. National Science Foundation. The opinions expressed in this paper are those of the authors and do not necessarily reflect the view of Templeton World Charity Foundation.

References

- [1] C. Jarzynski, *Phys. Rev. Lett.* **78**, 2690 (1997).
- [2] G.E. Crooks, *Phys. Rev. E* **60**, 2721 (1999).
- [3] G.E. Crooks, *J. Stat. Phys.* **90**, 1481 (1998).
- [4] H. Touchette and S. Lloyd, *Physica A* **331**, 140 (2004).
- [5] T. Sagawa and M. Ueda, *Phys. Rev. Lett.* **102**, 250602 (2009).
- [6] R. Dillenschneider and E. Lutz, *Phys. Rev. Lett.* **104**, 198903 (2010).
- [7] M. Esposito and C. Van den Broeck, *Phys. Rev. E* **82**, 011143 (2010).
- [8] K. Wiesner, M. Gu, E. Rieper, and V. Vedral, *Proc. Roy. Soc. A* **468**, 4058 (2012).
- [9] T. Sagawa and M. Ueda, *Phys. Rev. Lett.* **109**, 180602 (2012).
- [10] S. Still, D. A. Sivak, A. J. Bell, and G. E. Crooks, *Phys. Rev. Lett.* **109**, 120604 (2012).
- [11] A. Kolchinsky, D.H. Wolpert, ‘Dependence of dissipation on the initial distribution over states,’ arXiv: 1607.00956v2 (2016).
- [12] E.T. Jaynes, *Phys. Rev.* **106**, 620-630 (1957); *Phys. Rev.* **108**, 171-190 (1957).
- [13] J.M.R. Parrondo, J.M Horowitz, T. Sagawa, *Nat. Phys.* **11**, 131-139 (2015).
- [14] H-H. Hasegawa, J. Ishikawa, K. Takara, D.J. Driebe, *Phys. Lett. A* **374**, 10011004 (2010).
- [15] K. Takara, H-H. Hasegawa, D.J. Driebe, *Phys. Lett. A* **375**, 8892 (2010).
- [16] B. Gaveau and L. S. Schulman, *Phys. Lett. A* **229**, 347 (1997).
- [17] H. Ge, H. Qian, *Phys. Rev. E* **81**, 051133 (2010).
- [18] G.E. Crooks, *Phys. Rev. Lett.* **99**, 100602, (2007).
- [19] C. Esposito, C. van den Broeck, *Euro. Phys. Lett.* **95**, 40004 (2011).
- [20] Y. Guryanova, S. Popescu, A.J. Short, R. Silva, P. Skrzypczyk, *Nat. Comm.* **7**, 12049 (2016).
- [21] M. Horodecki, J. Oppenheim, *Nat. Comm.* **4**, 2059 (2013); arXiv:1111.3834.
- [22] F.G.S.L. Brandao, M. Horodecki, J. Oppenheim, J.M. Renes, R.W. Spekkens, *Phys. Rev. Lett.* **111**, 250404 (2013); arXiv: 1111.3882.

[23] C. Sparaciari, J. Oppenheim, T. Fritz, ‘A Resource Theory for Work and Heat,’ arXiv:1607.01302.

[24] Z-W. Liu, X. Hu, S. Lloyd *Phys. Rev. Lett.* **118**, 060502 (2017); arXiv: 1606.03723.

[25] To see that the optimum distribution generically has full support, suppose that the maximum free energy gain occurs for a distribution where $q_0(x_0) = 0$ for some x_0 . When $q_1(x_1) > 0$ for all x_1 , we see from equation (6) that perturbing $q_0(x_0) \rightarrow \delta > 0$ induces a large positive increase in the change of free energy, contradicting the assumption that the initial distribution gave the maximum free energy change.

Supplementary material

Proof of theorem 3: change in free energy is convex

Take $\tau = 1$ and consider the function $\Delta F(r_0) = D(r_0 \| q_0) - D(r_1 \| q_1)$. We show that ΔF is convex as a function of r_0 . Consider conditional probabilities $s(x_0|c)$, $c = 0, 1$ with $s(x_0) = \sum_c p(c)s(x_0|c)$. We want to show that

$$p(0)\Delta F(s(x_0|c=0)) + p(1)\Delta F(s(x_0|c=1)) - \Delta F(s(x_0)) \geq 0. \quad (S1)$$

Rewriting equation (S1) in terms of entropies, we see that the terms dependent on q_0, q_1 drop out, so that our goal is to show that

$$-S(X_0|C) + S(X_0) - S(X_1) + S(X_1|C) \geq 0, \quad (S2)$$

where $S(X_0|C)$ is the conditional entropy of X_0 given C , similarly for $S(X_1|C)$. But $S(X) - S(X|C)$ is the mutual information between X and C , and the data processing inequality for mutual information – mutual information does not increase under stochastic processes applied to the systems X and C separately – implies the desired result.

When $\tau_0 \geq \tau_1$, the same proof implies the convexity of $\Delta F(r_0) = \tau_0 D(r_0 \| q_0) - \tau_1 D(r_1 \| q_1)$.

Proof of theorem 4: generalized free energy increase

Consider a stochastic process as in the text. Look at the generalized free energy

$$G = \tau \sum_x p(x) \ln p(x) + \sum_j \beta_j \sum_x p(x) b_j(x) \equiv -\tau S + \vec{\beta} \cdot \vec{B}. \quad (S3)$$

Here S is the entropy, the $b_j(x)$ are the possible values of observables B_j , and β_j is a generalized temperature.

Let

$$G_0 = -\tau_0 S_0 + \vec{\beta}_0 \cdot \vec{B}_0 \quad (S4)$$

be the value of the generalized free energy at the beginning of the process, defined relative to the initial probability distribution $r_0(x_0)$, the inverse ‘temperatures’ β_j^0 , and the quantities, B_j^0 . Similarly, let

$$G_1 = -\tau_1 S_1 + \vec{\beta}_1 \cdot \vec{B}_1 \quad (S5)$$

be the quantity at the end of the process, defined relative to the final probability distribution $r_1(x_1)$, the ‘temperatures’ β_j^1 , and the quantities, B_j^1 . As noted, we do not require that the initial temperatures and the quantities B with which they are associated are the same as the final temperatures and quantities. For example, B_0^0 , τ_0 could be the Hamiltonian and temperature for the system at the beginning of the process, and B_0^1 , τ_1 could be a different Hamiltonian and inverse temperature at the end of the process.

Now use the method of Lagrange multipliers to extremize the generalized free energy increase $\Delta G = G_1 - G_0$ subject to constraints $\sum_{x_0} p(x_0) c_k^0(x_0) = \bar{c}_k^0$, $\sum_{x_1} p(x_1) c_k^1(x_1) = \bar{c}_k^1$ on the expectation values for observables C_k^0 and C_k^1 at the beginning and the end of the process. Introduce Lagrange multipliers γ_k^0 , γ_k^1 , and use the notation $\sum_k \gamma_k (\sum_x p(x) c_k(x) - \bar{c}_k) = \vec{\gamma} \cdot (\vec{C} - \vec{c})$. Define a Lagrangian

$$\mathcal{L} = \alpha S - \vec{\beta} \cdot B - \vec{\gamma} \cdot (\vec{C} - \vec{c}) - \lambda (\sum_x p(x) - 1), \quad (S6)$$

and the ‘difference’ Lagrangian

$$\Delta \mathcal{L} = \mathcal{L}_1 - \mathcal{L}_0. \quad (S7)$$

Extremizing $\Delta \mathcal{L}$ over initial probability distributions $r_0(x_0)$ subject to the constraints on input and output observables leads to Lagrange equations that take the form

$$\nabla_{r_0(x_0)} \Delta \mathcal{L} = 0, \quad \nabla_{\vec{\gamma}^0} \Delta \mathcal{L} = 0, \nabla_{\vec{\gamma}^1} \Delta \mathcal{L} = 0, \quad \nabla_{\lambda} \Delta \mathcal{L} = 0. \quad (S8)$$

It is instructive to write out the first of these Lagrange equations explicitly. Taking the derivative of $\Delta \mathcal{L}$ with respect to $p_0(x_0)$ yields

$$\begin{aligned} \frac{\partial}{\partial p_0(x_0)} \Delta \mathcal{L} &= \sum_{x_1} p(x_1 | x_0) (\tau_1 \ln p(x_1) + \sum_j \beta_j^1 b_j^1(x_1) + \sum_k \gamma_k^1 c_k^1(x_1)) \\ &\quad - \tau_0 \ln p(x_0) + \sum_j \beta_j^0 b_j^0(x_0) + \sum_k \gamma_k^0 c_k^0(x_0) - \lambda \\ &= 0. \end{aligned} \quad (S9)$$

Let $q_0(x_0)$ be an initial distribution that solves the Lagrange equations (S7), and let $r_0(x_0)$ be any other initial distribution that satisfies both initial and final constraints. Equation (S8) then implies that

$$\sum_{x_0} (r_0(x_0) - q_0(x_0)) \frac{\partial}{\partial q_0(x_0)} \Delta \mathcal{L} = 0. \quad (S10)$$

Combining equations (S9-10) with the definition of G in equation (S3) above yields the desired result, theorem (4) :

$$\Delta G(q_0) - \Delta G(r_0) = \tau_0 D(r_0 \| q_0) - \tau_1 D(r_1 \| q_1), \quad (S11)$$

where as above, $D(r \| q) = -\sum_x r(x) \ln q(x)/r(x)$ is the Kullback-Liebler divergence/relative entropy. Here $\Delta G(r_0)$, $\Delta G(q_0)$ are the changes in the free energy G over the process when the initial probability distributions are r_0 , q_0 respectively. The data processing inequality states that relative entropy is non-increasing in a stochastic process. Consequently, the right hand side of equation (S11) is non-negative as long as $\tau_1 \leq \tau_0$ showing that q_0 maximizes $\Delta G(q_0)$ subject to the constraints on the input and output distributions.

Quantum mechanical version

The quantum-mechanical version of a stochastic process is a completely positive map, which can be described in Kraus form as

$$\rho_0 \rightarrow \rho_1 = \sum_k A_k \rho_0 A_k^\dagger. \quad (S12)$$

Take $S(\rho) = -\text{tr} \rho \ln \rho$, and observables and constraints to correspond to Hermitian operators B_j , C_k with expectation values $\text{tr} B_j$, $\text{tr} C_k$. Defining Lagrangians as above with these definitions, demanding stationarity under variation in the initial state $\rho_0 \rightarrow \rho_0 + \delta \rho_0$, and using the cyclic property of the trace yields equation (S10), where the Kullback-Leibler divergences D are defined to be quantum K-L divergences

$$D(\rho \| \sigma) = -\text{tr} \rho \ln \sigma + \text{tr} \rho \ln \rho. \quad (S13)$$

The solution to the Lagrange equations always yields the optimal value of free energy increase when the output states ρ_1 have full rank for any input state ρ_0 . So theorems (1-4) (as well as the original K-W theorem) hold for quantum-mechanical systems as well.