# A Proof of the Explicit Formula for Product Moments of Multivariate Gaussian Random Variables

Iickho Song\*

School of Electrical Engineering Korea Advanced Institute of Science and Technology 291 Daehag Ro, Yuseong Gu, Daejeon 34141 Korea

#### Abstract

A detailed proof of a recent result on explicit formulae for the product moments  $E\{X_1^{a_1}X_2^{a_2}\cdots X_n^{a_n}\}$  of multivariate Gaussian random variables is provided in this note.

*Keywords:* jointly Gaussian random variables, Gaussian random vector, joint moment, multivariate Gaussian random variables, product moment

### 1. Introduction

5

Recently, explicit formulae for the product moments of multivariate Gaussian random variables have been derived in Song and Lee (2015). The formulae not only are simpler and more concise than any of the other ones but also allow us to promptly compute any term of the product moments.

Since the publication of the results, a huge number of requests has been received for the details of the proof. In this note, we provide a detailed proof of the main result of Song and Lee (2015).

 $<sup>^* \</sup>rm Corresponding author. Tel.: +82-42-350-3445. This study was supported by the National Research Foundation of Korea, under Grant NRF-2015R1A2A1A01005868, for which the author wishes to express his thanks.$ 

Email address: i.song@ieee.org (Iickho Song)

## 2. Proof of the Explicit Formula

#### 10 2.1. The Explicit Formula

Let  $\{X_i\}_{i=1}^n$  be multivariate Gaussian random variables with mean vector  $\boldsymbol{\mu} = (\mu_1, \mu_2, \cdots, \mu_n)$  and  $n \times n$  variance-covariance matrix  $\boldsymbol{\Phi} = [\varphi_{ij}]$ , where  $\mu_i = E\{X_i\}$  denotes the expected value of  $X_i$  and  $\varphi_{ij} = E\{X_iX_j\} - \mu_i\mu_j$  denotes the covariance between  $X_i$  and  $X_j$  if  $i \neq j$  and variance of  $X_i$  if i = j.

<sup>15</sup> One of the main results of Song and Lee (2015) is as follows:

**Theorem 1.** For the product moments

$$E\left\{\prod_{i=1}^{n} X_{i}^{a_{i}}\right\} = \sum_{\boldsymbol{l}\in S_{\boldsymbol{a}}} d_{\boldsymbol{a},\boldsymbol{l}} \left(\prod_{i=1}^{n} \prod_{j=i}^{n} \varphi_{ij}^{l_{ij}}\right) \left(\prod_{j=1}^{n} \mu_{j}^{L_{\boldsymbol{a},j}}\right)$$
(1)

of multivariate Gaussian random variables, we have

$$d_{\boldsymbol{a},\boldsymbol{l}} = \frac{\prod_{k=1}^{n} a_k!}{2^{M_{\boldsymbol{l}}} \left(\prod_{i=1}^{n} \prod_{j=i}^{n} l_{ij}!\right) \left(\prod_{j=1}^{n} L_{\boldsymbol{a},j}!\right)}$$
(2)

for  $l \in S_a$ , where  $M_l = \sum_{k=1}^n l_{kk}$ .

Here, the elements of the two sets  $\boldsymbol{a} = \{a_1, a_2, \cdots, a_n\}$  and  $\boldsymbol{l} = \{\{l_{ij}\}_{i=1}^n\}_{j=i}^n$ are nonnegative integers,

$$L_{a,k} = a_k - l_{kk} - \sum_{j=1}^n l_{jk}$$
(3)

with  $l_{ji} = l_{ij}$ , and  $S_a$  denotes the collection of all the sets l such that

$$\{L_{a,k} \ge 0\}_{k=1}^{n} \,. \tag{4}$$

2.2. Proof

Let us prove Theorem 1 by mathematical induction. First, it is clear from Lemma 2.2, Theorem 1, and Theorem 2 of Song and Lee (2015) that (2) holds for n = 1, 2, and 3, respectively. Next, assume that (2) is true when n = m - 1. Then, since

$$E\{X_1^{a_1}X_2^{a_2}\cdots X_m^{a_m}\} = E\{X_1^{a_1}X_2^{a_2}\cdots X_{m-1}^{a_{m-1}}\}E\{X_m^{a_m}\}$$
(5)

when  $\varphi_{1m} = \varphi_{2m} = \cdots = \varphi_{m-1,m} = 0$ , we get  $[d_{a_2,l_2}]_{l_{1m} \mapsto 0, l_{2m} \mapsto 0, \cdots, l_{m-1,m} \mapsto 0} = d_{a_1,l_1} d_{a_m,l_{mm}}$ , or

$$[d_{\boldsymbol{a}_{2},\boldsymbol{l}_{2}}]_{l_{1m}\mapsto 0,l_{2m}\mapsto 0,\cdots,l_{m-1,m}\mapsto 0} = \frac{\prod_{k=1}^{m} a_{k}!}{2^{M_{l_{2}}} \left(\prod_{i=1}^{m-1} \prod_{j=i}^{m-1} l_{ij}!\right) l_{mm}! \left(\prod_{j=1}^{m-1} L_{\boldsymbol{a}_{1},j}!\right) (a_{m} - 2l_{mm})!}$$
(6)

using (2) with n = m - 1 and

$$d_{a_1,l} = \frac{a_1!}{2^l l! (a_1 - 2l)!} \tag{7}$$

<sup>30</sup> (Eq. (5), Song and Lee (2015)) for  $l = 0, 1, \dots, \lfloor \frac{a_1}{2} \rfloor$ , where  $\mathbf{a}_1 = \{a_i\}_{i=1}^{m-1}$ ,  $\mathbf{a}_2 = \mathbf{a}_1 \cup \{a_m\}, \mathbf{l}_1 = \{l_{ij}\}$  with  $j = i, i+1, \dots, m-1$  for  $i = 1, 2, \dots, m-1$ , and  $\mathbf{l}_2 = \mathbf{l}_1 \cup \{l_{im}\}_{i=1}^m$ . Here, the symbol  $\mapsto$  denotes a substitution: For example,  $\alpha \mapsto \beta$  means the substitution of  $\alpha$  with  $\beta$ .

Next, employing the result

$$\frac{\partial}{\partial\varphi_{ij}}E\left\{\prod_{k=1}^{n}X_{k}^{a_{k}}\right\} = a_{i}a_{j}E\left\{X_{i}^{a_{i}-1}X_{j}^{a_{j}-1}\prod_{\substack{k=1\\k\neq i,j}}^{n}X_{k}^{a_{k}}\right\}$$
(8)

for  $(i, j) = (1, m), (2, m), \dots, (m - 1, m)$  in Price (1958), and then taking some steps (similar to those from (16) to (20) in Song and Lee (2015)), we will get

$$[d_{\boldsymbol{a}_2,\boldsymbol{l}_2}]_{l_{im}\mapsto l_{im}+1} = \frac{a_i!a_m! [d_{\boldsymbol{a}_2,\boldsymbol{l}_2}]_{l_{im}\mapsto 0, a_i\mapsto a_i-l_{im}-1, a_m\mapsto a_m-l_{im}-1}{(a_i-l_{im}-1)! (a_m-l_{im}-1)! (l_{im}+1)!}$$
(9)

and then, by substituting  $l_{im} + 1$  with  $l_{im}$  in (9), we have

$$d_{a_2,l_2} = \frac{a_i!a_m! \left[d_{a_2,l_2}\right]_{l_{im} \to 0, a_i \mapsto a_i - l_{im}, a_m \mapsto a_m - l_{im}}{(a_i - l_{im})! (a_m - l_{im})! (l_{im})!}$$
(10)

for  $i = 1, 2, \dots, m-1$ . Now, letting  $l_{2m} = 0$  in (10) with i = 1, we get

$$[d_{a_2,l_2}]_{l_{2m}\mapsto 0} = \frac{a_1!a_m! [d_{a_2,l_2}]_{l_{1m}\mapsto 0, l_{2m}\mapsto 0, a_1\mapsto a_1-l_{1m}, a_m\mapsto a_m-l_{1m}}{(a_1-l_{1m})! (a_m-l_{1m})! l_{1m}!},$$
(11)

which can be used into (10) with i = 2 to produce

$$d_{\boldsymbol{a}_{2},\boldsymbol{l}_{2}} = \frac{a_{2}!a_{m}!}{(a_{2}-l_{2m})!(a_{m}-l_{2m})!l_{2m}!} \times \frac{a_{1}!(a_{m}-l_{2m})!}{(a_{1}-l_{1m})!(a_{m}-l_{1m}-l_{2m})!l_{1m}!} \times [d_{\boldsymbol{a}_{2},\boldsymbol{l}_{2}}]_{l_{1m}\mapsto 0, l_{2m}\mapsto 0, a_{1}\mapsto a_{1}-l_{1m}, a_{2}\mapsto a_{2}-l_{2m}, a_{m}\mapsto a_{m}-l_{1m}-l_{2m}}.$$
 (12)

40 Subsequently, letting  $l_{3m} = 0$  in (12), we get

$$\begin{aligned} \left[d_{a_{2},l_{2}}\right]_{l_{3m}\mapsto0} &= \frac{a_{1}!a_{2}!a_{m}!}{l_{1m}!l_{2m}!\left(a_{1}-l_{1m}\right)!\left(a_{2}-l_{2m}\right)!\left(a_{m}-l_{1m}-l_{2m}\right)!} \\ &\times \left[d_{a_{2},l_{2}}\right]_{l_{km}\mapsto0 \text{ for } k=1,2,3, \ a_{1}\mapsto a_{1}-l_{1m},a_{2}\mapsto a_{2}-l_{2m},a_{m}\mapsto a_{m}-l_{1m}-l_{2m}}, \end{aligned}$$
(13)

which can be employed into (10) with i = 3 to produce

$$d_{a_{2},l_{2}} = \frac{a_{3}!a_{m}!}{(a_{3}-l_{3m})!(a_{m}-l_{3m})!l_{3m}!} \times \frac{a_{1}!a_{2}!(a_{m}-l_{3m})!}{l_{1m}!l_{2m}!(a_{1}-l_{1m})!(a_{2}-l_{2m})!(a_{m}-l_{1m}-l_{2m}-l_{3m})!} \times [d_{a_{2},l_{2}}]_{l_{km}\mapsto 0,a_{k}\mapsto a_{k}-l_{km}} \text{ for } k=1,2,3, a_{m}\mapsto a_{m}-l_{1m}-l_{2m}-l_{3m} .$$
(14)

From (14), it can be formulated

$$d_{\boldsymbol{a}_{2},\boldsymbol{l}_{2}} = \frac{\left(\prod_{k=1}^{i} a_{k}!\right) a_{m}!}{\left(\prod_{k=1}^{i} l_{km}!\right) \left\{\prod_{k=1}^{i} (a_{k} - l_{km})!\right\} \left(a_{m} - \sum_{k=1}^{i} l_{km}\right)!} \times [d_{\boldsymbol{a}_{2},\boldsymbol{l}_{2}}]_{l_{km} \mapsto 0, a_{k} \mapsto a_{k} - l_{km}} \text{ for } k=1,2,3, a_{m} \mapsto a_{m} - \sum_{k=1}^{i} l_{km}}$$
(15)

with i = 3. Next, letting  $l_{4m} = 0$  in (15), we will get  $[d_{a_2,l_2}]_{l_{4m}\mapsto 0}$  which can be employed into (10) with i = 4 to produce (15) with i = 4. After proceeding the <sup>45</sup> above procedures until i = m - 1, we will eventually get

$$d_{a_{2},l_{2}} = \frac{\left(\prod_{k=1}^{m-1} a_{k}!\right) a_{m}!}{\left(\prod_{k=1}^{m-1} l_{km}!\right) \left\{\prod_{k=1}^{m-1} (a_{k} - l_{km})!\right\} \left(a_{m} - \sum_{k=1}^{m-1} l_{km}\right)!} \times \left[d_{a_{2},l_{2}}\right]_{l_{km} \mapsto 0, a_{k} \mapsto a_{k} - l_{km} \text{ for } k=1,2,\cdots,m-1, a_{m} \mapsto a_{m} - \sum_{k=1}^{m-1} l_{km}} (16)$$

from  $[d_{a_2,l_2}]_{l_{m-1,m}\mapsto 0}$  (obtained by letting  $l_{m-1,m} = 0$  in (15) with i = m - 2)

and (10) with i = m - 1. Finally, by combining (6) into (16), we can get

$$d_{a_{2},l_{2}} = \frac{\left(\prod_{k=1}^{m-1} a_{k}!\right) a_{m}!}{\left(\prod_{k=1}^{m-1} l_{km}!\right) \left\{\prod_{k=1}^{m-1} (a_{k} - l_{km})!\right\} \left(a_{m} - \sum_{k=1}^{m-1} l_{km}\right)!} \\ \times \frac{\left\{\prod_{k=1}^{m-1} (a_{k} - l_{km})!\right\} \left(a_{m} - \sum_{k=1}^{m-1} l_{km}\right)!}{2^{M_{l_{2}}} \left(\prod_{k=1}^{m-1} \prod_{j=i}^{m-1} l_{kj}!\right) l_{mm}!} \\ \times \frac{1}{\prod_{j=1}^{m-1} \left([L_{a_{1},j}]_{a_{k} \mapsto a_{k} - l_{km} \text{ for } k=1,2,\cdots,m-1\right)!} \\ \times \frac{1}{\left(a_{m} - \sum_{k=1}^{m-1} l_{km} - 2l_{mm}\right)!} \\ = \frac{\prod_{k=1}^{m} a_{k}!}{2^{M_{l_{2}}} \left(\prod_{i=1}^{m} \prod_{j=i}^{m} l_{ij}!\right) \left(\prod_{j=1}^{m} L_{a_{2},j}!\right)},$$
(17)

which implies that (2) holds when n = m also.

## 3. Conclusion

We have provided a detailed proof of an elegant explicit formulae for product moments  $E\{X_1^{a_1}X_2^{a_2}\cdots X_n^{a_n}\}$  of multivariate Gaussian variables derived recently.

## References

55

Price, R., 1958. A useful theorem for nonlinear devices having Gaussian inputs. IRE Trans. Inform. Theory 4, 69–72.

Song. I., Lee, S., 2015, Explicit formulae for product moments of multivariate Gaussian random variables, *Stat.*, *Prob. Lett.* 100, 27-34.