

Inverse Moment Methods for Sufficient Forecasting using High-Dimensional Predictors*

Wei Luo, Lingzhou Xue[†] and Jiawei Yao

CUNY – Baruch College, Pennsylvania State University and Citadel LLC

Abstract

We consider forecasting a single time series using high-dimensional predictors in the presence of a possible nonlinear forecast function. The sufficient forecasting (Fan et al., 2016) used sliced inverse regression to estimate lower-dimensional sufficient indices for forecasting. However, Fan et al. (2016) is fundamentally limited to the inverse first-moment method, by assuming the restricted fixed number of factors, linearity condition for factors, and monotone effect of factors on the response. In this work, we study the inverse second-moment method using directional regression and the inverse third-moment method to extend the methodology and applicability of the sufficient forecasting. As the number of factors diverges with the dimension of predictors, the proposed method relaxes the distributional assumption of the predictor and enhances the capability of capturing the non-monotone effect of factors on the response. We not only provide a high-dimensional analysis of inverse moment methods such as exhaustiveness and rate of convergence, but also prove their model selection consistency. Our proposed methods are demonstrated in both simulation studies and an empirical study

*The authorship is on the alphabetical order. Three authors contributed equally to this paper. The authors thank Eric Aldrich, Panle Jia Barwick, Bing Li, Shanjun Li, Anna Mikusheva, Minchul Shin, Tim Vogelsang, Mark Watson, and seminar participants at Penn State University (Econometrics), New York University (Stern), Peking University (Statistics), Renmin University of China (Statistics and Big Data) and Cornell Institute for China Economic Research for their helpful comments and suggestions. Wei Luo's research is supported by CUNY – Baruch College. Lingzhou Xue's research is supported by the American Mathematical Society Simons Travel Grant and the National Science Foundation grant DMS-1505256.

[†]Corresponding author. Email: lzxue@psu.edu.

of forecasting monthly macroeconomic data from 1959 to 2016. During our theoretical development, we prove an invariance result for inverse moment methods, which make a separate contribution to the sufficient dimension reduction.

Key Words: factor model; forecasting; inverse moments; learning indices; principal components; regression; sufficient dimension reduction.

1 Introduction

Forecasting using high-dimensional predictors is an increasingly important research topic in statistics, biostatistics, macroeconomics and finance. A large body of literature has contributed to forecasting in a data rich environment, with various applications such as the forecasts of market prices, dividends and bond risks (Sharpe, 1964; Lintner, 1965; Ludvigson and Ng, 2009), macroeconomic outputs (Stock and Watson, 1989; Bernanke et al., 2005), macroeconomic uncertainty and fluctuations (Ludvigson and Ng, 2007; Jurado et al., 2015), and clinical outcomes based on massive genetic, genomic and imaging measurements. Motivated by principal component regression, the pioneering papers by Stock and Watson (2002*a,b*) systematically introduced the forecasting procedure using factor models, which has played an important role in macroeconomic analysis. Recently, Fan et al. (2016) further extended the analysis of Stock and Watson (2002*a,b*) to allow for a nonparametric nonlinear forecast function and multiple nonadditive forecasting indices. Following Fan et al. (2016), we consider the following factor model with a target variable y_{t+1} that we aim to forecast:

$$y_{t+1} = g(\phi_1' \mathbf{f}_t, \dots, \phi_L' \mathbf{f}_t, \epsilon_{t+1}), \quad (1.1)$$

$$x_{it} = \mathbf{b}_i' \mathbf{f}_t + u_{it}, \quad 1 \leq i \leq p, \quad 1 \leq t \leq T, \quad (1.2)$$

where x_{it} is the i -th high-dimensional predictor observed at time t , \mathbf{b}_i is a $K \times 1$ vector of factor loadings, \mathbf{f}_t is a $K \times 1$ vector of common factors driving both the predictor and the response, $g(\cdot)$ is an unknown forecast function that is possibly nonadditive and nonseparable, u_{it} is an idiosyncratic error, and ϵ_{t+1} is an independent stochastic error. Here, ϕ_1, \dots, ϕ_L , $\mathbf{b}_1, \dots, \mathbf{b}_p$ and $\mathbf{f}_1, \dots, \mathbf{f}_T$ are unobserved vectors.

Had the factors $\mathbf{f}_1, \dots, \mathbf{f}_t$ been observed, model (1.1) would be commonly adopted in the literature of sufficient dimension reduction. The linear space spanned by ϕ_1, \dots, ϕ_L , denoted by $\mathcal{S}_{y|\mathbf{f}}$, is the parameter of interest that is identifiable and known as the central subspace (Cook, 1998). Multiple methods have been proposed to estimate $\mathcal{S}_{y|\mathbf{f}}$, among which

a main family employ inverse moments, i.e. the moments of the conditional distribution $\mathbf{f}_t|y_t$, and are called the inverse regression methods. Representative members include sliced inverse regression (Li, 1991), which uses the inverse first moment $E(\mathbf{f}_t|y_{t+1})$, sliced average variance estimation (Cook and Weisberg, 1991) and directional regression (Li and Wang, 2007), which additionally use the inverse second moment $E(\mathbf{f}_t\mathbf{f}_t'|y_{t+1})$, and moreover, the inverse third-moment method (Yin and Cook, 2003), etc. Generally, methods that employ higher-order inverse moments can capture more comprehensive information from the data, which leads to more ability in exhaustive estimation, i.e. detecting all the directions in $\mathcal{S}_{y|\mathbf{f}}$. The price, on the other hand, is to estimate more moments and impose additional distributional assumptions on the factors; see Li (1991), Yin and Cook (2003), and Li and Wang (2007) for more details.

A commonly recognized limit of sufficient dimension reduction methods, including sliced inverse regression, is that they can only handle predictors with either a finite dimension or a diverging dimension that is dramatically smaller than the sample size (Zhu et al., 2006). Therefore, even though it is theoretically desirable to directly apply sufficient dimension reduction to forecasting, using x_t as the predictor and y_{t+1} as the response, none of the existing sufficient dimension reduction methods would be readily applicable. For this reason, it is necessary to reduce the dimension of the predictor prior to sufficient dimension reduction, for which adopting the factor model (1.2) is a reasonable choice. An alternative choice can be found in Jiang and Liu (2014) and Yu et al. (2016), etc.

Following this logic, Fan et al. (2016) introduced the sufficient forecasting scheme to use factor analysis in model (1.2) to estimate \mathbf{f}_t , and apply sliced inverse regression in model (1.1) with the estimated factors as the predictor. Such a combination provides a promising forecasting technique that not only extracts the underlying commonality of the high-dimensional predictor but also models the complex dependence between the predictor and the forecast target. Meanwhile, it allows the dimension of the predictor to diverge and even become much larger than the number of observations, which is intrinsically appealing to solving high-dimensional forecasting problems.

It is important to note that the consistency of the sequential procedure in Fan et al. (2016) is not granted as it may appear. Let $S_{y|\hat{\mathbf{f}}}$ be the central subspace based on the estimated factors $\hat{\mathbf{f}}_t$'s. Without additional assumptions, the two central subspaces $S_{y|\mathbf{f}}$ and $S_{y|\hat{\mathbf{f}}}$ may not coincide (Li and Yin, 2007). Thus, the naive method by applying existing dimension reduction methods to the estimated factors $\hat{\mathbf{f}}_t$'s may not necessarily lead to the consistent estimation of $S_{y|\mathbf{f}}$. Fan et al. (2016) effectively solved this issue by developing an important

invariance result between $E(\mathbf{f}_t|y_{t+1})$ and $E(\widehat{\mathbf{f}}_t|y_{t+1})$. See Proposition 1 and Equation (2.6) of Fan et al. (2016). This invariance result provides an essential theoretic foundation for using sliced inverse regression (Li, 1991) under Models (1.1)–(1.2). The method and theory of Fan et al. (2016) required three assumptions:

- (A1) The number of factors K is a fixed constant as $p, T \rightarrow \infty$.
- (A2) The matrix $\Sigma_{f|y} = \text{cov}\{E(\mathbf{f}_t|y_{t+1})\}$ satisfies that $\phi_1' \Sigma_{f|y} \phi_1, \dots, \phi_L' \Sigma_{f|y} \phi_L$ are positive and distinct.
- (B1) Linearity condition (Li, 1991): $E(\mathbf{b}'\mathbf{f}_t|\phi_1'\mathbf{f}_t, \dots, \phi_L'\mathbf{f}_t)$ is a linear function of ϕ_1, \dots, ϕ_L for any $\mathbf{b} \in \mathbb{R}^K$.

A fixed K facilitates the accurate estimation of factors and loadings, but it also narrows our attention to a fixed dimensional factor space to forecast y_{t+1} . It is advocated that a diverging K may find a better balance between estimating factors and forecasting (Lam and Yao, 2012; Li et al., 2013; Jurado et al., 2015); in other words, should a growing K be used, the sufficient forecasting would deliver a potentially more powerful forecast. In the meantime, a diverging K would also relax the linearity condition (B1), which would greatly enhance the applicability of the forecasting method: as $\mathcal{S}_{y|f}$ is unknown, the condition is commonly strengthened to that it is satisfied for basis matrices of any L -dimensional subspace of \mathbb{R}^K , which equivalently requires \mathbf{f}_t to follow an elliptical distribution and can be restrictive; however, when L is much smaller than K , the low-dimensional projection $E(\mathbf{b}'\mathbf{f}_t|\phi_1'\mathbf{f}_t, \dots, \phi_L'\mathbf{f}_t)$ from a high-dimensional random vector \mathbf{f}_t will always approximate to a linear function under fairly general regularity conditions (Hall and Li, 1993), and the ellipticity of \mathbf{f}_t is no longer needed.

Assumption (A2) can be restrictive as well: one can easily verify that $\phi' \Sigma_{f|y} \phi$ is zero if the factors have an elliptical distribution and the forecast function $g(\cdot)$ is symmetric along direction ϕ of the factors. The latter occurs, for instance, when the forecast target was investigated using squared factors (Bai and Ng, 2008; Ludvigson and Ng, 2007). When (A2) fails, Fan et al. (2016) cannot detect all the sufficient forecasting directions and will lead to sub-optimal forecasting performance. Referring to the literature of sufficient dimension reduction mentioned above, a natural way to relax (A2) is to employ higher-order inverse regression methods, such as directional regression and the inverse third-moment method, in the sufficient dimension reduction stage.

In this work, we follow the same spirit as in Fan et al. (2016) to conduct factor analysis and sufficient dimension reduction sequentially based on models (1.1) and (1.2). We allow

the number of factors K to diverge as p and T grows, and employ directional regression and the inverse third-moment method for sufficient dimension reduction, which use higher-order inverse moments and exhaustively estimate $S_{y|\mathbf{f}}$ under weaker assumptions. The proposed method is applicable for generally distributed predictor with diverging number of factors, and is capable of detecting non-monotone effect of the factors on the response. Hence it is more applicable and effective than Fan et al. (2016) in many cases. During our theoretical development, we also propose an invariance result, which makes separate contribution to the literature of surrogate sufficient dimension reduction.

The rest of this paper is organized as follows. We first study directional regression in the sufficient forecasting in Section 2, including the factor analysis in Subsection 2.1, an invariance result for sufficient dimension reduction in Subsection 2.2, the details of implementation in Subsection 2.3, the asymptotic results in Subsection 2.4, and a Bayesian information criterion (BIC) to select the dimension L of the central subspace in Subsection 2.5. In Section 3, we further incorporate the inverse third-moment method in the sufficient forecasting, and develop the corresponding theoretical results. Section 4 is devoted to the simulation studies and a real data example that illustrates the power of the proposed method. We leave all the proofs to Section 5.

2 Forecasting with directional regression

2.1 Factor analysis

To make forecast, we need to estimate the factor loadings \mathbf{B} and the error covariance matrix Σ_u . For ease of presentation, we first assume that the number of underlying factors K is growing as $p, T \rightarrow \infty$ but known. Consider the following constrained least squares problem:

$$\begin{aligned} (\widehat{\mathbf{B}}_K, \widehat{\mathbf{F}}_K) &= \arg \min_{(\mathbf{B}, \mathbf{F})} \|\mathbf{X} - \mathbf{B}\mathbf{F}'\|_F^2, \\ &\text{subject to } T^{-1}\mathbf{F}'\mathbf{F} = \mathbf{I}_K, \quad \mathbf{B}'\mathbf{B} \text{ is diagonal,} \end{aligned} \tag{2.1}$$

where $\mathbf{X} = (\mathbf{x}_1, \dots, \mathbf{x}_T)$, and $\|\cdot\|_F$ denotes the Frobenius norm of a matrix. The constraints $T^{-1}\mathbf{F}'\mathbf{F} = \mathbf{I}_K$ and that $\mathbf{B}'\mathbf{B}$ is diagonal is to address the issue of identifiability during the minimization. As they can always be satisfied for any $\mathbf{B}\mathbf{F}'$ after appropriate matrix operations on \mathbf{B} and \mathbf{F} , they impose no additional restrictions on the factor model (1.2).

It is a commonly known fact that the minimizers $\widehat{\mathbf{F}}_K$ and $\widehat{\mathbf{B}}_K$ of (2.1) are such that the columns of $\widehat{\mathbf{F}}_K/\sqrt{T}$ are the eigenvectors corresponding to the K largest eigenvalues of the $T \times T$ matrix $\mathbf{X}'\mathbf{X}$ and $\widehat{\mathbf{B}}_K = T^{-1}\mathbf{X}\widehat{\mathbf{F}}_K$. To simplify notation, let $\widehat{\mathbf{B}} = \widehat{\mathbf{B}}_K$ and $\widehat{\mathbf{F}} = \widehat{\mathbf{F}}_K$.

As both the dimension p of the predictor x_t and the number of factors K are diverging, it is necessary to regulate the magnitude of the factor loadings \mathbf{B} and the idiosyncratic error \mathbf{u}_t , so that the latter is negligible with respect to the former. We should also regulate the stationarity of the time series. In this paper, we adopt the following assumptions. For simplicity in notation, we let $\mathbf{B} = (\mathbf{b}_1, \dots, \mathbf{b}_p)'$, and $\|\mathbf{B}\|_{\max}$ be the maximum of the absolute values of all the entries in \mathbf{B} . Let \mathcal{F}_∞^0 and \mathcal{F}_T^∞ denote the σ -algebras generated by $\{(\mathbf{f}_t, \mathbf{u}_t, \epsilon_{t+1}) : t \leq 0\}$ and $\{(\mathbf{f}_t, \mathbf{u}_t, \epsilon_{t+1}) : t \geq T\}$ respectively.

Assumption 2.1 (Factors and Loadings).

(1) *There exists $b > 0$ such that $\sup_{p \in \mathbb{N}} \|\mathbf{B}\|_{\max} \leq b$, and there exist two positive constants c_1 and c_2 such that*

$$c_1 < p^{-1}\lambda_{\min}(\mathbf{B}'\mathbf{B}) < p^{-1}\lambda_{\max}(\mathbf{B}'\mathbf{B}) < c_2.$$

(2) *Identification: $T^{-1}\mathbf{F}'\mathbf{F} = \mathbf{I}_K$, and $\mathbf{B}'\mathbf{B}$ is a diagonal matrix with distinct entries.*

Assumption 2.2 (Data Generating Process). $\{\mathbf{f}_t\}_{t \geq 1}$, $\{\mathbf{u}_t\}_{t \geq 1}$ and $\{\epsilon_{t+1}\}_{t \geq 1}$ are three independent groups, and all of them are strictly stationary. The factor process satisfies that both $\{K^{-2}E\|\mathbf{f}_t\|^4 : p \in \mathbb{N}\}$ and $\{K^{-1}E(\|\mathbf{f}_t\|^2|y_{t+1}) : p \in \mathbb{N}\}$ are bounded sequences. In addition, $\alpha(T) = \sup_{A \in \mathcal{F}_\infty^0, B \in \mathcal{F}_T^\infty} |P(A)P(B) - P(AB)| < c\rho^T$ for $T \in \mathbb{Z}^+$ and some $\rho \in (0, 1)$.

Assumption 2.3 (Residuals and Dependence). *There exists a positive constant $M < \infty$ that does not depend on p or T , such that*

- (1) $E(\mathbf{u}_t) = \mathbf{0}$, and $E|u_{it}|^8 \leq M$.
- (2) $\|\boldsymbol{\Sigma}_u\|_1 \leq M$, and for every $i, j, t, s > 0$, $(pT)^{-1} \sum_{i,j,t,s} |E(u_{it}u_{js})| \leq M$
- (3) For every (t, s) , $E|p^{-1/2}(\mathbf{u}'_s\mathbf{u}_t - E(\mathbf{u}'_s\mathbf{u}_t))|^4 \leq M$.

Assumption 2.1 regulates the signal strength of the factors contained in the predictor through the order of the factor loadings, and Assumption 2.3 regulates the order of the idiosyncratic errors contained in the predictor. Together, they ensure that the former dominates the latter in the population level as p grows. Assumption 2.2 implies that the sample observations are only weakly dependent, so that the estimation accuracy grows with the sample size.

Under these assumptions, we have the following consistency result for estimating the factor loadings. Instead of the Frobenius norm used in (2.1), we use the spectral norm to

measure the magnitude of a matrix, defined as $\|\mathbf{A}\| = \lambda_{\max}^{1/2}(\mathbf{A}'\mathbf{A})$, the square root of the largest eigenvalue of $\mathbf{A}'\mathbf{A}$, for any matrix \mathbf{A} .

Theorem 2.1. *Suppose $K = o(\min\{p^{1/3}, T\})$. Let $\mathbf{\Lambda}_b = (\mathbf{B}'\mathbf{B})^{-1}\mathbf{B}'$ and $\widehat{\mathbf{\Lambda}}_b = (\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1}\widehat{\mathbf{B}}'$. Under Assumptions 2.1, 2.2 and 2.3, we have*

$$1) \|\widehat{\mathbf{B}} - \mathbf{B}\| = O_p(p^{1/2}(K^{3/2}p^{-1/2} + K^{1/2}T^{-1/2})),$$

$$2) \|\widehat{\mathbf{\Lambda}}_b - \mathbf{\Lambda}_b\| = O_p(p^{-1/2}(K^{3/2}p^{-1/2} + K^{1/2}T^{-1/2})).$$

Because the dimension p of the factor loadings \mathbf{B} is diverging, the estimation error $\widehat{\mathbf{B}} - \mathbf{B}$ accumulates as p grows. For a p -dimensional vector whose entries are constantly one, its spectral norm is $p^{1/2}$, which diverges to infinity. Thus, we should treat $p^{1/2}$ as the unit magnitude of the spectral norm of matrices with p rows, in which sense the statement 1) of Theorem 2.1 justifies the estimation consistency of the factor loadings \mathbf{B} . As the error term \mathbf{u}_t shrinks as p grows under Assumption 2.3, the convergence order of the factor loading estimation largely depends on p - a higher dimensional predictor means a more accurate estimation. The convergence order in this theorem can be further improved if we impose stronger assumptions on the negligibility of the error terms in the factor model (1.2).

Given $\widehat{\mathbf{B}}$, it is easy to see that $\widehat{\mathbf{f}}_t = \widehat{\mathbf{\Lambda}}_b\mathbf{B}\mathbf{f}_t + \widehat{\mathbf{\Lambda}}_b\mathbf{u}_t$. Thus, together with the negligibility of the error term \mathbf{u}_t , the consistency of $\widehat{\mathbf{B}}$ and $\widehat{\mathbf{\Lambda}}_b$ indicates the closeness between the true factors \mathbf{f}_t and the estimated factors $\widehat{\mathbf{f}}_t$, of which the latter will be used in the subsequent sufficient dimension reduction. The error covariance matrix $\mathbf{\Sigma}_u$ can be estimated by thresholding the sample covariance matrix of the estimated residual $x_t - \widehat{\mathbf{B}}\widehat{\mathbf{f}}_t$, denoted by $\widehat{\mathbf{\Sigma}}_u = (\widehat{\sigma}_{ij}^u)_{p \times p}$, as in Cai and Liu (2011), Xue et al. (2012), Fan et al. (2013) and Fan et al. (2016).

2.2 An invariance result

Using the estimated factors from factor analysis, we now apply directional regression to further produce a lower-dimensional sufficient predictor for forecasting. Before digging into more details about the estimation consistency, we focus on the population level, and temporarily assume an oracle scenario where \mathbf{B} is known a priori. This scenario simplifies the discussion, as it eliminates the estimation error introduced by estimating the factor loadings. We will return to the realistic case afterwards.

As pointed out in Fan et al. (2016), the inverse regression methods are not readily applicable to estimate the central subspace $\mathcal{S}_{y|\mathbf{f}}$, for the reason that the estimated factors $\widehat{\mathbf{f}}_t$

always contain an error term aside from the true factors \mathbf{f}_t , which can be asymptotically non-negligible in certain settings. To see this point, we apply an ordinary least square estimation in (1.2), and have

$$\widehat{\mathbf{f}}_t = \mathbf{f}_t + \mathbf{u}_t^*, \quad (2.2)$$

where $\mathbf{u}_t^* = \mathbf{\Lambda}_b \mathbf{u}_t$. Then \mathbf{u}_t^* is the price that we pay for the contamination of the original predictor \mathbf{x}_t by the error term \mathbf{u}_t . Such a price is intrinsic as it is inevitable whatever estimators of \mathbf{f}_t are used. The consequence is two-fold. First, as no distributional assumption is imposed on \mathbf{u}_t^* , the regularity conditions on $\widehat{\mathbf{f}}_t$ such as the linearity condition (B1) may fail, which causes inconsistency of the inverse regression methods in estimating the corresponding central subspace $\mathcal{S}_{y|\widehat{\mathbf{f}}}$. Second, even if $\mathcal{S}_{y|\widehat{\mathbf{f}}}$ can be estimated consistently, it needs not coincide with the central subspace $\mathcal{S}_{y|\mathbf{f}}$ of interest.

To address this issue, one may naturally search for suitable conditions that ensure the coincidence between the two central subspaces $\mathcal{S}_{y|\widehat{\mathbf{f}}}$ and $\mathcal{S}_{y|\mathbf{f}}$, which is equivalent to that the two spaces have equal dimension L and that any basis matrix (ϕ_1, \dots, ϕ_L) of $\mathcal{S}_{y|\mathbf{f}}$ satisfies

$$y_{t+1} \perp (\mathbf{f}_t + \mathbf{u}_t^*) \mid (\phi_1, \dots, \phi_L)'(\mathbf{f}_t + \mathbf{u}_t^*).$$

Such a study can be embedded in surrogate sufficient dimension reduction (Li and Yin, 2007), which aims to conduct sufficient dimension reduction when the predictor is contaminated by a measurement error. In particular, a direct application of Theorem 3.1 in Li and Yin (2007) implies that the coincidence of the central subspaces holds if both \mathbf{f}_t and \mathbf{u}_t^* are normally distributed, subject to the independence between \mathbf{f}_t and \mathbf{u}_t^* that we have already assumed. In that case, the resulting normality of $\widehat{\mathbf{f}}_t$ also makes $\mathcal{S}_{y|\widehat{\mathbf{f}}}$ estimable by the inverse regression methods like sliced inverse regression. Hence the central subspace $\mathcal{S}_{y|\mathbf{f}}$ of interest can be consistently estimated.

However, the normality of \mathbf{u}_t^* adopted in Li and Yin (2007) can be easily violated in practice, in which case the coincidence between the central subspaces becomes infeasible. It is important to notice that $\mathcal{S}_{y|\mathbf{f}}$ is the only parameter of interest, and its coincidence with $\mathcal{S}_{y|\widehat{\mathbf{f}}}$ is needed only when the latter serves as the intermediate parameter in the estimation procedure. Consequently, such coincidence can be relaxed if we manage to find other intermediate parameters, which naturally leads us to consider the inverse regression methods.

In all the inverse regression methods, the central subspace is characterized as the column space of certain positive semi-definite matrix parameters, called the kernel matrices. If we can manage to estimate the kernel matrices using the estimated factors $\widehat{\mathbf{f}}_t$ in place of \mathbf{f}_t ,

then so too is $\mathcal{S}_{y|\mathbf{f}}$. Naturally, this can be realized if we adopt suitable conditions on the predictor x_t , so that the kernel matrices are invariant of the change from \mathbf{f}_t to $\widehat{\mathbf{f}}_t$. Because the kernel matrices are constructed only by the inverse moments, rather than the entire joint distribution, we expect such conditions to be weaker than those required for the coincidence of the central subspaces. This point is demonstrated in the following theorem for directional regression.

Theorem 2.2. *Under model (1.2), the kernel matrix for directional regression,*

$$\mathbf{M}_{dr} = E\{2\text{var}(\mathbf{f}_t) - E[(\mathbf{f}_t - \mathbf{f}_s)(\mathbf{f}_t - \mathbf{f}_s)'|y_{t+1}, y_{s+1}]\}^2, \quad (2.3)$$

where (\mathbf{f}_s, y_{s+1}) is an independent copy of (\mathbf{f}_t, y_{t+1}) , is invariant if \mathbf{f}_t and \mathbf{f}_s are replaced by $\widehat{\mathbf{f}}_t$ and $\widehat{\mathbf{f}}_s$ in (2.3), respectively.

Because the true factors \mathbf{f}_t has identity covariance matrix, the form of the kernel matrix adopted in the theorem coincides with its original form in Li and Wang (2007). However, it does make a modification on the latter when the estimated factors $\widehat{\mathbf{f}}_t$ are used instead, which no longer have identity covariance matrix in the population level. From the proof of the theorem, one can easily see that such a modification is crucial, as it removes the effect of the estimation error \mathbf{u}_t^* from the kernel matrix estimation.

The coincidence between the column space of \mathbf{M}_{dr} and the central subspace $\mathcal{S}_{y|\mathbf{f}}$ requires both the linearity condition (B1) and the constant variance condition:

(B2) $\text{var}(\mathbf{f}_t | \phi_1' \mathbf{f}_t, \dots, \phi_L' \mathbf{f}_t)$ is degenerate.

Since the central subspace $\mathcal{S}_{y|\mathbf{f}}$ is unknown, same as (B1), (B2) is commonly strengthened to that it is satisfied for basis matrices of any L -dimensional subspace of \mathbb{R}^K . The strengthened conditions equivalently require the factors to be jointly normally distributed, which are again restrictive in practice. If one treat \mathbf{f}_t as the response and $(\phi_1' \mathbf{f}_t, \dots, \phi_L' \mathbf{f}_t)$ as the predictor in regression, then (B1) is the linearity assumption on the regression function and (B2) is the homoscedasticity assumption on the error term. In this sense, we follow the convention in the literature of regression to treat (B2) less worrisome than (B1) in practice.

A similar invariance result to Theorem 2.2 has been developed in Fan, Xue and Yao (2016) for sliced inverse regression where the inverse first moment is involved (see their equation (2.6)). As \mathbf{u}_t is allowed to be non-normally distributed, according to Li and Yin (2007), the two central subspaces $\mathcal{S}_{y|\widehat{\mathbf{f}}}$ and $\mathcal{S}_{y|\mathbf{f}}$ may differ, which means that the former may not be recovered by the corresponding kernel matrix in directional regression. This can also be

explained by the fact that when $\widehat{\mathbf{f}}_t$ is used as the predictor, the linearity condition (B1) and the constant variance condition (B2) are violated due to the arbitrariness of the distribution of \mathbf{u}_t^* , which makes directional regression inconsistent.

One can always argue that under the assumption of negligible error term \mathbf{u}_t^* relative to the true factors \mathbf{f} , the estimated factors $\widehat{\mathbf{f}}$ approximate to \mathbf{f} , which suggests the corresponding approximation between the central subspaces $\mathcal{S}_{y|\widehat{\mathbf{f}}}$ and $\mathcal{S}_{y|\mathbf{f}}$. Consequently, the central subspace of interest $\mathcal{S}_{y|\mathbf{f}}$ can be still consistently estimated through the intermediate parameter $\mathcal{S}_{y|\widehat{\mathbf{f}}}$, without using the invariance result. However, as the invariance result justifies the Fisher consistency of directional regression under the existence of potentially non-negligible measurement error \mathbf{u} , it is useful in more general settings. These settings include, for example, the case when the primary statistical interest is on estimating which linear combinations of x affect y , i.e. the factor loadings \mathbf{B} and the central subspace $\mathcal{S}_{y|\mathbf{f}}$, rather than the forecast function g . In this sense, this result itself makes an independent contribution to the literature of surrogate sufficient dimension reduction.

In reality, the hypothetical independent copies (\mathbf{f}_s, y_{s+1}) and (\mathbf{f}_t, y_{t+1}) do not exist in the observed data. Consequently, we estimate \mathbf{M}_{dr} using its equivalent form, which is derived by expanding (2.3),

$$\begin{aligned} \mathbf{M}_{dr} = & 2E\{\text{var}(\mathbf{f}_t) - E(\mathbf{f}_t\mathbf{f}'_t|y_{t+1})\} + 2E^2[E(\mathbf{f}_t|y_{t+1})E'(\mathbf{f}_t|y_{t+1})] \\ & + 2E[E'(\mathbf{f}_t|y_{t+1})E(\mathbf{f}_t|y_{t+1})] \cdot E[E(\mathbf{f}_t|y_{t+1})E'(\mathbf{f}_t|y_{t+1})]. \end{aligned} \quad (2.4)$$

The marginal covariance matrix $\text{var}(\widehat{\mathbf{f}}_t)$ and the conditional covariance matrix $E(\widehat{\mathbf{f}}_t\widehat{\mathbf{f}}'_t|y_{t+1})$ can be easily estimated by replacing \mathbf{B} with its estimate $\widehat{\mathbf{B}}$, slicing the support of y_{t+1} , and using the sample moments. By Theorem 2.1, the factor loadings B are consistently estimated by $\widehat{\mathbf{B}}$, so it is plausible that under suitable moment conditions and sufficiently large sample size, the leading eigenvectors of the resulting matrix span a consistent estimator of the central subspace $\mathcal{S}_{y|\mathbf{f}}$. In Subsection 2.4, we will justify the sufficiency of Assumptions 2.1 - 2.3 for such consistency, and give the corresponding convergence order of the estimation.

2.3 Implementation

In the literature of sufficient dimension reduction, it has been a common practice to estimate the inverse moments in the inverse regression methods using the slicing technique; that is, we partition the sample of y_{t+1} into H slices with equal sample proportion, and estimate the moments of the factors within each slice. In the population level, it corresponds

to partitioning the support of y_{t+1} into H slices, i.e. intervals, with equal probability, and using the slice indicator, denoted by y_{t+1}^D , as the new working response variable.

The slicing technique substantially simplifies the implementation of the inverse regression methods. Because the slice indicator y_{t+1}^D is a measurable function of the original response y_{t+1} , \mathbf{f}_t must affect y_{t+1}^D through y_{t+1} . Thus, the working central subspace $\mathcal{S}_{y^D|\mathbf{f}}$ is always a subspace of the central subspace of interest $\mathcal{S}_{y|\mathbf{f}}$, which means that no redundant directions of \mathbf{f}_t will be selected in the estimation.

Using the slice indicator y_{t+1}^D , the inverse moments $E(\widehat{\mathbf{f}}_t|y_{t+1}^D)$ and $E(\widehat{\mathbf{f}}_t\widehat{\mathbf{f}}_t'|y_{t+1}^D)$ can be easily estimated by the usual sample moments. As constrained in factor analysis, $\widehat{\mathbf{f}}_t$ has sample variance equal to \mathbf{I}_K , which we use to estimate its population variance $\text{var}(\widehat{\mathbf{f}}_t)$. Alternatively, one can also use the restriction that $\text{var}(\mathbf{f}_t) = \mathbf{I}_K$ to estimate $\text{var}(\widehat{\mathbf{f}}_t)$ by $\mathbf{I}_K + \widehat{\Sigma}_{\mathbf{u}^*}$, where $\widehat{\Sigma}_{\mathbf{u}^*}$ is the thresholding covariance estimator. An omitted simulation study shows that the two estimators perform similarly to each other, so we choose the former for simplicity. The kernel matrix estimator $\widehat{\mathbf{M}}_{dr}$, whose leading eigenvectors span an estimate of the central subspace, is then given by (2.3). In summary, the proposed estimator can be implemented using the following steps:

Algorithm 1 Forecasting with directional regression using factor models

- Step 1: Estimate the factor loadings \mathbf{B} and the factors \mathbf{f}_t by (2.1).
 - Step 2: For $i = 0, \dots, H$, let $y_{(i)/H}$ be the (i/H) th quantile of $\{y_1, \dots, y_T\}$, where $y_{(0)}$ is defined to be a constant that is less than the infimum of $\{y_1, \dots, y_T\}$. Let $y_{t+1}^D = i$ if $y_{t+1} \in (y_{(i)/H}, y_{(i+1)/H}]$, and $c_i = \sum_{t=1}^T I(y_{t+1} = i)$ be the number of observations in the i th slice of y_{t+1}^D . Estimate $E(\widehat{\mathbf{f}}_t|y_{t+1}^D = i)$ by $\sum_{l=1}^{c_i} \widehat{\mathbf{f}}_{i,l}/c_i$ and $E(\widehat{\mathbf{f}}_t\widehat{\mathbf{f}}_t'|y_{t+1}^D = i)$ by $\sum_{l=1}^{c_i} \widehat{\mathbf{f}}_{i,l}\widehat{\mathbf{f}}_{i,l}'/c_i$, where the subscript (i, l) corresponds to the l th observation in the i th slice of y_{t+1}^D . Estimate $\text{var}(\widehat{\mathbf{f}}_t)$ by \mathbf{I}_K .
 - Step 3: Estimate \mathbf{M}_{dr} by plugging the estimates in Step 2 into (2.4). Let $\widehat{\mathbf{M}}_{dr}$ be the resulting estimate. Estimate $\mathcal{S}_{y|\mathbf{f}}$ by the L eigenvectors of $\widehat{\mathbf{M}}_{dr}$ corresponding to the largest eigenvalues.
 - Step 4: Estimate $g(\cdot)$ with indices from Step 3, and forecast y_{t+1} .
-

An omitted simulation study shows that our estimate is robust to the choice of H , as long as the latter falls into a reasonable range, say three to ten. This phenomenon has also

been observed by multiple authors; see, for example, Li (1991) and Li and Wang (2007).

2.4 Asymptotic properties

For simplicity of the presentation, in this subsection, we assume both the dimension L of the central subspace $\mathcal{S}_{y|f}$ and the number of factors K to be known a priori, where the latter is a diverging sequence. The same asymptotic result can be developed similarly if K and L are unknown but consistently estimated. Consistent determination of K and L will be discussed later in Subsection 2.5, under which the result developed here can still be applied.

We first introduce some elementary result about the consistency of the eigen-decomposition of random matrices. The following concept characterizes a sequence of non-negligible random variables.

Definition 2.1. *A sequence of random variables $\{Z_T : T \in \mathbb{N}\}$ is called bounded below from 0 in probability, and written as $Z_T = O_P^+(1)$, if there exists a constant $c > 0$ such that $P(Z_T > c) \rightarrow 1$ as $T \rightarrow \infty$.*

This concept is a natural generalization of non-stochastic sequences that are bounded below from zero to a probabilistic version, much like the generalization from $O(1)$ to $O_P(1)$. In particular, it includes these non-stochastic sequences as a special case. For convenience in notations, we denote these non-stochastic sequence also by $O_P^+(1)$, if no ambiguity is caused. Another simple example of $O_P^+(1)$ is $c - \delta_T$, where c is an arbitrary positive constant and δ_T is an arbitrary sequence of random variables such that $\delta_T = o_P(1)$. Using this concept, the following result shows when the leading eigenvectors of a sequence of random matrices span a converging linear space. For any symmetric matrix A , we denote its smallest eigenvalue by $\lambda_{\min}(A)$.

Lemma 2.1. *Let $\{\widehat{\mathbf{M}}_T \in \mathbb{R}^{K \times K} : T \in \mathbb{N}\}$ be a sequence of symmetric random matrices and $U \in \mathbb{R}^{K \times L}$ and $V \in \mathbb{R}^{K \times (K-L)}$ be an orthonormal basis of \mathbb{R}^K . If*

$$(a) \lambda_{\min}(U' \widehat{\mathbf{M}}_T U) = O_P^+(1),$$

$$(b) \text{ there exists a non-stochastic sequence } \{c_T : T \in \mathbb{N}\} \text{ with } c_T \rightarrow 0, \text{ such that } \|U' \widehat{\mathbf{M}}_T V\| = O_P(c_T) \text{ and } \|V' \widehat{\mathbf{M}}_T V\| = O_P(c_T),$$

then the linear space spanned by the L leading eigenvectors of $\widehat{\mathbf{M}}_T$, denoted by \widehat{U}_T , consistently estimates the linear space spanned by U in the sense that the projection matrix of the former converges to that of the latter in probability; that is, $\|\widehat{U}_T \widehat{U}_T' - U U'\|_F = o_P(c_T)$.

If we let $\widehat{\mathbf{M}}_T$ be the sample kernel matrix $\widehat{\mathbf{M}}_{dr}$ of directional regression, and U be an orthonormal basis of the central subspace, then condition (a) of the lemma means that $\lambda_{\min}(U'\widehat{\mathbf{M}}_{dr}U) = O_P^+(1)$, which requires the rank of the kernel matrix \mathbf{M}_{dr} to be at least L , or equivalently, exhaustive estimation of the central subspace $\mathcal{S}_{y|\mathbf{f}}$ by directional regression. Based on Theorem 3 in Li and Wang (2007), the following result gives the corresponding sufficient conditions on the inverse moments.

Theorem 2.3. *The following two assumptions are equivalent:*

- a. *for any sequence of non-stochastic vectors $\{v_T \in \mathcal{S}_{y|\mathbf{f}} : \|v_T\| = 1\}$ and an independent copy (\mathbf{f}_s, y_{s+1}) of (\mathbf{f}_t, y_{t+1}) ,*

$$\text{var}[E\{v_T'(\mathbf{f}_t - \mathbf{f}_s)\}^2 | y_{t+1}, y_{s+1}]] = O_P^+(1).$$

- b. *For any sequence of non-stochastic vectors $\{v_T \in \mathcal{S}_{y|\mathbf{f}} : \|v_T\| = 1\}$,*

$$\max \{ \text{var}[E(v_T' \mathbf{f}_t | y_{t+1})], \text{var}[E\{(v_T' \mathbf{f}_t)^2 | y_{t+1}\}] \} = O_P^+(1).$$

Moreover, they imply the exhaustiveness of directional regression. That is, $\lambda_L(\mathbf{M}_{dr})$, the L th eigenvalue of \mathbf{M}_{dr} , is $O_P^+(1)$.

The statements in this theorem require that all the directions in the central subspace are captured in the first two inverse moments, $E(\mathbf{f}_t | y_{t+1})$ and $E(\mathbf{f}_t \mathbf{f}_t' | y_{t+1})$, which is satisfied, for example, when $\mathbf{f}_t | y_{t+1}$ is normally distributed. See also Cook and Lee (1999) for more details. Although one can always construct specific models in which the effect of \mathbf{f}_t is revealed only in higher-order inverse moments, the exhaustiveness of directional regression has been commonly recognized in applications. A detailed justification can also be found in Li and Wang (2007).

Because we estimate a sequence of central subspaces with diverging dimensions, in addition to the exhaustiveness of directional regression at each dimension K , we further require in this theorem that the weakest signal strength in the central subspace not to vanish as K grows. The condition can be relaxed if we conduct a more careful study about the order restriction in Lemma 2.1 and allow a slower convergence rate in estimating the central subspace. But considering the fact that L is fixed and the popularity of similar conditions in the literature of high-dimensional data analysis, we decide to adopt it here, and leave further relaxation to future work.

Theorem 2.4. *Under Assumptions 2.1, 2.2 and 2.3, the assumption in Theorem 2.3, the linearity condition (B1), and the constant variance condition (B2), if $K = o(\min(p^{1/3}, T^{1/2}))$, then the leading L eigenvectors of $\widehat{\mathbf{M}}_{dr}$, denoted by $\widehat{\boldsymbol{\phi}}_1, \dots, \widehat{\boldsymbol{\phi}}_L$, span a consistent estimator of the central subspace in the sense that*

$$\|(\widehat{\boldsymbol{\phi}}_1, \dots, \widehat{\boldsymbol{\phi}}_L)(\widehat{\boldsymbol{\phi}}_1, \dots, \widehat{\boldsymbol{\phi}}_L)' - (\boldsymbol{\phi}_1, \dots, \boldsymbol{\phi}_L)(\boldsymbol{\phi}_1, \dots, \boldsymbol{\phi}_L)'\|_F = O_P(K^{3/2}p^{-1/2} + KT^{-1/2}).$$

In connection with Theorem 2.1, the estimation error in sufficient dimension reduction, as justified in this theorem, can be divided into two parts. The first part, which is of order $O_P(K^{3/2}p^{-1/2})$, is inherited from factor analysis. This part represents the price we pay for estimating the factor loadings \mathbf{B} , and it depends on the dimension p of the original predictor. By contrast, the second part, which is of order $O_P(KT^{-1/2})$, does not depend on p and is newly generated in the sufficient dimension reduction stage. As can be easily seen from the proof of the theorem, it represents the price we pay for estimating the unknown inverse second moment involved in the kernel matrix. Therefore, this part would persist even if no error were generated in factor analysis.

2.5 Model selection

We now discuss how to determine the number of factors K and the dimension L of the central subspace $\mathcal{S}_{y|f}$. The problem is commonly called order determination in the literature of dimension reduction (Luo and Li, 2016).

The correct specification of the number of factors, K , is a fundamental issue of large factor models. Various consistent order-determination approaches have been established under the setting that K is fixed; see Bai and Ng (2008); Onatski (2010); Ahn and Horenstein (2013), etc. However, a growing number of empirical studies suggest that the number of factors may increase as the cross-sectional dimension N or time-series dimension T increases; see Ludvigson and Ng (2009) and Jurado et al. (2015), whose estimate of the number of factors explaining certain macroeconomic time series ranges from 1 to 10. Recently, Li et al. (2013) extended the analysis of Bai and Ng (2008) to estimate the number of factors that may increase with the cross-sectional size and time period, and prove the consistency of a modified procedure of Bai and Ng (2008). Specifically, it estimates K by

$$\hat{K} = \arg \min_{0 \leq K \leq K_{max}} \log\left(\frac{1}{pT} \|\mathbf{X} - T^{-1} \mathbf{X} \widehat{\mathbf{F}}_K \widehat{\mathbf{F}}_K'\|_F^2\right) + Kg(p, T),$$

where K_{max} is a prescribed upper bound that possibly increases with T and $\widehat{\mathbf{F}}_K$ corresponds to the solution to (2.1). $g(p, T)$ is a penalty function such that $g(p, T) = o(1)$ and $C_{p, T, K_{max}}^{-1} g(p, T) \rightarrow \infty$, where $C_{p, T, K_{max}} = O_p(\max\{K_{max}^6/p, K_{max}^4/T\})$. The specific choice of $g(p, T)$ does not affect asymptotic results. One example suggested by Bai and Ng (2008) as well as Li et al. (2013) is

$$g(p, T) = \frac{p + T}{pT} \log\left(\frac{pT}{p + T}\right).$$

Under Assumptions 2.1–2.3 and letting $K_{max} = o_p(\min\{p^{1/16}, T^{1/14}\})$, Li et al. (2013) showed that the data-driven \widehat{K} is a consistent estimator of K .

Next we discuss the choice of L . In the sufficient dimension reduction literature, multiple methods have been proposed to determine the dimension of the central subspace, including the sequential tests (Li 1991, Li and Wang 2007), the permutation test (Cook and Weisberg 1991), the bootstrap procedure (Ye and Weiss, 2003), the cross-validation method (Xia et al., 2002; Wang and Xia, 2008), the BIC type procedure (Zhu, Miao and Peng 2006), and the ladle estimator (Luo and Li, 2016). Among them, the BIC type procedure can be easily implemented in conjunction with the inverse regression methods, and reaches the desired consistency in high-dimensional cases. For a K -dimensional positive semi-definite matrix parameter \mathbf{M} of rank L and its sample estimator $\widehat{\mathbf{M}}$, let $\{\lambda_1, \dots, \lambda_K\}$ and $\{\widehat{\lambda}_1, \dots, \widehat{\lambda}_K\}$ be their eigenvalues in the descending order, respectively. We set a censoring constant $c \in (0, 1)$ that is invariant of K , and define the objective function $G : \{1, \dots, K_c\} \rightarrow \mathbb{R}$ to be

$$G(l) = (T/2) \sum_{i=1+\min(\tau, l)}^{K_c} \{\log(\widehat{\lambda}_i + 1) - \widehat{\lambda}_i\} - C_T l(2K - l + 1)/2, \quad (2.5)$$

in which K_c is nearest integer to cK and τ is the number of $\widehat{\lambda}_i$'s that are greater than zero. We then estimate L as the maximizer \widehat{L} of $G(\cdot)$. Compared with the original BIC type procedure in Zhu, Miao and Peng (2006), we introduce the censoring constant c to restrict the range of candidate dimensions. Because the number of factors K is diverging while the dimension of the central subspace L is fixed, this restriction is reasonable for all large samples. The restriction is indeed crucial for the consistency of the order-determination procedure revealed in the following theorem without introducing additional constraints on the order of K and $\|\widehat{\mathbf{M}} - \mathbf{M}\|$, which improves the original result in Zhu, Miao and Peng (2006).

Theorem 2.5. *Suppose $\lambda_L = O_P^+(1)$ and $\|\widehat{\mathbf{M}} - \mathbf{M}\| = o_P(1)$. If C_T satisfies that*

$$C_T K T^{-1} \rightarrow 0 \text{ and } \|\widehat{\mathbf{M}} - \mathbf{M}\|^2 = o_P(C_T K T^{-1}),$$

then \widehat{L} converges to L in probability.

A candidate of C_T can be $K^{-1}T\|\widehat{\mathbf{M}} - \mathbf{M}\|$. Referring to Theorem 2.4, if we apply the BIC-type procedure in conjunction with directional regression to detect the dimension of the central subspace $\mathcal{S}_{y|\mathbf{f}}$, then we can choose C_T to be $K^{1/2}p^{-1/2}T + T^{1/2}$. To further polish the procedure, we can incorporate a multiplicative constant in C_T , and tune its value in a data-driven manner such as cross-validation.

3 Sufficient forecasting with the inverse third moment

As mentioned in the Introduction, Yin and Cook (2003) employed the inverse third moment in their kernel matrix. In our context, the inverse third moment is

$$\mu_0^3(\mathbf{f}_t|y_{t+1}) = E[\{\mathbf{f}_t - E(\mathbf{f}_t|y_{t+1})\} \otimes \{\mathbf{f}_t - E(\mathbf{f}_t|y_{t+1})\} \{\mathbf{f}_t - E(\mathbf{f}_t|y_{t+1})\}' | y_{t+1}], \quad (3.1)$$

which can be treated as a $K^2 \times K$ matrix that contains $K(K+1)/2$ distinct rows. Let μ_t be the sub-matrix of $\mu_0^3(\mathbf{f}_t|y_{t+1})$ that contains these distinct rows. The kernel matrix \mathbf{M}_{tm} of the inverse third-moment method, where the subscript stands for ‘‘third moment’’, is $E(\mu_t' \mu_t)$. For the column space of \mathbf{M}_{tm} to be a subspace of the central subspace $\mathcal{S}_{y|\mathbf{f}}$, in addition to the linearity condition (B1) and the constant variance condition (B2), the distribution of the factors must also satisfy the symmetry condition (Yin and Cook, 2003):

$$(B3) \quad E(\mathbf{f}_t \otimes \mathbf{f}_t \mathbf{f}_t' | \phi_1' \mathbf{f}_t, \dots, \phi_L' \mathbf{f}_t) = 0.$$

Same as the linearity condition (B1) and the constant variance condition (B2), the symmetry condition (B3) is also satisfied when \mathbf{f}_t is normally distributed. Referring to the discussion below (B2), if we treat \mathbf{f}_t as the response and $(\phi_1' \mathbf{f}_t, \dots, \phi_L' \mathbf{f}_t)$ as the predictor in regression, then (B3) is implied by the symmetry of the error term, which is frequently adopted in the literature of regression. Thus the condition can be treated fairly general in practice.

Compared with directional regression, the inverse third-moment method incorporates higher-order inverse moments, so it captures additional information about the inverse conditional distribution $\mathbf{f}_t|y_{t+1}$. As justified in the following theorem, under the symmetry

condition (B3), it can serve as a useful complement to directional regression, if the latter fails to be exhaustive.

Theorem 3.1. *Suppose that the linearity condition (B1), the constant variance condition (B2), and the symmetry condition (B3) are satisfied. If, for any sequence of non-stochastic vectors $\{v_T \in \mathcal{S}_{y|\mathbf{f}} : \|v_T\| = 1\}$,*

$$E^2[\{v'_T \mathbf{f}_t - E(v'_T \mathbf{f}_t | y_{t+1})\}^3 | y_{t+1}] = O_P^+(1),$$

then the inverse third-moment method is exhaustive in the sense that $\lambda_L(\mathbf{M}_{tm})$, the L th eigenvalue of \mathbf{M}_{tm} , is $O_P^+(1)$.

As the inverse first moment is excluded from the kernel matrix, the inverse third-moment method cannot capture the corresponding information. In addition, it will also miss the directions of the factors that are associated with the response in a symmetric pattern, if any. Hence, it may fail to be exhaustive in applications. Because directional regression effectively uses the first two inverse moments and can detect symmetric pattern between the factors and the response, in the spirit of Ye and Weiss (2003), we can combine the two methods into one by using the kernel matrix $\mathbf{M}_{tm} + \mathbf{M}_{dr}$. As shown in the next corollary, compared with each individual method, their ensemble is exhaustive under more general conditions.

Corollary 3.1. *If, for any sequence of non-stochastic vectors $\{v_T \in \mathcal{S}_{y|\mathbf{f}} : \|v_T\| = 1\}$,*

$$\max \{ \text{var}\{E(v'_T \mathbf{f}_t | y_{t+1})\}, \text{var}[E\{(v'_T \mathbf{f}_t)^2 | y_{t+1}\}], \text{var}[E\{(v'_T \mathbf{f}_t)^3 | y_{t+1}\}] \} = O_P^+(1),$$

then $\lambda_L(\mathbf{M}_{dr} + \mathbf{M}_{tm})$, the L th eigenvalue of $\mathbf{M}_{dr} + \mathbf{M}_{tm}$, is $O_P^+(1)$.

To estimate \mathbf{M}_{tm} using the contaminated factors $\widehat{\mathbf{f}}_t$, similar to directional regression, we need to slightly modify the kernel matrix, which leads to the following invariance result.

Theorem 3.2. *Assume that the factor loadings \mathbf{B} are known in priori, then under model (1.2), the kernel matrix \mathbf{M}_{tm} is invariant of the replacement of \mathbf{f}_t by $\widehat{\mathbf{f}}_t$, if we modify $\mu_0^3(\mathbf{f}_t | y_{t+1})$ to be*

$$\mu^3(\mathbf{f}_t | y_{t+1}) = \mu_0^3(\mathbf{f}_t | y_{t+1}) - E(\mathbf{f}_t \otimes \mathbf{f}_t \mathbf{f}'_t).$$

It is easy to see that the symmetry condition (B3) implies $E(\mathbf{f}_t \otimes \mathbf{f}_t \mathbf{f}'_t) = 0$. Thus the modified inverse moment $\mu^3(\mathbf{f}_t | y_{t+1})$ coincides with its original form $\mu_0^3(\mathbf{f}_t | y_{t+1})$. However, the

two moments differ when the estimated factors $\widehat{\mathbf{f}}_t$ are used in place of \mathbf{f}_t . The modification made in the former removes the effect of the error term \mathbf{u}_t^* from the kernel matrix, so it is crucial to the invariance result. When $E(\mathbf{u}_t^* \otimes \mathbf{u}_t^* \mathbf{u}_t^{*'}) = 0$, which occurs, for example, if \mathbf{u}_t^* has a symmetric distribution, the effect of \mathbf{u}_t^* on $\mu_0^3(\mathbf{f}_t|y_{t+1})$ automatically vanishes, and the invariance result holds without any modification on the inverse moment.

Same as for directional regression, when \mathbf{u}_t^* is negligible with sufficiently fast convergence order, the kernel matrix \mathbf{M}_{tm} for the true factors \mathbf{f}_t can be approximated by using the estimated factors $\widehat{\mathbf{f}}_t$, in which case consistent estimation of the central subspace can be justified without the invariance result. However, the invariance result makes its own contribution to the surrogate sufficient dimension reduction literature, as it sheds light on consistent sufficient dimension reduction estimation using inverse third moment, even when the measurement error is non-negligible.

Same as in Li (1991) and Li and Wang (2007), Yin and Cook (2003) used the slicing strategy to estimate the kernel matrix \mathbf{M}_{tm} . In our context, we modify the estimations Steps 2 and 3 to:

Step 2*. For $i = 0, \dots, H - 1$, estimate $\mu_0^3(\widehat{\mathbf{f}}_t|y_{t+1} \in I_{h,T})$ by

$$\widehat{\mu}_0^3(\widehat{\mathbf{f}}_t|y_{t+1}^D = i) = \sum_{l=1}^{c_i} (\widehat{\mathbf{f}}_{i,l} - \sum_{l=1}^{c_i} \widehat{\mathbf{f}}_{i,l}/c_i) \otimes (\widehat{\mathbf{f}}_{i,l} - \sum_{l=1}^{c_i} \widehat{\mathbf{f}}_{i,l}/c_i) (\widehat{\mathbf{f}}_{i,l} - \sum_{l=1}^{c_i} \widehat{\mathbf{f}}_{i,l}/c_i)' / c_i, \quad (3.2)$$

and $\mu^3(\widehat{\mathbf{f}}_t|y_{t+1}^D = i)$ by $\widehat{\mu}^3(\widehat{\mathbf{f}}_t|y_{t+1}^D = i) = \widehat{\mu}_0^3(\widehat{\mathbf{f}}_t|y_{t+1}^D = i) - T^{-1} \sum_{t=1}^T \widehat{\mathbf{f}}_t \otimes \widehat{\mathbf{f}}_t \mathbf{f}_t'$.

Step 3*. Estimate \mathbf{M}_{tm} by $\widehat{\mathbf{M}}_{tm} = H^{-1} \sum_{i=0}^{H-1} \widehat{\mu}_i' \widehat{\mu}_i$, in which $\widehat{\mu}_i$ contains the $K(K+1)/2$ distinct rows of $\widehat{\mu}^3(\widehat{\mathbf{f}}_t|y_{t+1}^D = i)$. Estimate $\mathcal{S}_{y|\mathbf{f}}$ by the L eigenvectors of $\widehat{\mathbf{M}}_{dr} + \widehat{\mathbf{M}}_{tm}$ corresponding to the largest eigenvalues.

When the estimated factor loadings $\widehat{\mathbf{B}}$ are sufficiently close to the true factor loadings \mathbf{B} , the resulting estimator is consistent, as justified in the following theorem.

Theorem 3.3. *Under Assumptions 2.1, 2.2 and 2.3, the assumption in Corollary 3.1, the linearity condition (B1), the constant variance condition (B2), and the symmetry condition (B3), if $K = o(\min(p^{1/3}, T^{1/3}))$, then the leading L eigenvectors of $\widehat{\mathbf{M}}_{dr} + \widehat{\mathbf{M}}_{tm}$, denoted by $\widehat{\phi}_1, \dots, \widehat{\phi}_L$, span a consistent estimator of the central subspace $\mathcal{S}_{y|\mathbf{f}}$ in the sense that*

$$\|(\widehat{\phi}_1, \dots, \widehat{\phi}_L)(\widehat{\phi}_1, \dots, \widehat{\phi}_L)' - (\phi_1, \dots, \phi_L)(\phi_1, \dots, \phi_L)'\|_F = O_P(K^{3/2}(p^{-1/2} + T^{-1/2})).$$

Similar to the case of directional regression, the error in estimating the central subspace can be divided into two parts, depending on how it is generated. The first part comes

from the estimation of the factor loadings, which is of order $K^{3/2}p^{-1/2}$ and decreases as a consequence of the increasing preciseness of the factor analysis when p grows. The second part comes from the use of sample moments, which is of order $K^{3/2}T^{-1/2}$ and decreases as the sample size T grows. Compared with directional regression, the second part of the estimation error here has a larger scale. This is because a larger number of inverse moments need to be estimated, which makes the consistency of the estimation more demanding on the sample size. Again, the linearity condition (B1) can be relaxed to more general regularity conditions, with the price of introducing an additional error term that is negligible but with unknown convergence order to the estimation.

To determine L , we can still apply the BIC type procedure proposed in Subsection 2.5. Referring to the discussion below Theorem 2.5, we can choose the tuning parameter C_T in the procedure to be $K^{1/2}p^{-1/2}T + K^{1/2}T^{1/2}$.

4 Numerical Studies

4.1 Simulation Result

We now present a numerical example to illustrate the performance of the proposed forecasting method that uses directional regression in the sufficient dimension reduction stage. The data generating process is specified as the following:

$$\begin{aligned} y_{t+1} &= h(\boldsymbol{\phi}'_1 \mathbf{f}_t, \boldsymbol{\phi}'_2 \mathbf{f}_t) + \sigma \epsilon_{t+1}, \\ x_{it} &= \mathbf{b}'_i \mathbf{f}_t + u_{it}. \end{aligned}$$

where the number of factors K is taken to be 6 and we fix $\boldsymbol{\phi}_1 = (1, 1, 1, 0, 0, 0)/\sqrt{3}$, $\boldsymbol{\phi}_2 = (1, 0, 0, 0, 1, 3)/\sqrt{11}$. The factor loadings \mathbf{b}_i are independently sampled from $U[-1, 2]$. We generate the latent factors $f_{j,t}$ and the error terms u_{it} from two AR(1) processes, $f_{j,t} = \alpha_j f_{j,t-1} + e_{jt}$ and $u_{it} = \rho_i u_{i,t-1} + \nu_{it}$, with α_j, ρ_i drawn from $U[0.2, 0.8]$ and fixed during the simulation, and the noises e_{jt}, ν_{it} , are $N(0, 1)$. We set $\epsilon_{t+1} \sim N(0, 1)$ and $\sigma = 0.2$.

Table 1: Performance of estimated $\hat{\phi}$ using $R^2(\hat{\phi})$ (%)

Model I		SIR		DR	
p	T	$R^2(\hat{\phi}_1)$	$R^2(\hat{\phi}_2)$	$R^2(\hat{\phi}_1)$	$R^2(\hat{\phi}_2)$
100	100	75.0(21.3)	28.4(27.4)	82.9(14.8)	79.9(21.9)
100	200	88.7(10.4)	17.7(27.6)	94.5(5.4)	91.5(8.5)
100	500	95.9(3.6)	14.4(28.2)	98.4(1.4)	96.0(3.4)
200	100	75.9(21.4)	28.8(27.4)	83.3(16.0)	80.3(21.2)
500	200	95.9(3.4)	14.1(28.3)	98.4(1.5)	96.2(3.1)
500	500	96.1(3.4)	14.6(28.6)	98.3(1.5)	96.3(3.1)
Model II		SIR		DR	
100	100	95.8(3.5)	21.0(25.7)	95.8(3.5)	26.4(26.6)
100	200	97.8(1.8)	32.4(27.7)	97.9(1.8)	43.4(28.7)
100	500	99.1(0.7)	63.8(27.0)	99.1(0.7)	74.8(23.8)
200	100	95.8(3.2)	22.9(25.4)	95.8(3.2)	27.0(26.6)
500	200	99.1(0.7)	62.0(27.1)	99.1(0.7)	74.8(23.2)
500	500	99.1(0.7)	64.1(26.7)	99.1(0.7)	75.6(22.5)
Model III		SIR		DR	
100	100	33.4(26.7)	26.1(23.4)	83.0(19.7)	47.6(28.2)
100	200	34.8(27.3)	23.8(22.7)	94.9(4.1)	83.2(22.9)
100	500	33.0(28.1)	24.2(23.4)	98.4(1.4)	97.6(2.1)
200	100	34.3(28.5)	23.7(23.3)	83.9(19.1)	47.3(28.4)
500	200	32.5(28.3)	24.2(23.3)	98.3(1.4)	97.5(2.0)
500	500	31.7(28.4)	24.5(23.2)	98.4(1.3)	97.7(2.0)
Model IV		SIR		DR	
100	100	61.8(29.1)	31.3(26.0)	85.6(14.2)	79.1(23.5)
100	200	75.1(26.4)	41.6(27.9)	94.5(4.9)	93.5(5.2)
100	500	89.4(15.0)	67.8(27.4)	98.1(1.7)	97.7(1.9)
200	100	58.9(29.1)	33.4(25.8)	85.7(14.1)	80.2(22.6)
500	200	90.1(14.4)	72.2(27.2)	98.1(1.5)	97.8(1.8)
500	500	90.2(13.9)	70.9(26.9)	98.1(1.5)	98.0(1.6)

Notes: Median squared multiple correlation coefficients in percentage. The values in parentheses are standard deviations. The simulation is based on 1000 replications.

We consider four different choices of the link function $g(\cdot)$,

- Model I: $y_{t+1} = 0.4(\phi'_1 \mathbf{f}_t)^2 + 3 \sin(\phi'_2 \mathbf{f}_t/4) + \sigma \epsilon_{t+1}$;
- Model II: $y_{t+1} = 3 \sin(\phi'_1 \mathbf{f}_t/4) + 3 \sin(\phi'_2 \mathbf{f}_t/4) + \sigma \epsilon_{t+1}$;

- Model III: $y_{t+1} = 0.4(\phi'_1 \mathbf{f}_t)^2 + |\phi'_2 \mathbf{f}_t|^{1/2} + \sigma \epsilon_{t+1}$;
- Model IV: $y_{t+1} = (\phi'_1 \mathbf{f}_t)(\phi'_2 \mathbf{f}_t + 1) + \sigma \epsilon_{t+1}$.

The proposed inverse second-moment method using directional regression (DR) is then compared with the inverse first-moment method using sliced inverse regression (SIR) introduced in Fan et al. (2016) and the linear PC-estimator (principal components). In model I and III, at least one component is symmetric, which cannot be estimated well by SIR. In model II, the components are roughly monotone, making it favorable to SIR. Model IV contains the interaction component, which allows us to examine the ability of each method in detecting such nonlinear effect.

To gauge the quality of the estimated directions, we adopt the squared multiple correlation coefficient $R^2(\hat{\phi}) = \max_{\phi \in S_{y|\mathbf{f}}, \|\phi\|=1} (\phi' \hat{\phi})^2$, where $S_{y|\mathbf{f}}$ is the central space spanned by ϕ_1 and ϕ_2 . Note that one could ensure the true factors and loadings meet the identifiability conditions by calculating an invertible matrix \mathbf{H} such that $T^{-1} \mathbf{H} \mathbf{F}' \mathbf{F} \mathbf{H}' = \mathbf{I}_K$ and $\mathbf{H}^{-1} \mathbf{B}' \mathbf{B} \mathbf{H}^{-1}$ is diagonal. The central subspace is then understood as $\mathbf{H}^{-1} S_{y|\mathbf{f}}$. But here we still simply denote the rotated central space as $S_{y|\mathbf{f}}$ (see Fan et al. 2015).

Table 1 compares the estimation quality of SIR and DR under the aforementioned four models. The PC-estimator does not apply here since it produces only one directional estimate. It is evident that DR has substantial improvement over SIR in model I, III and IV, with higher R^2 and lower variance. This is not surprising as DR explores higher conditional moments and hence incorporates more information. Even in model II, where SIR works well, DR yields comparable results. We also observe that DR has outstanding performance in small samples, which makes it favorable in practical applications.

We next investigate the predicting power of DR through the lens of out-of-sample R^2 , i.e.,

$$R^2 = 1 - \frac{\sum_{t=T+1}^{T+n_T} (y_t - \hat{y}_t)^2}{\sum_{t=T+1}^{T+n_T} (y_t - \bar{y}_t)^2},$$

where for different training samples (p, T) we use a fixed length $n_T = 100$ of testing samples to evaluate the out-of-sample performance. \hat{y}_t is the predicted value using all information prior to t . The fitting is done by building an additive model for the extracted indices. In the case of PC-estimator, six smooth functions are constructed for the six estimated factors. In contrast, only two smooth functions are applied in the cases of SIR and DR. It is clear from Table 2 that DR enjoys great performance in almost all the cases. The PC-estimator is more robust than SIR in the presence of symmetric components, but fails to capture the

interaction effect in general.

Table 2: Performance of forecast using out-of-sample R^2 (%)

		Model I			Model II		
p	T	SIR	DR	PC	SIR	DR	PC
100	100	-2.9	51.3	5.0	95.2	95.2	94.5
100	200	3.4	70.3	12.6	95.5	95.5	95.9
100	500	8.4	88.5	21.3	95.8	95.8	96.3
200	100	-3.1	51.0	7.1	95.7	95.7	95.6
500	200	9.0	89.6	19.2	96.2	96.2	96.2
500	500	7.5	90.2	20.7	96.8	96.7	96.3
		Model III			Model IV		
p	T	SIR	DR	PC	SIR	DR	PC
100	100	-1.1	58.0	12.5	10.3	37.7	-15.8
100	200	6.3	81.9	20.0	21.2	48.5	-6.7
100	500	9.4	90.9	27.1	33.3	51.5	3.9
200	100	-0.5	60.7	11.5	9.4	39.3	-15.0
500	200	8.4	92.3	25.7	34.2	51.3	3.0
500	500	8.4	93.0	26.6	33.3	52.3	2.7

Notes: Out-of-sample median R^2 in percentage over 1000 replications.

4.2 Macro Index Forecast

This section analyzes how diffusion indices constructed by the proposed DR impact real-data forecasts. We use a monthly macro dataset consisting of 134 macroeconomic time series recently composed by McCracken and Ng (2016), which are classified into 8 groups : (1) output and income, (2) labor market, (3) housing, (4) consumption, orders and inventories, (5) money and credit, (6) bond and exchange rates, (7) prices, and (8) stock market. The dataset spans the period from 1959:01 to 2016:01. For a given target time series, we model the multi-step-ahead variable as:

$$y_{t+h}^h = g(\phi_1' \mathbf{f}_t, \dots, \phi_L' \mathbf{f}_t, \epsilon_{t+h}^h),$$

where $y_{t+h}^h = h^{-1} \sum_{i=1}^h y_{t+i}$ is the variable to forecast, as in Stock and Watson (2002), and L is taken to be 1 or 2.

Table 3: RMSE in Out of Sample Forecast (Median/Max/Min)

Group ($h = 1$)	SIR(1)	SIR(2)	DR(1)	DR(2)	NL-PC
Output and Income	1.03/1.61/0.96	1.02/1.13/0.94	0.99/1.19/0.92	1.02/1.14/0.90	1.21/1.38/1.05
Consumption	1.00/2.10/0.80	0.95/1.05/0.74	0.92/1.02/0.86	1.00/1.05/0.81	1.16/1.44/1.04
Labor market	1.02/2.27/0.71	1.00/1.21/0.42	0.97/1.13/0.52	0.98/1.16/0.42	1.21/1.53/0.46
Housing	1.04/1.32/0.64	0.92/1.08/0.52	0.83/1.04/0.50	0.79/0.94/0.44	0.83/0.97/0.49
Money and credit	0.94/1.04/0.86	0.97/1.05/0.90	0.96/1.10/0.86	1.04/1.24/0.92	1.14/1.41/1.07
Stock market	0.99/1.39/0.90	1.02/1.12/0.83	0.92/1.08/0.88	1.04/1.07/0.91	1.36/1.39/1.14
Interest rates	1.04/1.79/0.79	0.93/1.17/0.61	0.90/1.04/0.59	0.92/1.15/0.62	1.12/1.32/0.73
Prices	0.97/1.42/0.80	0.99/1.05/0.83	0.95/1.12/0.81	0.97/1.12/0.88	1.12/1.47/0.92
Group ($h = 6$)	SIR(1)	SIR(2)	DR(1)	DR(2)	NL-PC
Output and Income	1.07/1.47/0.93	0.97/1.23/0.81	0.99/1.18/0.89	1.05/1.27/0.95	1.28/1.52/0.97
Consumption	1.16/1.73/0.90	0.90/1.12/0.67	0.94/1.16/0.71	1.03/1.14/0.73	1.28/1.66/0.77
Labor market	1.15/2.02/0.68	0.89/1.22/0.39	0.90/1.26/0.48	0.98/1.39/0.43	1.24/1.42/0.45
Housing	0.96/1.29/0.66	0.85/0.95/0.51	0.73/0.89/0.50	0.69/0.86/0.47	0.78/1.02/0.55
Money and credit	0.95/3.51/0.76	1.01/3.65/0.83	0.99/1.52/0.76	1.02/1.74/0.78	1.23/2.90/0.92
Stock market	0.91/1.20/0.83	0.94/1.05/0.89	0.89/1.08/0.84	1.00/1.03/0.94	1.23/1.27/0.83
Interest rates	1.01/1.61/0.75	0.90/1.12/0.64	0.84/1.13/0.50	0.88/1.18/0.58	1.11/1.46/0.70
Prices	1.16/1.37/0.51	1.03/1.12/0.82	1.11/1.37/0.94	1.14/1.36/0.95	1.17/1.35/1.11
Group ($h = 12$)	SIR(1)	SIR(2)	DR(1)	DR(2)	NL-PC
Output and Income	1.24/1.67/0.79	1.01/1.45/0.76	0.99/1.22/0.76	1.01/1.36/0.86	1.17/1.34/0.92
Consumption	1.27/1.60/0.83	1.08/1.44/0.62	1.09/1.32/0.65	1.06/1.38/0.66	1.16/1.38/0.87
Labor market	1.07/1.76/0.67	0.83/1.40/0.41	0.91/1.44/0.54	0.89/1.41/0.46	1.13/1.39/0.56
Housing	0.85/1.35/0.59	0.69/0.93/0.46	0.67/0.91/0.40	0.68/0.83/0.36	0.89/1.16/0.54
Money and credit	1.14/2.03/0.41	1.03/2.16/0.80	1.05/1.52/0.85	1.00/1.40/0.82	1.20/1.69/0.87
Stock market	1.09/1.20/0.89	1.01/1.13/0.84	0.96/1.17/0.94	1.08/1.16/0.75	1.06/1.14/0.89
Interest rates	1.00/1.31/0.75	0.82/1.22/0.59	0.80/1.27/0.53	0.85/1.18/0.51	1.07/1.62/0.70
Prices	1.18/1.40/0.53	1.21/1.40/0.66	1.19/1.31/0.71	1.21/1.33/0.77	1.25/1.52/0.94

Notes: Out-of-sample mean squared error (MSE) relative to the linear diffusion index. In each group, the median, maximum and minimum of RMSE is reported. SIR(i) denotes sufficient forecasting using i indices, DR denotes sufficient directional forecasting, and NL-PC denotes a nonlinear additive model on all the estimated factors.

Forecasts of y_{t+h}^h are constructed based on a moving window with fixed length ($T = 120$) to account for timeliness. For each fixed window, the factors in the forecasting equation are estimated by the method of principal components using all time series except the target. As noted by McCracken and Ng (2016) 8 factors have good explanatory power in various

cases, so we set $K = 8$ throughout the exercise. For each method M , we calculate the mean squared out-of-sample forecasting error

$$\text{MSE}(M) = \frac{1}{m} \sum_{t=T+1}^{T+m} (y_t - \hat{y}_t)^2,$$

and report the relative MSE (RMSE) to the PC method,

$$\text{RMSE}(M) = \frac{\text{MSE}(M)}{\text{MSE}(\text{PC})},$$

which we evaluate on the last $m = 240$ months (20 years). The methods we consider here include $\text{SIR}(i)$, $\text{DR}(i)$ ($i = 1, 2$), where $\text{SIR}(i)$ denotes sufficient forecasting with $L = i$, and similar for DR. Both methods use an additive model in specifying the forecasting equation. We also impose an additive model to the estimated factors, denoted by NL-PC, to see how much we can leverage on the nonlinearity without projecting the principal components.

We report results in Table 3 for $h = 1, 6, 12$, on the maximum, minimum and median of RMSE in each broad sector. Several features are noteworthy. First, a nonlinear additive model built on estimated factors does not buy us more predictive power, except in the housing sector, where most of the nonlinear methods improve prediction accuracy. Second, the one-step-ahead out-of-sample forecast favors $\text{DR}(1)$, as we observe the median RMSEs are uniformly less than 1 and some of the reductions in RMSE are substantial. Moving from short horizon to long horizon changes predictability of the targets, but $\text{DR}(1)$ manages to improve the forecast over the PC method in many instances. Finally, as an illustration, we plot the out-of-sample R^2 for the 6-month-ahead forecast using $\text{DR}(1)$ and PC. Notably, macro time series in housing and labor market sectors have higher predictability than in rates and stock market sectors.

5 Proofs

5.1 Consistency of factor analysis

In this subsection, we give some basic theoretical results about the consistency of estimating the factors and factor loadings, summarized in the following four lemmas. They will be useful for the proof of theorems later. Again, we assume K to be known and diverging with p .




Figure 1: 6-month ahead forecasting out-of-sample R^2 for the 134 macroeconomic series organized into eight groups, using DR(1) and PC methods.

Let \mathbf{V} denote the $K \times K$ diagonal matrix consisting of the K largest eigenvalues of the sample covariance matrix $(pT)^{-1}\mathbf{X}'\mathbf{X}$ in descending order. Define a $K \times K$ matrix

$$\mathbf{H} = \mathbf{V}^{-1} \frac{\widehat{\mathbf{F}}'\mathbf{F}}{T} \frac{\mathbf{B}'\mathbf{B}}{p}, \quad (5.1)$$

where $\mathbf{F}' = (\mathbf{f}_1, \dots, \mathbf{f}_T)$ and $\widehat{\mathbf{F}}$ is estimated from principal component analysis. We first summarize some basic facts under Assumptions (2.1), (2.2) and (2.3) in the following lemma. The proof is omitted.

Lemma 5.1. 1) $\|\mathbf{B}\| = O_p(p^{1/2})$ and $\|\mathbf{B}\| = O_p^+(p^{1/2})$, 2) $\|\mathbf{f}_t\| = O_p(K^{1/2})$, 3) $\|\mathbf{V}\| = O_p^+(1)$, and 4) $\|\mathbf{H}\| = O_p(1)$.

We have the following results for factor estimates:

Lemma 5.2. For any $K = o(\min\{p^{1/3}, T\})$,

$$\frac{1}{T} \sum_{t=1}^T \|\widehat{\mathbf{f}}_t - \mathbf{H}\mathbf{f}_t\|^2 = O_p\left(\frac{K^3}{p} + \frac{K}{T}\right).$$

Proof of Lemma 5.2. This is a direct result from Li et al. (2013). Note that they used

$\|\mathbf{V}\widehat{\mathbf{f}}_t - \mathbf{V}\mathbf{H}\mathbf{f}_t\|$ as the summand. As $\|\mathbf{V}\| = O_p^+(1)$, it completes the proof. \square

Lemma 5.3. *Under the above assumptions, for any $K = o(\min\{p^{1/3}, T\})$, the $K \times K$ matrices*

- 1) $T^{-1}(\widehat{\mathbf{F}} - \mathbf{F}\mathbf{H}')'\mathbf{F} = O_p(\frac{K}{T} + \frac{K^2}{p})$.
- 2) $T^{-1}(\widehat{\mathbf{F}} - \mathbf{F}\mathbf{H}')'\widehat{\mathbf{F}} = O_p(\frac{K}{T} + \frac{K^3}{p})$.

Proof of Lemma 5.3. 1) The proof of the lemma closely follows Lemma B.2 in Bai (2003). By Theorem 1 of Bai (2003), we have the following identity,

$$\widehat{\mathbf{f}}_t - \mathbf{H}\mathbf{f}_t = \mathbf{V}^{-1}\left(\frac{1}{T}\sum_{s=1}^T\widehat{\mathbf{f}}_s\gamma_{st} + \frac{1}{T}\sum_{s=1}^T\widehat{\mathbf{f}}_s\zeta_{st} + \frac{1}{T}\sum_{s=1}^T\widehat{\mathbf{f}}_s\eta_{st} + \frac{1}{T}\sum_{s=1}^T\widehat{\mathbf{f}}_s\xi_{st}\right),$$

where $\gamma_{st} = p^{-1}E(\mathbf{u}'_s\mathbf{u}_t)$, $\zeta_{st} = p^{-1}\mathbf{u}'_s\mathbf{u}_t - \gamma_{st}$, $\eta_{st} = \mathbf{f}'_s\mathbf{B}'\mathbf{u}_t/p$ and $\xi_{st} = \mathbf{f}'_t\mathbf{B}'\mathbf{u}_s/p$. The identity leads to

$$\begin{aligned} T^{-1}\sum_{t=1}^T(\widehat{\mathbf{f}}_t - \mathbf{H}\mathbf{f}_t)\mathbf{f}'_t &= \mathbf{V}^{-1}\left[T^{-2}\sum_{t=1}^T\sum_{s=1}^T\widehat{\mathbf{f}}_s\mathbf{f}'_t\gamma_{st} + T^{-2}\sum_{t=1}^T\sum_{s=1}^T\widehat{\mathbf{f}}_s\mathbf{f}'_t\zeta_{st} \right. \\ &\quad \left. + T^{-2}\sum_{t=1}^T\sum_{s=1}^T\widehat{\mathbf{f}}_s\mathbf{f}'_t\eta_{st} + T^{-2}\sum_{t=1}^T\sum_{s=1}^T\widehat{\mathbf{f}}_s\mathbf{f}'_t\xi_{st}\right] \\ &= \mathbf{V}^{-1}(I + II + III + IV). \end{aligned}$$

We begin with term I , which can be written as

$$I = T^{-2}\sum_{t=1}^T\sum_{s=1}^T(\widehat{\mathbf{f}}_s - \mathbf{H}\mathbf{f}_s)\mathbf{f}'_t\gamma_{st} + T^{-2}\mathbf{H}\sum_{t=1}^T\sum_{s=1}^T\mathbf{f}_s\mathbf{f}'_t\gamma_{st}.$$

The first term is bounded by

$$T^{-1}\left(\frac{1}{T}\sum_{s=1}^T\|\widehat{\mathbf{f}}_s - \mathbf{H}\mathbf{f}_s\|^2\right)^{1/2}\left(\frac{1}{T}\sum_{t=1}^T\sum_{s=1}^T\|\mathbf{f}_t\|^2\gamma_{st}^2\right)^{1/2} = T^{-1}O_p\left(\frac{K^{3/2}}{\sqrt{p}} + \frac{K^{1/2}}{\sqrt{T}}\right)O_p(\sqrt{K}),$$

where $O_p(\sqrt{K})$ follows from $K^{-1}E(\|\mathbf{f}_t\|^2) \leq M$ and $\frac{1}{T}\sum_{t=1}^T\sum_{s=1}^T\gamma_{st}^2 \leq M$ by Lemma 1(i) of Bai and Ng (2002). Ignoring \mathbf{H} , the expectation of the second term of I is bounded by

$$T^{-1}E\|\mathbf{f}_t\|^2\left(T^{-1}\sum_{t=1}^T\sum_{s=1}^T|\gamma_{st}|\right) \leq O(T^{-1}K).$$

The second term is thus $O_p(T^{-1}K)$ and I is $O_p(\frac{K^2}{T\sqrt{p}} + \frac{K}{T^{3/2}} + \frac{K}{T}) = O_p(K/T)$.

For II , we write it as

$$II = T^{-2} \sum_{t=1}^T \sum_{s=1}^T (\hat{\mathbf{f}}_s - \mathbf{H}\mathbf{f}_s) \mathbf{f}'_t \zeta_{st} + T^{-2} \mathbf{H} \sum_{t=1}^T \sum_{s=1}^T \mathbf{f}_s \mathbf{f}'_t \zeta_{st}.$$

For the first term, we have

$$\|T^{-2} \sum_{t=1}^T \sum_{s=1}^T (\hat{\mathbf{f}}_s - \mathbf{H}\mathbf{f}_s) \mathbf{f}'_t \zeta_{st}\| \leq \left(\frac{1}{T} \sum_{s=1}^T \|\hat{\mathbf{f}}_s - \mathbf{H}\mathbf{f}_s\|^2\right)^{1/2} \left(\frac{1}{T} \sum_{s=1}^T \left\|\frac{1}{T} \sum_{t=1}^T \zeta_{st} \mathbf{f}_t\right\|^2\right)^{1/2}$$

Since

$$\frac{1}{T} \sum_{t=1}^T \zeta_{st} \mathbf{f}_t = \frac{1}{\sqrt{p}} \frac{1}{T} \sum_{t=1}^T \left(\frac{1}{\sqrt{p}} \sum_{k=1}^p [u_{ks} u_{kt} - E(u_{ks} u_{kt})]\right) \mathbf{f}_t = O_p\left(\left(\frac{K}{p}\right)^{1/2}\right),$$

the first term is $O_p(\frac{K^{3/2}}{\sqrt{p}} + \frac{K^{1/2}}{\sqrt{T}}) \cdot O_p(\sqrt{\frac{K}{p}}) = O_p(\frac{K^2}{p} + \frac{K}{\sqrt{pT}})$. The second term can be written as $\frac{\mathbf{H}}{\sqrt{pT}} (\frac{1}{T} \sum_{t=1}^T z_t \mathbf{f}_t)$ with $z_t = \frac{1}{\sqrt{pT}} \sum_{s=1}^T \sum_{k=1}^p [u_{ks} u_{kt} - E(u_{ks} u_{kt})] \mathbf{f}_s$. Since $K^{-1} E\|z_t\|^2 \leq M$, $E\|z_t \mathbf{f}_t\| \leq (E\|z_t\|^2 E\|\mathbf{f}_t\|^2)^{1/2} \leq KM$. This implies that $\frac{1}{T} \sum_{t=1}^T z_t \mathbf{f}_t = O_p(K)$, which makes the second term $O_p(\frac{K}{\sqrt{pT}})$. Thus $II = O_p(\frac{K^2}{p} + \frac{K}{\sqrt{pT}})$.

For III , we similarly write it as

$$III = T^{-2} \sum_{t=1}^T \sum_{s=1}^T (\hat{\mathbf{f}}_s - \mathbf{H}\mathbf{f}_s) \mathbf{f}'_t \eta_{st} + T^{-2} \mathbf{H} \sum_{t=1}^T \sum_{s=1}^T \mathbf{f}_s \mathbf{f}'_t \eta_{st}.$$

Now, $T^{-2} \mathbf{H} \sum_{t=1}^T \sum_{s=1}^T \mathbf{f}_s \mathbf{f}'_t \eta_{st} = \mathbf{H} (\frac{1}{T} \sum_{s=1}^T \mathbf{f}_s \mathbf{f}'_s) \frac{1}{pT} \sum_{t=1}^T \sum_{k=1}^p \mathbf{b}_k \mathbf{f}'_t u_{kt} = O_p(\frac{K}{\sqrt{pT}})$, where we used the fact that $\frac{1}{T} \sum_{s=1}^T \mathbf{f}_s \mathbf{f}'_s = \mathbf{I}_K$ and $E\|\frac{1}{K\sqrt{pT}} \sum_{t=1}^T \sum_{k=1}^p \mathbf{b}_k \mathbf{f}'_t u_{kt}\|^2 \leq M$. Consider the first term,

$$T^{-2} \sum_{t=1}^T \sum_{s=1}^T (\hat{\mathbf{f}}_s - \mathbf{H}\mathbf{f}_s) \mathbf{f}'_t \eta_{st} \leq \left(\frac{1}{T} \sum_{s=1}^T \|\hat{\mathbf{f}}_s - \mathbf{H}\mathbf{f}_s\|^2\right)^{1/2} \left(\frac{1}{T} \sum_{s=1}^T \left\|\frac{1}{T} \sum_{t=1}^T \mathbf{f}'_t \eta_{st}\right\|^2\right)^{1/2},$$

where the second part can be rewritten as

$$\left(\frac{1}{T} \sum_{s=1}^T \left\|\frac{1}{pT} \sum_{t=1}^T \mathbf{f}'_t \sum_{k=1}^p \mathbf{f}'_s \mathbf{b}_k u_{kt}\right\|^2\right)^{1/2} = \frac{1}{\sqrt{pT}} \left(\frac{1}{T} \sum_{s=1}^T \|\mathbf{f}'_s\| \frac{1}{\sqrt{pT}} \sum_{t=1}^T \sum_{k=1}^p \mathbf{b}_k \mathbf{f}'_t u_{kt}\right)^{1/2}$$

which is $O_p(\frac{K^{3/2}}{\sqrt{pT}})$. So $T^{-2} \sum_{t=1}^T \sum_{s=1}^T (\hat{\mathbf{f}}_s - \mathbf{H}\mathbf{f}_s) \mathbf{f}'_t \eta_{st} = O_p(\frac{K^{3/2}}{\sqrt{p}} + \frac{K^{1/2}}{\sqrt{T}}) \cdot O_p(\frac{K^{3/2}}{\sqrt{pT}}) = O_p(\frac{K^3}{p\sqrt{T}} +$

$\frac{K^2}{T\sqrt{p}}$). Thus $III = O_p(\frac{K^3}{p\sqrt{T}} + \frac{K^2}{T\sqrt{p}} + \frac{K}{\sqrt{pT}})$.

The proof for IV is similar to that of III . Hence,

$$I + II + III + IV = O_p\left(\frac{K}{T} + \frac{K^2}{p} + \frac{K}{\sqrt{pT}} + \frac{K^3}{p\sqrt{T}} + \frac{K^2}{T\sqrt{p}}\right).$$

Since $K = o(\min\{p^{1/3}, T\})$, we have $I + II + III + IV = O_p(\frac{K}{T} + \frac{K^2}{p})$. Thus,

$$T^{-1}(\widehat{\mathbf{F}} - \mathbf{F}\mathbf{H}')'\mathbf{F} = O_p\left(\frac{K}{T} + \frac{K^2}{p}\right)$$

2)

$$\begin{aligned} T^{-1}(\widehat{\mathbf{F}} - \mathbf{F}\mathbf{H}')'\widehat{\mathbf{F}} &= T^{-1}(\widehat{\mathbf{F}} - \mathbf{F}\mathbf{H}')'\mathbf{F}\mathbf{H}' + T^{-1}(\widehat{\mathbf{F}} - \mathbf{F}\mathbf{H}')'(\widehat{\mathbf{F}} - \mathbf{F}\mathbf{H}') \\ &= O_p\left(\frac{K}{T} + \frac{K^2}{p} + \frac{K}{T} + \frac{K^3}{p}\right) \\ &= O_p\left(\frac{K}{T} + \frac{K^3}{p}\right) \end{aligned}$$

This completes the proof. □

Lemma 5.4. *Suppose $K = o(\min\{p^{1/3}, T\})$. For the factor identification matrix, we have*

$$\|\mathbf{H} - \mathbf{I}_K\| = O_p\left(\frac{K^3}{p} + \frac{K}{T}\right)$$

Proof of Lemma 5.4. Suppose that $T^{-1}(\widehat{\mathbf{F}} - \mathbf{F}\mathbf{H}')'\mathbf{F} = O_p(a_1)$ and $T^{-1}(\widehat{\mathbf{F}} - \mathbf{F}\mathbf{H}')'\widehat{\mathbf{F}} = O_p(a_2)$. Following the same argument in Bai and Ng (2013), it's easy to see that

$$\|\mathbf{H} - \mathbf{I}_K\| = O_p(a_1 + a_2). \tag{5.2}$$

This completes the proof. □

5.2 Proof of Theorem 2.1

1) According to the arguments in Lemma 6.4 of Fan et.al. (2015), $\|\widehat{\mathbf{B}} - \mathbf{B}\| \leq \|\widehat{\mathbf{B}} - \mathbf{B}\mathbf{H}'\| + \|\mathbf{B}(\mathbf{H}' - \mathbf{I}_K)\|$. The square of the first term is bounded by

$$\begin{aligned} \|\widehat{\mathbf{B}} - \mathbf{B}\mathbf{H}'\|^2 &\leq 2 \sum_{i=1}^p \left\| \frac{1}{T} \sum_{t=1}^T x_{it} (\widehat{\mathbf{f}}_t - \mathbf{H}\mathbf{f}_t) \right\|^2 + 2 \sum_{i=1}^p \left\| \frac{1}{T} \sum_{t=1}^T \mathbf{H}\mathbf{f}_t u_{it} \right\|^2 \\ &\leq O_p\left(p\left(\frac{K^3}{p} + \frac{K}{T}\right) + \frac{pK}{T}\right) = O_p\left(p\left(\frac{K^3}{p} + \frac{K}{T}\right)\right). \end{aligned}$$

The second term

$$\|\mathbf{B}(\mathbf{H}' - \mathbf{I}_K)\| \leq \|\mathbf{B}\| \|\mathbf{H} - \mathbf{I}_K\| = O_p((p)^{1/2}(\frac{K^3}{p} + \frac{K}{T}))$$

is smaller than the first term as long as $K = o(\min\{p^{1/3}, T\})$.

2) By Assumption 2.1 and simple algebra, we have $\|(\mathbf{B}'\mathbf{B})^{-1}\| = O_P(p^{-1})$, which, together with 1), implies that $\|(\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1}\| = O_P(p^{-1})$. Also, by 1) and simple algebra, we have $\|\widehat{\mathbf{B}}'\widehat{\mathbf{B}} - \mathbf{B}'\mathbf{B}\| = O_p(p(K^{3/2}p^{-1/2} + K^{1/2}T^{-1/2}))$. Thus,

$$\begin{aligned} \|(\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1} - (\mathbf{B}'\mathbf{B})^{-1}\| &= \|(\mathbf{B}'\mathbf{B})^{-1}(\mathbf{B}'\mathbf{B} - \widehat{\mathbf{B}}'\widehat{\mathbf{B}})(\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1}\| \\ &\leq \|(\mathbf{B}'\mathbf{B})^{-1}\| \|\widehat{\mathbf{B}}'\widehat{\mathbf{B}} - \mathbf{B}'\mathbf{B}\| \|(\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1}\| \\ &= O_P(p^{-1}(K^{3/2}p^{-1/2} + K^{1/2}T^{-1/2})), \end{aligned}$$

Together with 1), it implies

$$\begin{aligned} \|\widehat{\Lambda}_b - \Lambda_b\| &= \|(\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1}\widehat{\mathbf{B}}' - (\mathbf{B}'\mathbf{B})^{-1}\mathbf{B}'\| \\ &= \|(\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1}\widehat{\mathbf{B}}' - (\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1}\mathbf{B}' + (\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1}\mathbf{B}' - (\mathbf{B}'\mathbf{B})^{-1}\mathbf{B}'\| \\ &\leq \|(\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1}\widehat{\mathbf{B}}' - (\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1}\mathbf{B}'\| + \|(\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1}\mathbf{B}' - (\mathbf{B}'\mathbf{B})^{-1}\mathbf{B}'\| \\ &\leq \|(\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1}\| \|\widehat{\mathbf{B}} - \mathbf{B}\| + \|(\widehat{\mathbf{B}}'\widehat{\mathbf{B}})^{-1} - (\mathbf{B}'\mathbf{B})^{-1}\| \|\mathbf{B}'\| \\ &= O_P(p^{-1/2}(K^{3/2}p^{-1/2} + K^{1/2}T^{-1/2})). \end{aligned}$$

This completes the proof. □

5.3 Proof of Theorem 2.2

By (2.4), \mathbf{M}_{dr} is invariant of the change from \mathbf{f}_t to $\widehat{\mathbf{f}}_t$, if we can show the same invariance result for the centered inverse moments $E(\mathbf{f}_t|y_{t+1})$ and $E(\mathbf{f}_t\mathbf{f}'_t|y_{t+1}) - \text{var}(\mathbf{f}_t)$. By (1.1) and (1.2), $\{\mathbf{u}_t\}_{t \geq 1}$ is independent of $\{(\mathbf{f}_t, y_{t+1}) : t \geq 1\}$, and $E(\mathbf{u}_t^*) = 0$, which implies that

$$E(\widehat{\mathbf{f}}_t|y_{t+1}) = E(\mathbf{f}_t|y_{t+1}) + E(\mathbf{u}_t^*|y_{t+1}) = E(\mathbf{f}_t|y_{t+1}) + E(\mathbf{u}_t^*) = E(\mathbf{f}_t|y_{t+1}). \quad (5.3)$$

To show the invariance of $E(\mathbf{f}_t\mathbf{f}'_t|y_{t+1}) - \text{var}(\mathbf{f}_t)$, we first note that

$$\begin{aligned} E(\widehat{\mathbf{f}}_t\widehat{\mathbf{f}}'_t|y_{t+1}) &= E[(\mathbf{f}_t + \mathbf{u}_t^*)(\mathbf{f}_t + \mathbf{u}_t^*)'|y_{t+1}] \\ &= E(\mathbf{f}_t\mathbf{f}'_t|y_{t+1}) + E(\mathbf{f}_t\mathbf{u}_t^{*'}|y_{t+1}) + E(\mathbf{u}_t^*\mathbf{f}'_t|y_{t+1}) + E(\mathbf{u}_t^*\mathbf{u}_t^{*'}|y_{t+1}) \\ &= E(\mathbf{f}_t\mathbf{f}'_t|y_{t+1}) + E(\mathbf{f}_t|y_{t+1})E(\mathbf{u}_t^{*'}) + E(\mathbf{u}_t^*)E(\mathbf{f}'_t|y_{t+1}) + E(\mathbf{u}_t^*\mathbf{u}_t^{*'}) \\ &= E(\mathbf{f}_t\mathbf{f}'_t|y_{t+1}) + \Sigma_{\mathbf{u}^*}, \end{aligned}$$

where $\Sigma_{\mathbf{u}^*} = I_K + (\mathbf{B}'\mathbf{B})^{-1}\mathbf{B}'\Sigma_u\mathbf{B}(\mathbf{B}'\mathbf{B})^{-1}$. On the other hand, by simple algebra, $\text{var}(\widehat{\mathbf{f}}_t) = E[(\mathbf{f}_t + \mathbf{u}_t^*)(\mathbf{f}_t + \mathbf{u}_t^*)'] - \text{var}(\mathbf{f}_t) = \text{var}(\mathbf{f}_t) + \Sigma_{\mathbf{u}^*}$. Hence we have

$$E(\widehat{\mathbf{f}}_t\widehat{\mathbf{f}}'_t|y_{t+1}) - \text{var}(\widehat{\mathbf{f}}_t) = E(\mathbf{f}_t\mathbf{f}'_t|y_{t+1}) - \text{var}(\mathbf{f}_t).$$

Combined with (5.3), it can be readily seen that \mathbf{M}_{dr} is invariant if \mathbf{f}_t is replaced by $\widehat{\mathbf{f}}_t$. This completes the proof. \square

5.4 Proof of Lemma 2.1

For simplicity in notations, we denote $\widehat{\mathbf{M}}_T$ by $\widehat{\mathbf{M}}$. For any semi-orthogonal matrix $W \in \mathbb{R}^{K \times L}$, write $W = UA + VB$, where $A \in \mathbb{R}^{L \times L}$ and $B \in \mathbb{R}^{(K-L) \times L}$. Since (U, V) is orthogonal, $A'A + B'B = \mathbf{I}_L$. Let a_1, \dots, a_L be the eigenvalues of $(A'A)^{1/2}$, then $a_i \in [0, 1]$ for each $i = 1, \dots, L$. Denote Δ_A as the diagonal matrix with the i th diagonal element to be a_i , and $V_l, V_r \in \mathbb{R}^{L \times L}$ be the orthogonal matrices such that $A = V_l\Delta_AV_r$. Let $\text{tr}(\cdot)$ be the trace of a square matrix. We have

$$\begin{aligned} \text{tr}(W'\widehat{\mathbf{M}}W) &= \text{tr}(A'U'\widehat{\mathbf{M}}UA + A'U'\widehat{\mathbf{M}}VB + B'V'\widehat{\mathbf{M}}UA + B'V'\widehat{\mathbf{M}}VB) \\ &= \text{tr}(A'U'\widehat{\mathbf{M}}UA) + h(W, \widehat{\mathbf{M}}) = \text{tr}(\Delta'_A V_l' U' \widehat{\mathbf{M}} U V_l \Delta_A) + h(W, \widehat{\mathbf{M}}), \quad (5.4) \end{aligned}$$

where $h(W, \widehat{\mathbf{M}}) = 2tr(A'U'\widehat{\mathbf{M}}VB) + tr(B'V'\widehat{\mathbf{M}}VB)$. Since $A'A + B'B = \mathbf{I}_L$, the eigenvalues of $B'B$ fall into $[0, 1]$, so too are the eigenvalues of BB' , which means that $\mathbf{I}_{K-L} - BB'$ is positive semi-definite. Also, we have $tr(A'A) \leq tr(\mathbf{I}_L) = L$. By simple matrix algebra, we have

$$\begin{aligned} tr^2(A'U'\widehat{\mathbf{M}}VB) &\leq tr(A'U'\widehat{\mathbf{M}}VB(B'V'\widehat{\mathbf{M}}UA)) \leq tr(A'U'\widehat{\mathbf{M}}Vtr(BB')\mathbf{I}_{K-L}V'\widehat{\mathbf{M}}UA)) \\ &\leq tr(A'U'\widehat{\mathbf{M}}VV'\widehat{\mathbf{M}}UA)tr(BB') = tr(A'U'\widehat{\mathbf{M}}VV'\widehat{\mathbf{M}}UA)tr(B'B) \\ &\leq tr(A'\|U'\widehat{\mathbf{M}}V\|^2\mathbf{I}_LA)tr(\mathbf{I}_L - A'A) = L\|V'\widehat{\mathbf{M}}U\|^2tr(\mathbf{I}_L - A'A) \\ &\leq (L - \sum_{i=1}^L a_i^2)O_P(c_T^2). \end{aligned}$$

Thus $tr(A'U'\widehat{\mathbf{M}}VB) = (L - \sum_{i=1}^L a_i^2)^{1/2}O_P(c_T)$. Similarly, we also have

$$tr(B'V'\widehat{\mathbf{M}}VB) \leq tr(B'\|V'\widehat{\mathbf{M}}V\|\mathbf{I}_{K-L}B) = tr(B'B)\|V'\widehat{\mathbf{M}}V\| = (L - \sum_{i=1}^L a_i^2)O_P(c_T).$$

Thus $(L - \sum_{i=1}^L a_i^2)^{-1/2}h(W, \widehat{\mathbf{M}}) = O_P(c_T)$ uniformly for (A, B) . Denote $tr(\Delta'_A V'_i U' \widehat{\mathbf{M}} U V_i \Delta_A)$ by $f(a_1, \dots, a_L, V_i)$. For each V_i , we write $V'_i U' \widehat{\mathbf{M}} U V_i = (b_{ij})$ and let $(\widehat{\lambda}_1, \dots, \widehat{\lambda}_L)$ be its eigenvalues in the descending order. Then since V_i is orthogonal, by condition (a), $\widehat{\lambda}_L = O_P^+(1)$. As $f(a_1, \dots, a_L, V_i)$ can be written as $\sum_i b_{ii} a_i^2$, for each fixed V_i , it has the unique maximizer $(1, \dots, 1)$ with the corresponding partial derivative

$$\partial f / \partial a_i |_{(a_1, \dots, a_L)} = 2b_{ii} = 2e'_i (V'_i U' \widehat{\mathbf{M}} U V_i) e_i \geq 2\widehat{\lambda}_L = O_P^+(1)$$

for $i = 1, \dots, L$, where $e_i \in \mathbb{R}^L$ has 1 on the i th component and 0 otherwise. On the other hand, fixing A , we maximize $tr(W' \widehat{\mathbf{M}} W)$ over B under the constraint that $A'A + B'B = \mathbf{I}_L$, and denote the resulting function as $g(a_1, \dots, a_L, V_i)$. By (5.4), $g(a_1, \dots, a_L, V_i) = f(a_1, \dots, a_L, V_i) + O_P(c_T)$ uniformly for (A, B) . Since \widehat{U}_T maximizes $tr(W' \widehat{\mathbf{M}} W)$, denote $(\widehat{a}_1, \dots, \widehat{a}_L, \widehat{V}_i)$ as the corresponding maximizer of $g(a_1, \dots, a_L, V_i)$, we have

$$f(\widehat{a}_1, \dots, \widehat{a}_L, \widehat{V}_i) - f(1, \dots, 1, \widehat{V}_i) = g(\widehat{a}_1, \dots, \widehat{a}_L, \widehat{V}_i) - g(1, \dots, 1, \widehat{V}_i) + (L - \sum_{i=1}^L \widehat{a}_i^2)^{1/2} O_P(c_T).$$

Since $f(\widehat{a}_1, \dots, \widehat{a}_L, \widehat{V}_i) - f(1, \dots, 1, \widehat{V}_i) \leq 0$ and $g(\widehat{a}_1, \dots, \widehat{a}_L, \widehat{V}_i) - g(1, \dots, 1, \widehat{V}_i) \geq 0$, it implies

$$f(\widehat{a}_1, \dots, \widehat{a}_L, \widehat{V}_i) - f(1, \dots, 1, \widehat{V}_i) = (L - \sum_{i=1}^L \widehat{a}_i^2)^{1/2} O_P(c_T).$$

Fixing V_l at \widehat{V}_l , since f is uniquely maximized at $(1, \dots, 1)$ with all the partial derivatives to be $O_P^+(1)$, we have $\widehat{a}_i = 1 + O_P(c_T^2)$ for $i = 1, \dots, L$. Thus

$$\begin{aligned}
\|\widehat{U}_T \widehat{U}_T' - UU'\|_F &= \|UAA'U' + UAB'V' + VBA'U' + VBB'V' - UU'\|_F \\
&\leq \|UAA'U' - UU'\|_F + 2\|UAB'V'\|_F + \|VBB'V'\|_F \\
&= \|\mathbf{I}_L - A'A\|_F + 2tr^{1/2}(AB'BA') + \|B'B\|_F \\
&= \{\sum_{i=1}^L (1 - \widehat{a}_i^2)^2\}^{1/2} + 2tr^{1/2}(AA' - AA'AA') + \|\mathbf{I}_L - A'A\|_F \\
&= 2\{\sum_{i=1}^L (1 - \widehat{a}_i^2)^2\}^{1/2} + 2\{\sum_{i=1}^L \widehat{a}_i^2 (1 - \widehat{a}_i^2)\}^{1/2} = O_P(c_T).
\end{aligned}$$

This completes the proof. \square

5.5 Proof of Theorem 2.3

The proof can be similarly derived following the proof of Theorem 3 in Li and Wang (2007). Thus, we omit the details for space consideration. \square

5.6 Proof of Theorem 2.4

For simplicity, we assume that the number of observations in each sample slice is the same, i.e. $c_{h-1} \equiv c$ for $h = 1, \dots, H$, without loss of generality. For each $h = 1, \dots, H$, Let $I_{h,T}$ be the h th sample slice of y_{t+1} , i.e. $y_{t+1} \in I_h$ if and only if $y_{t+1}^D = h - 1$, and I_h be the probabilistic limit of $I_{h,T}$ as $T \rightarrow \infty$. For $l = 1, \dots, c$, let $\mathbf{f}_{h,l}^* = \mathbf{f}_{h,l} + \mathbf{\Lambda}_b u_{h,l}$, and denote it by \mathbf{f}_t^* if no slice number is specified. By expanding (2.3) and using the invariance result in Theorem 2.2, we can write as \mathbf{M}_{dr} as $\sum_{h=1}^H \sum_{g=1}^H \mathbf{M}_{h,g}^2 / H^2$, in which

$$\begin{aligned}
\mathbf{M}_{h,g} &= 2\text{var}(\mathbf{f}_t^*) - E(\mathbf{f}_t^* \mathbf{f}_t^{*'} | y_{t+1} \in I_h) - E(\mathbf{f}_t^* \mathbf{f}_t^{*'} | y_{t+1} \in I_g) \\
&\quad + E(\mathbf{f}_t^* | y_{t+1} \in I_h) E(\mathbf{f}_t^{*' | y_{t+1} \in I_g) + E(\mathbf{f}_t^* | y_{t+1} \in I_g) E(\mathbf{f}_t^{*' | y_{t+1} \in I_h). \quad (5.5)
\end{aligned}$$

Similarly, we can write $\widehat{\mathbf{M}}_{dr}$ as $\sum_{h=1}^H \sum_{g=1}^H \widehat{\mathbf{M}}_{h,g}^2 / H^2$, in which

$$\begin{aligned}
\widehat{\mathbf{M}}_{h,g} &= 2\mathbf{I}_K - \sum_{l=1}^c \widehat{\mathbf{f}}_{h,l} \widehat{\mathbf{f}}_{h,l}' / c - \sum_{l=1}^c \widehat{\mathbf{f}}_{g,l} \widehat{\mathbf{f}}_{g,l}' / c \\
&\quad + (\sum_{l=1}^c \widehat{\mathbf{f}}_{h,l} / c) (\sum_{l=1}^c \widehat{\mathbf{f}}_{g,l} / c)' + (\sum_{l=1}^c \widehat{\mathbf{f}}_{g,l} / c) (\sum_{l=1}^c \widehat{\mathbf{f}}_{h,l} / c)'. \quad (5.6)
\end{aligned}$$

We next study the consistency in estimating $\mathbf{M}_{h,g}$. In the estimation of $E(\mathbf{f}_t^* \mathbf{f}_t^{*'} | y_{t+1} \in I_h)$, the error comes in three ways, first, by replacing \mathbf{f}_t^* by $\widehat{\mathbf{f}}_t$, and second, by using the sample slice $I_{h,T}$ instead of its probabilistic limit I_h , and third, by using the sample moment instead of the true moment on each sample slice.

First, we study the effect of replacing \mathbf{f}_t^* by $\widehat{\mathbf{f}}_t$. Let δ_h be such that $\sum_{l=1}^c \widehat{\mathbf{f}}_{h,l} \widehat{\mathbf{f}}_{h,l}' / c = \sum_{l=1}^c \mathbf{f}_{h,l}^* \mathbf{f}_{h,l}^{*'} / c + \delta_h$. Then

$$\begin{aligned}
\|\delta_h\| &= \left\| \sum_{l=1}^c (\widehat{\mathbf{f}}_{h,l} \widehat{\mathbf{f}}_{h,l}' - \mathbf{f}_{h,l}^* \mathbf{f}_{h,l}^{*'}) / c \right\| \\
&\leq \left\| \sum_{l=1}^c (\widehat{\Lambda}_b \mathbf{B} \mathbf{f}_{h,l} \mathbf{f}_{h,l}' \mathbf{B}' \widehat{\Lambda}_b' - \mathbf{f}_{h,l} \mathbf{f}_{h,l}') / c \right\| + 2 \left\| \sum_{l=1}^c (\widehat{\Lambda}_b \mathbf{B} \mathbf{f}_{h,l} u_{h,l}' \widehat{\Lambda}_b' - \mathbf{f}_{h,l} u_{h,l}' \Lambda_b') / c \right\| \\
&\quad + \left\| \sum_{l=1}^c (\widehat{\Lambda}_b u_{h,l} u_{h,l}' \widehat{\Lambda}_b' - \Lambda_b u_{h,l} u_{h,l}' \Lambda_b') / c \right\| \\
&\leq \|\widehat{\Lambda}_b \mathbf{B} - \mathbf{I}_K\| \left\| \sum_{l=1}^c \mathbf{f}_{h,l} \mathbf{f}_{h,l}' / c \right\| (2 + \|\widehat{\Lambda}_b \mathbf{B} - \mathbf{I}_K\|) \\
&\quad + 2 \left\| \sum_{l=1}^c \mathbf{f}_{h,l} u_{h,l}' / c \right\| \{ \|\widehat{\Lambda}_b \mathbf{B} - \mathbf{I}_K\| (\|\Lambda_b\| + \|\widehat{\Lambda}_b - \Lambda_b\|) + \|\widehat{\Lambda}_b - \Lambda_b\| \} \\
&\quad + \|\widehat{\Lambda}_b - \Lambda_b\| \left\| \sum_{l=1}^c u_{h,l} u_{h,l}' / c \right\| (2\|\Lambda_b\| + \|\widehat{\Lambda}_b - \Lambda_b\|) \\
&\equiv I + II + III,
\end{aligned} \tag{5.7}$$

in which the second inequality is derived, for example, by

$$\begin{aligned}
&\left\| \sum_{l=1}^c (\widehat{\Lambda}_b \mathbf{B} \mathbf{f}_{h,l} \mathbf{f}_{h,l}' \mathbf{B}' \widehat{\Lambda}_b' - \mathbf{f}_{h,l} \mathbf{f}_{h,l}') / c \right\| \\
&= \left\| \sum_{l=1}^c (\widehat{\Lambda}_b \mathbf{B} \mathbf{f}_{h,l} \mathbf{f}_{h,l}' \mathbf{B}' \widehat{\Lambda}_b' - \widehat{\Lambda}_b \mathbf{B} \mathbf{f}_{h,l} \mathbf{f}_{h,l}' + \widehat{\Lambda}_b \mathbf{B} \mathbf{f}_{h,l} \mathbf{f}_{h,l}' - \mathbf{f}_{h,l} \mathbf{f}_{h,l}') / c \right\| \\
&\leq \left\| \sum_{l=1}^c (\widehat{\Lambda}_b \mathbf{B} \mathbf{f}_{h,l} \mathbf{f}_{h,l}' \mathbf{B}' \widehat{\Lambda}_b' - \widehat{\Lambda}_b \mathbf{B} \mathbf{f}_{h,l} \mathbf{f}_{h,l}') / c \right\| + \left\| \sum_{l=1}^c (\widehat{\Lambda}_b \mathbf{B} \mathbf{f}_{h,l} \mathbf{f}_{h,l}' - \mathbf{f}_{h,l} \mathbf{f}_{h,l}') / c \right\| \\
&\leq \|\widehat{\Lambda}_b \mathbf{B} - \mathbf{I}_K + \mathbf{I}_K\| \left\| \sum_{l=1}^c \mathbf{f}_{h,l} \mathbf{f}_{h,l}' / c \right\| \|\widehat{\Lambda}_b \mathbf{B} - \mathbf{I}_K\| + \left\| \sum_{l=1}^c \mathbf{f}_{h,l} \mathbf{f}_{h,l}' / c \right\| \|\widehat{\Lambda}_b \mathbf{B} - \mathbf{I}_K\| \\
&\leq \|\widehat{\Lambda}_b \mathbf{B} - \mathbf{I}_K\| \left\| \sum_{l=1}^c \mathbf{f}_{h,l} \mathbf{f}_{h,l}' / c \right\| \|\widehat{\Lambda}_b \mathbf{B} - \mathbf{I}_K\| + 2 \left\| \sum_{l=1}^c \mathbf{f}_{h,l} \mathbf{f}_{h,l}' / c \right\| \|\widehat{\Lambda}_b \mathbf{B} - \mathbf{I}_K\|.
\end{aligned}$$

For ease of presentation, we assume that for each $l = 1, \dots, L$, $\phi_l' \mathbf{f}_t = \mathbf{f}_{l,t}$. Then for any $i = 1, \dots, K$ and $j = L + 1, \dots, K$ such that $j \neq i$, $E(\mathbf{f}_{i,t} \mathbf{f}_{j,t}' | y_{t+1} \in I_{h,T}) = 0$ and $E(\mathbf{f}_{j,t}^2 | y_{t+1} \in I_{h,T}) = 1$. Thus

$$\begin{aligned}
\left\| \sum_{l=1}^c \mathbf{f}_{h,l} \mathbf{f}_{h,l}' / c \right\| &\leq \|E(\mathbf{f}_t \mathbf{f}_t' | y_{t+1} \in I_{h,T})\| + \left\| \sum_{l=1}^c \mathbf{f}_{h,l} \mathbf{f}_{h,l}' / c - E(\mathbf{f}_t \mathbf{f}_t' | y_{t+1} \in I_{h,T}) \right\| \\
&\leq \|E(\mathbf{f}_t \mathbf{f}_t' | y_{t+1} \in I_{h,T})\| + \left\| \sum_{l=1}^c \mathbf{f}_{h,l} \mathbf{f}_{h,l}' / c - E(\mathbf{f}_t \mathbf{f}_t' | y_{t+1} \in I_{h,T}) \right\|_F \\
&= \max\{1, \|E\{(\mathbf{f}_{1,t}, \dots, \mathbf{f}_{L,t})(\mathbf{f}_{1,t}, \dots, \mathbf{f}_{L,t})' | y_{t+1} \in I_{h,T}\}\| \} + O_P(KT^{-1/2}) \\
&\leq \max\{1, \|HE\{(\mathbf{f}_{1,t}, \dots, \mathbf{f}_{L,t})(\mathbf{f}_{1,t}, \dots, \mathbf{f}_{L,t})'\}\| \} + O_P(KT^{-1/2}) \\
&= H + O_P(KT^{-1/2}) = O_P(1),
\end{aligned}$$

in which the first inequality is based on the triangle inequality and the first equality is based on Assumption 2.2 and the central limit theorem (Billingsley, 1999, Theorem 19.2) By Lemma 5.1 and Lemma 2.1, we have

$$\|\widehat{\Lambda}_b \mathbf{B} - \mathbf{I}_K\| \leq \|\widehat{\Lambda}_b - \Lambda_b\| \|\mathbf{B}\| = O_P(K^{3/2}p^{-1/2} + K^{1/2}T^{-1/2}).$$

Thus $I = O_P(K^{3/2}p^{-1/2} + K^{1/2}T^{-1/2})$. Since $\Lambda_b \Lambda'_b = (\mathbf{B}'\mathbf{B})^{-1}$, again by Assumption 2.1, we have $\|\Lambda_b\| = \lambda_{max}^{1/2}((\mathbf{B}'\mathbf{B})^{-1}) = O(p^{-1/2})$. Let $tr(\cdot)$ be the trace of a square matrix. By Assumption 2.3, and the independence between $u_{h,l}$ and $\mathbf{f}_{h,l}$, and Markov's inequality, we have

$$\begin{aligned} \|\sum_{l=1}^c u_{h,l} \mathbf{f}'_{h,l}/c\|^2 &\leq tr(\sum_{l=1}^c u_{h,l} \mathbf{f}'_{h,l}/c \sum_{l=1}^c \mathbf{f}_{h,l} u'_{h,l}/c) \\ &= \sum_{i=1}^p (\sum_{l=1}^c u_{i,h,l} \mathbf{f}'_{h,l}/c \sum_{l=1}^c \mathbf{f}_{h,l} u'_{i,h,l}/c) \\ &\leq c^{-2} \sum_{i=1}^p \|\sum_{t=1}^T u_{it} \mathbf{f}_t\|^2 = O_P(KpT^{-1}). \end{aligned}$$

Thus $II = O_P(K^{1/2}T^{-1/2}(K^{3/2}p^{-1/2} + K^{1/2}T^{-1/2})) = O_P(K^{3/2}p^{-1/2} + K^{1/2}T^{-1/2})$. By Assumption 2.3 again, $\|\sum_{l=1}^c u_{h,l} u'_{h,l}/c\| = O_P(p^{1/2})$, thus $III = O_P(p^{-1/2}(K^{3/2}p^{-1/2} + K^{1/2}T^{-1/2}))$. In summary, we have

$$\|\delta_h\| = O_P(K^{3/2}p^{-1/2} + K^{1/2}T^{-1/2}).$$

Next, we study the effect of using the sample slices instead of the true slices, and further using the sample moments instead of the true moments on each sample slice. We have

$$\begin{aligned} &\|\sum_{l=1}^c \mathbf{f}_{h,l}^* \mathbf{f}_{h,l}^{*'} / c - E(\mathbf{f}_t^* \mathbf{f}_t^{*'} | y_{t+1} \in I_h)\| \\ &= \|\sum_{l=1}^c \mathbf{f}_{h,l}^* \mathbf{f}_{h,l}^{*'} / c - E(\mathbf{f}_t^* \mathbf{f}_t^{*'} | y_{t+1} \in I_{h,T})\| \\ &\quad + \|E(\mathbf{f}_t^* \mathbf{f}_t^{*'} | y_{t+1} \in I_{h,T}) - E(\mathbf{f}_t^* \mathbf{f}_t^{*'} | y_{t+1} \in I_h)\| \\ &= c^{-1/2} O_P(E^{1/2}(\sum_{i=1}^K \sum_{j=1}^K \mathbf{f}_{it}^2 \mathbf{f}_{jt}^2 | y_{t+1} \in I_{h,T})) \\ &\quad + O_P(\|(\mathbf{f}_{1t}, \dots, \mathbf{f}_{Lt})'(\mathbf{f}_{1t}, \dots, \mathbf{f}_{Lt})\|) O_P(I(y_{t+1} \in I_{h,T}) - I(y_{t+1} \in I_h)) \\ &= O_P(KT^{-1/2}) + O_P(T^{-1/2}). \end{aligned}$$

in which the order of the second term is due to Bahadur representation theorem. Note that the first term above represents the estimation error due to the use of sample moments, which would persist even if B were known a priori.

Denote $\Delta_{0,h}$ as $\sum_{l=1}^c \widehat{\mathbf{f}}_{h,l} \widehat{\mathbf{f}}'_{h,l}/c - E(\mathbf{f}_t^* \mathbf{f}_t^{*'} | y_{t+1} \in I_h)$. Then combining these results together, we have

$$\|\Delta_{0,h}\| = O_P(K^{3/2}p^{-1/2} + K^{1/2}T^{-1/2} + KT^{-1/2}) = O_P(K^{3/2}p^{-1/2} + KT^{-1/2}). \quad (5.8)$$

Using similar arguments, we have

$$\begin{aligned} \|\sum_{t=1}^T \widehat{\mathbf{f}}_t \widehat{\mathbf{f}}'_t / T - \text{var}(\mathbf{f}_t^*)\| &= O_P(K^{3/2}p^{-1/2} + KT^{-1/2}), \\ \|\sum_{l=1}^c \widehat{\mathbf{f}}_{h,l}/c - \sum_{l=1}^c \mathbf{f}_{h,l}^*/c\| &= O_P(K^2p^{-1/2}T^{-1/2} + KT^{-1}), \\ \|\sum_{l=1}^c \mathbf{f}_{h,l}^*/c - E(\mathbf{f}_t^* | y_{t+1} \in I_h)\| &= O_P(K^{1/2}T^{-1/2}), \end{aligned}$$

in which the latter two equations imply that

$$\|\sum_{l=1}^c \widehat{\mathbf{f}}_{h,l}/c - E(\mathbf{f}_t^* | y_{t+1} \in I_h)\| = O_P(K^2p^{-1/2}T^{-1/2} + K^{1/2}T^{-1/2}). \quad (5.9)$$

Denote $\Delta_{1,h}$ as $\sum_{l=1}^c \widehat{\mathbf{f}}_{h,l}/c - E(\mathbf{f}_t^* | y_{t+1} \in I_h)$. By plugging (5.8) and (5.9) into (5.5) and (5.6), we have

$$\begin{aligned} \widehat{\mathbf{M}}_{h,g} &= \mathbf{M}_{h,g} - 2\mathbf{\Lambda}_b \Sigma_u \mathbf{\Lambda}'_b + [E(\mathbf{f}_t^* | y_{t+1} \in I_h) \Delta'_{1,g} + \Delta_{1,g} E(\mathbf{f}_t^{*'} | y_{t+1} \in I_h)] \\ &\quad + [E(\mathbf{f}_t^* | y_{t+1} \in I_g) \Delta'_{1,h} + \Delta_{1,h} E(\mathbf{f}_t^{*'} | y_{t+1} \in I_g)] \\ &\quad + (\Delta_{1,h} \Delta'_{1,g} + \Delta_{1,g} \Delta'_{1,h} - \Delta_{0,h} - \Delta_{0,g}) \\ &\equiv \mathbf{M}_{h,g} + \Delta_{2,h,g}, \end{aligned} \quad (5.10)$$

By Assumption 2.3, $\|\Sigma_u\| = \lambda_{\max}(\Sigma_u) \leq \text{tr}(\Sigma_u) \leq \|\Sigma_u\|_1 = O(1)$. Thus, $\|\mathbf{\Lambda}_b \Sigma_u \mathbf{\Lambda}'_b\| \leq \|\mathbf{\Lambda}_b\|^2 \|\Sigma_u\| = O(p^{-1})$. Also, since $E(\mathbf{f}_t | y_{t+1} \in I_h) = (E(\mathbf{f}_{1t}, \dots, \mathbf{f}_{Lt} | y_{t+1} \in I_h), 0, \dots, 0)'$, we have

$$\begin{aligned} \|E(\mathbf{f}_t^* | y_{t+1} \in I_h)\| &= \{\sum_{i=1}^K E^2(\mathbf{f}_{it} | y_{t+1} \in I_h)\}^{1/2} = \{\sum_{i=1}^L E^2(\mathbf{f}_{it} | y_{t+1} \in I_h)\}^{1/2} \\ &\leq [\sum_{i=1}^L E(\mathbf{f}_{it}^2 | y_{t+1} \in I_h)]^{1/2} \leq \{\sum_{i=1}^L HE(\mathbf{f}_{it}^2)\}^{1/2} = O(1). \end{aligned} \quad (5.11)$$

Thus $\|\Delta_{2,h,g}\| = O_P(K^{3/2}p^{-1/2} + KT^{-1/2})$. By taking the square of both sides of (5.10) and averaging over (h, g) , we have

$$\begin{aligned} \widehat{\mathbf{M}}_{dr} &= \mathbf{M}_{dr} + \sum_{l=1}^H \sum_{g=1}^H [\mathbf{M}_{h,g} \Delta_{2,h,g} + \Delta_{2,h,g} \mathbf{M}_{h,g} + \Delta_{2,h,g}^2] \\ &\equiv \mathbf{M}_{dr} + \Delta_3. \end{aligned} \quad (5.12)$$

Since $\|E(\mathbf{f}_t \mathbf{f}'_t | y_{t+1} \in I_h)\| \leq \|HE(\mathbf{f}_t \mathbf{f}'_t)\| = O_P(1)$, together with (5.11), we have

$$\begin{aligned} \|\mathbf{M}_{h,g}\| &\leq 2\|\mathbf{I}_K\| + \|E(\mathbf{f}_t \mathbf{f}'_t | y_{t+1} \in I_h)\| + \|E(\mathbf{f}_t \mathbf{f}'_t | y_{t+1} \in I_g)\| \\ &\quad + \|E(\mathbf{f}_t | y_{t+1} \in I_h)\|_F^2 + \|E(\mathbf{f}_t | y_{t+1} \in I_g)\|_F^2 \\ &= 2 + O_P(1) + O_P(1) + \sum_{i=1}^L E(\mathbf{f}_{it}^2 | y_{t+1} \in I_h) + \sum_{i=1}^L E(\mathbf{f}_{it}^2 | y_{t+1} \in I_g) \\ &\leq O_P(1) + 2H \sum_{i=1}^L E(\mathbf{f}_{it}^2) = O_P(1). \end{aligned}$$

Thus $\|\Delta_3\| = O_P(K^{3/2}p^{-1/2} + KT^{-1/2})$. Let $\Phi = (\phi_1, \dots, \phi_L)$ be arranged such that each ϕ_i is an eigenvector of \mathbf{M}_{dr} and the corresponding eigenvalues $\{\lambda_1, \dots, \lambda_L\}$ have an non-increasing order. Let Ψ be such that (Φ, Ψ) is an orthogonal matrix. Treating (Φ, Ψ) as (U, V) and $\widehat{\mathbf{M}}_{dr}$ as $\widehat{\mathbf{M}}$ in Lemma 2.1, and $c_T = K^{3/2}p^{-1/2} + KT^{-1/2}$, we next verify conditions (a) and (b) in Lemma 2.1. Let $\widehat{\phi}_{\min}$ be the eigenvector of $\Phi' \widehat{\mathbf{M}}_{dr} \Phi$ that corresponds to $\lambda_{\min}(\Phi' \widehat{\mathbf{M}}_{dr} \Phi)$. By (5.12), we have

$$\lambda_{\min}(\Phi' \widehat{\mathbf{M}}_{dr} \Phi) = \widehat{\phi}'_{\min} \Phi' \mathbf{M}_{dr} \Phi \widehat{\phi}_{\min} + \widehat{\phi}'_{\min} \Phi' \Delta_3 \Phi \widehat{\phi}_{\min} \geq \lambda_L - \|\Phi \widehat{\phi}_{\min}\|^2 \|\Delta_3\| = \lambda_L + o_P(1).$$

By the condition in the theorem, we have $\lambda_L = O_P^+(1)$, which means $\lambda_{\min}(\Phi' \widehat{\mathbf{M}}_{dr} \Phi) = O_P^+(1)$. Thus condition (a) in Lemma 2.1 is satisfied. Since

$$\|\Phi' \widehat{\mathbf{M}}_{dr} \Psi\| = \|\Phi' \Delta_3 \Psi\| \leq \|\Delta_3\| = O_P(c_T),$$

and similarly $\|\Psi' \widehat{\mathbf{M}}_{dr} \Psi\| = O_P(c_T)$, Lemma 2.1 implies $\|\widehat{\Phi} \widehat{\Phi}' - \Phi \Phi'\|_F = O_P(c_T)$. This completes the proof. \square

5.7 Proof of Theorem 2.5

For simplicity in notations, we write the first term and the second term on the right-hand side of (2.5) as $W(l)$ and $Q(l)$, respectively, so that $G(l) = W(l) + Q(l)$. By Weyl's Theorem, for each $i = 1, \dots, K_c$, $|\widehat{\lambda}_i - \lambda_i| \leq \|\widehat{\mathbf{M}} - \mathbf{M}\| = o_P(1)$. Since $\lambda_i = O_P^+(1)$ when $i \leq L$, we have $\widehat{\lambda}_i = O_P^+(1)$. Thus $\tau > L$ in probability, and we can assume that $\tau > L$ without loss of generality. For any $l < L$,

$$W(L) - W(l) = -(T/2) \sum_{i=l+1}^L \{\log(\widehat{\lambda}_i + 1) - \widehat{\lambda}_i\} = O_P^+(T).$$

Since $Q(L) - Q(l) = O(C_T K) = o(T)$, we have $G(L) - G(l) > 0$ in probability. Since L is fixed, it further implies that

$$P(G(L) > \max\{G(l) : l = 1, \dots, L-1\}) \rightarrow 1. \quad (5.13)$$

For any $l > L$ such that $l < K_c$, if $\tau > l-1$, then $W(l) - W(l-1) = -(T/2)\{\log(\widehat{\lambda}_l + 1) - \widehat{\lambda}_l\}$. Otherwise, $W(l) - W(l-1) = 0$. Since the function $f : \mathbb{R}^+ \rightarrow \mathbb{R}$, $f(x) = \log(x+1) - x$ is monotone decreasing and $f'(0) = 0$ and $f''(0) = -1$, we have

$$\begin{aligned} \max\{W(l) - W(l-1) : L < l \leq K_c\} &= -(T/2)\{\log(\widehat{\lambda}_{L+1} + 1) - \widehat{\lambda}_{L+1}\} \\ &= O_P(T\|\widehat{\mathbf{M}} - \mathbf{M}\|^2) = o_P(C_T K). \end{aligned}$$

Since $Q(l-1) - Q(l) = C_T(K-l+1) \geq C_T(K-K_c) \geq C_T K(1-c)/2$, we have

$$\begin{aligned} P(G(L) > \max\{G(l) : L < l \leq K_c\}) \\ &\geq P(\max\{G(l) - G(l-1) : L < l \leq K_c\} < 0) \\ &\geq P(\max\{W(l) - W(l-1) : L < l \leq K_c\} < C_T K(1-c)/2) \rightarrow 1. \end{aligned} \quad (5.14)$$

Combined with (5.13), $P(G(L) = \max\{G(l) : l = 1, \dots, K_c\}) \rightarrow 1$. This completes the proof. \square

References

- Ahn, S. C. and Horenstein, A. R. (2013), ‘Eigenvalue ratio test for the number of factors’, *Econometrica* **81**(3), 1203–1227.
- Bai, J. and Ng, S. (2008), ‘Forecasting economic time series using targeted predictors’, *Journal of Econometrics* **146**, 304–317.
- Bernanke, B. S., Boivin, J. and Eliasch, P. (2005), ‘Measuring the effects of monetary policy: a factor-augmented vector autoregressive (favar) approach’, *The Quarterly Journal of Economics* **120**(1), 387–422.
- Billingsley, P. (1999), *Convergence of Probability Measures*, 2nd edn, John Wiley & Sons.

- Cai, T. and Liu, W. (2011), ‘Adaptive thresholding for sparse covariance matrix estimation’, *Journal of the American Statistical Association* **106**(494), 672–684.
- Cook, R. D. (1998), *Regression Graphics*, Wiley, New York.
- Cook, R. D. and Lee, H. (1999), ‘Dimension reduction in binary response regression’, *Journal of the American Statistical Association* **94**(448), 1187–1200.
- Cook, R. D. and Weisberg, S. (1991), ‘Comment on “sliced inverse regression for dimension reduction (with discussion)”’, *Journal of the American Statistical Association* **86**(414), 328–332.
- Fan, J., Liao, Y. and Mincheva, M. (2013), ‘Large covariance estimation by thresholding principal orthogonal complements (with discussion)’, *Journal of the Royal Statistical Society: Series B* **75**(4), 603–680.
- Fan, J., Xue, L. and Yao, J. (2016), ‘Sufficient forecasting using factor models’, *Journal of Econometrics*, to appear .
- Fan, J., Xue, L. and Zou, H. (2016), ‘Multi-task quantile regression under the transnormal model’, *Journal of the American Statistical Association* **111**(516), 1726–1735.
- Hall, P. and Li, K.-C. (1993), ‘On almost linearity of low dimensional projections from high dimensional data’, *The Annals of Statistics* **21**(2), 867–889.
- Jiang, B. and Liu, J. S. (2014), ‘Variable selection for general index models via sliced inverse regression’, *The Annals of Statistics* **42**(5), 1751–1786.
- Jurado, K., Ludvigson, S. C. and Ng, S. (2015), ‘Measuring uncertainty’, *The American Economic Review* **105**(3), 1177–1216.
- Lam, C. and Yao, Q. (2012), ‘Factor modeling for high-dimensional time series: inference for the number of factors’, *The Annals of Statistics* **40**(2), 694–726.
- Li, B. and Wang, S. (2007), ‘On directional regression for dimension reduction’, *Journal of the American Statistical Association* **102**(479), 997–1008.
- Li, B. and Yin, X. (2007), ‘On surrogate dimension reduction for measurement error regression: an invariance law’, *The Annals of Statistics* **35**(5), 2143–2172.

- Li, H., Li, Q. and Shi, Y. (2013), ‘Determining the number of factors when the number of factors can increase with sample size’, *Manuscript* .
- Li, K.-C. (1991), ‘Sliced inverse regression for dimension reduction (with discussion)’, *Journal of the American Statistical Association* **86**(414), 316–327.
- Lintner, J. (1965), ‘The valuation of risk assets and the selection of risky investments in stock portfolios and capital budgets’, *The Review of Economics and Statistics* **47**(1), 13–37.
- Ludvigson, S. and Ng, S. (2007), ‘The empirical risk return relation: A factor analysis approach’, *Journal of Financial Economics* **83**(1), 171–222.
- Ludvigson, S. and Ng, S. (2009), ‘Macro factors in bond risk premia’, *Review of Financial Studies* **22**(12), 5027–5067.
- Luo, W. and Li, B. (2016), ‘Combining eigenvalues and variation of eigenvectors for order determination’, *Biometrika* **103**(4), 875–887.
- McCracken, M. W. and Ng, S. (2016), ‘Fred-md: A monthly database for macroeconomic research’, *Journal of Business & Economic Statistics* **34**(4), 574–589.
- Onatski, A. (2010), ‘Determining the number of factors from empirical distribution of eigenvalues’, *The Review of Economics and Statistics* **92**(4), 1004–1016.
- Sharpe, W. F. (1964), ‘Capital asset prices: a theory of market equilibrium under conditions of risk’, *Journal of Finance* **19**(3), 425–442.
- Stock, J. H. and Watson, M. W. (1989), ‘New indexes of coincident and leading economic indicators’, *NBER Macroeconomics Annual* **4**, 351–409.
- Stock, J. H. and Watson, M. W. (2002a), ‘Forecasting using principal components from a large number of predictors’, *Journal of the American Statistical Association* **97**(460), 1167–1179.
- Stock, J. H. and Watson, M. W. (2002b), ‘Macroeconomic forecasting using diffusion indexes’, *Journal of Business & Economic Statistics* **20**(2), 147–162.
- Wang, H. and Xia, Y. (2008), ‘Sliced regression for dimension reduction’, *Journal of the American Statistical Association* **103**(482), 811–821.

- Xia, Y., Tong, H., Li, W. and Zhu, L.-X. (2002), ‘An adaptive estimation of dimension reduction space (with discussion)’, *Journal of the Royal Statistical Society Series B* **64**(3), 363–410.
- Xue, L., Ma, S. and Zou, H. (2012), ‘Positive definite ℓ_1 penalized estimation of large covariance matrices’, *Journal of the American Statistical Association* **107**(500), 1480–1491.
- Ye, Z. and Weiss, R. E. (2003), ‘Using the bootstrap to select one of a new class of dimension reduction methods’, *Journal of the American Statistical Association* **98**(464), 968–979.
- Yin, X. and Cook, R. D. (2003), ‘Estimating central subspaces via inverse third moments’, *Biometrika* **90**(1), 113–125.
- Yu, Z., Dong, Y. and Zhu, L.-X. (2016), ‘Trace pursuit: A general framework for model-free variable selection’, *Journal of the American Statistical Association* **111**(514), 813–821.
- Zhu, L., Miao, B. and Peng, H. (2006), ‘On sliced inverse regression with high-dimensional covariates’, *Journal of the American Statistical Association* **101**(474), 630–643.