arXiv:1705.01005v1 [cond-mat.str-el] 2 May 2017

$CP^{1|1}$ nonlinear sigma model vs strong electron correlations

E.A. Kochetov¹ and A. $Ferraz^2$

¹International Institute of Physics - UFRN, Natal, Brazil;

Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, 141980 Dubna, Russia

²International Institute of Physics - UFRN, Department of Experimental and Theoretical Physics - UFRN, Natal, Brazil

The nonlinear sigma model targeted on the coset supermanifold $CP^{1|1} = SU(2|1)/U(1|1)$ is derived in an attempt to describe the quasiclassical low-energy effective action for the doped t - Jmodel at the SUSY point, J = 2t. In spite of the fact that the supermanifold $CP^{1|1}$ indeed appears as the phase space of the strongly correlated electrons, the canonical $CP^{1|1}$ nonlinear sigma model (NLSM) is unable to capture the physics of strong correlations displayed by the SUSY t - J model at any finite doping. This is due to the fact that, in this regime, the doping itself cannot be self-consistently incorporated into the $CP^{1|1}$ NLSM.

Keywords: t - J model of strongly correlated electrons, su(2|1) coherent state, $CP^{1|1}$ nonlinear sigma model

I. INTRODUCTION: t - J MODEL OF STRONGLY CORRELATED ELECTRONS

As is well known, a d - dimensional quantum antiferromagnetic (AF) Heisenberg model in the large-spin limit at zero temperature can be mapped onto a d + 1 classical NLSM targeted on the projected space $CP^1 = SU(2)/U(1)$. This mapping is fully controlled by 1/s expansion. In 1d, a topological term also emerges to discriminate between the spectrum of the low-energy excitations for integer and half-integer spin values s [1]. This term produces an interference between the different topological sectors for a half-integer s. As a result, gapless low energy excitations emerge out of that. For integer s, the topological term does not affect the path integral and the model displays gapful excitations instead. Although the semiclassical approach is strictly speaking valid for a large s only, the qualitative distinction between the integer and half-integer chains holds true down to the smallest possible spin magnitude, s = 1/2. In other words the large-s expansion gives a qualitatively correct picture down to the physical value of s = 1/2.

To include doping, a natural guess might be that a doped quantum AF model, in the low-energy quasiclassical limit, would admit a mapping onto the $CP^{1|1}$ NLSM which in its turn can be viewed as a natural extension of the CP^1 NLSM to incorporate extra fermionic degrees of freedom to describe the doped holes. The aim of the present paper is to show that this is not what happens for a doped quantum AF described by the t - J model. Because of the strong correlations inherent of the hopping lattice electrons due to a large on-site Coulomb repulsion, strongly competing phases emerge at nonzero doping. These are not captured by the semiclassics that substantially destroys strong correlations. This in turn results in qualitatively different physics for large versus small values of the su(2|1)representation index.

To set the stage, let us start with some definitions. The t - J model of correlated electrons is a lattice model on the restricted $3^{\mathcal{N}}$ - dimensional electronic Hilbert space (\mathcal{N} -is a number of the lattice sites), where the occurrence of two electrons on the same lattice site is strictly forbidden. This restriction comes from the strong on-site Coulomb repulsion between the electrons which drive the strong correlations. Explicitly, the t - J model Hamiltonian reads

$$H_{t-J} = -t \sum_{ij\sigma} \tilde{c}_{i\sigma}^{\dagger} \tilde{c}_{j\sigma} + J \sum_{ij} (\vec{Q}_i \cdot \vec{Q}_j - \frac{1}{4} \tilde{n}_i \tilde{n}_j), \tag{1}$$

where $\tilde{c}_{i\sigma} = c_{i\sigma}(1 - n_{i,-\sigma})$ is the Gutzwiller projected electron operator (to avoid the on-site double occupancy), $\vec{Q}_i = \sum_{\sigma,\sigma'} \tilde{c}^{\dagger}_{i\sigma} \vec{\tau}_{\sigma\sigma'} \tilde{c}_{i\sigma'}, \vec{\tau}^2 = 3/4$, is the electron spin operator and $\tilde{n}_i = n_{i\uparrow} + n_{i\downarrow} - 2n_{i\uparrow}n_{i\downarrow}$. It contains the hopping term $\sim t$ and the spin exchange term $\sim J$. The summation is extended over the nearest neighbour (nn) sites of a *d*-dimensional bipartite lattice, $L = A \oplus B$. Here if $i \in A$ then all its nn sites belong to the sublattice *B*. For the hole-doped cuprates, $t \approx 0.5$ ev and $J \approx t/3$.

The implementation of the no double occupancy (NDO) constraint drives the system towards the strong coupling regime[2]. This is a key point that makes the problem essentially non-perturbative. Because of the NDO constraint, the hopping term cannot be taken as a bare free interaction to derive a perturbation expansion in J/t. In fact, it can be diagonalized only in 1d. In higher dimensions, the hopping constrained electrons exhibit a nontrivial physics (e.g., the Nagaoka phase to describe a single hole doped into a 2d lattice of the hopping constrained electrons [3].) In the t-J model, the effective charge and spin degrees of freedom are entangled due to those strong electron correlations.

In view of the non-perturbative nature of strong electron correlations any reliable approach is significantly important. In particular, a considerable simplification is provided by the supersymmetric variant of the t - J model. Namely, in d = 1, the t - J model is exactly solvable at the supersymmetric (SUSY) point, J = 2t [4]. At this point, it can be brought into a bilinear form of the generators of the su(2|1) superalgebra (also called spl(2, 1) in the literature) in the 3d fundamental degenerate irreducible representation (irrep). Such a system exhibits a global SU(2|1) supersymmetry which makes it solvable by Bethe ansatz in 1d. Away from the SUSY point, the symmetry group reduces to $SU(2) \times U(1)$ which leads to the separate conservation of the total electron spin and the total number of doped vacancies as is physically appropriate for a doped quantum Heisenberg model. Note that the $SU(2) \times U(1)$ group appears as an even subgroup of SU(2|1).

A complementary theoretical method which is not based on a standard perturbation theory can also be established if we explore the effects of quantum and thermal fluctuations in the semiclassical limit. It proves useful in describing the physics at large values of the relevant representation indices of a global symmetry group. A natural question then arises as to whether the t - J model admits a reliable semiclassical treatment that truly preserves strong correlations. After all, this is precisely the case at half-filling ($\tilde{n}_i = 1$) at which the t - J model reduces to the SU(2) invariant antiferromagnetic (AF) Heisenberg spin model.

The su(2|1) algebraic approach provides two possible generalizations of the standard t - J Hamiltonian to include particles with spin higher than 1/2, which is a necessary step to properly formulate a semiclassical expansion. One possibility might be to consider N electron orbitals at each site, which would correspond to the fundamental representation of the su(N|1) superalgebra instead of su(2|1). One can then consider a fully antisymmetric (slave-boson) [5], fully symmetric (slave-fermion), or mixed (L-shaped) representations of su(N|1) [6]. There appears an overall multiplier $\sim N$ in a corresponding path-integral action, which formally justifies a subsequent 1/N expansion. It is important to emphasize, however, that the uniform saddle point, whenever it is stable, describes a Fermi liquid up to very small values of doping [5]. This is a qualitatively incorrect picture, since at small doping strong correlation are known to display a manifestly non-Fermi liquid behaviour. Therefore the large-N expansion does not converge to the physical value N = 2. As a result, strong correlations are suppressed within the proposed 1/N expansion.

An alternative procedure is to interpret the holes to be sites which have spin s = q - 1/2 where $q \in k/2$ and k is an integer.[7]. In view of that, the sites without a "hole" acquire a spin (q - 1/2) + 1/2 = q. The latter possibility amounts to considering the more general higher-spin (q, q) representation of su(2|1) rather than the (q = 1/2, q = 1/2)fundamental one. A total action is proportional to the representation index q >> 1, which implies that a large-qexpansion may be applicable in that case. However, there is a severe technical problem in explicitly carrying out the 1/q expansion. In the su(2|1) coherent-state path integral approach, the spin-charge entanglement is encoded in the SU(2|1) invariant path-integral measure: it does not admit a decomposition into a product of a pure SU(2)spin and U(1) fermionic pieces. The bosonic and fermionic fields are intertwined in the SU(2|1) invariant measure in a nontrivial manner which makes, except in a few trivial instances, an explicit computation of the pertinent path integral rather problematic [8].

To simplify the matter, we restrict ourselves to the case of the maximal possible global symmetry of the t - JHamiltonian exhibited at the SUSY point, J = 2t. Although this simplifies the model considerably, strong electron correlations are still at work. The spin-charge entanglement due to strong correlations manifests itself by means of the even (representing spin) and odd (representing charge) generators closed into a unique superalgebra, su(2|1), so that the charge and spin degrees of freedom transform themselves under the SU(2|1) action through each other. Technically, the semiclassical approach to the SUSY t - J model follows the steps similar to those encoded in the Haldane conjecture in dealing with the quasiclassical Heisenberg su(2) quantum spin model. However, the su(2|1)superalgebra brings into consideration a few qualitative new features. In contrast with the su(2) spin case for which any irrep is equivalent to its conjugate, the su(2|1) superalgebra admits two sets of inequivalent conjugate representations. One can either place the su(2|1) generators in *different* representations associated with two different sublattices, or use a single irrep all over the full lattice.

We start with the semiclassical effective theory for the SUSY spin model based on the so-called alternating group representations. In this case the fundamental and the conjugate su(2|1) representations are placed on different neighbouring sites. Such a model has earlier been argued to produce a canonical $CP^{1|1}$ NLSM. It was written down in Ref.[9] although no explicit derivation based on a microscopical model was provided in their work.

Within the second option, we fix a classical ground state to be realized by the generalized AF Neel superspin configuration. We explicitly derive the relevant quasiclassical effective action to specify the entering coupling constants. Such a model turns out to be given by the same $CP^{1|1}$ NLSM.

However both approaches are shown to fail to accommodate the strongly correlated electron states at finite doping. As a result, the physics exposed by the canonical NLSM is unable to resolve the theoretical challenge of properly describing strongly correlated metallic state in the vicinity of an insulating regime as a function of doping. Although the present derivation may indeed be relevant for some physical models, the generalized AF ansatz does not admit an interpretation in terms of the Hubbard operators at finite doping. This by far rules out the possibility that the canonical $CP^{1|1}$ NLSM may be considered as a relevant low-energy quasiclassical action to describe the physics of the t - J model at finite doping.

A novel point that helps us to derive the continuum limit of the SUSY t-J model is that we express the lattice action

entirely in terms of the projectors onto the underlying classical phase space – the su(2|1) coherent-state manifold. This allows us to simplify the calculations and in addition to provide a universal approach that produces a straightforward generalization to treat other models with Hamiltonians constructed out of the generators of Lie (super)algebras. The overlap of the projectors is directly related to the important geometric structure – the Kaehler (super)potential of the target (super)manifold. In the continuum limit, this overlap reduces to the metric function in terms of which the corresponding NLSM emerges. Such an approach is quite a general one: It basically involves only one essential ingredient that determines the local geometry of the coherent-state manifold – the Kaehler (super)potential.

II. SUPERSETTING

In this Section, we briefly review the basic facts concerning the SU(2|1) supergroup and the supersymmetric t - J model.

A. su(2|1) superalgebra & su(2|1) coherent states

Our conventions for the SU(2|1) supergroup are summarized as follows. The SU(2|1) supergroup in the fundamental representation is the group of $(2 + 1) \times (2 + 1)$ unitary, unimodular supermatrices with the Hermitian conjugate operation. It is generated by even and odd generators, $\{B, Q_3, Q_+, Q_-\}$ and $\{W_+, W_-, V_+, V_-\}$, respectively, which satisfy the following commutation rules [10]:

$$\begin{split} [Q_3,Q_{\pm}] &= \pm Q_{\pm}, [Q_+,Q_-] = 2Q_3, [B,Q_{\pm}] = [B,Q_3] = 0, \\ [B,V_{\pm}] &= \frac{1}{2}V_{\pm}, [B,W_{\pm}] = -\frac{1}{2}W_{\pm}, [Q_3,V_{\pm}] = \pm \frac{1}{2}V_{\pm}, [Q_3,W_{\pm}] = \pm \frac{1}{2}W_{\pm}, \\ [Q_{\pm},V_{\mp}] &= V_{\pm}, [Q_{\pm},W_{\mp}] = W_{\pm}, [Q_{\pm},V_{\pm}] = [Q_{\pm},W_{\pm}] = 0, \\ \{V_{\pm},V_{\pm}\} &= \{W_{\pm},W_{\pm}\} = \{W_{\pm},W_{\mp}\} = 0, \{V_{\pm},W_{\pm}\} = \pm Q_{\pm}, \{V_{\pm},W_{\mp}\} = -Q_3 \pm B. \end{split}$$

The second Casimir operator takes the form

$$K_2 = \vec{Q}^2 - B^2 + \frac{1}{2}(V_+W_- - V_-W_+ + W_+V_- - W_-V_+).$$

Let $|b, q, q_3\rangle$ stand for a vector of any su(2|1) abstract representation, where b, q and q_3 denote the eigenvalues of the operators B and Q_3 , respectively, whereas q is the quantum number that labels the eigenvalue of the \vec{Q}^2 which is actually q(q+1). A typical SU(2|1) coherent state reads

$$|z,\xi,\theta\rangle = \mathcal{N} \exp(-\theta W_{-} - \xi V_{-} + zQ_{-})|b,q,q\rangle,$$

where $(z,\xi,\theta) \in SU(2|1)/U(1) \times U(1)$. We are however interested in the so-called atypical (degenerate) b = q representation that happens to be relevant for the t - J model. This is specified by

$$W_{-}|q,q,q\rangle = 0$$

and is called the (q,q) representation with dimension 4q + 1. This representation is spanned by 2q + 1 vectors $\{|q,q,q_3\rangle, -q \leq q_3 \leq q\}$ of the even (bosonic) and by 2q vectors $\{|q+1/2,q-1/2,q_3\rangle, -q+1/2 \leq q_3 \leq q-1/2\}$ that correspond to the odd (fermionic) sectors. Both the second and third order Casimir operators are zero in this representation. The coherent state reduces in the (q,q) representation to

$$|z,\xi\rangle = (1+|z|^2 + \bar{\xi}\xi)^{-q} e^{-\xi V_- + zQ_-} |q,q,q\rangle,$$
(2)

where z and ξ are even and odd Grassmann parameters, respectively, to be viewed upon as local coordinates on the supermanifold $CP^{1|1} = SU(2|1)/U(1|1)$. Classical images of the su(2|1) generators are found to be $(A^c \equiv \langle z, \xi | A | z, \xi \rangle)$:

$$Q_3^c = -q(1-|z|^2)w, \quad (Q^+)^c = 2qzw, \quad (Q^-)^c = 2q\bar{z}w, B^c = q(1+|z|^2+2\bar{\xi}\xi)w, \quad (V^+)^c = -2qz\bar{\xi}w, \quad (V^-)^c = 2q\bar{\xi}w, (W^+)^c = -2q\xi w, \quad (W^-)^c = -2q\bar{z}\xi w, \quad w = (1+|z|^2+\bar{\xi}\xi)^{-1}.$$
(3)

There is also the second atypical 4q + 1-dimensional representation (-q, q) which happens to be conjugate to the (q, q), i.e., $(q, q)^* = (-q, q)$. These irreps are inequivalent.

The su(2|1) generators in the lowest atypical representation (q = 1/2, q = 1/2) can be identified with the Hubbard operators [11], $X_i^{\sigma 0}$, defined by

$$X_i^{\sigma 0} = c_{i\sigma}^{\dagger} (1 - n_{i,-\sigma}), \quad n_{i\sigma} n_{i,-\sigma} = 0,$$

where $c_{i\sigma}$ is the annihilation operator of an electron at site *i* with spin $\sigma = \pm$, and $n_{i\sigma} \equiv c_{i\sigma}^{\dagger} c_{i\sigma}$. In terms of these operators, the t - J Hamiltonian becomes

$$H_{t-J} = -t \sum_{ij\sigma} X_i^{\sigma 0} X_j^{0\sigma} + J \sum_{ij} \left(\vec{Q}_i \vec{Q}_j - n_i n_j / 4 \right)$$
(4)

with J > 0 and the sum restricted to nearest neighbor sites. Here the local electron spin operator

$$\vec{Q}_i = \frac{1}{2} \sum_{\sigma\sigma'} X_i^{\sigma 0} \vec{\tau}_{\sigma\sigma'} X_i^{0\sigma'},$$

with $\vec{\tau} = (\tau^1, \tau^2, \tau^3)$ being the Pauli matrices. Electron on-site number operator $n_i = X_i^{++} + X_i^{--}$.

If $|0\rangle$ stands for a doped state (a hole) and $|\sigma\rangle$ denotes a state occupied by an electron with the spin projection σ , Hubbard operators take the form

$$X^{\sigma 0} = |\sigma\rangle \langle 0|, \quad X^{\sigma \sigma'} = |\sigma\rangle \langle \sigma'|, \tag{5}$$

 $X^{\sigma 0}$ being a fermionic operator and $X^{\sigma \sigma'}$ corresponding to the bosonic degrees of freedom.

It is clear that there are eight linearly independent operators since by definition

$$X^{00} + \sum_{\sigma} X^{\sigma\sigma} = I.$$

The X-operators are closed into the su(2|1) superalgebra in the 3d degenerate (1/2, 1/2) representation in the following way, [11]

$$\begin{aligned} Q_3 &= \frac{1}{2}(X^{++} - X^{--}), \quad Q_+ = X^{+-}, \quad Q_- = X^{-+}, \quad B = \frac{1}{2}(X^{++} + X^{--}) + X^{00} \\ V_+ &= X^{0-}, \quad V_- = -X^{0+}, \quad W_+ = X^{+0}, \quad W_- = X^{-0}. \end{aligned}$$

The even (bosonic) states $|1/2, 1/2, 1/2\rangle$ and $|1/2, 1/2, -1/2\rangle$ are identified with the spin up and spin down states, $|+\rangle$ and $|-\rangle$, respectively, whereas the odd (fermionc) state $|1,0,0\rangle$, with the doped state $|0\rangle$. In contrast with the (1/2, 1/2) irrep, the conjugate one, (-1/2, 1/2), does not provide a representation in terms of the Hubbard operators.

B. SUSY t - J model

It turns out that at J = 2t this model exhibits in any dimension a global SU(2|1) supersymmetry. This means that $H_{SUSY} = \sum_{ij} g_{\mu\nu} T_i^{\mu} T_j^{\nu}$, $T^{\mu} \in su(2|1)$, where $\nu, \mu = 1, 2, ..., 8$, and summation over the group indices is implied. It is crucial that $g_{\mu\nu}$ appears as the SU(2|1) invariant (under the adjoint action) tensor, with $g_{\mu\nu} = str(T_{\mu}T_{\nu})$. Therefore any continuum version of this interaction should also maintain that symmetry.

It can be shown that

$$H_{SUSY} = 2t \sum_{\langle ij \rangle} \left[\vec{Q}_i \vec{Q}_j - B_i B_j + \frac{1}{2} (V_i^+ W_j^- - V_i^- W_j^+ + W_i^+ V_j^- - W_i^- V_j^+) \right]$$
(6)

$$=t\sum_{\langle ij\rangle} \left[X_i^{0-} X_j^{-0} + X_i^{0+} X_j^{+0} - X_i^{+0} X_j^{0+} - X_i^{-0} X_j^{0-} + X_i^{+-} X_j^{-+} + X_i^{-+} X_j^{+-} - X_i^{++} X_j^{--} - X_i^{--} X_j^{++} \right].$$
(7)

Comparing eq. (6) with the representation of the second Casimir operator K_2 immediately proves that H_{SUSY} displays global SU(2|1) invariance.

This symmetry can be examined in terms of a more standard set of the SU(2|1) group generators. Let us define

$$E^{\alpha\beta} = X^{\alpha\beta}, \ \alpha \neq \beta, \quad E^{\alpha\alpha} = (-1)^{|\alpha|} X^{\alpha\alpha} - 1/2, \quad \alpha, \beta = \pm, 0,$$

the Z_2 grading being defined so that $|\alpha| = 0$ for $\alpha = \pm$ and $|\alpha| = 1$ if $\alpha = 0$. With these definitions

$$H_{SUSY} = t \sum_{\langle ij \rangle} E_i^{\alpha\beta} E_j^{\beta\alpha} (-1)^{|\beta|}, \tag{8}$$

which again coincides with the quadratic Laplace-Casimir element.

The coherent state in the (1/2, 1/2) irrep takes the form

$$|z,\xi\rangle = (1+|z|^2 + \bar{\xi}\xi)^{-1/2}(|+\rangle + \xi|0\rangle + z|-\rangle)$$
(9)

and the operator $\hat{Q}(\bar{z}, \bar{\xi}; z, \xi) = |z, \xi\rangle \langle z, \xi| - 1$, in the 3*d* basis, $|\pm\rangle$, $|0\rangle$, displays the following matrix representation:

$$Q(\bar{z},\bar{\xi};z,\xi) = (1+|z|^2 + \bar{\xi}\xi)^{-1} \begin{pmatrix} -(|z|^2 + \bar{\xi}\xi) & \bar{z} & -\bar{\xi} \\ z & -(1+\bar{\xi}\xi) & -z\bar{\xi} \\ -\xi & -\xi\bar{z} & -(1+2\bar{\xi}\xi+|z|^2) \end{pmatrix}$$
$$= \begin{pmatrix} Q_z - B & Q^- & -V^- \\ Q^+ & -Q_z - B & V^+ \\ W^+ & W^- & -2B \end{pmatrix}^c$$

Note that str Q = 0. As a result,

$$\langle H_{SUSY} \rangle := H_{SUSY}^c = t \sum_{\langle ij \rangle} Q_i^{\beta\alpha} Q_j^{\alpha\beta} (-1)^{|\beta|} = t \sum_{\langle ij \rangle} str(Q_i Q_j).$$
(10)

Explicitly, we have that

$$H_{SUSY}^{c} = t \sum_{\langle ij \rangle} w_{i}^{-1} w_{j}^{-1} \left[\bar{z}_{i} z_{j} + \bar{z}_{j} z_{i} - |z_{i}|^{2} - |z_{j}|^{2} - \xi_{i} \bar{\xi}_{j} (1 + \bar{z}_{i} z_{j}) - \xi_{j} \bar{\xi}_{i} (1 + \bar{z}_{j} z_{i}) \right] = t \sum_{\langle ij \rangle} |\langle z_{i} \xi_{i} | z_{j} \xi_{j} \rangle|^{2}.$$
(11)

where $w_i := (1 + |z_i|^2 + \bar{\xi}_i \xi_i)$. In fact $|\langle z_i \xi_i | z_j \xi_j \rangle|^2 = str P_i P_j$, where the projection operator $P_i = |z_i \xi_i \rangle \langle z_i \xi_j|$. It can be checked that str P = 1and

$$\sum_{\langle ij\rangle} str(Q_i Q_j) = \sum_{\langle ij\rangle} str(P_i P_j) + const.$$
(12)

The $str(P_iP_i)$ is directly related to the Kaehler superpotential of the target superspace,

$$str(P_iP_j) = \exp\Gamma_{ij}, \quad \Gamma_{ij} = F_{ij} + F_{ji} - F_{ii} - F_{jj},$$

where the $CP^{1|1}$ Kaehler superpotential $F_{ij} = \log(1 + \bar{z}_i z_j + \bar{\xi}_i \xi_j)$. In the continuum limit $F_{ij} + F_{ji} - F_{ii} - F_{jj} = \delta^2 F_{ii} + o(a^2)$, a being the lattice spacing, and with the second derivatives of F emerging. Naturally from this definition those derivatives manifest themselves as the entries of the $CP^{1|1}$ metric in terms of which the nonlinear $CP^{1|1}$ sigma model is going to be derived.

III. CP^{1|1} NONLINEAR SIGMA MODEL: PRELIMINARY REMARKS

Depending on a possible choice of the classical ground state, one can arrive at different variants of relevant field theories to incorporate small quantum corrections. To address a quasiclassical (q >> 1) behavior of the SUSY t - Jmodel (6) one may adopt two possible ansatzes. First, since there are two inequivalent atypical representations (q, q)and (-q,q), one may arrange the operators in (6) in an alternating manner: the irrep (q,q) of the su(2|1) superalgebra is placed on sites $i \in A$, whereas its conjugate, $(q, q)^* = (-q, q)$, are defined on sites $i \in B$, with $L = A \oplus B$. A tensor

product of the two is decomposed into a direct sum of the SU(2|1) irreps that contains a singlet. We briefly comment on this option at the end of the present Section.

Another route could be to assume that, classically (i.e., in the $q \to \infty$ limit), the system minimizes its energy by making the nearest neighbour site superspins acquire an "antiparallel" orientation. This option seems to formally provide a direct extension of the Haldane ansatz for the SU(2) spins to the SU(2|1) generators.

Since in both cases our destination is the $CP^{1|1}$ sigma model, let us keep in mind its abstract definition. A canonical nonlinear $CP^{1|1}$ -valued σ -model can formally be described by a virtue of the map $\varphi: \Sigma \to CP^{1|1} = SU(2|1)/U(1|1)$, where Σ stands for a space-time manifold with local coordinates $x^{\mu}, \mu = 0, 1, ..., d(x^0 := t)$, while the $CP^{1|1}$ denotes a N = 1 superextension of a complex projecive line CP^1 to the superspace with local coordinates $(z, \xi) := \Phi^{\alpha}, \alpha = 1, 2$. The Kähler potential $F = 2q \log(1 + \bar{z}z + \bar{\xi}\xi)$, with 2q being integer, defines on $CP^{1|1}$ the Berry connection a and the Berry curvature da, where the external derivative $d = \delta + \bar{\delta}, \ \delta = dz\partial_z + d\xi\partial_{\xi}, \ \text{and } \bar{\delta} = d\bar{z}\partial_{\bar{z}} + d\bar{\xi}\partial_{\bar{\xi}}$. (All derivatives are understood to be the left ones). Explicitly, $a = \frac{i}{2}(\delta - \bar{\delta})F = \frac{i}{2}(dzF_z + d\xi F_{\xi} - d\bar{z}F_{\bar{z}} - d\bar{\xi}F_{\bar{\xi}}) = \frac{iq}{1+|z|^2+\bar{\xi}\xi}(\bar{z}dz - zd\bar{z} - d\bar{\xi}\xi + \bar{\xi}d\xi) =: a_zdz + a_{\bar{z}}d\bar{z} + d\xi a_{\xi} + a_{\bar{\xi}}d\bar{\xi}$. The irrep (1/2, 1/2) is specified by setting q = 1/2. The pull-back of the one-form a is given by $\varphi^* a$, $(\varphi^* a)_{\mu} = a_z \partial_{\mu} z + ...$

Under the SU(2|1) action, $F \to F + \Lambda(z,\xi) + \overline{\Lambda(z,\xi)}$, which means that the symplectic one-form *a* undergoes the U(1) gauge transformation,

$$a \to a + d\psi, \quad \psi = \frac{i}{2}(\Lambda - \bar{\Lambda})$$

The associated invariant canonical supersymplectic two-form is

$$\Omega^{(2)} = da = -i\delta\bar{\delta}F = -i(F_{z\bar{z}}dz \wedge d\bar{z} - F_{z\bar{\xi}}dz \wedge d\bar{\xi} - F_{\xi\bar{z}}d\xi \wedge d\bar{z} + F_{\xi\bar{\xi}}d\xi \wedge d\bar{\xi}), \tag{13}$$

where

$$\begin{split} F_{z\bar{z}} &= \frac{2q}{(1+|z|^2+\bar{\xi}\xi)^2}(1+\bar{\xi}\xi), \quad F_{z\bar{\xi}} = -\bar{z}\xi \, \frac{2q}{(1+|z|^2+\bar{\xi}\xi)^2}, \quad F_{\bar{z}\xi} = \bar{\xi}z \, \frac{2q}{(1+|z|^2+\bar{\xi}\xi)^2}, \\ F_{\xi\bar{\xi}} &= -\frac{2q}{1+|z|^2}, \end{split}$$

belong to the exterior algebra on $CP^{1|1}$. The latter is a bi-graded $Z \times Z_2$ algebra, where the Z-gradation is the usual gradation of de Rham complexes, while Z_2 -gradation is a natural gradation of Grassmann algebra. For any two superforms β_1 and β_2 on $CP^{1|1}$, one has $\beta_1 \wedge \beta_2 = (-)^{a_1a_2+b_1b_2}\beta_2 \wedge \beta_1$, where $a_i(b_i)$ is the degree of β_i with respect to the $Z(Z_2)$ gradation. Hence, $dz \wedge d\bar{z} = -d\bar{z} \wedge dz$, $dz \wedge d\bar{\xi} = -d\bar{\xi} \wedge dz$, $d\xi \wedge d\bar{\xi} = d\bar{\xi} \wedge d\xi$. In terms of the coherent states

$$a = i\langle z, \xi | d | z, \xi \rangle, \quad da = id\langle z, \xi | d | z, \xi \rangle = i \operatorname{str}(P d P \wedge d P).$$

In view of this the supersymplectic two-form can be written in the form

$$\Omega^{(2)} = -ig_{\alpha\bar\beta}d\Phi^\alpha \wedge d\bar\Phi^\beta, \quad \Phi = (\Phi^1, \Phi^2) := (z, \xi),$$

and the SU(2|1) invariant metric supertensor then reads,

$$g(\bar{\Phi},\Phi) = g_{\alpha\bar{\beta}}d\Phi^{\alpha}d\bar{\Phi}^{\beta} = F_{z\bar{z}}dzd\bar{z} - F_{z\bar{\xi}}dzd\bar{\xi} - F_{\xi\bar{z}}d\xi d\bar{z} + F_{\xi\bar{\xi}}d\xi d\bar{\xi},$$
(14)

where differentials of the odd coordinates are understood to anticommute among themselves, and commute with differentials of the even coordinates, with the latter exhibiting the usual behavior.

The action functional for the SU(2|1) invariant σ -model can then be written in the form (D = 1 + d):

$$S_{g} = -\frac{1}{2g^{2}} \int_{\Sigma} d^{D}x \, str(\varphi^{*}g) = -\frac{1}{2g^{2}} \int_{\Sigma} d^{D}x \left[\partial_{\mu} \Phi \, g \partial_{\mu} \bar{\Phi} \right] = -\frac{1}{2g^{2}} \int_{\Sigma} d^{D}x \left[F_{z\bar{z}} \partial_{\mu} z \partial_{\mu} \bar{z} - F_{z\bar{\xi}} \partial_{\mu} z \partial_{\mu} \bar{\xi} - F_{\xi\bar{z}} \partial_{\mu} \bar{z} \partial_{\mu} \xi + F_{\xi\bar{\xi}} \partial_{\mu} \xi \partial_{\mu} \bar{\xi} \right] = -\frac{1}{2g^{2}} \int_{\Sigma} d^{D}x \left[\frac{1}{(1+|z|^{2}+\bar{\xi}\xi)^{2}} \left((1+\bar{\xi}\xi) \partial_{\mu} z \partial_{\mu} \bar{z} + \bar{z}\xi \partial_{\mu} z \partial_{\mu} \bar{\xi} - z\bar{\xi} \partial_{\mu} \bar{z} \partial_{\mu} \xi \right) - \frac{1}{1+|z|^{2}} \partial_{\mu} \xi \partial_{\mu} \bar{\xi} \right],$$
(15)

where the space-time Σ is assumed to be a flat euclidean manifold.

It is also well known that a topological (metric independent) SU(2|1) invariant Wess–Zumino term can be added to S. In general, it can be defined as follows. Consider a map $\varphi : \Sigma \to M$, of a D space-time surface Σ into an ordinary manifold (not a supermanifold) M. Let us assume that a closed D+1-form, $\Omega^{(D+1)}$, with $d\Omega^{(D+1)} = 0$, can be picked up on M. In case it defines a trivial cohomology class, one has globally $\Omega^{(D+1)} = d\Omega^{(D)}$ for a certain D-form on M. In that case one simply defines

$$S_{WZ} = \int_{\Sigma} \varphi^* \Omega^{(D)}$$

If, however, $\Omega^{(D+1)}$ is not globally exact, one may either consider a set of coordinate patches Σ_a , so that $\Sigma = \bigcup_a \Sigma_a$, and a field $\Omega_a^{(D)}$ defined in each patch such that $\Omega^{(D+1)} = d\Omega_a^{(D)}$, or one may try to extend φ to a certain $\tilde{\varphi} : \tilde{\Sigma} \to M$, such that $\partial \tilde{\Sigma} = \Sigma$ and, consequently

$$S_{WZ} = \int_{\tilde{\Sigma}} \tilde{\varphi}^* \Omega^{(D+1)}.$$

In both cases the theory is well-defined, provided $\Omega^{(D+1)}$ defines an integral cohomology class, i.e., with the period of the form being an integer number,

$$\frac{1}{2\pi} \int_{V_{D+1}} \Omega^{(D+1)} = N,$$

where V_{D+1} stands for a D+1 oriented closed manifold in M.

The generalization of the above formalism to the supersetting is however not straightforward. First, any superform that involves $d\xi$ or/and $d\bar{\xi}$ defines a trivial cohomology. This observation seems to simplify the consideration. What makes it more involved is the fact that superforms cannot in general be integrated. This might be a sort of disaster, if it were not for the fact that there is a natural mapping $\varphi : \Sigma \to CP^{1|1}$, As a matter of fact, relevant integrals can be defined via the pull-back φ^* :

$$\int_{\varphi(\Sigma)} d\xi \wedge d\bar{\xi} := \int_{\Sigma} \varphi^* (d\xi \wedge d\bar{\xi}) = \int_{\Sigma} \partial_\mu \xi \partial_\nu \bar{\xi} \, dx^\mu \wedge dx^\nu. \tag{16}$$

For example, the 2d t - J continuum model can be viewed upon as a mapping $\varphi : (x, t) = S^2 \to CP^{1|1}$, with the space-time (x, t) being compactified into S^2 . We are therefore looking for a closed SU(2|1)-invariant 3-forms on $CP^{1|1}$ so that $\Omega^{(3)} = d\Omega^{(2)}$ globally. One easily finds that there exists four trivially closed forms on $CP^{1|1}$ generated by the functions $F_{z\bar{z}\xi} + F_{z\bar{z}\bar{\xi}}$, $F_{\xi\bar{\xi}z} + F_{\xi\bar{\xi}\bar{z}} + F_{z\bar{z}\bar{\xi}}$ and $F_{z\bar{\xi}\xi} + F_{z\bar{z}\bar{\xi}}$, respectively. These in turn yield

$$\Omega_1^{(2)} = -iF_{z\bar{z}}dz \wedge d\bar{z}, \quad \Omega_2^{(2)} = -iF_{\xi\bar{\xi}}d\xi \wedge d\bar{\xi}, \quad \Omega_3^{(2)} = -iF_{z\bar{\xi}}dz \wedge d\bar{\xi}, \quad \Omega_4^{(2)} = -iF_{\bar{z}\xi}d\bar{z} \wedge d\xi$$

The only SU(2|1)-invariant combination of all these choices is the symplectic superform $\Omega^{(2)}$.

As a result, the Wess-Zumino action takes the form

$$S_{WZ} =: S_{\theta} = (i\theta/4\pi q) \int_{CP^{1|1}} \Omega^{(2)},$$
 (17)

with the coefficient θ of the topological term being determined at microscopical level. Explicitly, this reads

$$S_{\theta} = (\theta/2\pi) \int_{\Sigma} \frac{dx^{\mu} \wedge dx^{\nu}}{(1+|z|^{2}+\bar{\xi}\xi)^{2}} \left[(1+\bar{\xi}\xi)z_{\mu}\bar{z}_{\nu} + z\bar{\xi}\bar{z}_{\mu}\xi_{\nu} - \bar{z}\xi z_{\mu}\bar{\xi}_{\nu} \right] \\ + (\theta/2\pi) \int_{\Sigma} \frac{dx^{\mu} \wedge dx^{\nu}}{1+|z|^{2}} \xi_{\mu}\bar{\xi}_{\nu}.$$
(18)

If one discards the fermionic degrees of freedom, thereby considering a conventional $2d \sigma$ -model as a mapping $\varphi: CP^1 \to CP^1$, the above Wess-Zumino term reduces to a topological invariant [12]

$$S_{\theta} \to S_{\theta} = (\theta/2\pi) \int_{CP^{1}} \frac{dx^{\mu} \wedge dx^{\nu}}{(1+|z|^{2})^{2}} z_{\mu} \bar{z}_{\nu} = (\theta/2\pi) \int_{CP^{1}} \varphi^{*} \frac{dz \wedge d\bar{z}}{(1+|z|^{2})^{2}}$$
$$= (\theta/2\pi) (deg\phi) \int_{CP^{1}} \frac{dz \wedge d\bar{z}}{(1+|z|^{2})^{2}} = i\theta N,$$

using

$$i \int_{CP^1} \frac{dz \wedge d\bar{z}}{(1+|z|^2)^2} = 2\pi.$$

At $\theta = 2\pi q$ this is the familiar Haldane's result for the topological term of the spin-q 2d continuum AF action [13]. As a result of that, the total SU(2|1) invariant action takes the form

$$S = S_{g} + S_{\theta} = -\frac{1}{2g^{2}} \int_{\Sigma} d^{2}x \, str(\varphi^{*}g) + (i\theta/4\pi q) \int_{CP^{1|1}} \Omega^{(2)} = \\ -\frac{1}{2g^{2}} \int_{\Sigma} d^{2}x \left[\frac{1}{(1+|z|^{2}+\bar{\xi}\xi)^{2}} \left((1+\bar{\xi}\xi)\partial_{\mu}z\partial_{\mu}\bar{z} + \bar{z}\xi\partial_{\mu}z\partial_{\mu}\bar{\xi} - z\bar{\xi}\partial_{\mu}\bar{z}\partial_{\mu}\xi \right) - \frac{1}{1+|z|^{2}}\partial_{\mu}\xi\partial_{\mu}\bar{\xi} \right] \\ + (\theta/2\pi) \int_{\Sigma} \frac{dx^{\mu} \wedge dx^{\nu}}{(1+|z|^{2}+\bar{\xi}\xi)^{2}} \left[(1+\bar{\xi}\xi)z_{\mu}\bar{z}_{\nu} + z\bar{\xi}\bar{z}_{\mu}\xi_{\nu} - \bar{z}\xi z_{\mu}\bar{\xi}_{\nu} \right] + (\theta/2\pi) \int_{\Sigma} \frac{dx^{\mu} \wedge dx^{\nu}}{1+|z|^{2}}\xi_{\mu}\bar{\xi}_{\nu}.$$
(19)

The 2D NLSM given by Eq.(19) has been argued to yield a continuum limit of the alternating su(2|1) quantum superspin chain with alternating fundamental and conjugate fundamental representations [9]. In this reference, the action (19) is written down through the homogeneous (gauge dependent) coordinates, whereas we employ the inhomogeneous variables. They are related to each other by a change of variables to explicitly resolve the gauge redundancy.

The SU(2|1) alternating chain is closely related with the properties of the percolation cluster boundaries. It is gapless, and flows to the fixed point that is described by a logarithmic conformal field theory at central charge c = 0. This theory corresponds to the strong coupling regime of a $CP^{1|1}$ sigma model [9]. However, the model (19) cannot describe strongly correlated electrons. The point is that strong electron correlations are encoded into the properties of the fundamental (1/2, 1/2) irrep of su(2|1). Generators of this representation can be identified with the Hubbard operators (4). In contrast, the conjugate irrep (-1/2, 1/2) of the su(2|1) superalgebra does not admit a description in terms of the Hubbard operators. In particular, the slave-fermion representation of the (-1/2, 1/2) generators necessarily involves "fermions" that anticommute to -1: $\{f_i, f_i^+\} = -1, i \in B$. Physically, those fermions cannot represent doped holes. Consequently, the (-1/2, 1/2) representation cannot accommodate physical electrons at any finite doping.

IV. CP^{1|1} NONLINEAR SIGMA MODEL: EXPLICIT DERIVATION

Let us now turn to our second option. In dealing with a quasiclassical limit of the SUSY t - J model (6) we now start with a generalized AF ansatz. It implies that on both even and odd sublattices the superspins predominantly point in opposite "directions". In the quasiclassical limit, $q \to \infty$, this is supposed to yield an exact classical ground state, whereas in its vicinity one can derive a low-energy effective action to incorporates small quantum/thermal fluctuations. Such a theory is supposed to be fully controlled by a 1/q expansion.

As far as we know, a relevant theory has never been derived explicitly in this case. Presumably it is due to the fact that the standard Haldane's ansatz $\vec{S}_i^c \to -\vec{S}_i^c$, $i \in B$ (see Appendix A) cannot be generalized straightforwardly to the superspins. There is no such change of variables in a relevant SU(2|1) coherent-state path-integral action that can change the sign of all the superspins simultaneously on one of the sublattices.

To proceed with that, we employ instead the projectors on the coherent-state manifold. This enables us to rewrite the lattice action entirely in terms of the projectors: $H_{SUSY}^{cl} = str \sum_{ij} P_i P_j$ and $a_i = i str(dP_i)$. This simplifies our calculations greatly providing in addition a universal approach to quasiclassically treat the model Hamiltonians which display global invariance under a Lie (super)group action. The overlap of the projectors is directly related to the important geometric structure – the Kaehler (super)potential of the target (super)manifold. In the continuum limit, this overlap reduces to the metric function in terms of which emerges the NLSM in question.

In case there is projector P_i on the sublattice A, on sublattice B one has to change $P_i \to 1 - P_i$, $i \in B$. Indeed, $Q^c = strQP \to strQ(1-P) = -Q^c$. Assuming the semi-classical Neel superspin configuration, we can then make the ansatz

$$H_{SUSY}^{c} = \sum_{\langle ij \rangle} str(P_i P_j) \rightarrow \sum_{\langle ij \rangle} str(1 - P_i P_j) = \sum_{\langle ij \rangle} (1 - \exp \Gamma_{ij}).$$
(20)

As we show in the Appendix A, this procedure at half-filling is equivalent to the Haldane ansatz for the AF Heisenberg model. It then follows that the continuum limit to describe the theory around a classical Neel state is determined by

9

only one ingredient, the SU(2|1) covariant Kaehler superpotential. Away from the SUSY point, this approach does not hold, however.

To explicitly derive the relevant sigma model let us start by writing down a lattice Euclidean action that enters the path-integral representation of the t - J partition function,

$$S = i \sum_{i} \oint a_{i} - \int H^{cl}_{SUSY}(t) dt.$$

The first piece in this action is a topological term: it emerges as a sum of the line integrals of the site dependent su(2|1) symplectic one-forms $a_i = i\langle z_i, \xi_i | d | z_i, \xi_i \rangle$ and involves no metric. It is commonly referred to as the Berry phase term or, the Wess-Zumino term or, the 1D Chern-Simons term. The second part is a metric dependent term.

Let us first turn to the evaluation of a continuum form of that second part of the action. For a general magnitude of the representation index q we have $H_{SUSY}^{cl} = 2q H_{q=1/2}^{cl}$, with parameter t being replaced by $t \to t/2q$. Low-energy continuum limit then implies that $2q \to \infty$. In view of eqs. (10,20) we get in 1D

$$H_{q=1/2}^{cl} =: H^{cl} = -t \sum_{i} \Gamma_{ii+1} + o(a^2),$$
(21)

where *a* stands for the lattice spacing. Here $\Gamma_{ii+1} := F(\bar{z}_i, \bar{\xi}_i | z_{i+1}, \xi_{i+1}) + F(\bar{z}_{i+1}, \bar{\xi}_{i+1} | z_i, \xi_i) - F(\bar{z}_i, \bar{\xi}_i | z_i, \xi_i) - F(\bar{z}_i, \bar{\xi}_i, \xi_i) - F(\bar{z}_i, \bar{\xi}_i, \xi_i) - F(\bar{z}_i, \bar{\xi}_i, \xi_i) - F(\bar$

To proceed, we impose the ansatz

$$z_i \to z_i + \delta z_i, \quad \xi_i \to \xi_i + \delta \xi_i, \quad i \in A; \qquad z_i \to z_i - \delta z_i, \quad \xi_i \to \xi_i - \delta \xi_i, \quad i \in B,$$
 (22)

where $\delta z_i \sim a$, $\delta \xi_i \sim a$ are small AF field fluctuations. In the continuum limit, $a \to 0$, eq. (21) takes the form

$$H^{cl} = t \sum_{i} a^{2} (\Phi'_{i} g(\bar{\Phi}_{i}, \Phi_{i}) \bar{\Phi}'_{i}) + 4t \sum_{i} (\delta \Phi_{i} g(\bar{\Phi}_{i}, \Phi_{i}) \delta \bar{\Phi}_{i}) + o(a^{2}),$$
(23)

where the superfields $\Phi = (z, \xi)$, $\bar{\Phi}^t = (\bar{z}, \bar{\xi})$, and g is the SU(2|1) invariant metric supertensor explicitly given by (14).

Let us now turn to the topological term. As already mentioned, on sublattice B we let $P \rightarrow 1 - P$. Because of this the one-form a_i changes its sign on sublattice B. Hence we get

$$i\sum_{i} \oint (a_i + \delta a_i) = i\sum_{i} \oint a_i = i\sum_{i \in A} \oint a_i - i\sum_{i \in B} \oint a_i + i\sum_{i \in A} \oint \delta a_i - i\sum_{i \in B} \oint \delta a_i,$$
(24)

where variation δa_i is produced by change (22). The second piece of eq. (24) that arises from the changes of the Berry phase in each of the sublattices yield

$$i\sum_{i\in A}\oint \delta a_i - i\sum_{i\in B}\oint \delta a_i = \sum_i \int dt \left[\delta \Phi_i g(\bar{\Phi}_i, \Phi_i)\dot{\bar{\Phi}}_i - \dot{\Phi}_i g(\bar{\Phi}_i, \Phi_i)\delta\bar{\Phi}_i\right].$$

This term along with eq. (23) give the following contributions to the action

$$\sum_{i} \int dt \left[\left(\delta \Phi_{i} g(\bar{\Phi}_{i}, \Phi_{i}) \dot{\bar{\Phi}}_{i} \right) - \left(\dot{\Phi}_{i} g(\bar{\Phi}_{i}, \Phi_{i}) \delta \bar{\Phi}_{i} \right) + ta^{2} \left(\Phi_{i}^{\prime} g(\bar{\Phi}_{i}, \Phi_{i}) \bar{\Phi}_{i}^{\prime} \right) + 4t \left(\delta \Phi_{i} g(\bar{\Phi}_{i}, \Phi_{i}) \delta \bar{\Phi}_{i} \right) \right] + o(a^{2}).$$

Extremizing the action with respect to $\delta \Phi$ and $\delta \bar{\Phi}$ produces the metric dependent contribution

$$-q\sum_{i}\int dt\left(\frac{\dot{\Phi}_{i}g_{i}\dot{\bar{\Phi}}_{i}}{4qt}+a^{2}4qt\Phi_{i}'g_{i}\bar{\Phi}_{i}'\right).$$

Note that we have restored the full q dependence. In the continuum limit this contributes to the total action in the form

$$S_g = -\frac{1}{2g^2} \int dx dt \left(\frac{\dot{\Phi}g(\bar{\Phi}, \Phi)\dot{\bar{\Phi}}}{c} + c\Phi'g(\bar{\Phi}, \Phi)\bar{\Phi}' \right), \qquad g^2 = 1/q, \quad c = 4qat.$$
(25)

At c = 1 eq. (25) goes over into the Lorentz invariant action targeted on $CP^{1|1}$:

$$S_{g} = -\frac{1}{2g^{2}} \int dx dt \left(\partial_{\mu} \Phi g(\bar{\Phi}, \Phi) \partial_{\mu} \bar{\Phi} \right) -\frac{1}{2g^{2}} \int dx dt \left[F_{z\bar{z}} \partial_{\mu} z \partial_{\mu} \bar{z} - F_{z\bar{\xi}} \partial_{\mu} z \partial_{\mu} \bar{\xi} - F_{\xi\bar{z}} \partial_{\mu} \bar{z} \partial_{\mu} \xi + F_{\xi\bar{\xi}} \partial_{\mu} \xi \partial_{\mu} \bar{\xi} \right],$$
(26)

which exactly coincides with Eq. (15) if we take $g^2 = 1/q$.

Let us now focus on the evaluation of the topological invariant S_{θ} that arises from the first piece of the variation of the Berry phase term,

$$\oint a_i - \oint a_{i+1} =: \delta \oint a_i$$

Here a small change in the integral is due to the following transformations of the coordinates:

$$\delta z_i = z_{i+1} - z_i = z(x_i + a) - z(x_i) = az'(x_i) + O(a^2); \quad \delta \xi_i = a\xi'(x_i) + O(a^2). \tag{27}$$

This transformation is generated by the infinitesimal vector field

$$\delta X = \delta z \partial_z + \delta \bar{z} \partial_{\bar{z}} + \delta \xi \partial_{\xi} + \delta \bar{\xi} \partial_{\bar{\xi}}.$$

As is well known a variation of a line integral due to a local one-parameter group of diffeomorphisms generated by a vector field δX gives

$$\delta \oint a_i = \oint \mathcal{L}_{\delta X_i} a_i, \tag{28}$$

where \mathcal{L}_X stands for a Lie derivative with respect to the vector field X. By making use of the identity

$$\mathcal{L}_X = i_X \, d + d \, i_X$$

one can then transform (28) into

$$\delta \oint a_i = \oint i_{\delta X_i} \, da_i + \oint d \, (i_{\delta X_i} a_i), \tag{29}$$

where i_X denotes the interior product. The last term in (29) drops out thanks to Stokes' theorem and we are left only with the first term. To evaluate it we need to know how the interior derivative acts on da given by eq. (13). Explicitly we get

$$i_{\delta X_i}(F_{\bar{z}_i z_i} dz_i \wedge d\bar{z}_i) = F_{\bar{z}_i z_i}(\delta z_i d\bar{z}_i - dz_i \delta \bar{z}_i), \quad i_{\delta X_i}(F_{z_i \bar{\xi}_i} dz_i \wedge d\bar{\xi}_i) = F_{z_i \bar{\xi}_i}(\delta z_i d\bar{\xi}_i - dz_i \delta \bar{\xi}_i),$$

$$i_{\delta X_i}(F_{\xi_i\bar{z}_i}d\xi_i \wedge d\bar{z}_i) = F_{\xi_i\bar{z}_i}(\delta\xi_i d\bar{z}_i - d\xi_i \delta\bar{z}_i), \quad i_{\delta X_i}(F_{\xi_i\bar{\xi}_i}d\xi_i \wedge d\bar{\xi}_i) = F_{\xi_i\bar{\xi}_i}(\delta\xi_i d\bar{\xi}_i - d\xi_i \delta\bar{\xi}_i).$$

Here $\delta z_i = z'_i a$, $dz_i = \dot{z}_i dt$ and similarly for the variations of ξ_i . The first equation tells us that in the continuum limit

$$\sum_{i} F_{\bar{z}_i z_i} (\delta z_i d\bar{z}_i - dz_i \delta \bar{z}_i) = \int F_{\bar{z}z} (z' \dot{\bar{z}} - \dot{z} \bar{z}') dx \wedge dt = \int F_{\bar{z}z} dz \wedge d\bar{z},$$

where z = z(x,t) and $dz = z'dx + \dot{z}dt$. Similar relations hold for the remaining parts of the expressions and we finally end up with the following contribution to the action coming from the first piece of the topological term

$$S_{\theta} = i \sum_{i \in A} \oint a_i - i \sum_{i \in B} \oint a_i = \frac{i}{2} \int_{CP^{1|1}} \Omega^{(2)}.$$
 (30)

Comparing this with eq. (17), gives $\theta = 2\pi q$. Note that factor 1/2 arises because of the doubling of the lattice spacing.

In summary, we have derived the 2D $CP^{1|1}$ NLSM to describe the low-energy physics of the 1d quantum su(2|1) superspin chain (7) in the form

$$S = S_g + S_\theta$$

$$= -\frac{1}{2g^2} \int dx dt [F_{z\bar{z}}\partial_{\mu}z\partial_{\mu}\bar{z} - F_{z\bar{\xi}}\partial_{\mu}z\partial_{\mu}\bar{\xi} - F_{\xi\bar{z}}\partial_{\mu}\bar{z}\partial_{\mu}\xi + F_{\xi\bar{\xi}}\partial_{\mu}\xi\partial_{\mu}\bar{\xi}] + \frac{i\theta}{4\pi q} \int_{CP^{1|1}} \Omega^{(2)}, \quad g^2 = 1/q, \quad \theta = 2\pi q, \quad (31)$$

which agrees with eq. (19), the phenomenological parameters being now explicitly specified. In particular, the θ term ensures that the model (31) contains low-energy gapless excitations in accordance with the exact solution for
the alternating superspin chain obtained by Bethe ansatz [14].

However, once again, the ansatz (20) cannot describe the Hubbard operators on both sublattices simultaneously as given by Eq.(7). On the sublattice B, we get $B_i^c \to -B_i^c$, which is inconsistent with the definition of the su(2|1)generator B in terms of Hubbard operators: $B = \frac{1}{2}(1 + X^{00})$. The eigenvalues of the operator X^{00} are either 0 and 1 and they describe either an absence of the doping or a doped hole, respectively. This change of sign of B implies that the doping becomes "negative" and this is physically inappropriate. At half-filling, however, the operator Bbecomes a c-number that takes on the values of +q and -q on the two different sublattices. This does not affect the effective action that reduces in this case to the CP^1 NLSM. However, in the presence of a nonzero doping the proper quasiclassical ground state has a more complicated structure than that displayed by a pure Neel state. As a result, the Neel state is not a good reference state, in those cases.

As a matter of fact, an exact diagonalization of a two-site cluster indicates that the AF ground state of the SUSY t-J model (7) is a quartet, rather than a Neel state or a singlet [15]. This result can be visualized in terms of the following su(2|1) diagram in the Clebsch-Gordon series $3 \otimes 3 = 5 \oplus 4$. At half filling, this quartet reduces to a SU(2) spin singlet. It is quite a nontrivial problem to figure out whether a proper large-q generalization of such a state may serve as a reference state for the CP^1 Haldane action, which is a necessary condition for the quartet state to represent a proper reference state away from half filling.

Another alternative possibility might be to consider the low energy effective action of the t - J model as being represented by the spinless fermions interacting with the CP^1 purely spin sigma model via the emergent gauge field. The gauge field appears as a U(1) connection of the corresponding magnetic monopole bundle [8]. At half-filling, such a theory automatically reduces to the CP^1 NLSM as required. However, the emergent SU(2|1) invariant measure intertwines the spin and charge degrees of freedom in a highly nontrivial manner, which is again a manifestation of strong correlations. As a result, no nontrivial computations are available so far within this approach.

However, even in the case one happens to guess correctly an appropriate reference state there is still an issue as to whether strong correlations that are built into the theory at q = 1/2 survive at the large -q limit. Our guess is that it does not persist. Indeed, a state with one hole and a state with two holes acquire spins q - 1/2 and q - 1, respectively. Those states can hardly be distinguished from each other in the large-q limit. The crucial no doubly occupied constraint that encodes the essence of strong correlations seems to be relaxed in this limit.

V. CONCLUSION

We consider the t - J model of strongly correlated electrons at the SUSY point, J = 2t. The constrained electron operators that describe the correlated electrons can be identified with the generators of the fundamental (1/2, 1/2)3d representation of the su(2|1) superalgebra given by the canonical Hubbard operators. However, to investigate the issue as to whether the quasiclassics may capture or not the physics of strong correlations in a reliable way, we make use of the higher (q, q) representation for large magnitudes of the superspin q. Starting from the generalized AF Neel configuration of the superspins on a bipartite lattice, we derive the low-energy effective action – the canonical $CP^{1|1}$ NLSM.

We then show that although this derivation may indeed be relevant for some physical models, the generalized AF ansatz does not admit an interpretation in terms of the Hubbard operators at finite doping. This by far rules out the possibility that the canonical $CP^{1|1}$ NLSM may be considered as a relevant low-energy quasiclassical action to describe the physics of the t - J model at finite doping.

The proposed derivation of the NLSM is based on the representation of a classical image of the SUSY t - J model Hamiltonian in terms of the projectors on the SU(2|1) coherent-state manifold. Such an approach appears as quite a general one: It basically involves only one essential ingredient that determines a local geometry of the coherent-state manifold – a Kaehler (super)potential.

VI. APPENDIX A

Let us show how this technique works in the case of the AF Heisenberg model. The su(2) algebra is spanned by the operators S^+, S^-, S_z with the commutation relations $[S_z, S^{\pm}] =$ $\pm S^{\pm}, [S^+, S^-] = 2S_z$. In the fundamental representati (s = 1/2), we define

$$S^+ = |+\rangle\langle -|, S^- = |-\rangle\langle +|, S_z = \frac{1}{2}(|+\rangle\langle +|-|-\rangle\langle -|),$$

with the $|\pm\rangle$ being the spin-up and spin-down states, respectively.

The su(2) normalized coherent state in this irrep is [16]

$$|z\rangle = (1+|z|^2)^{-1/2}e^{zS^-}|+\rangle = (1+|z|^2)^{-1/2}(|+\rangle+z|-\rangle),$$

where z is a complex number – a local coordinate on $SU(2)/U(1) = CP^1 = S^2$.

The corresponding covariant symbols (the su(2) cs averages) are

$$S_{c}^{+} = trPS^{+} = \frac{z}{1+|z|^{2}}, \quad S_{c}^{-} = trPS^{-} = \frac{\bar{z}}{1+|z|^{2}}, \quad S_{c}^{z} = trPS^{z} = \frac{1}{2}\frac{1-|z|^{2}}{1+|z|^{2}}, \tag{32}$$

where the projection operator $P := |z\rangle \langle z|$.

A classical image of the AF Heisenberg model

$$H = J \sum_{ij} \vec{S}_i \vec{S}_j, \quad J > 0, \tag{33}$$

reads

$$H^{c} = trPH = J \sum_{ij} \vec{S}_{i}^{c} \vec{S}_{j}^{c} = J \sum_{ij} |\langle z_{i} | z_{j} \rangle|^{2} = J \sum_{ij} tr(P_{i}P_{j}).$$
(34)

The Neel ground state implies staggered classical spin moments so that two neighboring-site spins point in the opposite directions: $\vec{S}_i^c, i \in A, -\vec{S}_i^c, i \in B, A \bigoplus B = L$

$$H_{AF}^{c} = -J \sum_{ij} \vec{S}_{i}^{c} \vec{S}_{j}^{c} = -J \sum_{ij} \left(S_{zi}^{c} S_{zj}^{c} + \frac{1}{2} (S_{ci}^{+} S_{cj}^{-} + S_{ci}^{-} S_{cj}^{+}) \right).$$
(35)

Here the vectors \vec{S}_i^c are given by Eqs.(32) in which the z_i do not depend on time (quantum fluctuations are ignored). Equation (35) follows upon the change $z_i \to -1/\bar{z}_i$, $i \in B$. Under such a change of variables $P_i \to 1 - P_i$ so that

$$H_{AF}^{c} = J \sum_{ij} tr P_{i}(1 - P_{j}) = J \sum_{ij} (1 - tr P_{i}P_{j}) = J \sum_{ij} (1 - e^{\Gamma_{ij}}),$$
(36)

where $\Gamma_{ij} = F_{ij} + F_{ji} - F_{ii} - F_{jj}$, and the CP^1 Kaehler potential reads $F_{ij} = 2s \log(1 + \bar{z}_i z_j)$. Small quantum fluctuation around the AF classical ground state, can be incorporated along the lines depicted in the preceding Section to arrive at the CP^1 NLSM given by Eq.(31) where one sets $\xi_i = \bar{\xi}_i \equiv 0$.

Let us now turn to the conjugate representation. All relevant quantities are marked by the "tilde" sign. The irrep conjugated to the fundamental one is generated by the operators

$$\tilde{S}^{+} = -S^{-}, \quad \tilde{S}^{-} = -S^{+}, \quad \tilde{S}_{z} = -S_{z}.$$
 (37)

One therefore gets

$$H_{AF}^{c} \to \tilde{H}_{AF}^{c} = -J \sum_{ij} \left(S_{zi}^{c} S_{zj}^{c} + \frac{1}{2} (S_{ci}^{+} S_{cj}^{+} + S_{ci}^{-} S_{cj}^{-}) \right).$$
(38)

Upon the operation of complex conjugation, $z \to \bar{z}$, this reduces back to Eq.(35).

In the irrep conjugated to the fundamental one we get

$$|\tilde{z}\rangle = (1+|z|^2)^{-1/2}e^{-\bar{z}S^+}|-\rangle = (1+|z|^2)^{-1/2}(|-\rangle - \bar{z}|+\rangle).$$

This yields

$$\tilde{P}(\bar{z},z) = |\tilde{z}\rangle \langle \tilde{z}| = 1 - P(\bar{z},z)$$
(39)

The su(2) case is special in that any su(2) irrep is equivalent to its conjugate representation. Therefore there must be an operator U_g , $g \in SU(2)$ such that $(U_g^{\dagger}PU_g)(\bar{z}, z) = P(\bar{g}z, gz) = \tilde{P}^*(\bar{z}, z)$. One can easily find a required SU(2)action on CP^1 in the form g(z) = -1/z. As a result, $P^*(\overline{g(z)}, g(z)) = \tilde{P}(\bar{z}, z)$, and we get back to Eq.(36). Note that $trP_i = tr(1 - P_i) = 1$ which indicates that all the states on sublattices A and B can be properly normalized. To summarize, we see that in this case the Neel state can be arranged on the neighboring sites in a self-consistent manner.

VII. APPENDIX B

Let us briefly comment on the SU(2|1) lattice action in the conjugate irrep. In the su(2|1) fundamental irrep, conjugate to the (q, q) one, we get

$$\tilde{Q^+} = -Q^-, \ \tilde{Q^-} = -Q^+, \ \tilde{Q}_3 = -Q_3, \ \tilde{B} = -B,$$

 $\tilde{V}_- = -W_+, \ \tilde{V}_+ = W_-, \ \tilde{W}_+ = V_-, \ \tilde{W}_- = -V_+,$

where the "tilde" sign indicates the conjugate irrep. It can be checked that the "tilde" operators fulfil the su(2|1) commutation relations.

In this representation, the su(2|1) coherent state takes the form

$$\langle \tilde{z}, \xi \rangle = \mathcal{N} \langle -1/2, 1/2, -1/2 \rangle e^{-\bar{\xi}W_+ - \bar{z}Q_+},$$
(40)

which yields

$$|\tilde{z,\xi}\rangle = (1+|z|^2+\bar{\xi}\xi)^{-1/2}(|-1/2,1/2,-1/2\rangle-z|-1/2,1/2,1/2\rangle+\xi|-1,0,0\rangle),$$
(41)

The classical images of the su(2|1) generators are then found to be:

$$\begin{split} (\tilde{Q^+})^c &= -(Q^-)^c, \, (\tilde{Q^-})^c = -(Q^+)^c, \, (\tilde{Q_3})^c = -(Q_3)^c, \, (\tilde{B})^c = -B^c, \\ (\tilde{V_-})^c &= -W_+^c, \, (\tilde{V_+})^c = W_-^c, \, (\tilde{W_+})^c = V_-^c, \, (\tilde{W_-})^c = -V_+^c, \end{split}$$

where $\tilde{A^c} = \langle \tilde{z,\xi} | \tilde{A} | \tilde{z,\xi} \rangle$ and the quantities A^c are given by Eqs.(3).

The lattice Euclidean action becomes

$$S = i \sum_{i} \oint a_{i} - \int H^{cl}_{SUSY}(t) dt$$

where we now have

$$H_{SUSY}^{cl} = t \sum_{\langle ij \rangle} str(Q_i \tilde{Q_j}).$$
(42)

It can easily be seen that $\hat{Q}_j \neq -Q_j$. Besides, \tilde{a}_j -the Berry phase in the conjugate representation- coincides with a_j . As a result, our approach developed in Section IV cannot be applied in this case. In view of this observation, it remains unclear whether the SUSY superspin model based on the alternating group representations and the one based on the generalized AF Neel superspin representation lead, in the low-energy limit, to formally identical actions. This is presumably true in the large-q limit, however, a full comparison of the microscopic derivation of the both models is needed to draw a precise conclusion.

- [1] F.D.M. Haldane, Phys. Rev. Lett. 50 (1983) 1153; ibid. 61 (1988) 1029.
- [2] A. Ferraz, and E. Kochetov, Eur. Phys. Lett. 109 (2015) 37003
- [3] Y. Nagaoka, Phys. Rev. **147**, (1966) 392.
- [4] P. Schlottmann, Phys. Rev. B 36 (1987) 5177;
- P.A. Bares, G. Blatter, and M. Ogata, Phys. Rev. B 44 (1991) 130.
- [5] M. Grilli, and G. Kotliar, Phys. Rev. Lett. 64 (1990) 1170.
- [6] P. Coleman, C. Pepin, and J. Hopkinson, Phys. Rev. B 63 (2001) 140411(R).
- [7] C.L. Kane, P.A. Lee, and N. Read, Phys. Rev. B 39 (1989) 6880.
- [8] A. Ferraz, and E. Kochetov, Nucl. Phys. B 853 (2011) 710.
- [9] N. Read and H. Saleur, Nucl. Phys. B 613 (2001) 409.
- [10] M. Scheunert, W. Nahm, and V. Rittenberg, J. Math. Phys. 18 (1977) 155.
- [11] P. Wiegmann, Phys. Rev. Lett. 60 (1988) 821.

- [12] For the 3d t J continuum model, a relevant mapping is $\varphi : (x_1, x_2, t) = S^3 \to CP^{1|1}$. A natural choice is then $\Omega^{(4)} = da \wedge da$ - the second Chern character – its Chern-Simons form serving as a Wess-Zumino term, $S_{WZ} = \int_{CP^{1|1}} a \wedge da$. Being restricted to the Hopf map $S^3 \to CP^1$, which corresponds to the continuum limit for a 2d quantum AF, it vanishes in accordance with the direct large-spin limit calculations.
- [13] It is interesting to note that formally the same θ -term ensures (in the path-integral context) absence of tunneling for half-integer spins, provided a system has time reversal (T) invariance (Kramers' theorem). More specifically, a change of variables $z(t) \to (Tz)(t) = -1/\bar{z}(\tau - t)$ in the quasiclassical path-integral representation of a transition amplitude \mathcal{K} between the T-related coherent states results in the relation $\mathcal{K} = \mathcal{K} e^{i\theta}$ which means that \mathcal{K} vanishes for half-integer s.
- [14] F. Essler, H.Frahm, H. Saleur, Nucl. Phys. B 712 (2005) 513.
- [15] A. Foerster, and M. Karowski, Nucl. Phys. B 396 (1993) 611.
- [16] M. Stone, K-S. Park, and A. Garg, J. Math. Phys. 41 (2000) 8025.