

Solving (most) of a set of quadratic equalities: Composite optimization for robust phase retrieval

John C. Duchi Feng Ruan
Stanford University

December 11, 2021

Abstract

We develop procedures, based on minimization of the composition $f(x) = h(c(x))$ of a convex function h and smooth function c , for solving random collections of quadratic equalities, applying our methodology to phase retrieval problems. We show that the prox-linear algorithm we develop can solve phase retrieval problems—even with adversarially faulty measurements—with high probability as soon as the number of measurements m is a constant factor larger than the dimension n of the signal to be recovered. The algorithm requires essentially no tuning—it consists of solving a sequence of convex problems—and it is implementable without any particular assumptions on the measurements taken. We provide substantial experiments investigating our methods, indicating the practical effectiveness of the procedures and showing that they succeed with high probability as soon as $m/n \geq 2$ when the signal is real-valued.

1 Introduction

We wish to solve the following problem: we have a set of m vectors $a_i \in \mathbb{C}^n$ and nonnegative scalars $b_i \in \mathbb{R}_+$, $i = 1, \dots, m$, and wish to find a vector $x \in \mathbb{C}^n$ such that

$$b_i = |\langle a_i, x \rangle|^2 \text{ for most } i \in \{1, \dots, m\}. \quad (1)$$

As stated, this is a combinatorial problem that is, in the worst case, NP-hard [20]. Yet it naturally arises in a number of real-world situations, including phase retrieval [21, 22, 24], in which one receives measurements of the form

$$b_i = |\langle a_i, x_\star \rangle|^2$$

for known measurement vectors $a_i \in \mathbb{C}^n$, while $x_\star \in \mathbb{C}^n$ is unknown. The problem in phase retrieval arises due to limitations of optical sensors, where one illuminates an object x_\star , which yields diffraction pattern Ax_\star , but sensors may measure only the amplitudes $b = |Ax_\star|^2$, where $|\cdot|^2$ denotes the elementwise squared-magnitude [38]. In the case in which some measurements may be corrupted, the problem is even more challenging.

An alternative objective for the problem (1) is an exact penalty formulation [26], which replaces the equality constraint $b_i = |\langle a_i, x \rangle|^2$ with a non-differentiable cost measuring the error $b_i - |\langle a_i, x \rangle|^2$, yielding the formulation

$$\underset{x}{\text{minimize}} \quad f(x) := \frac{1}{m} \sum_{i=1}^m ||\langle a_i, x \rangle|^2 - b_i| = \frac{1}{m} \| |Ax|^2 - b \|_1. \quad (2)$$

This objective is a natural replacement of the equality constrained problem (1), and the ℓ_1 -loss handles gross errors in the measurements b_i in a relatively benign way (as is well-known in the

statistics and optimization literature on ℓ_1 -based losses and median estimators). Moreover, in the case when $b_i = |\langle a_i, x_\star \rangle|^2$ for all i , it is clear that taking $\iota = \sqrt{-1}$ to be the imaginary unit, then the set $\{e^{\iota\theta} x_\star \mid \theta \in [0, 2\pi)\}$ globally minimizes $f(x)$, though it is only possible to recover x_\star up to its phase (or sign flip in the real case). Candès, Strohmer and Voroninski [9] and Eldar and Mendelson [19], as well as results we discuss later in the paper, show (roughly) that $f(x)$ stably identifies x_\star , in that $f(x)$ grows very quickly as $\text{dist}(x, X_\star) = \inf\{\|x - x_\star\|_2 \mid x_\star \in X_\star\}$, where X_\star denotes the global minimum of f , grows. The objective is, unfortunately, non-smooth, non-convex—not even locally convex near x_\star , as is clear in the special case when $x \in \mathbb{R}$ and $f(x) = |x^2 - 1|$, so that a local analysis based on convexity is impossible—and at least $f(x)$ *a priori* seems difficult to minimize.

Nonetheless, the objective (2) enjoys a number of structural properties that, as we explore below, make solving problem (1) tractable as long as the measurement vectors a_i are sufficiently random. In particular, we can write f as the composition $f(x) = h(c(x))$ of a convex function h and smooth function c , a structure known in the optimization literature to be amenable to efficient algorithms [23, 7, 18]. This compositional structure lends itself nicely to the *prox-linear* algorithm, a variant of the Gauss-Newton procedure, which we describe briefly here for the real case. The composite optimization problem, which Fletcher and Watson [23] originally develop (see also [6, 37]) and a number of researchers [7, 8, 18, 17] have studied further, is to minimize

$$\underset{x}{\text{minimize}} \quad f(x) := h(c(x)) \quad \text{subject to } x \in X \quad (3)$$

where the function $h : \mathbb{R}^m \rightarrow \mathbb{R}$ is convex, $c : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is smooth, and X is a convex set. Extended to the complex case, this general form encompasses our objective (2), where we take $h(z) = \frac{1}{m} \|z\|_1$ and $c(x) = [|\langle a_i, x \rangle|^2 - b_i]_{i=1}^m$. Using the common idea of most optimization schemes—trust region, gradient descent, Newton’s method—to build a simpler to optimize local model of the objective and repeatedly minimize this model, we can replace $h(c(x))$ in problem (3) by linearizing only c . This immediately gives a convex surrogate and leads to the prox-linear algorithm developed by Burke and Ferris [8, 7], among others [18, 16, 17]. Fixing $x \in \mathbb{R}^n$, for any $y \in \mathbb{R}^n$ we define the local “linearization” of f at x by

$$f_x(y) := h(c(x) + \nabla c(x)^T(y - x)), \quad (4)$$

where $\nabla c(x) \in \mathbb{R}^{n \times m}$ denotes the Jacobian transpose of c at x . This function is evidently convex in y , and the prox-linear algorithm proceeds iteratively x_1, x_2, \dots by minimizing the regularized models

$$x_{k+1} = \underset{x \in X}{\text{argmin}} \left\{ f_{x_k}(x) + \frac{1}{2\alpha_k} \|x - x_k\|_2^2 \right\}, \quad (5)$$

where $\alpha_k > 0$ is a stepsize. If h is L -Lipschitz and ∇c is β -Lipschitz, then choosing any stepsize $\alpha \leq \frac{1}{\beta L}$ guarantees that the method (5) is a descent method and finds approximate stationary points of the problem (3) [16, 18]. The case in which $c : \mathbb{C}^n \rightarrow \mathbb{R}^m$ requires a bit more elaboration based on the Wirtinger calculus, which we address later.

We briefly summarize our main contribution as follows. Broadly, this work provides a general method for robust non-convex modeling, focusing carefully on the problem (2); our work carries on a line of work identifying statistical scenarios that nominally yield non-convex optimization problems yet admit computationally efficient estimation and optimization procedures [28, 31, 15, 1]. In this direction, our work develops analytic and statistical tools to analyze a collection of optimization and modeling approaches beyond gradient-based (and Riemannian gradient-based) procedures, using both non-smooth and non-convex models while leveraging statistical structure. More precisely, we show how to apply prox-linear method (5) to any measurement matrix A with no tuning parameters

except that the stepsize satisfies $\alpha \leq (\frac{1}{m} \|A^H A\|_{\text{op}})^{-1}$. Each iteration requires solving a QP in n variables, which is efficiently solvable using standard convex programming approaches. We show that—with extremely high probability under appropriate random measurement models—our prox-linear method exhibits local quadratic convergence to the signal as soon as the number of measurements m/n is greater than some numerical constant, meaning we must solve only $\log_2 \log_2 \frac{1}{\epsilon}$ such convex problems to find an estimate \hat{x} of x_\star such that $\text{dist}(x, X_\star) \leq \epsilon$. In practice, this is 5 convex quadratic programs. Our procedure applies both in the noiseless setting and when a (constant but random) fraction of the measurements are even adversarially corrupted.

1.1 Related work and approaches to phase retrieval

Our work should be viewed in the context of the recent and successful collection of work on phase retrieval. A natural strategy for problem (1), when we wish to find x satisfying $|\langle a_i, x \rangle|^2 = b_i$ for all $i \in [m]$, is to lift the problem into a semidefinite program (SDP) by setting $X = xx^H$, relaxing the rank one constraint, and solving

$$\underset{X}{\text{minimize}} \quad \text{tr}(X) \quad \text{subject to} \quad X \succeq 0, \quad \text{tr}(X a_i a_i^H) = b_i.$$

This is the approach that a number of convex approaches to phase retrieval take [14, 12, 9, 10, 45]. The resulting SDP is computationally challenging for large n , as it requires storing and manipulating an $n \times n$ matrix variable. Moreover, computation times to achieve ϵ -accurate solutions to this problem generally scale on the order of $n^3/\text{poly}(\epsilon)$, where $\text{poly}(\epsilon)$ denotes a polynomial in ϵ .

These difficulties have led a number of researchers to consider non-convex approaches to the phase retrieval problem that—as we do—maintain only a vector $x \in \mathbb{C}^n$, rather than forming a full matrix $X \in \mathbb{C}^{n \times n}$. We necessarily give only a partial overview, focusing on recent work on provably convergent schemes. Early work in computational approaches to phase retrieval is based on (non-convex) alternating projection approaches, notably those by Gerchberg and Saxton [24] and Fineup [22]. Motivated by the challenges of convex approaches and the success of alternating minimization [24, 22], Netrapalli et al. [33] develop an algorithm (AltMinPhase) that alternates between minimizing $\|Ax - Cb\|_2$ in x and in C over diagonal matrices of phases (signs) with modulus 1. Their algorithm is elegant, but the analysis requires resampling a new measurement matrix A and measurements b in each iteration. More recently, Candès et al. [13] develop Wirtinger flow, a gradient-based method that performs a careful modification of gradient descent on the objective

$$F(x) := \frac{1}{2m} \sum_{i=1}^m (|\langle a_i, x \rangle|^2 - b_i)^2,$$

where $x \in \mathbb{C}^n$ may be complex. Wang et al. [46] build on this work by attacking a modification of this objective, showing how to perform a generalized descent method on

$$F(x) := \frac{1}{2m} \sum_{i=1}^m \left(|\langle a_i, x \rangle| - \sqrt{b_i} \right)^2,$$

and providing arguments for the convergence of their method. Wang et al. achieve striking empirical results when the measurements and signals are real-valued, achieving better than 50% perfect signal recovery when the measurement ratio $m/n = 2$, which is essentially at the threshold for injectivity of the real-valued measurements $b = (Ax_\star)^2$. Zhang et al. [47] also study a variant of Wirtinger flow based on median estimates that handles some outliers. Unfortunately, these procedures rely fairly strongly on Gaussianity assumptions, and their gradient descent approaches require subsampling

schemes (to select “good” terms in the sum); these procedures have parameters chosen carefully to reflect Gaussianity in the measurement matrices A , and it is not always clear how to extend them to non-Gaussian measurements.

1.2 Our contributions and outline

In this paper, we focus on prox-linear methods, the iterations (5) for the non-smooth non-convex problem (2). In addition to being (to us at least) aesthetically pleasing, as we minimize the natural objective (2), our approach yields a number of theoretical and practical benefits.

In the literature on signal recovery from phaseless measurements, *stability* of the reconstruction of a signal is of paramount importance. To solve the phase retrieval problem at all, one requires injectivity of the measurements $b = |Ax|^2$, which for real $A \in \mathbb{R}^{m \times n}$ in general position necessitates $m \geq 2n - 1$ and for complex $A \in \mathbb{C}^{m \times n}$ in general position necessitates $m \geq 4n - 2$ (cf. [2]). Stability makes this injectivity more robust. Consider the real-valued case first. Eldar and Mendelson [19] say that a measurement matrix $A \in \mathbb{R}^{m \times n}$ is $\lambda \geq 0$ stable if

$$\|(Ax)^2 - (Ay)^2\|_1 \geq \lambda \|x - y\|_2 \|x + y\|_2 \quad \text{for all } x, y \in \mathbb{R}^n. \quad (6a)$$

Such conditions, which hold with high probability for suitable designs A , are also common in semidefinite relaxation approaches to phase retrieval; cf. Candès et al. [9, Lemma 3.2]. (See also the paper [4].) This condition means that distant signals $x, x' \in \mathbb{R}^n$ cannot be confused in the measurement domain $\{(Ay)^2 \mid y \in \mathbb{R}^n\} \subset \mathbb{R}_+^m$ because A does a good job of separating them; the more stable a measurement matrix, the “easier” the recovery problem should be. In the case in which x is complex, the stability condition (6a) becomes

$$\||Ax|^2 - |Ay|^2\|_1 \geq \lambda \inf_{\theta} \|x - e^{i\theta}y\|_2 \cdot \sup_{\theta} \|x - e^{i\theta}y\|_2 \quad \text{for all } x, y \in \mathbb{C}^n, \quad (6b)$$

a slightly stronger condition. We provide stability guarantees for both situations for general classes of random matrices by adapting Mendelson’s “small ball” techniques [32] (Sec. 3.1). Most literature on non-convex approaches to phase retrieval requires such a stability condition—and usually more because of the quadratic objectives often used—to guarantee signal recovery. In contrast, our procedure requires essentially *only* the stability condition (6), a mild bound on the operator norm $\frac{1}{m} \|A\|_{\text{op}}^2$, and an initialization within some constant factor of $\|x_\star\|_2$ to guarantee both fast convergence and exact signal recovery.

With this in mind, in Section 2 we develop purely optimization-based deterministic results, which build off of classical results on composite optimization, which rely on the stability condition (6). By identifying the conditions required for fast convergence and recovery, we can then spend the remainder of the paper showing how various measurement models guarantee sufficient conditions for our convergence results. In particular, in Section 3, we show how a number of sensing matrices A suffice to guarantee convergence and signal recovery in the noiseless setting, that is, when $b = |Ax_\star|^2$. In Section 4, we extend these results to the case when a constant fraction of the measurements b_i may be arbitrarily corrupted, showing that stability and a somewhat stronger condition on $\|A\|_{\text{op}}$ are still sufficient to guarantee signal recovery; again, these results hold for our basic algorithm with no tuning parameters.

In the final sections of the paper, we provide a substantial empirical evaluation of our proposed algorithms. While our method in principle requires no tuning—it solves a sequence of explicit *convex* problems—there is some art in developing efficient methods for the solution of the sequence of convex optimization problems we solve. In Section 5, we describe these implementation details, and

Data: Initializer x_0 , stepsize α , tolerance $\epsilon > 0$, convex $h : \mathbb{R}^m \rightarrow \mathbb{R}$, suitably smooth
 $c : \mathbb{C}^n \rightarrow \mathbb{R}^m$
set $k = 0$
repeat
 $x_{k+1} = \operatorname{argmin}_{x \in X} \{f_{x_k}(x) + \frac{1}{2\alpha} \|x - x_k\|_2^2\}$ where $f_x(y) = h(c(x) + \operatorname{Re}(\nabla c(x)^H(y - x)))$
 $k = k + 1$
until $\|x_{k-1} - x_k\|_2 \leq \alpha\epsilon$;
return x_k

Algorithm 1: Prox-linear algorithm for problem (3)

in Section 6 we provide experimental evidence of the success of our proposed approach. In reasonably high-dimensional settings ($n \geq 1000$), with real-valued random Gaussian measurements our method achieves perfect signal recovery in about 80-90% of cases even when $m/n = 2$. The method also handles outlying measurements well, substantially improving state-of-the-art performance, and we give applications with measurement matrices that demonstrably fail all of our conditions but for which the method is still straightforward to implement and empirically successful.

Notation We collect our common notation here. We let $\|\cdot\|$ and $\|\cdot\|_2$ denote the usual vector ℓ_2 -norm, and for a matrix $A \in \mathbb{C}^{m \times n}$, $\|A\|_{\text{op}}$ denotes its ℓ_2 -operator norm. The notation A^H means the Hermitian conjugate (conjugate transpose) of $A \in \mathbb{C}^{m \times n}$. For $a \in \mathbb{C}$, $\operatorname{Re}(a)$ denotes its real part and $\operatorname{Im}(a)$ its imaginary part. We take $\langle \cdot, \cdot \rangle$ to be the standard inner product on whatever space it applies; for $u, v \in \mathbb{C}^n$, this is $\langle u, v \rangle = u^H v$, while for $A, B \in \mathbb{C}^{m \times n}$, this is $\langle A, B \rangle = \operatorname{tr}(A^H B)$. Let $\operatorname{quant}_\alpha(\{c_i\})$ denote the α -quantile of a vector $c \in \mathbb{R}^m$, that is, if $c_{(1)} \leq c_{(2)} \leq \dots \leq c_{(m)}$, the α th quantile linearly interpolates $c_{(\lfloor m\alpha \rfloor)}$ and $c_{(\lceil m\alpha \rceil)}$. For a random variable X , $\operatorname{quant}_\alpha(X)$ denotes its α th quantile.

2 Composite Optimization, Algorithm and Convergence Analysis

We begin our development by providing convergence guarantees—under appropriate conditions—for the prox-linear algorithm (the iteration (5)) applied to the composite optimization problem (3), which we recall is to

$$\underset{x}{\text{minimize}} \quad f(x) := h(c(x)) \quad \text{subject to } x \in X,$$

where $h : \mathbb{R}^m \rightarrow \mathbb{R}$ is convex and $c : \mathbb{C}^n \rightarrow \mathbb{R}^m$ is appropriately smooth. In our context, as c is a real-valued complex function, it cannot have an ordinary complex derivative, so some care is required; we use the Wirtinger Calculus, also known as the $\mathbb{C}\mathbb{R}$ -calculus, referring the interested reader to the survey of Kreutz-Delgado [29] for more (see especially [29, Eq. (31)]). In brief, however, because c is real-valued, we let $\nabla c(x)$ denote the Hermitian conjugate of the Jacobian of c , which allows treatment c as a mapping from \mathbb{R}^{2n} to \mathbb{R} , so that $\nabla c(x) \in \mathbb{C}^{n \times m}$ satisfies

$$c(y) = c(x) + \operatorname{Re}(\nabla c(x)^H(y - x)) + O(\|y - x\|^2)$$

as $y \rightarrow x$. We summarize the procedure in Alg. 1 for further reference. In our application to quadratic constraints and phase retrieval, $h(z) = \frac{1}{m} \|z\|_1$ and $c(x) = |Ax|^2 - b$, so that the iteration (5) is the solution of a quadratic problem. (We describe this in more detail in the next section.)

A number of researchers have studied convergence and stopping conditions for Algorithm 1, showing that it converges to stationary points [7], as well as demonstrating that the stopping

condition $\|x_k - x_{k-1}\|_2 \leq \alpha\epsilon$ holds after $O(\epsilon^{-2})$ iterations and guarantees approximate stationarity [16, 18]. Algorithm 1 and the iteration (5) sometimes enjoy fast (local) convergence rates as well. To describe this phenomenon, define the distance function $\text{dist}(x, S) := \inf_y \{\|x - y\|_2 \mid y \in S\}$. We say that h has *weak sharp minima* if it grows linearly away from its minima, meaning $h(z) \geq \inf_z h(z) + \lambda \text{dist}(z, \text{argmin } h)$ for some $\lambda > 0$. Under this condition (with an additional transversality condition between c and $\text{argmin } h$), Burke and Ferris [8] show that convergence of the prox-linear algorithm near points in $X_\star := \{x : c(x) \in \text{argmin } h\}$ is quadratic, because the model (4) of f is quadratically good, but $h(c(x))$ grows linearly away from X_\star . These assumptions can be weakened to growth of $h \circ c$ along its minimizing set [16, Thm. 7.2]. We build from this elegant development—though our problems do not precisely satisfy the weak sharp minima conditions because of outliers—to show how the prox-linear algorithm provides an effective, easy-to-implement, and elegant approach to problems involving solution of quadratic equalities, specifically focusing on phase retrieval.

2.1 Quadratic convergence and the prox-linear method for phase retrieval

We turn now to an analysis of the prox-linear algorithm for phase retrieval problems, providing conditions on the function f sufficient for quadratic convergence. We introduce two conditions on the function $f(x)$ and its linearized form $f_x(y)$ that suffice for this desideratum; as we show in the sequel, these conditions hold with extremely high probability for a number of random measurement models. These conditions are the keystones of our analysis of the (robust) phase retrieval problem.

As motivation for our first condition, consider the phase retrieval problem. If the measurement matrix A satisfies conditions (6) and the measurements b_i are noiseless with $b = |Ax_\star|^2$, then for $f(x) = \frac{1}{m} \| |Ax|^2 - b \|_1$ we have $f(x) - f(x_\star) \geq \lambda \text{dist}(x, X_\star) \|x_\star\|_2$ for the set of signals $X_\star = \{e^{i\theta} x_\star \mid \theta \in \mathbb{R}\}$. When the measurements have noise or outliers, this may still hold, prompting us to define the following

Condition C1. *There exists $\lambda > 0$ such that for all $x \in \mathbb{R}^n$ (or $x \in \mathbb{C}^n$ in the complex case)*

$$f(x) - f(x_\star) \geq \lambda \text{dist}(0, X_\star) \text{dist}(x, X_\star),$$

where X_\star denotes the set of global minima of f .

This condition is a close cousin of Burke and Ferris’s sharp minima condition [8], though it does not require that $c(x_\star) \in \text{argmin}_z h(z)$; based on their work, it is intuitive that it should prove useful in establishing fast convergence of the prox-linear algorithm. The second condition, which is essentially automatically satisfied for the linear approximation (4), is a requirement that the linearized function $f_x(y)$ is quadratically close to $f(y)$.

Condition C2. *There exists $L < \infty$ such that for all $x, y \in \mathbb{R}^n$ (or $x, y \in \mathbb{C}^n$ in the complex case)*

$$|f(y) - f_x(y)| \leq \frac{L}{2} \|x - y\|_2^2.$$

Locally, Condition C2 holds for any composition $f(x) = h(c(x))$ of a convex h with smooth c , but the phase retrieval objective (2) satisfies the bound globally. Indeed, for $a, x, y \in \mathbb{C}^n$ we have

$$|\langle a, y \rangle|^2 = |\langle a, x \rangle|^2 + 2\text{Re}(\langle x, a \rangle \langle a, y - x \rangle) + |\langle a, y - x \rangle|^2,$$

so the linearization (4) of f around $x \in \mathbb{C}^n$ is

$$f_x(y) = \frac{1}{m} \sum_{i=1}^m \left| |\langle a_i, x \rangle|^2 - b_i + 2\text{Re}(\langle x, a_i \rangle \langle a_i, y - x \rangle) \right|. \quad (7)$$

Letting $A = [a_1 \ \cdots \ a_m]^H$ denote the measurement matrix, then using the preceding expansion of $|\langle a, y \rangle|^2$, we have immediately by the triangle inequality that

$$\begin{aligned} -(x - y)^H \left(\frac{1}{m} A^H A \right) (x - y) + f_x(y) &\leq \frac{1}{m} \sum_{i=1}^m ||\langle a_i, y \rangle|^2 - b_i| = f(y) \\ &\leq f_x(y) + (x - y)^H \left(\frac{1}{m} A^H A \right) (x - y). \end{aligned}$$

That is, Condition C2 holds with $L = 2 \left\| \frac{1}{m} A^H A \right\|_{\text{op}}$:

$$|f(y) - f_x(y)| \leq \frac{1}{m} \left\| A^H A \right\|_{\text{op}} \|x - y\|_2^2. \quad (8)$$

Given Conditions C1 and C2, we turn to convergence guarantees for the prox-linear Algorithm 1, which in our case requires solving a sequence of convex quadratic programs. An implementation of Alg. 1 that solves iteration (5) exactly may be computationally challenging. Thus, we allow inaccuracy in the solutions, assuming there exists a sequence of additive accuracy parameters $\epsilon_k \geq 0$ such that the iterates x_k satisfy

$$f_{x_k}(x_{k+1}) + \frac{L}{2} \|x_{k+1} - x_k\|_2^2 \leq \inf_x \left\{ f_{x_k}(x) + \frac{L}{2} \|x - x_k\|_2^2 \right\} + \epsilon_k. \quad (9)$$

We have the following theorem, whose proof we provide in Section 2.2.

Theorem 1. *Let Conditions C1 and C2 hold. Assume that in each step of Algorithm 1, we solve the intermediate optimization problem to accuracy ϵ_k , and define the relative error measures $\beta_k = \frac{2\epsilon_k}{\lambda \text{dist}(0, X_\star)^2}$. Then*

$$\frac{\text{dist}(x_k, X_\star)}{\text{dist}(0, X_\star)} \leq \frac{\lambda}{2L} \max \left\{ \left(\frac{2L}{\lambda} \cdot \frac{\text{dist}(x_0, x_\star)}{\text{dist}(0, X_\star)} \right)^{2^k}, \max_{0 \leq j < k} \left(\frac{4L}{\lambda} \cdot \beta_j \right)^{2^{k-j-1}} \right\}.$$

If $\epsilon_k = 0$ in the solution quality inequality (9), then

$$\frac{\text{dist}(x_k, X_\star)}{\text{dist}(0, X_\star)} \leq \frac{\lambda}{L} \left(\frac{L}{\lambda} \cdot \frac{\text{dist}(x_0, x_\star)}{\text{dist}(0, X_\star)} \right)^{2^k}.$$

Theorem 1 motivates our approach for the remainder of the paper: we can guarantee exact, accurate, and fast solutions to the phase retrieval problem under the three conditions

1. Stability (Condition C1),
2. Quadratic approximation (Condition C2), via an upper bound on $\left\| A^H A \right\|_{\text{op}}$ and application of inequality (8) and
3. An initializer x_0 of the iterations that is good enough, meaning that it satisfies the constant relative error guarantee $\text{dist}(x_0, X_\star) \leq \text{dist}(0, X_\star) \frac{\lambda}{L}$.

In the coming sections, we show that each of these three conditions holds in both noiseless measurement models (Section 3) and with adversarially perturbed measurements b_i (Section 4).

Before continuing, we provide a few additional remarks regarding Theorem 1. First, if ϵ_k are very small because we solve the intermediate steps (5) to (near) machine precision, then for all intents

and purposes about five iterations suffice for machine precision accurate solutions. Quadratic convergence is also achievable with errors in inequality (9); if the minimization accuracies decrease quickly enough that $\epsilon_k \leq 2^{-2^k}$, then we certainly still have quadratic convergence. More broadly, Theorem 1 shows that the accuracy of solution in iteration j need not be very high to guarantee high accuracy reconstruction of the signal x_* ; only in the last few iterations is moderate to high accuracy necessary. If it is computationally cheap, it is thus advantageous—as we explore in our experimental work—to solve early iterations of the prox-linear method inaccurately.

2.2 Proof of Theorem 1

We prove the result in two steps: we first provide a per-iteration progress guarantee, and then we use this guarantee to show quadratic convergence.

Let $x_* \in X_*$ be any global optimum of the objective $f(x)$. The function $x \mapsto f_{x_k}(x) + \frac{L}{2} \|x - x_k\|_2^2$ is L -strongly convex in x . If we define $x_{*,k+1}$ to be the exact minimizer of $f_{x_k}(x) + \frac{L}{2} \|x - x_k\|_2^2$, the standard optimality conditions for strongly convex minimization imply

$$\begin{aligned} f_{x_k}(x_{k+1}) + \frac{L}{2} \|x_{k+1} - x_k\|_2^2 &\leq f_{x_k}(x_{*,k+1}) + \frac{L}{2} \|x_{*,k+1} - x_k\|_2^2 + \epsilon_k \\ &\leq f_{x_k}(x_*) + \frac{L}{2} \|x_* - x_k\|_2^2 - \frac{L}{2} \|x_* - x_{*,k+1}\|_2^2 + \epsilon_k. \end{aligned}$$

Using the approximation Condition C2, so that $\frac{L}{2} \|x_k - x_{k+1}\|_2^2 + f_{x_k}(x_{k+1}) \geq f(x_{k+1})$ and $f_{x_k}(x_*) \leq f(x_*) + \frac{L}{2} \|x_k - x_*\|_2^2$, we have by substituting in the preceding inequality that

$$\begin{aligned} f(x_{k+1}) &\leq f_{x_k}(x_*) + \frac{L}{2} \|x_* - x_k\|_2^2 - \frac{L}{2} \|x_* - x_{*,k+1}\|_2^2 + \epsilon_k \\ &\leq f(x_*) + L \|x_k - x_*\|_2^2 - \frac{L}{2} \|x_* - x_{*,k+1}\|_2^2 + \epsilon_k. \end{aligned}$$

Summarizing, we get for all $x_* \in X_*$ and $k \in \mathbb{N}$,

$$f(x_{k+1}) - f(x_*) + \frac{L}{2} \|x_* - x_{*,k+1}\|_2^2 \leq L \|x_k - x_*\|_2^2 + \epsilon_k \quad (10)$$

By applying the stability Condition C1, we immediately obtain the progress guarantee

$$\lambda \|x_*\|_2 \text{dist}(x_{k+1}, X_*) + \frac{L}{2} \|x_{*,k+1} - x_*\|_2^2 \leq L \|x_k - x_*\|_2^2 + \epsilon_k \quad (11)$$

We now transform the guarantee (11) into one involving only x_k , x_{k+1} , and x_* , rather than $x_{*,k+1}$, by bounding the difference between $x_{*,k+1}$ and x_{k+1} . The L -strong convexity of $f_{x_k}(\cdot) + \frac{L}{2} \|\cdot - x_k\|_2^2$ implies that

$$\begin{aligned} \epsilon_k + f_{x_k}(x_{*,k+1}) + \frac{L}{2} \|x_{*,k+1} - x_k\|_2^2 &\geq f_{x_k}(x_{k+1}) + \frac{L}{2} \|x_{k+1} - x_k\|_2^2 \\ &\geq f_{x_k}(x_{*,k+1}) + \frac{L}{2} \|x_{*,k+1} - x_k\|_2^2 + \frac{L}{2} \|x_{k+1} - x_{*,k+1}\|_2^2, \end{aligned}$$

whence we obtain $\frac{L}{2} \|x_{*,k+1} - x_{k+1}\|_2^2 \leq \epsilon_k$. Using the standard quadratic inequality $\|x_{k+1} - x_*\|_2^2 \leq 2 \|x_{k+1} - x_{*,k+1}\|_2^2 + 2 \|x_{*,k+1} - x_*\|_2^2$, we thus have by expression (11) that, for all $x_* \in X_*$,

$$\lambda \|x_*\|_2 \text{dist}(x_{k+1}, X_*) + \frac{L}{4} \|x_{k+1} - x_*\|_2^2 \leq L \|x_k - x_*\|_2^2 + 2\epsilon_k \quad (12)$$

Now, taking infimum over $x_\star \in X_\star$ over both sides for Eq (12), we find that,

$$\lambda \operatorname{dist}(0, X_\star) \operatorname{dist}(x_{k+1}, X_\star) \leq L \operatorname{dist}(x_k, X_\star)^2 + 2\epsilon_k.$$

Dividing each side by $\lambda \operatorname{dist}(0, X_\star)^2$ yields

$$\frac{\operatorname{dist}(x_{k+1}, X_\star)}{\operatorname{dist}(0, X_\star)} \leq \frac{L}{\lambda} \frac{\operatorname{dist}(x_k, X_\star)^2}{\operatorname{dist}(0, X_\star)^2} + \frac{2\epsilon_k}{\lambda \operatorname{dist}(0, X_\star)^2}. \quad (13)$$

Inductively applying inequality (13) when $\epsilon_k = 0$ yields the second statement of the theorem. When $\epsilon_k > 0$, a brief technical lemma shows the convergence rate:

Lemma 2.1. *Let the sequences $a_k \geq 0$, $\epsilon_k \geq 0$ satisfy $a_k \leq \kappa a_{k-1}^2 + \epsilon_{k-1}$. Then*

$$a_k \leq \max \left\{ (2\kappa)^{2^k-1} a_0^{2^k}, \max_{0 \leq j < k} (2\kappa)^{2^{k-j}-1} (2\epsilon_j)^{2^{k-j}-1} \right\}.$$

Proof The proof is by induction. For a_1 , we certainly have $a_1 \leq 2\kappa \vee 2\epsilon_0$ because both sequences are non-negative. For the general case, assume the result holds for a_k , where k is arbitrary. Then

$$\begin{aligned} a_{k+1} &\leq \kappa a_k^2 + \epsilon_k \leq \max \{ 2\kappa a_k^2, 2\epsilon_k \} \\ &\leq 2\kappa (2\kappa)^{2^{k+1}-2} a_0^{2^{k+1}} \vee \max_{0 \leq j < k} \left\{ 2\kappa (2\kappa)^{2^{k-j}-2} (2\epsilon_j)^{2^{k-j}-2} \right\} \vee 2\epsilon_k \\ &= \max \left\{ (2\kappa)^{2^{k+1}-1} a_0^{2^{k+1}}, \max_{0 \leq j < k+1} (2\kappa)^{2^{k-j}-1} (2\epsilon_j)^{2^{k-j}-1} \right\} \end{aligned}$$

as desired. □

Applying Lemma 2.1 in inequality (13) with $\kappa = \frac{L}{\lambda}$ and $a_k = \frac{\operatorname{dist}(x_k, X_\star)}{\operatorname{dist}(0, X_\star)}$ yields the theorem.

3 Noiseless Phase Retrieval Problem

We begin our discussion of the phase retrieval problem by considering the noiseless case, that is, when the observations $b_i = |\langle a_i, x_\star \rangle|^2$. Throughout this section and the remainder of the paper, the vectors a_i are assumed to be independent and identically distributed copies of a random vector $a \in \mathbb{C}^n$. Based on our discussion after Theorem 1, we show that (i) the objective (2) is stable C1, (ii) is quadratically approximable C2, and (iii) that we have a good initializer x_0 . In the coming three sections, we address each of these in turn, providing progressively stronger assumptions that are sufficient for each condition to hold with high probability as soon as the number of measurements $m/n > c$, where c is a numerical constant. In Section 3.4 we provide a summary theorem that encapsulates our results. For readability, we defer proofs to Sec. A.

3.1 Stability

Our first step is to provide conditions under which stability holds. We divide our discussion of stability conditions into the real and complex cases, as the real case is considerably easier.

3.1.1 Stability for real-valued vectors

With the stability condition in mind, we make the following assumption on the random measurement vectors $a \in \mathbb{R}^n$:

Assumption A1. *There exist constants $\kappa_{\text{st}} > 0$ and $p_0 > 0$ such that for all $u, v \in \mathbb{R}^n$, $\|u\|_2 = \|v\|_2 = 1$, we have*

$$\mathbb{P}(|\langle a, u \rangle| \wedge |\langle a, v \rangle| \geq \kappa_{\text{st}}) \geq p_0.$$

Intuitively, Assumption A1 says that the measurement vectors $a \in \mathbb{R}^n$ have sufficient support in all directions $u, v \in \mathbb{R}^n$. One simple sufficient condition for this is a type of small-ball condition [32], as follows, which makes it clear that Assumption A1 requires no light tails: just that the probability of one of $|\langle a, u \rangle|$ and $|\langle a, v \rangle|$ being small is small.

Example 1 (Stability by the small-ball method): Assume that we may choose positive but small enough $\varepsilon > 0$ that

$$\sup_{\|u\|_2=1} \mathbb{P}(|\langle a, u \rangle| < \varepsilon) < \frac{1}{2}.$$

Then by the union bound, the choice $p_0 = 1 - 2 \sup_{\|u\|_2} \mathbb{P}(|\langle a, u \rangle| < \varepsilon) > 0$ and $\kappa_{\text{st}} = \varepsilon$ immediately yields

$$\begin{aligned} \mathbb{P}(|\langle a, u \rangle| \wedge |\langle a, v \rangle| \geq \kappa_{\text{st}}) &= 1 - \mathbb{P}(|\langle a, u \rangle| < \varepsilon \text{ or } |\langle a, v \rangle| < \varepsilon) \\ &\geq 1 - \mathbb{P}(|\langle a, u \rangle| < \varepsilon) - \mathbb{P}(|\langle a, v \rangle| < \varepsilon) \geq p_0. \end{aligned}$$

As a further specialization, if $a \sim \mathcal{N}(0, I_n)$ is an isotropic Gaussian, then the choice $\kappa_{\text{st}} = .31$ yields $\mathbb{P}(|\langle a, u \rangle| \leq \kappa_{\text{st}}) \leq \frac{1}{4}$, so we may take $\kappa_{\text{st}} = .31$ and $p_0 = \frac{1}{2}$. \diamond

As we note in the discussion preceding Condition C1, for the objective (2) it is immediate that in the noiseless case that,

$$f(x) - f(\pm x_\star) = f(x) = \frac{1}{m} \|(Ax)^2 - (Ax_\star)^2\|_1 = \frac{1}{m} \sum_{i=1}^m |\langle a_i, x - x_\star \rangle \langle a_i, x + x_\star \rangle|.$$

Thus, if we can show the stability condition (6a)—equivalently, that $\sum_{i=1}^m |\langle a_i, u \rangle \langle a_i, v \rangle| \geq \lambda m$ for $u, v \in \mathbb{S}^{n-1}$ —then Condition C1 holds for $X_\star = \{\pm x_\star\}$. To that end, we provide the following guarantee, which we prove in Sec. A.1.

Proposition 1. *Let Assumption A1 hold. There exists a numerical constant $c < \infty$ such that for any $t \geq 0$,*

$$\mathbb{P}\left(\frac{1}{m} \sum_{i=1}^m |\langle a_i, u \rangle \langle a_i, v \rangle| \geq \kappa_{\text{st}}^2 \cdot \left(p_0 - c\sqrt{\frac{n}{m}} - \sqrt{2t}\right) \text{ for all } u, v \in \mathbb{S}^{n-1}\right) \geq 1 - 2e^{-mt}.$$

Proposition 1 immediately yields the following corollary, which shows that the stability condition C1 holds with high probability for $m/n \gtrsim 1$.

Corollary 3.1. *Let Assumption A1 hold. Then there exists a numerical constant $c < \infty$ such that if $mp_0^2 \geq cn$, then*

$$\mathbb{P}\left(f(x) - f(x_\star) \geq \frac{1}{2} \kappa_{\text{st}}^2 p_0 \|x - x_\star\|_2 \|x + x_\star\|_2 \text{ for all } x \in \mathbb{R}^n\right) \geq 1 - 2 \exp\left(-\frac{mp_0^2}{32}\right).$$

Eldar and Mendelson [19] establish the stability Condition C1 under more restrictive assumptions on the distribution of the measurement vectors $\{a_i\}_{i=1}^m$. Concretely, they require that the distribution of a is subgaussian (see Definition 3.1 to come) and isotropic (meaning that $\mathbb{E}[aa^T] = I_n$). As Example 1 makes clear, our result only requires weaker small ball assumptions without any restrictions on tails or the covariance structure of the random vector a .

3.1.2 Complex Case

We now investigate conditions sufficient for stability of A in the measurement vectors a_i are complex-valued. In this case, the argument is not quite so simple, as stability for complex vectors requires more uniform notions of function growth. Accordingly, we make the following two assumptions on the random vectors $a_i \in \mathbb{C}^n$. The first is a small-ball type assumption, while the second requires that the vectors a_i are appropriately uniform in direction. To fully state our assumptions, we require an additional definition on the sub-Gaussianity of random vectors. We define this in terms of Orlicz norms (following [44, 43, Ch. 2.2]).

Definition 3.1. *The random vector $a \in \mathbb{C}^n$ is σ^2 -sub-Gaussian if for all $v \in \mathbb{C}^n$, $\|v\|_2 = 1$,*

$$\mathbb{E} \left[\exp \left(\frac{|\langle a, v \rangle|^2}{\sigma^2} \right) \right] \leq e.$$

With this definition, we now provide assumptions on the random measurement vector $a \in \mathbb{C}^n$.

Assumption A2. *There exists a non-increasing function $h : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $h(0) = 0$ such that for $0 \leq \epsilon$,*

$$\mathbb{P}(\|a\|_2 \leq \epsilon\sqrt{n}) \leq h(\epsilon).$$

We also require that the a , when normalized, are sufficiently uniform.

Assumption A3. *Let $w = \sqrt{n}a / \|a\|_2$. The random vector w is σ^2 -sub-Gaussian (Definition 3.1). In addition, for any matrix $X \in \mathbb{C}^{n \times n}$ with rank at most 2,*

$$\mathbb{E} [|w^H X w|] \geq \tau^2 \|X\|_{\text{Fr}}.$$

Assumption A3 may seem somewhat challenging to verify, but it holds for any rotationally symmetric distribution, and moreover, in this case we have that $\tau \geq c\sigma$ for a numerical constant $c > 0$.

Example 2 (Rotationally symmetric measurements): Let the measurement vectors a_i be rotationally symmetric, so that for unitary $U \in \mathbb{C}^{n \times n}$, the distribution of Ua is identical to a . We show that Assumption A3 holds. In this case, $w = \sqrt{n}a / \|a\|_2$ is uniform on the radius- \sqrt{n} sphere in \mathbb{C}^n , and standard results in convex geometry [3] show w is $O(1)$ -sub-Gaussian (Definition 3.1). As a is rotationally symmetric, w is also equal in distribution to $\sqrt{n}z / \|z\|_2$ where z is standard complex normal; thus, we have for any rank 2 or less Hermitian X that

$$\mathbb{E}[|w^H X w|] = n \mathbb{E}[|z^H X z|] \mathbb{E}[1/\|z\|_2^2] \stackrel{(i)}{\geq} \mathbb{E}[|z^H X z|] \stackrel{(ii)}{\geq} \frac{2\sqrt{2}}{\pi} \|X\|_{\text{Fr}}$$

where inequality (i) is a consequence of $\mathbb{E}[1/\|z\|_2^2] \geq \frac{1}{n}$ and inequality (ii) is a calculation (see Lemma C.2 in Appendix C.3) as X is rank 2. \diamond

With these assumptions in place, we have the following stability guarantee for the random matrix A . We defer the proof to Appendix A.2.

Proposition 2. *Let Assumptions A2 and A3 hold. Let $c > 0$ be chosen such that $h(c) \leq \frac{1}{2(1+c)} \frac{\tau^4}{\sigma^4}$. There exist numerical constants $c_0 > 0$ and $c_1 < \infty$ such that with probability at least*

$$1 - \exp \left(-c_0 m \frac{\tau^4}{\sigma^4} + c_1 n \log \frac{\sigma^2}{\tau^2} \right),$$

we have

$$\frac{1}{m} \left\| |Ax|^2 - |Ay|^2 \right\|_1 \geq \frac{c^2 \tau^2}{4} \cdot \inf_{\theta} \|x - e^{i\theta} y\|_2 \cdot \sup_{\theta} \|x - e^{i\theta} y\|_2$$

simultaneously for all $x, y \in \mathbb{C}^n$.

The proposition shows that if the ratio between the growth constant τ and the sub-Gaussian constant σ in Assumption A3 is bounded, as it is in the case of rotationally invariant vectors a (Example 2), then we have the complex stability guarantee (6b) as soon as $m \gtrsim n$ with extremely high probability.

3.2 Quadratic Approximation

With the stability condition C1 in place, we turn to a discussion of the approximation condition C2. As implied by the estimate in inequality (8), the quadratic approximation condition is satisfied with parameter $L = 2\|\frac{1}{m}A^H A\|_{\text{op}}$. To control this quantity, we require that the rows of the matrix $A \in \mathbb{C}^{m \times n}$ be sufficiently light-tailed.

Assumption A4. *The random vector $a \in \mathbb{C}^n$ is σ^2 -sub-Gaussian.*

It is of course possible to bound $\|A\|_{\text{op}}$ when the rows have heavier tails using appropriate symmetrization techniques and matrix Khintchine inequalities (cf. [44, Sec. 2.6]); the extension is clear, so we do not address such issues.

Certainly, not all measurement vectors a_i satisfy Assumption A4. Using that $\mathbb{E}[e^{\lambda Z^2}] = [1 - 2\lambda]_+^{-\frac{1}{2}}$ for $Z \sim \mathcal{N}(0, 1)$, it holds for standard real normal vectors $a_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, I_n)$ or complex normal vectors $a_i \stackrel{\text{iid}}{\sim} \frac{1}{\sqrt{2}}(\mathcal{N}(0, I_n) + i\mathcal{N}(0, I_n))$ with $\sigma^2 = \frac{2e^2}{e^2 - 1} \approx 2.313$. Similarly, it also holds for a_i uniform on \mathbb{S}^{n-1} with $\sigma^2 = O(1) \cdot \frac{1}{n}$. In practice it may be useful to apply our algorithm to the transformed data $\{a_i / \|a_i\|_2\}_{i=1}^m$ and $\{b_i / \|a_i\|_2^2\}_{i=1}^m$, which (in the noiseless case or case with infrequent but arbitrary corruptions of b_i) is likely to make the problem better conditioned. There are two heuristic motivations for this: first, those measurement vectors with larger magnitudes $\|a_i\|$ tend to place a higher weight in the optimization problem (2), and thus normalization can make the observations “comparable” to each other; second, normalization guarantees the measurement vectors $\{a_i\}_{i=1}^m$ satisfy Assumption A4, yielding easier verification of Condition C2. (It may be more challenging to verify Assumption A1, but if the a_i are sufficiently isotropic this presents no special difficulties.)

Standard results guarantee that the random matrices A have well-behaved singular vectors whenever Assumption A4 holds; we provide one such result due to Vershynin [44, Thm. 39, Eq. (25)] with constants that are achievable by tracing his proof.

Lemma 3.1. *Let Assumption A4 hold and $\Sigma = \mathbb{E}[aa^H]$. Then for all $t \geq 0$,*

$$\mathbb{P}\left(\left\|\frac{1}{m}A^H A - \Sigma\right\|_{\text{op}} \geq 11\sigma^2 \max\left\{\sqrt{\frac{4n}{m}} + t, \frac{4n}{m} + t\right\}\right) \leq \exp(-mt).$$

Moreover, $\|\Sigma\|_{\text{op}} \leq \sigma^2$ and $\mathbb{E}[|\langle a, v \rangle|^k] \leq \Gamma(\frac{k}{2} + 1)e\sigma^k$ for all $k \geq 0$.

Thus, we have the following corollary of Lemma 3.1, which guarantees that Condition C2 holds with high probability for $m/n \gtrsim 1$.

Corollary 3.2. *Let Assumption A4 hold. Then there exists a numerical constant $c < \infty$ such that whenever $m \geq cn$*

$$\mathbb{P}\left(|f_x(y) - f(y)| \leq 2\sigma^2 \|x - y\|_2^2 \text{ holds uniformly for } x, y \in \mathbb{R}^n\right) \geq 1 - \exp\left(-\frac{m}{c}\right).$$

Proof Assume $m \geq cn$ for c large enough, and choose t small enough in Lemma 3.1 that $11\sigma^2 \sqrt{4n/m} + t \leq \sigma^2$. Applying the triangle inequality to $\|\frac{1}{m}A^H A - \Sigma\|_{\text{op}} \leq \|\frac{1}{m}A^H A\|_{\text{op}} + \sigma^2$ yields the result by equation (8). \square

Data: Measurement matrix $A \in \mathbb{C}^{m \times n}$ and signals $b = |Ax_\star|^2$

begin

Set $\hat{r}^2 = \frac{1}{m} \sum_{i=1}^m b_i$

Select indices $\mathcal{I}_{\text{sel}} := \{i \in [m] : b_i \leq \frac{1}{2}\hat{r}^2\}$

Construct directional estimates \hat{d} by

$$X^{\text{init}} := \sum_{i \in \mathcal{I}_{\text{sel}}} a_i a_i^H, \quad \hat{d} = \underset{d \in \mathbb{S}^{n-1}}{\text{argmin}} d^H X^{\text{init}} d$$

return (\hat{r}, \hat{d})

end

Algorithm 2: Initialization procedure for non-noisy data

3.3 Initialization

The last ingredient in achieving strong convergence guarantees for our prox-linear procedure for phase retrieval is to provide a good initialization. There are a number of initialization strategies in the literature [13, 46, 47] based on spectral techniques, which work as follows. We decompose the initialization into two steps: we (i) find an estimate of the direction $d_\star := x_\star / \|x_\star\|_2$, and (ii) estimate the magnitude $r_\star := \|x_\star\|_2$. The latter is easy: assuming that $\mathbb{E}[aa^H] = I_n$, one simply uses $\hat{r}^2 = m^{-1} \sum_{i=1}^m b_i^2$, which is unbiased and tightly concentrated. The former, the direction estimate, is somewhat trickier.

Wang, Giannakis, and Eldar [46] provide an empirically excellent initialization whose heuristic justification is as follows. First, for random vectors a_i in high dimensions, we expect a_i to usually be orthogonal to the direction d_\star . Thus, by extracting the smallest magnitude $b_i = |\langle a_i, x_\star \rangle|^2$, we have the vectors a_i that are “most” orthogonal to the direction d_\star ; letting \mathcal{I}_{sel} be these small indices, the eigenvector corresponding to the smallest eigenvalue (for simplicity, we simply call this the smallest eigenvector) of $\sum_{i \in \mathcal{I}_{\text{sel}}} a_i a_i^H$ should be close to the direction d_\star . A variant of this procedure is to note that $\frac{1}{m} \sum_{i=1}^m a_i a_i^H \approx I_n$ when the a_i are isotropic, so that the largest eigenvector of $\sum_{i \notin \mathcal{I}_{\text{sel}}} a_i a_i^H$ should also be close to d_\star . This initialization strategy has the added benefit that—unlike the original spectral initialization schemes developed by Candès et al. [13], which rely on eigenvectors of $\sum_{i=1}^m b_i a_i a_i^H$ that may not concentrate at sub-Gaussian rates (as the sum involves fourth moments of random vectors)—the sums $\sum_i a_i a_i^H$ are tightly concentrated.

Unfortunately, we believe Wang et al.’s proof that this initialization works contains a mistake (note that they consider only the case where $\{a_i\}_{i=1}^m$ are real): letting $U \in \mathbb{R}^{n \times (n-1)}$ be an orthogonal matrix whose columns are all orthogonal to d_\star , in the proof of Lemma 2 (Eqs. (68)–(70) in [46]) they assert that $|\mathcal{I}_{\text{sel}}^c|^{-1} \sum_{i \in \mathcal{I}_{\text{sel}}^c} U^H a_i a_i U \rightarrow I_{n-1}$ when the a_i are uniform on $\sqrt{n}\mathbb{S}^{n-1}$. This is not true (nor does appropriate normalization by n or $n-1$ make it true), as it ignores the subtle effects of conditioning in the construction of \mathcal{I}_{sel} . In spite of this issue, the initialization they propose works remarkably well, and as we show presently, it provably provides a good estimate \hat{d} of d_\star . We include the initialization procedure in Algorithm 2.

With the previous discussion in mind, we provide a general assumption that is sufficient for Alg. 2 to return a direction and magnitude estimate sufficiently accurate for phase retrieval.

Assumption A5. *For some $\epsilon_0 \in (0, 1]$ and $\mathfrak{p}_0(d_\star) > 0$, the following hold.*

(i) For all $\epsilon \in (0, \epsilon_0]$, the following continuity and directional likelihood conditions hold:

$$\mathbb{P} \left(|\langle a, d_\star \rangle|^2 \in \left[\frac{1-\epsilon}{2}, \frac{1+\epsilon}{2} \right] \right) \leq \kappa\epsilon \quad \text{and} \quad \mathbb{P} \left(|\langle a, d_\star \rangle|^2 \leq \frac{1-\epsilon}{2} \right) \geq \mathbf{p}_0(d_\star) > 0.$$

(ii) There exist functions $\phi : [0, \epsilon_0] \rightarrow \mathbb{R}_+$ and $\Delta : [0, \epsilon_0] \rightarrow \mathbb{C}^{n \times n}$ such that

$$\mathbb{E} \left[aa^H \mid |\langle a, d_\star \rangle|^2 \leq \frac{1-\epsilon}{2} \right] = I_n - \phi(\epsilon) d_\star d_\star^H + \Delta(\epsilon) \quad \text{for } \epsilon \in [0, \epsilon_0].$$

(iii) $\mathbb{E}[aa^H] = I_n$.

Assumption A5 on its face seems complex, but each of its components is not too stringent. Part (i) says that $|\langle a, d_\star \rangle|^2$ has no point mass at $|\langle a, d_\star \rangle|^2 = \frac{1}{2}$ and that $|\langle a, d_\star \rangle|^2$ has reasonable probability of being smaller than $\frac{1}{2}$. Part (iii) simply states that in expectation, a is isotropic (and a rescaling of a can guarantee this). Part (ii) is the most subtle and essential for our derivation; it says that $a \in \mathbb{R}^n / \mathbb{C}^n$ is reasonably isotropic, even if we condition on $|\langle a, d_\star \rangle|$ being near zero for some direction d_\star , so that most mass of aa^H is distributed uniformly in the orthogonal directions $I_n - d_\star d_\star^H$. The error terms ϕ and Δ allow non-trivial latitude in this condition, so that Assumption A5 holds for more than just Gaussian vectors. That said, for concreteness we provide the following example.

Example 3 (Gaussian vectors and conditional directions): We consider the standard cases that $a_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, I_n)$, or the complex counterpart $a_i \stackrel{\text{iid}}{\sim} \frac{1}{\sqrt{2}} (\mathcal{N}(0, I_n) + i\mathcal{N}(0, I_n))$, showing that such a_i satisfy Assumption A5 for any $\epsilon_0 \in (0, 1)$ with residual error $\Delta \equiv 0$. Clearly Part (iii) holds. For Part (i), note that $|\langle a, d_\star \rangle|^2$ is χ_1^2 -distributed with density $f(t) = e^{-t/2} (2\pi t)^{-\frac{1}{2}}$. Integrating the density using its upper bound, we may set $\kappa \leq f(\frac{1-\epsilon_0}{2}) = \exp(\frac{\epsilon_0-1}{4}) / \sqrt{\pi(1-\epsilon_0)}$ and $\mathbf{p}_0(d_\star) = \mathbb{P}(\chi_1^2 \leq \frac{1-\epsilon_0}{2}) \geq \frac{1}{2} \sqrt{1-\epsilon_0}$. Part (ii) is all that remains. By the rotational invariance of $a \sim \mathcal{N}(0, I_n)$, we see for any $t \in \mathbb{R}_+$ that

$$\mathbb{E}[aa^H \mid |\langle a, d_\star \rangle|^2 \leq t] = I_n - d_\star d_\star^H + \mathbb{E}[|\langle a, d_\star \rangle|^2 \mid |\langle a, d_\star \rangle|^2 \leq t] d_\star d_\star^H.$$

We claim the following lemma, whose proof we provide in Appendix C.1.

Lemma 3.2. *Let Z be a continuous random variable with a density symmetric about zero and decreasing on \mathbb{R}_+ . Then for any $c \in \mathbb{R}$ we have $\mathbb{E}[Z^2 \mid Z^2 \leq c^2] \leq \frac{c^2}{3}$.*

Thus, setting $\phi(\epsilon) = 1 - \mathbb{E}[|\langle a, d_\star \rangle|^2 \mid |\langle a, d_\star \rangle|^2 \leq \frac{1-\epsilon}{2}] \geq 1 - \frac{(1-\epsilon)^2}{6} \geq \frac{5}{6}$, we see that part (ii) of Assumption A5 holds with $\phi(\epsilon) \geq \frac{5}{6}$ and $\Delta(\epsilon) \equiv 0$. \diamond

We now state our main proposition of this section.

Proposition 3. *Let Assumptions A4 and A5 hold and let ϵ_0 and $\mathbf{p}_0(d_\star)$ be as in Assumption A5. Define the error measure*

$$\nu(\epsilon) := \left[1 + \frac{4e(1 + \log \frac{1}{1 \wedge \kappa\epsilon})\sigma^2\kappa}{\mathbf{p}_0(d_\star)} + \frac{2\kappa(1+\epsilon)}{\mathbf{p}_0(d_\star)^2} \right] \epsilon.$$

There exists a numerical constant $c > 0$ such that the following holds. Let (\hat{r}, \hat{d}) be the estimated magnitude and direction of Alg. 2 and define $x_0 = \hat{r}\hat{d}$. For any $\epsilon \in [0, \epsilon_0]$, if

$$c \cdot \frac{m}{n} \geq \frac{\sigma^4 \log^2 \mathbf{p}_0(d_\star)}{\mathbf{p}_0(d_\star)\epsilon^2} \vee \frac{1}{(\kappa\epsilon)^2}$$

then

$$\frac{\text{dist}(x_0, X_\star)}{\|x_\star\|_2} \leq \sqrt{2(1+\epsilon)} \frac{\|\Delta(\epsilon)\|_{\text{op}} + \nu(\epsilon)}{\left[\phi(\epsilon) - \|\Delta(\epsilon)\|_{\text{op}} - \nu(\epsilon)\right]_+} + \epsilon \quad (14)$$

with probability at least

$$1 - \exp\left(-\frac{cm\epsilon^2}{\sigma^4}\right) - 2\exp(-cm\epsilon^2\kappa^2) - \exp\left(-\frac{mp_0(d_\star)^2}{2}\right) - \exp\left(-\frac{cm\kappa\epsilon^2}{\sigma^4 \log^2 p_0(d_\star)}\right).$$

We prove Proposition 3 in two parts. In the first part (Sec. A.3), we define a number of events and proceed conditionally, showing that if each of the events occurs then the conclusion (14) holds. In the second part (Sec. A.4) we show that the events occur with high probability.

We provide a few remarks to make the result clearer. Let us make the simplifying assumptions that the constants in Assumption A5 are absolute (which is true for Gaussian measurements), that is, that $\sigma^2 = O(1)$ and $p_0(d_\star) = \Omega(1)$ (it is no loss of generality to assume $\kappa \geq 1$). Then for numerical constants $c > 0, C < \infty$ we have for any $\epsilon \in [0, \epsilon_0]$ that the error measure $\nu(\epsilon) \leq C\epsilon \log \frac{1}{\epsilon}$, and with probability at least $1 - 5\exp(-cm\epsilon^2)$ that

$$\frac{m}{n} \geq \frac{C}{\epsilon^2} \text{ implies } \frac{\text{dist}(x_0, X_\star)}{\|x_\star\|_2} \leq C \frac{\|\Delta(\epsilon)\|_{\text{op}} + \nu(\epsilon)}{\left[\phi(\epsilon) - \|\Delta(\epsilon)\|_{\text{op}} - \nu(\epsilon)\right]_+} + \epsilon.$$

Here, we see three competing terms. The first two, the separation $\phi(\epsilon)$ and error $\Delta(\epsilon)$, arise from the conditional expectation of Assumption A5(ii), where $\mathbb{E}[aa^H \mid \langle a, d_\star \rangle^2 \leq \frac{1-\epsilon}{2}] = I_n - \phi(\epsilon)d_\star d_\star^H + \Delta(\epsilon)$. This is intuitive: the larger the separation $\phi(\epsilon)$ from uniformity in the conditional expectation of aa^H , the easier it is for spectral initialization to succeed; larger error $\Delta(\epsilon)$ will hide the directional signal d_\star . The last term is the error $\nu(\epsilon) \lesssim \epsilon \log \frac{1}{\epsilon}$, which approaches 0 nearly as quickly as $\epsilon \rightarrow 0$.

In the case that the error $\Delta(\epsilon) = 0$ and gap $\phi(\epsilon)$ is bounded away from zero, which holds for elliptical distributions with identity covariance—the Gaussian distribution and uniform distribution on the sphere being the primary examples—we thus see that as soon as $\frac{m}{n} \gtrsim \epsilon^{-2}$ we have relative error

$$\frac{\text{dist}(x_0, X_\star)}{\|x_\star\|_2} \lesssim \epsilon \log \frac{1}{\epsilon} \text{ with probability } \geq 1 - 5\exp(-cm\epsilon^2). \quad (15)$$

That is, we can construct an arbitrarily good initialization with large enough sample size. (This proves that the initialization scheme of Wang et al. [46] also succeeds with high probability.) On the other hand, when $\Delta(\epsilon) \neq 0$ for all $\epsilon \in [0, \epsilon_0]$, then Proposition 3 cannot guarantee arbitrarily good initialization: the error term $\|\Delta(\epsilon)\|_{\text{op}}$ is never zero. However, if it is small enough, we still achieve initializers that are within *constant* relative distance of x_\star , which is good enough for Theorem 1.

3.4 Summary and success guarantees

We now have guarantees of stability, quadratic approximation, and good initialization for appropriate measurement matrices $A \in \mathbb{R}^{m \times n}$ when the observations $b = |Ax_\star|^2$. We provide a summary theorem showing that our composite optimization procedure works as soon as the sample size is large enough. In stating the theorem, we assume that each of Assumptions A1, A4, and A5 holds with all of their constants actually numerical constants. The one somewhat technical assumption we require is that relating the sub-Gaussian parameter σ^2 , the stability parameters κ_{st} and p_0 (and in the complex case τ and the function $h(\cdot)$), and the error $\Delta(\epsilon)$ and directional separation constants $\phi(\epsilon)$. Denote the stability constant $\nu = \kappa_{\text{st}}^2 p_0$ in the real case and $\nu = \left(h^{-1}\left(\frac{\tau^4}{2(1+\epsilon)\sigma^4}\right)\tau\right)^2$.

Then, in particular, if we assume that for a suitably small numerical constant $c > 0$, we have

$$\|\Delta(\epsilon)\|_{\text{op}} \leq c \frac{\nu}{\sigma^2}, \quad \phi(\epsilon) \geq \frac{1}{2} \|\Delta(\epsilon)\|_{\text{op}}, \quad \text{and} \quad \phi(\epsilon) \geq \phi_\star > 0$$

for all $\epsilon \in [0, \epsilon_0]$. We then have the following theorem, which follows by combining our convergence Theorem 1 with Proposition 1, Corollary 3.2, and Proposition 3.

Theorem 2. *Let the conditions of the preceding paragraph hold. There exists a numerical constant $c > 0$ such that if $\frac{n}{m} < c$, then (i) the initializer x_0 returned by Alg. 2 satisfies $\text{dist}(x_0, X_\star) \leq \frac{\nu}{8\sigma^2} \|x_\star\|_2$ and (ii) assuming no error in the minimization steps of the prox-linear method (Alg. 1),*

$$\text{dist}(x_k, X_\star) \leq \|x_\star\|_2 2^{-2^k}$$

for all $k \in \mathbb{N}$ with probability at least $1 - e^{-cm}$.

We make two brief summarizing remarks. First, it is necessary to have $m \gtrsim n$ to achieve exact recovery of the signal, as the parameter $x_\star \in \mathbb{C}^n$ has $2n$ unknowns and we have m equations (indeed, $m \geq 4n - 2$ is necessary for injectivity of the measurements in the complex case [2]). Thus, the sample complexity Theorem 2 specifies is optimal to within numerical constants. Second, Theorem 2 shows that the prox-linear algorithm exhibits local quadratic convergence to the signal x_\star , which is in contrast to the local linear convergence of other non-convex methods based on gradients and generalized gradients [13, 15, 46]. In contrast, however, each iteration of our algorithm requires solving a structured convex quadratic program, which is somewhat more expensive than the typical gradient iterations; as we demonstrate in our experiments (Section 6), this means our methods are about four times slower in overall run time than the best gradient-based methods, though their recovery properties are better.

4 Phase retrieval with outliers

The objective (2) is the analogue of the least-absolute deviation estimator—the median in \mathbb{R} —so in analogy with our understanding of robustness [27], it is natural to expect it should be robust to outliers. We show that this is indeed the case, and the prox-linear method we develop is effective. For simplicity in our development, we assume for this section that all measurements and signals are real-valued, and we consider the following corruption model: let $\{\xi_i\} \subset \mathbb{R}$ be an arbitrary sequence, and given the m measurement vectors a_i , we observe

$$b_i = \begin{cases} \langle a_i, x_\star \rangle^2 & \text{if } i \in \mathcal{I}^{\text{in}} \\ \xi_i & \text{if } i \in \mathcal{I}^{\text{out}}, \end{cases}$$

where $\mathcal{I}^{\text{out}} \subset [m]$ and $\mathcal{I}^{\text{in}} \subset [m]$ denote the outliers and inliers, respectively. We assume there is a pre-specified measurement failure probability $p_{\text{fail}} \in [0, \frac{1}{2})$, and $|\mathcal{I}^{\text{out}}| = p_{\text{fail}} m$, and the indices $i \in \mathcal{I}^{\text{out}}$ are chosen randomly. That is, measurement failures are random, though the noise sequence ξ_i may depend on a_i (even adversarially), as we specify presently. We assume no prior knowledge of which indices $i \in [m]$ actually satisfy $i \in \mathcal{I}^{\text{out}}$, or even of p_{fail} .

We consider the following two models for errors:

Model M1. *The measurement vectors $\{a_i\}_{i=1}^m$ are independent of all the values $\{\xi_i\}_{i=1}^m$.*

Model M2. *The inlying measurement vectors $\{a_i\}_{i \in \mathcal{I}^{\text{in}}}$ are independent of the values $\{\xi_i\}_{i \in \mathcal{I}^{\text{out}}}$ of the corrupted observations.*

Model M1 requires independence between the noise and measurements: the adversary may only corrupt ξ_i without observing a_i . Model M2 relaxes this, allowing arbitrary dependence between the corrupted data and the measurement vectors a_i for $i \in \mathcal{I}^{\text{out}}$. This is natural in scenarios in which the corruption may depend on the measurement a_i .

The arbitrary corruption causes some technical challenges, but we may still follow the outline in our analysis of phase retrieval without noise in Sec. 3. As we show in Section 4.1, the objective $f(x)$ is still stable (Condition C1) as long as the measurement vectors are light-tailed, though Gaussianity is unnecessary. The quadratic approximation conditions (Condition C2) are completely identical to those in Sec. 3.2, so we ignore them. Thus, as long as $\|A\|_{\text{op}}$ is not too large (meaning $f_x(y) \approx f(y)$) and we can find a good initializer, the prox-linear iterations (5) will converge quadratically to x_\star . Finding a good initializer x_0 is somewhat trickier, but in Section 4.2 we provide a spectral method, inspired by Wang et al. [46], that works with high probability as soon as $m/n \geq C$ for some numerical constant C . We defer our arguments to Appendix B.

4.1 Stability

The outlying indices, even when corruptions are chosen adversarially, have limited effect on the growth and identification behavior of $f(x) = \frac{1}{m} \|(Ax)^2 - b\|_1$. In particular, for p_{fail} smaller than a numerical constant, which we can often specify, the stability condition C1 holds with high probability whenever m/n is large. More precisely, we have the following proposition, which applies to independent σ^2 -sub-Gaussian measurements. (See Sec. B.1 for a proof.)

Proposition 4. *Let Assumption A4 hold and $\kappa_{\text{st}} = \inf_{u,v \in \mathbb{S}^{n-1}} \mathbb{E}[|\langle a, u \rangle \langle a, v \rangle|]$. Then under either of the models M1 or M2, there are numerical constants $c > 0$ and $C < \infty$ such that*

$$f(x) - f(x_\star) \geq \left(\kappa_{\text{st}} - 2p_{\text{fail}} - C\sigma^2 \sqrt[3]{\frac{n}{m}} - C\sigma^2 t \right) \|x - x_\star\|_2 \|x + x_\star\|_2 \quad \text{for all } x \in \mathbb{R}^n$$

with probability at least $1 - 2e^{-cm} - 2e^{-mt^2}$.

We continue our running example of Gaussian random variables to motivate the proposition.

Example 4 (Gaussian vectors): We claim that for $a \sim \mathcal{N}(0, I_n)$ we have

$$\kappa_{\text{st}} := \inf_{u,v \in \mathbb{S}^{n-1}} \mathbb{E}[|\langle a, u \rangle \langle a, v \rangle|] = \frac{2}{\pi}.$$

Let $Z_u = \langle a, u \rangle$, $Z_v = \langle a, v \rangle$, and let X, Y be independent $\mathcal{N}(0, 1)$. Then

$$\inf_{u,v \in \mathbb{S}^{n-1}} \mathbb{E}[|Z_u Z_v|] = \inf_{\rho \in [0,1]} \left\{ f(\rho) := \mathbb{E}[|\rho X^2 - (1-\rho)Y^2|] \right\}.$$

The function $f(\cdot)$ is convex and symmetric around $\frac{1}{2}$. Thus $f(\rho) \geq f(1/2) = 2/\pi$. \diamond

In the Gaussian measurement case, whenever $p_{\text{fail}} < \frac{1}{\pi} \approx .318$, there is a numerical constant $\lambda > 0$ such that we have the stability $f(x) - f(x_\star) \geq \lambda \|x - x_\star\| \|x + x_\star\|$ for as long as m/n is larger than some numerical constant.

4.2 Initialization

The last ingredient for achieving strong convergence guarantees for the prox-linear algorithm for phase retrieval is to provide a good initialization $x_0 \approx x_\star$. The strategies in the noiseless setting

Data: Measurement matrix $A \in \mathbb{R}^{m \times n}$ and corrupted signals b

begin

Define indices $\mathcal{I}_{\text{sel}} := \{i \in [m] : b_i \leq \text{med}(\{b_i\}_{i=1}^m)\}$.

Construct directional and norm estimates \hat{d} and \hat{r} by

$$X^{\text{init}} := \frac{1}{m} \sum_{i=1}^m a_i a_i^T \mathbf{1}_{\{i \in \mathcal{I}_{\text{sel}}\}}, \quad \hat{d} = \underset{d \in \mathbb{S}^{n-1}}{\text{argmin}} d^T X^{\text{init}} d$$

$$\hat{r}^2 := \underset{r}{\text{argmin}} G(r) := \frac{1}{m} \sum_{i=1}^m |b_i - r \langle a_i, \hat{d} \rangle|^2.$$

return (\hat{r}, \hat{d})

end

Algorithm 3: Initialization procedure with outliers

in Section 3 will fail because of corruptions. With this in mind, we present Algorithm 3, which provides an initializer in corrupted problems.

Before turning to the analysis, we provide some intuition for the algorithm. We must construct two estimates: an estimate \hat{d} of the direction $d_\star = x_\star / \|x_\star\|_2$ and an estimate \hat{r} of the radius, or magnitude, of the signal $r_\star = \|x_\star\|_2$. For the former, a variant of the spectral initialization (Alg. 2) suffices. If we take the $\mathcal{I}_{\text{sel}} \subset [m]$ to be the set of indices \mathcal{I}_{sel} corresponding to the smallest (say) b_i , in either model M1 or M2 the indices $i \in \mathcal{I}^{\text{out}}$ are independent of the measurements a_i , so we expect as in Section 3.3 that $X^{\text{init}} = |\mathcal{I}_{\text{sel}}|^{-1} \sum_{i \in \mathcal{I}_{\text{sel}}} a_i a_i^T = z I_n - z' d_\star d_\star^T + \Delta$, where z, z' are random positive constants and Δ is an error matrix coming from both randomness in the a_i and the corruptions. As long as the error Δ is small, the minimum eigenvector of X^{init} should be approximately d_\star . Once we have a good initializer $\hat{d} \approx d_\star$, a natural idea to estimate r_\star is to pretend that \hat{d} is the direction of the signal, substitute the variable $x = \sqrt{r} \hat{d}$ into the objective (2), and solve for r to get a robust estimate of the signal strength $\|x_\star\|$. As we show presently, this procedure succeeds with high probability (and the estimate \hat{r} is good even when the data are non-Gaussian).

Let us make these ideas precise. First, we show that our estimate \hat{r} of $\|x_\star\|_2$ is accurate (See Sec. B.2 for a proof).

Proposition 5. *Let Assumption A4 hold and $\mathbb{E}[aa^T] = I_n$. Let $\delta \in [0, 1]$ and $p_{\text{fail}} \in [0, \frac{1}{2}]$. There exist numerical constants $0 < c, C < \infty$ such that if \hat{d} is an estimate of d_\star for which*

$$\delta := \frac{C\sigma^2}{1 - 2p_{\text{fail}}} \text{dist}(\hat{d}, \{\pm d_\star\}) \leq 1, \quad (16)$$

then with probability at least $1 - 2e^{-cm(1-2p_{\text{fail}})^2/\sigma^4}$ all minimizers \hat{r}^2 of $G(r) = \frac{1}{m} \sum_{i=1}^m |b_i - r \langle a_i, \hat{d} \rangle|^2$, defined in Alg. 3, satisfy $\hat{r}^2 \in [1 \pm \delta] \|x_\star\|_2^2$.

Given Proposition 5, finding a good initialization of x_\star reduces to finding a good estimate \hat{d} of the direction $d_\star = x_\star / \|x_\star\|_2$. To make this precise, let $r_\star := \|x_\star\|$ and assume that $\delta = \frac{C\sigma^2}{1-2p_{\text{fail}}} \text{dist}(\hat{d}, \{\pm d_\star\}) \leq 1$ as in Proposition 5; assume also the relative error guarantee $|\hat{r} - r_\star| \leq \delta r_\star$. Using the triangle inequality and Proposition 5, for $x_0 = \hat{r} \hat{d}$ we have

$$\text{dist}(x_0, X_\star) \leq \hat{r} \text{dist}(\hat{d}, \{\pm d_\star\}) + |r_\star - \hat{r}| \leq r_\star \left[(1 + \delta) \text{dist}(\hat{d}, \{\pm d_\star\}) + \delta \right] \leq 2\delta r_\star = \frac{C\sigma^2}{1 - 2p_{\text{fail}}} \text{dist}(\hat{d}, \{\pm d_\star\}) r_\star,$$

as claimed. We turn to the directional estimate; to make the analysis cleaner we make the normality

Assumption A6. The measurement vectors $a_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, I_n)$.

To state our guarantee on \hat{d} , we require additional notation for quantiles of Gaussian and χ^2 -random variables. Let $W \sim \mathcal{N}(0, 1)$ and define the constant q_{fail} and its associated χ^2 -quantile w_q^2 by

$$\mathbb{P}(W^2 \leq w_q^2) = q_{\text{fail}} := \frac{1}{2(1 - p_{\text{fail}})} + \frac{1 - 2p_{\text{fail}}}{4(1 - p_{\text{fail}})} = \frac{3 - 2p_{\text{fail}}}{4(1 - p_{\text{fail}})} < 1. \quad (17)$$

Second, define the constant $\delta_q = 1 - \mathbb{E}[W^2 \mid W^2 \leq w_q^2]$. We have the following guarantee.

Proposition 6. Let \hat{d} be the smallest eigenvector of $X^{\text{init}} := \frac{1}{m} \sum_{i=1}^m a_i a_i^T \mathbf{1}\{i \in \mathcal{I}_{\text{sel}}\}$, as in Alg. 3. Let the constant $M = 0$ if Model M1 holds and $M = 1$ if Model M2 holds. There are numerical constants $0 < c, C < \infty$ such that for $t \geq 0$, with probability at least

$$1 - \exp(-mt) - \exp(-c(1 - 2p_{\text{fail}})m) - \exp\left(-c \frac{m\delta_q^2}{w_q^2}\right)$$

we have

$$\text{dist}(\hat{d}, \{\pm d_\star\}) \leq \frac{C\sqrt{\frac{n}{m} + t}}{\left[(1 - 2p_{\text{fail}})\delta_q - C\sqrt{\frac{n}{m} + t} - Mp_{\text{fail}}\right]_+}.$$

For intuition, we provide a few simplifications of Proposition 6 by bounding the quantities w_q and δ_q defined in Eq. (17). Using the conditional expectation bound in Lemma 3.2 and a more careful calculation for Gaussian random variables (see Lemma C.1 in the appendices) we have

$$\mathbb{E}[a_{i,1}^2 \mid a_{i,1}^2 \leq w_q^2] \leq \min\left\{\frac{w_q^2}{3}, 1 - \frac{1}{2}w_q^2\mathbb{P}(a_{i,1}^2 \geq w_q^2)\right\} < 1,$$

so $\delta_q \geq \max\{1 - \frac{w_q^2}{3}, \frac{1}{2}w_q^2\mathbb{P}(W^2 \geq w_q^2)\}$. For $p_{\text{fail}} \leq \frac{3}{8}$, we may take $w_q^2 \leq 2.71$ and $\delta_q > \frac{1}{11}$. More generally, a standard Gaussian calculation that $\Phi^{-1}(p) \geq \sqrt{|\log(1 - p)|}$ as $p \rightarrow 1$ shows that $w_q^2 \geq \log \frac{8(1 - p_{\text{fail}})}{1 - 2p_{\text{fail}}}$ as $p_{\text{fail}} \rightarrow \frac{1}{2}$, so $\delta_q \geq \frac{1 - 2p_{\text{fail}}}{8(1 - p_{\text{fail}})} \log \frac{8(1 - p_{\text{fail}})}{1 - 2p_{\text{fail}}}$. Under Model M1, then, as long as the sample is large enough and $p_{\text{fail}} < \frac{1}{2}$, we can achieve constant accuracy in the directional estimate $\text{dist}(\hat{d}, \{\pm d_\star\})$ with probability of failure e^{-cm} . Under the more adversarial noise model M2, we require a bit more; more precisely, we must have $(1 - 2p_{\text{fail}})\delta_q - p_{\text{fail}} > 0$ to achieve accurate estimates. A numerical calculation shows that if $p_{\text{fail}} < \frac{1}{4}$, then this condition holds, so that under Model M2 we can achieve constant accuracy in the directional estimate $\text{dist}(\hat{d}, \{\pm d_\star\})$ with high probability.

4.3 Summary and success guarantees

With the guarantees of stability, quadratic approximation, and good initialization for suitably random matrices $A \in \mathbb{R}^{m \times n}$, we can provide a theorem showing that the prox-linear approach to the composite optimization phase retrieval objective succeeds with high probability. Roughly, once m/n is larger than a numerical constant, the prox-linear method with noisy initialization succeeds with exponentially high probability, even with outliers. Combining the convergence Theorem 1 with Propositions 4, 5, 6, and Corollary 3.2, we have the following theorem.

Theorem 3. Let Assumptions A6 hold. There exist numerical constants $c > 0$ and $C < \infty$ such that the following hold for any $t \geq 0$. Let the independent outliers Model M1 hold and $p_{\text{fail}} < \frac{1}{\pi}$ or

the adversarial outliers Model M2 hold and $p_{\text{fail}} < \frac{1}{4}$. Let x_0 be the initializer returned by Alg. 3, and assume the iterates x_k of Alg. 1 are generated without error. Then

$$\text{dist}(x_0, X_\star) \leq \frac{C}{(1 - 2p_{\text{fail}})^2} \sqrt{\frac{n}{m} + t \cdot \|x_\star\|_2} \quad \text{and} \quad \text{dist}(x_k, X_\star) \leq \|x_\star\|_2 2^{-2^k} \quad \text{for all } k \in \mathbb{N}$$

with probability at least $1 - 4e^{-mt} - e^{-c(1-2p_{\text{fail}})^2 m}$.

Thus, we see that the method succeeds with high probability as long as the sample size is large enough, though there is non-trivial degradation when p_{fail} is large.

5 Optimization methods

In practice, we require some care to solve the sub-problems (5), to minimize $f_x(y) + \frac{L}{2} \|x - y\|_2^2$, at large scale. In this section, we describe the three schemes we use to solve the sub-problems (one is simply using the industrial **Mosek** solver). We evaluate these more carefully in the experimental section to come.

To fix notation, let $\phi(\cdot)$ be the elementwise square operator. Recalling that $f(x) = \frac{1}{m} \|\phi(Ax) - b\|_1$, we perform a few simplifications to more easily describe the schemes we use to solve the prox-linear sub-problems. Assuming we begin an iteration at point x_0 , defining the diagonal matrix $D = 2 \text{diag}(Ax_0) \in \mathbb{R}^{m \times m}$ and setting $c = b + \phi(Ax_0)$, we recall inequality (8) and have

$$f(y) \leq f_{x_0}(y) + \frac{1}{m} (y - x_0)^T A^T A (y - x_0) \leq \frac{1}{m} \|DAy - c\|_1 + \|m^{-1} A^T A\|_{\text{op}} \|y - x_0\|_2^2.$$

Rewriting this with appropriately rescaled diagonal matrices D and vector c , implementing Algorithm 1 becomes equivalent to solving a sequence of optimization problems of the form

$$\underset{x}{\text{minimize}} \quad \|DAx - c\|_1 + \frac{1}{2} \|x\|_2^2. \quad (18)$$

In small scale scenarios, the problem (18) is straightforward to solve via standard interior point method software; we use **Mosek** via the **Convex.jl** toolbox [42]. We do not describe this further. In larger-scale scenarios, we use more specialized methods, which we now describe.

5.1 Graph splitting methods for the prox-linear sub-problem

When the matrices are large, we use a variant of the Alternating Directions Method of Multipliers (ADMM) procedure known as the proximal operator graph splitting (POGS) method of Parikh and Boyd [34, 35], which minimizes objectives of the form $f(x) + g(y)$ subject to a linear constraint $Bx = y$. We experimented with a number of specialized first-order and interior-point methods for solving problem (18), but in our experience, POGS offers the empirically best performance. Let us describe the POGS method. Let the matrix $B = DA$ for shorthand; evidently, problem (18) has precisely this form and is equivalent to

$$\underset{x \in \mathbb{R}^n, y \in \mathbb{R}^m}{\text{minimize}} \quad \|y - c\|_1 + \frac{1}{2} \|x\|_2^2 \quad \text{subject to } Bx = y.$$

The POGS method iterates to solve problem (18) as follows. Introduce dual variables $\lambda_k \in \mathbb{R}^n$ and $\nu_k \in \mathbb{R}^m$ associated to x and y , and consider the iterations

$$\begin{aligned}
x_{k+\frac{1}{2}} &= \underset{x}{\operatorname{argmin}} \left\{ \frac{1}{2} \|x\|_2^2 + \frac{\rho}{2} \|x - (x_k - \lambda_k)\|_2^2 \right\} \\
y_{k+\frac{1}{2}} &= \underset{y}{\operatorname{argmin}} \left\{ \|y - c\|_1 + \frac{\rho}{2} \|y - (y_k - \nu_k)\|_2^2 \right\} \\
\begin{bmatrix} x_{k+1} \\ y_{k+1} \end{bmatrix} &= \begin{bmatrix} I_n & B^T \\ B & -I_m \end{bmatrix}^{-1} \begin{bmatrix} I_n & B^T \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x_{k+\frac{1}{2}} + \lambda_k \\ y_{k+\frac{1}{2}} + \nu_k \end{bmatrix} \\
\lambda_{k+1} &= \lambda_k + (x_{k+\frac{1}{2}} - x_{k+1}), \quad \nu_{k+1} = \nu_k + (y_{k+\frac{1}{2}} - y_{k+1}).
\end{aligned} \tag{19}$$

Each of the steps of the method (19) is trivial except for the matrix inversion, or the “graph projection” step, which projects the pair $(x_{k+\frac{1}{2}} + \lambda_k, y_{k+\frac{1}{2}} + \nu_k)$ to the set $\{x, y : Bx = y\}$. The first two updates amount to

$$x_{k+\frac{1}{2}} = \frac{\rho}{1+\rho}(x_k - \lambda_k) \quad \text{and} \quad y_{k+\frac{1}{2}} = c + \operatorname{sgn}(y_k - \nu_k - c) \odot [|y_k - \nu_k - c| - 1/\rho]_+$$

where \odot denotes elementwise multiplication and each operation is element-wise. The matrix B is tall, so setting $v_k = x_{k+\frac{1}{2}} + \lambda_k + B^T(y_{k+\frac{1}{2}} + \nu_k) \in \mathbb{R}^n$, then the solution of the system

$$\begin{bmatrix} I_n & B^T \\ B & -I_n \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} v_k \\ 0_n \end{bmatrix} \quad \text{is} \quad x_{k+1} = (I_n + B^T B)^{-1} v_k \quad \text{and} \quad y_{k+1} = Bx_{k+1}. \tag{20}$$

In this iteration, it is straightforward to cache the matrix $(I_n + B^T B)^{-1}$, or a Cholesky factorization of the matrix, so that we can repeatedly compute the multiplication (20) in time $n^2 + nm$.

Following Parikh and Boyd [35], we define the primal and dual residuals

$$r_{k+1}^{\text{pri}} := \begin{bmatrix} x_{k+1} - x_{k+\frac{1}{2}} \\ y_{k+1} - y_{k+\frac{1}{2}} \end{bmatrix} \quad \text{and} \quad r_{k+1}^{\text{dual}} := \rho \begin{bmatrix} x_k - x_{k+1} \\ y_k - y_{k+1} \end{bmatrix}.$$

These residuals define a natural stopping criterion [35], where one terminates the iteration (19) once the residuals satisfy

$$\|r_k^{\text{pri}}\|_2 < \epsilon (\sqrt{n} + \max\{\|x_k\|_2, \|y_k\|_2\}) \quad \text{and} \quad \|r_k^{\text{dual}}\|_2 < \epsilon (\sqrt{n} + \max\{\|\lambda_k\|_2, \|\nu_k\|_2\}) \tag{21}$$

for some $\epsilon > 0$, which must be specified. In our case, the quadratic convergence guarantees of Theorem 1 suggest a strategy of decreasing ϵ across iterative solutions of the sub-problem (18). That is, we begin with some $\epsilon = \epsilon_0$. We perform iterations of the prox-linear Algorithm 1 using the POGS iteration (19) (until the residual criterion (21) holds) to solve the inner problem (18). Periodically, in the outer prox-linear iterations of Alg. 1, we decrease ϵ by some large multiple.

In our experiments with the POGS method for sub-problem solutions, we perform two phases. In the first phase, we iterate the prox-linear method (Alg. 1) until either $k = 25$ or $\|x_k - x_{k+1}\|_2 \leq \delta / \|(1/m)A^T A\|_{\text{op}}$, where $\delta = 10^{-3}$, using accuracy parameter $\epsilon = 10^{-5}$ for POGS (usually, the iterations terminate well-before $k = 25$). We then decrease this parameter to $\epsilon = 10^{-8}$, and begin the iterations again.

5.2 Conjugate gradient methods for sub-problems

In a number of scenarios, it is possible to multiply by the matrix A quickly, though the direct computation $(I_n + A^T D^2 A)^{-1}$ in expression (20) (recall $B = DA$) may be difficult. For example, if A is structured (say, a Fourier or Hadamard transform matrix or sparse), computing multiplication by $I_n + A^T D^2 A$ quickly is possible. This suggests [40, Part VI] using conjugate gradient methods.

We make this more explicit here to mesh with our experiments to come. Let H_n be an $n \times n$ orthogonal matrix for which computing the multiplication $H_n v$ is efficient (e.g. a Hadamard or discrete cosine transform matrix). We assume that the measurement matrix A takes the form of repeated randomized measurements under H_n , that is,

$$A = [H_n S_1 \quad H_n S_2 \quad \cdots \quad H_n S_k]^T \quad (22)$$

where $S_l \in \mathbb{R}^{n \times n}$ are (random) diagonal matrices, so that $A \in \mathbb{R}^{m \times n}$ with $m = kn$. In this case, the expensive part of the system (20) is the solution of $(I + A^T D^2 A)x = v$. This is a positive definite system, and it is possible to compute the multiplication $(I + A^T D^2 A)x$ in time $O(kn + kT_{\text{mult}})$, where T_{mult} is the time to multiply an n -vector by the matrix H_n and H_n^T . When S is a random sign matrix and H_n is a Hadamard or FFT matrix, the matrix $A^T D^2 A$ is well-conditioned, as the analysis of random (subsampled) Hadamard and Fourier transforms shows [41]. Thus, in our experiments with structured matrices we use the conjugate gradient method *without* preconditioning [40], iterating until we have relative error $\|(I + A^T D^2 A)x - v\| \leq \epsilon_{\text{cg}} / \|v\|$ where $\epsilon_{\text{cg}} = 10^{-6}$.

6 Experiments

We perform a number of simulations as well as experiments on real images to evaluate our method and compare with other state-of-the-art (non-convex) methods for real-valued phase retrieval problems. To standardize notation and remind the reader, in each experiment, we generate data via (variants) of the following process. We take a measurement matrix $A \in \mathbb{R}^{m \times n}$, from one of a few distributions and generate a signal $x_\star \in \mathbb{R}^n$ either by drawing $x_\star \sim \mathcal{N}(0, I_n)$ or by taking $x_\star \in \{-1, 1\}^n$ uniformly at random. We receive observations of the form

$$b_i = \langle a_i, x_\star \rangle^2,$$

and with probability $p_{\text{fail}} \in [0, \frac{1}{2}]$, we corrupt the measurements b_i .

In our experiments, to more carefully isolate the relative performance of the iterative algorithms, rather than initialization used, we compare three initializations. The first two rely on Gaussianity of the measurement matrix A .

- (i) **Big:** The initialization of Wang et al. [46]. Defining

$$\mathcal{I}_0 := \left\{ i \in [m] : b_i / \|a_i\|_2^2 \geq \text{quant}_{5/6}(\{b_i / \|a_i\|_2^2\}) \right\} \quad \text{and} \quad X^{\text{init}} := \sum_{i \in \mathcal{I}_0} \frac{1}{\|a_i\|_2^2} a_i a_i^T,$$

we set the direction $\hat{d} = \arg\max_{\|d\|_2=1} d^T X^{\text{init}} d$ and $x_0 = (\frac{1}{m} \sum_{i=1}^m b_i)^{\frac{1}{2}} \hat{d}$.

- (ii) **Median:** The initialization of Zhang et al. [47]. We set $\hat{r}^2 = \text{quant}_{\frac{1}{2}}(\{b_i\}) / .455$ and define

$$\mathcal{I}_0 := \{i \in [m] : |b_i| \leq 9\lambda_0\} \quad \text{and} \quad X^{\text{init}} := \sum_{i \in \mathcal{I}_0} b_i a_i a_i^T.$$

We then set the direction $\hat{d} = \arg\max_{\|d\|_2=1} d^T X^{\text{init}} d$ and $x_0 = \hat{r} \hat{d}$.

(iii) **Small:** The outlier-aware initialization we describe in Sec. 4.2.

Our convergence results in Section 2 rely on four quantities: the quality of the initialization, $\text{dist}(x_0, X_\star)$; the stability parameter λ such that $f(x) - f(x_\star) \geq \lambda \|x - x_\star\|_2 \|x + x_\star\|_2$; the quadratic upper bound guaranteed by our linearized models $f_x(y)$, so that $|f_x(y) - f(y)| \leq \frac{L}{2} \|x - y\|_2^2$; and the accuracy to which we solve the optimization problems. The random matrix A governs the middle two quantities—stability λ and closeness L —and we take $L = \frac{2}{m} \|A\|_{\text{op}}^2$. Thus, in our experiments we directly vary the initialization scheme to generate x_0 and the accuracy to which we solve the sub-problems (5), that is,

$$x_{k+1} = \underset{x}{\text{argmin}} \left\{ \frac{1}{m} \sum_{i=1}^m |\langle a_i, x_k \rangle^2 + 2\langle a_i, x_k \rangle \langle a_i, x - x_k \rangle - b_i| + \frac{L}{2} \|x - x_k\|_2^2 \right\}.$$

We perform each of our experiments on a server with a 16 core 2.6 GhZ Intel Xeon processor with 128 GB of RAM using `julia` as our language, with `OpenBLAS` as the BLAS library. We restrict each optimization method to use 4 of the cores (`OpenBLAS` is multi-threaded).

6.1 Simulations with zero noise and Gaussian matrices

For our first collection of experiments, we evaluate the performance of our method for recovering random signals $x_\star \in \mathbb{R}^n$ using Gaussian measurement matrices $A \in \mathbb{R}^{m \times n}$, varying the number of measurements m over the ten values $m \in \{1.8n, 1.9n, \dots, 2.7n\}$. (Taking $m \geq 2.7n$ yielded 100% exact recovery in all the methods we experiment with.) We assume a noiseless measurement model, so that for $A = [a_1 \ \dots \ a_m]^T \in \mathbb{R}^{m \times n}$, we have $b_i = \langle a_i, x_\star \rangle^2$ for each $i \in [m]$. We perform experiments with dimensions $n \in \{400, 600, 800, 1000, 1500, 2000, 3000\}$, and within each experiment we vary a number of problem parameters, including initialization scheme and the algorithm we use to solve the sub-problems (5).

To serve as our baseline for comparison, we use Truncated Amplitude Flow (TAF) [46], given that it outperforms other non-convex iterative methods for phase retrieval, including Wirtinger Flow [13], Truncated Wirtinger Flow [15], and Amplitude Flow [46]. Setting $\psi_i = \sqrt{b_i}$, TAF tries to minimize the loss $\frac{1}{2m} \sum_{i=1}^m (\psi_i - |\langle a_i, x \rangle|)^2$ via a carefully designed (generalized) gradient method. For convenience, we replicate the method (as described by Wang et al. [46]) in Alg. 4.

Data: Initializer $x_0 \in \mathbb{R}^n$, i.i.d. standard Gaussian matrix $A \in \mathbb{R}^{m \times n}$, $\psi_i = |\langle a_i, x_\star \rangle|$, and parameters $\gamma = .7$ and $\alpha = .6$

for $k = 0, 1, \dots, K - 1$ **do**

 Set $\mathcal{I}_k = \{i \in [m] : |\langle a_i, x_k \rangle| \geq \frac{1}{1+\gamma} \psi_i\}$
 Update $x_{k+1} = x_k - \frac{\alpha}{m} \sum_{i \in \mathcal{I}_k} \left(\langle a_i, x_k \rangle - \psi_i \frac{\langle a_i, x_k \rangle}{|\langle a_i, x_k \rangle|} \right) a_i$

end

return x_K

Algorithm 4: Truncated amplitude flow

6.1.1 Low-dimensional experiments with accurate prox-linear steps

We begin by describing our experiments with dimensions $n \in \{400, 600, 800\}$. For each of these, we solve the iterative sub-problems to machine precision, using `Mosek` and the Julia package `Convex.jl` [42]. We plot representative results for $n = 400$ and $n = 800$ in Figure 1, which summarize 400 independent experiments, and we generate the true x_\star by taking $x_\star \in \{-1, 1\}^n$




Figure 1. Zero-noise experiments in dimensions $n = 400$ and $n = 800$. (a) Fraction of times (in 400 experiments) that one method succeeded while other failed for $n = 400$. (b) Fraction of successful recoveries for $n = 400$. (c) Fraction of successful recoveries for $n = 800$.

uniformly at random. (In separate experiments, we drew $x_\star \sim \mathcal{N}(0, I_n)$, and the results were essentially identical.) In these figures, we use the “big” initialization (i) (the initialization of Wang et al., as it yields the best empirical performance. Following Wang et al. [46], we perform 1000 iterations of TAF (Alg. 4), and within each experiment, both TAF and the prox-linear method use identical initializer and data matrix A . We run the prox-linear method until sequential updates x_k and x_{k+1} satisfy $\|x_k - x_{k+1}\|_2 \leq \epsilon = 10^{-5}$, which in *every* successful experiment we perform (with these data settings) requires 6 or fewer iterations. We declare an experiment successful if the output \hat{x} of the algorithm satisfies

$$\text{dist}(\hat{x}, X_\star) = \min \{ \|\hat{x} - x_\star\|_2, \|\hat{x} - x_\star\|_2 \} \leq \epsilon_{\text{acc}} \|x_\star\|_2 \quad \text{where} \quad \epsilon_{\text{acc}} = 10^{-5}. \quad (23)$$

In Fig. 1(a), we plot the number of times that one method succeeds (out of the 400 experiments) while the other does not as a function of the ratio m/n . We see that for these relatively small dimensional problems, the prox-linear method has mildly better performance for m/n small than does truncated amplitude flow. In Fig. 1(b), we plot the fraction of successful runs of the algorithms, again against m/n for $n = 400$, and in Fig. 1(c) we plot the fraction of successful runs for $n = 800$. Even when $m/n = 2$, the prox-linear method has success rate of around .6.

6.1.2 Medium dimensional experiments with inaccurate prox-linear steps

We now shift to a description of our experiments in dimensions $n \in \{1000, 1500, 2000, 3000\}$, again without noise in the measurements; we perform between 100 and 400 experiments for each dimension in this regime, and we use the “big” initialization (i) [46]. In this case, we use Parikh and Boyd’s proximal operator graph splitting (POGS) method [34] for the prox-linear steps, as we describe in Section 5.1. We perform two phases of the prox-linear method: the first performing POGS until the residual errors (21) are less than $\epsilon = 10^{-5}$ within the prox-linear steps, the second to accuracy $\epsilon = 10^{-8}$. We apply the prox-linear method to the matrix A/\sqrt{m} with data b/m , which is equivalent but numerically more stable because A/\sqrt{m} and I_n are comparable (see the recommendations [34]). In these accuracy regimes, the time for solution of the prox-linear method and that required for 1000 iterations of truncated amplitude flow are comparable; TAF requires about 1.5 seconds while the two-phase prox-linear method requires around 4 seconds in dimension $n = 1000$, and TAF requires about 5 seconds while the two-phase prox-linear method requires around 20 seconds in dimension $n = 2000$, each with $m = 2n$.




Figure 2. Zero-noise experiments. (a) Fraction of trials (of 400) in which one method succeeds while other fails for $n = 1000$. (b) Fraction of trials (of 100) in which one method succeeds while other fails for $n = 3000$. (c) Fraction of successful recoveries for $n = 3000$.

We provide two sets of plots for these results, where again we measure success by the criterion (23), $\|x \pm x_\star\|_2 \leq \epsilon_{\text{acc}} = 10^{-5}$. In the first, Fig. 2, we show performance of prox-linear against TAF specifically for dimensions $n = 1000$ and 3000 . In Fig. 2(a), we plot the number of trials in which one method succeeds (out of 400 experiments) while the other does not as a function of the ratio m/n . In Fig. 2(b) we plot the number of trials in which the prox-linear or TAF method succeeds, while the other does not, for $n = 3000$ with x_\star chosen either $\mathcal{N}(0, I_n)$ or uniform in $\{\pm 1\}^n$ (∇ and $+$ markers, respectively). We ignore ratios $m/n \geq 2.2$ as both methods succeed in all of our trials. Out of 100 trials, there is only *one* (with $m/n = 2$) in which TAF succeeds but the prox-linear method does not. In Fig. 2(c), we plot the fraction of successful runs of the algorithms, again against m/n for $n = 3000$. For these larger problems, there is a substantial gap in recovery probability between prox-linear method and TAF, where with $m/n = 2$ the prox-linear method achieves recovery more than 78% (± 4), 88% (± 6), and 91% (± 6) of the time, with 95% confidence intervals, for $n = 1000, 2000$, and 3000 , respectively.

6.2 Phase retrieval with outlying measurements

One of the advantages we claim for the objective (2) is that it is robust to outliers. Zhang et al. [47] develop a method, which they term median-truncated Wirtinger flow, for handling outlying measurements, which (roughly) sets the index set \mathcal{I}_k used for updates in Alg. 4 to be a set of measurements near the median values of the b_i . Their algorithm requires a number of parameters that strongly rely on the assumptions that $a_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, I_n)$; in contrast, the objective (2) and prox-linear method we investigate are straightforward to implement without any particular assumptions on A (and require no parameter tuning, relying on an explicit sequence of convex optimizations).

Nonetheless, to make comparisons between the algorithms as fair as possible, we implement their procedure and perform experiments in dimension $n \in \{100, 200\}$ with i.i.d. standard normal data matrices A . We perform 100 experiments as follows. Within each experiment, we evaluate each $m \in \{1.8n, 2n, 3n, 4n, 6n, 8n\}$ and failure probability $p_{\text{fail}} \in \{0, .01, .02, \dots, .29, .3\}$. For fixed m, n , we draw a data matrix $A \in \mathbb{R}^{m \times n}$, then choose $x_\star \in \mathbb{R}^n$ either by drawing $x_\star \sim \text{Uni}(\{-1, 1\}^n)$ or $x_\star \sim \mathcal{N}(0, I_n)$. We then generate $b_i = \langle a_i, x_\star \rangle^2$ for $i \in [m]$. For our experiments with $n = 100$, we simply set $b_i = 0$, which is more difficult for our initialization strategy, as it corrupts a large fraction of the vectors a_i used to initialize x_0 . For our experiments with $n = 200$, we draw b_i from a Cauchy distribution. Each problem setting (b_i Cauchy vs. zeroing and x_\star discrete or normal)




Figure 3. Probability of success for median-truncated Wirtinger flow (left) and the prox-linear method (right) for different initializations, where dimension $n = 100$. Horizontal axis indexes p_{fail} while vertical axis indexes the measurement ratio m/n . Each pixel represents the fraction of successful recoveries (defined as $\text{dist}(\hat{x}, X_*) / \|x_*\|_2 \leq 10^{-5}$) over 100 experiments. (a) & (b): “Big” initialization, with \hat{r} estimated by Alg. 3. (c) & (d): “Median” initialization. (e) & (f): “Small” initialization.

yields qualitatively similar results.

Figures 3, 4, and 5 summarize our results. In Figures 3 and 4, we display success rates of the median truncated Wirtinger flow (the left column in each figure) and our composite optimization-based procedure (right column) for a number of initializations; each plot represents results of 100 experiments. Within each plot, a white square indicates that 100% of trials were successful, meaning the signal is recovered to accuracy $\text{dist}(\hat{x}, X_*) \leq 10^{-5} \|x_*\|_2$, while black squares indicate 0% success rates. Within each row of the figure, we present results for the two methods using the same initialization scheme. Figure 3 gives results with $n = 100$, using precise (Mosek-based) solutions of the prox-linear updates, while Figure 4 gives results with $n = 200$ using the POGS-based updates (recall step (19)), with identical parameters as in the previous section. It is clear from the figures that the composite objective yields better recovery.

In Figure 5, we present a different view into the behavior of the prox-linear and MTWF methods. In the left two plots, we show the recovery probability (over 100 trials) for our composite optimization method (Fig. 5(a)) and MTWF (Fig. 5(c)). The success probability for the composite method is higher. In Fig. 5(b), we plot the average number of *iterations* (along with standard error bars) the prox-linear method performs when the dimension $n = 100$ and we use accurate sub-problem solves. We give iteration counts only for those trials that result in successful recovery; the iteration counts on unsuccessful trials are larger (indeed, if the method is not converging rapidly, this serves as a proxy for failure). In the high measurement regime, $m/n \geq 2.5$ or so, we




Figure 4. Probability of success for median-truncated Wirtinger flow (left) and the prox-linear method (right), using the approximate proximal graph splitting method for sub-problem solves. Dimension $n = 200$. (a) & (b): “Big” initialization, with \hat{r} estimated by Alg. 3. (c) & (d): “Small” initialization.

see that if $p_{\text{fail}} = 0$ no more than 7 iterations are required: this is the quadratic convergence of the method. (Indeed, for $m/n = 8$, for $p_{\text{fail}} \leq .15$ each execution of the prox-linear method uses precisely 5 iterations, never more, and never fewer.) In Fig. 5(d), we show the number of matrix multiplications by the inverse matrix $(I_n + A^T D A)^{-1}$ the method uses (recall the update (19) in the proximal graph operator splitting method). We see that for well-conditioned problems and those with little noise, the methods require relatively few matrix multiplications, while for more outliers and when m/n shrinks, there is a non-trivial increase.

6.3 Recovery of real images

Our final collection of experiments investigates recovery of real-world images using more specialized measurement matrices. In this case, we let $H_n \in \{-1, 1\}^{n \times n} / \sqrt{n}$ denote a normalized Hadamard matrix, where the multiplication $H_n v$ requires time $n \log n$, and H_n satisfies $H_n = H_n^T$ and $H_n^2 = I_n$. For some $k \in \mathbb{N}$, we then take k i.i.d. diagonal sign matrices $S_1, \dots, S_k \in \text{diag}(\{-1, 1\}^n)$, uniformly at random, and define $A = [H_n S_1 \ H_n S_2 \ \dots \ H_n S_k]^T \in \mathbb{R}^{kn \times n}$, as in expression (22). We note that this matrix explicitly *does not* satisfy the stability conditions that we use for our theoretical guarantees. Indeed, letting e_1 and e_2 be the first standard basis vectors, we have $H_n S(e_1 + e_2) \perp H_n S(e_1 - e_2)$ no matter the sign matrix S ; there are similarly pathological vectors for FFT, discrete cosine, and other structured matrices. Nonetheless, we perform experiments with this structured A matrix as follows.¹

Given an image X represented as a matrix, we define $x_\star = \text{Vec}(X)$, the vectorized representation of X . (In the case of colored images, where $X \in \mathbb{R}^{n_1 \times n_2 \times 3}$ because of the 3 RGB channels, we vectorize the channels as well.) We then set $b = \phi(Ax)$, where $\phi(\cdot)$ denotes elementwise squaring, and corrupt a fraction $p_{\text{fail}} \in [0, .2]$ of the measurements b_i by zeroing them. We then follow our standard experimental protocol, initializing x_0 by the “small” initialization scheme (iii), with the slight twist that now we use the proximal operator graph splitting method (POGS) with conjugate gradient methods (Sec. 5.2) to solve the graph projection step (20).

We first give results on a collection of 500 images of handwritten digits (using $k = 3$), available

¹We do not compare to other methods designed for outliers because we could not set the constants the methods require in such a way as to yield good performance.




Figure 5. Comparison of median-truncated Wirtinger flow and composite minimization with “Median” initialization. Horizontal axis of each plot indexes $p_{\text{fail}} \in [0, .3]$. (a) Proportion of successful solves for prox-linear method. (b) Number of iterations of prox-linear step (5) over *successful* solves (with error bars). (c) Proportion of successful solves for median-truncated Wirtinger flow method. (d) Number of matrix multiplications of the form $x \mapsto (I + A^T D^2 A)^{-1} A x$ for the prox-linear method with proximal graph solves.




Figure 6. Example recoveries of digits. Left digit is true digit, middle is initialization, right is recovered image.






Figure 7. Performance of composite optimization scheme using conjugate gradient method and proximal graph splitting to solve sub-problems (Sections 5.1–5.2).

on the website for the book [25]. We provide example results of the execution of our procedure in Fig. 6, which shows that while there is signal in the initialization, there is substantial work that the prox-linear method must perform to recover the images. For each of the 500 images, we vary $p_{\text{fail}} \in \{0, .025, .05, \dots, .175, .2\}$, then execute the prox-linear method. We plot summary results in Fig. 7, which in the blue curve with square markers gives the probability of successful recovery of the digit (left axis) versus p_{fail} (horizontal axis). The right axis indexes the number of matrix multiplications the method executes until completion (black line with circular marks). We see that in spite of the demonstrated failure of the matrix A to satisfy our stability assumptions, we have frequent recovery of the images to accuracy 10^{-3} or better.


Finally, we perform experiments with eight real color images with sizes up to 1024×1024 (yielding $n = 2^{22}$ -dimensional problems), where we use $k = 3$ random Hadamard sensing matrices. The prox-linear method successfully recovers each of the 8 images to relative accuracy at least 10^{-4} , and performs an average of 15100 matrix-vector multiplications (i.e. fast Hadamard transforms) over the eight experiments, with a standard deviation of 2600 multiplications. To give a sense of the importance of different parts of our procedure, and the relative accuracies to which we solve sub-problems (5), we display one example in Fig. 8. In this example, we perform phase retrieval on an image of RNA nanoparticles in cancer cells [36]. In Fig. 8(a), we show the result of initialization, which displays non-trivial structure, though is clearly noisy. In Fig. 8(b), we show the result of 10 steps of the prox-linear method, solving each step using POGS until the residual errors (21) are below $\epsilon = 10^{-3}$. There are clear artifacts in this image. We then perform one refinement step with higher accuracy ($\epsilon = 10^{-7}$), which results in Fig. 8(c). This is indistinguishable, at least to our eyes, from the original image (Fig. 8(d)).




(a)



(b)



(c)



(d)

Figure 8. Reconstruction of RNA image ($n = 2^{22}$) from $m = 3n$ measurements. (a) Initialization x_0 of prox-linear method. (b) Result of 10 inaccurate ($\epsilon = 10^{-3}$) solutions of prox-linear step (5). (c) Result of *one* additional accurate ($\epsilon = 10^{-7}$) solution of prox-linear step (5). (d) Original image.

Acknowledgments We thank Dima Drusvyatskiy and Courtney Paquette for a number of inspirational and motivating conversations. JCD and FR were partially supported by NSF-CAREER Award 1553086 and the Toyota Research Institute. FR was additionally supported by a Stanford Graduate Fellowship.

A Proofs for noiseless phase retrieval

In this appendix, we collect the proofs of the propositions and results in Section 3.

Because we use it multiple times in what follows, we state a standard eigenvector perturbation result, a variant of the Davis-Kahane sin- Θ theorem.

Lemma A.1 (Stewart and Sun [39], Theorem 3.6). *For vectors $u, v \in \mathbb{S}^{n-1}$, define the angle $\theta(u, v) = \cos^{-1} |\langle u, v \rangle|$. Let $X \in \mathbb{C}^{n \times n}$ be symmetric, Δ a symmetric perturbation, and $Z = X + \Delta$, and define $\text{gap}(X) = \lambda_1(X) - \lambda_2(X)$ to be the eigengap of X . Let v_1 and u_1 be the first eigenvectors of X and Z , respectively. Then*

$$\sqrt{1 - |\langle u_1, v_1 \rangle|^2} = |\sin \theta(u_1, v_1)| \leq \frac{\|\Delta\|_{\text{op}}}{[\text{gap}(X) - \|\Delta\|_{\text{op}}]_+}.$$

We will also have occasion to use the following one-sided variant of Bernstein's inequality.

Lemma A.2 (One-sided Bernstein inequality). *Let X_i be non-negative random variables with $\text{Var}(X_i) \leq \sigma^2$. Then*

$$\mathbb{P}\left(\frac{1}{m} \sum_{i=1}^m X_i \leq \frac{1}{m} \sum_{i=1}^m \mathbb{E}[X_i] - t\right) \leq \exp\left(-\frac{mt^2}{2\sigma^2}\right).$$

A.1 Proof of Proposition 1

Our proof uses Mendelson's "small-ball" techniques for concentration [32] along with control over a particular VC-dimension condition. If we define $h_i(u, v) := \kappa_{\text{st}}^2 \mathbf{1}\{|\langle a_i, u \rangle| \wedge |\langle a_i, v \rangle| \geq \kappa_{\text{st}}\}$, then we certainly have

$$\frac{1}{m} \sum_{i=1}^m |\langle a_i, u \rangle \langle a_i, v \rangle| \geq \frac{1}{m} \sum_{i=1}^m h_i(u, v) \quad \text{for all } u, v \in \mathbb{R}^n.$$

We now control the class $\mathcal{F} := \{a \mapsto f(a) = |\langle a, u \rangle| \wedge |\langle a, v \rangle| \mid u, v \in \mathbb{R}^n\}$ by its VC-dimension:

Lemma A.3. *There exists some numerical constant $C > 0$, such that, for any $c \geq 0$, the VC-dimension of the collection of sets*

$$\mathcal{G} := \{\{x \in \mathbb{R}^n \text{ s.t. } |\langle x, u \rangle| \wedge |\langle x, v \rangle| \geq c\} \mid u \in \mathbb{R}^n, v \in \mathbb{R}^n\}$$

is upper bounded by Cn .

Proof The collection of half planes

$$\mathcal{G}_{\text{plane}} := \{\{x \in \mathbb{R}^n \text{ s.t. } \langle u, x \rangle + b \geq 0\} \mid u \in \mathbb{R}^n, b \in \mathbb{R}\}$$

has VC-dimension at most $n + 2$ [43, Lemma 2.6.15]. We have the containment

$$\mathcal{G} \subset \{(G_1 \cup G_2) \cap (G_3 \cup G_4) \mid G_1, G_2, G_3, G_4 \in \mathcal{G}_{\text{plane}}\},$$

and so standard results on preservation of VC-dimension under set operations [43, Lemma 2.6.17], imply that the VC-dimension of \mathcal{G} is at most C times the VC-dimension of $\mathcal{G}_{\text{plane}}$ for some numerical constant $C > 0$. \square

The associated thresholds $a \mapsto 1 \{ |\langle a, u \rangle| \wedge |\langle a, v \rangle| \geq \kappa_{\text{st}} \}$ likewise have VC-dimension at most Cn as u, v vary. Applying standard VC-concentration inequalities [5, 43, Ch. 2.6] we immediately obtain that

$$\mathbb{P} \left(\sup_{u, v \in \mathbb{S}^{n-1}} \frac{1}{\kappa_{\text{st}}} \left| \frac{1}{m} \sum_{i=1}^m h_i(u, v) - \mathbb{E}[h_i(u, v)] \right| \geq c \sqrt{\frac{n}{m}} + t \right) \leq 2 \exp \left(-\frac{mt^2}{2} \right)$$

for all $t \geq 0$, where $c < \infty$ is a numerical constant. Substitute $t^2 \mapsto 2t$ to achieve the result.

A.2 Proof of Proposition 2

We begin with a small lemma relating the Frobenius norm of certain rank-two matrices to the distances between vectors.

Lemma A.4. *Let $X = xx^H - yy^H \in \mathbb{C}^{n \times n}$. Then $\|X\|_{\text{Fr}} \geq \inf_{\theta} \|x - e^{i\theta}y\|_2 \sup_{\theta} \|x - e^{i\theta}y\|_2$.*

We defer the proof of the lemma to Section A.2.1.

To prove the proposition, we define the function $F : \mathbb{C}^{m \times n} \times \mathbb{C}^{n \times n}$, for $Z = [z_1 \ \cdots \ z_m]^H$, by

$$F(Z, X) = \frac{1}{m} \sum_{i=1}^m |\langle z_i, X z_i \rangle|. \quad (24)$$

Now, for a constant $c > 0$ to be chosen, define the truncated variables

$$z_i := \sqrt{n} a_i / \|a_i\|_2 \mathbf{1} \{ \|a_i\|_2 \geq c\sqrt{n} \}. \quad (25)$$

Then we have that for any $A \in \mathbb{C}^{m \times n}$ that

$$F(A, xx^H - yy^H) \geq c^2 F(Z, xx^H - yy^H).$$

As the function F is homogenous in its second argument, based on Lemma A.4, the result of the theorem will hold if we can show that (with suitably high probability)

$$F(Z, X) \geq \frac{\tau^2}{4} \quad \text{for all rank 2 or less } X \in \mathbb{C}^{n \times n} \text{ with } \|X\|_{\text{Fr}} = 1. \quad (26)$$

It is, of course, no loss of generality to assume that X is Hermitian in inequality (26).

With our desired inequality (26) in mind, we present a covering-number-based argument to prove the theorem. The first step in this direction is to lower bound the expectation of $F(Z, X)$. Let $X \in \mathcal{S}_r$, $X = \sum_{j=1}^r \lambda_j u_j u_j^H$. Then we have that for $w_i = \sqrt{n} a_i / \|a_i\|_2$ and our choice of z_i that

$$\mathbb{E} [|z_i^H X z_i|] = \mathbb{E} [|w_i^H X w_i|] - \mathbb{E} [|w_i^H X w_i| \mathbf{1} \{ \|a_i\|_2 \leq c\sqrt{n} \}].$$

Now, we have

$$\mathbb{E} [|w_i^H X w_i| \mathbf{1} \{ \|a_i\|_2 \leq \epsilon\sqrt{n} \}] \stackrel{(i)}{\leq} \sqrt{\mathbb{E} [|w_i^H X w_i|^2]} \sqrt{h(c)} \stackrel{(ii)}{\leq} \left(\sum_{j=1}^r \lambda_j^2 \right)^{\frac{1}{2}} \mathbb{E} \left[\sum_{i=1}^r |\langle u_i, w_i \rangle|^4 \right]^{\frac{1}{2}} \sqrt{h(c)}$$

where the inequalities follow (i) by Cauchy-Schwartz and Assumption A3 (for the small-ball probability bound $h(c)$) and (ii) by Cauchy-Schwartz, respectively. As w_i is a σ^2 -sub-Gaussian vector, standard results on sub-Gaussians imply that $\mathbb{E} [|\langle u, w_i \rangle|^4] \leq (1 + e)\sigma^4$ for any unit vector u , and

thus the final expectation has bound $\sum_{i=1}^r \mathbb{E}[\langle u_i, w_i \rangle^4] \leq (1+e)r\sigma^4$, and thus we obtain for any choice of c in our construction (25) of z_i and any rank r matrix X that

$$\mathbb{E}[|z_i^H X z_i|] \geq \mathbb{E}[|w_i^H X w_i|] - \sqrt{1+e} \|X\|_{\text{Fr}} \sigma^2 \sqrt{rh(c)}.$$

In particular, as we consider only rank $r = 2$ matrices, Assumption A3 allows us to choose c in the definition (25) small enough that

$$h(c) \leq \frac{1}{2(1+e)} \cdot \frac{\tau^4}{\sigma^4},$$

and then

$$\mathbb{E}[|z_i^H X z_i|] \geq \mathbb{E}[|w_i^H X w_i|] - \tau^2 \|X\|_{\text{Fr}} / 2 \geq \frac{\tau^2}{2} \|X\|_{\text{Fr}}. \quad (27)$$

The remainder of our argument now proceeds by a covering argument, which we begin with a lemma on the continuity properties of F .

Lemma A.5. *Let Hermitian matrices $X, Y \in \mathbb{C}^{n \times n}$ have rank at most r and eigen-decompositions $X = U\Lambda U^H$ and $Y = V\Sigma V^H$, and let $Z = [z_1 \ \cdots \ z_m]^H \in \mathbb{C}^{m \times n}$. Then*

$$|F(Z, X) - F(Z, Y)| \leq \frac{2\sqrt{r}}{m} \|Z\|_{\text{op}}^2 \left[\|U - V\|_{1,2} + \|\Lambda - \Sigma\|_{\text{Fr}} \right].$$

We provide the proof of Lemma A.5 in Section A.2.2.

Based on Lemma A.5, we develop a particular covering set of the $n \times n$ Hermitian matrices of rank r , following a construction due to Candès and Plan [11, Thm. 2.3]. Let $\mathcal{S}_r \subset \{X \in \mathbb{C}^{n \times n} \mid X \text{ Hermitian, rank } r\}$ be the set of Hermitian rank r matrices with $\|X\|_{\text{Fr}} = 1$, and let $\mathcal{O}_{n,r} = \{U \in \mathbb{C}^{n \times r} \mid U^H U = I_r\}$ denote the set of $n \times r$ unitary matrices; for each $\epsilon > 0$ there exists an ϵ -cover $\overline{\mathcal{O}}_{n,r}$ of $\mathcal{O}_{n,r}$ in $\|\cdot\|_{1,2}$ -norm of cardinality at most $(3/\epsilon)^{2nr}$. Similarly, there exists an ϵ -cover $D \subset \mathbb{R}^r$ of all vectors $v \in \mathbb{R}^r$ in ℓ_2 -norm of cardinality at most $(3/\epsilon)^r$. Now, let $\overline{\mathcal{S}}_r = \{U \text{diag}(d)U^H \mid U \in \overline{\mathcal{O}}_{n,r}, d \in D\}$ be a subset of the rank- r Hermitian matrices. Then $\text{card}(\overline{\mathcal{S}}_r) \leq (3/\epsilon)^{(2n+1)r}$, and we have the following immediate consequence of Lemma A.5.

Lemma A.6. *For any $\epsilon > 0$, there exists a set $\overline{\mathcal{S}}_r \subset \mathcal{S}_r$ of cardinality at most $\text{card}(\overline{\mathcal{S}}_r) \leq (3/\epsilon)^{(2n+1)r}$ such that*

$$\inf_{X \in \mathcal{S}_r} F(Z, X) \geq \min_{X \in \overline{\mathcal{S}}_r} F(Z, X) - \frac{4\sqrt{r}}{m} \|Z\|_{\text{op}}^2 \epsilon.$$

For our final step, we control the variance of $F(Z, Y)$, so that we can apply the one-sided Bernstein inequality (Lemma A.2) and then a covering argument. For $X \in \mathcal{S}_r$, then, we have

$$\text{Var}(z_i^H X z_i) \leq \mathbb{E}[|z_i^H X z_i|^2] = \mathbb{E}\left[\left|\sum_{j=1}^r \lambda_j |\langle z_i, u_j \rangle|^2\right|^2\right] \leq \|X\|_{\text{Fr}}^2 \sum_{j=1}^r \mathbb{E}[|\langle z_i, u_j \rangle|^4] \leq r(1+e) \|X\|_{\text{Fr}}^2 \sigma^2$$

by Cauchy-Schwartz and that $\mathbb{E}[|\langle z_i, u_j \rangle|^4] \leq (1+e)\sigma^4$ by sub-Gaussian moment bounds. Thus, Lemma A.2 implies that for any fixed $X \in \mathcal{S}_r$ we have

$$\mathbb{P}\left(F(Z, X) \leq \frac{\tau^2}{2} - \sqrt{2(1+e)r} \cdot \sigma^2 t\right) \leq e^{-mt^2}$$

for all $t \geq 0$. Now, let \mathcal{E}_t be the event that $\frac{1}{m} \|Z\|_{\text{op}}^2 \leq \varepsilon_{n,m}(t) := \sigma^2 + 11\sigma^2 \max\{\sqrt{4n/m} + t, 4n/m + t\}$. Then by Lemma A.6 and Lemma 3.1 with $r = 2$, we have for any K and ϵ that

$$\begin{aligned} \mathbb{P}\left(\inf_{X \in \mathcal{S}_r} F(Z, X) \leq K\right) &\leq \mathbb{P}\left(\min_{X \in \mathcal{S}_r} F(Z, X) - \frac{4}{m} \|Z\|_{\text{op}}^2 \epsilon \leq K\right) \\ &\leq \mathbb{P}\left(\min_{X \in \mathcal{S}_r} F(Z, X) - \frac{4}{m} \|Z\|_{\text{op}}^2 \epsilon \leq K, \mathcal{E}_t\right) + \mathbb{P}(\mathcal{E}_t) \\ &\leq \mathbb{P}\left(\min_{X \in \mathcal{S}_r} F(Z, X) - 4\varepsilon_{n,m}(t)\epsilon \leq K\right) + e^{-mt} \\ &\leq \left(\frac{3}{\epsilon}\right)^{(2n+1)r} \sup_{X \in \mathcal{S}_r} \mathbb{P}(F(Z, X) - 4\varepsilon_{n,m}(t)\epsilon \leq K) + e^{-mt}. \end{aligned}$$

Letting $K = \frac{\tau^2}{2} - 2\sqrt{(1+e)}\sigma^2 t - 4\varepsilon_{n,m}(t)\epsilon$, we thus see that

$$\mathbb{P}\left(\inf_{X \in \mathcal{S}_2} F(Z, X) \leq \frac{\tau^2}{2} - 2\sqrt{(1+e)}\sigma^2 t - 4\varepsilon_{n,m}(t)\epsilon\right) \leq \left(\frac{3}{\epsilon}\right)^{2(2n+1)} e^{-mt^2} + e^{-mt}.$$

Summarizing, we have the following lemma.

Lemma A.7. *There exist numerical constants $C_1 \leq 2\sqrt{(1+e)}$ and $C_2 \leq 44$ such that*

$$\begin{aligned} \mathbb{P}\left(\inf_{X \in \mathcal{S}_2} F(Z, X) \leq \frac{\tau^2}{2} - C_1\sigma^2 t - C_2\sigma^2 \left(\max\left\{\sqrt{\frac{n}{m}} + t, \frac{n}{m} + t\right\} \epsilon + \epsilon\right)\right) \\ \leq \exp\left(-mt^2 + 2(2n+1)\log \frac{3}{\epsilon}\right) + e^{-mt}. \end{aligned}$$

Rewriting this slightly by taking $\epsilon, t \lesssim \frac{\tau^2}{\sigma^2}$, we see that there exist numerical constants $c_0 > 0$ and $c_1 < \infty$ such that

$$\mathbb{P}\left(\inf_{X \in \mathcal{S}_2} F(Z, X) \leq \frac{\tau^2}{4}\right) \leq \exp\left(-c_0 m \frac{\tau^4}{\sigma^4} + c_1 n \log \frac{\sigma^2}{\tau^2}\right).$$

Now, returning to inequality (26), we see we have that for $z_i = \sqrt{n}a_i / \|a_i\|_2 \mathbf{1}\{\|a_i\|_2 \geq c\sqrt{n}\}$ chosen as in (25) and $c > 0$ chosen small enough so that inequality (27) holds, we have

$$\inf_{X \in \mathcal{S}_2} F(Z, X) \geq \frac{\tau^2}{4}$$

with probability at least $1 - e^{-c_m \frac{\tau^4}{\sigma^4} + c_1 n \log \frac{\sigma^2}{\tau^2}}$. Comparing with inequality (26) gives the theorem.

A.2.1 Proof of Lemma A.4

For $X = xx^H - yy^H$ that

$$\begin{aligned} \|X\|_{\text{Fr}}^2 &= \|x\|_2^4 + \|y\|_2^4 - 2|\langle x, y \rangle|^2 \\ &= \|x\|_2^4 + \|y\|_2^4 - 2(\text{Re}(\langle x, y \rangle)^2 + \text{Im}(\langle x, y \rangle)^2) \\ &= \left(\sqrt{2\|x\|^4 + 2\|y\|^4} - 2\sqrt{\text{Re}(\langle x, y \rangle)^2 + \text{Im}(\langle x, y \rangle)^2}\right) \left(\sqrt{2\|x\|^4 + 2\|y\|^4} + 2\sqrt{\text{Re}(\langle x, y \rangle)^2 + \text{Im}(\langle x, y \rangle)^2}\right). \end{aligned}$$

By using that

$$\|x - e^{i\theta}y\|^2 = \|x\|^2 + \|y\|^2 - 2(\cos\theta \cdot \operatorname{Re}(\langle x, y \rangle) - \sin\theta \cdot \operatorname{Im}(\langle x, y \rangle)),$$

and that the concavity of $\sqrt{\cdot}$ implies $\sqrt{2a^4 + 2b^4} \geq a^2 + b^2$, the final expression above has lower bound

$$\begin{aligned} & \left(\|x\|^2 + \|y\|^2 - 2\sqrt{\operatorname{Re}(\langle x, y \rangle)^2 + \operatorname{Im}(\langle x, y \rangle)^2} \right) \left(\|x\|^2 + \|y\|^2 + 2\sqrt{\operatorname{Re}(\langle x, y \rangle)^2 + \operatorname{Im}(\langle x, y \rangle)^2} \right) \\ &= \inf_{\theta} \|x - e^{i\theta}y\|^2 \cdot \sup_{\theta} \|x - e^{i\theta}y\|^2. \end{aligned}$$

A.2.2 Proof of Lemma A.5

We have

$$\begin{aligned} m|F(Z, X) - F(Z, Y)| &\leq \sum_{i=1}^m \left| \sum_{j=1}^r (\lambda_j |\langle u_j, z_i \rangle|^2 - \sigma_j |\langle v_j, z_i \rangle|^2) \right| \\ &\leq \sum_{i=1}^m \left| \sum_{j=1}^r (\lambda_j (|\langle u_j, z_i \rangle|^2 - |\langle v_j, z_i \rangle|^2)) \right| + \sum_{i=1}^m \left| \sum_{j=1}^r ((\lambda_j - \sigma_j) |\langle v_j, z_i \rangle|^2) \right|. \end{aligned} \quad (28)$$

We bound each of the terms in expression (28) in term. For the first, we note that for any complex $u, v, z \in \mathbb{C}^n$, we have

$$|\langle u, z \rangle|^2 - |\langle v, z \rangle|^2 \leq |z^H u u^H z - z^H v v^H z + u^H z z^H v - v^H z z^H u| = |\langle u - v, z \rangle \langle z, u + v \rangle|,$$

as $u^H z z^H v - v^H z z^H u$ is purely imaginary. As a consequence, we have

$$\begin{aligned} \sum_{i=1}^m \left| \sum_{j=1}^r \lambda_j (|\langle u_j, z_i \rangle|^2 - |\langle v_j, z_i \rangle|^2) \right| &= \sum_{i=1}^m \left| \sum_{j=1}^r \lambda_j \langle u_j u_j^H - v_j v_j^H, z_i z_i^H \rangle \right| \\ &\leq \sum_{j=1}^r |\lambda_j| \sum_{i=1}^m |\langle z_i, u_j - v_j \rangle| |\langle z_i, u_j + v_j \rangle| \\ &\leq \sum_{j=1}^r |\lambda_j| \left(\sum_{i=1}^m |\langle z_i, u_j - v_j \rangle|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m |\langle z_i, u_j + v_j \rangle|^2 \right) \\ &\leq 2 \sum_{j=1}^r |\lambda_j| \|Z^T Z\|_{\text{op}} \|u_j - v_j\|_2 \leq 2 \|\lambda\|_1 \|Z^T Z\|_{\text{op}} \|U - V\|_{1,2}, \end{aligned}$$

where we have used that $\|u_j\|_2 = \|v_j\|_2 = 1$ so $\|u_j + v_j\|_2 \leq 2$. For the second term in inequality (28), we have

$$\sum_{i=1}^m \left| \sum_{j=1}^r ((\lambda_j - \sigma_j) |\langle v_j, z_i \rangle|^2) \right| \leq \sum_{j=1}^r |\lambda_j - \sigma_j| \sum_{i=1}^m |\langle v_j, z_i \rangle|^2 \leq \|\lambda - \sigma\|_1 \|Z^T Z\|_{\text{op}}.$$

Noting that for vectors $\lambda, \sigma \in \mathbb{R}^r$ we have $\|\lambda\|_1 \leq \sqrt{r} \|\lambda\|_2$, we obtain the lemma.

A.3 Proof of Proposition 3: deterministic part

Our proof of Proposition 3 eventually reduces to applying eigenvector perturbation results to the random matrices X^{init} . To motivate our approach, note that

$$X^{\text{init}} := \frac{1}{|\mathcal{I}_{\text{sel}}|} \sum_{i \in \mathcal{I}_{\text{sel}}} a_i a_i^H = I_n - \phi(\epsilon) \cdot d_\star d_\star^H + \Delta, \quad (29)$$

where Δ is a random error term that we will show has small norm. From the quantity (29) we can apply eigenvector perturbation arguments to derive that the directional estimate $\hat{d} = \arg\min_{d \in \mathbb{S}^{n-1}} d^H X^{\text{init}} d$ satisfies $\hat{d} \approx d_\star$. This will hold so long as $\|\Delta\|_{\text{op}}$ is small because there is substantial separation in the eigenvalues of I_n and $I_n - \phi(\epsilon) d_\star d_\star^H$.

With this goal in mind, we define two index sets that with high probability surround \mathcal{I}_{sel} . Let

$$\mathcal{I}_{-\epsilon} := \left\{ i \in [m] \mid |\langle a_i, d_\star \rangle|^2 < \frac{1-\epsilon}{2} \right\} \quad \text{and} \quad \mathcal{I}_{+\epsilon} := \left\{ i \in [m] \mid |\langle a_i, d_\star \rangle|^2 \leq \frac{1+\epsilon}{2} \right\}.$$

We now define events \mathcal{E}_1 through \mathcal{E}_5 , showing that conditional on these five, the result of the proposition holds. These events roughly guarantee (\mathcal{E}_1) that $\sum_{i=1}^m a_i a_i^H$ is well-behaved, (\mathcal{E}_2 and \mathcal{E}_3) that $\mathcal{I}_{+\epsilon} \setminus \mathcal{I}_{-\epsilon}$ is small, and (\mathcal{E}_4 and \mathcal{E}_5) that most of the vectors a_i for indices $i \in \mathcal{I}_{\text{sel}}$ are close enough to uniform on the subspace perpendicular to d_\star that we have a good directional estimate. Now, let $q \in [1, \infty]$ and let $1/p + 1/q = 1$, so that p is its conjugate. Recalling the definition of the error $\Delta(\epsilon)$ in Assumption A5.(ii), we define

$$\begin{aligned} \mathcal{E}_1 &:= \left\{ \frac{1}{m} \|A^H A\|_{\text{op}} \in [1-\epsilon, 1+\epsilon] \right\}, \quad \mathcal{E}_2 := \{ |\mathcal{I}_{+\epsilon}| \leq |\mathcal{I}_{-\epsilon}| + 2\epsilon \kappa m \}, \\ \mathcal{E}_3 &:= \left\{ |\mathcal{I}_{-\epsilon}| \geq \frac{1}{2} m p_0(d_\star) \right\}, \quad \mathcal{E}_4 := \left\{ \left\| \frac{1}{m} \sum_{i \in \mathcal{I}_{+\epsilon} \setminus \mathcal{I}_{-\epsilon}} a_i a_i^H \right\|_{\text{op}} \leq 4q\sigma^2(\kappa\epsilon)^{\frac{1}{p}} \right\} \\ \mathcal{E}_5 &:= \left\{ \left\| \frac{1}{|\mathcal{I}_{-\epsilon}|} \sum_{i \in \mathcal{I}_{-\epsilon}} a_i a_i^H - (I_n - \phi(\epsilon) d_\star d_\star^H) \right\|_{\text{op}} \leq \|\Delta(\epsilon)\|_{\text{op}} + \epsilon \right\}. \end{aligned} \quad (30)$$

We prove the result of the proposition when each \mathcal{E}_i occurs. Decompose the matrix X^{init} into

$$\begin{aligned} X^{\text{init}} &= \underbrace{I_n - \phi(\epsilon) d_\star d_\star^H}_{:=Z_0} \\ &+ \underbrace{\left[\frac{1}{|\mathcal{I}_{-\epsilon}|} \sum_{i \in \mathcal{I}_{-\epsilon}} a_i a_i^H - (I_n - \phi(\epsilon) d_\star d_\star^H) \right]}_{:=Z_1} + \underbrace{\frac{1}{|\mathcal{I}_{-\epsilon}|} \sum_{i \in \mathcal{I}_{\text{sel}} \setminus \mathcal{I}_{-\epsilon}} a_i a_i^H}_{:=Z_2} - \underbrace{\left(\frac{1}{|\mathcal{I}_{-\epsilon}|} - \frac{1}{|\mathcal{I}_{\text{sel}}|} \right) \sum_{i \in \mathcal{I}_{\text{sel}}} a_i a_i^H}_{:=Z_3}. \end{aligned}$$

We bound the operator norms of Z_1, Z_2, Z_3 in turn. On the event \mathcal{E}_5 , we have

$$\|Z_1\|_{\text{op}} \leq \|\Delta(\epsilon)\|_{\text{op}} + \epsilon. \quad (31)$$

We turn to the error matrix Z_2 . On the event \mathcal{E}_1 , we evidently have $\hat{r} \in \|x_\star\|_2 (1 \pm \epsilon)$ by definition of $\hat{r}^2 = \frac{1}{m} \|A x_\star\|_2^2$, so that

$$\mathcal{I}_{-\epsilon} \subset \mathcal{I}_{\text{sel}} \subset \mathcal{I}_{+\epsilon}.$$

Using that the summands $a_i a_i^H$ are all positive semidefinite, we thus obtain the upper bound

$$\|Z_2\|_{\text{op}} = \frac{1}{|\mathcal{I}_{-\epsilon}|} \left\| \sum_{i \in \mathcal{I}_{\text{sel}} \setminus \mathcal{I}_{-\epsilon}} a_i a_i^H \right\|_{\text{op}} \leq \frac{1}{|\mathcal{I}_{-\epsilon}|} \left\| \sum_{i \in \mathcal{I}_{+\epsilon} \setminus \mathcal{I}_{-\epsilon}} a_i a_i^H \right\|_{\text{op}} \stackrel{(i)}{\leq} \frac{1}{\mathfrak{p}_0(d_\star)} \cdot 4q\sigma^2(\kappa\epsilon)^{\frac{1}{p}}, \quad (32)$$

where in inequality (i) we use that $2|\mathcal{I}_{-\epsilon}| \geq \mathfrak{p}_0(d_\star)m$ on \mathcal{E}_3 and $\|\sum_{i \in \mathcal{I}_{+\epsilon} \setminus \mathcal{I}_{-\epsilon}} a_i a_i^H\|_{\text{op}} \leq 4qm\sigma^2(\kappa\epsilon)^{\frac{1}{p}}$ on \mathcal{E}_4 . Lastly, we provide an upper bound on $\|Z_3\|_{\text{op}}$. Again using that $\mathcal{I}_{-\epsilon} \subset \mathcal{I}_{\text{sel}} \subset \mathcal{I}_{+\epsilon}$ on event \mathcal{E}_1 , we have

$$\left| \frac{1}{|\mathcal{I}_{-\epsilon}|} - \frac{1}{|\mathcal{I}_{\text{sel}}|} \right| \leq \frac{1}{|\mathcal{I}_{-\epsilon}|} - \frac{1}{|\mathcal{I}_{+\epsilon}|} = \frac{|\mathcal{I}_{+\epsilon}| - |\mathcal{I}_{-\epsilon}|}{|\mathcal{I}_{-\epsilon}||\mathcal{I}_{+\epsilon}|} \leq \frac{2\epsilon\kappa}{\mathfrak{p}_0(d_\star)^2 m},$$

where in the last inequality, we use that $|\mathcal{I}_{+\epsilon}| - |\mathcal{I}_{-\epsilon}| \leq 2\epsilon\kappa m$ on \mathcal{E}_2 and that $|\mathcal{I}_{+\epsilon}| \geq |\mathcal{I}_{-\epsilon}| \geq \mathfrak{p}_0(d_\star)m/2$ on \mathcal{E}_3 . Thus, by the definition of \mathcal{E}_1 , we have

$$\|Z_3\|_{\text{op}} \leq \frac{2\epsilon\kappa}{\mathfrak{p}_0(d_\star)^2} \left\| \frac{1}{m} \sum_{i=1}^m a_i a_i^H \right\|_{\text{op}} \leq \frac{2\epsilon(1+\epsilon)\kappa}{\mathfrak{p}_0(d_\star)^2}. \quad (33)$$

Combining inequalities (31), (32) and (33) on the error matrices Z_i , the triangle inequality gives

$$\|Z_1 + Z_2 + Z_3\|_{\text{op}} \leq \|\Delta(\epsilon)\|_{\text{op}} + \underbrace{\left[\epsilon + \frac{4q\sigma^2(\kappa\epsilon)^{\frac{1}{p}}}{\mathfrak{p}_0(d_\star)} + \frac{2\epsilon(1+\epsilon)\kappa}{\mathfrak{p}_0(d_\star)^2} \right]}_{:=\nu(\epsilon)}.$$

This implies equality (29) with error bound $\|\Delta\|_{\text{op}} \leq \|\Delta(\epsilon)\|_{\text{op}} + \nu(\epsilon)$. Recall the definition of $Z_0 = I_n - \phi(\epsilon)d_\star d_\star^H$, which has smallest eigenvector d_\star and eigengap $\phi(\epsilon)$. Lastly, we simplify $\nu(\epsilon)$ by a specific choice of p and q in the definition (30) of \mathcal{E}_4 . Without loss of generality, we assume $\kappa\epsilon < 1$ (recall Assumption A5 on κ), and define $p = 1 + \frac{1}{\log \frac{1}{\kappa\epsilon}}$ and $q = 1 + \log \frac{1}{\kappa\epsilon}$. Using that for any $z < 0$ we have $\exp(\frac{z}{1-1/z}) \leq e^{z+1}$, we have $(\kappa\epsilon)^{\frac{1}{p}} \leq e\kappa\epsilon$, allowing us to bound $q(\kappa\epsilon)^{\frac{1}{p}} \leq e(1 + \log \frac{1}{\kappa\epsilon})(\kappa\epsilon)$ in the error term $\nu(\epsilon)$.

We now apply the eigenvector perturbation inequality of Lemma A.1. Using that, for $\theta \in \mathbb{R}$, $\|u - e^{i\theta}v\|_2^2 \geq 2-2|\langle u, v \rangle| \geq 2-2|\langle u, v \rangle|^2$ for $\|u\|_2 = \|v\|_2 = 1$, a minor rearrangement of Lemma A.1 applied to $X^{\text{init}} = Z_0 + \Delta$ for $\Delta = Z_1 + Z_2 + Z_3$ yields

$$\text{dist}\left(\hat{d}, \frac{1}{r}X_\star\right) = \inf_{\theta \in [0, 2\pi]} \left\| \hat{d} - e^{i\theta}d \right\|_2 \leq \frac{\sqrt{2}(\|\Delta(\epsilon)\|_{\text{op}} + \nu(\epsilon))}{\left[\phi(\epsilon) - (\|\Delta(\epsilon)\|_{\text{op}} + \nu(\epsilon)) \right]_+}.$$

Finally, using that $x_0 = \hat{r}\hat{d}$ and defining $r_\star = \|x_\star\|$, we have by the triangle inequality that

$$\text{dist}(x_0, X_\star) \leq \hat{r} \text{dist}\left(\hat{d}, \frac{1}{r}X_\star\right) + |r_\star - \hat{r}| \stackrel{(i)}{\leq} \left[\sqrt{1+\epsilon} \text{dist}(\hat{d}, d_\star) + \epsilon \right] r_\star,$$

where inequality (i) uses \mathcal{E}_1 , which we recall implies $(1-\epsilon)\|x_\star\|_2^2 \leq \hat{r}^2 \leq (1+\epsilon)\|x_\star\|_2^2$, where $\epsilon \in [0, 1]$. This is the claimed consequence (14) in Proposition 3.

A.4 Proof of Proposition 3: high probability events

It remains to demonstrate that each of the events $\mathcal{E}_1, \dots, \mathcal{E}_5$ (recall definition (30)) holds with high probability, to which we dedicate the remainder of this argument in the next series of lemmas, each of which argues that one of the five events occurs with high probability. Before the statement of each lemma, we recall the corresponding event whose high probability we wish to demonstrate. Throughout, $c > 0$ and $C < \infty$ denote numerical constants whose values may change.

We begin with $\mathcal{E}_1 := \{\frac{1}{m}\|A^H A\|_{\text{op}} \in [1 \pm \epsilon]\}$.

Lemma A.8. *We have $\mathbb{P}(\mathcal{E}_1) \geq 1 - \exp(-cm\epsilon^2/\sigma^4)$ for m large enough that $m/n \geq \sigma^4/(c\epsilon^2)$.*

Proof Set $t = c\frac{\epsilon^2}{\sigma^4}$ in Lemma 3.1, noting that we must have $\sigma^2 \geq 1$ because $\mathbb{E}[aa^H] = I_n$ and $\|\mathbb{E}[aa^H]\|_{\text{op}} \leq \sigma^2$ (recall the final part of Lemma 3.1). Moreover, $\epsilon \in [0, 1]$ by assumption. Then taking c small enough, once we have $\frac{n}{m} \leq c\frac{\epsilon^2}{\sigma^4}$ we obtain the result. \square

The event $\mathcal{E}_2 := \{|\mathcal{I}_{-\epsilon}| \geq |\mathcal{I}_{+\epsilon}| - 2\kappa\epsilon m\}$ likewise holds with high probability.

Lemma A.9. *We have $\mathbb{P}(\mathcal{E}_2) \geq 1 - \exp(-2\epsilon^2\kappa^2 m)$.*

Proof We always have that

$$\mathcal{I}_{-\epsilon} \subset \mathcal{I}_{+\epsilon} \quad \text{and} \quad \mathcal{I}_{+\epsilon} \setminus \mathcal{I}_{-\epsilon} = \left\{ i \in [m] : \frac{1}{2}(1 - \epsilon) \leq |\langle a_i, d_\star \rangle|^2 \leq \frac{1}{2}(1 + \epsilon) \right\}.$$

Therefore, the difference in cardinalities of $|\mathcal{I}_{-\epsilon}|$ and $|\mathcal{I}_{+\epsilon}|$ is

$$\frac{1}{m} (|\mathcal{I}_{+\epsilon}| - |\mathcal{I}_{-\epsilon}|) = \frac{1}{m} \sum_{i=1}^m 1_{\{1 - \epsilon \leq 2|\langle a_i, d_\star \rangle|^2 \leq 1 + \epsilon\}}.$$

The right hand side is an average of i.i.d. Bernoulli random variables with means bounded by $\kappa\epsilon$ by Assumption A5(i). Hoeffding's inequality gives the result that $\mathbb{P}(|\mathcal{I}_{+\epsilon}| > |\mathcal{I}_{-\epsilon}| + 2\kappa m\epsilon) \leq e^{-2\kappa^2\epsilon^2 m}$. \square

Lemma A.10. *The event $\mathcal{E}_3 := \{|\mathcal{I}_{-\epsilon}| \geq \frac{1}{2}m\mathbf{p}_0(d_\star)\}$ satisfies $\mathbb{P}(\mathcal{E}_3) \geq 1 - \exp(-\frac{1}{2}m\mathbf{p}_0(d_\star)^2)$.*

Proof As in Lemma A.9, this result is immediate from Hoeffding's inequality. We have $|\mathcal{I}_{-\epsilon}| = \sum_{i=1}^m 1_{\{|\langle a_i, d_\star \rangle|^2 \leq (1 - \epsilon)/2\}}$, an i.i.d sum of Bernoullis with $\mathbb{P}(|\langle a_i, d_\star \rangle|^2 \leq (1 - \epsilon)/2) \geq \mathbf{p}_0(d_\star)$ by Assumption A5(i). Hoeffding's inequality gives the result. \square

Showing that events \mathcal{E}_4 and \mathcal{E}_5 in Eq. (30) each happen with high probability requires a little more work. We begin with \mathcal{E}_4 , defined in terms of a conjugate pair $p, q \geq 1$ with $1/p + 1/q = 1$, as $\mathcal{E}_4 := \{\|\sum_{i \in \mathcal{I}_{+\epsilon} \setminus \mathcal{I}_{-\epsilon}} a_i a_i^H\|_{\text{op}} \leq 4q\sigma^2(\kappa\epsilon)^{\frac{1}{p}}\}$.

Lemma A.11. *If $m/n > c^{-1}(\kappa\epsilon)^{-\frac{2}{p}}$, then $\mathbb{P}(\mathcal{E}_4) \geq 1 - \exp(-cm(\kappa\epsilon)^{\frac{2}{p}})$.*

It is no loss of generality to assume that $\kappa\epsilon \leq 1$ by Assumption A5, so $(\kappa\epsilon)^{\frac{2}{p}} \geq (\kappa\epsilon)^2$.

Proof The $\{a_i\}_{i=1}^m$ are independent σ^2 -sub-Gaussian random vectors by Assumption A4, and for any random variable B_i with $|B_i| \leq 1$, which may depend on a_i , it is clear that the collection $\{B_i a_i\}_{i=1}^m$ are mutually independent and still satisfy Definition 3.1. To that end, define the Bernoulli

variables $B(a) = 1\{|\langle a, d_\star \rangle|^2 \in [\frac{1-\epsilon}{2}, \frac{1+\epsilon}{2}]\}$ (letting $B_i = B(a_i) = 1\{i \in \mathcal{I}_{+\epsilon} \setminus \mathcal{I}_{-\epsilon}\}$ for shorthand). Then Lemma 3.1 implies for a numerical constant $C < \infty$ that

$$\mathbb{P}\left(\left\|\frac{1}{m}\sum_{i=1}^m a_i a_i^H B_i - \mathbb{E}[a a^H B(a)]\right\|_{\text{op}} \geq C\sigma^2 \max\left\{\sqrt{\frac{n}{m} + t}, \frac{n}{m} + t\right\}\right) \leq e^{-mt}. \quad (34)$$

Now, note by Hölder's inequality that

$$\begin{aligned} \mathbb{E}[\langle v, a \rangle^2 B(a)] &\leq \mathbb{E}[\langle v, a \rangle^{2q}]^{\frac{1}{q}} \mathbb{P}\left(|\langle a, d_\star \rangle|^2 \in \left[\frac{1-\epsilon}{2}, \frac{1+\epsilon}{2}\right]\right)^{\frac{1}{p}} \\ &\leq (\sigma^{2q} \Gamma(q+1) e)^{\frac{1}{q}} (\kappa\epsilon)^{\frac{1}{p}} \leq q e^{1/q} \sigma^2 (\kappa\epsilon)^{\frac{1}{p}} \end{aligned}$$

where we have applied Assumption A5(i) and Lemma 3.1 to bound $\mathbb{E}[\langle a, d_\star \rangle^{2q}]$. Using the triangle inequality and substituting $t = \frac{1}{4C}(\kappa\epsilon)^{\frac{2}{p}}$ into inequality (34), we find that for any $q \in (1, \infty)$ and $1/p + 1/q = 1$, if $\frac{n}{m} \leq \frac{1}{4C}(\kappa\epsilon)^{\frac{2}{p}}$ we have

$$\mathbb{P}\left(\left\|\frac{1}{m}\sum_{i=1}^m a_i a_i^H B_i - \mathbb{E}[a a^H B(a)]\right\|_{\text{op}} \geq q\sigma^2 (\kappa\epsilon)^{\frac{1}{p}}\right) \leq \exp\left(-\frac{m(\kappa\epsilon)^{\frac{2}{p}}}{4C}\right).$$

Applying the triangle inequality and that $1 + e^{1/q} < 4$ gives the result. \square

The final high probability guarantee is the most complex, and applies to the event $\mathcal{E}_5 := \{\|\frac{1}{|\mathcal{I}_{-\epsilon}|} \sum_{i \in \mathcal{I}_{-\epsilon}} a_i a_i^H - (I_n - \phi(\epsilon) d_\star d_\star^H)\|_{\text{op}} \leq \|\Delta(\epsilon)\|_{\text{op}} + \epsilon\}$.

Lemma A.12. *Let $\mathcal{E}_3 = \{|\mathcal{I}_{-\epsilon}| \geq \frac{1}{2} m \mathbf{p}_0(d_\star)\}$ as in Eq. (30). Then*

$$c \frac{m}{n} \geq \frac{\sigma^4 \log^2 \mathbf{p}_0(d_\star)}{\mathbf{p}_0(d_\star) \epsilon^2}$$

implies $\mathbb{P}(\mathcal{E}_5 \mid \mathcal{E}_3) \geq 1 - \exp(-c \frac{m \mathbf{p}_0(d_\star) \epsilon^2}{\sigma^4 \log^2 \mathbf{p}_0(d_\star)})$.

Proof For notational simplicity, define the following shorthand:

$$E_{d_\star} := \mathbb{E}\left[aa^H \mid |\langle a, d_\star \rangle|^2 \leq \frac{1-\epsilon}{2}\right] = I_n - \phi(\epsilon) d_\star d_\star^H + \Delta(\epsilon),$$

where the equality uses Assumption A5(ii). The main idea of the proof is to show the following crucial fact: define the new sub-Gaussian parameter $\tau^2 = \sigma^2 \log \frac{e}{\mathbf{p}_0(d_\star)} \geq 1$. Then there exists a numerical constant $1 \leq C < \infty$ such that for all $t \geq 0$,

$$\mathbb{P}\left(\left\|\frac{1}{|\mathcal{I}_{-\epsilon}|} \sum_{i \in \mathcal{I}_{-\epsilon}} a_i a_i^H - E_{d_\star}\right\|_{\text{op}} \geq C\tau^2 \max\left\{\sqrt{\frac{n}{|\mathcal{I}_{-\epsilon}|} + t}, \frac{n}{|\mathcal{I}_{-\epsilon}|} + t\right\} \mid \mathcal{I}_{-\epsilon}\right) \leq \exp(-|\mathcal{I}_{-\epsilon}|t). \quad (35)$$

Suppose that the bound (35) holds. On the event \mathcal{E}_3 , we have that $|\mathcal{I}_{-\epsilon}| \geq \frac{1}{2} m \mathbf{p}_0(d_\star)$, and so choosing

$$t = \frac{\epsilon^2}{4C^2 \tau^4} < 1, \quad \text{and letting} \quad \frac{m}{n} \geq \frac{2C^2 \tau^4}{\mathbf{p}_0(d_\star) \epsilon^2}$$

yields that $\frac{n}{|\mathcal{I}_{-\epsilon}|} + t < 1$ on \mathcal{E}_3 and

$$\mathbb{P} \left(\left\| \frac{1}{|\mathcal{I}_{-\epsilon}|} \sum_{i \in \mathcal{I}_{-\epsilon}} a_i a_i^H - E_{d_\star} \right\|_{\text{op}} \geq \epsilon \mid \mathcal{I}_{-\epsilon} \right) \leq \exp \left(-\frac{m \mathbf{p}_0(d_\star) \epsilon^2}{8C^2 \tau^4} \right).$$

By the definition of E_{d_\star} and the triangle inequality we have

$$\begin{aligned} \mathbb{P} \left(\left\| \frac{1}{|\mathcal{I}_{-\epsilon}|} \sum_{i \in \mathcal{I}_{-\epsilon}} a_i a_i^H - (I_n - \phi(\epsilon) d_\star d_\star^H) \right\|_{\text{op}} \geq \|\Delta(\epsilon)\|_{\text{op}} + \epsilon \mid \mathcal{I}_{-\epsilon} \right) \\ \leq \mathbb{P} \left(\left\| \frac{1}{|\mathcal{I}_{-\epsilon}|} \sum_{i \in \mathcal{I}_{-\epsilon}} a_i a_i^H - E_{d_\star} \right\|_{\text{op}} \geq \epsilon \mid \mathcal{I}_{-\epsilon} \right). \end{aligned}$$

The lemma follows from the fact that \mathcal{E}_3 is measurable with respect to the indices $\mathcal{I}_{-\epsilon}$.

Now, we show the key inequality (35). The main idea is to show that, conditioning on the set $\mathcal{I}_{-\epsilon}$, the distribution $\{a_i\}_{i \in \mathcal{I}_{-\epsilon}}$ is still conditionally independent and sub-Gaussian. To do so, we introduce a bit of (more or less standard) notation. For a random variable X , let $\mathcal{L}(X)$ denote the law of distribution of X . Using the independence of the vectors a_i , we have the fact that for any *fixed* subset $\mathcal{I} \subset [m]$, the collection $\{a_i\}_{i \in \mathcal{I}}$ is independent of $\{a_i\}_{i \notin \mathcal{I}}$. Therefore, using the definition that $\mathcal{I}_{-\epsilon} = \{i \in [m] : |\langle a_i, d_\star \rangle|^2 \leq \frac{1-\epsilon}{2}\}$, we have the key identity

$$\mathcal{L} \left(\frac{1}{|\mathcal{I}_{-\epsilon}|} \sum_{i \in \mathcal{I}_{-\epsilon}} a_i a_i^H \mid \mathcal{I}_{-\epsilon} = \mathcal{I} \right) \stackrel{\text{dist}}{=} \mathcal{L} \left(\frac{1}{|\mathcal{I}|} \sum_{i \in \mathcal{I}} a_i a_i^H \mid \max_{i \in \mathcal{I}} |\langle a_i, d_\star \rangle|^2 \leq \frac{1-\epsilon}{2} \right).$$

This implies that, conditioning on $\mathcal{I}_{-\epsilon} = \mathcal{I}$, the vectors $\{a_i\}_{i \in \mathcal{I}}$ are still conditionally independent, and their conditional distribution is identical to the law $\mathcal{L}(a \mid |\langle a, d_\star \rangle|^2 \leq \frac{1-\epsilon}{2})$. The claimed inequality (35) will thus follow by the matrix concentration inequality in Lemma 3.1, so long as we can demonstrate appropriate sub-Gaussianity of the conditional law $\mathcal{L}(a \mid |\langle a, d_\star \rangle|^2 \leq \frac{1-\epsilon}{2})$.

Indeed, let us temporarily assume that $a \mid |\langle a, d_\star \rangle|^2 \leq \frac{1-\epsilon}{2}$ is τ^2 -sub-Gaussian, let \mathcal{J} denote all subsets $\mathcal{I} \subset [m]$ such that $|\mathcal{I}| \geq \frac{1}{2} m \mathbf{p}_0(d_\star)$, and define the shorthand $E_{d_\star} = \mathbb{E}[a a^H \mid |\langle a, d_\star \rangle|^2 \leq \frac{1-\epsilon}{2}]$. Then by summing over \mathcal{J} , we have on the event \mathcal{E}_3 that for a numerical constant $C < \infty$,

$$\begin{aligned} \mathbb{P} \left(\left\| \frac{1}{|\mathcal{I}_{-\epsilon}|} \sum_{i \in \mathcal{I}_{-\epsilon}} a_i a_i^H - E_{d_\star} \right\|_{\text{op}} \geq C \tau^2 \max \left\{ \sqrt{\frac{n}{|\mathcal{I}_{-\epsilon}|}} + t, \frac{n}{|\mathcal{I}_{-\epsilon}|} + t \right\} \mid \mathcal{E}_3 \right) \\ = \sum_{\mathcal{I} \in \mathcal{J}} \mathbb{P} \left(\left\| \frac{1}{|\mathcal{I}|} \sum_{i \in \mathcal{I}} a_i a_i^H - E_{d_\star} \right\|_{\text{op}} \geq C \tau^2 \max \left\{ \sqrt{\frac{n}{|\mathcal{I}|}} + t, \frac{n}{|\mathcal{I}|} + t \right\} \mid \mathcal{I}_{-\epsilon} = \mathcal{I} \right) \mathbb{P}(\mathcal{I} = \mathcal{I}_{-\epsilon} \mid \mathcal{E}_3) \\ = \sum_{\mathcal{I} \in \mathcal{J}} \mathbb{P} \left(\left\| \frac{1}{|\mathcal{I}|} \sum_{i \in \mathcal{I}} a_i a_i^H - E_{d_\star} \right\|_{\text{op}} \geq C \tau^2 \max \left\{ \sqrt{\frac{n}{|\mathcal{I}|}} + t, \frac{n}{|\mathcal{I}|} + t \right\} \mid \max_{i \in \mathcal{I}} |\langle a_i, d_\star \rangle|^2 \leq \frac{1-\epsilon}{2} \right) \mathbb{P}(\mathcal{I} = \mathcal{I}_{-\epsilon} \mid \mathcal{E}_3) \\ \stackrel{(i)}{\leq} \sum_{\mathcal{I} \in \mathcal{J}} e^{-|\mathcal{I}|t} \cdot \mathbb{P}(\mathcal{I} = \mathcal{I}_{-\epsilon} \mid \mathcal{E}_3) \leq e^{-\frac{1}{2} m \mathbf{p}_0(d_\star) t}, \end{aligned}$$

where inequality (i) is an application of Lemma 3.1. This is evidently inequality (35) with appropriate choice of τ^2 .

We thus show that $\mathcal{L}(a \mid |\langle a, d_\star \rangle|^2 \leq \frac{1-\epsilon}{2})$ is subgaussian with parameter $\tau^2 = \sigma^2 \log \frac{e}{\mathbf{p}_0(d_\star)}$ by bounding the conditional moment generating function. Let $\lambda \in [1, \infty]$ and λ' be conjugate, so that $1/\lambda + 1/\lambda' = 1$. Then by Hölder's inequality, for any $v \in \mathbb{S}^{n-1}$ we have

$$\begin{aligned} \mathbb{E} \left[\exp \left(\frac{|\langle a, v \rangle|^2}{\lambda \sigma^2} \right) \mid |\langle a, v \rangle|^2 \leq \frac{1-\epsilon}{2} \right] &= \frac{\mathbb{E} \left[\exp \left(\frac{|\langle a, v \rangle|^2}{\lambda \sigma^2} \right) \mathbf{1} \{ |\langle a, v \rangle|^2 \leq \frac{1-\epsilon}{2} \} \right]}{\mathbb{P} (|\langle a, v \rangle|^2 \leq \frac{1-\epsilon}{2})} \\ &\leq \frac{(\mathbb{E}[\exp(\frac{|\langle a, v \rangle|^2}{\sigma^2})])^{\frac{1}{\lambda}} \mathbb{P} (|\langle a, v \rangle|^2 \leq \frac{1-\epsilon}{2})^{\frac{1}{\lambda'}}}{\mathbb{P} (|\langle a, v \rangle|^2 \leq \frac{1-\epsilon}{2})} \\ &= \mathbb{E} \left[\exp \left(\frac{|\langle a, v \rangle|^2}{\sigma^2} \right) \right]^{\frac{1}{\lambda}} \mathbb{P} \left(|\langle a, v \rangle|^2 \leq \frac{1-\epsilon}{2} \right)^{-\frac{1}{\lambda}} \leq \left(\frac{e}{\mathbf{p}_0(d_\star)} \right)^{\frac{1}{\lambda}}, \end{aligned}$$

where the final inequality uses the σ^2 -sub-Gaussianity of a . Set $\lambda = \log \frac{e}{\mathbf{p}_0(d_\star)}$ to see that conditional on $|\langle a, d_\star \rangle|^2 \leq \frac{1-\epsilon}{2}$, the vector a is $\sigma^2 \log \frac{e}{\mathbf{p}_0(d_\star)}$ -sub-Gaussian, as desired. \square

B Proofs for phase retrieval with outliers

In this section, we collect the proofs of the various results in Section 4. Before providing the proofs, we state one inequality that we use frequently that will be quite useful. Let $W_i \in \{0, 1\}$ satisfy $W_i = 1$ if $i \in \mathcal{I}^{\text{out}}$ and $W_i = 0$ otherwise, so that W indexes the outlying measurements. Because W_i are independent of the a_i vectors and $\sum_i W_i = p_{\text{fail}} m$, Lemma 3.1 implies for all $t \geq 0$ that

$$\mathbb{P} \left(\left\| \frac{1}{m} \sum_{i=1}^m W_i a_i a_i^T - p_{\text{fail}} \mathbb{E}[a_i a_i^T] \right\|_{\text{op}} \geq C \sigma^2 \max \left\{ \sqrt{\frac{n}{m}} + t, \frac{n}{m} + t \right\} \right) \leq e^{-mt}. \quad (36)$$

B.1 Proof of Proposition 4

Recalling the set \mathcal{I}^{out} of outlying indices, we evidently have

$$\begin{aligned} f(x) - f(x_\star) &= \frac{1}{m} \sum_{i=1}^m |\langle a_i, x \rangle^2 - \langle a_i, x_\star \rangle^2| + \frac{1}{m} \sum_{i \in \mathcal{I}^{\text{out}}} (|\langle a_i, x \rangle^2 - \xi_i| - |\langle a_i, x \rangle^2 - \langle a_i, x_\star \rangle^2|) - f(x_\star) \\ &= \frac{\|(Ax)^2 - (Ax_\star)^2\|_1}{m} + \frac{1}{m} \sum_{i \in \mathcal{I}^{\text{out}}} (|\langle a_i, x \rangle^2 - \xi_i| - |\langle a_i, x \rangle^2 - \langle a_i, x_\star \rangle^2| - |\langle a_i, x_\star \rangle^2 - \xi_i|) \\ &\geq \frac{\|(Ax)^2 - (Ax_\star)^2\|_1}{m} - \frac{2}{m} \sum_{i \in \mathcal{I}^{\text{out}}} |\langle a_i, x \rangle^2 - \langle a_i, x_\star \rangle^2|. \end{aligned} \quad (37)$$

Now, we note the trivial fact that if we define $W_i = 1$ for $i \in \mathcal{I}^{\text{out}}$ and $W_i = 0$ for $i \in \mathcal{I}^{\text{in}}$, then

$$\begin{aligned} \sum_{i \in \mathcal{I}^{\text{out}}} |\langle a_i, x \rangle^2 - \langle a_i, x_\star \rangle^2| &= \sum_{i=1}^m W_i |(x - x_\star)^T a_i a_i^T (x + x_\star)| \\ &\leq \|A^T \text{diag}(W) A\|_{\text{op}} \|x - x_\star\|_2 \|x + x_\star\|_2. \end{aligned}$$

Inequality (36) shows that the matrix $\frac{1}{m} \sum_{i=1}^m W_i a_i a_i^T$ is well concentrated.

Now, let \mathcal{E} denote the event that $\frac{1}{m}\|A^T \text{diag}(W)A\|_{\text{op}} \leq p_{\text{fail}} + C\sigma^2\sqrt{n/m+t}$, where t is chosen so that $n/m+t \leq 1$. Returning to inequality (37), we have

$$f(x) - f(x_\star) \geq \frac{1}{m} \|(Ax)^2 - (Ax_\star)^2\|_1 - 2(p_{\text{fail}} + C\sigma^2\sqrt{n/m+t}) \|x - x_\star\|_2 \|x + x_\star\|_2$$

for all $x \in \mathbb{R}^n$ with probability at least $1 - e^{-mt}$ by inequality (36). We finish with the following lemma, which is a minor sharpening of Theorem 2.4 of Eldar and Mendelson [19] so that we have sharp concentration in all dimensions n . We provide a proof for completeness in Section C.2.

Lemma B.1. *Let a_i be independent σ^2 -sub-Gaussian vectors, and define $\kappa_{\text{st}}(u, v) := \mathbb{E}[\langle a, u \rangle \langle a, v \rangle]$ for $u, v \in \mathbb{R}^n$. Then there exist numerical constants $c > 0$ and $C < \infty$ such that*

$$\frac{1}{m} \sum_{i=1}^m |\langle a_i, u \rangle \langle a_i, v \rangle| \geq \kappa_{\text{st}}(u, v) - C\sigma^2 \sqrt[3]{\frac{n}{m}} - \sigma^2 t \quad \text{for all } u, v \in \mathbb{S}^{n-1}$$

with probability at least $1 - e^{-cmt^2} - e^{-cm}$ when $m/n \geq C$.

Noting that $|\langle a_i, x \rangle^2 - \langle a_i, x_\star \rangle^2| = |\langle a_i, x - x_\star \rangle \langle a_i, x + x_\star \rangle|$ and substituting the result of the lemma into the preceding display, we have

$$f(x) - f(x_\star) \geq \left(\kappa_{\text{st}} - 2p_{\text{fail}} - C\sigma^2 \sqrt[3]{\frac{n}{m}} - C\sigma^2 t \right) \|x - x_\star\|_2 \|x + x_\star\|_2$$

uniformly in x with probability at least $1 - 2e^{-cmt^2} - e^{-cm}$.

B.2 Proof of Proposition 5

We first state a lemma providing a deterministic bound on the errors of the minimizing radius.

Lemma B.2. *Let*

$$\delta = \frac{6 \|x_\star\|_2^2 \|A^T A\|_{\text{op}}}{\sum_{i \in \mathcal{I}^{\text{in}}} \langle a_i, x_\star \rangle^2 - \sum_{i \in \mathcal{I}^{\text{out}}} \langle a_i, x_\star \rangle^2} \text{dist}(\hat{d}, \{\pm d_\star\}). \quad (38)$$

If $\delta \leq 1$, then all minimizers of $G(\cdot)$ belong to the set $[1 \pm \delta] \|x_\star\|_2^2$.

Temporarily assuming the conclusions of the lemma, let us show that the random quantities in the bound (38) are small with high probability. We apply the matrix concentration inequality (36) to see that for a numerical constant $C < \infty$ and all $t \in [0, 1 - \frac{n}{m}]$, we have

$$\frac{1}{m} \sum_{i \in \mathcal{I}^{\text{in}}} \langle a_i, v \rangle^2 \geq \left[(1 - p_{\text{fail}}) - C\sigma^2 \sqrt{\frac{n}{m} + t} \right] \quad \text{and} \quad \frac{1}{m} \sum_{i \in \mathcal{I}^{\text{out}}} \langle a_i, v \rangle^2 \leq \left[p_{\text{fail}} + C\sigma^2 \sqrt{\frac{n}{m} + t} \right]$$

with probability at least $1 - e^{-mt}$ for all vectors $v \in \mathbb{S}^{n-1}$, and $\frac{1}{m}\|A^T A\|_{\text{op}} \leq \sigma^2(1 + C\sqrt{\frac{n}{m} + t})$ with the same probability. That is, for $t \in [0, 1 - \frac{n}{m}]$ with probability at least $1 - 2e^{-mt}$ we have that δ in expression (38) satisfies

$$\delta \leq \frac{6\sigma^2}{1 - 2p_{\text{fail}} - C\sigma^2 \sqrt{\frac{n}{m} + t}} \text{dist}(\hat{d}, \{\pm d_\star\}).$$

If we assume that $\frac{n}{m} \leq c(1-2p_{\text{fail}})^2/\sigma^4$ and replace t with $c(1-2p_{\text{fail}})^2/\sigma^4$ for small enough constant c , we find that

$$\delta \leq \frac{C\sigma^2}{1-2p_{\text{fail}}} \text{dist}(\hat{d}, \{\pm d_\star\})$$

with probability at least $1 - 2e^{-cm(1-2p_{\text{fail}})^2/\sigma^4}$, where C is a numerical constant. This is our desired result.

Proof We define a few pieces of notation for shorthand. Let

$$\hat{\sigma}^2 := \frac{1}{m} \|A^T A\|_{\text{op}} \quad \text{and} \quad \hat{\sigma}_{\mathcal{I}^{\text{out}}}^2 := \frac{1}{m} \sum_{i \in \mathcal{I}^{\text{in}}} \langle a_i, x_\star \rangle^2 - \frac{1}{m} \sum_{i \in \mathcal{I}^{\text{out}}} \langle a_i, x_\star \rangle^2,$$

and define the functions $\hat{g}(\delta) = G((1+\delta) \|x_\star\|_2^2)$, equivalently

$$\hat{g}(\delta) := \frac{1}{m} \sum_{i=1}^m |b_i - \|x_\star\|_2^2 \langle a_i, \hat{d} \rangle (1+\delta)|,$$

and a slightly more accurate counterpart

$$g(\delta) := \frac{1}{m} \sum_{i=1}^m |b_i - (1+\delta) \langle a_i, x_\star \rangle^2| = \frac{1}{m} \sum_{i \notin \mathcal{I}^{\text{in}}} |\langle a_i, x_\star \rangle^2 - (1+\delta) \langle a_i, x_\star \rangle^2| + \frac{1}{m} \sum_{i \in \mathcal{I}^{\text{out}}} |b_i - (1+\delta) \langle a_i, x_\star \rangle^2|.$$

Note that if δ minimizes $\hat{g}(\delta)$, then $(1+\delta) \|x_\star\|_2^2$ minimizes $G(r)$. By inspection we find that the subgradients of g with respect to δ are

$$\partial_\delta g(\delta) = \frac{1}{m} \sum_{i \in \mathcal{I}^{\text{in}}} \text{sgn}(\delta) \langle a_i, x_\star \rangle^2 - \frac{1}{m} \sum_{i \in \mathcal{I}^{\text{out}}} \text{sgn}((1+\delta) \langle a_i, x_\star \rangle^2 - b_i) \langle a_i, x_\star \rangle^2,$$

where $\text{sgn}(t) = 1$ if $t > 0$, -1 if $t < 0$, and $\text{sgn}(0) = [-1, 1]$. Evidently, for $\delta > 0$ we have $g'(\delta) \geq \hat{\sigma}_{\mathcal{I}^{\text{out}}}^2$ and $g'(-\delta) \leq -\hat{\sigma}_{\mathcal{I}^{\text{out}}}^2$, so that

$$g(\delta) \geq \hat{\sigma}_{\mathcal{I}^{\text{out}}}^2 |\delta| + g(0). \quad (39)$$

Now, we consider the gaps between \hat{g} and g : for $\delta \in [-1, 1]$, we have the gap

$$\begin{aligned} |g(\delta) - \hat{g}(\delta)| &\leq \frac{(1+\delta) \|x_\star\|_2^2}{m} \sum_{i=1}^m |\langle a_i, d_\star \rangle^2 - \langle a_i, \hat{d} \rangle^2| \\ &\leq \frac{(1+\delta) \|x_\star\|_2^2}{m} \|A^T A\|_{\text{op}} \|d_\star - \hat{d}\|_2 \|d_\star + \hat{d}\|_2 \leq 4 \|x_\star\|_2^2 \hat{\sigma}^2 \text{dist}(d_\star, \{\pm \hat{d}\}), \end{aligned}$$

where we have used the triangle inequality and Cauchy-Schwarz. Thus we obtain

$$\hat{g}(\delta) - \hat{g}(0) \geq g(\delta) - g(0) + \hat{g}(\delta) - g(\delta) + g(0) - \hat{g}(0) \geq \hat{\sigma}_{\mathcal{I}^{\text{out}}}^2 |\delta| - 6 \|x_\star\|_2^2 \hat{\sigma}^2 \text{dist}(\hat{d}, \{\pm d_\star\}),$$

where we have applied inequality (39). Rearranging, we have that if $\hat{g}(\delta) \leq \hat{g}(0)$ we must have

$$|\delta| \leq \frac{6 \|x_\star\|_2^2 \hat{\sigma}^2}{\hat{\sigma}_{\mathcal{I}^{\text{out}}}^2} \text{dist}(\hat{d}, \{\pm d_\star\}).$$

By convexity, any minimizer of \hat{g} must thus lie in the above region, which gives the result when we recall that minimizers δ of \hat{g} are equivalent to minimizers $(1+\delta) \|x_\star\|_2^2$ of G . \square

B.3 Proof of Proposition 6

We introduce a bit of notation before giving the proof proper. Recall that $\mathcal{I}^{\text{out}} \subset [m]$ denotes the outliers, or failed measurements, and $\mathcal{I}^{\text{in}} = [m] \setminus \mathcal{I}^{\text{out}}$ the set of i such that $b_i = \langle a_i, x_\star \rangle^2$ (the inliers). Recalling the selected set of indices \mathcal{I}_{sel} , we define the shorthand

$$\mathcal{I}_{\text{sel}}^{\text{in}} := \mathcal{I}^{\text{in}} \cap \mathcal{I}_{\text{sel}} \quad \text{and} \quad \mathcal{I}_{\text{sel}}^{\text{out}} := \mathcal{I}^{\text{out}} \cap \mathcal{I}_{\text{sel}}.$$

for the chosen inliers and outliers.

We decompose the matrix X^{init} into four matrices, each of which we control to guarantee that $X^{\text{init}} \approx I_n - cd_\star d_\star^T$ for some constant c , thus guaranteeing $\hat{d} \approx d_\star$. Let $P = d_\star d_\star^T$ and $P_\perp = I_n - d_\star d_\star^T$ be the projection operator onto the span of d_\star and its orthogonal complement. Then we may decompose the matrix X^{init} into the four parts

$$\begin{aligned} X^{\text{init}} = & \underbrace{\frac{1}{m} \sum_{i=1}^m P a_i a_i^T P \mathbf{1}_{\{i \in \mathcal{I}_{\text{sel}}^{\text{in}}\}}}_{:=Z_0} + \underbrace{\frac{1}{m} \sum_{i=1}^m [P a_i a_i^T P_\perp + P_\perp a_i a_i^T P] \mathbf{1}_{\{i \in \mathcal{I}_{\text{sel}}^{\text{in}}\}}}_{:=Z_1} \\ & + \underbrace{\frac{1}{m} \sum_{i=1}^n P_\perp a_i a_i^T P_\perp \mathbf{1}_{\{i \in \mathcal{I}_{\text{sel}}^{\text{in}}\}}}_{:=Z_2} + \underbrace{\frac{1}{m} \sum_{i=1}^m a_i a_i^T \mathbf{1}_{\{i \in \mathcal{I}_{\text{sel}}^{\text{out}}\}}}_{:=Z_3}. \end{aligned} \quad (40)$$

Let us briefly motivate this decomposition. We expect that Z_0 should be small because we choose indices \mathcal{I}_{sel} by taking the smallest values of b_i , which should be least correlated with d_\star (recall the $P = d_\star d_\star^T$). We expect Z_1 to be small because of the independence of the vectors $P a_i$ and $P_\perp a_i$ for Gaussian measurement vectors, and Z_3 to be small because $\mathcal{I}_{\text{sel}}^{\text{out}}$ should be not too large. This leaves Z_2 , which (by Gaussianity) we expect to be some multiple of $I_n - d_\star d_\star^T$, which will then allow us to apply eigenvector perturbation guarantees using the eigengap of the matrix X^{init} .

The rotational invariance of the Gaussian means that it is no loss of generality to assume that $d_\star = e_1$, the first standard basis vector, so for the remainder of the argument we assume this without further comment. This means that we may decompose a_i as $a_i = [a_{i,1} \ a_{i,2} \ \cdots \ a_{i,n}]^T = [a_{i,1} \ a_{i,\setminus 1}^T]^T$, which we will do without further comment for the remainder of the proof.

We now present four lemmas, each controlling one of the terms Z_l in the expansion (40). We defer proofs of each of the lemmas to the end of this argument. We begin by considering the Z_0 term, which (because P is rank one) satisfies

$$Z_0 = \underbrace{\left(\frac{1}{m} \sum_{i=1}^m \langle a_i, d_\star \rangle^2 \mathbf{1}_{\{i \in \mathcal{I}_{\text{sel}}^{\text{in}}\}} \right)}_{:=z_0} d_\star d_\star^T.$$

Recalling the definition (17) of the constants δ_q and w_q in the statement of the proposition, we have the following lemma.

Lemma B.3. *Define the random quantities $z_0 = \frac{1}{m} \sum_{i=1}^m \langle a_i, d_\star \rangle^2 \mathbf{1}_{\{i \in \mathcal{I}_{\text{sel}}^{\text{in}}\}}$ and $z_2 = \frac{1}{m} |\mathcal{I}_{\text{sel}}^{\text{in}}|$. Then for $t \in [0, 1]$,*

$$\mathbb{P}(z_2 \geq z_0 + (1 - 2p_{\text{fail}})(1 - t)\delta_q) \geq 1 - \exp\left(-\frac{1}{10}(1 - 2p_{\text{fail}})m\right) - 2\exp\left(-\frac{m}{4}\right) - \exp\left(-\frac{mt^2\delta_q^2}{4w_q^2}\right).$$

See Sec. B.3.1 for a proof of the lemma. We thus see that it is likely that z_0 is substantially smaller than the rough fraction of inlying indices selected.

We now argue that Z_1 is likely to be small because it is the sum of products of independent vectors.

Lemma B.4. *For $t \geq 0$ we have*

$$\mathbb{P} \left(\|Z_1\|_{\text{op}} \geq 2\sqrt{\frac{n}{m}} + t \right) \leq \exp \left(-\frac{mt^2}{8} \right) + \exp \left(-\frac{m}{2} \right).$$

See Section B.3.2 for a proof. We can also show that Z_2 is well-behaved in the sense that it is approximately a scaled multiple of $(I - d_\star d_\star^T)$.

Lemma B.5. *Let z_2 be the random quantity $z_2 := \frac{1}{m}|\mathcal{I}_{\text{sel}}^{\text{in}}|$. There exists a numerical constant C such that for $t \in [0, 1]$ we have*

$$\mathbb{P} \left(\left\| Z_2 - z_2(I_n - d_\star d_\star^T) \right\|_{\text{op}} \geq C \left(\sqrt{\frac{n}{m}} + t \right) \right) \leq \exp(-mt).$$

See Section B.3.3 for a proof of the lemma.

Finally, we control the size of the error matrix Z_3 in the expansion (40), which corresponds to the contribution of the outlying measurements a_i that are included in the initialization matrix X^{init} . We provide two slightly different guarantees, depending on the model (strength) of adversarial noise ξ assumed.

Lemma B.6. *Define the random quantity $z_3 := \frac{1}{m}|\mathcal{I}^{\text{out}} \cap \mathcal{I}_{\text{sel}}| \leq p_{\text{fail}}$. There is a numerical constant C such that the following hold.*

(i) *Under the independent noise model M1, for all $t \in [0, 1]$,*

$$\mathbb{P} \left(\|Z_3 - z_3 I_n\|_{\text{op}} \geq C \sqrt{\frac{n}{m}} + t \right) \leq \exp(-mt).$$

(ii) *Under the adversarial noise model M2, for all $t \in [0, 1]$,*

$$\mathbb{P} \left(\|Z_3\|_{\text{op}} \geq p_{\text{fail}} + C \sqrt{\frac{n}{m}} + t \right) \leq \exp(-mt).$$

See Section B.3.4 for a proof.

With these four lemmas in hand, we can prove the result of the proposition by applying the eigenvector perturbation Lemma A.1. The expansion (40), coupled with the four lemmas defining the constants $z_0 = \frac{1}{m} \sum_{i \in \mathcal{I}_{\text{sel}}^{\text{in}}} \langle a_i, d_\star \rangle^2$, $z_2 = \frac{1}{m}|\mathcal{I}_{\text{sel}}^{\text{in}}|$, and $z_3 = \frac{1}{m}|\mathcal{I}_{\text{sel}}^{\text{out}}|$, guarantees that

$$\begin{aligned} X^{\text{init}} &= Z_0 + Z_1 + Z_2 + Z_3 = z_0 d_\star d_\star^T + z_2(I_n - d_\star d_\star^T) + z_3 I_n + \Delta \\ &= (z_2 + z_3)I_n - (z_2 - z_0)d_\star d_\star^T + \Delta, \end{aligned}$$

where the perturbation $\Delta \in \mathbb{R}^{n \times n}$ satisfies

$$\|\Delta\|_{\text{op}} \leq \|Z_1\|_{\text{op}} + \left\| Z_2 - z_2(I_n - d_\star d_\star^T) \right\|_{\text{op}} + \begin{cases} \|Z_3 - z_3 I_n\|_{\text{op}} & \text{under model M1} \\ \|Z_3\|_{\text{op}} & \text{under model M2.} \end{cases}$$

On the event that $z_2 > z_0$, the minimal eigenvector of $(z_2 + z_3)I_n - (z_2 - z_0)d_\star d_\star^T$ is d_\star with eigengap $z_2 - z_0$. Applying Lemma A.1 gives that $\hat{d} = \operatorname{argmin}_{d \in \mathbb{S}^{n-1}} d^T X^{\text{init}} d$ satisfies

$$2^{-\frac{1}{2}} \operatorname{dist}(\hat{d}, \{\pm d_\star\}) \leq \sqrt{1 - \langle \hat{d}, d_\star \rangle^2} \leq \frac{\|\Delta\|_{\text{op}}}{\left[z_2 - z_0 - \|\Delta\|_{\text{op}}\right]_+}. \quad (41)$$

Applying Lemmas B.4 and B.5, we have for some numerical constant $C < \infty$ that

$$\|Z_1\|_{\text{op}} + \|Z_2 - z_2(I_n - d_\star d_\star^T)\|_{\text{op}} \leq C \sqrt{\frac{n}{m} + t} \quad \text{with probability} \geq 1 - e^{-mt} - e^{-m/2}$$

for any $t \geq 0$. We now consider the two noise models in turn. Under Model M1, Lemma B.6 then implies that $\|\Delta\|_{\text{op}} \leq C \sqrt{\frac{n}{m} + t}$ with probability at least $1 - e^{-mt} - e^{-m/2}$. Recalling Lemma B.3, we have for the constants w_q^2 and $\delta_q > 0$ defined in the lemma that $z_2 \geq z_0 + \frac{1-2p_{\text{fail}}}{2} \delta_q$ with probability at least $1 - \exp(-c(1 - 2p_{\text{fail}})m) - \exp(-cm\delta_q^2/w_q^2)$, where $c > 0$ is a numerical constant. That is, the perturbation inequality (41) implies that under Model M1, we have with probability at least $1 - \exp(-mt) - \exp(-c(1 - 2p_{\text{fail}})m) - \exp(-cm\delta_q^2/w_q^2)$ that

$$2^{-\frac{1}{2}} \operatorname{dist}(\hat{d}, \{\pm d_\star\}) \leq \frac{C \sqrt{n/m + t}}{\left[(1 - 2p_{\text{fail}})\delta_q - C \sqrt{n/m + t}\right]_+},$$

where $0 < c, C < \infty$ are numerical constants. Under Model M2, we can bound $\|Z_3\|_{\text{op}}$ by $p_{\text{fail}} + C \sqrt{n/m + t}$ with probability $1 - e^{-mt} - e^{-m/2}$ (recall Lemma B.6), so that with the same probability as above, we have

$$2^{-\frac{1}{2}} \operatorname{dist}(\hat{d}, \{\pm d_\star\}) \leq \frac{p_{\text{fail}} + C \sqrt{n/m + t}}{\left[(1 - 2p_{\text{fail}})\delta_q - C \sqrt{n/m + t} - p_{\text{fail}}\right]_+} \quad \text{under Model M2.}$$

This is the proposition.

B.3.1 Proof of Lemma B.3

As noted earlier, it is no loss of generality to assume that $d_\star = e_1$, the first standard basis vector, so that using our definitions of $z_0 = \frac{1}{m} \sum_{i \in \mathcal{I}_{\text{sel}}^{\text{in}}} \langle a_i, d_\star \rangle^2$ and $z_2 = \frac{1}{m} |\mathcal{I}_{\text{sel}}^{\text{in}}|$, we have

$$z_2 - z_0 = \frac{1}{m} \sum_{i=1}^m (1 - a_{i,1}^2) 1_{\{i \in \mathcal{I}_{\text{sel}}^{\text{in}}\}}. \quad (42)$$

Given that we choose in indices \mathcal{I}_{sel} by a median, it is helpful to have the following median concentration result.

Lemma B.7. *Let $\{W_i\}_{i=1}^m \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$. Fix $p \in (0, 1)$ and choose $w_q \geq 0$ so that $q := \mathbb{P}(W^2 \leq w_q^2) = 2(1 - \Phi(w_q)) > p$. Then*

$$\mathbb{P}(\operatorname{quant}_p(\{W_i^2\}_{i=1}^m) \geq w_q^2) \leq \exp\left(-\frac{m(q-p)^2}{2(q-p)/3 + 2q(1-q)}\right).$$

Proof Note that $\text{quant}_p(\{W_i^2\}_{i=1}^m) \geq w_q^2$ if and only if $\frac{1}{m} \sum_{i=1}^m 1\{W_i^2 \leq w_q^2\} \leq p$. Using that $\text{Var}(1\{W_i^2 \leq w_q^2\}) = q(1-q)$, Bernstein's inequality applied to the i.i.d. sum $\sum_{i=1}^m (1\{W_i^2 \leq w_q^2\} - q)$ gives the result. \square

We now control the median of the perturbed vector $b \in \mathbb{R}^m$. Since we have $|\mathcal{I}^{\text{out}}| \leq p_{\text{fail}}m$, we have deterministic result

$$\text{med}(\{b_i\}_{i=1}^m) \leq \text{quant}_{\frac{1}{2(1-p_{\text{fail}})}}(\{\langle a_i, x_\star \rangle^2\}_{i \in \mathcal{I}^{\text{in}}}),$$

so by upper bounding the right hand quantity we can upper bound $\text{med}(\{b_i\})$, which in turn allows us to control \mathcal{I}_{sel} . By the definition of w_q and q_{fail} in Eq. (17), which satisfies $\delta = q_{\text{fail}} - \frac{1}{2(1-p_{\text{fail}})} = \frac{1-2p_{\text{fail}}}{4(1-p_{\text{fail}})}$, Lemma B.7 with $q = q_{\text{fail}}$ and the fact that $|\mathcal{I}^{\text{in}}| = (1-p_{\text{fail}})m$ then implies

$$\begin{aligned} \mathbb{P}\left(\text{med}(\{b_i\}_{i=1}^m) \geq w_q^2 \|x_\star\|_2^2\right) &\leq \mathbb{P}\left(\text{quant}_{(2(1-p_{\text{fail}}))^{-1}}(\{\langle a_i, x_\star \rangle^2\}_{i \in \mathcal{I}^{\text{in}}}) \geq w_q^2 \|x_\star\|_2^2\right) \\ &\leq \exp\left(-\frac{m(1-p_{\text{fail}})\delta^2}{2\delta/3 + 2\delta(1-\delta)}\right) = \exp\left(-\frac{3(1-2p_{\text{fail}})m}{4(2+6(1-\delta))}\right) \leq \exp\left(-\frac{3}{32}(1-2p_{\text{fail}})m\right). \end{aligned} \quad (43)$$

We now consider the indices i that are inliers for which $\langle a_i, d_\star \rangle^2$ is small; again letting $w_q \geq 0$ be defined as in the quantile (17), we define

$$\mathcal{I}_q^{\text{in}} := \{i \in \mathcal{I}^{\text{in}} \mid \langle a_i, d_\star \rangle^2 \leq w_q^2\} = \{i \in \mathcal{I}^{\text{in}} \mid \langle a_i, x_\star \rangle^2 \leq w_q^2 \|x_\star\|_2^2\}.$$

Lemma B.8. *Let the set of inliers $\mathcal{I}_q^{\text{in}}$ be defined as above, and let $\delta_q = 1 - \mathbb{E}[W^2 \mid W^2 \leq w_q^2]$ for $W \sim \mathcal{N}(0, 1)$. Then for all $t \geq 0$ we have*

$$\mathbb{P}\left(\frac{1}{|\mathcal{I}_q^{\text{in}}|} \sum_{i \in \mathcal{I}_q^{\text{in}}} (1 - a_{i,1}^2) \leq \delta_q - t \mid \mathcal{I}_q^{\text{in}}\right) \leq \exp\left(-\frac{2|\mathcal{I}_q^{\text{in}}|t^2}{w_q^2}\right).$$

We defer the proof of Lemma B.8, continuing on to give the proof of Lemma B.3.

We now integrate out the conditioning in Lemma B.8. Recalling the definition (17) of w_q in terms of p_{fail} , we have that $q_{\text{fail}} = \frac{3-2p_{\text{fail}}}{4(1-p_{\text{fail}})} > \frac{1}{2}$ for $p_{\text{fail}} \in [0, 1/2)$. By Hoeffding's inequality we have $\mathbb{P}(|\mathcal{I}_q^{\text{in}}| \leq m/8) \leq e^{-m/4}$ because $|\mathcal{I}^{\text{in}}| \geq (1-p_{\text{fail}})m \geq m/2$, whence we obtain

$$\begin{aligned} \mathbb{P}\left(\frac{1}{|\mathcal{I}_q^{\text{in}}|} \sum_{i \in \mathcal{I}_q^{\text{in}}} (1 - a_{i,1}^2) \leq \delta_q - t\right) &\leq \exp\left(-\frac{mt^2}{4w_q^2}\right) \mathbb{P}(|\mathcal{I}_q^{\text{in}}| \geq m/8) + \mathbb{P}(|\mathcal{I}_q^{\text{in}}| \leq m/8) \\ &\leq \exp\left(-\frac{mt^2}{4w_q^2}\right) + \exp\left(-\frac{m}{4}\right). \end{aligned}$$

Using the notation δ_q in Lemma B.8, for $t \in [0, 1]$ we define the event

$$\mathcal{E} := \left\{ \text{med}(\{b_i\}_{i=1}^m) \leq w_q^2 \|x_\star\|_2^2, |\mathcal{I}_q^{\text{in}}| \geq \frac{m}{8}, \frac{1}{|\mathcal{I}_q^{\text{in}}|} \sum_{i \in \mathcal{I}_q^{\text{in}}} (1 - a_{i,1}^2) \geq (1-t)\delta_q \right\}.$$

We immediately find that

$$\mathbb{P}(\mathcal{E}) \geq 1 - \exp\left(-\frac{1}{10}(1-2p_{\text{fail}})m\right) - 2\exp\left(-\frac{m}{4}\right) - \exp\left(-\frac{mt^2\delta_q^2}{4w_q^2}\right) \quad (44)$$

by the preceding display and inequality (43). Recalling the set $\mathcal{I}_q^{\text{in}} = \{i \in \mathcal{I}^{\text{in}} : \langle a_i, d_\star \rangle^2 \leq w_q^2\}$, we have on the event \mathcal{E} that the selected inliers satisfy $\mathcal{I}_{\text{sel}}^{\text{in}} \subset \mathcal{I}_q^{\text{in}}$ (because $\text{med}(b) \leq w_q^2 \|x_\star\|_2^2$). Because of our selection mechanism with $\mathcal{I}_{\text{sel}}^{\text{in}}$ as the smallest b_i in the sample, we have that

$$\frac{1}{|\mathcal{I}_{\text{sel}}^{\text{in}}|} \sum_{i \in \mathcal{I}_{\text{sel}}^{\text{in}}} (1 - a_{i,1}^2) \geq \frac{1}{|\mathcal{I}_q^{\text{in}}|} \sum_{i \in \mathcal{I}_q^{\text{in}}} (1 - a_{i,1}^2).$$

Moreover, on the event \mathcal{E} the rightmost sum is positive, and using that $|\mathcal{I}_{\text{sel}}^{\text{in}}| \geq |\mathcal{I}^{\text{in}}| + |\mathcal{I}_{\text{sel}}| - m \geq (\frac{1}{2} - p_{\text{fail}})m$, we obtain that on \mathcal{E} we have

$$\frac{1}{m} \sum_{i \in \mathcal{I}_{\text{sel}}^{\text{in}}} (1 - a_{i,1}^2) \geq \frac{1 - 2p_{\text{fail}}}{2|\mathcal{I}_{\text{sel}}^{\text{in}}|} \sum_{i \in \mathcal{I}_{\text{sel}}^{\text{in}}} (1 - a_{i,1}^2) \geq \frac{1 - 2p_{\text{fail}}}{2|\mathcal{I}_q^{\text{in}}|} \sum_{i \in \mathcal{I}_q^{\text{in}}} (1 - a_{i,1}^2) \geq \frac{1 - 2p_{\text{fail}}}{2} \delta_q.$$

Recalling expression (42) thus gives Lemma B.3.

Proof of Lemma B.8 Fix any set of indices $\mathcal{I}_0 \subset [m]$, and note that by Hoeffding's inequality for bounded random variables we have

$$\mathbb{P} \left(\frac{1}{|\mathcal{I}_0|} \sum_{i \in \mathcal{I}_0} a_{i,1}^2 - \mathbb{E}[a_{i,1}^2 \mid a_{i,1}^2 \leq w_q^2] \geq t \mid a_{i,1}^2 \leq w_q^2 \text{ for } i \in \mathcal{I}_0 \right) \leq \exp \left(-\frac{2|\mathcal{I}_0|t^2}{w_q^2} \right)$$

for $t \geq 0$. Recalling the definition $\delta_q := 1 - \mathbb{E}[W^2 \mid W^2 \leq w_q^2]$ for $W \sim \mathcal{N}(0, 1)$, we thus find that for any index set $\mathcal{I}_0 \subset [m]$, we have

$$\begin{aligned} & \mathbb{P} \left(\frac{1}{|\mathcal{I}_q^{\text{in}}|} \sum_{i \in \mathcal{I}_q^{\text{in}}} (1 - a_{i,1}^2) \leq \delta_q - t \mid \mathcal{I}_q^{\text{in}} = \mathcal{I}_0 \right) \\ &= \mathbb{P} \left(\frac{1}{|\mathcal{I}_0|} \sum_{i \in \mathcal{I}_0} (1 - a_{i,1}^2) \leq \delta_q - t \mid a_{i,1}^2 \leq w_q^2 \text{ for } i \in \mathcal{I}_0 \right) \leq \exp \left(-\frac{2|\mathcal{I}_0|t^2}{w_q^2} \right). \end{aligned}$$

Noticing that the random vectors $\{a_i\}_{i \in \mathcal{I}_0}$ are independent of $\{a_i\}_{i \notin \mathcal{I}_0}$, we have for any measurable set $\mathcal{C} \subset \mathbb{R}$ that

$$\mathbb{P} \left(\sum_{i \in \mathcal{I}_q^{\text{in}}} (1 - a_{i,1}^2) \in \mathcal{C} \mid \mathcal{I}_q^{\text{in}} = \mathcal{I}_0 \right) = \mathbb{P} \left(\sum_{i \in \mathcal{I}_0} (1 - a_{i,1}^2) \in \mathcal{C} \mid a_{i,1}^2 \leq w_q^2 \text{ for } i \in \mathcal{I}_0 \right).$$

Combining the preceding two displays yields Lemma B.8. \square

B.3.2 Proof of Lemma B.4

By our assumption (w.l.o.g.) that $d_\star = e_1$, we have

$$\|Z_1\|_{\text{op}} = \left\| \frac{1}{m} \sum_{i=1}^n a_{i,1} a_{i,\setminus 1} \mathbf{1}_{\{i \in \mathcal{I}_{\text{sel}}^{\text{in}}\}} \right\|_2 \quad (45)$$

Letting $\{a'_i\}_{i=1}^m \stackrel{\text{iid}}{\sim} \mathcal{N}(0, I_n)$ be an independent collection of vectors, the collections $\{a_i\}_{i \in \mathcal{I}^{\text{in}}}$ and $\{\xi_i\}_{i \in \mathcal{I}^{\text{out}}}$ are independent, as are $a_{i,1}$ and $a_{i,\setminus 1}$ for each i . Thus, because $\mathcal{I}_{\text{sel}}^{\text{in}} \subset \mathcal{I}^{\text{in}}$, that for any measurable set $\mathcal{C} \subset \mathbb{R}^{n-1}$ we have

$$\mathbb{P} \left(\sum_{i \in \mathcal{I}_{\text{sel}}^{\text{in}}} a_{i,1} a_{i,\setminus 1} \in \mathcal{C} \mid \{a_{i,1}\}_{i \in \mathcal{I}^{\text{in}}}, \{\xi_i\}_{i \in \mathcal{I}^{\text{out}}} \right) = \mathbb{P} \left(\sum_{i \in \mathcal{I}_{\text{sel}}^{\text{in}}} a_{i,1} a'_{i,\setminus 1} \in \mathcal{C} \mid \{a_{i,1}\}_{i \in \mathcal{I}^{\text{in}}}, \mathcal{I}_{\text{sel}}^{\text{in}} \right). \quad (46)$$

Now, we use the standard result [30] that if f is an L -Lipschitz function with respect to the ℓ_2 -norm, then for any standard Gaussian vector Z we have $\mathbb{P}(f(Z) - \mathbb{E}[f(Z)] \geq t) \leq \exp(-\frac{t^2}{2L^2})$ for all $t \geq 0$. Thus, defining the random Lipschitz constant $\widehat{L}^2 = \frac{1}{m} \sum_{i \in \mathcal{I}^{\text{in}}} a_{i,1}^2$ we obtain

$$\mathbb{E} \left[\left\| \frac{1}{m} \sum_{i=1}^m a_{i,1} a'_{i,\setminus 1} 1_{\{i \in \mathcal{I}_{\text{sel}}^{\text{in}}\}} \right\|_2^2 \mid \mathcal{I}_{\text{sel}}^{\text{in}}, \{a_{i,1}\}_{i \in \mathcal{I}^{\text{in}}} \right] = \frac{1}{m^2} \sum_{i \in \mathcal{I}_{\text{sel}}^{\text{in}}} a_{i,1}^2 (n-1) \leq \frac{n-1}{m} \widehat{L}^2,$$

and thus for $t \geq 0$

$$\mathbb{P} \left(\left\| \frac{1}{m} \sum_{i=1}^m a_{i,1} a'_{i,\setminus 1} 1_{\{i \in \mathcal{I}_{\text{sel}}^{\text{in}}\}} \right\|_2 \geq \widehat{L} \sqrt{\frac{n-1}{m}} + t \mid \{a_{i,1}\}_{i \in \mathcal{I}^{\text{in}}}, \mathcal{I}_{\text{sel}}^{\text{in}} \right) \leq \exp \left(-\frac{mt^2}{2\widehat{L}^2} \right).$$

Moreover, \widehat{L} is $\{a_{i,1}\}_{i \in \mathcal{I}^{\text{in}}}$ -measurable, and $\mathbb{P}(\widehat{L} \geq 2) \leq \exp(-m/2)$, again by the Lipschitz concentration of Gaussian random variables. We thus apply Eq. (46) and obtain

$$\begin{aligned} & \mathbb{P} \left(\left\| \frac{1}{m} \sum_{i \in \mathcal{I}_{\text{sel}}^{\text{in}}} a_{i,1} a_{i,\setminus 1} \right\|_2 \geq 2\sqrt{\frac{n-1}{m}} + t \right) \\ & \leq \mathbb{P} \left(\left\| \frac{1}{m} \sum_{i \in \mathcal{I}_{\text{sel}}^{\text{in}}} a_{i,1} a_{i,\setminus 1} \right\|_2 \geq 2\sqrt{\frac{n-1}{m}} + t \mid \widehat{L} \leq 2 \right) \mathbb{P}(\widehat{L} \leq 2) + \mathbb{P}(\widehat{L} \geq 2) \\ & \leq \exp \left(-\frac{mt^2}{8} \right) + \exp \left(-\frac{m}{2} \right). \end{aligned}$$

Recalling the equality (46) on $\|Z_1\|_{\text{op}}$ shows that the previous display gives the lemma.

B.3.3 Proof of Lemma B.5

Our first observation is simply the definition of the projection operator P_{\perp} , which gives

$$\|Z_{m,2} - z_2(I_n - d_{\star} d_{\star}^T)\|_{\text{op}} = \left\| \frac{1}{m} \sum_{i \in \mathcal{I}_{\text{sel}}^{\text{in}}} a_{i,\setminus 1} a_{i,\setminus 1}^T - z_2 I_{n-1} \right\|_{\text{op}}.$$

As in the proof of Lemma B.4, we let $\{a'_i\}_{i=1}^m \stackrel{\text{iid}}{\sim} \mathcal{N}(0, I_n)$ be an independent copy of the a_i . The independence of $\{a_i\}_{i \in \mathcal{I}^{\text{in}}}$ and $\{\xi_i\}_{i \in \mathcal{I}^{\text{out}}}$ imply that for any measurable set \mathcal{C} we have

$$\mathbb{P} \left(\sum_{i \in \mathcal{I}_{\text{sel}}^{\text{in}}} (a_{i,\setminus 1} a_{i,\setminus 1}^T - z_2 I_{n-1}) \in \mathcal{C} \mid \{a_{i,1}\}_{i \in \mathcal{I}^{\text{in}}}, \{\xi_i\}_{i \in \mathcal{I}^{\text{out}}} \right) = \mathbb{P} \left(\sum_{i \in \mathcal{I}_{\text{sel}}^{\text{in}}} (a'_{i,\setminus 1} a_{i,\setminus 1}'^T - I_{n-1}) \in \mathcal{C} \mid \mathcal{I}_{\text{sel}}^{\text{in}} \right).$$

As $\{a'_{i,\setminus 1}\}_{i=1}^m$ are i.i.d. Gaussian vectors, Lemma 3.1 implies that for $t \in [0, 1]$:

$$\mathbb{P} \left(\left\| \frac{1}{m} \sum_{i=1}^m (a'_{i,\setminus 1} a'_{i,\setminus 1}^T - I_{n-1}) \mathbf{1}\{i \in \mathcal{I}_{\text{sel}}^{\text{in}}\} \right\|_{\text{op}} \geq C \sqrt{\frac{n}{m}} + t \mid \mathcal{I}_{\text{sel}}^{\text{in}} \right) \leq \exp(-mt)$$

for some constant C . The claim of the lemma follows by integration of the above inequality.

B.3.4 Proof of Lemma B.6

For notational convenience, let us denote the selected outlying indices by $\mathcal{I}_{\text{sel}}^{\text{out}} = \mathcal{I}^{\text{out}} \cap \mathcal{I}_{\text{sel}}$.

We begin our proof by considering Model M1, in which case the proof is essentially identical to Lemma B.5 (see Sec. B.3.3). Indeed, we have the identity

$$\|Z_3 - z_3 I_n\|_{\text{op}} = \left\| \frac{1}{m} \sum_{i=1}^m (a_i a_i^T - I_n) \mathbf{1}\{i \in \mathcal{I}_{\text{sel}}^{\text{out}}\} \right\|_{\text{op}},$$

and so following the completely parallel route of introducing the independent sample $\{a'_i\}_{i=1}^m \stackrel{\text{iid}}{\sim} \mathcal{N}(0, I_n)$, we have for any $t \geq 0$ that

$$\begin{aligned} \mathbb{P}(\|Z_3 - z_3 I_n\|_{\text{op}} \geq t) &= \mathbb{E} \left[\mathbb{P}(\|Z_3 - z_3 I_n\|_{\text{op}} \geq t \mid \mathcal{I}_{\text{sel}}^{\text{out}}) \right] \\ &= \mathbb{E} \left[\mathbb{P} \left(\left\| \frac{1}{m} \sum_{i=1}^m (a'_i a'_i{}^T - I_n) \mathbf{1}\{i \in \mathcal{I}_{\text{sel}}^{\text{out}}\} \right\|_{\text{op}} \geq t \mid \mathcal{I}_{\text{sel}}^{\text{out}} \right) \right]. \end{aligned}$$

Applying Lemma 3.1 to the vectors $a'_i \mathbf{1}\{i \in \mathcal{I}_{\text{sel}}^{\text{out}}\}$ gives the result.

In the case in which we assume Model M2, we have the deterministic upper bound

$$\|Z_3\|_{\text{op}} = \left\| \frac{1}{m} \sum_{i=1}^m a_i a_i^T \mathbf{1}\{i \in \mathcal{I}_{\text{sel}}^{\text{out}}\} \right\|_{\text{op}} \leq \left\| \frac{1}{m} \sum_{i=1}^m a_i a_i^T \mathbf{1}\{i \in \mathcal{I}^{\text{out}}\} \right\|_{\text{op}}.$$

As $a_i \mathbf{1}\{i \in \mathcal{I}^{\text{out}}\}$ is an $O(1)$ -sub-Gaussian vector, Lemma 3.1 implies

$$\mathbb{P} \left(\left\| \frac{1}{m} \sum_{i=1}^m (a_i a_i^T - I_n) \mathbf{1}\{i \in \mathcal{I}^{\text{out}}\} \right\|_{\text{op}} \geq C \sqrt{\frac{n}{m}} + t \mid \mathcal{I}_{\text{sel}}^{\text{out}} \right) \leq \exp(-mt).$$

Since $|\mathcal{I}^{\text{out}}|/m = p_{\text{fail}}$, the desired claim follows via the triangle inequality.

C Technical results

C.1 Proof of Lemma 3.2

Let $p(z)$ denote the density of z and $p_c(z') = p(z') / \int_{-c}^c p(z) dz$ for $z' \in [-c, c]$. Let $F_c(z) = \int_{-c}^z p_c(z') dz'$ be the CDF of p_c , and define the mapping $T : [-c, c] \rightarrow [-c, c]$ by

$$T(u) = F_c^{-1} \left(\frac{u + c}{2c} \right) = z \text{ for the } z \text{ s.t. } F_c(z) = \frac{u + c}{2c}.$$

Evidently, for $Z \sim P_c$ we have $T(Z) \sim \text{Uni}[-c, c]$, and by the symmetry and monotonicity properties of p and p_c we have $|T(z)| \geq |z|$ for $z \in [-c, c]$. In particular, letting $U \sim \text{Uni}[-c, c]$, we have

$$\text{Var}(U) = \mathbb{E}[U^2] = \mathbb{E}[T(Z)^2 \mid |Z| \leq c] \geq \mathbb{E}[Z^2 \mid |Z| \leq c].$$

Using that $\text{Var}(U) = \frac{1}{2c} \int_{-c}^c u^2 du = \frac{c^2}{3}$ gives the result.

C.2 Proof of Lemma B.1

The proof is an essentially standard concentration and covering number argument, with a few minor wrinkles. First, we note that

$$\text{Var}(|\langle u, a_i \rangle \langle v, a_i \rangle|) \leq \mathbb{E}[|\langle u, a_i \rangle \langle v, a_i \rangle|^2] \stackrel{(i)}{\leq} \sup_{u \in \mathbb{S}^{n-1}} \mathbb{E}[\langle u, a_i \rangle^4] \leq 2e\sigma^4,$$

where inequality (i) is Cauchy-Schwarz and the final inequality follows Lemma 3.1. Thus, the lower tail Bernstein's inequality (Lemma A.2) applied to the positive random variables $|\langle u, a_i \rangle \langle v, a_i \rangle| \geq 0$ with variance bounded by $2e\sigma^4$ implies that

$$\mathbb{P}\left(\frac{1}{m} \sum_{i=1}^m |\langle u, a_i \rangle \langle v, a_i \rangle| - \kappa_{\text{st}}(u, v) \leq -t\right) \leq \exp\left(-\frac{mt^2}{4e\sigma^4}\right). \quad (47)$$

Define $Z_{u,v} := \frac{1}{m} \sum_{i=1}^m |\langle u, a_i \rangle \langle v, a_i \rangle|$ for shorthand. Using that for vectors $w, x \in \mathbb{R}^n$ we have $\sum_{i=1}^m |\langle a_i, w \rangle \langle a_i, x \rangle| \leq \|Aw\|_2 \|Ax\|_2$ by Cauchy-Schwarz, we see that for any $u, v, u', v' \in \mathbb{S}^{n-1}$,

$$\begin{aligned} |Z_{u,v} - Z_{u',v'}| &\leq \frac{1}{m} \sum_{i=1}^m |\langle a_i, u \rangle \langle a_i, v \rangle - \langle a_i, u' \rangle \langle a_i, v' \rangle| \\ &\leq \frac{1}{m} \sum_{i=1}^m (|\langle a_i, u - u' \rangle \langle a_i, v \rangle| + |\langle a_i, u' \rangle \langle a_i, v - v' \rangle|) \\ &\leq \frac{1}{m} \|A(u - u')\|_2 \|Av\|_2 + \frac{1}{m} \|A(v - v')\|_2 \|Au\|_2 \leq \frac{\|A\|_{\text{op}}^2}{m} (\|u - u'\|_2 + \|v - v'\|_2). \end{aligned}$$

Now, let \mathcal{N} be an ϵ -cover of the sphere \mathbb{S}^{n-1} , which we use to control $\inf_{u,v \in \mathbb{S}^{n-1}} \{Z_{u,v} - \mathbb{E}[Z_{u,v}]\}$. Using the previous display, we obtain by the definition of an ϵ -cover argument that

$$\mathbb{P}\left(\inf_{u,v \in \mathbb{S}^{n-1}} \{Z_{u,v} - \kappa_{\text{st}}(u, v)\} \leq -t\right) \leq \mathbb{P}\left(\min_{u,v \in \mathcal{N}} \{Z_{u,v} - \kappa_{\text{st}}(u, v)\} \leq -t + \frac{2\|A^T A\|_{\text{op}}}{m} \epsilon\right).$$

From Lemma 3.1, we know that $\|A^T A\|_{\text{op}}$ is well-concentrated, and thus by considering the event that $\|A^T A\|_{\text{op}} \gtrsim m\sigma^2$ that for some numerical constant $C < \infty$ we have

$$\begin{aligned} &\mathbb{P}\left(\inf_{u,v \in \mathbb{S}^{n-1}} \{Z_{u,v} - \kappa_{\text{st}}(u, v)\} \leq -t\right) \\ &\leq \mathbb{P}\left(\min_{u,v \in \mathcal{N}} \{Z_{u,v} - \kappa_{\text{st}}(u, v)\} \leq -t + 2C\sigma^2\epsilon\right) + \mathbb{P}\left(\|A^T A\|_{\text{op}} \geq Cm\sigma^2\right) \\ &\leq \text{card}(\mathcal{N})^2 \exp\left(-\frac{cm[t - C\sigma^2\epsilon]_+^2}{\sigma^4}\right) + \exp(-m), \end{aligned} \quad (48)$$

where the first term comes from Bernstein's inequality (47) and the second from Lemma 3.1.

Now, let us assume that \mathcal{N} is an ϵ -cover of \mathbb{S}^{n-1} with minimal cardinality, which by standard volume arguments [44, Lemma 2] satisfies $N(\epsilon) := \text{card}(\mathcal{N}) \leq (1 + 2/\epsilon)^n$ for $\epsilon > 0$. Noting that $(1 + 2/\epsilon)^{2n} \leq \exp(n/\epsilon)$ for $\epsilon \leq \epsilon_0 := .21398$, we may replace inequality (48) with

$$\mathbb{P}\left(\inf_{u,v \in \mathbb{S}^{n-1}} \{Z_{u,v} - \kappa_{\text{st}}(u, v)\} \leq -t\right) \leq \exp\left(\frac{n}{\epsilon} - \frac{cm[t - C\sigma^2\epsilon]_+^2}{\sigma^4}\right) + e^{-m}$$

valid for all $\epsilon \leq \epsilon_0$. Now, if we set $\epsilon = \sqrt[3]{n/m}$ and define $t = \sigma^2 \bar{t} + (C+1)\sigma^2 \epsilon$, then we find that if $m/n \geq \epsilon_0^{-3}$, we have

$$\mathbb{P} \left(\inf_{u,v \in \mathbb{S}^{n-1}} \{Z_{u,v} - \kappa_{\text{st}}(u,v)\} \leq -\sigma^2 \bar{t} - (C+1)\sigma^2 \sqrt[3]{\frac{n}{m}} \right) \leq \exp \left(cm \bar{t}^2 \right) + e^{-m}.$$

This is the desired result.

C.3 Properties of Gaussian Random Variable

Lemma C.1. *Let $W \sim \mathcal{N}(0,1)$. Then for all $c \in \mathbb{R}_+$*

$$\mathbb{E} [W^2 \mid W^2 \leq c] \leq 1 - \frac{1}{2}c\mathbb{P}(W^2 > c).$$

Proof By the law of total probability, we have

$$\mathbb{E}[W^2] = \mathbb{E}[W^2 \mid W^2 \leq c]\mathbb{P}(W^2 \leq c) + \mathbb{E}[W^2 \mid W^2 > c]\mathbb{P}(W^2 > c).$$

Using that $\mathbb{E}[W^2] = 1$ yields

$$1 - \mathbb{E}[W^2 \mid W^2 \leq c] = \mathbb{P}(W^2 > c) (\mathbb{E}[W^2 \mid W^2 > c] - \mathbb{E}[W^2 \mid W^2 \leq c]).$$

For $t \in [0, c]$, we have $\mathbb{P}(W^2 \in [c-t, c]) \leq \mathbb{P}(W^2 \in [0, t])$, so that for such t we have $\mathbb{P}(W^2 \geq c-t \mid W^2 \leq c) \leq \mathbb{P}(W^2 \leq t \mid W^2 \leq c)$, or $\mathbb{P}(W^2 \geq t \mid W^2 \leq c) + \mathbb{P}(W^2 \geq c-t \mid W^2 \leq c) \leq 1$. Performing the standard change of variables to compute $\mathbb{E}[W^2 \mid W^2 \leq c]$, we thus obtain

$$\begin{aligned} \mathbb{E}[W^2 \mid W^2 \leq c] &= \int_0^c \mathbb{P}(W^2 \geq t \mid W^2 \leq c) dt \\ &= \int_0^c \frac{1}{2} [\mathbb{P}(W^2 \geq t \mid W^2 \leq c) + \mathbb{P}(W^2 \geq c-t \mid W^2 \leq c)] dt \\ &\leq \int_0^c \frac{1}{2} dt = \frac{c}{2}. \end{aligned}$$

Using that $\mathbb{E}[W^2 \mid W^2 > c] \geq c$ gives the lemma. \square

Lemma C.2. *Let $X \in \mathbb{C}^{n \times n}$ be rank-2 and Hermitian and $z = \frac{1}{\sqrt{2}}(\mathcal{N}(0, I_n) + i\mathcal{N}(0, I_n))$ be standard complex normal. Then $\mathbb{E}[|z^H X z|] \geq \frac{2\sqrt{2}}{\pi} \|X\|_{\text{Fr}}$.*

Proof Without loss of generality, we assume that $\|X\|_{\text{Fr}} = 1$, so that $X = \lambda_1 u_1 u_1^H + \lambda_2 u_2 u_2^H$ for $\lambda_1^2 + \lambda_2^2 = 1$ and u_i orthonormal. Thus, if Z_1, Z_2 are standard normal, we have

$$z^H X z = \lambda_1 |\langle z, u_1 \rangle|^2 + \lambda_2 |\langle z, u_2 \rangle|^2 \stackrel{\text{dist}}{=} \lambda_1 Z_1^2 + \lambda_2 Z_2^2.$$

Without loss of generality, we assume $\lambda_1 \geq 0$, so that by inspection we have for $\lambda = \lambda_1$ that $\mathbb{E}[\lambda_1 Z_1^2 + \lambda_2 Z_2^2] \geq \mathbb{E}[\lambda Z_1^2 - \sqrt{1-\lambda^2} Z_2^2]$. Letting $f(\lambda) = \mathbb{E}[\lambda Z_1^2 - \sqrt{1-\lambda^2} Z_2^2]$, we have that

$$f'(\lambda) = \mathbb{P} \left(\frac{\lambda}{\sqrt{1-\lambda^2}} Z_1^2 \geq Z_2^2 \right) - \mathbb{P} \left(\frac{\lambda}{\sqrt{1-\lambda^2}} Z_1^2 \leq Z_2^2 \right) = \begin{cases} > 0 & \text{if } \lambda > \frac{1}{\sqrt{2}} \\ = 0 & \text{if } \lambda = \frac{1}{\sqrt{2}} \\ < 0 & \text{if } \lambda < \frac{1}{\sqrt{2}}. \end{cases}$$

In particular, f is quasi-convex and minimized at $\lambda = 1/\sqrt{2}$, and thus

$$\frac{1}{\|X\|_{\text{Fr}}} \mathbb{E}[|z^H X z|] \geq \frac{1}{\sqrt{2}} \mathbb{E}[|Z_1^2 - Z_2^2|] = \frac{2\sqrt{2}}{\pi},$$

where the final equality is a calculation. □

References

- [1] S. Balakrishnan, M. J. Wainwright, and B. Yu. Statistical guarantees for the EM algorithm: From population to sample-based analysis. *Annals of Statistics*, 45(1):77–120, 2017.
- [2] R. Balan, P. Casazza, and D. Edidin. On signal reconstruction without phase. *Applied and Computational Harmonic Analysis*, 20:345–356, 2006.
- [3] K. Ball. An elementary introduction to modern convex geometry. In S. Levy, editor, *Flavors of Geometry*, pages 1–58. MSRI Publications, 1997.
- [4] A. Bandeira, J. Cahill, D. Mixon, and A. Nelson. Saving phase: Injectivity and stability for phase retrieval. *Applied and Computational Harmonic Analysis*, 37(1):106–125, 2014.
- [5] P. L. Bartlett and S. Mendelson. Rademacher and Gaussian complexities: Risk bounds and structural results. *Journal of Machine Learning Research*, 3:463–482, 2002.
- [6] D. P. Bertsekas. Approximation procedures based on the method of multipliers. *Journal of Optimization Theory and Applications*, 23(4):487–510, 1977.
- [7] J. Burke. Descent methods for composite nondifferentiable optimization problems. *Mathematical Programming*, 33:260–279, 1985.
- [8] J. Burke and M. Ferris. A Gauss-Newton method for convex composite optimization. *Mathematical Programming*, 71:179–194, 1995.
- [9] E. Candès, T. Strohmer, and V. Voroninski. PhaseLift: exact and stable signal recovery from magnitude measurements via convex programming. *Communications on Pure and Applied Mathematics*, 66(8), 2013.
- [10] E. J. Candès and X. Li. Solving quadratic equations via PhaseLift when there are about as many equations as unknowns. *Foundations of Computational Mathematics*, 14(5):1017–1026, 2014.
- [11] E. J. Candès and Y. Plan. Tight oracle inequalities for low-rank matrix recovery from a minimal number of noisy random measurements. *IEEE Transactions on Information Theory*, 57(4):2342–2359, 2011.
- [12] E. J. Candès, Y. Eldar, T. Strohmer, and V. Voroninski. Phase retrieval via matrix completion. *SIAM Journal on Imaging Sciences*, 6(1):199–225, 2013.
- [13] E. J. Candès, X. Li, and M. Soltanolkotabi. Phase retrieval via Wirtinger flow: Theory and algorithms. *IEEE Transactions on Information Theory*, 61(4):1985–2007, 2015.

- [14] A. Chai, M. Moscoso, and G. Papanicolaou. Array imaging using intensity-only measurements. *Inverse Problems*, 27(1):015005, 2011.
- [15] Y. Chen and E. Candès. Solving random quadratic systems of equations is nearly as easy as solving linear systems. *Communications on Pure and Applied Mathematics*, To appear, 2015.
- [16] D. Drusvyatskiy and A. Lewis. Error bounds, quadratic growth, and linear convergence of proximal methods. *Mathematics of Operations Research*, To appear, 2018.
- [17] D. Drusvyatskiy and C. Paquette. Efficiency of minimizing compositions of convex functions and smooth maps. *arXiv:1605.00125 [math.OC]*, 2016.
- [18] D. Drusvyatskiy, A. Ioffe, and A. Lewis. Nonsmooth optimization using Taylor-like models: error bounds, convergence, and termination criteria. *arXiv:1610.03446 [math.OC]*, 2016. URL <https://arxiv.org/abs/1610.03446>.
- [19] Y. Eldar and S. Mendelson. Phase retrieval: Stability and recovery guarantees. *Applied and Computational Harmonic Analysis*, 36(3):473–494, 2014.
- [20] M. Fickus, D. G. Mixon, A. A. Nelson, and Y. Wang. Phase retrieval from very few measurements. *Linear Algebra and its Applications*, 449:475–499, 2014.
- [21] J. R. Fienup. Reconstruction of an object from the modulus of its Fourier transform. *Optics Letters*, 3(1):27–29, 1978.
- [22] J. R. Fienup. Phase retrieval algorithms: a comparison. *Applied Optics*, 21(15):2758–2769, 1982.
- [23] R. Fletcher and G. A. Watson. First and second order conditions for a class of nondifferentiable optimization problems. *Mathematical Programming*, 18:291–307, 1980.
- [24] R. W. Gerchberg and W. O. Saxton. A practical algorithm for the determination of phase from image and diffraction. *Optik*, 35:237–246, 1972.
- [25] T. Hastie, R. Tibshirani, and J. Friedman. *The Elements of Statistical Learning*. Springer, second edition, 2009.
- [26] J. Hiriart-Urruty and C. Lemaréchal. *Convex Analysis and Minimization Algorithms I*. Springer, New York, 1993.
- [27] P. J. Huber and E. M. Ronchetti. *Robust Statistics*. John Wiley and Sons, second edition, 2009.
- [28] R. H. Keshavan, A. Montanari, and S. Oh. Matrix completion from a few entries. *IEEE Transactions on Information Theory*, 56(6):2980–2998, 2010.
- [29] K. Kreutz-Delgado. The complex gradient operator and the \mathbb{CR} -calculus. *arXiv:0906.4835 [math.OC]*, 2009.
- [30] M. Ledoux. *The Concentration of Measure Phenomenon*. American Mathematical Society, 2001.
- [31] P.-L. Loh and M. J. Wainwright. Regularized M-estimators with nonconvexity: Statistical and algorithmic theory for local optima. *Journal of Machine Learning Research*, 16:559–616, 2013.

- [32] S. Mendelson. Learning without concentration. In *Proceedings of the Twenty Seventh Annual Conference on Computational Learning Theory*, 2014.
- [33] P. Netrapalli, P. Jain, and S. Sanghavi. Phase retrieval using alternating minimization. In *Advances in Neural Information Processing Systems 26*, 2013.
- [34] N. Parikh and S. Boyd. Proximal algorithms. *Foundations and Trends in Optimization*, 1(3): 123–231, 2013.
- [35] N. Parikh and S. Boyd. Block splitting for distributed optimization. *Mathematical Programming Computation*, 6:77–102, 2014.
- [36] F. Pi, H. Zhang, and P. Guo. RNA nanoparticles in cancer cells. National Cancer Institute Visuals Online, 2016. URL <https://visualsonline.cancer.gov/details.cfm?imageid=11167>. Accessed April 1, 2017.
- [37] B. T. Polyak. On the Bertsekas’ method for minimization of composite functions. In A. Bensoussan and J. L. Lions, editors, *International Symposium on Systems Optimization and Analysis*, volume 14 of *Lecture Notes in Control and Information Sciences*, pages 178–186. Springer, 1979.
- [38] Y. Schechtman, Y. C. Eldar, O. Cohen, H. N. Chapman, J. Miao, and M. Segev. Phase retrieval with application to optical imaging. *IEEE Signal Processing Magazine*, pages 87–109, May 2015.
- [39] G. W. Stewart and J.-G. Sun. *Matrix Perturbation Theory*. Academic Press, 1990.
- [40] L. N. Trefethen and D. Bau III. *Numerical Linear Algebra*. SIAM, 1997.
- [41] J. A. Tropp. Improved analysis of the subsampled randomized Hadamard transform. *Advances in Adaptive Data Analysis*, 3(1–2):115–126, 2011. Special issue on Sparse Representation of Data and Images.
- [42] M. Udell, K. Mohan, D. Zeng, J. Hong, S. Diamond, and S. Boyd. Convex optimization in Julia. In *First Workshop on High Performance Technical Computing in Dynamic Languages*, pages 18–28. IEEE, 2014.
- [43] A. W. van der Vaart and J. A. Wellner. *Weak Convergence and Empirical Processes: With Applications to Statistics*. Springer, New York, 1996.
- [44] R. Vershynin. Introduction to the non-asymptotic analysis of random matrices. In *Compressed Sensing: Theory and Applications*, chapter 5, pages 210–268. Cambridge University Press, 2012.
- [45] I. Waldspurger, A. d’Aspremont, and S. Mallat. Phase recovery, maxcut and complex semidefinite programming. *Mathematical Programming, Series A*, 149(1–2):47–81, 2015.
- [46] G. Wang, G. B. Giannakis, and Y. C. Eldar. Solving systems of random quadratic equations via truncated amplitude flow. *arXiv:1605.08285 [stat.ML]*, 2016.
- [47] H. Zhang, Y. Chi, and Y. Liang. Provable non-convex phase retrieval with outliers: Median truncated Wirtinger flow. In *Proceedings of the 33rd International Conference on Machine Learning*, 2016.