

Exact solution of the Schwarzian theory

Vladimir V. Belokurov*

*Lomonosov Moscow State University, Leninskie gory 1, Moscow, 119991,
Russia and Institute for Nuclear Research of the Russian Academy of Sciences,
60th October Anniversary Prospect 7a, Moscow, 117312, Russia*

Evgeniy T. Shavgulidze†

Lomonosov Moscow State University, Leninskie gory 1, Moscow, 119991, Russia

The explicit evaluation of the partition function in the Schwarzian theory is presented.

The Schwarzian theory [1] is the basic element of various physical models including the SYK model and the two-dimensional dilaton gravity (see, e.g., [2], [3], [4], [5], and references therein). The action of the theory is

$$I = -\frac{1}{g^2} \int_0^{2\pi} \left[\mathcal{S}_\phi(\tau) + \frac{1}{2}(\phi')^2(\tau) \right] d\tau. \quad (1)$$

Here,

$$\mathcal{S}_\phi(t) \equiv \left(\frac{\phi''(t)}{\phi'(t)} \right)' - \frac{1}{2} \left(\frac{\phi''(t)}{\phi'(t)} \right)^2 \quad (2)$$

is the Schwarzian derivative, $\phi \in \text{Diff}^3([0, 2\pi])$, and $\phi'(0) = \phi'(2\pi)$.

It is convenient to rewrite the action in the form

$$I = -\frac{1}{\sigma^2} \int_0^1 [\mathcal{S}_\varphi(t) + 2\pi^2 \dot{\varphi}^2(t)] dt, \quad (3)$$

where

$$\sigma = \sqrt{2\pi}g, \quad t = \frac{1}{2\pi}\tau, \quad \varphi(t) = \frac{1}{2\pi}\phi(\tau),$$

$$\varphi \in \text{Diff}^3([0, 1]), \quad \dot{\varphi}(0) = \dot{\varphi}(1).$$

The functional integral for the partition function

$$Z(g) = \int_{\dot{\varphi}(0)=\dot{\varphi}(1)} \exp\{-I\} d\varphi \quad (4)$$

diverges [1]. However, as we will see later on (eq. (17)), the integral

$$Z_\alpha(g)$$

$$= \int_{\text{Diff}^3([0,1])} \exp\{-I\} \exp\left\{ \frac{-2[\pi^2 - \alpha^2]}{\sigma^2} \int_0^1 \dot{\varphi}^2(t) dt \right\} d\varphi$$

$$= \int_{\dot{\varphi}(0)=\dot{\varphi}(1)} \exp\left\{ \frac{1}{\sigma^2} \int_0^1 [\mathcal{S}_\varphi(t) + 2\alpha^2 \dot{\varphi}^2(t)] dt \right\} d\varphi \quad (5)$$

converges for $0 \leq \alpha < \pi$. Therefore, let us evaluate the integral (5) first.

The measure

$$\mu_\sigma(X) = \int_X \exp\left\{ \frac{1}{\sigma^2} \int_0^1 \mathcal{S}_\varphi(t) dt \right\} d\varphi \quad (6)$$

is quasi-invariant, and the Radon - Nikodym derivative of the measure is [6], [7]

$$\frac{d\mu_\sigma^f}{d\mu_\sigma}(\varphi) = \frac{1}{\sqrt{\dot{f}(0)\dot{f}(1)}}$$

$$\times \exp\left\{ \frac{1}{\sigma^2} \left[\frac{\ddot{f}(0)}{\dot{f}(0)} \dot{\varphi}(0) - \frac{\ddot{f}(1)}{\dot{f}(1)} \dot{\varphi}(1) \right] + \frac{1}{\sigma^2} \int_0^1 \mathcal{S}_f(\varphi(t)) \dot{\varphi}^2 dt \right\}, \quad (7)$$

where

$$\mu_\sigma^f(X) = \mu_\sigma(f \circ X).$$

Here, we have used the well known property of the Schwarzian derivative:

$$\mathcal{S}_{f \circ \varphi}(t) = \mathcal{S}_f(\varphi(t)) \dot{\varphi}^2(t) + \mathcal{S}_\varphi(t), \quad (f \circ \varphi)(t) = f(\varphi(t)).$$

Take the function f to be

$$f(t) = \frac{1}{2} \left[\frac{1}{\tan \frac{\alpha}{2}} \tan \left(\alpha(t - \frac{1}{2}) \right) + 1 \right]. \quad (8)$$

In this case,

$$\mathcal{S}_f(t) = 2\alpha^2, \quad \dot{f}(0) = \dot{f}(1) = \frac{\alpha}{\sin \alpha},$$

$$-\frac{\ddot{f}(0)}{\dot{f}(0)} = -\frac{\ddot{f}(1)}{\dot{f}(1)} = 2\alpha \tan \frac{\alpha}{2}. \quad (9)$$

Now we have the following functional integrals equality:

$$\begin{aligned} & \frac{\alpha}{\sin \alpha} \int_{\dot{\varphi}(0)=\dot{\varphi}(1)} F(\varphi) \mu_{\sigma}(d\varphi) \\ &= \int_{\dot{\varphi}(0)=\dot{\varphi}(1)} F(f(\varphi)) \exp \left\{ -\frac{4\alpha}{\sigma^2} \tan \frac{\alpha}{2} \dot{\varphi}(0) \right\} \\ & \times \exp \left\{ \frac{1}{\sigma^2} \int_0^1 [\mathcal{S}_{\varphi}(t) + 2\alpha^2 \dot{\varphi}^2(t)] dt \right\} d\varphi. \end{aligned} \quad (10)$$

The next step is the choice of the function F . Let it be

$$F(f(\varphi)) = \exp \left\{ \frac{4\alpha}{\sigma^2} \tan \frac{\alpha}{2} \dot{\varphi}(0) \right\}. \quad (11)$$

To find $F(\varphi)$ from the previous equation, note that for $u(t) = f(\varphi(t))$

$$\dot{\varphi}(0) = \frac{1}{\dot{f}(0)} \dot{u}(0).$$

Then

$$F(u) = \exp \left\{ \frac{8 \sin^2 \frac{\alpha}{2}}{\sigma^2} \dot{u}(0) \right\}. \quad (12)$$

Thus for the regularized partition function we have

$$\begin{aligned} & Z_{\alpha}(g) \\ &= \frac{\alpha}{\sin \alpha} \int_{\dot{\varphi}(0)=\dot{\varphi}(1)} \exp \left\{ \frac{8 \sin^2 \frac{\alpha}{2}}{\sigma^2} \dot{\varphi}(0) \right\} \mu_{\sigma}(d\varphi). \end{aligned} \quad (13)$$

Under the substitution

$$\varphi(t) = \frac{\int_0^t \exp\{\xi(\eta)\} d\eta}{\int_0^1 \exp\{\xi(\eta)\} d\eta}, \quad (14)$$

the measure $\mu_{\sigma}(d\varphi)$ turns into the Wiener measure [6], [7]

$$w_{\sigma}(d\xi) = \exp \left\{ -\frac{1}{2\sigma^2} \int_0^1 \dot{\xi}^2(t) dt \right\} d\xi. \quad (15)$$

In this case,

$$\xi(t) = \ln \dot{\varphi}(t) - \ln \dot{\varphi}(0), \quad \xi \in C([0, 1]), \quad (16)$$

and $\xi(0) = \xi(1) = 0$.

Now $Z_{\alpha}(g)$ is written as

$$\begin{aligned} & Z_{\alpha}(g) = \frac{\alpha}{\sin \alpha} \\ & \times \int_{\xi(0)=\xi(1)=0} \exp \left\{ \frac{8 \sin^2 \frac{\alpha}{2}}{\sigma^2} \frac{1}{\int_0^1 \exp\{\xi(\eta)\} d\eta} \right\} w_{\sigma}(d\xi). \end{aligned} \quad (17)$$

The singularity at $\alpha = \pi$ is canceled out in the ratio

$$\frac{Z_{\alpha}(g)}{Z_{\alpha}(1)},$$

and we can remove the regularization there

$$\frac{Z(g)}{Z(1)} = \lim_{\alpha \rightarrow \pi-0} \frac{Z_{\alpha}(g)}{Z_{\alpha}(1)}. \quad (18)$$

To evaluate the integral

$$\int_{\xi(0)=\xi(1)=0} \exp \left\{ \frac{8}{\sigma^2} \frac{1}{\int_0^1 \exp\{\xi(\eta)\} d\eta} \right\} w_{\sigma}(d\xi), \quad (19)$$

we use the following equation:

$$\begin{aligned} & \int_{\xi(0)=\xi(1)=0} \exp \left\{ \frac{-2\beta^2}{\sigma^2(\beta+1)} \frac{1}{\int_0^1 \exp\{\xi(\eta)\} d\eta} \right\} w_{\sigma}(d\xi) \\ &= \frac{1}{\sqrt{2\pi}\sigma} \exp \left\{ -\frac{4(\ln(\beta+1))^2}{2\sigma^2} \right\}. \end{aligned} \quad (20)$$

The proof of the more general formula will be given in the forthcoming paper (also, see [7]). For the integral (19)

$$(\beta+1) = -1.$$

Thus the final result is

$$\frac{Z(g)}{Z(1)} = \frac{\exp\{-\pi\}}{g} \exp \left\{ \frac{\pi}{g^2} \right\}. \quad (21)$$

It is interesting to compare the one-loop results for $Z(g)$ in [1] with the eq. (21). Note that the power of the constant g in the denominator is determined by the number of gauge fixing conditions. The one-loop result for the orbit $Diff(S^1)/U(1)$ (eq. (3.45) in [1]) has the same form as our exact result (21).

Unlike its compact subgroup $U(1)$, the group $SL(2, \mathbf{R})$ is noncompact. Therefore, integrating over the quotient

space $Diff^3([0, 1])/SL(2, \mathbf{R})$ we get the finite result for the partition function in the Schwarzian theory.

We define the Schwarzian partition function as a limit

$$Z_{Schw}(g) = \lim_{\alpha \rightarrow \pi-0} \frac{Z_\alpha(g)}{V_\alpha(g)}. \quad (22)$$

Here, $Z_\alpha(g)$ is given by the eqs. (5), (17), and $V_\alpha(g)$ is the regularized volume of the group $SL(2, \mathbf{R})$

$$V_\alpha(g) = \int_{SL(2, \mathbf{R})} \exp \left\{ -\frac{2[\pi^2 - \alpha^2]}{\sigma^2} \int_0^1 \dot{\varphi}^2(t) dt \right\} d\mu_H. \quad (23)$$

Note that the functional measure in the eq. (5) and the Haar measure $d\mu_H$ on the group $SL(2, \mathbf{R})$ in the eq. (23) are regularized in the same manner.

To perform the integration over the group $SL(2, \mathbf{R})$ in the eq. (23) we choose the representation [8]

$$\varphi_z(t) = -\frac{i}{2\pi} \ln \frac{e^{i2\pi t} + z}{\bar{z} e^{i2\pi t} + 1}, \quad z = \rho e^{i\theta}, \quad \rho < 1. \quad (24)$$

In this case, the Haar measure is [8]

$$\mu_H(dz) = \frac{4\rho d\rho d\theta}{(1 - \rho^2)^2}. \quad (25)$$

The integral

$$\int_0^1 \dot{\varphi}_z^2(t) dt = \int_0^1 \frac{(1 - |z|^2)^2 dt}{(e^{i2\pi t} + z)^2 (e^{-i2\pi t} + \bar{z})^2} = \frac{1 + \rho^2}{1 - \rho^2} \quad (26)$$

does not depend on θ . And the regularized volume of the group has the form

$$V_\alpha(g) = \int_0^1 \exp \left\{ -\frac{[\pi^2 - \alpha^2]}{\pi g^2} \frac{(1 + \rho^2)}{(1 - \rho^2)} \right\} \frac{8\pi \rho d\rho}{(1 - \rho^2)^2}$$

$$= \exp \left\{ -\frac{[\pi^2 - \alpha^2]}{\pi g^2} \right\} \frac{4\pi^2 g^2}{[\pi^2 - \alpha^2]}. \quad (27)$$

Thus we can evaluate the Schwarzian partition function

$$Z_{Schw}(g) = \frac{1}{4\pi g^3} \exp \left\{ \frac{\pi}{g^2} \right\}. \quad (28)$$

Note that the one-loop result in [1], [2] has the same form as the exact partition function (28) obtained by the direct functional integration.

* vvelokurov@yandex.ru

† shavgulidze@bk.ru

- [1] D. Stanford and E. Witten, "Fermionic Localization of the Schwarzian Theory", arXiv:1703.04612v1 [hep-th].
- [2] J. Maldacena and D. Stanford, "Remarks on the Sachdev-Ye-Kitaev model", Phys. Rev. D **94**, 106002 (2016), arXiv:1604.07818 [hep-th].
- [3] D. Bagrets, A. Altland and A. Kamenev, "Sachdev-Ye-Kitaev Model as Liouville Quantum Mechanics", Nucl.Phys. B **911**, 191 (2016), arXiv:1607.00694 [cond-mat.str-el].
- [4] D. J. Gross and V. Rosenhaus, "A Generalization of Sachdev-Ye-Kitaev", JHEP **02**, 093 (2017), arXiv:1610.01569 [hep-th].
- [5] G. Mandal, P. Nayak and S. R. Wadia, "Coadjoint orbit action of Virasoro group and two-dimensional quantum gravity dual SYK/tensor models", arXiv:1702.04266v2 [hep-th].
- [6] E.T. Shavgulidze, "Some properties of quasi-invariant measures on groups of diffeomorphisms of the circle", Russ. J. Math. Phys. **7**, 464 (2000).
- [7] V.V. Belokurov and E.T. Shavgulidze, "Extraordinary Properties of Functional Integrals and Groups of Diffeomorphisms", Phys. of Part. and Nuclei, **48**, 267 (2017) [Fiz. Elem. Chastits At. Yadra **48**, 194 (2017)].
- [8] S. Lang, "SL₂($\mathbf{R}$)". Addison-Wesley Publishing. 1975.