

Typical ranks of semi-tall real 3-tensors

Toshio SUMI*, Mitsuhiro MIYAZAKI† and Toshio SAKATA‡

Abstract

Let m , n and p be integers with $3 \leq m \leq n$ and $(m-1)(n-1) + 1 \leq p \leq (m-1)m$. We showed in previous papers that if $p \geq (m-1)(n-1) + 2$, then typical ranks of $p \times n \times m$ -tensors over the real number field are p and $p+1$ if and only if there exists a nonsingular bilinear map $\mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}^{mn-p}$. We also showed that the “if” part also valid in the case where $p = (m-1)(n-1) + 1$. In this paper, we consider the case where $p = (m-1)(n-1) + 1$ and show that the typical ranks of $p \times n \times m$ -tensors over the real number field are p and $p+1$ in several cases including the case where there is no nonsingular bilinear map $\mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}^{mn-p}$. In particular, we show that the “only if” part of the above mentioned fact does not valid for the case $p = (m-1)(n-1) + 1$.

Keywords: tensor rank, typical rank, tall tensor, semi-tall tensor, Bezout’s theorem, determinantal variety

MSC:15A69, 14P10, 14M12, 13C40

1 Introduction

Tensor rank is a subject which is widely studied in both pure and applied mathematics. A high dimensional array of datum is called a tensor in the field of data analysis. Precisely, let $N_1, \dots, N_d$ be positive integers. A d -dimensional array datum $T = (t_{i_1 \dots i_d})_{1 \leq i_j \leq N_j, 1 \leq j \leq d}$ is called a d -way tensor or simply a d -tensor of format $N_1 \times \dots \times N_d$. For a set S , the set of $N_1 \times \dots \times N_d$ tensors with entries in S is denoted by $S^{N_1 \times \dots \times N_d}$.

Let K be a field and V_j an N_j -dimensional vector space over K with fixed basis $v_{j1}, \dots, v_{jN_j}$ for $1 \leq j \leq d$. Then there is a one to one correspondence between $K^{N_1 \times \dots \times N_d}$ and $V_1 \otimes \dots \otimes V_d$ by $(m_{i_1 \dots i_d}) \leftrightarrow \sum_{i_1 \dots i_d} m_{i_1 \dots i_d} v_{1i_1} \otimes \dots \otimes$

*Faculty of Arts and Science, Kyushu University, Fukuoka, Japan

†Department of Mathematics, Kyoto University of Education, Kyoto, Japan

‡Emeritus professor, Kyushu University, Fukuoka, Japan

v_{di_d} . A non-zero tensor corresponding to an element of $V_1 \otimes \cdots \otimes V_d$ of the form $\mathbf{a}_1 \otimes \cdots \otimes \mathbf{a}_d$ is called a rank 1 tensor. For a general tensor T of format $N_1 \times \cdots \times N_d$, the rank of T , denoted by $\text{rank} T$, is by definition the minimum integer r such that T can be expressed as a sum of r rank 1 tensors, where we set the empty sum to be zero. Thus, the rank is a measure of the complexity of a tensor. Further, for a 2-tensor, i.e., a matrix, the rank is identical with the one defined in linear algebra.

However, for the case where $d \geq 3$, the behavior of rank is much more complicated than the matrix case. In the matrix case, the rank is the maximum size of non-zero minors. Thus, if K is an infinite field, the set of $m \times n$ matrices with rank $\min\{m, n\}$ form a Zariski dense open subset of $K^{m \times n}$. However, there are non-empty Euclidean open subsets of $\mathbb{R}^{2 \times 2 \times 2}$ such that one consists of rank 2 tensors and the other one consists of rank 3 tensors. In particular, it is not possible to characterize the rank of a tensor by vanishing and/or non-vanishing of polynomials.

Let m, n, p be positive integers. If the set of rank r tensors of format $p \times n \times m$ over $\mathbb{R}$ contains a non-empty Euclidean open subset of $\mathbb{R}^{p \times n \times m}$, we say that r is a typical rank of $p \times n \times m$ tensors over $\mathbb{R}$. The set of typical ranks of $p \times n \times m$ tensors over $\mathbb{R}$ is denoted as $\text{trank}_{\mathbb{R}}(p, n, m)$ or simply $\text{trank}(p, n, m)$.

If the base field is $\mathbb{C}$, the set of tensors of format $p \times n \times m$ with rank at most r contains a non-empty Zariski open set if and only if its Zariski closure is $\mathbb{C}^{p \times n \times m}$ (cf., Chevalley's Theorem, see e.g., [Har92, p. 39]). Therefore, there exists exactly one "typical rank of $p \times n \times m$ tensors over $\mathbb{C}$ ". This is called the generic rank of $p \times n \times m$ tensors over $\mathbb{C}$ and denoted as $\text{grank}_{\mathbb{C}}(p, n, m)$ or simply $\text{grank}(p, n, m)$.

It is fairly easy to show that $\text{grank}(p, n, m) = \min \text{trank}(p, n, m)$ (see e.g., [SSM16, Chapter 6]). Further $r \geq \text{grank}(p, n, m)$ if and only if the r -th higher secant variety of the image of Segre embedding $\mathbb{P}_{\mathbb{C}}\mathbb{C}^p \times \mathbb{P}_{\mathbb{C}}\mathbb{C}^n \times \mathbb{P}_{\mathbb{C}}\mathbb{C}^m \rightarrow \mathbb{P}_{\mathbb{C}}\mathbb{C}^{p \times n \times m}$ is the whole space $\mathbb{P}_{\mathbb{C}}\mathbb{C}^{p \times n \times m}$, where $\mathbb{P}_K V$ denotes the projective space consisting of one dimensional subspaces of the K -vector space V . Thus, by counting the dimensions, we see that $\text{grank}(p, n, m) \geq \lceil \frac{mnp}{m+n+p-2} \rceil$.

Suppose that $3 \leq m \leq n \leq p$. Then $p \geq \lceil \frac{mnp}{m+n+p-2} \rceil$ if and only if $p \geq (m-1)(n-1) + 1$. Catalisano, Geramita, and Gimigliano [CGG02] (see also [CGG08]) proved that if $(m-1)(n-1) + 1 \leq p \leq mn$, then $\text{grank}(p, n, m) = p$. Thus, $\min \text{trank}(p, n, m) = p$ in these cases. ten Berge [tB00] called a $p \times n \times m$ -tensor with $(m-1)n < p < mn$ a tall array or a tall tensor and proved that $\text{trank}(p, n, m) = \{p\}$ for these cases (see [SSM16, Chapter 6] for another proof). Here we define a $p \times n \times m$ -tensor a semi-tall tensor if $(m-1)(n-1) + 1 \leq p \leq (m-1)n$. We [SSM13, SMS15, SMS17]

studied the plurality of typical ranks of semi-tall tensors and proved that if $(m-1)(n-1)+2 \leq p \leq (m-1)n$, then $\text{trank}(p, n, m) = \{p, p+1\}$ if there exists a nonsingular bilinear map $\mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}^{mn-p}$ and $\text{trank}(p, n, m) = \{p\}$ otherwise, where a bilinear map $\varphi: V_1 \times V_2 \rightarrow W$ is nonsingular if $\varphi(x, y) = 0$ implies $x = 0$ or $y = 0$.

We also showed in [SMS17] that the former part of the above mentioned result also valid in the case where $p = (m-1)(n-1)+1$. Therefore, the latter part of the above mentioned result in the case where $p = (m-1)(n-1)+1$ is left open. In this paper, we treat the case where $p = (m-1)(n-1)+1$ and show that $\text{trank}(p, n, m) = \{p, p+1\}$ in several cases. In particular, we show that the latter part does not valid in the case where $p = (m-1)(n-1)+1$.

2 Preliminaries

Let K be a field and $T = (t_{ijk}) \in K^{\ell \times m \times n}$. For $1 \leq k \leq n$, we set $T_k = (t_{ijk}) \in K^{\ell \times m}$ and denote $T = (T_1; \dots; T_n)$. For $P \in \text{GL}(\ell, K)$ and $Q \in \text{GL}(m, K)$, we set $PTQ = (PT_1Q; \dots; PT_nQ)$. Note $\text{rank}PTQ = \text{rank}T$ by the definition of rank.

We first state the definition of the typical rank over $\mathbb{R}$.

Definition 2.1 If the set of rank r tensors over $\mathbb{R}$ of format $\ell \times m \times n$ contains a non-empty Euclidean open subset of $\mathbb{R}^{\ell \times m \times n}$, then we say r is a typical rank of $\ell \times m \times n$ tensors over $\mathbb{R}$. We denote the set of typical ranks of $\ell \times m \times n$ tensors over $\mathbb{R}$ by $\text{trank}_{\mathbb{R}}(\ell, m, n)$ or simply $\text{trank}(\ell, m, n)$.

By the definition of the rank, we see the following fact.

Lemma 2.2 *Let n_1, n_2 and n_3 be positive integers. Then $\text{trank}(n_{i_1}, n_{i_2}, n_{i_3}) = \text{trank}(n_1, n_2, n_3)$ for any permutation i_1, i_2, i_3 of 1, 2, 3.*

Definition 2.3 For $T = (T_1; \dots; T_n) \in K^{\ell \times m \times n}$, we set

$$\text{fl}_1(T) := (T_1, \dots, T_n) \in K^{\ell \times mn}$$

and

$$\text{fl}_2(T) := \begin{pmatrix} T_1 \\ \vdots \\ T_n \end{pmatrix} \in K^{\ell n \times m}$$

and call flattenings of T .

By the correspondence $K^{\ell \times m \times n} \leftrightarrow V_1 \otimes V_2 \otimes V_3$, where V_1 (resp. V_2, V_3) is a vector space over K of dimension ℓ (resp. m, n) with fixed basis, flattenings correspond to natural isomorphisms $V_1 \otimes V_2 \otimes V_3 \rightarrow V_1 \otimes (V_2 \otimes V_3)$ and $V_1 \otimes V_2 \otimes V_3 \rightarrow (V_1 \otimes V_3) \otimes V_2$. In particular, $\text{rank}(\text{fl}_1(T)) \leq \text{rank} T$ and $\text{rank}(\text{fl}_2(T)) \leq \text{rank} T$.

Definition 2.4 For $M = (\mathbf{m}_1, \dots, \mathbf{m}_n) = \begin{pmatrix} \mathbf{m}^{(1)} \\ \vdots \\ \mathbf{m}^{(\ell)} \end{pmatrix} \in K^{\ell \times n}$, we set

$$M_{\leq j} := (\mathbf{m}_1, \dots, \mathbf{m}_j), \quad j < M := (\mathbf{m}_{j+1}, \dots, \mathbf{m}_n), \quad M^{\leq i} := \begin{pmatrix} \mathbf{m}^{(1)} \\ \vdots \\ \mathbf{m}^{(i)} \end{pmatrix} \text{ and}$$

$${}^{i <} M := \begin{pmatrix} \mathbf{m}^{(i+1)} \\ \vdots \\ \mathbf{m}^{(\ell)} \end{pmatrix}.$$

Definition 2.5 Let R be a commutative ring and $M \in R^{\ell \times n}$. We denote by $I_t(M)$ the ideal of R generated by t -minors of R .

3 A condition of an $n \times p \times m$ -tensor to be of rank p

From now on, let m, n and p be integers with $3 \leq m \leq n$ and $(m-1)(n-1) + 1 \leq p \leq mn$. We set $u = mn - p$.

Fact 3.1 (1) $\text{grank}(p, n, m) = p$. In particular, $\min \text{trank}(p, n, m) = p$ [CGG02].

(2) If $p > (m-1)n$, then $\text{trank}(p, n, m) = \{p\}$ [tB00].

(3) Suppose $p \leq (m-1)n$. If there exists a nonsingular bilinear map $\mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}^u$, then $\text{trank}(p, n, m) = \{p, p+1\}$. Moreover, if $(m-1)(n-1) + 2 \leq p$, then the converse also hold true [SSM13, SMS15, SMS17].

Therefore, the case where $p = (m-1)(n-1) + 1$ and there is no nonsingular bilinear map $\mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}^u$ is still left open. In the following, we consider the case where $p = (m-1)(n-1) + 1$ and study if there are plural typical ranks of $p \times n \times m$ tensors over $\mathbb{R}$.

Before concentrating on the case where $p = (m-1)(n-1) + 1$, we state notations and a criterion of an $n \times p \times m$ tensor to be of rank p in the case

where $(n-1)(m-1)+1 \leq p \leq (m-1)n$. Note that $\text{trank}(p, n, m) = \text{trank}(n, p, m)$.

Definition 3.2 We set $\mathcal{V} := \{T \in \mathbb{R}^{n \times p \times m} \mid \text{fl}_2(T)^{\leq p} \text{ is nonsingular.}\}$, $\mathcal{O} := \{Y \in \mathbb{R}^{u \times n \times m} \mid {}_{p<} \text{fl}_1(Y) \text{ is nonsingular.}\}$, $\sigma: \mathcal{V} \rightarrow \mathbb{R}^{u \times p}$, $\sigma(T) = ({}^{p<} \text{fl}_2(T))(\text{fl}_2(T)^{\leq p})^{-1}$, $\nu: \mathcal{O} \rightarrow \mathbb{R}^{u \times p}$, $\nu(Y) = -({}_{p<} \text{fl}_1(Y))^{-1}(\text{fl}_1(Y)^{\leq p})$, $\tau: \mathbb{R}^{u \times p} \rightarrow \mathcal{V}$, $\tau(W) = \text{fl}_2^{-1} \begin{pmatrix} E_p \\ W \end{pmatrix}$ and $\mu: \mathbb{R}^{u \times p} \rightarrow \mathcal{O}$, $\mu(W) = \text{fl}_1^{-1}(W, -E_u)$.

Remark 3.3

$$\begin{array}{ccc} \mathbb{R}^{n \times p \times m} \supset \mathcal{V} & & \\ & \tau \swarrow \searrow \sigma & \\ & & \mathbb{R}^{u \times p} \\ & \mu \swarrow \searrow \nu & \\ \mathbb{R}^{u \times n \times m} \supset \mathcal{O} & & \end{array}$$

and $\sigma(\tau(W)) = \nu(\mu(W)) = W$ for $W \in \mathbb{R}^{u \times p}$.

Definition 3.4 Let $\mathbf{x} = (x_1, \dots, x_m)$ be a row vector of indeterminates, i.e., $x_1, \dots, x_m$ are independent indeterminates. For $A = (A_1; \dots; A_m) \in \mathbb{R}^{u \times n \times m}$, we set $M(\mathbf{x}, A) := x_1 A_1 + \dots + x_m A_m \in \mathbb{R}[x_1, \dots, x_m]^{u \times n}$.

Definition 3.5 Let $\mathbf{a}$ and $\mathbf{b}$ be column vectors with entries in $\mathbb{R}$ of dimension m and n respectively. We set $\psi(\mathbf{a}, \mathbf{b}) := (\mathbf{a} \otimes_{\text{Kr}} \mathbf{b})^{\leq p}$, where $\otimes_{\text{Kr}}$ denotes the

Kronecker product, i.e., if $\mathbf{a} = \begin{pmatrix} a_1 \\ \vdots \\ a_m \end{pmatrix}$, then $\psi(\mathbf{a}, \mathbf{b}) = \begin{pmatrix} a_1 \mathbf{b} \\ \vdots \\ a_{m-2} \mathbf{b} \\ a_{m-1} \mathbf{b}^{\leq p-(m-2)n} \end{pmatrix}$.

For $Y \in \mathbb{R}^{u \times n \times m}$, we define $U(Y)$ to be the vector subspace of $\mathbb{R}^p$ generated by $\{\psi(\mathbf{a}, \mathbf{b}) \mid M(\mathbf{a}^\top, Y)\mathbf{b} = \mathbf{0}\}$.

Lemma 3.6 For $T \in \mathcal{V}$, $\text{rank} T = p$ if and only if $\dim U(\mu(\sigma(T))) = p$.

Proof Set $\mu(\sigma(T)) = W = (W_1; \dots; W_m)$ and $\ell = u - n$. Then by [SMS17, Theorem 6.5 (1) $\iff$ (3)], we see that $\text{rank} T = p$ if and only if there are $B = (\mathbf{b}_1, \dots, \mathbf{b}_p) \in \mathbb{R}^{n \times p}$ and $p \times p$ diagonal matrices $D_1, \dots, D_m$ such that

$$(*) \left\{ \begin{array}{ll} D_k = \text{Diag}(d_{1k}, \dots, d_{pk}) & \text{for } 1 \leq k \leq m, \\ (d_{j1}W_1 + \dots + d_{jm}W_m)\mathbf{b}_j = \mathbf{0} & \text{for } 1 \leq j \leq p \text{ and} \\ \left(\begin{array}{c} BD_1 \\ \vdots \\ BD_{m-2} \\ B^{\leq n-\ell} D_{m-1} \end{array} \right) & \text{is nonsingular.} \end{array} \right.$$

First suppose that there are $B = (\mathbf{b}_1, \dots, \mathbf{b}_p) \in \mathbb{R}^{n \times p}$ and $D_1, \dots, D_m$ which satisfy (*). If we set $\mathbf{a}_j = (d_{j1}, \dots, d_{jm})^\top$ for $1 \leq j \leq p$, then

$$M(\mathbf{a}_j^\top, W)\mathbf{b}_j = \mathbf{0}$$

for $1 \leq j \leq p$ and

$$(\psi(\mathbf{a}_1, \mathbf{b}_1), \dots, \psi(\mathbf{a}_p, \mathbf{b}_p)) = \begin{pmatrix} BD_1 \\ \vdots \\ BD_{m-2} \\ B^{\leq n-\ell} D_{m-1} \end{pmatrix}$$

is nonsingular. Therefore $\dim U(\mu(\sigma(T))) = p$.

Conversely, assume that $\dim U(\mu(\sigma(T))) = p$. Then there are $\mathbf{a}_1, \dots, \mathbf{a}_p \in \mathbb{R}^m$ and $\mathbf{b}_1, \dots, \mathbf{b}_p \in \mathbb{R}^n$ such that $\psi(\mathbf{a}_1, \mathbf{b}_1), \dots, \psi(\mathbf{a}_p, \mathbf{b}_p)$ are linearly independent. Set $\mathbf{a}_j = (d_{j1}, \dots, d_{jm})^\top$ for $1 \leq j \leq p$, $D_k = \text{Diag}(d_{1k}, \dots, d_{pk})$ for $1 \leq k \leq m$ and $B = (\mathbf{b}_1, \dots, \mathbf{b}_p) \in \mathbb{R}^{n \times p}$. Then it is easily verified that B and $D_1, \dots, D_m$ satisfy (*). ■

4 Determinantal varieties and Bezout's theorem

From now on, we consider the case where $p = (m-1)(n-1) + 1$. Then $u = m + n - 2$.

Definition 4.1 We set

$$A_k := \begin{pmatrix} O_{(k-1) \times n} \\ E_n \\ O_{(m-k-1) \times n} \end{pmatrix} \in \mathbb{R}^{u \times n}$$

for $1 \leq k \leq m-1$ and

$$A_m := \begin{pmatrix} O_{(m-1) \times (n-1)} & -\mathbf{e}_1 \\ E_{n-1} & \mathbf{0} \end{pmatrix} \in \mathbb{R}^{u \times n},$$

where $\mathbf{e}_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \in \mathbb{R}^{m-1}$ and $A = (A_1; \dots; A_m) \in \mathbb{R}^{u \times n \times m}$.

The next fact is the key lemma of this paper.

Lemma 4.2 *Let y be an indeterminate and $a_1, \dots, a_{m-1} \in \mathbb{C}$. Then the following conditions are equivalent, where $V_a(I)$ denotes the affine variety defined by an ideal I .*

$$(1) (a_1, \dots, a_{m-1}, -1) \in V_a(I_n(M(\mathbf{x}, A))).$$

$$(2) y^{m-1} - a_{m-1}y^{m-2} - \dots - a_2y - a_1 \text{ is a factor of } y^u + 1.$$

In order to prove this lemma, we need some preparation. First we make the following

Definition 4.3 Let y be an indeterminate and $\{\mu_t\}_{t \geq 1}$ an infinite sequence of complex numbers. We set $\mathcal{J}(\{\mu_t\}_{t \geq 1}) = \{f(y) \in \mathbb{C}[y] \mid f(y) = \sum_k c_k y^k, \sum_k c_k \mu_{k+t} = 0 \text{ for any } t \geq 1\}$.

It is easily verified that $\mathcal{J}(\{\mu_t\}_{t \geq 1})$ is an ideal of $\mathbb{C}[y]$.

Now let $a_1, \dots, a_{m-1} \in \mathbb{C}$. Set $\lambda_t = 0$ for $1 \leq t \leq m-2$, $\lambda_{m-1} = 1$ and

$$\lambda_{m-1+s} = \det \begin{pmatrix} a_{m-1} & a_{m-2} & \cdots & a_1 & & & \\ -1 & \ddots & \ddots & & \ddots & & \\ & \ddots & \ddots & \ddots & & a_1 & \\ & & \ddots & \ddots & \ddots & \vdots & \\ & & & \ddots & \ddots & a_{m-2} & \\ & & & & -1 & a_{m-1} \end{pmatrix}$$

for $s \geq 1$, where the right hand side is an $s \times s$ -determinant (some a_i 's may not appear for small s).

By the first row expansion, we see the following

Lemma 4.4 *For $t \geq m$, we have $\lambda_t = \sum_{k=1}^{m-1} a_{m-k} \lambda_{t-k}$.*

Set $h(y) = y^{m-1} - a_{m-1}y^{m-2} - \dots - a_2y - a_1$. By the above lemma, we see that $h(y) \in \mathcal{J}(\{\lambda_t\}_{t \geq 1})$. Further, since $\lambda_t = 0$ for $1 \leq t \leq m-2$ and $\lambda_{m-1} = 1$, there is no polynomial in $\mathcal{J}(\{\lambda_t\}_{t \geq 1})$ whose degree is less than $m-1$ except the zero polynomial, i.e., $\mathcal{J}(\{\lambda_t\}_{t \geq 1})$ is generated by $h(y)$.

Set

$$N = M((a_1, \dots, a_{m-1}, -1), A) = \begin{pmatrix} a_1 & & & & & 1 \\ & a_2 & & & & \\ & & \ddots & & & \\ & & & \ddots & & \\ & & & & \ddots & \\ a_{m-1} & & & & & \\ -1 & & & & & a_1 \\ & & & & & a_2 \\ & & & & & \vdots \\ & & & & -1 & a_{m-1} \end{pmatrix}$$

and for integers $c_1, \dots, c_n$ with $1 \leq c_1 < \dots < c_n \leq u$, we denote by $[c_1, \dots, c_n]_N$ the maximal minor of N consisting of the c_1 -th, $\dots$, c_n -th rows of N .

Now we state the following

Lemma 4.5 *Under the notation above, the following conditions are equivalent.*

- (1) $\text{rank} N < n$.
- (2) $[i, m, m+1, \dots, u]_N = 0$ for $1 \leq i \leq m-1$.
- (3) $\lambda_{u+t} = 0$ for $1 \leq t \leq m-2$ and $\lambda_{u+m-1} = -1$.
- (4) $\lambda_{u+t} = -\lambda_t$ for $t \geq 1$.
- (5) $y^u + 1 \in \mathcal{J}(\{\lambda_t\}_{t \geq 1})$.

Proof Let

$$U = \begin{pmatrix} a_{m-1} & a_{m-2} & \cdots & a_1 & & 1 \\ -1 & & \ddots & & & \\ & \ddots & & \ddots & & \\ & & \ddots & \ddots & \ddots & a_1 \\ & & & \ddots & \ddots & \vdots \\ & & & & \ddots & a_{m-2} \\ & & & & -1 & a_{m-1} \end{pmatrix}$$

be a $u \times u$ matrix. Then $\det(t \leq U) = \lambda_{u+m-1-t} + \delta_{0,t}$ for $0 \leq t \leq u-1$, where $\delta_{0,t}$ is the Kronecker's delta.

(1) $\implies$ (3): Since $_{m-2}U = N$ and $\text{rank} N < n$ by assumption, we see that $\det(t \leq U) = 0$ for $0 \leq t \leq m-2$. Thus, we see that $\lambda_{u+m-1-t} + \delta_{0,t} = 0$ for $0 \leq t \leq m-2$.

(3) $\implies$ (2): We see by the first row expansions of $\det({}_t^{\leq}U)$ and $[t+1, m, \dots, u]$ and the assumption that

$$\begin{aligned}
0 &= \lambda_{u+m-1-t} + \delta_{0,t} \\
&= \sum_{k=1}^{m-1} a_{m-k} \lambda_{u+m-1-t-k} + \delta_{0,t} \\
&= \sum_{k=m-t-1}^{m-1} a_{m-k} \lambda_{u+m-1-t-k} + \delta_{0,t} \\
&= \sum_{s=1}^{t+1} a_{t+2-s} \lambda_{u+1-s} + \delta_{0,t} \\
&= [t+1, m, \dots, u]_N
\end{aligned}$$

for $0 \leq t \leq m-2$.

(2) $\implies$ (1) follows from the fact that the last $n-1$ rows of N are linearly independent, (3) $\iff$ (4) follows from the facts that $\lambda_1 = \dots = \lambda_{m-2} = 0$ and $\lambda_{m-1} = 1$ and Lemma 4.4 and (4) $\iff$ (5) follows from the definition of $\mathcal{J}(\{\lambda_t\}_{t \geq 1})$. ■

Since $(a_1, \dots, a_{m-1}, -1) \in V_a(I_n(M(\mathbf{x}, A)))$ if and only if $\text{rank} N < n$ and $y^u + 1 \in \mathcal{J}(\{\lambda_t\}_{t \geq 1})$ if and only if $h(y)$ divides $y^u + 1$, we see Lemma 4.2 by Lemma 4.5.

Now we recall the following facts about determinantal varieties (see e.g. [Har92, p.151 and p.243]).

Fact 4.6 *Let X be a $u \times n$ matrix of indeterminates. Then the projective variety in $\mathbb{P}_{\mathbb{C}}\mathbb{C}^{u \times n}$ defined by $I_n(X)$ has degree $\binom{u}{n-1}$ and codimension $u - n + 1$.*

Note that $\binom{u}{n-1} = \binom{u}{m-1}$ and $u - n + 1 = m - 1$ since $u = m + n - 2$. Note also that there are $\binom{u}{m-1}$ monic factors of $y^u + 1$ of degree $m - 1$ in $\mathbb{C}[y]$.

In view of this fact, we make the following

Definition 4.7 For $B = (B_1; \dots; B_m) \in \mathbb{C}^{u \times n \times m}$, we set $\varphi_B: \mathbb{C}^m \rightarrow \mathbb{C}^{u \times n}$, $(\alpha_1, \dots, \alpha_m) \mapsto \alpha_1 B_1 + \dots + \alpha_m B_m$.

Then by Lemma 4.2, Fact 4.6 and Bezout's theorem, we see the following

Corollary 4.8 *Let $\mathbb{P}_{\mathbb{C}}(\text{Im} \varphi_A)$ be the linear subspace of $\mathbb{P}_{\mathbb{C}}\mathbb{C}^{u \times n}$ defined by $\text{Im} \varphi_A$. Then $\mathbb{P}_{\mathbb{C}}(\text{Im} \varphi_A)$ and $V_p(I_n(X))$ intersect transversely at $\binom{u}{m-1}$ distinct points, where $V_p(I)$ denotes the projective variety defined by the homogeneous ideal I .*

By the implicit function theorem, we see the following fact.

Corollary 4.9 *There is a Euclidean open neighborhood $\mathcal{U}$ of A in $\mathbb{R}^{u \times n \times m}$ such that if $B \in \mathcal{U}$, then φ_B is injective and the number of real points of $V_p(I_n(M(\mathbf{x}, B))) \subset \mathbb{P}_{\mathbb{C}}\mathbb{C}^m$ is the number of real monic polynomials of degree $m-1$ which divide $y^u + 1$, where we say a point of a complex projective space is real if all possible ratios of its homogeneous coordinates are real numbers.*

We denote the number of real monic polynomials of degree $m-1$ which divides $y^u + 1$ by $\alpha(m, n)$. Then we see the following

Lemma 4.10

$$\alpha(m, n) = \begin{cases} \binom{u/2}{(m-1)/2} & \text{if } m \text{ and } n \text{ are odd,} \\ \binom{(u-1)/2}{(m-2)/2} & \text{if } m \text{ is even and } n \text{ is odd,} \\ \binom{(u-1)/2}{(m-1)/2} & \text{if } m \text{ is odd and } n \text{ is even and} \\ 0 & \text{if } m \text{ and } n \text{ are even.} \end{cases}$$

By replacing $\mathcal{U}$ to a smaller neighborhood if necessary, we may assume that $\text{rank}(M(\mathbf{a}, B)_{\leq n-1}) = n-1$ for any $B \in \mathcal{U}$ and $\mathbf{a} \in \mathbb{R}^m \setminus \{\mathbf{0}\}$ (cf. [SMS17, Corollary 4.20]). Then we have the following fact.

Lemma 4.11 *Suppose $B \in \mathcal{U}$. Then $\#\{[\psi(\mathbf{a}, \mathbf{b})] \in \mathbb{P}_{\mathbb{R}}\mathbb{R}^p \mid M(\mathbf{a}, B)\mathbf{b} = \mathbf{0}, \mathbf{a} \in \mathbb{R}^m \setminus \{\mathbf{0}\}, \mathbf{b} \in \mathbb{R}^n \setminus \{\mathbf{0}\}\} = \alpha(m, n)$, where $[\mathbf{x}]$ denotes the point of $\mathbb{P}_{\mathbb{R}}\mathbb{R}^p$ defined by $\mathbf{x} \in \mathbb{R}^p \setminus \{\mathbf{0}\}$.*

5 Plural typical ranks of some formats of 3-tensors

In this section, we show that in certain formats of 3-tensors, there are plural typical ranks. We use the notation of the previous section.

First we recall the following fact.

Fact 5.1 ([SMS17, Proposition 2.4, Lemma 3.5, Theorems 7.3 and 8.1])

If $m-1$ and $n-1$ are not bit-disjoint, then $\text{trank}(n, p, m) = \{p, p+1\}$, where two positive integers are bit-disjoint if there are no 1's in the same place of their binary notation.

Example 5.2 $\text{trank}(n, p, m) = \{p, p+1\}$ in the following cases.

- (1) Both m and n are even.

- (2) $m = 5$ and $n \equiv 5, 6, 7, 8 \pmod{8}$.
- (3) $m = 6$ and $n \equiv 2, 4, 5, 6, 7, 8 \pmod{8}$.
- (4) $m = 7$ and $n \equiv 3, 4, 5, 6, 7, 8 \pmod{8}$.
- (5) $m = 8$ and $n \equiv 2, 3, 4, 5, 6, 7, 8 \pmod{8}$.
- (6) $m = 9$ and $n \equiv 9, 10, 11, 12, 13, 14, 15, 16 \pmod{16}$.

Set $A'' = (A_2; \dots; A_{m-2}; A_m; -A_{m-1}; -A_1)$. Then there is a permutation matrix $P \in \text{GL}(u, \mathbb{R})$ such that ${}_P\text{fl}_1(PA'') = -E_u$. Set $A' = PA''$ and $W_0 = \text{fl}_1(A')_{\leq p}$. Further set $\rho: \mathbb{R}^{u \times n \times m} \rightarrow \mathbb{R}^{u \times n \times m}$, $B = (B_1; \dots; B_m) \mapsto P(B_2; \dots; B_{m-2}; B_m; -B_{m-1}; -B_1)$ and $\mathcal{U} = \mu^{-1}(\rho(\mathcal{U}))$, where μ is the map defined in Definition 3.2. Then $\mathcal{U}$ is an open neighborhood of W_0 . Further, we see the following fact by Lemma 4.11.

Lemma 5.3 *If $T \in \sigma^{-1}(\mathcal{U})$, then $\#\{[\psi(\mathbf{a}, \mathbf{b})] \in \mathbb{P}_{\mathbb{R}}\mathbb{R}^p \mid M(\mathbf{a}, \mu(\sigma(T)))\mathbf{b} = \mathbf{0}, \mathbf{a} \in \mathbb{R}^m \setminus \{\mathbf{0}\}, \mathbf{b} \in \mathbb{R}^n \setminus \{\mathbf{0}\}\} = \alpha(m, n)$.*

Since $\sigma^{-1}(\mathcal{U})$ contains $\tau(W_0)$, $\sigma^{-1}(\mathcal{U})$ is not an empty set. Therefore, we see by Lemma 3.6 that if $\alpha(m, n) < p$, then there exists a non-empty Euclidean open subset $\sigma^{-1}(\mathcal{U})$ of $\mathbb{R}^{n \times p \times m}$ consisting of tensors of rank larger than p . Further, since we see by [SMS17, Theorem 8.1] that typical ranks of $n \times p \times m$ tensors are less than or equal to $p + 1$, we see the following fact.

Lemma 5.4 *If $\alpha(m, n) < p$, then $\text{trank}(n, p, m) = \{p, p + 1\}$.*

Now we state the following

Theorem 5.5 *Suppose $3 \leq m \leq n$ and $p = (m - 1)(n - 1) + 1$. Then $\text{trank}(n, p, m) = \{p, p + 1\}$ in the following cases.*

- (1) $m = 3$ or $m = 4$.
- (2) $m = 5$ and $n \leq 26$ or $n = 28$.
- (3) $m = 6$ and $n \leq 34$.
- (4) $m = 7$ and $n \leq 16$.
- (5) $m = 8$ and $n \leq 16$.

Proof By Lemma 4.10 and computation, we see that $\alpha(m, n) < p$ in the following cases: (1) $m = 3$ or $m = 4$, (2) $m = 5$ and $n \leq 26$ or $n = 28$, (3) $m = 6$ and $n \leq 34$, (4) $m = 7$ and $n \leq 12$, (5) $m = 8$ and $n \leq 14$ and (6) $m = 9$ and $n = 10$. Thus, we see the result by Example 5.2 and Lemma 5.4. ■

Remark 5.6 Set $m = 3$ and $n = 5$. Then $m - 1$ and $n - 1$ are bit-disjoint. However, by Theorem 5.5, we see that $\text{trank}(n, p, m) = \{p, p + 1\}$. Thus the converse of Fact 5.1 does not valid.

References

- [CGG02] M. V. Catalisano, A. V. Geramita, and A. Gimigliano, *Ranks of tensors, secant varieties of Segre varieties and fat points*, Linear Algebra Appl. **355** (2002), 263–285.
- [CGG08] M. V. Catalisano, A. V. Geramita, and A. Gimigliano, *On the ideals of secant varieties to certain rational varieties*, J. Algebra **319** (2008), no. 5, 1913–1931.
- [Har92] Joe Harris, *Algebraic geometry, A first course.*, Graduate Texts in Mathematics, vol. 133, Springer-Verlag, New York, 1992,
- [SSM16] Toshio Sakata, Toshio Sumi and Mitsuhiro Miyazaki, *Algebraic and Computational Aspects of Real Tensor Ranks*, SpringerBriefs in Statistics, Springer, 2016.
- [SMS15] Toshio Sumi, Mitsuhiro Miyazaki and Toshio Sakata, *Typical ranks of $m \times n \times (m - 1)n$ tensors with $3 \leq m \leq n$ over the real number field*, Linear Multilinear Algebra **63** (2015), no. 5, 940–955.
- [SMS17] Toshio Sumi, Mitsuhiro Miyazaki and Toshio Sakata, *Typical ranks for 3-tensors, nonsingular bilinear maps and determinantal ideals*, Journal of Algebra, **471** 409–453 (Feb. 2017)
- [SSM13] Toshio Sumi, Toshio Sakata and Mitsuhiro Miyazaki, *Typical ranks for $m \times n \times (m - 1)n$ tensors with $m \leq n$* , Linear Algebra and its Application, **438** 953–958 (Jan. 2013).
- [tB00] Jos M. F. ten Berge, *The typical rank of tall three-way arrays*, Psychometrika **65** (2000), no. 4, 525–532.