

Prediction Measures in Nonlinear Beta Regression Models

Patrícia Leone Espinheira¹ Luana C. Meireles da Silva¹ Alisson de Oliveira Silva¹

Raydonal Ospina^{1,#}

¹ Departamento de Estatística, Universidade Federal de Pernambuco, Cidade Universitária, Recife/PE, 50740-540, Brazil.

E-mail: raydonal@de.ufpe.br

Abstract

Nonlinear models are frequently applied to determine the optimal supply natural gas to a given residential unit based on economical and technical factors, or used to fit biochemical and pharmaceutical assay nonlinear data. In this article we propose PRESS statistics and prediction coefficients for a class of nonlinear beta regression models, namely P^2 statistics. We aim at using both prediction coefficients and goodness-of-fit measures as a scheme of model select criteria. In this sense, we introduce for beta regression models under nonlinearity the use of the model selection criteria based on robust pseudo- R^2 statistics. Monte Carlo simulation results on the finite sample behavior of both prediction-based model selection criteria P^2 and the pseudo- R^2 statistics are provided. Three applications for real data are presented. The linear application relates to the distribution of natural gas for home usage in São Paulo, Brazil. Faced with the economic risk of too overestimate or to underestimate the distribution of gas has been necessary to construct prediction limits and to select the best predicted and fitted model to construct best prediction limits it is the aim of the first application. Additionally, the two nonlinear applications presented also highlight the importance of considering both goodness-of-predictive and goodness-of-fit of the competitive models.

Keywords: Nonlinear beta regression; PRESS; prediction coefficient; pseudo- R^2 , power prediction.

1 Introduction

Ferrari and Cribari-Neto (2004) introduced a regression model in which the response is beta-distributed, its mean being related to a linear predictor through a link function. The linear predictor includes independent variables and regression parameters. Their model also includes a precision parameter whose reciprocal can be viewed as a dispersion measure. In the standard formulation of the beta regression model it is assumed that the precision is constant across observations. However, in many practical situations this assumption does not hold. Smithson and Verkuilen (2006) consider a beta regression specification in which dispersion is not constant, but is a function of covariates and unknown parameters and (Simas et al., 2010) introduces the class of nonlinear beta regression models. Parameter estimation is carried out by maximum likelihood (ML) and standard asymptotic hypothesis testing can be easily

performed. Practitioners can use the **betareg** package, which is available for the R statistical software (<http://www.r-project.org>), for fitting beta regressions. Cribari-Neto and Zeileis (2010) provide an overview of varying dispersion beta regression modeling using the **betareg** package. Diagnostic tools and improve ML estimation were accomplished in Espinheira et al. (2008a,b); Ospina et al. (2006); Chien (2011) and others. Inference for beta regression also have been developed in an Bayesian context (Figueroa-Zúñiga et al. (2013); Brascum et al. (2007) and Cepeda-Cuervo and Gamerman (2005).)

Recently Espinheira et al. (2014) build and evaluated bootstrap-based prediction intervals for the class of beta regression models with varying dispersion. However, a prior approach it is necessary, namely: the selection of the model with the best predictive ability, regardless of the goodness-of-fit. Indeed, the model selection is a crucial step in data analysis, since all inferential performance is based on the selected model. Bayer and Cribari-Neto (2017) evaluated the performance of different model selection criteria in samples of finite size in a beta regression model, such as Akaike Information Criterion (AIC) (Akaike, 1973), Schwarz Bayesian Criterion (SBC) (Schwarz, 1978) and various approaches based on pseudo- R^2 such as the coefficient of determination adjusted R_{FC}^2 proposed by Ferrari and Cribari-Neto (2004) and the version based on log-likelihood functions, namely by R_{LR}^2 . Indeed, the authors proposed two new model selection criteria and a fast two step model selection scheme considering both mean and dispersion submodels, for beta regression models with varying dispersion.

However, these methods do not offer any insight about the quality of the predictive values in agreement with the findings of Spiess and Neumeyer (2010) for nonlinear models. In this context, Allen (1974), proposed the PRESS (Predictive Residual Sum of Squares) criterion, that can be used as a measure of the predictive power of a model. The PRESS statistic is independent from the goodness-of-fit of the model, since, that its calculation is made by leaving out the observations that the model is trying to predict (Palmer and O'Connell, 2009). The PRESS statistics can be viewed as a sum of squares of external residuals (Bartoli, 2009). Thus, similarly of the approach of R^2 , Mediavilla et al. (2008) proposed a coefficient of prediction based on PRESS namely P^2 . The P^2 statistic can be used to select models from a predictive perspective adding important information about the predictive ability of the model in various scenarios.

Our chief goal in this paper is to propose versions of the PRESS statistics and the coefficients of prediction P^2 associated, for the linear and nonlinear beta regression models. As a second contribution, in especial to beta regression under nonlinearity, we evaluate the behavior of R_{LR}^2 (Bayer and Cribari-Neto, 2017) and R_{FC}^2 (Ferrari and Cribari-Neto, 2004) measures both when the model is correctly specified and when under model misspecification. The results of the simulations showed as the prediction coefficients can be useful in detecting misspecifications, or indicate difficulties on to estimate beta regression models when the data are close to the boundaries of the standard unit interval. Finally, the real data applications are the last and important contribution. Here we provide guidance for researchers in choosing and interpreting the measures proposed. In fact, based on these applications we can shown how it is important to consider both coefficients of prediction and coefficients of determination to build models more useful to describe the data.

2 The P^2 statistic measure

Consider the linear model, $Y = X\beta + \varepsilon$ where Y is a vector $n \times 1$ of responses, X is a known matrix of covariates of dimension $n \times p$ of full rank, β is the parameter vector of dimension $p \times 1$ and ε is a vector $n \times 1$ of errors. We have the least squares estimators: $\hat{\beta} = (X^\top X)^{-1} X^\top y$, the residual $e_t = y_t - x_t^\top \hat{\beta}$ and the predicted value $\hat{y}_t = x_t^\top \hat{\beta}$, where $x_t^\top = (x_{t1}, \dots, x_{tp})$, and $t = 1, \dots, n$. Let $\hat{\beta}_{(t)}$ be the least squares estimate of β without the t th observation and $\hat{y}_{(t)} = x_t^\top \hat{\beta}_{(t)}$ be the predicted value of the case deleted,

such that $e_{(t)} = y_t - \hat{y}_{(t)}$ is the prediction error or external residual. Thus, for multiple regression, the classis statistic

$$PRESS = \sum_{t=1}^n e_{(t)}^2 = \sum_{t=1}^n (y_t - \hat{y}_{(t)})^2, \quad (1)$$

which can be rewritten as $PRESS = \sum_{t=1}^n (y_t - \hat{y}_{(t)})^2 / (1 - h_{tt})^2$, where h_{tt} is the t th diagonal element of the projection matrix $X(X^\top X)^{-1}X^\top$.

Now, let y_1, \dots, y_n be independent random variables such that each y_t , for $t = 1, \dots, n$, is beta distributed, beta-distributed, denoted by $y_t \sim \mathcal{B}(\mu_t, \phi_t)$, i.e., each y_t has density function given by

$$f(y_t; \mu_t, \phi_t) = \frac{\Gamma(\phi_t)}{\Gamma(\mu_t \phi_t) \Gamma((1 - \mu_t) \phi_t)} y_t^{\mu_t \phi_t - 1} (1 - y_t)^{(1 - \mu_t) \phi_t - 1}, \quad 0 < y_t < 1, \quad (2)$$

where $0 < \mu_t < 1$ and $\phi_t > 0$. Here, $E(y_t) = \mu_t$ and $\text{Var}(y_t) = V(\mu_t)/(1 + \phi_t)$, where $V(\mu_t) = \mu_t(1 - \mu_t)$. Simas et al. (2010) proposed the class of nonlinear beta regression models in which the mean of y_t and the precision parameter can be written as

$$g(\mu_t) = \eta_{1t} = f_1(x_t^\top; \beta) \quad \text{and} \quad h(\phi_t) = \eta_{2t} = f_2(z_t^\top; \gamma), \quad (3)$$

where $\beta = (\beta_1, \dots, \beta_k)^\top$ and $\gamma = (\gamma_1, \dots, \gamma_q)^\top$ are, respectively, $k \times 1$ and $q \times 1$ vectors of unknown parameters ($\beta \in \mathbb{R}^k$; $\gamma \in \mathbb{R}^q$), η_{1t} and η_{2t} are the nonlinear predictors, $x_t^\top = (x_{t1}, \dots, x_{tk_1})$ and $z_t^\top = (z_{t1}, \dots, z_{tq_1})$ are vectors of covariates (i.e., vectors of independent variables), $t = 1, \dots, n$, $k_1 \leq k$, $q_1 \leq q$ and $k + q < n$. Both $g(\cdot)$ and $h(\cdot)$ are strictly monotonic and twice differentiable link functions. Furthermore, $f_i(\cdot)$, $i = 1, 2$, are differentiable and continous functions, such that the matrices $J_1 = \partial \eta_1 / \partial \beta$ and $J_2 = \partial \eta_2 / \partial \gamma$ have full rank (their ranks are equal to k and q , respectively). The parameters that index the model can be estimated by maximum likelihood (ML). In the Appendix, we present the log-likelihood function, the score vector and Fisher's information matrix for the nonlinear beta regression model.

In the nonlinear beta regression model, the ML estimator $\hat{\beta}$ can be viewed as the least squares estimator of β (see Appendix) obtained by regressing

$$\check{y} = \hat{\Phi}^{1/2} \hat{W}^{1/2} u_1 \quad \text{on} \quad \check{J}_1 = \hat{\Phi}^{1/2} \hat{W}^{1/2} J_1, \quad (4)$$

with $\Phi = \text{diag}(\phi_1, \dots, \phi_n)$, $J_1 = \partial \eta_1 / \partial \beta$. Here, matrix W and u_1 are given in (15)–(17) in the Appendix. Thus, the prediction error is $\check{y}_t - \hat{y}_{(t)} = \hat{\phi}_t^{1/2} \hat{w}_t^{1/2} u_{1,t} - \hat{\phi}_t^{1/2} \hat{w}_t^{1/2} J_{1t}^\top \hat{\beta}_{(t)}$, in which J_{1t}^\top is the t th row of the J_1 matrix. Using the ideas proposed by (Pregibon, 1981) we have that $\hat{\beta}_{(t)} = \hat{\beta} - \{(J_1^\top \hat{\Phi} \hat{W} J_1)^{-1} J_{1t} \hat{\phi}_t^{1/2} \hat{w}_t^{1/2} r_t^\beta\} / (1 - h_{tt}^*)$, where r_t^β is the weighted 1 residual (Espinheira et al., 2008a) defined as

$$r_t^\beta = \frac{y_t^* - \hat{\mu}_t^*}{\sqrt{\hat{v}_t}}, \quad (5)$$

where, $y_t^* = \log\{y_t / (1 - y_t)\}$, $\mu_t^* = \psi(\mu_t \phi_t) - \psi((1 - \mu_t) \phi_t)$ and v_t is given in 15 in the Appendix. Hence, we can write $\check{y}_t - \hat{y}_{(t)} = r_t^\beta / (1 - h_{tt}^*)$, where h_{tt}^* is the t th diagonal element of projection matrix

$$H^* = (\hat{W} \hat{\Phi})^{1/2} J_1 (J_1 \hat{\Phi} \hat{W} J_1)^{-1} J_1^\top (\hat{\Phi} \hat{W})^{1/2}.$$

Finally, for the nonlinear beta regressions models the classic PRESS statistic based on (1) is given by

$$PRESS = \sum_{t=1}^n (\tilde{y}_t - \hat{y}_{(t)})^2 = \sum_{t=1}^n \left(\frac{r_t^\beta}{1 - h_{tt}^*} \right)^2. \quad (6)$$

Note that the t th observation in (6) is not used in fitting the regression model to predict y_t , then both the external predicted values $\hat{y}_{(t)}$ and the external residuals $e_{(t)}$ are independent of y_t . This fact enables the PRESS statistic to be a true assessment of the prediction capabilities of the regression model regardless of the overall model fit quality. Additionally, when the predictors in (3) are linear functions of the parameters, i.e., $g(\mu_t) = x_t^\top \beta$ and $h(\phi_t) = z_t^\top \gamma$, the expression in (6) also represent the PRESS statistic for a class of linear beta regression models with $p = k + q$ unknown regression parameters.

Considering the same approach to construct the determination coefficient R^2 for linear models, we can think in a prediction coefficient based on PRESS, namely

$$P^2 = 1 - \frac{PRESS}{SST_{(t)}}, \quad (7)$$

where $SST_{(t)} = \sum_{t=1}^n (y_t - \bar{y}_{(t)})^2$ and $\bar{y}_{(t)}$ is the arithmetic average of the $y_{(t)}$, $t = 1, \dots, n$. It can be shown that $SST_{(t)} = (n/(n-p))^2 SST$, wherein p is the number of model parameters and SST is the Total Sum of Squares for the full data. For a class of beta regressions models with varying dispersion, $SST = \sum_{t=1}^n (\tilde{y}_t - \bar{\tilde{y}})^2$, $\bar{\tilde{y}}$ is the arithmetic average of the $\tilde{y}_t = \hat{\phi}_t^{1/2} \hat{w}_t^{1/2} u_{1,t}$, $t = 1, \dots, n$. It is noteworthy that the measures R^2 and P^2 are distinct, since that the R^2 propose to measure the model fit quality and the P^2 measure the predictive power of the model.

Cook and Weisberg (1982) suggest other versions of PRESS statistic based on different choices of residuals. Thus, we present another version of PRESS statistic and P^2 measure by considering the combined residual proposed by Espinheira et al. (2017). In this way,

$$PRESS_{\beta\gamma} = \sum_{t=1}^n \left(\frac{r_{p,t}^{\beta\gamma}}{1 - h_{tt}^*} \right)^2 \quad \text{and} \quad P_{\beta\gamma}^2 = 1 - \frac{PRESS_{\beta\gamma}}{SST_{(t)}}, \quad (8)$$

respectively, where

$$r_t^{\beta\gamma} = \frac{(y_t^* - \hat{\mu}_t^*) + \hat{a}_t}{\sqrt{\hat{\zeta}_t}}, \quad a_t = \mu_t(y_t^* - \mu_t^*) + \log(1 - y_t) - \psi((1 - \mu_t)\phi_t) + \psi(\phi_t) \quad (9)$$

and $\zeta_t = (1 + \mu_t)^2 \psi'(\mu_t \phi_t) + \mu_t^2 \psi'((1 - \mu_t)\phi_t) - \psi'(\phi_t)$.

Note that, P^2 and $P_{\beta\gamma}^2$ given in (7) and (8), respectively, are not positive quantifiers. In fact, the $PRESS/SST_{(t)}$ is a positive quantity, thus the P^2 and the $P_{\beta\gamma}^2$ are measures that take values in $(-\infty; 1]$. The closer to one the better is the predictive power of the model.

In order to check the model goodness-of-fit with linear or nonlinear predictors for a class of beta regression, we evaluate the R_{FC}^2 defined as the square of the sample coefficient of correlation between $g(y)$ and $\hat{\eta}_1$ (Ferrari and Cribari-Neto, 2004), and its penalized version based on Bayer and Cribari-Neto (2017) given by $R_{FC_c}^2 = 1 - (1 - R_{FC}^2)(n - 1)/(n - (k_1 + q_1))$, where k_1 and q_1 are, respectively, the number of covariates of the mean submodel and dispersion submodel.

We also evaluate two version of pseudo- R^2 based on likelihood ratio. The first one proposed by Nagelkerke (1991): $R_{LR}^2 = 1 - (L_{null}/L_{fit})^{2/n}$, where L_{null} is the maximum likelihood achievable (satu-

rated model) and L_{fit} is the achieved by the model under investigation. The second one is a proposal of Bayer and Cribari-Neto (2017) that takes account the inclusion of covariates both in the mean submodel and in the precision submodel, given by:

$$R_{LR_c}^2 = 1 - (1 - R_{LR}^2) \left(\frac{n-1}{n - (1+\alpha)k_1 - (1-\alpha)q_1} \right)^\delta,$$

where $\alpha \in [0, 1]$ and $\delta > 0$. Based on simulation results obtained in Bayer and Cribari-Neto (2017) we choose in this work the values $\alpha = 0.4$ and $\delta = 1$. Therefore, penalized versions of P^2 and $P_{\beta_\gamma}^2$ are respectively now given by: $P_c^2 = 1 - (1 - P^2)(n-1)/(n - (k_1 + q_1))$ and $P_{\beta_{\gamma_c}}^2 = 1 - (1 - P_{\beta_\gamma}^2)(n-1)/(n - (k_1 + q_1))$.

3 Simulation

In this section we simulate several different data generating processes to evaluate the performance of the predictive measures. The Monte Carlo experiments were carried out using both fixed and varying dispersion beta regressions as data generating processes. All results are based on 10,000 Monte Carlo replications.

Linear models: Table 1 shows the mean values of the predictive statistics obtained by simulation of fixed dispersion beta regression model that involves a systematic component for the mean given by

$$\log \left(\frac{\mu_t}{1 - \mu_t} \right) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3} + \beta_4 x_{t4} + \beta_5 x_{t5}, \quad t = 1, \dots, n, \quad (10)$$

The covariate values were independently obtained as random draws of the following distributions: $X_{ti} \sim U(0, 1)$, $i = 2, \dots, 5$ and were kept fixed throughout the experiment. The precisions, the sample sizes and the range of mean response are, respectively, $\phi = (20, 50, 148, 400)$, $n = (40, 80, 120, 400)$, $\mu \in (0.005, 0.12)$, $\mu \in (0.90, 0.99)$ and $\mu \in (0.20, 0.88)$. Under the model specification given in (10) we investigate the performances of the statistics by omitting covariates. In this case, we considered the Scenarios 1, 2 and 3, in which are omitted, three, two and one covariate, respectively. In a fourth scenario the estimated model is correctly specified (true model).

The results in Table 1 show that the mean values of all statistics increase as covariates are included in the model and the value of ϕ increases. On the other hand, as the size of the sample increases, the misspecification of the model is evidenced by lower values of the statistics (Scenarios 1, 2 and 3). It shall be noted that the values of all statistics are considerably larger when $\mu \in (0.20, 0.88)$. Additionally, its values approaching one when the estimated model is closest to the true model. For instance, in Scenario 4 for $n = 40$, $\phi = 150$ the values of P^2 and R_{LR}^2 are, respectively, 0.936 and 0.947.

The behavior of the statistics for finite samples changes substantially when $\mu \in (0.90; 0.99)$. It is noteworthy the reduction of its values, revealing the difficulty in fitting the model and make prediction when $\mu \approx 1$ (The log-likelihood of the model tends to no longer limited). Indeed, in this range of μ is more difficult to make prediction than to fit the model. For example, in Scenario 1, when three covariates are omitted from the model, $n = 40$ and $\phi = 150$ the P^2 value equals to 0.071, whereas the R_{LR}^2 value is 0.243. Similar results were obtained for $n = 80, 120$. Even under true specification (Scenario 4) the model predictive power is more affected than the model quality of fit by the fact of $\mu \approx 1$. For instance, when $n = 120$ and $\phi = 50$ we have $P_{\beta_\gamma}^2 = 0.046$ and $R_{LR}^2 = 0.565$. The same difficulty in obtaining

predictions and in fitting the regression model occurs when $\mu \in (0.005, 0.12)$. Once again the greatest difficulty lies on the predictive sense.

Figure 1 present the boxplots of the 10,000 replications of the statistics: $P_{\beta\gamma}^2$, $P_{\beta\gamma_c}^2$, R_{LR}^2 , $R_{LR_c}^2$, R_{FC}^2 and $R_{FC_c}^2$ when the model is correctly specified (scenario 4), $n = 40$ and $\phi = 150$. In all boxplots the “side point” represents the mean value of the replications of the statistics. In the panel (a) we present the boxplots when $\mu \approx 0$. In the panel (b) we present the boxplots when μ is scattered on the standard unit interval and in the panel (c) we present the boxplots for $\mu \approx 1$. This figure shows that the means and the medians of all statistics are close. We also can notice based on the Figure 1 that both prediction power and goodness-of-fit of the model are affected when μ is close to the boundaries of the standard unit interval. However, it is noteworthy the great difficult to make prediction. Additionally, is possible to notice that the versions of R^2 displays similar behavior. In it follows we shall investigate the empirical distributions behaviour of the statistics proposed.




Figure 1: Model estimated correctly: $g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3} + \beta_4 x_{t4} + \beta_5 x_{t5}$. $\mu \in (0.20, 0.88)$; $\beta = (-1.9, 1.2, 1.0, 1.1, 1.3)^\top$; $\mu \in (0.90, 0.99)$; $\beta = (1.8, 1.2, 1, 1.1, 0.9)^\top$; $\mu \in (0.005, 0.12)$; $\beta = (-1.5, -1.2, -1.0, -1.1, -1.3)^\top$.

In Figures 2 and 3 we consider $\mu \in (0.20, 0.88)$. In Figure 2 the model is estimated correctly, $n = 40$ and $\phi = (20, 50, 150, 400)$. We notice that the prediction power distortions increase as the precision parameter increases, as expected. In Figure 3 we consider a misspecification problem (three omitted covariates). For illustration, we consider only $\phi = 50$ and $n = 40, 80, 120, 400$. It is important to notice that here the performance of the prediction measures does not deteriorate when the sample size is increased. Based on these figures we can reach some conclusions. First, the model precision affects both its predict power and goodness-of-fit. Second, for this range of μ the performance of the statistics are similar revealing the correct specification of the model (Figure 2). Third, when three covariates are omitted, with the increasing of sample size the replications values of the statistics tend being concentrated at small values due to the misspecification problem (Figure 3).

In what follows, we shall report simulation results on the finite-sample performance of the statistics when the dispersion modeling is neglected. To that end, the true data generating process considers varying dispersion, but a fixed dispersion beta regression is estimated. We also used different covariates in the mean and precision submodels. The samples sizes are $n = 40, 80, 120$. We generated 20 values for each covariate and replicated them to get covariate values for the three sample sizes (once, twice and three times, respectively).

Table 1: Values of the statistics. True model versus misspecification models (omitted covariates (Scenarios 1, 2 and 3)).

n	Estimated model	Scenario 1			Scenario 2			Scenario 3			Scenario 4		
		$g(\mu_t) = \beta_1 + \beta_2 x_{t2}$			$g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3}$			$g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3} + \beta_4 x_{t4}$			$g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3} + \beta_4 x_{t4} + \beta_5 x_{t5}$		
		$\mu \in (0.20, 0.88);$			$\beta = (-1.9, 1.2, 1.0, 1.1, 1.3)^\top.$								
$\phi \rightarrow$		20	50	150	20	50	150	20	50	150	20	50	150
40	P^2	0.307	0.363	0.393	0.393	0.463	0.502	0.506	0.602	0.656	0.694	0.847	0.936
	P_c^2	0.270	0.329	0.361	0.342	0.418	0.461	0.450	0.557	0.617	0.649	0.825	0.927
	$P_{\beta\gamma}^2$	0.304	0.357	0.385	0.402	0.473	0.512	0.512	0.609	0.663	0.694	0.847	0.936
	$P_{\beta\gamma_c}^2$	0.267	0.322	0.352	0.352	0.429	0.472	0.456	0.564	0.625	0.649	0.825	0.927
	R_{LR}^2	0.296	0.358	0.391	0.394	0.473	0.515	0.518	0.620	0.675	0.723	0.869	0.947
	$R_{LR_c}^2$	0.258	0.324	0.358	0.344	0.429	0.475	0.463	0.577	0.638	0.682	0.849	0.939
80	P^2	0.286	0.346	0.379	0.368	0.445	0.488	0.506	0.584	0.643	0.666	0.833	0.930
	P_c^2	0.267	0.329	0.363	0.343	0.423	0.468	0.450	0.561	0.624	0.643	0.821	0.925
	$P_{\beta\gamma}^2$	0.282	0.339	0.370	0.377	0.455	0.498	0.512	0.590	0.650	0.666	0.833	0.930
	$P_{\beta\gamma_c}^2$	0.264	0.322	0.353	0.353	0.433	0.478	0.456	0.569	0.632	0.643	0.821	0.925
	R_{LR}^2	0.291	0.356	0.391	0.385	0.468	0.513	0.518	0.614	0.672	0.706	0.860	0.943
	$R_{LR_c}^2$	0.273	0.339	0.375	0.361	0.447	0.494	0.463	0.593	0.655	0.686	0.851	0.939
120	P^2	0.279	0.340	0.374	0.360	0.439	0.483	0.469	0.578	0.639	0.656	0.833	0.928
	P_c^2	0.267	0.329	0.363	0.343	0.424	0.470	0.450	0.563	0.626	0.641	0.821	0.925
	$P_{\beta\gamma}^2$	0.275	0.333	0.365	0.369	0.449	0.493	0.475	0.585	0.646	0.656	0.833	0.928
	$P_{\beta\gamma_c}^2$	0.263	0.322	0.354	0.353	0.435	0.480	0.457	0.570	0.634	0.641	0.821	0.925
	R_{LR}^2	0.290	0.355	0.390	0.382	0.467	0.513	0.501	0.612	0.671	0.700	0.860	0.942
	$R_{LR_c}^2$	0.278	0.344	0.380	0.366	0.453	0.500	0.483	0.598	0.660	0.686	0.851	0.939
		$\mu \in (0.90, 0.99);$			$\beta = (1.8, 1.2, 1, 1.1, 0.9)^\top.$								
$\phi \rightarrow$		20	50	150	20	50	150	20	50	150	20	50	150
40	P^2	0.119	0.061	0.071	0.139	0.062	0.072	0.171	0.072	0.156	0.149	0.089	0.213
	P_c^2	0.071	0.010	0.021	0.067	-0.016	-0.006	0.076	-0.034	0.059	0.023	-0.045	0.097
	$P_{\beta\gamma}^2$	0.119	0.061	0.071	0.139	0.062	0.071	0.171	0.072	0.155	0.148	0.089	0.213
	$P_{\beta\gamma_c}^2$	0.072	0.010	0.020	0.068	-0.016	-0.007	0.076	-0.034	0.058	0.022	-0.045	0.097
	R_{LR}^2	0.164	0.196	0.243	0.221	0.266	0.336	0.271	0.374	0.466	0.444	0.593	0.774
	$R_{LR_c}^2$	0.119	0.153	0.203	0.157	0.205	0.281	0.188	0.303	0.405	0.362	0.533	0.741
80	P^2	0.093	0.036	0.044	0.112	0.038	0.046	0.149	0.045	0.120	0.123	0.056	0.175
	P_c^2	0.070	0.011	0.019	0.077	0.000	0.008	0.103	-0.006	0.073	0.063	-0.007	0.119
	$P_{\beta\gamma}^2$	0.094	0.036	0.043	0.113	0.038	0.045	0.149	0.045	0.119	0.122	0.056	0.175
	$P_{\beta\gamma_c}^2$	0.070	0.011	0.018	0.078	0.000	0.008	0.104	-0.006	0.072	0.063	-0.008	0.119
	R_{LR}^2	0.158	0.190	0.240	0.211	0.253	0.327	0.268	0.356	0.451	0.416	0.571	0.760
	$R_{LR_c}^2$	0.136	0.169	0.221	0.180	0.224	0.301	0.229	0.321	0.422	0.376	0.542	0.744
120	P^2	0.085	0.028	0.035	0.102	0.030	0.038	0.141	0.036	0.107	0.114	0.046	0.162
	P_c^2	0.069	0.012	0.018	0.079	0.005	0.013	0.111	0.002	0.076	0.075	0.004	0.125
	$P_{\beta\gamma}^2$	0.085	0.028	0.034	0.103	0.030	0.037	0.141	0.036	0.107	0.113	0.046	0.162
	$P_{\beta\gamma_c}^2$	0.069	0.012	0.018	0.080	0.005	0.012	0.112	0.002	0.076	0.075	0.004	0.125
	R_{LR}^2	0.156	0.188	0.239	0.207	0.249	0.324	0.268	0.349	0.447	0.406	0.565	0.756
	$R_{LR_c}^2$	0.142	0.175	0.226	0.186	0.230	0.306	0.242	0.327	0.428	0.380	0.545	0.745
		$\mu \in (0.005, 0.12);$			$\beta = (-1.5, -1.2, -1.0, -1.1, -1.3)^\top.$								
$\phi \rightarrow$		20	50	150	20	50	150	20	50	150	20	50	150
40	P^2	0.128	0.063	0.056	0.108	0.059	0.028	0.153	0.070	0.202	0.149	0.090	0.212
	P_c^2	0.081	0.013	0.005	0.033	-0.020	-0.053	0.056	-0.036	0.111	0.023	-0.044	0.096
	$P_{\beta\gamma}^2$	0.128	0.063	0.057	0.107	0.055	0.026	0.153	0.071	0.203	0.150	0.090	0.212
	$P_{\beta\gamma_c}^2$	0.081	0.013	0.006	0.032	-0.023	-0.055	0.056	-0.036	0.112	0.025	-0.044	0.097
	R_{LR}^2	0.199	0.215	0.254	0.265	0.349	0.379	0.326	0.415	0.548	0.442	0.595	0.774
	$R_{LR_c}^2$	0.156	0.172	0.214	0.204	0.295	0.327	0.249	0.348	0.496	0.360	0.535	0.741
80	P^2	0.105	0.040	0.032	0.083	0.043	0.012	0.128	0.038	0.165	0.123	0.057	0.174
	P_c^2	0.081	0.015	0.006	0.047	0.005	-0.027	0.081	-0.013	0.121	0.064	-0.007	0.119
	$P_{\beta\gamma}^2$	0.104	0.040	0.032	0.081	0.039	0.010	0.128	0.038	0.166	0.124	0.057	0.175
	$P_{\beta\gamma_c}^2$	0.081	0.015	0.007	0.045	0.001	-0.030	0.081	-0.013	0.121	0.065	-0.007	0.119
	R_{LR}^2	0.197	0.211	0.251	0.253	0.340	0.372	0.311	0.394	0.534	0.416	0.572	0.760
	$R_{LR_c}^2$	0.176	0.191	0.231	0.223	0.314	0.347	0.274	0.362	0.509	0.376	0.543	0.743
120	P^2	0.097	0.033	0.024	0.074	0.037	0.006	0.118	0.028	0.153	0.114	0.046	0.162
	P_c^2	0.081	0.016	0.007	0.050	0.012	-0.020	0.088	-0.006	0.123	0.075	0.004	0.125
	$P_{\beta\gamma}^2$	0.096	0.032	0.024	0.072	0.034	0.004	0.118	0.028	0.153	0.115	0.046	0.162
	$P_{\beta\gamma_c}^2$	0.081	0.016	0.007	0.048	0.009	-0.022	0.087	-0.006	0.124	0.076	0.004	0.125
	R_{LR}^2	0.195	0.209	0.250	0.247	0.337	0.370	0.304	0.388	0.530	0.407	0.565	0.755
	$R_{LR_c}^2$	0.181	0.196	0.237	0.228	0.320	0.354	0.280	0.367	0.513	0.381	0.546	0.744




Figure 2: Model estimated correctly: $g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3} + \beta_4 x_{t4} + \beta_5 x_{t5}$. $\mu \in (0.20, 0.88)$; $\beta = (-1.9, 1.2, 1.0, 1.1, 1.3)^\top$.

This was done so that the intensity degree of nonconstant dispersion

$$\lambda = \frac{\phi_{\max}}{\phi_{\min}} = \frac{\max_{t=1, \dots, n} \{\phi_t\}}{\min_{t=1, \dots, n} \{\phi_t\}}, \quad (11)$$

would remain constant as the sample size changes. The numerical results were obtained using the following beta regression model $g(\mu_t) = \log(\mu_t/(1 - \mu_t)) = \beta_1 + \beta_i x_{ti}$, and $\log(\phi_t) = \gamma_1 + \gamma_i z_{ti}$, $x_{ti} \sim U(0, 1)$, $z_{ti} \sim U(-0.5, 0.5)$, $i = 2, 3, 4, 5$, and $t = 1, \dots, n$ under different choices of parameters (Scenarios): **Scenario 5:** $\beta = (-1.3, 3.2)^\top$, $\mu \in (0.22, 0.87)$, $[\gamma = (3.5, 3.0)^\top; \lambda \approx 20]$, $[\gamma = (3.5, 4.0)^\top; \lambda \approx 50]$ and $[\gamma = (3.5, 5.0)^\top; \lambda \approx 150]$. **Scenario 6:** $\beta = (-1.9, 1.2, 1.6, 2.0)^\top$, $\mu \in (0.24, 0.88)$, $[\gamma = (2.4, 1.2, -1.7, 1.0)^\top; \lambda \approx 20]$, $[\gamma = (2.9, 2.0, -1.7, 2.0)^\top; \lambda \approx 50]$ and $[\gamma = (2.9, 2.0, -1.7, 2.8)^\top; \lambda \approx 100]$. **Scenarios 7 and 8 (Full models):** $\beta = (-1.9, 1.2, 1.0, 1.1, 1.3)^\top$, $\mu \in (0.20, 0.88)$, $[\gamma = (3.2, 2.5, -1.1, 1.9, 2.2)^\top; \lambda \approx 20]$, $[\gamma = (3.2, 2.5, -1.1, 1.9, 3.2)^\top; \lambda \approx 50]$, and $[\gamma = (3.2, 2.5, 1.1, 1.9, 4.0)^\top; \lambda \approx 100]$. All results were obtained using 10,000 replicates Monte Carlo replications.

Table 2 contain the values of the statistics. We notice that for each scenario the prediction power measure not present high distortion when we increase intensity degree of nonconstant dispersion. However, in the case of misspecification the statistics display smaller values in comparison with Scenario 8 (True specification), in which as greater is λ greater are the values of the statistics, as expected. Other important impression lies in the fact that the values of the R^2_{FC} are considerably smaller than the values of the others statistics, in special when λ increases.

That is a strong evidence that the R^2_{FC} does not perform well under nonconstant dispersion models.




Figure 3: Omitted covariates. Estimated model: $g(\mu_t) = \beta_1 + \beta_2 x_{t2}$. Correct model: $g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3} + \beta_4 x_{t4} + \beta_5 x_{t5}$; $\mu \in (0.20, 0.88)$; $\beta = (-1.9, 1.2, 1.0, 1.1, 1.3)^\top$

In fact, under nonconstant dispersion models the better performances are of the P^2 statistics, both in identifying wrong and correct specifications.

Table 2: Values of the statistics. Misspecified models, ϕ fixed: Scenarios 5, 6 and 7 versus Scenario 8 (correct specification).

	Scenario 5			Scenario 6			Scenario 7			Scenario 8		
True models	$g(\mu_t) = \beta_1 + \beta_2 x_{t2}$			$g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3} + \beta_4 x_{t4}$			$g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3} + \beta_4 x_{t4} + \beta_5 x_{t5}$			$g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3} + \beta_4 x_{t4} + \beta_5 x_{t5}$		
	$h(\phi_t) = \gamma_1 + \gamma_2 z_{t2} + \gamma_3 z_{t3}$			$h(\phi_t) = \gamma_1 + \gamma_2 z_{t2} + \gamma_3 z_{t3} + \gamma_4 z_{t4}$			$h(\phi_t) = \gamma_1 + \gamma_2 z_{t2} + \gamma_3 z_{t3} + \gamma_4 z_{t4} + \gamma_5 z_{t5}$			$h(\phi_t) = \gamma_1 + \gamma_2 z_{t2} + \gamma_3 z_{t3} + \gamma_4 z_{t4} + \gamma_5 z_{t5}$		
	$g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3}$			$g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3} + \beta_4 x_{t4}$			$g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3} + \beta_4 x_{t4} + \beta_5 x_{t5}$			$g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3} + \beta_4 x_{t4} + \beta_5 x_{t5}$		
Estimated models										$h(\phi_t) = \gamma_1 + \gamma_2 z_{t2} + \gamma_3 z_{t3} + \gamma_4 z_{t4} + \gamma_5 z_{t5}$		
$\lambda \rightarrow$	20	50	100	20	50	100	20	50	100	20	50	100
P^2	0.759	0.718	0.674	0.545	0.565	0.523	0.638	0.624	0.529	0.885	0.906	0.914
P^2_c	0.739	0.695	0.647	0.493	0.515	0.469	0.585	0.569	0.460	0.851	0.878	0.888
$P^2_{\beta\gamma}$	0.758	0.716	0.671	0.546	0.567	0.528	0.637	0.624	0.530	0.885	0.906	0.913
$P^2_{\beta\gamma_c}$	0.738	0.693	0.643	0.494	0.517	0.474	0.584	0.568	0.460	0.851	0.878	0.888
R^2_{LR}	0.782	0.743	0.700	0.580	0.611	0.577	0.670	0.653	0.554	0.796	0.816	0.840
$R^2_{LR_c}$	0.764	0.722	0.675	0.532	0.567	0.529	0.622	0.602	0.488	0.735	0.761	0.792

Figure 4 summarizes the predictive power performance for each measure. The graphs show that the mean and median performance of R^2_{FC} and $R^2_{FC_c}$ are significantly worse than the performance of the

other measures. The comparison among the best measures indicate that the median of P^2 and R_{LR}^2 performance are significantly better than the measures based on pseudo- R^2 and besides reveal some asymmetry of the statistics when the intensity degree of nonconstant dispersion levels increasing. These findings hold in all observation scenarios.




Figure 4: True model: $g(\mu_t) = \beta_1 + \beta_2 x_t$, $h(\phi_t) = \gamma_1 + \gamma_2 z_t$.

Nonlinear models: In it follows we shall present Monte Carlo experiments for the class of nonlinear beta regression models. To that end we shall use the starting values scheme for the estimation by maximum likelihood proposed by Espinheira et al. (2017). The numerical results were obtained using the following beta regression model as data generating processes:

$$\log\left(\frac{\mu_t}{1-\mu_t}\right) = \beta_1 + x_{t2}^{\beta_2} + \beta_3 \log(x_{t3} - \beta_4) + \frac{x_{t3}}{\beta_5}, \quad t = 1, \dots, n$$

$x_{t2} \sim U(1, 2)$, $x_{t3} \sim U(4.5, 34.5)$ and ϕ were kept fixed throughout the experiment. The precisions and the sample sizes are $\phi = (20, 50, 150, 400)$, $n = (20, 40, 60, 200, 400)$. Here, the vector of the parameters of the submodel of mean is $\beta = (1.0, 1.9, -2.0, 3.4, 7.2)^\top$ that produce approximately a range of values for the mean given by $\mu \in (0.36, 0.98)$. To evaluate the performances of statistics on account of nonlinearity negligence we consider the following model specification: $\log\left(\frac{\mu_t}{1-\mu_t}\right) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3}$. All results are based on 10,000 Monte Carlo replications and for each replication, we generated the response values as $y_t \sim \mathcal{B}(\mu_t, \phi_t)$, $t = 1, \dots, n$.

Table 3 contains numerical results for the fixed dispersion beta regression model as data generating processes. Here, we compared the performances of the statistics both under incorrect specifications and under correct specification of the nonlinear beta regression model. The results presented in this table

reveal that the P^2 and $P_{\beta\gamma}^2$ statistics outperform the R^2 statistics in identifying more emphatically the misspecification. We must emphasize that the response mean is scattered on the standard unit interval.

Thus, we should not have problems to make prediction and the smaller values of P^2 statistics in comparison with the values of the R^2 statistics is due to the better performance of the statistics based on residuals in identifying misspecification problems. For example, fixing the precision on $\phi = 400$, for $n = 20$, we have values of P^2 , $P_{\beta\gamma}^2$, R_{LR}^2 and R_{FC}^2 equal to 0.576, 0.601, 0.700, 0.637, respectively. For $n = 40$ and $n = 60$ the values of the statistics are 0.568, 0.593, 0.698, 0.634 and 0.562, 0.588, 0.698, 0.633, respectively. We can also notice that the values of the penalized versions of the statistics tend to be greater as the sample size increasing, what it makes sense.

Figure 5 summarizes the predictive power measure performance with boxplots over the Monte Carlo replics. The boxplots clearly show the statistical significance of the performance differences between the measures. The outperformance of the P^2 and $P_{\beta\gamma}^2$ statistics in identifying misspecification is more clear when we analyzed the plot. When the sample size increases, the distributions of the statistics based on residuals tend been concentrated in small values. For the other hand, the distributions of the R^2 statistics tend been concentrated at the same values, considerably greater than the values of the P^2 and $P_{\beta\gamma}^2$ statistics.

Figure 6 summarizes the empirical distribution behavior of predictive power measure when $n = 60$. The graphs show that the median performance of R_{LR}^2 is significantly worse than the performance of the P^2 and $P_{\beta\gamma}^2$ measures. However, under true specification the statistics perform equally well and as the precision of the model increases the values of the statistics tend being concentrated close to one. Also, we notice that the performance comparison among different levels of ϕ shows a systematic increase of the power performance.

Table 3: Values of the statistics. True model: $g(\mu_t) = \beta_1 + x_{t2}^{\beta_2} + \beta_3 \log(x_{t3} - \beta_4) + \frac{x_{t3}}{\beta_5}$, $x_{t2} \sim U(1, 2)$, $x_{t3} \sim U(4.5, 34.5)$, $\beta = (1.0, 1.9, -2.0, 3.4, 7.2)^\top$, $\mu \in (0.36, 0.98)$, $t = 1, \dots, n$, ϕ fixed. Misspecification: $: g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3}$ (omitted nonlinearity).

Estimated Model	With misspecification: $g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3}$												Correctly		
	20				40				60				60		
	20	50	150	400	20	50	150	400	20	50	150	400	50	150	400
P^2	0.485	0.535	0.564	0.576	0.438	0.508	0.550	0.568	0.420	0.496	0.543	0.562	0.849	0.936	0.975
P_c^2	0.388	0.448	0.483	0.497	0.391	0.467	0.513	0.532	0.388	0.469	0.518	0.539	0.835	0.930	0.973
$P_{\beta\gamma}^2$	0.502	0.556	0.588	0.601	0.456	0.531	0.575	0.593	0.439	0.520	0.568	0.588	0.849	0.936	0.975
$P_{\beta\gamma_c}^2$	0.409	0.473	0.511	0.526	0.411	0.492	0.539	0.559	0.409	0.494	0.545	0.566	0.835	0.930	0.973
R_{LR}^2	0.578	0.647	0.684	0.700	0.563	0.639	0.681	0.698	0.557	0.636	0.680	0.698	0.883	0.953	0.982
$R_{LR_c}^2$	0.499	0.581	0.625	0.643	0.526	0.608	0.654	0.673	0.533	0.616	0.662	0.681	0.863	0.945	0.979
R_{FC}^2	0.486	0.574	0.619	0.637	0.448	0.556	0.612	0.634	0.437	0.550	0.609	0.633	0.879	0.951	0.981
$R_{RC_c}^2$	0.389	0.494	0.548	0.569	0.402	0.519	0.580	0.604	0.407	0.526	0.588	0.613	0.867	0.946	0.979

Nonlinearity on dispersion model: The last simulations consider two nonlinear submodels both to mean and dispersion, namely:

$$\log\left(\frac{\mu_t}{1 - \mu_t}\right) = \beta_1 + x_t^{\beta_2} \quad \text{and} \quad \log(\phi_t) = \gamma_1 + z_t^{\gamma_2}.$$




Figure 5: Misspecification: omitted nonlinearity. Estimated model: $g(\mu_t) = \beta_1 + \beta_2 x_{t2} + \beta_3 x_{t3}$. True model: $g(\mu_t) = \beta_1 + x_{t2}^{\beta_2} + \beta_3 \log(x_{t3} - \beta_4) + \frac{x_{t3}}{\beta_5}$, $x_{t2} \sim U(1, 2)$, $x_{t3} \sim U(4.5, 34.5)$, $\beta = (1.0, 1.9, -2.0, 3.4, 7.2)^\top$, $t = 1, \dots, n$, $\phi = 150$, $\mu \in (0.36, 0.98)$.




Figure 6: Model correctly specified. True model: $g(\mu_t) = \beta_1 + x_{t2}^{\beta_2} + \beta_3 \log(x_{t3} - \beta_4) + \frac{x_{t3}}{\beta_5}$, $t = 1, \dots, n$, $x_{t2} \sim U(1, 2)$, $x_{t3} \sim U(4.5, 34.5)$, $\beta = (1.0, 1.9, -2.0, 3.4, 7.2)^\top$. Range of values for mean $\mu \in (0.36, 0.98)$.

We fixed: $n = 400$, $\beta = (-1.1, 1.7)^\top$, $x_t \sim U(0.3, 1.3)$; ($\mu \in (0.28, 0.61)$), $z_t \sim U(0.5, 1.5)$ and we varying γ such that $\gamma = (2.6, 3.0)^\top$; $\lambda \approx 25$, $\gamma = (1.6, 3.1)^\top$; $\lambda \approx 29$, $\gamma = (0.9, 3.2)^\top$; $\lambda \approx 35$ and $\gamma = (-0.3, 3.9)^\top$; $\lambda \approx 100$, $t = 1, \dots, n$. In Figure 7 we present the boxplots of the P^2 , $P^2_{\beta\gamma}$ and R^2_{LR} under negligence of nonlinearity, that is the estimated model is $\log\left(\frac{\mu_t}{1-\mu_t}\right) = \beta_1 + \beta_2 x_t$ and $\log(\phi_t) =$

$\gamma_1 + \gamma_2 z_t$. Based on this figure we notice that once again the statistics based on residuals outperform the R^2 statistics since that the values of the P^2 and $P_{\beta\gamma}^2$ are considerably smaller than the values of the R_{LR}^2 statistic, in especial when the nonconstant dispersion is more several (when λ increases).




Figure 7: Misspecified model: $g(\mu_t) = \beta_1 + \beta_2 x_t$, $h(\phi_t) = \gamma_1 + \gamma_2 z_t$. True model: $g(\mu_t) = \beta_1 + x_t^{\beta_2}$, $h(\phi_t) = \gamma_1 + z_t^{\gamma_2}$, $x_t \sim U(0.3, 1.3)$, $z_t \sim U(0.5, 1.5)$, $\beta = (-1.1, 1.7)^\top$, $t = 1, \dots, n$, $\mu \in (0.28, 0.61)$, $n = 400$.

Figure 8 summarizes the predictive power measure performance with boxplots over the Monte Carlo replics. Here, we evaluated the empirical distribution of the R_{FC}^2 statistics under nonconstant dispersion for two models estimated correctly. The plots reveals evidences that the pseudo- R^2 measures are not a good statistics for model selection when the dispersion varying along the observations. It is clear that P^2 and $P_{\beta\gamma}^2$ measures become more powerful as the λ increases.

4 Applications

In what follows we shall present an application based on real data.

Application I: The application relates to the distribution of natural gas for home usage (e.g., in water heaters, ovens and stoves) in São Paulo, Brazil. Such a distribution is based on two factors: the simultaneity factor (F) and the total nominal power of appliances that use natural gas, computed power Q_{max} . Using these factors one obtains an indicator of gas release in a given tubulation section, namely: $Q_p = F \times Q_{max}$. The simultaneity factor assumes values in $(0, 1)$, and can be interpreted as the probability of simultaneous appliances usage. Thus, based on F the company that supplies the gas decides how much gas to supply to a given residential unit.




Figure 8: True model: $g(\mu_t) = \beta_1 + x_t^{\beta_2}$, $h(\phi_t) = \gamma_1 + z_t^{\gamma_2}$, $x_t \sim U(0.3, 1.3)$, $z_t = x_t$, $\beta = (-1.1, 1.7)^\top$, $\gamma = (2.3, 5.3)^\top$, $n = 400$.


The data were analyzed by Zerbinatti (2008), obtained from the Instituto de Pesquisas Tecnológicas (IPT) and the Companhia de Gás de São Paulo (COMGÁS). The response variable (y) are the simultaneity factors of 42 valid measurements of sampled households, and the covariate is the computed power (x_1). The simultaneity factors ranged from 0.02 to 0.46, being the median equals 0.07. Zerbinatti (2008) modeled such data and concluded that the best performing model was the beta regression model based on logit link function for the mean and logarithmic form (log) of computed power used as covariate. However, the author shows that the beta regression model can underpredict the response. Thus, Espinheira et al. (2014) argue that it is important to have at disposal prediction intervals that can be used with beta regressions. To that end, the authors built and evaluated bootstrap-based prediction intervals for the response for the class of beta regression models. They applied the approach to the data on simultaneity factor. However, an important step in this case was the selection of the model with the best predictive power. To reach this aim the authors used a simplified version of PRESS statistic given by $PRESS = \sum_{t=1}^{42} (y_t - \hat{y}_{(t)})^2 / 42$ which selected the same model of Zerbinatti (2008). Here we aim at selecting the better predictive model to the data on simultaneity factor using the P^2 and $P^2_{\beta\gamma}$ statistics. We also consider the R^2_{LR} and R^2_{RC} as measures of goodness-of-fit model. The response is the simultaneity factor and the covariate X_2 is the log of computed power. We adjusted four beta regression models, considering constant and nonconstant dispersion and logit or log-log link function for μ . For the varying dispersion model we used the log link function for ϕ . The values of the statistics are presented in Table 4. Here, we should emphasize that the model predictive power is better when the measures P^2 and $P^2_{\beta\gamma}$ are close to one.

The Table 4 displays two important informations. First, we notice that by the R^2 measures the models equally fits well. Second, the P^2 and $P^2_{\beta\gamma}$ measures lead to the same conclusions, selecting the beta regression model with link log-log for the mean submodel and link log for the dispersion submodel, as the best model to make prediction to the data on simultaneity factor. The maximum likelihood parameter estimates are $\hat{\beta}_1 = -0.63$, $\hat{\beta}_2 = -0.31$, $\hat{\gamma}_1 = 3.81$ and $\hat{\gamma}_2 = 0.77$. Furthermore, the estimative of intensity of nonconstant dispersion is $\hat{\lambda} = 21.16$ (see (16)), such that $\hat{\phi}_{\max} = 242.39$ and $\hat{\phi}_{\min} = 11.45$. Selected among the candidates the best model in a predictive perspective, we still can use the PRESS statistic to identifying which observations are more difficult to predict. In this sense, we plot the individual components of $PRESS_{\beta\gamma_t}$ versus the observations index and we added a horizontal line at $3 \sum_{t=1}^n PRESS_{\beta\gamma_t} / n$ and singled out points that considerably exceeded this threshold.

Table 4: Values of the statistics from the candidate models. Data on simultaneity factor

Candidate models				
Mean	$\log(\mu_t/(1 - \mu_t)) =$	$-\log(-\log(\mu_t)) =$	$\log(\mu_t/(1 - \mu_t)) =$	$-\log(-\log(\mu_t)) =$
submodel	$\beta_1 + \beta_2 x_{t1}$	$\beta_1 + \beta_2 x_{t1}$	$\beta_1 + \beta_2 x_{t1}$	$\beta_1 + \beta_2 x_{t1}$
Dispersion	–	–	$\log(\phi_t) =$	$\log(\phi_t) =$
submodel			$\gamma_1 + \gamma_2 x_{t1}$	$\gamma_1 + \gamma_2 x_{t1}$
P^2	0.66	0.42	0.70	0.88
P_c^2	0.64	0.39	0.68	0.87
$P_{\beta\gamma}^2$	0.65	0.42	0.70	0.88
$P_{\beta\gamma_c}^2$	0.64	0.39	0.68	0.87
R_{LR}^2	0.72	0.70	0.74	0.74
$R_{LR_c}^2$	0.69	0.65	0.70	0.70
R_{FC}^2	0.69	0.72	0.70	0.72
$R_{FC_c}^2$	0.67	0.70	0.67	0.70

Figure 9 shows that the cases 3, 11, 33 and 33 arise as the observations with more predictive difficulty and are worthy of further investigation.

**Figure 9:** PRESS plot. Data on simultaneity factor

Application II: The second application consider a nonlinear beta regression used to modeling the proportion of killed grasshopper (y) at an assays on a grasshopper *Melanopus sanguinipes* with the insecticide carbofuran and the synergist piperonyl butoxide. This model was proposed by Espinheira et al. (2017) after careful building the scheme of starting values to iterative process of maximum likelihood estimation and after a meticulous residual analysis. Our aim here is applies both predictive power statistics and goodness-of-fit statistics to confirm or not the choice of the model made by residual analysis. The covariates are the dose of the insecticide (x_1) and the dose of the synergist (x_2). The data can be found in McCullagh and Nelder (1989, p. 385). Additionally, $y \in [0.04, 0.84]$, $\mu_y = 0.4501$ and the median of the response is equal to 0.4967.

The model selected with its estimates and respective p-values is present in Espinheira et al. (2017) and is given by $\log(\mu_t/1 - \mu_t) = \beta_1 + \beta_2 \log(x_{t1} - \beta_3) + \beta_4 \frac{x_{t2}}{x_{t2} + \beta_5}$ and $\sqrt{\phi_t} = \gamma_1 + \gamma_2 x_{t1} + \gamma_3 x_{t2}$, $t = 1, \dots, 15$. Now, by using residual analysis as diagnostic tools several linear beta regression models are compared with nonlinear model. After competition the nonlinear model: $\log(\mu_t/1 - \mu_t) = \beta_1 + \beta_2 \log(x_{t1} + 1.0) + \beta_3 x_{t2}$ $\sqrt{\phi_t} = \gamma_1 + \gamma_2 x_{t1}$ $t = 1, \dots, 15$ was selected. The estimatives of parameters are $\hat{\beta}_1 = -4.25$; $\hat{\beta}_2 = 1.79$; $\hat{\beta}_3 = 0.04$; $\hat{\gamma}_1 = 1.19$ and $\hat{\gamma}_2 = 0.19$. The values of the P^2 and R^2 measures for the two candidate models are present Table 5. Based on this table we can note that the nonlinear model outperforms the linear model under all statistics, that is, the nonlinear model has both better predictive power and better goodness-of-fit.

With aim in identifying observations for which to make prediction can be a hard task we plot values of PRESS statistic versus indices of the observations. In Figure 10 it is noteworthy how the case 14 is strongly singled out. In fact, this case was also singled out in plots of residual analysis made by Espinheira et al. (2017). However, the observation 14 is not an influential case, in sense of to affect inferential results. Besides, the choose model was capable to estimated well this case.

Thus, we confirm by the model selection measures that the beta nonlinear model proposed by Espinheira et al. (2017) is a suitable alternative to modeling of the data of insecticide carbofuran and the synergist piperonyl butoxide McCullagh and Nelder (1989, p. 385).

Table 5: Values of the statistics from the candidate models. Data on insecticide.

Candidate models		
	Linear Models	Nolinear models
Mean	$\log(\mu_t/(1 - \mu_t)) =$	$\log(\mu_t/(1 - \mu_t)) =$
submodel	$\beta_1 + \beta_2 \log(x_{t1} + 1.0) + \beta_3 x_{t2}$	$\beta_1 + \beta_2 \log(x_{t1} - \beta_3) + \beta_4 \frac{x_{t2}}{x_{t2} + \beta_5}$
Dispersion	$\sqrt{\phi_t} =$	$\sqrt{\phi_t} =$
submodel	$\gamma_1 + \gamma_2 x_{t1}$	$\gamma_1 + \gamma_2 x_{t1} + \gamma_3 x_{t2} + \gamma_4 (x_{t1} x_{t2})$
P^2	0.89	0.99
P_c^2	0.85	0.99
$P_{\beta\gamma}^2$	0.89	0.99
$P_{\beta\gamma_c}^2$	0.86	0.99
R_{LR}^2	0.83	0.99
$R_{LR_c}^2$	0.70	0.99
R_{FC}^2	0.79	0.97
$R_{FC_c}^2$	0.71	0.94

Application III: In the latter application we will use the dataset about available chlorine fraction after weeks of manufacturing from an investigation performed at Proctor & Gamble. A certain product must have a fraction of available chlorine equal to 0.50 at the time of manufacturing. It is known that chlorine fraction of the product decays with time. Eight weeks after the production, before the product is consumed, in theory there is a decline to a level 0.49.

The theory related to the problem indicates that the available chlorine fraction (y) decays according to a nonlinear function of the number of weeks (x) after fabrication of the product and unknown parameters (Draper and Smith, 1981 p. 276), given by

$$\eta_t = \beta_1 + (0.49 - \beta_1) \exp\{\beta_2(x_t - 8)\}. \quad (12)$$




Figure 10: PRESS plots. Data on insecticide.

The level 0.49 depends on several uncontrolled factors, as for example warehousing environments or handling facilities. Thus, the predictions based on theoretical model can be not reliable.

Cartons of the product were analyzed over a period aiming answer some questions like as: “*When should warehouse material be scrapped?*” or “*When should store stocks be replaced?*” According to knowledgeable chemists an equilibrium asymptotic level of available chlorine should be expected somewhere close to 0.30.

From predictor based on (12) we can note that when $x = 8$ the nonlinear model provides a true level for the available chlorine fraction (no error), wherein $\eta = 0.49$. We consider a new logit nonlinear beta regression model. We replaced the deterministic value 0.49 by an additional parameter at the predictor. Thus, the new nonlinear predictor for mean submodel is given by $\eta_t = \beta_1 + (\beta_3 - \beta_1)\exp\{-\beta_2(x_t - 8)\}$, $t = 1, \dots, 42$. Here the available chlorine fraction ranged from 0.38 to 0.49, being the mean and median are approximately equal 0.42. We investigated some competitive models. Table 6 the results of final candidates models and its statistics. The findings reveals that the model $\log(\mu_t/(1 - \mu_t)) = \beta_1 + (\beta_3 - \beta_1)\exp\{\beta_2(x_{t1} - 8)\}$ and $\log(\phi_t) = \gamma_1 + \gamma_2 \log x_{t1} + \exp\{\gamma_3(x_{t1} - 8)\}$ is the best performer in sense that it displays the higher statistics values. To estimate this model was necessary to build a starting values procedure for log-likelihood maximization as proposed by Espinheira et al. (2017). Since that we have more parameters than covariates firstly we used the theoretical information about the asymptotic level and found a initial guess to β_1 equal to 0.30. Thus, based on equation we took some values to y and x_1 and found a initial guess to β_2 , $\beta_2^{(0)} = 0.02$. Then we carried out the scheme of starting values to be used in nonlinear beta regression maximum likelihood estimation (Espinheira et al., 2017). The parameters estimatives are $\hat{\beta}_1 = -0.45963$ $\hat{\beta}_2 = -0.04166$ $\hat{\beta}_3 = 0.09479$ $\hat{\gamma}_1 = 13.18335$ $\hat{\gamma}_2 = -0.05413$ $-\hat{\gamma}_3 = 2.63158$. It is importance emphasize that the β_2 estimative conduce to a level of chlorine fraction equal to $0.4896 \approx 0.49$ that for this dataset confirm the theory that there is a decline to a level 0.49.

5 Conclusion

In this paper we develop the P^2 and $P_{\beta\gamma}^2$ measures based on two versions of PRESS statistics for the class of beta regression models. The P^2 coefficient consider the PRESS statistic based on ordinary residual

Table 6: Values of the statistics from the candidate models. Data on chlorine fraction

	Candidate models			
	$\log(\mu_t/(1-\mu_t)) = \beta_1 + (\beta_3 - \beta_1)\exp\{\beta_2(x_{t1} - 8)\}$	$-\log(-\log(\mu_t)) = \beta_1 + (\beta_3 - \beta_1)\exp\{\beta_2(x_{t1} - 8)\}$	$\log(\mu_t/(1-\mu_t)) = \beta_1 + (\beta_3 - \beta_1)\exp\{\beta_2(x_{t1} - 8)\}$	$-\log(-\log(\mu_t)) = \beta_1 + (\beta_3 - \beta_1)\exp\{\beta_2(x_{t1} - 8)\}$
η_{1t}				
η_{2t}	–	–	$\log(\phi_t) = \gamma_1 + \gamma_2 \log x_{t1} + \exp\{\gamma_3(x_{t1} - 8)\}$	$\log(\phi_t) = \gamma_1 + \gamma_2 \log x_{t1} + \exp\{\gamma_3(x_{t1} - 8)\}$
P^2	0.88	0.87	0.96	0.95
P_c^2	0.87	0.86	0.96	0.95
$P_{\beta\gamma}^2$	0.88	0.87	0.96	0.95
$P_{\beta\gamma_c}^2$	0.87	0.86	0.96	0.95
R_{LR}^2	0.87	0.87	0.88	0.88
$R_{LR_c}^2$	0.85	0.85	0.86	0.86

obtained from the Fisher’s scoring iterative algorithm for estimating β whereas $P_{\beta\gamma}^2$ is based on a new residual which is a combination of ordinaries residuals from the Fisher’s scoring iterative algorithm for estimating β and γ . We have presented the results of Monte Carlo simulations carried out to evaluate the performance of predictive coefficients. Additionally, to access the goodness-of-fit model we used the R_{LR}^2 and R_{FC}^2 measures. We consider different scenarios including misspecification of omitted covariates and negligence of varying dispersion, simultaneous increase in the number of covariates in the two submodels (mean and dispersion) and and negligence of nonlinearity.

In general form, the coefficients P^2 and $P_{\beta\gamma}^2$ perform similar and both enable to identify when the model are not reliable to predict or when is more difficult to make prediction. In particular it is noteworthy that when the response values are close to one or close to zero the power predictive of the model is substantially affected even under correct specification. In these situations, the R^2 statistics also revel that the model does not fit well.

Other important conclusion is about the bad performance of the R_{FC}^2 for beta regression models with varying dispersion, in sense that even when the model is well specified the values of this statistic tend to be too smaller than the values of the others statistics.

Finally, three empirical applications were performed and yield to a relevant information. In cases that the R^2 statistics evaluate the candidate models quite equally the predictive measures were decisive to choose the best between the candidate models. But as suggestion, to selected a model even in a predictive sense way it is also important to use goodness-of-fit measures and our recomendation for the class of nonlinear models is to use the version of R_{LR}^2 considered by Bayer and Cribari-Neto (2017) as the more appropriated model select criteria to linear beta regression models with varying dispersion.

Acknowledgement

This work was supported in part by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Fundação de Amparo à Ciência e Tecnologia de Pernambuco (FACEPE).

Conflict of Interest

The authors have declared no conflict of interest.

Appendix

Fisher's scoring iterative algorithm: In what follows we shall present the score function and Fisher's information for β and γ in nonlinear beta regression models (Simas et al., 2010). The log-likelihood function for model (2) is given by $\ell(\beta, \gamma) = \sum_{t=1}^n \ell_t(\mu_t, \phi_t)$, and $\ell_t(\mu_t, \phi_t) = \log \Gamma(\phi_t) - \log \Gamma(\mu_t \phi_t) - \log \Gamma((1 - \mu_t)\phi_t) + (\mu_t \phi_t - 1) \log y_t + \{(1 - \mu_t)\phi_t - 1\} \log(1 - y_t)$. The score function for β is

$$U_\beta(\beta, \gamma) = J_1^\top \Phi T(y^* - \mu^*), \quad (13)$$

where $J_1 = \partial \eta_1 / \partial \beta$ (an $n \times k$ matrix), $\Phi = \text{diag}\{\phi_1, \dots, \phi_n\}$, the t th elements of y^* and μ^* being given in (7). Also, $T = \text{diag}\{1/g'(\mu_1), \dots, 1/g'(\mu_n)\}$. The score function for γ can be written as $U_\gamma(\beta, \gamma) = J_2^\top H a$, where $J_2 = \partial \eta_2 / \partial \gamma$ (an $n \times q$ matrix), a_t is given in (9) and $H = \text{diag}\{1/h'(\phi_1), \dots, 1/h'(\phi_n)\}$. The components of Fisher's information matrix are

$$K_{\beta\beta} = J_1^\top \Phi W J_1^\top, \quad K_{\beta\gamma} = K_{\gamma\beta}^\top = J_1^\top C T H J_2^\top \quad \text{and} \quad K_{\gamma\gamma} = J_2^\top D J_2^\top. \quad (14)$$

Here, $W = \text{diag}\{w_1, \dots, w_n\}$, where

$$w_t = \phi_t v_t [1/\{g'(\mu_t)\}^2] \quad \text{and} \quad v_t = \{\psi'(\mu_t \phi_t) + \psi'((1 - \mu_t)\phi_t)\}. \quad (15)$$

Also, $C = \text{diag}\{c_1, \dots, c_n\}$; $c_t = \phi_t \{\psi'(\mu_t \phi_t) \mu_t - \psi'((1 - \mu_t)\phi_t)(1 - \mu_t)\}$, $D = \text{diag}\{d_1, \dots, d_n\}$; $d_t = \xi_t / (h'(\mu_t))^2$ and $\xi_t = \{\psi'(\mu_t \phi_t) \mu_t^2 + \psi'((1 - \mu_t)\phi_t)(1 - \mu_t)^2 - \psi'(\phi_t)\}$. To propose PRESS statistics for a beta regression we shall based on Fisher iterative maximum likelihood scheme and weighted least square regressions. Fisher's scoring iterative scheme used for estimating β , both to linear and nonlinear regression model, can be written as

$$\beta^{(m+1)} = \beta^{(m)} + (K_{\beta\beta}^{(m)})^{-1} U_\beta^{(m)}(\beta). \quad (16)$$

Where $m = 0, 1, 2, \dots$ are the iterations which are carried out until convergence. The convergence happens when the difference $|\beta^{(m+1)} - \beta^{(m)}|$ is less than a small, previously specified constant.

From (13), (14) and (16) it follows that the m th scoring iteration for β , in the class of linear and nonlinear regression model, can be written as $\beta^{(m+1)} = \beta^{(m)} + (J_1^\top \Phi^{(m)} W^{(m)} J_1)^{-1} J_1^\top \Phi^{(m)} T^{(m)}(y^* - \mu^{*(m)})$, where the t th elements of the vectors y^* and μ^* are given in (5). It is possible rewrite this equation in terms of weighted least squares estimator as $\beta^{(m+1)} = (J_1^\top \Phi^{(m)} W^{(m)} J_1)^{-1} \Phi^{(m)} J_1^\top W^{(m)} u_1^{(m)}$. Here, $u_1^{(m)} = J_1 \beta^{(m)} + W^{-1(m)} T^{(m)}(y^* - \mu^{*(m)})$. Upon convergence,

$$\hat{\beta} = (J_1^\top \hat{\Phi} \hat{W} J_1)^{-1} \hat{\Phi} J_1^\top \hat{W} u_1 \quad \text{where} \quad u_1 = J_1 \hat{\beta} + \hat{W}^{-1} \hat{T}(y^* - \hat{\mu}^*). \quad (17)$$

Here, \hat{W} , \hat{T} , \hat{H} and \hat{D} are the matrices W , T , H and D , respectively, evaluated at the maximum likelihood estimates. We note that $\hat{\beta}$ in (17) can be viewed as the least squares estimates of β obtained by regressing $\hat{\Phi}^{1/2} \hat{W}^{1/2} u_1$ on $\hat{\Phi}^{1/2} \hat{W}^{1/2} J_1$.

References

- Akaike, H. (1973). Information theory and an extension of the maximum likelihood principle. In *Second International Symposium on Information Theory (Tsahkadsor, 1971)*, pp. 267–281. Budapest: Akadémiai Kiadó.
- Allen, D. M. (1974). The relationship between variable selection and data augmentation and a method for prediction. *Technometrics* 16, 125–127.
- Bartoli, A. (2009). On computing the prediction sum of squares statistic in linear least squares problems with multiple parameter or measurement sets. *International Journal of Computer Vision* 85(2), 133–142.
- Bayer, F. M. and F. Cribari-Neto (2017). Model selection criteria in beta regression with varying dispersion. *Communications in Statistics, Simulation and Computation* 46, 720–746.
- Brascum, A. J., E. O. Johnson, and M. C. Thurmond (2007). Bayesian beta regression: applications to household expenditures and genetic distances between foot-and-mouth disease viruses. *Australian and New Zealand Journal of Statistics* 49(3), 287–301.
- Cepeda-Cuervo, E. and D. Gamerman (2005). Bayesian methodology for modeling parameters in the two parameter exponential family. *Estadística* 57, 93–105.

- Chien, L.-C. (2011). Diagnostic plots in beta-regression models. *Journal of Applied Statistics* 38(8), 1607–1622.
- Cook, R. D. and S. Weisberg (1982). *Residuals and Influence in Regression*. Chapman and Hall.
- Cribari-Neto, F. and A. Zeileis (2010). Beta regression in R. *Journal of Statistical Software* 34(2), 1–24.
- Espinheira, P., S. Ferrari, and F. Cribari-Neto (2008a). On beta regression residuals. *Journal of Applied Statistics* 35(4), 407–419.
- Espinheira, P. L., S. Ferrari, and F. Cribari-Neto (2014). Bootstrap prediction intervals in beta regressions. *Computational Statistics* 29(5), 1263–1277.
- Espinheira, P. L., S. L. P. Ferrari, and F. Cribari-Neto (2008b). Influence diagnostics in beta regression. *Computational Statistics and Data Analysis* 52, 4417–4431.
- Espinheira, P. L., E. G. Santos, and F. Cribari-Neto (2017). On nonlinear beta regression residuals. *Biometrical Journal* $n/a(n/a)$, $n/a-n/a$.
- Ferrari, S. and F. Cribari-Neto (2004). Beta regression for modelling rates and proportions. *Journal of Applied Statistics* 31(7), 799–815.
- Figuro-Zúñiga, J. I., R. B. Arellano-Valle, and S. L. Ferrari (2013). Mixed beta regression: A bayesian perspective. *Computational Statistics and Data Analysis* 61, 137–147.
- McCullagh, P. and J. A. Nelder (1989). *Generalized Linear Models* (2 ed.). London: Chapman and Hall.
- Mediavilla, F., L. F., and V. A. Shah (2008). A comparison of the coefficient of predictive power, the coefficient of determination and aic for linear regression. In K. JE (Ed.), *Decision Sciences Institute, Atlanta*, pp. 1261–1266.
- Nagelkerke, N. (1991). A note on a general definition of the coefficient of determination. *Biometrika* 78(3), 691–692.
- Ospina, R., F. Cribari-Neto, and K. L. Vasconcellos (2006). Improved point and interval estimation for a beta regression model. *Computational Statistics & Data Analysis* 51(2), 960 – 981.
- Palmer, P. B. and D. G. O’Connell (2009, sep). Regression analysis for prediction: Understanding the process. *Cardiopulmonary Physical Therapy Journal* 20(3), 23–26.
- Pregibon, D. (1981, 07). Logistic regression diagnostics. *The Annals of Statistics* 9(4), 705–724.
- Schwarz, G. (1978). Estimating the dimension of a model. *Annals of Statistics* 6(2), 461–464.
- Simas, A. B., W. Barreto-Souza, and A. V. Rocha (2010). Improved estimators for a general class of beta regression models. *Computational Statistics & Data Analysis* 54(2), 348–366.
- Smithson, M. and J. Verkuilen (2006). A Better Lemon Squeezer? Maximum-Likelihood Regression With Beta-Distributed Dependent Variables. *Psychological Methods* 11(1), 54–71.
- Spiess, A.-N. and N. Neumeyer (2010). An evaluation of r^2 as an inadequate measure for nonlinear models in pharmacological and biochemical research: a monte carlo approach. *BMC Pharmacology* 10(1), 6.
- Zerbinatti, L. (2008). Predição de fator de simultaneidade através de modelos de regressão para proporções contínuas. Msc thesis, University of São Paulo.