

Probing the No-Scale \mathcal{F} - $SU(5)$ One-Parameter Model via Gluino Searches at the LHC2

Tianjun Li,^{1,2} James A. Maxin,^{3,4} and Dimitri V. Nanopoulos^{5,6,7}

¹*CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,
Chinese Academy of Sciences, Beijing 100190, China*

²*School of Physical Sciences, University of Chinese Academy of Sciences, No.19A Yuquan Road, Beijing 100049, China*

³*Department of Chemistry and Physics, Louisiana State University, Shreveport, Louisiana 71115 USA*

⁴*Department of Physics and Engineering Physics,
The University of Tulsa, Tulsa, OK 74104 USA*

⁵*George P. and Cynthia W. Mitchell Institute for Fundamental Physics and Astronomy,
Texas A&M University, College Station, TX 77843, USA*

⁶*Astroparticle Physics Group, Houston Advanced Research Center (HARC), Mitchell Campus, Woodlands, TX 77381, USA*

⁷*Academy of Athens, Division of Natural Sciences,
28 Panepistimiou Avenue, Athens 10679, Greece*

In our recent paper entitled “The return of the King: No-Scale \mathcal{F} - $SU(5)$ ”, we showed that the model space supporting the most favorable phenomenology should have been probed in 2016 at the LHC2, with an even further reach into this region of the model in 2017–18. This ideal realm of the one-parameter version of No-Scale \mathcal{F} - $SU(5)$ yields a 1.9–2.3 TeV gluino mass at the very same point where the light Higgs boson mass enters its rather narrow experimentally determined range of $m_h = 125.09 \pm 0.24$ GeV. Given the recent results reported at Moriond 2017 for 36 fb^{-1} of luminosity collected in 2016 at the 13 TeV LHC2, we now update the status of the No-Scale \mathcal{F} - $SU(5)$ model space in light of the gluino mass exclusion limits presented by the ATLAS and CMS Collaborations. We illustrate that a resolution could be reached soon as to whether supersymmetry lives in this most critical region of the model space.

PACS numbers: 11.10.Kk, 11.25.Mj, 11.25.-w, 12.60.Jv

The ATLAS and CMS Collaborations at the LHC2 recorded 36 fb^{-1} of luminosity in 2016 at a center-of-mass energy of 13 TeV. In the search for supersymmetry (SUSY), the LHC2 in 2016 probed gluino (\tilde{g}) masses beyond 2 TeV, with current constraints in the multijet and multiple b-jet search regions residing at 1.92 TeV for ATLAS [1] and 1.96 TeV for CMS [2]. It is therefore expected that the LHC2 in 2017 will have at least a reach to a 2.0–2.3 TeV gluino mass. The 13 TeV beams will soon reenergize and begin recording proton-proton collision data again in June 2017, therefore, we use this present short pause in the SUSY search while the LHC2 prepares to launch its 2017 run to evaluate the model space of one prominent supersymmetric Grand Unified Theory (GUT) model against the full compilation of 2016 data statistics.

The supersymmetric GUT model No-Scale flipped $SU(5)$ with extra vector-like *flippon* multiplets, dubbed \mathcal{F} - $SU(5)$ for short, was shown in “The return of the King: No-Scale \mathcal{F} - $SU(5)$ ” [3] to possess the curious characteristic of generating an $m_{\tilde{g}} = 1.9 - 2.3$ TeV gluino mass at the very point in the model space where the light Higgs boson mass enters into its experimentally viable 1σ range of $m_h = 125.09 \pm 0.24$ GeV [4, 5]. In GUTs with gravity mediated SUSY breaking, otherwise known as the Minimal Supergravity (mSUGRA) models, the SUSY breaking soft terms can be represented by four universal terms: gaugino mass $M_{1/2}$, scalar mass M_0 , trilinear soft term A_0 , and the low-energy ratio of the Higgs vacuum expectation values (VEVs) $\tan\beta$, in addition to the sign

of the Higgs bilinear mass term μ . The most elemental No-Scale scenario requires $M_0 = A_0 = B_\mu = 0$ at the ultimate high-energy unification scale $M_{\mathcal{F}}$, where B_μ is the SUSY breaking soft term for the $\mu H_d H_u$ term in the superpotential. All the SUSY breaking soft terms therefore evolve down to low-energy from the single non-zero parameter $M_{1/2}$ at the $M_{\mathcal{F}}$ scale, hence, a true one-parameter model. Consequently, the entire SUSY spectrum will be proportional to $M_{1/2}$ at the leading order, yielding an invariance of the bulk “internal” physical properties under an overall rescaling. The complete SUSY spectrum thus maintains a piecewise smooth linear functional dependence upon $M_{1/2}$. A gluino mass of $m_{\tilde{g}} = 1.9 - 2.3$ TeV will therefore correlate to only a single specific confined range of light Higgs boson masses, which in the case of No-Scale \mathcal{F} - $SU(5)$ happens to be its observed value. In addition to reproducing the observed light Higgs boson mass, the $m_{\tilde{g}} = 1.9 - 2.3$ TeV gluino mass range under analysis here further produces the 9-year WMAP [6] and Planck 2015 [7] precision relic density 1σ measurements, while also conforming with empirical data for rare-decay processes, direct dark matter detection cross-sections, and proton decay lifetimes [3].

The accompanying vector-like multiplets, which we denote with the characterful term *flippons* when the multiplets are explicitly implemented within No-Scale flipped $SU(5)$, induce a few distinctive phenomenological signatures. As is known with No-Scale Supergravity, the light stau will be the Lightest Supersymmetric Particle (LSP) if the supersymmetry breaking scale is the tradi-




FIG. 1: The ATLAS 95% CL expected and observed exclusion limits of Ref. [1] with the linear model space of No-Scale \mathcal{F} - $SU(5)$ superimposed. The green strip of model space is consistent with the $\pm 1\sigma$ uncertainties on the light Higgs boson mass, top quark mass, and cold dark matter relic density. The yellow strip is that region of the model space satisfying the $\pm 2\sigma$ uncertainties on these same experiments.

tional GUT scale around 2.0×10^{16} GeV. Interestingly, the lightest neutralino with a Bino dominant component becomes the LSP in our model since the supersymmetry breaking scale is around the string scale, and then we can obtain the correct dark matter density via the light stau and LSP neutralino coannihilation [8, 9]. Moreover, we run the Renormalization Group Equations (RGEs) from the string scale to the $SU(3)_C \times SU(2)_L$ unification scale M_{32} , *i.e.*, around the traditional GUT scale. Therefore, at the traditional GUT scale, we shall have non-Universal supersymmetry breaking soft terms, unlike mSUGRA or the Constrained Minimal Supersymmetric Standard Model (CMSSM) [8, 9]. In addition, the one-loop beta function for the $SU(3)_C$ gauge symmetry receives a null value after inclusion of the flippons, *i.e.*, $b_3 = 0$, and thus the gaugino mass M_3 mirrors the vanishing b_3 via flat RGE evolution, as the conventional low-energy logarithmic

mic mass enhancement is suppressed. The outcome is a scaled-down gluino mass reduced below the first/second generation squarks, bottom squarks, and heavy stop, preserving a positive mass delta with only the light stop \tilde{t}_1 . Given this rather unique SUSY mass spectrum of $M(\tilde{t}_1) < M(\tilde{g}) < M(\tilde{q})$, where $\tilde{q} = \{\tilde{u}, \tilde{d}, \tilde{c}, \tilde{s}, \tilde{b}_1, \tilde{b}_2, \tilde{t}_2\}$, a compulsory gluino decay through an on-shell light stop of $\tilde{g} \rightarrow \tilde{t}_1 t \rightarrow t\bar{t} + \tilde{\chi}_1^0$ is imposed, triggering a hard 4-top multijet event via pair-produced gluinos and/or first/second generation squarks per $\tilde{q} \rightarrow q\tilde{g}$. The 4-top 100% branching fraction ensures a prolific accumulation of events with large numbers of hadronic jets, including tagged b-jets. This No-Scale \mathcal{F} - $SU(5)$ characteristic SUSY event topology compels a search region at the LHC2 inclusive of final states with multijet events and multiple b-jets. For a more detailed background of the No-Scale \mathcal{F} - $SU(5)$ model, please see Refs. [3, 10–13] and references therein.




FIG. 2: The CMS 95% CL expected and observed exclusion limits of Ref. [2] with the linear model space of No-Scale \mathcal{F} - $SU(5)$ superimposed. The green strip of model space is consistent with the $\pm 1\sigma$ uncertainties on the light Higgs boson mass, top quark mass, and cold dark matter relic density. The yellow strip is that region of the model space satisfying the $\pm 2\sigma$ uncertainties on these same experiments.

The analysis of the multijet and multiple b-jet SUSY search regions for the 36 fb^{-1} 2016 run at the LHC2 have been completed by ATLAS [1] and CMS [2]. Both studies focus on gluino pair-production leading to a 4-top signature by means of $\tilde{g} \rightarrow t\bar{t} + \tilde{\chi}_1^0$, as annotated in FIG. 1 and FIG. 2. The ATLAS 95% CL expected and observed exclusion limits of Ref. [1] are reproduced here in FIG. 1, showing the $\pm 1\sigma$ boundaries enclosing the expected and observed limits, where ATLAS refers to this search region as the Gtt simplified model. The CMS 95% CL expected and observed exclusion limits of Ref. [2] are also reproduced here in FIG. 2, displaying the $\pm 1\sigma$ uncertainties surrounding the expected and observed limits, where CMS refers to this search region as the T1tttt simplified model. We superimpose on the ATLAS and CMS exclusion plots in FIGs. 1–2 the linear model space of No-Scale

\mathcal{F} - $SU(5)$ as described in the prior paragraphs. We distinguish between the regions of the model space that satisfy the $\pm 1\sigma$ and $\pm 2\sigma$ experimental uncertainties for a set of experiments that comprise the light Higgs boson mass, top quark mass, and total cold dark matter relic density. The light Higgs boson mass central value and $\pm 1\sigma$ uncertainty is $m_h = 125.09 \pm 0.24 \text{ GeV}$ [4, 5], where for the $\pm 2\sigma$ limits we double the 1σ experimental uncertainty and add/subtract from the observed central value. For the top quark mass, we apply the PDG 2016 1σ world average of $m_t = 173.21 \pm 0.874 \text{ GeV}$ [14], also doubling the 1σ uncertainty for the 2σ limits. The relic density constraint we employ exploits both the 9-year WMAP [6] and Planck 2015 [7] measurements, using the WMAP9 1σ lower limit as our minimum value and the Planck 2015 1σ upper limit as our maximum value. This pro-

vides a relic density constraint of $\Omega h^2 = 0.1166 \pm 0.0032$. In FIGs. 1–2 the region of the model space satisfying the $\pm 1\sigma$ experimental constraints $\{m_h = 125.09 \pm 0.24, m_t = 173.21 \pm 0.874, \Omega h^2 = 0.1166 \pm 0.0032\}$ is highlighted in green, whereas all the points in the model space consistent with the $\pm 2\sigma$ experimental constraints $\{m_h = 125.09 \pm 0.48, m_t = 173.21 \pm 1.75, \Omega h^2 = 0.1166 \pm 0.0064\}$ are accentuated in yellow, which of course will further encompass the green $\pm 1\sigma$ strip as well.

The upper 1σ limit on the ATLAS and CMS exclusion limits delineated in FIGs. 1–2 lands at about 2030 GeV. Evidently, according to Ref. [1] the observed limits were significantly weaker than expected attributable to mild excesses observed in the relevant signal regions. The looser than anticipated limits are clearly exhibited in FIGs. 1–2, certainly right on the cusp of that precise region of No-Scale \mathcal{F} - $SU(5)$ compatible with the $\pm 1\sigma$ experimental uncertainties on the light Higgs boson mass, top quark mass, and relic density, as this region of No-Scale \mathcal{F} - $SU(5)$ was expected to be mostly excluded at 1σ in the 36 fb^{-1} 2016 run but could not due to the mild excesses noted above. Consequently, this realm of No-Scale \mathcal{F} - $SU(5)$ satisfying the $\pm 1\sigma$ empirical uncertainties remains viable and it is reasonable to conclude the region was probed in 2016 and surely the LHC2 will reach deeply into this region in 2017–18. We thus can confidently project that the presence of SUSY in this rather phenomenologically favorable dominion of No-Scale \mathcal{F} - $SU(5)$ should be adjudicated in the forthcoming couple of years, either to the delight or dismay of SUSY enthusiasts near and far.

ACKNOWLEDGMENTS

The primary numerical results for this project were computed at the Tandy Supercomputing Center, using dedicated resources provided by The University of Tulsa. Other supporting data for the theoretical background have been obtained via the HPC Cluster of ITP-CAS. This research was supported in part by the Projects 11475238 and 11647601 supported by the National Natural Science Foundation of China and Key Research Program of Frontier Sciences CAS (TL), and by DOE grant DE-FG02-13ER42020 (DVN).

- transverse momentum and multiple b-jets at $\sqrt{s} = 13$ TeV proton-proton collisions with the ATLAS detector,” (2017), ATLAS-CONF-2017-021.
- [2] A. M. Sirunyan et al. (CMS), “Search for supersymmetry in multijet events with missing transverse momentum in proton-proton collisions at 13 TeV,” (2017), 1704.07781.
- [3] T. Li, J. A. Maxin, and D. V. Nanopoulos, “The return of the King: No-Scale \mathcal{F} - $SU(5)$,” Phys. Lett. **B764**, 167 (2017), 1609.06294.
- [4] G. Aad et al. (ATLAS Collaboration), “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC,” Phys.Lett. **B716**, 1 (2012), 1207.7214.
- [5] S. Chatrchyan et al. (CMS Collaboration), “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC,” Phys.Lett. **B716**, 30 (2012), 1207.7235.
- [6] G. Hinshaw et al. (WMAP), “Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Parameter Results,” Astrophys. J. Suppl. **208**, 19 (2013), 1212.5226.
- [7] P. A. R. Ade et al. (Planck), “Planck 2015 results. XIII. Cosmological parameters,” Astron. Astrophys. **594**, A13 (2016), 1502.01589.
- [8] T. Li, J. A. Maxin, D. V. Nanopoulos, and J. W. Walker, “The Golden Point of No-Scale and No-Parameter \mathcal{F} - $SU(5)$,” Phys. Rev. **D83**, 056015 (2011), 1007.5100.
- [9] T. Li, J. A. Maxin, D. V. Nanopoulos, and J. W. Walker, “The Golden Strip of Correlated Top Quark, Gaugino, and Vectorlike Mass In No-Scale, No-Parameter \mathcal{F} - $SU(5)$,” Phys. Lett. **B699**, 164 (2011), 1009.2981.
- [10] T. Li, J. A. Maxin, D. V. Nanopoulos, and J. W. Walker, “The Ultrahigh jet multiplicity signal of stringy no-scale \mathcal{F} - $SU(5)$ at the $\sqrt{s} = 7$ TeV LHC,” Phys.Rev. **D84**, 076003 (2011), 1103.4160.
- [11] T. Li, J. A. Maxin, D. V. Nanopoulos, and J. W. Walker, “A Higgs Mass Shift to 125 GeV and A Multi-Jet Supersymmetry Signal: Miracle of the Flippons at the $\sqrt{s} = 7$ TeV LHC,” Phys.Lett. **B710**, 207 (2012), 1112.3024.
- [12] T. Li, J. A. Maxin, D. V. Nanopoulos, and J. W. Walker, “No-Scale \mathcal{F} - $SU(5)$ in the Light of LHC, Planck and XENON,” Jour.Phys. **G40**, 115002 (2013), 1305.1846.
- [13] T. Leggett, T. Li, J. A. Maxin, D. V. Nanopoulos, and J. W. Walker, “Confronting Electroweak Fine-tuning with No-Scale Supergravity,” Phys.Lett. **B740**, 66 (2015), 1408.4459.
- [14] C. Patrignani et al. (Particle Data Group), “Review of Particle Physics,” Chin. Phys. **C40**, 100001 (2016).

[1] T. A. collaboration (ATLAS), “Search for production of supersymmetric particles in final states with missing